Precision Editing of Cyclophilin A Generates Cyclosporine and Voclosporin Resistant Cellular Therapies

Short Title: CNI-Resistant Immune Effector Cells

Holly Wobma[1,2], Francesca Alvarez Calderon[2-4], Jiayi Dong[2,3], Kayleigh Omdahl[2,3], Xianliang Rui[3], Elisa J. Rojas Palato[3], Rene S. Bermea[2,5], Alexandre Albanese[3], Franziska Wachter[2-4], Marlana Winschel[3], Katherine A. Michaelis[1,2], William A. Burns[3], Gillian Selig[3], Lorenzo Cagnin[3], Victor Tkachev[2,6], Susan E. Prockop[2-4], Peter A. Nigrovic[1,2,7], Bruce R. Blazar[8], Ulrike Gerdemann[2-4]* and Leslie S. Kean[2-4]*

[1] Division of Immunology, Boston Children's Hospital, Boston, MA
[2] Harvard Medical School, Boston, MA
[3] Division of Hematology/Oncology, Boston Children's Hospital, Boston, MA
[4] Department of Pediatric Oncology, Dana-Farber Cancer Institute, Boston, MA
[5] Division of Pulmonary and Critical Care Medicine, Massachusetts General Hospital, Boston, MA
[6] Center for Transplantation Sciences, Massachusetts General Hospital, Boston, MA
[7] Division of Rheumatology, Inflammation, and Immunity, Brigham and Women's Hospital, Boston, MA
[8] Division of Pediatric Blood & Marrow Transplantation & Cellular Therapy, University of Minnesota, Minneapolis, MN

*co-senior author

Corresponding Author:

Leslie Kean, MD, PhD

Leslie.Kean@childrens.harvard.edu

Karp Research Building, 8th Floor, Room 08215

Phone: 617-919-1370

Correspondence should be addressed to leslie.kean@childrens.harvard.edu

Abstract

Recipients of allogeneic transplants or patients with autoimmune disease require immune

suppression, often with calcineurin inhibitors. There is an expanding repertoire of immune

effector cell therapies, including CD19 CAR-T cells and viral-specific T cells, deployed in these

patients; however, ongoing calcineurin inhibition may be detrimental to cell therapy function. We

developed a CRISPR/Cas9-based approach to engineer dual cyclosporine/voclosporin resistant

cell therapies by targeting *PPIA* (encoding cyclophilin A), a critical binding partner for both

drugs. Because Cyclophilin A has homeostatic functions in T cells, a complete knock-out is

detrimental to cell viability. We thus targeted its C-terminus, disrupting drug binding while

leaving the majority of the protein intact. C-terminal editing was stable throughout expansion

and preserved Cyclophilin A expression. Edited CD19 CAR-T cells retained effector function in

the presence of cyclosporine and voclosporin, including proliferation, cytokine production, and

target cell killing, resulting in improved survival in murine models of CD19+ leukemia. Edited

CMV-specific T cells also demonstrated preserved antigen-specific proliferation and cytokine

production in the presence of these drugs. C-terminal editing of Cyclophilin A offers a promising

avenue for developing next-generation cell therapies for patients receiving calcineurin inhibitors.

Introduction

Immunosuppression is the cornerstone of management of allo- and auto-immunity. Calcineurin

inhibitors (CNIs) are a key medication class used to prevent graft-vs-host-disease after

hematopoietic cell transplantation (HCT),[1] to prevent graft rejection after solid organ

transplantation (SOT),[2] and to treat autoimmune disease.[3] Standard CNIs include cyclosporine

(CsA) and tacrolimus, both widely used in HCT and SOT. Next-generation CNIs with more

favorable pharmacokinetic properties are also being developed, including the CsA derivative,

voclosporin (VCS), which is FDA approved for patients with lupus nephritis, expanding

application of this medication class.[4]

A key attribute of CNIs is their ability to inhibit lymphocyte proliferation, persistence, and

function. While these characteristics form the basis of their efficacy, they can be

disadvantageous for patients with allo- or auto-immune disease experiencing complications

such as relapse and infection, who are increasingly being treated with adoptive immune effector

cells (IEC). These IECs include chimeric antigen receptor (CAR)-T cells and viral-specific T cells

(VSTs), each of which may be inhibited by CNIs. Managing the delivery of IECs in these

patients is complex, as CNI withdrawal risks exacerbating allo- or auto-immunity.[5] Engineering

CNI-resistance is one way to optimize the potential of IEC therapies.

Calcineurin is a phosphatase that becomes active with binding of calcium/calmodulin

following T cell receptor (TCR) activation. Calcineurin mediates T cell activity in two ways: 1) it

dephosphorylates the transcription factor NFAT, which can then enter the nucleus and induce

expression of genes related to T cell activation (pro-inflammatory cytokines, co-stimulatory

proteins),[6] and 2) it removes an inhibitory phosphate from LCK that helps maintain upstream

TCR signaling. Both mechanisms further contribute to immune synapse formation and

enhanced cytotoxicity.[7] Additionally, calcineurin dephosphorylates synaptopodin in the kidney,

resulting in podocyte instability and increased proteinuria in lupus.[8] Thus, the therapeutic benefit

of CNIs derives from inhibition of T cell activation, with disease- and organ-specific effects also

observed.

CNIs rely on immunophilins to bind calcineurin and exert their inhibitory effect.

Immunophilins are a family of peptidyl-prolyl isomerases that are highly evolutionarily conserved

and widely expressed.[9] They can be divided into 'FK506 binding proteins' (FKBP) and

cyclophilins. Since tacrolimus and CsA/VCS can bind to calcineurin only after pairing with

FKBP12 and Cyclophilin A (CypA), respectively,[10] eliminating their expression represents a

rational approach for engineering CNI resistance. Indeed, tacrolimus resistance has been

achieved by knocking-out FKBP12.[11-15] In this case, knock-out also impacts the mTOR pathway,

conferring resistance to rapamycin that is inseparable from resistance to tacrolimus. Unlike for

FKBP12, knock-out of CypA in T cells has not been reported. This is likely due to the

widespread homeostatic functions of CypA in immune cells such that complete deletion may be

incompatible with T cell survival and function.[16-18]

To engineer CsA/VCS-resistant IECs, we therefore employed selective CRISPR/Cas9

editing to induce a frameshift in the last exon of *PPIA* (encoding CypA)*,* thereby disrupting the

CNI interaction site, and thus the inhibitory effect of CNIs on calcineurin, while leaving the core

CypA structure intact. Here, we demonstrate this targeted approach leads to resistance to both

CsA and VCS in otherwise fully functional CAR-T cells and VSTs.

Results

C-Terminal PPIA editing retains CypA expression

To confer CNI resistance to T cells, we selectively disrupted the C-terminal region of CypA,

which is critical for the interaction of CsA/VCS-CypA with calcineurin A.[19] This was achieved via

targeted CRISPR/Cas9 editing of the last exon (Exon 5) of *PPIA* in T cells from healthy donors

(**Fig. 1A**), a strategy that maintained CypA protein expression and T cell viability. As a

comparator, we edited the immediate upstream exon (Exon 4, **Fig. 1A**), an approach predicted

to lead to complete gene knock-out (NMDetective-A[20] score 0.78, with scores >0.52

corresponding to translation products predicted to undergo non-sense mediated decay

(NMD)).[20] With the Exon 4 edit, we observed high levels of early cell death compared to controls

(**Fig. 1B)**, rapid loss of the edited T cells during cell culture **(Fig. 1C),** and consequently,

minimal proliferation of edited cells **(Fig. 1D).** Notably, the Exon 5 approach was associated with

a low NMDetective-A score (0.23), low early cell death, and comparable growth kinetics to

PPIA^{WT} cells **(Fig. 1B-D),** including an average of 3.9 population doublings in the first 10 days

after CRISPR editing, supporting the feasibility of this engineering approach. Given the

promising results with our Exon 5 edit, and the predicted C-terminal disruption, we henceforth

refer to cells with the Exon 5 edit as '*PPIA^{△C}*'.

PPIA^{△C} cells demonstrated a high editing efficiency 89.5 (1.3)% measured on day 3 (**Fig.**

2A-B), predominantly comprised of a single nucleotide insertion (+1 base pair (bp) frameshift) in

58 (12)% of cells, followed by a single nucleotide deletion (-1 bp frameshift) in 26 (8%) of cells

(**Fig. 2C**). This editing efficiency and indel distribution was highly consistent across donors, and

edited cells demonstrated a stable editing efficiency and indel distribution over 3-4 weeks in

culture (**Fig. 2C**), indicating compatibility with cell survival. As hypothesized, CypA protein

expression was preserved when assessed at day 10, as demonstrated by flow cytometry,

Western blot, and confocal imaging (**Fig. 2D-F).** To confirm long term CypA protein expression

after *PPIA^{△C}* editing, Jurkat cells containing the dominant +1 bp indel were expanded for >8

weeks in culture demonstrated preserved editing efficiency (98% +1 bp indel) and CypA protein

expression (**Suppl. Fig. 1A**).

Based on the two predominant indels, we modeled predicted *PPIA^{△C}* protein structures

using AlphaFold Colab[22,23] and aligned them with the wildtype protein (*PPIA^{WT}* shown in purple;

Fig. 2G). As expected, the two predicted *PPIA^{△C}* protein structures align around important

residues for CypA function (pink)[19,24] but vary near Arg148 (green),[19] an amino acid critical for

the interaction of CypA with the CsA-calcineurin complex.[19] For the +1 bp indel protein, the

regional loss of structural homology was due to a frameshift after amino acid 132, with the

protein only minimally truncated (159 amino acids vs. wildtype of 165). For the -1 bp indel, the

frameshift led to a premature stop codon at amino acid 135 and thus a protein truncated prior to

Arg148.

To verify that cells with $PPIA^{\Delta C}$ retained wildtype T cell transcription patterns as well as

functional TCR signaling pathways – critical for any IEC therapy – we performed bulk RNA-seq

on purified CD8+ and CD4+ T cells expressing either $PPIA^{WT}$ (mock edited) or $PPIA^{\Delta C}$ at

baseline and upon stimulation with anti-CD3/CD28 Dynabeads. Unlike the significant

transcriptional differences between edited and wildtype cells previously reported for gene-

addition approaches to CsA resistance[25], the transcriptomes of the $PPIA^{WT}$ and $PPIA^{\Delta C}$ CD8+ T

cell populations were highly similar, with no differentially expressed genes, either at baseline or

after stimulation **(Fig. 2H-I, Suppl. Fig. 1B)**. This included comparable induction of NFAT

pathway related genes (CCL3, CCL4, IL2, TNF-α, IFN-γ) upon stimulation in both $PPIA^{WT}$ and

$PPIA^{\Delta C}$ cells (**Fig. 2H-I**). Similar bulk RNA-seq results were obtained from CD4+ T cells (**Fig.**

2J). Finally, $PPIA^{\Delta C}$ cells adopted similar maturation patterns (naïve, central memory, effector

memory) to $PPIA^{WT}$ cells after bead stimulation (**Suppl. Fig. 1C**).

$PPIA^{\Delta C}$ T cells are resistant to CsA and VCS

To determine whether $PPIA^{\Delta C}$ cells were resistant to CsA and VCS, we performed a mixed

lymphocyte reaction comparing division of $PPIA^{WT}$ and $PPIA^{\Delta C}$ responder T cells in the presence

of increasing doses of each agent (**Fig. 3**). This dose-escalation importantly revealed that lower

CsA doses are inhibitory *in vitro* relative to those commonly targeted in patients. As shown in

Fig. 3C, $PPIA^{WT}$ responder T cells exhibited a dose-dependent suppression of proliferation,

while $PPIA^{\Delta C}$ responder T cells were substantially resistant to CsA/VCS suppression across all

doses. For example, at a dose of 25 ng/mL CsA, $PPIA^{WT}$ responder CD4+ T cells demonstrated

69 (8)% suppression of division, whereas $PPIA^{\Delta C}$ responder CD4+ T cells demonstrated less

suppression (28 (12)%, p < 0.0001). Similarly, at a dose of 10 ng/mL VCS, $PPIA^{WT}$ responder

CD4+ T cells demonstrated 84 (5)% suppression of division by drug, whereas $PPIA^{\triangle C}$ responder

CD4+ T cells demonstrated less suppression (42 (8)%, p < 0.0001). These results were similar

for CD8+ responder T cells. In contrast, and consistent with the dependency of tacrolimus and

rapamycin on the alternative immunophilin, FKBP12, division of $PPIA^{\triangle C}$ and $PPIA^{WT}$ responder T

cells were equally suppressible by tacrolimus and rapamycin (**Fig. 3D**).

$PPIA^{\triangle C}$ CD19 CAR-T cell activation and proliferation is resistant to CsA and VCS

Having established CsA and VCS resistance in primary T cells, we next sought to validate drug

resistance in representative IEC therapies, beginning with CD19 CAR-T cells. Current FDA-

approved CD19 CAR-T cell products contain either 41BB or CD28 costimulatory domains

(denoted BBζ and 28ζ, respectively).[26] Given the distinct signaling pathway these costimulatory

molecules confer, we first evaluated the ability of CsA to inhibit proliferation of BBζ and 28ζ

CAR-T cells in $PPIA^{WT}$ CAR-T using CD19+ NALM6 $\beta 2M^{KO}$ cells as stimulators. As shown in

Fig. 4, both BBζ and 28ζ were suppressible by CsA but exhibited different drug sensitivities,

with BBζ-CAR-T more sensitive to CsA inhibition than 28ζ-CAR-T cells. For example, at a low

effector-to-target (E:T) ratio (1:1), there was 6 (1)% suppression of BBζ CD8+ CAR-T cell

division at the lowest CsA dose (12.5 ng/mL) and 74 (5)% suppression at the highest dose (100

ng/mL). For 28ζ CD8+ CAR-T cells at the 1:1 E:T ratio, there was no suppression at either drug

dose (6 (1)% and -6 (6)% respectively). However, high E:T ratios (10:1) rendered 28ζ CAR-T cells

inhibitable by CsA, with 46 (7)% suppression at a CsA dose of 25 ng/mL and 81 (9)%

suppression at a dose of 50 ng/mL. Trends were similar when CD4+ CAR-T cells were

assessed (**Suppl. Fig. 2**).

Next, we evaluated the proliferative potential of CRISPR/Cas9 $PPIA^{\triangle C}$ edited 28ζ CAR-T

cells in response to stimulation with NALM6 $B2M^{KO}$ cells (E:T 10:1). In vitro the $PPIA^{\triangle C}$ edit in

CAR-T cells led to near complete resistance to inhibition of proliferation by CsA and VCS (**Fig.**

5A-B and **Suppl. Fig. 3).** For example, at a CsA dose of 50 ng/mL, there was 45 (8)%

suppression of division of CD4+ $PPIA^{WT}$ CAR-T cells vs. 3 (1)% suppression of $PPIA^{\triangle C}$ cells (p <

0.0001). Similar findings were shown across different doses when cells were co-cultured in the

presence of VCS – for example, at a dose of 10 ng/mL, there was 41 (5)% suppression of

division of CD4+ $PPIA^{WT}$ CAR-T cells vs. 1 (2)% suppression of $PPIA^{\triangle C}$ cells (p < 0.0001).

Similar results were found for CD8+ CAR-T cells. To assess if $PPIA^{\triangle C}$ 28ζ CAR-T cells could

overcome inhibition of proliferation by CsA *in vivo*, NSG mice were injected with NALM6 cells to

provide an antigenic stimulus (**Fig. 5C**). As observed *in vitro*, there was marked inhibition of division

of $PPIA^{WT}$ CAR-T cells in mice treated with CsA, relative to mice treated with PBS (47 $\pm$ 10%

inhibition), whereas there was no inhibition of division by CsA for the $PPIA^{\triangle C}$ CAR-T cells (-24 $\pm$

17% inhibition, with the negative value indicating preserved expansion during the experiment, p <

0.0001).

To further define the impact of drug and drug resistance on CAR-T cell activation, we

evaluated cytokine production in response to a mitogen (PMA/ionomycin) or exposure to the

CAR target (NALM6 cells). Without addition of CsA, the fraction of cytokine-producing cells was

similar between $PPIA^{\triangle C}$ and $PPIA^{WT}$ CAR-T cells (**Suppl. Fig. 4**); however, in the presence of

100 ng/ml CsA, $PPIA^{\triangle C}$ CAR-T cells demonstrated sustained production of IFN-γ, TNF-α, and

IL-2 when compared with $PPIA^{WT}$ cells in both the CD4+ and CD8+ fractions. For example, in

the setting of mitogen stimulation (**Fig. 5D**), there was 38 (8)% suppression of IFN-γ induction in

$PPIA^{WT}$ CD8+ T cells in the presence of CsA, and less suppression of IFN-γ in the presence of

CsA for the $PPIA^{\triangle C}$ CAR-T cells (9 (4)% p < 0.0001). Similar results were observed for TNF-α,

with 80 (10)% suppression of TNF-α induction in $PPIA^{WT}$ CD8+ CAR-T cells versus 33 (3)% (p <

0.0001) for the $PPIA^{\triangle C}$ CAR-T cells. For IL-2, there was 89% (10%) suppression of IL-2 induction

in $PPIA^{WT}$ CD4+ CAR-T cells, versus 50 (13)% (p < 0.0001) for the $PPIA^{\triangle C}$ CAR-T cells. These

patterns were recapitulated when edited CAR-T cells were co-cultured with NALM6 cells as a

source of stimulation more specifically through the CAR (**Fig. 5E**).

While CsA inhibition of the NFAT pathway is commonly thought to mediate immune

suppression through disrupting cytokine production as above, there are many additional NFAT-

inducible genes that contribute to T cell activation related to costimulation and effector function. We

thus sought to determine the impact of CsA on CAR-T cell activation more broadly. As seen in **Fig.**

5F, after two day co-culture with NALM6 cells (4:1 E:T), both CD4+ and CD8+ CD19 CAR-T cells

induced upregulation of CD25, CD70, and ICOS, while CD4+ CAR-T cells also induced CD40L and

OX40, and CD8+ CAR-T cells increased expression of granzyme B (GZMB) and perforin (PRF1).

This induction was uniformly suppressed in the presence of 100 ng/mL CsA in $PPIA^{WT}$ CAR-T cells

but minimally – or not at all – in $PPIA^{\triangle C}$ CAR-T cells.

$PPIA^{\triangle C}$ CD19 CAR-T cells have superior tumor killing in the setting of CsA

We next evaluated CD19 CAR-T cell cytotoxicity towards two CD19 high-expressing

leukemia/lymphoma cell lines – wildtype NALM6 cells (NALM6 WT) and Raji cells (**Suppl. Fig. 5A**).

In a 3-day cytotoxicity assay, there was CsA-mediated suppression of CAR-T cell cytotoxicity

towards both cell lines (**Fig. 6A**). To assess if $PPIA^{\triangle C}$ CAR-T cells were also able to kill leukemia

cells with low CD19 antigen density, we assessed day 3 %killing of NALM6 cells with low CD19

expression (NALM6 CD19^{lo} (**Fig. 6A**)). In fact, in this experiment, $PPIA^{\triangle C}$ CAR-T cells overcame

CsA-induced suppression even by the 24-hour mark, and this enhanced cytotoxicity persisted

through day 3, as reflected by lower live NALM6 CD19^{lo} tumor burden over time (**Fig. 6B**). The long-

term cytotoxic advantage of $PPIA^{\triangle C}$ CAR-T cells in the presence of CsA was also observed when

they were cocultured with NALM6 WT cells for up to 7 days, with tumor rechallenge half-way

through (**Suppl Fig. 5B**). This sustained cytotoxic effect reflects a combination of preserved effector

function and proliferation of $PPIA^{\triangle C}$ CAR-T cells in the presence of CsA.

Next, we evaluated the *in vivo* anti-leukemic efficacy of $PPIA^{\triangle C}$ CAR-T cells in a xenograft

model (**Fig. 6C**). NSG mice received daily subcutaneous injections of PBS or CsA, leading to an

average trough of 350 ng/mL for the latter. CsA significantly inhibited tumor cell clearance by

$PPIA^{WT}$ CD19 CAR-T cells, as reflected by flux over time (**Fig. 6D**). For example, on Day +12, mice

treated with $PPIA^{WT}$ CAR-T cells had a flux of 2.1 (0.9) $\times$ 10^9 photons/s for those also receiving CsA

vs. 5.2 (4.5) $\times$ 10^8 photons/s for those receiving sham injections (p = 0.0001). By contrast, for mice

treated with $PPIA^{\triangle C}$ CAR-T cells, flux was comparable in groups receiving CsA vs. sham injections

(7.3 (4.0) $\times$ 10^8 vs. 5.7 (3.8) $\times$ 10^8 photons/s, respectively; p = 1.0) (**Fig. 6E-F).** Mice receiving CsA

had a significantly increased median survival if they were treated with $PPIA^{\triangle C}$ CAR-T cells vs.

$PPIA^{WT}$ CAR-T cells (**Fig. 6G**; p = 0.004). Notably, the $PPIA^{\triangle C}$ CAR-T cell + CsA group had similar

survival to the CAR-T cell groups receiving sham injections, whereas the $PPIA^{WT}$ CAR-T cell + CsA

group survival was not significantly different from mice that were untreated (Tumor Only and Tumor

Only + CsA groups). These experiments demonstrate a clear benefit of $PPIA^{\triangle C}$ CAR-T cells in

preserving *in vitro* and *in vivo* cytotoxic function, culminating in increased survival in a murine

leukemia model.

$PPIA^{\triangle C}$ CMV VSTs are resistant to CsA and VCS

We also evaluated the impact of the $PPIA^{\triangle C}$ edit on an additional IEC therapy - VSTs targeting

the cytomegalovirus (CMV) pp65 protein.[28] As shown in **Fig. 7A**, we generated CMV/pp65

specific VSTs that were 13-18% NLV-CMV dextramer positive after a 10-day exposure to pp65

PepMix. After re-stimulation with pp65 PepMix loaded PHA blasts, $PPIA^{\triangle C}$ VSTs demonstrated

CD8+ T cell-specific proliferation (**Fig. 7B**) that was better maintained in the presence of CsA

and VCS compared to $PPIA^{WT}$ cells (**Fig. 7C**). For example, for CD8+ VSTs, there was 100

(4)% suppression of division in the $PPIA^{WT}$ cells vs. 22 (10)% suppression of division with

$PPIA^{\triangle C}$ VSTs in the setting of 25 ng/mL CsA (p < 0.0001) and 101 (2)% suppression vs. 9

(15)%, respectively, in the setting of 5 ng/mL VCS (p < 0.0001). Additionally, as observed with

PPIA$^{\triangle C}$ CAR-T cells, a higher percentage of *PPIA*$^{\triangle C}$ VSTs produced IFN-γ, TNF-α, and IL-2

when stimulated with PMA/ionomycin or pp65 PepMix with CsA present (**Fig. 7D-E**).

Discussion

Here we report a targeted CRISPR/Cas9-based approach to generate CsA- and VCS-resistant

IECs by modifying the C-terminus region of CypA protein while preserving *PPIA* expression.

Gene edited CD19 CAR-T cells and VSTs preserved effector and proliferative functions during

CsA/VCS therapy, with important clinical implications for IEC use in immunosuppressed

patients. Our approach contains the key design feature of targeting the last exon of the *PPIA*

gene. This achieves two critical goals. The first is that edits in this region escape NMD, which is

generally not triggered when premature stop codons are found in the last exon.[20] By avoiding

NMD, we preserve translation of the majority of the CypA protein, an enzyme with several

critical functions in T cells. The second goal was to specifically induce a frameshift in the C-

terminus of CypA. This region is important for the interaction of CsA-CypA with calcineurin A

while being separate from the catalytic region of CypA, a domain critical for the performance of

its homeostatic functions as a peptidyl-prolyl isomerase.[29]

Our approach can be contrasted with prior strategies to engineer CNI-resistant cells. A

gene knock-out strategy targeting the tacrolimus- and rapamycin-dependent immunophilin,

FKBP12, has been previously published.[11-15,30] A similar knock-out approach has not been

published for *PPIA/*CypA. This may be due to intolerance of a full *PPIA/*CypA knock-out in T

cells, given the critical homeostatic functions to which CypA contributes. Indeed, in the Genome

Aggregation Database (gnomAD), which contains genetic information from >70,000 individuals,

the 'probability of loss-of-function intolerance' (pLI) score (ranges from 0 to 1 with 1 being loss-

of-function intolerant) is 0.98 for *PPIA*[31], implying even haploinsufficiency of this protein may be

selected against in the human population. Our results with the Exon 4 editing approach were

consistent with this, with negative selection of the edit in cultured T cells. Both FKBP12 and

CypA are conserved proteins with homeostatic functions. Most studies have described

FKBP12's role in calcium signaling in cardiomyocytes and neurons;[32,33] with its role in immunity

attributed to inhibition of TGF-β signaling.[18] CypA has been demonstrated to have multiple key

roles in normal T cell function: it is highly expressed in T cells and functions in protein folding,

cell survival, TCR signaling, infection, and inflammation.[9,24,34-39]

An alternative CsA-resistance approach to *PPIA* gene editing has been described, which

is to disrupt CsA-CypA docking with calcineurin A by gene addition of a modified calcineurin A

(mCNA).[25,40] However, a limitation observed with this strategy is that cells expressing the mCNA

have disrupted downstream NFAT signaling as well as other transcriptomic differences

compared to unmodified cells potentially impacting effector function.[25,40] For this reason, we

focused on selective gene modification of *PPIA*/CypA, to avoid both the loss of viability with a

total *PPIA* gene deletion and the abnormal NFAT signaling observed with the mCNA addition.

As predicted, we observed preserved cell survival and expansion, with stability of the Exon 5

gene edit for >3 weeks in primary T cells, and no detrimental impact on T cell activation or

calcineurin signaling.

As we consider application of the *PPIA*$^{\triangle C}$ edit to IEC therapies, the differential impact of

CsA on 28ζ versus BBζ CAR-T cells deserves discussion. It has been previously demonstrated

that strong CD28 co-stimulation may enable T cells to overcome CNI-based suppression,

possibly due to decreased NFAT export from the nucleus and enhanced PKCθ signaling.[41-44]

We confirmed this also applies to CAR-T cells, as BBζ CAR-T cells were more sensitive to CsA-

mediated inhibition than 28ζ CAR-T cells. However, the latter were still sensitive to CsA when

cultured with decreased tumor antigen stimulation and could be rendered resistant to CsA/VCS

by the *PPIA*$^{\triangle C}$ edit. These CsA and VCS resistant CAR-T cells may be considered for post-HCT

patients with early B-cell malignancy relapse, HCT or SOT patients with post-transplant

lymphoproliferative disorder, or patients with lupus nephritis, who are increasingly treated with

VCS.[45,46]

We also highlight the application of the *PPIA*$^{\triangle C}$ edit to VSTs. Viral reactivation is a

leading cause of morbidity and mortality in HCT and SOT patients.[47,48] In addition to triggering

organ dysfunction, there is a bidirectional relationship between viremia and graft-vs-host

disease/graft rejection.[49] Reduction in immune suppression may be used to treat viral

reactivation and disease, but further increases the risk of alloreactivity, and traditional antiviral

therapies may be insufficient to clear the viral infection and carry significant toxicities.[50] The

demonstration of substantial clinical responses with VSTs not resistant to

immunosuppression[51,52] provides a strong rationale for the utility of VSTs to treat otherwise

refractory viral diseases. The addition of CsA/VCS resistance holds the promise of increasing

the proliferation, peak expansion, and cytokine production of these cells in HCT and SOT

patients treated with CsA or VCS, which may further increase their efficacy.

In conclusion, we have developed a novel strategy for engineering T cell resistance to

CsA and VCS. This strategy can be incorporated into diverse cellular products for patients with

allo- and auto-immune disease who require ongoing immune suppression with these agents.

Materials and Methods

Immune Cell Isolation

Peripheral blood mononuclear cells (PBMCs) were isolated from healthy human donors using a

Ficoll gradient. A Pan-T cell isolation kit (Miltenyi) was used to isolate T cells. Cells were

cultured in X-VIVO15 (Lonza) media supplemented with 10% fetal bovine serum, 1%

penicillin/streptomycin, 2 mM of Glutamax, and 50 μM β-mercaptoethanol. IL-2 (R&D) was

added at 100 IU/mL for initial T-cell stimulation with 1:1 anti-CD3/28 Dynabeads (Gibco) and

lowered to 25 IU/mL once Dynabeads were removed. No cytokines were added during

functional assays.

CAR-T Cell Production

Plasmids encoding lentiviral constructs for CD19-CD28-CD3ζ and CD19-41BB-CD3ζ CARs

were designed through VectorBuilder. Viral packaging was performed as described.[53] Isolated T

cells were stimulated with anti-CD3/CD28 Dynabeads for 2 days prior to transduction.

Transduction was performed via spinoculation using Lentiboost (Mayflower Bioscience).[53]

Viral-Specific T Cell Production

PBMCs from HLA-A*0201 bearing CMV seropositive donors (n=3) were resuspended with 50

ng/million cells of pp65 PepMix (JPT). After a 60-minute incubation, cells were plated at 1

million cells/mL in a 24-well plate in media containing IL-4 and IL-7 (R&D; both at 10 ng/mL).[54]

Fresh media and cytokines were provided every three days. CMV specificity of expanded CD8+

T cells was assessed using the HLA-A*0201-specific dextramer NLVPMVATV (Immudex).

CRISPR/Cas9 Editing

T cells were CRISPR/Cas9 edited after two days of anti-CD3/CD28 Dynabead stimulation, and

CAR-T cells were edited two days after lentiviral transduction. CMV-pp65 peptide stimulated

PBMCs were edited three days after exposure to pp65 PepMix. CRISPR/Cas9 editing was

performed as described.[55] Guide RNAs (gRNA) were made by combining tracrRNA (IDT) with

negative control cRNA (IDT) or cRNAs for *PPIA* (Exon 4: ATCCTAAAGCATACGGGTCC; Exon

5: GTGTTTGGCAAAGTGAAAGA). DNA was sequenced by Genewiz. Synthego ICE Analysis

v3.0 was used to determine editing efficiency.

DNA and Protein Modelling

Frameshift variants were aligned in TCoffee[56] and viewed using Mview.[57] Structural predictions

were made using AlphaFold Colab,[22] with pdb sequences compared to wildtype CypA

(PDB:3k0m), viewed in Pymol v.2.5.5.

CypA Protein Validation

For flow cytometry, cells were fixed with BD Cytofix/Cytoperm and stained with CoraLite®594-

conjugated CypA monoclonal antibody (Proteintech; 1:50). For Western blot, membranes were

labelled with rabbit CypA polyclonal antibody (1:1000; Fisher; PA1-025) followed by an anti-

rabbit HRP secondary antibody (1:5000) using 3% BSA blocking buffer. The membrane was

stripped and re-stained with actin polyclonal antibody (PA5-78715). For confocal imaging, cells

were fixed with BD Cytofix/Cytoperm, stained with the same polyclonal CypA antibody at a

1:100 dilution followed by an AlexaFluor488 donkey anti-rabbit secondary at 1:250 and imaged

on a Zeiss 880 laser scanning confocal microscope.

RNA-seq

T cells from day 6 after CRISPR editing were either CD8 or CD4 selected using a magnetic

column (Miltenyi). Cells were kept unstimulated or re-stimulated with anti-CD3/CD28 beads for 4

hours and processed for bulk RNA-seq. Paired-end next generation sequencing was performed

using the Illumina HiSeq platform. Adapter and other Illumina-specific sequences were trimmed

using the Trimmomatic package (v0.39).[58] An alignment index was created using the Kallisto

package (v0.50.1)[59] and the GRCh38 cDNA reference transcriptome. Only "protein coding" and

"miRNA" transcripts were retained. Transcripts were normalized to 10,000 counts/sample and

transformed with the "numpy.log1p" function.[60] Clustermaps were created via the Seaborn

package (v0.13.2) with the linkage method set to "average" and distance set to "Euclidean".[61]

Drug Preparation

CsA and VCS (Millipore-Sigma) powders were reconstituted in dimethyl sulfoxide and diluted for

in vitro assays. Tacrolimus (Prograf 5 mg/mL) and rapamycin (LC laboratories) were also used

for *in vitro* assays. SandImmune® (Novartis) was used for *in vivo* proliferation assays.

Mixed Lymphocyte Reaction (MLR)

CRISPR/Cas9 edited T cell 'responders' were labelled with Cell Trace Violet (CTV;

Thermofisher) per manufacturer's instructions (modification: 1 μL/10 million cells). In a 96-well

U-bottom plate, 100,000 edited responder T cells were plated with 300,000 HLA-A*02 disparate,

irradiated (30 Gy) 'stimulator' PBMCs in a total volume of 200 μL. Cyclosporine, voclosporin,

tacrolimus, and rapamycin were added at indicated doses. Assays were read on day 5 after

staining for CD4 (L200), CD8 (RPA-T8), and HLA-A*02 (BB7.2). The %suppression by drug was

calculated by $100\% \times (1 - \frac{divided_{drug}}{divided_{no\ drug}})$.[62]

CAR-T and VST Proliferation Assays

Edited CAR-T cells, 10-12 days after initial stimulation, were stained with CTV and co-cultured

with a CD19 positive NALM6 (ATCC) line knocked-out for Beta-2-microglobulin[53] to focus on

CAR-driven proliferation (vs. TCR driven allo-proliferation). Cells were collected for analysis on

day 3. Edited VSTs were kept in culture for 10-12 days after initial stimulation, then labeled with

CTV, and re-stimulated by co-culture with pp65 PepMix loaded onto autologous PHA blasts.

Samples were collected for analysis on day 4.

CAR-T and VST Cytokine Stimulation Assay & NFAT inducible proteins

Protein Transport Inhibitor Cocktail (eBioscience) was added to prevent cytokine secretion. For

stimulation, either pp65 pepMix was added (for VSTs), NALM6 cells (for CAR-T), or

PMA/ionomycin as part of the Cell Stimulation Cocktail (eBioscience; for both VST & CAR-T

cells). Intracellular staining was performed after 4 hours using: FITC-anti-human TNF-α

(Mab11), BV421-anti-human IL-2 (MQ1-17H12), and BV711-anti-Human IFN-γ (B27). 25 ng/mL

of CsA was added for VST assays and 100 ng/mL for CAR-T cell assays. The %suppression

was calculated as $100\% \times (1 - \frac{\%postive_{drug}}{\%postive_{no\ drug}})$ for each cytokine. In separate studies to assess

expression of NFAT inducible proteins, CD19-CAR-T cells were co-culture with NALM6 cells at

a 4:1 ratio for 2 days and surface stained for CD25 (BC96), CD70 (Ki-24), ICOS (ISA-3), 41BB

(4B4-1), OX40 (ACT35), CD40L (24-31), and intracellularly stained for PRF1 (B-D48), GZMB

(QA16A02).

CAR-T Cytotoxicity Assay

Edited CAR-T cells were co-cultured with luciferase positive NALM6 WT, NALM6 CD19^{lo} (kindly

obtained from Dr. Robbie Majzner)[27], or Raji cells (ATCC) at the indicated E:T ratios and

durations. %Killing on Day 3 was calculated as $100\% \times (1 - \frac{NALM6_{coculture}}{NALM6_{alone}})$, where the numerator

and denominator reflect bioluminescent signal. For multi-day experiments, separate plates were

created for individual day read-outs, allowing the experiment to continue. Read-outs were

performed by adding luciferin reagent from the Bright-Glo Luciferase Assay System (Promega)

and measuring luminescence on a EnSpire Multimode Plate Reader (PerkinElmer).

In vivo studies

Nine-week-old female NOD.Cg-*Prkdc^{scid} Il2rg^{tm1Wjl}*/SzJ (NSG) mice were injected with 500,000

NALM6 CD19^{lo} cells on Day 0 by tail vein. On Day +2, mice started receiving either daily

subcutaneous injections of 15 mg/kg CsA or PBS (sham). On Day +4, they were injected with

250,000 CAR+ *PPIA^{WT}* or *PPIA$^{\triangle C}$* CD19-CAR-T cells. Bioluminescence imaging was performed on

Days +4, +8, +12, and +15 using D-luciferin (Perkin Elmer) at 150 mg/kg injected subcutaneously

and an IVIS imaging instrument. Radiance was calculated using the Living Image software. Mice

reached experimental end point when they lost $\geq$ 20% of body weight, developed a body

conditioning score of $\leq$ 2, developed clinical symptoms of respiratory distress, or developed

neurological symptoms of leukemic infiltration into the central nervous system (gait imbalance,

weakness, hind-limb paralysis). A total of n = 9-15 mice were used per group, as indicated, across

two experiments for statistical power.

For the *in vivo* proliferation study, healthy NSG mice were injected by tail vein with 5×10^5

NALM6 WT cells on Day 0 and 6×10^5 CD19-CAR-T cells on Day +4 after initial labelling with CTV.

As with prior, CsA or PBS was subcutaneously injected daily, starting on Day +2. On Day +8, blood

samples were collected via retro-orbital bleed and stained for murine and human CD45 to assess

for CAR-T cell division. The %suppression by drug was calculated by 100% $\times$

$(1 - \frac{Geometric\ Mean\ CTV_{no\ drug}}{Geometric\ Mean\ CTV_{drug}})$.

Data Analysis

Flow cytometry data were analyzed using FlowJo v.10.7.1. Data were graphed using GraphPad

Prism v.10, and statistical analysis performed with one-way or two-way ANOVA with multiple

comparisons (post-hoc tests specified with each Figure).

Data Availability Statement

Data can be requested from the authors by reaching out to Leslie.kean@childrens.harvard.edu. The RNA-seq data will be deposited on a public server upon publication.

Acknowledgments

This work was supported by National Institutes of Health R01-HL095791 (LSK), P01-HL158504 (LSK), U19-AI051731 (LSK), the Lupus Research Alliance (LSK, PAN), P30AR070253 (PAN), R01AR073201 (PAN), R01AR075906 (PAN), R01 HL11879 (BRB), R37 AI34495 (BRB), 2P01 CA065493 (BRB), NIAID T32AI007512-36 (HW), NICHD K12HD052896 Career Development Award (FAC), ASTCT New Investigator Award (VT), Be the Match Foundation/CIBMTR Amy Strelzer Manasevit Research Grant (VT).

Authorship Contributions

HW, FAC, UG, and LSK designed experiments. HW, JD, KO, XR, EJRP, AA, FW, MW, KAM, GS, WB and LC performed research. HW, RSB, AA, FAC, UG, and LSK analyzed data. HW, UG, and LSK wrote the manuscript. All authors critically reviewed and approved the content of the manuscript.

Declarations of Interest

HW, SEP, and LSK have equity in Regatta Bio. HW and LSK are inventors on a submitted (pending) patent related to the topic of this paper (PCT/US2024/054075). PAN is a consultant for Alkermes, Apollo, Century Therapeutics, Merck, Monte Rosa, Novartis, Pfizer, and has received grants from BMS and Pfizer. He holds equity in Edelweiss Immune, Inc. BRB reports research funding from Carisma Therapeutics and consulting fees from BlueRock Therapeutics, Atara Biotherapeutics, Sandoz, Legend Biotech, GentiBio Inc. and Affyxell Co., Ltd. UG possesses intellectual property rights related to AlloVir, including interests in royalties. SEP receives support for the conduct of clinical trials through Boston Children's Hospital from Atara, and Jasper, is an inventor of intellectual property related to development of third party viral specific T cells program with all rights assigned to Memorial Sloan Kettering Cancer Center, receives honoraria from Pierre Fabre as well as consulting fees from Atara Biotherapeutics, Ensomo, HEOR, Pierre Fabre, and VOR. She serves on the DSMB for Stanford University and NYBC. LSK is on the scientific advisory board for HiFiBio. She reports research funding from Tessera Therapeutics, EMD Serono, Gilead Pharmaceuticals, and Regeneron Pharmaceuticals. She reports consulting fees from Vertex and Santa Ana Bio. LSK reports grants and personal fees from Bristol Myers Squibb. Her conflict-of-interest with Bristol Myers Squibb is managed under an agreement with Harvard Medical School.

Keywords

Calcineurin inhibitor, cyclosporine, voclosporin, cyclophilin A, immune effector cell, CAR T cell

References

1. Hamilton, B.K. (2018). Current approaches to prevent and treat GVHD after allogeneic stem cell transplantation. Hematology Am Soc Hematol Educ Program *2018*, 228-235. 10.1182/asheducation-2018.1.228.

2. Nelson, J., Alvey, N., Bowman, L., Schulte, J., Segovia, M.C., McDermott, J., Te, H.S., Kapila, N., Levine, D.J., Gottlieb, R.L., Oberholzer, J., et al. (2022). Consensus recommendations for use of maintenance immunosuppression in solid organ transplantation: Endorsed by the American College of Clinical Pharmacy, American Society of Transplantation, and International Society for Heart and Lung Transplantation: An executive summary. Pharmacotherapy *42*, 594-598. 10.1002/phar.2718.

3.   Rovin, B.H., Teng, Y.K.O., Ginzler, E.M., Arriens, C., Caster, D.J., Romero-Diaz, J.,
Gibson, K., Kaplan, J., Lisk, L., Navarra, S., Parikh, S.V., et al. (2021). Efficacy and
safety of voclosporin versus placebo for lupus nephritis (AURORA 1): a double-blind,
randomised, multicentre, placebo-controlled, phase 3 trial. Lancet *397*, 2070-2080.
10.1016/S0140-6736(21)00578-X.
4.   Abdel-Kahaar, E., and Keller, F. (2023). Clinical Pharmacokinetics and
Pharmacodynamics of Voclosporin. Clin Pharmacokinet *62*, 693-703. 10.1007/s40262-
023-01246-2.
5.   Reshef, R., Vardhanabhuti, S., Luskin, M.R., Heitjan, D.F., Hadjiliadis, D., Goral, S.,
Krok, K.L., Goldberg, L.R., Porter, D.L., Stadtmauer, E.A., and Tsai, D.E. (2011).
Reduction of immunosuppression as initial therapy for posttransplantation
lymphoproliferative disorder(bigstar). Am J Transplant *11*, 336-347. 10.1111/j.1600-
6143.2010.03387.x.
6.   Martinez-Martinez, S., and Redondo, J.M. (2004). Inhibitors of the calcineurin/NFAT
pathway. Curr Med Chem *11*, 997-1007. 10.2174/0929867043455576.
7.   Dutta, D., Barr, V.A., Akpan, I., Mittelstadt, P.R., Singha, L.I., Samelson, L.E., and
Ashwell, J.D. (2017). Recruitment of calcineurin to the TCR positively regulates T cell
activation. Nat Immunol *18*, 196-204. 10.1038/ni.3640.
8.   Faul, C., Donnelly, M., Merscher-Gomez, S., Chang, Y.H., Franz, S., Delfgaauw, J.,
Chang, J.M., Choi, H.Y., Campbell, K.N., Kim, K., Reiser, J., et al. (2008). The actin
cytoskeleton of kidney podocytes is a direct target of the antiproteinuric effect of
cyclosporine A. Nat Med *14*, 931-938. 10.1038/nm.1857.
9.   Nigro, P., Pompilio, G., and Capogrossi, M.C. (2013). Cyclophilin A: a key player for
human disease. Cell Death Dis *4*, e888. 10.1038/cddis.2013.410.
10. Breuder, T., Hemenway, C.S., Movva, N.R., Cardenas, M.E., and Heitman, J. (1994).
Calcineurin is essential in cyclosporin A- and FK506-sensitive yeast strains. Proc Natl
Acad Sci U S A *91*, 5372-5376. 10.1073/pnas.91.12.5372.
11. Amini, L., Wagner, D.L., Rossler, U., Zarrinrad, G., Wagner, L.F., Vollmer, T.,
Wendering, D.J., Kornak, U., Volk, H.D., Reinke, P., and Schmueck-Henneresse, M.
(2021). CRISPR-Cas9-Edited Tacrolimus-Resistant Antiviral T Cells for Advanced
Adoptive Immunotherapy in Transplant Recipients. Mol Ther *29*, 32-46.
10.1016/j.ymthe.2020.09.011.
12. De Angelis, B., Dotti, G., Quintarelli, C., Huye, L.E., Zhang, L., Zhang, M., Pane, F.,
Heslop, H.E., Brenner, M.K., Rooney, C.M., and Savoldo, B. (2009). Generation of
Epstein-Barr virus-specific cytotoxic T lymphocytes resistant to the immunosuppressive
drug tacrolimus (FK506). Blood *114*, 4784-4791. 10.1182/blood-2009-07-230482.
13. Maldini, C.R., Messana, A.C., Bendet, P.B., Camblin, A.J., Musenge, F.M., White, M.L.,
Rocha, J.J., Coholan, L.J., Karaca, C., Li, F., Yan, B., et al. (2024). Immunosuppressant
therapy averts rejection of allogeneic FKBP1A-disrupted CAR-T cells. Mol Ther *32*,
3485-3503. 10.1016/j.ymthe.2024.06.022.
14. Peter, L., Wendering, D.J., Schlickeiser, S., Hoffmann, H., Noster, R., Wagner, D.L.,
Zarrinrad, G., Munch, S., Picht, S., Schulenberg, S., Moradian, H., et al. (2022).
Tacrolimus-resistant SARS-CoV-2-specific T cell products to prevent and treat severe
COVID-19 in immunosuppressed patients. Mol Ther Methods Clin Dev *25*, 52-73.
10.1016/j.omtm.2022.02.012.
15. Dragon, A.C., Bonifacius, A., Lienenklaus, S., Verboom, M., Gerhards, J.P., Ius, F.,
Hinze, C., Hudecek, M., Figueiredo, C., Blasczyk, R., and Eiz-Vesper, B. (2025).
Depletion of alloreactive B cells by drug-resistant chimeric alloantigen receptor T cells to
prevent transplant rejection. Mol Ther. 10.1016/j.ymthe.2025.01.009.
16. Aghdasi, B., Ye, K., Resnick, A., Huang, A., Ha, H.C., Guo, X., Dawson, T.M., Dawson,
V.L., and Snyder, S.H. (2001). FKBP12, the 12-kDa FK506-binding protein, is a

physiologic regulator of the cell cycle. Proc Natl Acad Sci U S A *98*, 2425-2430.
10.1073/pnas.041614198.

17. Dubois, S., Shou, W., Haneline, L.S., Fleischer, S., Waldmann, T.A., and Muller, J.R.
(2003). Distinct pathways involving the FK506-binding proteins 12 and 12.6 underlie IL-
2-versus IL-15-mediated proliferation of T cells. Proc Natl Acad Sci U S A *100*, 14169-
14174. 10.1073/pnas.2335979100.

18. Wang, T., Li, B.Y., Danielson, P.D., Shah, P.C., Rockwell, S., Lechleider, R.J., Martin, J.,
Manganaro, T., and Donahoe, P.K. (1996). The immunophilin FKBP12 functions as a
common inhibitor of the TGF beta family type I receptors. Cell *86*, 435-444.
10.1016/s0092-8674(00)80116-6.

19. Huai, Q., Kim, H.Y., Liu, Y., Zhao, Y., Mondragon, A., Liu, J.O., and Ke, H. (2002).
Crystal structure of calcineurin-cyclophilin-cyclosporin shows common but distinct
recognition of immunophilin-drug complexes. Proc Natl Acad Sci U S A *99*, 12037-
12042. 10.1073/pnas.192206699.

20. Lindeboom, R.G.H., Vermeulen, M., Lehner, B., and Supek, F. (2019). The impact of
nonsense-mediated mRNA decay on genetic disease, gene editing and cancer
immunotherapy. Nat Genet *51*, 1645-1651. 10.1038/s41588-019-0517-5.

21. Cradick, T.J., Qiu, P., Lee, C.M., Fine, E.J., and Bao, G. (2014). COSMID: A Web-based
Tool for Identifying and Validating CRISPR/Cas Off-target Sites. Mol Ther Nucleic Acids
3, e214. 10.1038/mtna.2014.64.

22. Mirdita, M., Schutze, K., Moriwaki, Y., Heo, L., Ovchinnikov, S., and Steinegger, M.
(2022). ColabFold: making protein folding accessible to all. Nat Methods *19*, 679-682.
10.1038/s41592-022-01488-1.

23. Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O.,
Tunyasuvunakool, K., Bates, R., Zidek, A., Potapenko, A., Bridgland, A., et al. (2021).
Highly accurate protein structure prediction with AlphaFold. Nature *596*, 583-589.
10.1038/s41586-021-03819-2.

24. Dorfman, T., Weimann, A., Borsetti, A., Walsh, C.T., and Gottlinger, H.G. (1997). Active-
site residues of cyclophilin A are crucial for its incorporation into human
immunodeficiency virus type 1 virions. J Virol *71*, 7110-7113. 10.1128/JVI.71.9.7110-
7113.1997.

25. Zhang, Y., Fang, H., Wang, G., Yuan, G., Dong, R., Luo, J., Lyu, Y., Wang, Y., Li, P.,
Zhou, C., Yin, W., et al. (2023). Cyclosporine A-resistant CAR-T cells mediate
antitumour immunity in the presence of allogeneic cells. Nat Commun *14*, 8491.
10.1038/s41467-023-44176-0.

26. Feucht, J., and Sadelain, M. (2020). Function and evolution of the prototypic CD28zeta
and 4-1BBzeta chimeric antigen receptors. Immunooncol Technol *8*, 2-11.
10.1016/j.iotech.2020.09.001.

27. Majzner, R.G., Rietberg, S.P., Sotillo, E., Dong, R., Vachharajani, V.T., Labanieh, L.,
Myklebust, J.H., Kadapakkam, M., Weber, E.W., Tousley, A.M., Richards, R.M., et al.
(2020). Tuning the Antigen Density Requirement for CAR T-cell Activity. Cancer Discov
10, 702-723. 10.1158/2159-8290.CD-19-0945.

28. Prockop, S.E., Hasan, A., Doubrovina, E., Dahi, P.B., Rodriguez-Sanchez, I., Curry, M.,
Mauguen, A., Papanicolaou, G.A., Su, Y., Yao, J., Arcila, M., et al. (2023). Third-party
cytomegalovirus-specific T cells improved survival in refractory cytomegalovirus viremia
after hematopoietic transplant. J Clin Invest *133*. 10.1172/JCI165476.

29. Zydowsky, L.D., Etzkorn, F.A., Chang, H.Y., Ferguson, S.B., Stolz, L.A., Ho, S.I., and
Walsh, C.T. (1992). Active site mutants of human cyclophilin A separate peptidyl-prolyl
isomerase activity from cyclosporin A binding and calcineurin inhibition. Protein Sci *1*,
1092-1099. 10.1002/pro.5560010903.

30.   Ricciardelli, I., Brewin, J., Lugthart, G., Albon, S.J., Pule, M., and Amrolia, P.J. (2013).
Rapid generation of EBV-specific cytotoxic T lymphocytes resistant to calcineurin
inhibitors for adoptive immunotherapy. Am J Transplant *13*, 3244-3252.
10.1111/ajt.12475.
31.   Chen, S., Francioli, L.C., Goodrich, J.K., Collins, R.L., Kanai, M., Wang, Q., Alfoldi, J.,
Watts, N.A., Vittal, C., Gauthier, L.D., Poterba, T., et al. (2024). A genomic mutational
constraint map using variation in 76,156 human genomes. Nature *625*, 92-100.
10.1038/s41586-023-06045-0.
32.   Guo, T., Cornea, R.L., Huke, S., Camors, E., Yang, Y., Picht, E., Fruen, B.R., and Bers,
D.M. (2010). Kinetics of FKBP12.6 binding to ryanodine receptors in permeabilized
cardiac myocytes and effects on Ca sparks. Circ Res *106*, 1743-1752.
10.1161/CIRCRESAHA.110.219816.
33.   Hoeffer, C.A., Tang, W., Wong, H., Santillan, A., Patterson, R.J., Martinez, L.A., Tejada-
Simon, M.V., Paylor, R., Hamilton, S.L., and Klann, E. (2008). Removal of FKBP12
enhances mTOR-Raptor interactions, LTP, memory, and perseverative/repetitive
behavior. Neuron *60*, 832-845. 10.1016/j.neuron.2008.09.037.
34.   Anto, N.P., Arya, A.K., Muraleedharan, A., Shaik, J., Nath, P.R., Livneh, E., Sun, Z.,
Braiman, A., and Isakov, N. (2022). Cyclophilin A associates with and regulates the
activity of ZAP70 in TCR/CD3-stimulated T cells. Cell Mol Life Sci *80*, 7.
10.1007/s00018-022-04657-9.
35.   Daneri-Becerra, C., Valeiras, B., Gallo, L.I., Lagadari, M., and Galigniana, M.D. (2021).
Cyclophilin A is a mitochondrial factor that forms complexes with p23 - correlative
evidence for an anti-apoptotic action. J Cell Sci *134*. 10.1242/jcs.253401.
36.   Bauer, K., Kretzschmar, A.K., Cvijic, H., Blumert, C., Loffler, D., Brocke-Heidrich, K.,
Schiene-Fischer, C., Fischer, G., Sinz, A., Clevenger, C.V., and Horn, F. (2009).
Cyclophilins contribute to Stat3 signaling and survival of multiple myeloma cells.
Oncogene *28*, 2784-2795. 10.1038/onc.2009.142.
37.   Colgan, J., Asmal, M., Neagu, M., Yu, B., Schneidkraut, J., Lee, Y., Sokolskaja, E.,
Andreotti, A., and Luban, J. (2004). Cyclophilin A regulates TCR signal strength in CD4+
T cells via a proline-directed conformational switch in Itk. Immunity *21*, 189-201.
10.1016/j.immuni.2004.07.005.
38.   Maneix, L., Iakova, P., Lee, C.G., Moree, S.E., Lu, X., Datar, G.K., Hill, C.T., Spooner,
E., King, J.C.K., Sykes, D.B., Saez, B., et al. (2024). Cyclophilin A supports translation of
intrinsically disordered proteins and affects haematopoietic stem cell ageing. Nat Cell
Biol *26*, 593-603. 10.1038/s41556-024-01387-x.
39.   Yurchenko, V., Constant, S., Eisenmesser, E., and Bukrinsky, M. (2010). Cyclophilin-
CD147 interactions: a new target for anti-inflammatory therapeutics. Clin Exp Immunol
160, 305-317. 10.1111/j.1365-2249.2010.04115.x.
40.   Brewin, J., Mancao, C., Straathof, K., Karlsson, H., Samarasinghe, S., Amrolia, P.J., and
Pule, M. (2009). Generation of EBV-specific cytotoxic T cells that are resistant to
calcineurin inhibitors for the treatment of posttransplantation lymphoproliferative disease.
Blood *114*, 4792-4803. 10.1182/blood-2009-07-228387.
41.   Isakov, N., and Altman, A. (2012). PKC-theta-mediated signal delivery from the
TCR/CD28 surface receptors. Front Immunol *3*, 273. 10.3389/fimmu.2012.00273.
42.   Diehn, M., Alizadeh, A.A., Rando, O.J., Liu, C.L., Stankunas, K., Botstein, D., Crabtree,
G.R., and Brown, P.O. (2002). Genomic expression programs and the integration of the
CD28 costimulatory signal in T cell activation. Proc Natl Acad Sci U S A *99*, 11796-
11801. 10.1073/pnas.092284399.
43.   June, C.H., Ledbetter, J.A., Gillespie, M.M., Lindsten, T., and Thompson, C.B. (1987). T-
cell proliferation involving the CD28 pathway is associated with cyclosporine-resistant

interleukin 2 gene expression. Mol Cell Biol *7*, 4472-4481. 10.1128/mcb.7.12.4472-
4481.1987.

44. Yokosuka, T., Kobayashi, W., Sakata-Sogawa, K., Takamatsu, M., Hashimoto-Tane, A.,
Dustin, M.L., Tokunaga, M., and Saito, T. (2008). Spatiotemporal regulation of T cell
costimulation by TCR-CD28 microclusters and protein kinase C theta translocation.
Immunity *29*, 589-601. 10.1016/j.immuni.2008.08.011.

45. Dall'Era, M., Kalunian, K., Solomons, N., Truman, M., Hodge, L.S., Yap, E., and
Askanase, A.D. (2024). Comparison of a voclosporin-based triple immunosuppressive
therapy to high-dose glucocorticoid-based immunosuppressive therapy: a propensity
analysis of the AURA-LV and AURORA 1 studies and ALMS. Lupus Sci Med *11*.
10.1136/lupus-2024-001319.

46. Arriens, C., Teng, Y.K.O., Ginzler, E.M., Parikh, S.V., Askanase, A.D., Saxena, A.,
Gibson, K., Caster, D.J., Atsumi, T., Lisk, L., Randhawa, S., et al. (2023). Update on the
Efficacy and Safety Profile of Voclosporin: An Integrated Analysis of Clinical Trials in
Lupus Nephritis. Arthritis Care Res (Hoboken) *75*, 1399-1408. 10.1002/acr.25007.

47. Hill, J.A., Mayer, B.T., Xie, H., Leisenring, W.M., Huang, M.L., Stevens-Ayers, T.,
Milano, F., Delaney, C., Sorror, M.L., Sandmaier, B.M., Nichols, G., et al. (2017). The
cumulative burden of double-stranded DNA virus detection after allogeneic HCT is
associated with increased mortality. Blood *129*, 2316-2325. 10.1182/blood-2016-10-
748426.

48. Limaye, A.P., Bakthavatsalam, R., Kim, H.W., Randolph, S.E., Halldorson, J.B., Healey,
P.J., Kuhr, C.S., Levy, A.E., Perkins, J.D., Reyes, J.D., and Boeckh, M. (2006). Impact
of cytomegalovirus in organ transplant recipients in the era of antiviral prophylaxis.
Transplantation *81*, 1645-1652. 10.1097/01.tp.0000226071.12562.1a.

49. Cantoni, N., Hirsch, H.H., Khanna, N., Gerull, S., Buser, A., Bucher, C., Halter, J., Heim,
D., Tichelli, A., Gratwohl, A., and Stern, M. (2010). Evidence for a bidirectional
relationship between cytomegalovirus replication and acute graft-versus-host disease.
Biol Blood Marrow Transplant *16*, 1309-1314. 10.1016/j.bbmt.2010.03.020.

50. Avery, R.K., Arav-Boger, R., Marr, K.A., Kraus, E., Shoham, S., Lees, L., Trollinger, B.,
Shah, P., Ambinder, R., Neofytos, D., Ostrander, D., et al. (2016). Outcomes in
Transplant Recipients Treated With Foscarnet for Ganciclovir-Resistant or Refractory
Cytomegalovirus Infection. Transplantation *100*, e74-80.
10.1097/TP.0000000000001418.

51. Mahadeo, K.M., Baiocchi, R., Beitinjaneh, A., Chaganti, S., Choquet, S., Dierickx, D.,
Dinavahi, R., Duan, X., Gamelin, L., Ghobadi, A., Guzman-Becerra, N., et al. (2024).
Tabelecleucel for allogeneic haematopoietic stem-cell or solid organ transplant
recipients with Epstein-Barr virus-positive post-transplant lymphoproliferative disease
after failure of rituximab or rituximab and chemotherapy (ALLELE): a phase 3,
multicentre, open-label trial. Lancet Oncol *25*, 376-387. 10.1016/S1470-2045(23)00649-
6.

52. Haque, T., Wilkie, G.M., Jones, M.M., Higgins, C.D., Urquhart, G., Wingate, P., Burns,
D., McAulay, K., Turner, M., Bellamy, C., Amlot, P.L., et al. (2007). Allogeneic cytotoxic
T-cell therapy for EBV-positive posttransplantation lymphoproliferative disease: results of
a phase 2 multicenter clinical trial. Blood *110*, 1123-1131. 10.1182/blood-2006-12-
063008.

53. Kaminski, J., Fleming, R.A., Alvarez-Calderon, F., Winschel, M.B., McGuckin, C., Ho,
E.E., Eng, F., Rui, X., Keskula, P., Cagnin, L., Charles, J., et al. (2024). B-cell-directed
CAR-T cell therapy activates CD8+ cytotoxic CARneg bystander T-cells in non-human
primates and patients. Blood. 10.1182/blood.2023022717.

54. Papadopoulou, A., Gerdemann, U., Katari, U.L., Tzannou, I., Liu, H., Martinez, C.,
Leung, K., Carrum, G., Gee, A.P., Vera, J.F., Krance, R.A., et al. (2014). Activity of

broad-spectrum T cells as treatment for AdV, EBV, CMV, BKV, and HHV6 infections
after HSCT. Sci Transl Med *6*, 242ra283. 10.1126/scitranslmed.3008825.

55.   Rupp, L.J., Schumann, K., Roybal, K.T., Gate, R.E., Ye, C.J., Lim, W.A., and Marson, A.
(2017). CRISPR/Cas9-mediated PD-1 disruption enhances anti-tumor efficacy of human
chimeric antigen receptor T cells. Sci Rep *7*, 737. 10.1038/s41598-017-00462-8.

56.   Di Tommaso, P., Moretti, S., Xenarios, I., Orobitg, M., Montanyola, A., Chang, J.M.,
Taly, J.F., and Notredame, C. (2011). T-Coffee: a web server for the multiple sequence
alignment of protein and RNA sequences using structural information and homology
extension. Nucleic Acids Res *39*, W13-17. 10.1093/nar/gkr245.

57.   Brown, N.P., Leroy, C., and Sander, C. (1998). MView: a web-compatible database
search or multiple alignment viewer. Bioinformatics *14*, 380-381.
10.1093/bioinformatics/14.4.380.

58.   Bolger, A.M., Lohse, M., and Usadel, B. (2014). Trimmomatic: a flexible trimmer for
Illumina sequence data. Bioinformatics *30*, 2114-2120. 10.1093/bioinformatics/btu170.

59.   Bray, N.L., Pimentel, H., Melsted, P., and Pachter, L. (2016). Near-optimal probabilistic
RNA-seq quantification. Nat Biotechnol *34*, 525-527. 10.1038/nbt.3519.

60.   Harris, C.R., Millman, K.J., van der Walt, S.J., Gommers, R., Virtanen, P., Cournapeau,
D., Wieser, E., Taylor, J., Berg, S., Smith, N.J., Kern, R., et al. (2020). Array
programming with NumPy. Nature *585*, 357-362. 10.1038/s41586-020-2649-2.

61.   Waskom, M.L. (2021). seaborn: statistical data visualization. J. Open Source Softw. *6*,
3021. https://doi.org/10.21105/joss.03021.

62.   Rui, X., Alvarez Calderon, F., Wobma, H., Gerdemann, U., Albanese, A., Cagnin, L.,
McGuckin, C., Michaelis, K.A., Naqvi, K., Blazar, B.R., Tkachev, V., et al. (2024). Human
OX40L-CAR-T(regs) target activated antigen-presenting cells and control T cell
alloreactivity. Sci Transl Med *16*, eadj9331. 10.1126/scitranslmed.adj9331.

Figure 1. CRISPR/Cas9 editing of Exon 5 but not Exon 4 facilitates expansion of edited
cells A) Schematic of the *PPIA* gene containing 5 exons, the largest of which are Exon 4 and
Exon 5. CRISPR/Cas9 gRNA cut sites are indicated, and guide sequences are provided in the
Methods. B) Significantly lower live T cell counts relative to the number of electroporated cells
(input) after CRISPR/Cas9 targeting of Exon 4, two days after gene editing. Control, $PPIA^{WT}$
cells were edited with a non-targeting guide. Groups compared via One-way ANOVA with
Tukey's post-hoc test. C) Change of Exon 4 and Exon 5 editing rates over time (relative to date
of CRISPR/Cas9 RNP nucleofection) as determined by Sanger sequencing on Days 3 (first time
point of assessing editing rate), 7 and 10. D) The degree of expansion of CRISPR-Cas9 edited
cells was calculated based on the live cell count multiplied by % edited cells relative to the Day
0 input cell count. The * at Day 3 reflects the comparison between $PPIA^{WT}$ and the Exon 4 edit.
The remaining Day 3 comparisons were not significantly different (ns). Experiments performed
with n = 3 donors per group; data represent mean (SD); error bars are not visible when smaller
than the marker size. *<0.05, **<0.01, ***<0.001, ****<0.0001
811
Figure 2. Characterizing the $PPIA^{\triangle C}$ edit and associated protein A) CRISPR editing at a
specific locus in Exon 5 leads to a consistent indel distribution of mostly +1 > -1 bp indels. B-C) The
overall editing efficiency and indel distribution are stable over 3-4 weeks (n = 3-9). D-F) CypA protein
expression 10 days after CRISPR/Cas9 editing by (D) flow cytometry and (E) Western blot using
actin as a loading control (after stripping the membrane and re-staining) and a Precision Plus
Protein™ Kaleidoscope™ ladder with corresponding protein sizes indicated on the left; no image
enhancement software was used. (F) Confocal imaging also confirming intracellular presence of
CypA. Scale bar represents 10 μm. G) Predicted 3D protein structure based on AlphaFold Colab
where the edited protein (rose or tangerine) is overlayed on the wildtype protein (purple). Catalytic
residues important to homeostatic function are highlighted in pink (R55, F60, M61, Q63, A101,
F113, W121, L122, and H126). An important residue for CsA binding is shown in green (R148; see
black arrow). Another black arrow points towards a β-strand absent in the edited proteins. The
corresponding amino acid sequences in Exon 5 are shown underneath as well as the CRISPR cut
site after amino acid 132. H) Comparison of the log fold change (LFC) in expression (unstimulated
vs. stimulated) in $PPIA^{WT}$ and $PPIA^{\triangle C}$ CD8+ T cells, analyzed by a simple linear regression model.
p<0.0001. Several NFAT-inducible genes are shown with purple markers. H-I) Normalized gene
expression based on bulk RNA-seq in $PPIA^{WT}$ and $PPIA^{\triangle C}$ (I) CD8+ T cells and (J) CD4+ T cells,
highlighting genes related to TCR activation and the NFAT signaling pathway. n = 3 donors. *<0.05,
<0.01, *<0.001, ****<0.0001
Figure 3. $PPIA^{\triangle C}$ T cell division is resistant to CsA and VCS suppression. A) Schematic of a
mixed lymphocyte reaction suppression assay in which 'stimulator' PBMCs are in co-culture with
'responder' $PPIA^{WT}$ or $PPIA^{\triangle C}$ T cells at a 3:1 ratio B) Gating strategy of assessing proliferating cells
in the mixed lymphocyte reaction: Stimulator and responder cells were HLA-A*02 disparate allowing
separation of populations on flow cytometry. Responder cells were gated on CD4+ and CD8+ T cell
subsets and proliferation was evaluated based on CTV dilution within these populations. C-D)
Division of Responder CD4+ and CD8+ T cells was assessed based on CTV dilution relative to
a "Responder only" negative control group across a (C) CsA and VCS dose escalation and (D)
tacrolimus and rapamycin dose escalation. Percent suppression by drug was calculated by
normalizing data to the drug-free (positive) control (see Methods). Black bars show the mean. 2-
way ANOVA with Sidak post-hoc test was used for statistical analysis. n = 6 replicates. *<0.05,
<0.01, *<0.001, ****<0.0001
Figure 4. BBζ CAR-T cells are more sensitive to inhibition by CsA than 28ζ CAR-T cells.
Division of CAR-T cells in response to CD19+ β2M^{KO} NALM6 stimulator cells was assessed

based on CTV dilution on Day 3, relative to unstimulated CAR-T cells (negative control). The
percent of divided cells was normalized to the drug-free (positive) control after subtracting non-
specific proliferation in the CAR-only (no tumor) negative controls for each construct (see
Methods. Panels are shown for different E:T ratios of A) 1:1 B) 2:1 C) 4:1 and D) 10:1 with a
CsA dose titration. Data shown reflects CAR-T cells gated on CD8. Black bars show the mean.
2-way ANOVA with Sidak post-hoc test used for statistical analysis. *<0.05, **<0.01, ***<0.001,
****<0.0001

Figure 5. $PPIA^{\Delta C}$ CAR-T cells exhibit preserved division, cytokine production, induction of
NFAT-inducible genes in the presence of drug. A) Edited CD19 CAR-T cells were stimulated by
$\beta 2M^{KO}$ NALM6 cells at a 10:1 E:T ratio. Division of CAR-T cells was assessed based on CTV
dilution on Day 3, comparing to unstimulated CAR-T cells (negative control). The percent of
divided cells was normalized to the drug-free (positive) control (see Methods). B) Data shown
for CD4+ and CD8+ CAR-T cell division across a CsA and VCS dose titration. Black bars show
the mean. n = 6 replicates. 2-way ANOVA with Sidak post-hoc test used for statistical analysis.
C) CTV labelled CD19 CAR-T cells were injected into immune deficient (NSG) mice previously
inoculated with human CD19+ NALM6 cells. Mice received daily subcutaneous injections of
CsA or PBS. Four days after infusion, CAR-T cell division was assessed based on CTV dilution
after gating on murine CD45-, human CD45+ cells. A two-tailed T-test was used for statistical
comparison (n = 5 per group). D-E) To assess cytokine production, $PPIA^{WT}$ or $PPIA^{\Delta C}$ CAR-T
cells were stimulated for 4 hours in the presence of protein transport inhibitor and either (D)
PMA/ionomycin or (E) CD19+ NALM6 cells (E:T of 2:1), and stained for intracellular IFN-γ, TNF-
α, and IL-2. 100 ng/mL CsA was used for experiments. The % Suppression was calculated as
described in Methods. F) CD19 CAR-T cells were co-cultured with NALM6 cells at a 4:1 ratio for 2
days and either surface or intracellularly stained for the indicated proteins in addition to CD4 or CD8
for initial gating.

Figure 6. $PPIA^{\Delta C}$ CAR-T cells exhibit enhanced cytotoxicity towards leukemia cells *in vitro*
and *in vivo*. A) Three-day cytotoxicity assay for which CD19 CAR-T cells were co-cultured with
luciferase positive NALM6 WT, Raji, or NALM6 CD19^{lo} cells at the indicated E:T ratios. B) The
NALM6 CD19^{lo} experiment shown in (A) now shown as luminescence over time as a reflection
of remaining tumor burden. RLU – Relative Light Units on the luminometer. Group comparisons
were made by 2-way ANOVA with multiple comparisons. C) Schematic for *in vivo* assay of
CAR-T cell killing of a NALM6 CD19^{lo} leukemia model. D) Absolute Flux over time, where Flux
saturates the detector in the 10^{10} range, such that it no longer tracks proportionally with
increasing tumor burden. E-F) Representative (E) images of mice in different experimental
groups from Day 12 with corresponding (F) Flux analyzed by One-Way ANOVA with multiple
comparisons. G) Kaplan-Meier survival curve, with curves compared using a log-rank test with
Holm-Sidak correction for multiple comparisons. In vivo studies include 9-15 mice per group,
pooling two experiments for Flux, and 5-10 mice per group for survival studies. *<0.05, **<0.01,
<0.001, *<0.0001

Figure 7. $PPIA^{\Delta C}$ VSTs exhibit preserved division and cytokine production in the presence of
drug. A) Schematic for making CRISPR/Cas9 edited CMV specific VSTs from healthy donor
PBMCs, of which 13-18% stained positive for HLA-A*02/NLVPMVATV after a first stimulation. B)
Representative example of CD8 specific T cell proliferation after restimulation of VST culture with
pp65-peptamer loaded PHA-blasts. C) % Suppression of $PPIA^{WT}$ and $PPIA^{\Delta C}$ VST proliferation in
the presence of 25 and 50 ng/ml CsA and 5 and 10 ng/ml VCS. D-E) VSTs were restimulated with
(D) PMA/ionomycin or (E) PepMix and stained for intracellular cytokine production. 25 ng/mL CsA

was used for these experiments. % Suppression was calculated as described in Methods. A 2-way

ANOVA was used for statistical analysis. *<0.05, **<0.01, ***<0.001, ****<0.0001

A

PPIA gene

Exon 1 · Intron · Exon 2 · Exon 3 · Exon 4 (Exon 4 gRNA cut site) · Exon 5 (Exon 5 gRNA cut site)

1000 · 2000 · 3000 · 4000

B

Live Cells Rel. to Input

PPIA^{WT} · Exon 5 · Exon 4

C

%Frameshift

Day 3 · Day 7 · Day 10

**** **** ****

D

Edited Live Cells Rel. to Input

Days

- *PPIA^{WT}*
- Exon 5
- Exon 4

A

B

C Top Indels Over Time

D

E

F *PPIA^WT*   *PPIA^ΔC*

G

| *PPIA^WT* | *PPIA^ΔC (major: +1bp)* | *PPIA^ΔC (minor: -1bp)* |

Exon 5 Sequence

	122		132		165
Wild Type	LDGKHVVFGKV	KEGMNIVEAMERFGSRNGKTSKKITIAD	C	GQLE	
+1 Asn	LDGKHVVFGKV	NRRHEYCGGHGALWVQEWQDQQ	EDHH	C	-----
+1 Lys	LDGKHVVFGKV	KRRHEYCGGHGALWVQEWQDQQ	EDHH	C	-----
-1	LDGKHVVFGKV	KKA	-----------------------		

CRISPR cut site

H CD8+ T cells

y = 0.90x + 0.01
R² = 0.87

I CD8+ T cells

J CD4+ T cells

A

CMV

pp65-spanning 15mers

+

PBMCs

CRISPR

IL-4, IL-7

13-18%

PPIA^{ΔC}
PPIA^{WT}
Unstained

CMV Dextramer (NLV...)

Proliferation/Cytokine Assays

10 day expansion

B **PHA Blast Loaded Peptamers**

CD4

CD8

CTV

CTV

+ PepMix
- PepMix

C **CD8 Responders**

%Suppression by CsA

CsA (ng/mL)

****   ****

25   50

CD8 Responders

%Suppression by VCS

VCS (ng/mL)

****   ****

5   10

PPIA^{WT}
PPIA^{ΔC}

D **CD8 VST**

% Suppression by CsA

*   ***   ***

PPIA^{WT} PPIA^{ΔC} PPIA^{WT} PPIA^{ΔC} PPIA^{WT} PPIA^{ΔC}

E **CD8 VST**

% Suppression by CsA

***   **   *

PPIA^{WT} PPIA^{ΔC} PPIA^{WT} PPIA^{ΔC} PPIA^{WT} PPIA^{ΔC}

IFN-γ
TNF-α
IL-2