1 **Glucocorticoid receptor and RUNX transcription factors cooperatively drive**
2 **CD8 T cells dysfunction in human cancer**
3

Christopher J Ward[1$], Soura Chakraborty[1$], Sanu K Shaji[1], Clara Veiga-Villauriz[1], Qiuchen

Zhao[1], Xi Chen[2] and Bidesh Mahata[1]*

1. Department of Pathology, University of Cambridge, Cambridge, CB2 1QP, UK

2. Shenzhen Key Laboratory of Gene Regulation and Systems Biology, Department of

Biology, School of Life Sciences, Southern University of Science and Technology, Shenzhen,

China

[$] Equal contribution

*To whom correspondence should be addressed: bm562@cam.ac.uk

Abstract

Glucocorticoids (GCs) are potent modulators of immune responses; however, the

mechanisms by which GCs regulate gene expression in human CD8 T cells remain

incompletely defined. Here, we delineate how physiological cortisol signalling shapes the

transcriptional and chromatin landscapes of primary human CD8 T cells. We identify a

substantial cohort of GC-responsive genes that are co-regulated through the cooperative

activity of the glucocorticoid receptor (GR) and RUNX transcription factors. Integrative RNA

sequencing and ChIP sequencing analyses identified genome-wide cortisol-responsive

immunoregulatory genes. Genetic deletion of the glucocorticoid receptor (GR, encoded by

NR3C1) abolished cortisol-induced gene expression changes, confirming GR-dependency.

Notably, GR chromatin occupancy in cortisol-treated CD8 T cells was strongly enriched at

RUNX transcription factor (TF) motifs rather than canonical glucocorticoid response elements

(GREs). Co-immunoprecipitation assays validated a ligand-dependent physical interaction

between GR and RUNX. Single-cell transcriptomic analyses of tumour infiltrating CD8 T cells

revealed significant enrichment of cortisol-responsive genes, indicating an active GC

signalling (response) within the tumour microenvironment. GR-RUNX dual controlled genes

were enriched in tumour-infiltrating CD8 T cells across multiple cancer types, including lung

adenocarcinoma, head and neck squamous carcinoma, pancreatic cancer, and breast cancer.

We found GR-RUNX co-regulated genes are mostly expressed in the exhausted CD8 T cell

population of different solid tumours These results suggest that local cortisol signalling within

tumour microenvironments drives CD8 T cell dysfunction through GR-RUNX TF cooperation.

Collectively, our findings identify RUNX TF as a critical mediator of GR signalling in human

CD8 T cells and reveal a novel mechanism by which endogenous glucocorticoids influence

antitumour immunity, which could be therapeutically targetable.

.

Introduction

Glucocorticoids (GCs) are steroid hormones widely recognised for their potent immunomodulatory and anti-inflammatory properties. Synthetic glucocorticoids, such as dexamethasone, betamethasone and prednisone, are extensively utilised in clinical settings to treat autoimmune disorders, inflammatory diseases, and haematological malignancies due to their ability to suppress immune cell activation and cytokine production[1,2]. Endogenous glucocorticoids, primarily cortisol in humans, also play critical roles in regulating immune homeostasis and responses under physiological and stress conditions[3]. Chronic or dysregulated GC signalling has been implicated in impaired antitumour immunity and cancer progression[2,4,5-7]. Despite extensive studies on synthetic GC-mediated immunosuppression, the precise molecular mechanisms by which physiological cortisol signalling modulates human CD8 T cell functions, particularly within tumour microenvironments, remain incompletely understood.

The glucocorticoid receptor (GR), encoded by the *NR3C1* gene, is a ligand-activated transcription factor belonging to the nuclear receptor superfamily. Upon ligand binding, GR translocate into the nucleus, binds chromatin at glucocorticoid response elements (GREs), and regulates target gene expression through direct DNA binding or tethering interactions with other transcription factors[8]. Although classical GRE-dependent mechanisms have been extensively characterised in various cell types, emerging evidence suggests that GR can also regulate gene expression through interactions with non-canonical transcriptional partners. GR can modulate gene expression in immune cells through protein–protein interactions with other transcription factors such as NF-κB and AP-1[9]. This "tethering" or "composite" model of gene regulation allows GR to repress or activate a distinct subset of genes indirectly, without binding DNA at traditional GRE motifs. For example, GR can bind to NF-κB p65 subunits, preventing NF-κB from inducing proinflammatory cytokine transcription. Conversely, it can cooperate with AP-1 to enhance the expression of anti-inflammatory mediators. Such interactions enable GR to modulate gene expression programs in a highly context-dependent manner[10,11]. Identification of these non-canonical GR partners in specific immune subsets is crucial for understanding the complex regulatory networks orchestrated by glucocorticoids in a physiological or pathological contexts of immune regulation.

CD8 T cells are central players in adaptive immunity, mediating cytotoxic responses against infected cells (such as virus infected) or cancer cells. Effective CD8 T cell responses are critical for tumour control and elimination. However, persistent antigen exposure and immunosuppressive signals within tumour microenvironments frequently drive CD8 T cell

dysfunction or exhaustion, limiting their antitumour efficacy[12,13]. In the context of CD8 T cells,

GCs have been repeatedly shown to suppress T cell effector programmes, including the

upregulation of checkpoint inhibitory proteins (CTLA-4, PD-1, LAG-3, TIM3) and the

downregulation of co-stimulatory molecules (CD2, CD28 & 41BB) in a context dependent

manner. Recent studies indicate that the local glucocorticoid signalling may contribute

significantly to immune suppression within tumour microenvironment by promoting T cell

dysfunction[2,4,5-7], yet the underlying transcriptional mechanisms remain largely unexplored.

RUNX transcription factors represent a family of evolutionarily conserved proteins crucially

involved in lineage specification, differentiation, proliferation, and functional regulation of

various immune cell subsets[14]. RUNX proteins (all three paralogues RUNX1-3) bind DNA

through conserved Runt domains (>95% sequence identity) at core consensus motif

YGYGGTY to regulate gene expression programs essential for hematopoietic development

and immune function. Among the RUNX family members (RUNX1-3), RUNX2 is

predominantly known for its roles in osteogenesis. However, recent evidence reveals its

broader involvement in immune regulation and cancer biology[15]. RUNX1 and RUNX3 are

critical regulators of CD8 T cell development and function. During thymopoiesis, they

collaborate to repress CD4 and enforce CD8 lineage commitment[16,17]. In mature CD8 cytotoxic

T lymphocytes (CTLs), RUNX3 dominates. It directly binds to the enhancers (e.g., *Cd8a*

enhancer E8I) to maintain CD8 expression[16,17] and activating effector genes like *Ifng, Gzmb*

(granzyme B), and *Prf1* (perforin) [16,17]. Runx3 deficiency severely impairs cytolytic activity and

proliferation[16,17], while also promoting tissue-resident memory (T_{RM}) CD8 T cells by

upregulating tissue-retention genes and suppressing recirculation[18]. Runx1 supports

thymocyte transitions and mature T cell homeostasis but plays a secondary role in peripheral

CTLs[19]. Runx2, though largely restricted to early thymic development, contributes to CD8

memory cell maintenance[1,18]. Together, Runx3 emerges as the central driver of CD8 T cell

identity, effector function, and tissue residency, while Runx1 and Runx2 provide

developmental and memory-related support. The potential interaction between GR signaling

and RUNX transcription factors in CD8 T cells has not been previously investigated.

In this study, we aimed to elucidate how physiological cortisol signalling modulates human

CD8 T cell function at the molecular level. Using integrative RNA sequencing (RNA-seq) and

ChIP sequencing (ChIP-seq) approaches combined with genetic perturbations and

biochemical validations, we demonstrate that cortisol profoundly reshapes the transcriptional

landscape of primary human CD8 T cells through ligand-dependent GR-RUNX cooperation.

We further provide single-cell transcriptomic evidence that this novel GR-RUNX regulatory

axis operates within tumour-infiltrating CD8 T cells across multiple cancer types. Our findings

identify RUNX as a critical non-canonical partner of GR in human CD8 T cells and suggest a previously unrecognised mechanism by which local endogenous glucocorticoid signalling within tumour influence antitumour immunity, offering new therapeutic opportunities aimed at enhancing cancer immunotherapy efficacy by targeting context-specific GR interactions.

Results

Cortisol reshapes the transcriptional landscape and signalling pathways in human activated CD8 T cells

To investigate how physiological cortisol influences the transcriptional profile of human CD8 T cells, we performed RNA-seq on *in vitro* activated primary CD8 T cells treated with cortisol (100 nM, 48 hours) or vehicle control (Figure 1a). Differential gene expression analysis identified 332 significantly regulated genes (170 upregulated, 162 downregulated) (Figure 1b). Among the most strongly induced genes were *TSC22D3, IL7R, FKBP5 and PIK3IP1*. Notably, *PIK3IP,* a negative regulator of PI3Ks, *and* amphiregulin (*AREG*), a cytokine implicated in tissue repair and immunosuppression, emerged as a novel cortisol-induced gene in CD8 T cells. A complete list of these differentially expressed genes can be accessed in the Supplementary Table S1. Gene set enrichment analysis (GSEA) using the Hallmark gene set collection revealed distinct patterns of pathway regulation following glucocorticoid treatment (Figure 1c, Supplementary Table S2). Significantly upregulated pathways include E2F targets, MYC targets, G2M checkpoint, and mTORC1 signalling, indicating enhanced cell cycle regulation and metabolic reprogramming. Conversely, downregulated pathways comprise inflammatory response, TNFα signalling via NF-κB, IL6/JAK/STAT3 signalling, KRAS signalling (both up and down gene sets), apoptosis, and IL2/STAT5 signalling. These data suggest cortisol enforces a metabolic checkpoint state in CD8 T cells, limiting inflammatory effector functions while simultaneously promoting cell survival and proliferation-associated pathways.

To confirm GR-dependency of these transcriptional changes, we employed CRISPR-Cas9-mediated knockout of the glucocorticoid receptor gene (*NR3C1*) in primary human CD8 T cells (Fig 1d). Loss of GR expression abolished cortisol-induced upregulation of target genes (e.g., *TSC22D3*, *IL7R, FKBP5, PIK3IP1 and AREG*), confirming direct GR involvement in mediating cortisol-driven transcriptional reprogramming (Figure 1e). A complete list of differentially expressed genes in NR3C1 knocked out CD8 T cell can be found in Supplementary Table S1.

In CD8 T cells, these 123 genes get upregulated in hydrocortisone treatment but downregulated when the glucocorticoid receptor gene (*NR3C1*) is knocked out. The Venn

diagram analysis demonstrated that 47 genes were exclusively upregulated by cortisol in a GR-independent manner (Mock HC vs. Veh), while 54 genes were uniquely upregulated in GR-KO cells independent of the ligand cortisol (Figure 1f). 123 genes were commonly upregulated in both conditions, suggesting GR-dependent mechanisms of cortisol-induced gene activation. Similarly, we found 85 genes get downregulated in cortisol treated CD8 T cell and get upregulated in *NR3C1* KO CD8 T cells. These findings identify cortisol induced true upregulated and downregulated genes in CD8 T cells. These findings also indicate that while a substantial proportion of cortisol-regulated genes require functional GR signalling, a subset of genes are expressed in a cortisol or GR-independent mechanisms.

To gain further insight into the functional implications of cortisol-induced transcriptional changes, we examined differentially expressed genes encoding transcription factors and various receptors (Figure 1g). The heatmap analysis revealed distinct expression patterns of transcription factors regulated by cortisol treatment in both GR-sufficient and GR-deficient conditions. Several transcription factors, including *ZBTB16, ELK3, TSC22D3 and CEBPB*, showed pronounced regulation following cortisol treatment. The receptor expression analysis identified multiple classes of differentially expressed receptors, including cytokine receptors (e.g., IL10RA, CCR7, IL7R, CXCR4), transporters (e.g., SLC3A2, SLC1A5), cell surface and adhesion receptors (e.g., PTGIR, ITGB2, ADGRG5), and nuclear receptors VDR and TSC22D3. Among these, several interleukin receptors showed predominantly GR-dependent regulation, while others exhibited mixed patterns of GR-dependent and independent expression changes. These findings highlight the complex interplay between GR-dependent and independent mechanisms in mediating the effects of cortisol on the transcriptional landscape of CD8 T cells, with potential implications for understanding glucocorticoid actions on immune function.

Together, these results demonstrate that physiological cortisol reshapes the transcriptional landscape of human CD8 T cells through GR-dependent regulation of immunomodulatory genes and pathways, highlighting a novel role for glucocorticoids in controlling T cell functionality via metabolic checkpoint enforcement.

Genome-wide chromatin occupancy reveals the direct target genes of GR and its noncanonical association with RUNX transcription factors

To elucidate the molecular mechanisms underlying cortisol-mediated transcriptional reprogramming in activated CD8 T cells, we performed chromatin immunoprecipitation followed by sequencing (ChIP-seq) to map genome-wide glucocorticoid receptor (GR) binding sites upon cortisol treatment. Primary human CD8 T cells from healthy donors were activated

in vitro and treated with cortisol (100 nM, 48 hours) or vehicle (Figure 2a). GR-bound

chromatin regions were identified by ChIP-seq. Cortisol treatment markedly increased GR

occupancy across the genome, identifying a total of 4871 high-confidence GR binding peaks

compared to vehicle-treated controls (Figure 2b, Supplementary Table S3). Motif enrichment

analysis of these GR-bound genomic regions surprisingly revealed no significant enrichment

of canonical glucocorticoid response elements (GREs), however, showed a striking

enrichment of non-canonical enrichment of RUNX transcription factor binding motifs upon

cortisol treatment (Figure 2b). Specifically, RUNX motifs were present in approximately 12.6%

of GR-bound peaks in cortisol-treated cells. Other transcription factor motifs such as TEAD

and NFAT also showed modest enrichment following cortisol stimulation (Figure 2b). These

data suggest that cortisol-induced GR chromatin occupancy in CD8 T cells preferentially

occurs at genomic loci harbouring non-canonical binding to other TFs sites, particularly RUNX

binding sites rather than canonical glucocorticoid response elements (GREs). In depth ChIP-

seq analysis of CD8 T cell identifies 1103 common peaks between two individual CD8 T cells

treated with hydrocortisone (Figure 2c), and 344 common peaks between vehicle-treated and

hydrocortisone treated CD8 T cell (Figure 2d). We found 81 direct targets of glucocorticoid by

integrating GR occupied genes (obtained from ChIP-seq) and glucocorticoid induced

upregulated genes (Figure 2e). We also observed out of 81 direct targets of GR 50 genes are

co-regulated by GR-RUNX (Figure 2f). Notably, genes exhibiting GR occupancy enriched with

RUNX motifs included critical immunoregulatory targets such as TSC22D3, IL7R, FKBP5,

PIK3IP1 and FAM53B, all significantly upregulated upon cortisol stimulation (Figure 2g).

Further genomic annotation analysis indicated that cortisol-induced GR binding predominantly

localised to intronic (31%) and promoter-proximal regions (31%), contrasting with classical

GRE-driven enhancer occupancy typically observed in other cellular contexts (Figure 2h).

ChIP-seq analysis also revealed the following genes that directly controlled by the GR binding

to the promoter or locus control region of the target genes. The list of GR direct target genes

can be found in Supplementary Table S4.

Ligand-dependent interaction between GR and RUNX3 orchestrates cortisol-induced

gene regulation in CD8 T cells

Given the prominent enrichment of RUNX motifs at GR-bound genomic regions (Figure 2), we

hypothesised that GR might physically interact with RUNX transcription factors in a ligand-

dependent manner. To test this hypothesis, first we compared the expression level of RUNX

transcription factor in CD8 T cells and found RUNX3 is predominantly expressed

(Supplementary Figure S1a). Next, we performed co-immunoprecipitation (co-IP) experiments

in HEK293T cells transiently expressing GR and RUNX3 in the presence or absence of cortisol

stimulation (100 nM, 36 hours). Immunoblot analysis demonstrated that RUNX3 robustly co-

precipitated with GR only upon cortisol treatment, confirming ligand-dependent formation of a
GR-RUNX2 protein complex (Figure 3a).

To assess the functional relevance of this interaction in primary human CD8 T cells, we
integrated our GR ChIP-seq, RNA-seq with previously publicly available RUNX target genes
datasets that were reported based on RUNX ChIP-seq and RNA-seq. First, we integrated
RUNX1 target genes and the genes that were cortisol responsive and enriched with RUNX 1
motif. Notably, 12 genes displayed robust cortisol-induced expression coupled with proximal
GR-RUNX motif occupancy, suggesting direct transcriptional co-regulation by GR and RUNX
TFs (Figure 3b). Heatmap view in the right-hand-side panel of Figure 3b shows their GC and
GR dependency. Next, we integrated RUNX2 target genes and distinguished the genes that
are upregulated (45 genes) or downregulated (7 genes) by the cortisol induced GR-RUNX
complex (Figure 3c, left panels). Heatmap (right panel) shows the expression of these genes
in different conditions. to show their dependence on GC and GR. We observed 45 genes co-
upregulated by GR-RUNX3 and 5 genes co-downregulated by GR-RUNX3 in CD8 T cells.
(Supplementary Figure 1b). All the gene names corresponding to this figure have been listed
in supplementary table S5.

Collectively, these results demonstrate that cortisol induces a physical and functional
cooperation between GR and RUNX3 in CD8 T cells. This ligand-dependent interaction
expands the regulatory repertoire of glucocorticoids beyond classical GRE-mediated
transcriptional control, highlighting a previously unrecognised molecular mechanism by which
endogenous glucocorticoids modulate human CD8 T cell function.

GR-dependent cortisol-induced gene signature is enriched in tumour-infiltrating CD8 T
cells across multiple cancer types
Given our findings that cortisol signalling profoundly reshapes the transcriptional landscape of
human CD8 T cells via GR-RUNX cooperation, we next explored whether this regulatory axis
is relevant in the tumour microenvironment (TME). To address this, we analysed publicly
available single-cell scRNA-seq datasets from tumour-infiltrating lymphocytes derived from
patients with breast cancer (Figure 4a), lung adenocarcinoma (Figure 4b), head and neck
squamous cell carcinoma (HNSCC) (Figure 4c) and pancreatic adenocarcinoma (Figure 4d).
We examined whether the cortisol-responsive gene signature identified in our primary human
CD8 T cell model (including key genes such as *TSC22D3, IL7R, FKBP5, PIK3IP1* and *AREG*)
was enriched within tumour-infiltrating CD8 T cells compared to matched tissue or peripheral
blood CD8 T cells. Strikingly, single-cell transcriptomic analyses revealed a consistent and
significant enrichment of cortisol-induced genes in tumour-infiltrating CD8 T cells across all

four cancer types examined (Figure 4a-d). Furthermore, tumour-infiltrating CD8 T cells
consistently expressed elevated levels of GR (*NR3C1*) compared to circulating counterparts,
suggesting enhanced sensitivity to local glucocorticoid signalling within the tumour
microenvironment.
262
These observations indicate that endogenous glucocorticoid signalling actively operates
within tumour microenvironments to modulate CD8 T cell function, potentially contributing to
an immunosuppressive state that favours tumour immune evasion. Collectively, these findings
underscore the clinical relevance of GR-mediated transcriptional regulation in shaping
immune responses within tumours and highlight the potential therapeutic value of targeting
GR to reinvigorate antitumour immunity.
269

Elevated expression of RUNX-GR co-regulated genes in tumour-infiltrating CD8 T cells
across cancer types
We investigated the expression patterns and functional significance of RUNX-GR co-regulated
genes in CD8 T cells across multiple cancer types. Single-cell transcriptomic profiling was
performed on CD8 T cells isolated from tumours and matched adjacent normal tissue or blood
in four major malignancies: triple-negative breast cancer (TNBC), lung cancer, head and neck
squamous cell carcinoma (HNSC), and pancreatic ductal adenocarcinoma (PDAC) (Figure
5a-d). RUNX-GR co-regulated genes exhibited significantly elevated expression in tumour-
infiltrating CD8 T cells compared to their non-tumour counterparts across all examined cancer
types (Figure 5a-d). The dot plots display both the proportion of cells expressing these genes
(represented by dot size) and the relative expression intensity (indicated by colour intensity),
with both parameters markedly increased in tumour-infiltrating populations. Whilst we
employed RUNX2-GR regulated genes for our analysis, it is important to note that this gene
set encompasses RUNX1-GR regulated genes, effectively representing broader RUNX-GR
regulation. This approach is substantiated by the established functional redundancy among
RUNX1, RUNX2, and RUNX3, which share a common core DNA-binding motif and frequently
regulate overlapping target genes.
287

To assess the functional relevance of this expression pattern, we calculated module scores
using the AddModuleScore method for individual CD8 T cells (Figure 5e-g). Higher module
scores, indicative of elevated RUNX-GR co-regulated gene expression, strongly correlated
with increased expression of established T cell exhaustion markers. This correlation was
consistent across TNBC (Figure 5e), lung cancer (Figure 5f), HNSC (Figure 5g) and PDAC
(Figure 5h), highlighting a conserved link between RUNX-GR transcriptional activity and the
exhausted phenotype in tumour-infiltrating CD8 T cells.

These findings indicate that RUNX-GR dependent gene regulation is particularly prominent within the tumour microenvironment, suggesting its potential role in shaping the phenotype of tumour-infiltrating CD8 T cells.

Single-cell trajectory analysis reveals an association between cortisol-responsive gene expression and CD8 T cell dysfunction in cancer

To investigate how cortisol signalling affects CD8 T cell differentiation within the tumour microenvironment, we performed single-cell trajectory inference analysis on CD8 T cells isolated from lung tumours and paired peripheral blood samples from patients with lung cancer. Using the Monocle algorithm, pseudo-temporal trajectories were constructed based on single-cell transcriptomic profiles (Supplementary Figure S2A). Cells expressing high levels of cortisol-responsive genes (TSC22D3, IL7R, FKBP5) were predominantly positioned along trajectory branches associated with progression towards terminal exhaustion or dysfunction, characterised by increased expression of inhibitory receptors, such as *PDCD1, CTLA4, HAVCR2, TIGIT, LAYN, and TOX* (Supplementary Figure S2B). In contrast, cells exhibiting lower cortisol-responsive gene signatures were primarily associated with early activation states. These results suggest that glucocorticoid signalling within the tumour microenvironment is activated shortly after CD8 T cells infiltrate the tumour, potentially driving them toward dysfunctional differentiation states.

Discussion

In this study, we reveal that physiological cortisol signalling fundamentally remodels the transcriptional and chromatin landscapes of human CD8 T cells through a previously unrecognised, ligand-dependent cooperation between the GR and RUNX transcription factors. By integrating RNA sequencing, ChIP sequencing, genetic perturbation, biochemical validation, and single-cell transcriptomic analyses of tumour-infiltrating lymphocytes, we identify a novel GR-RUNX regulatory axis that significantly modulates CD8 T cell function within the tumour microenvironment.

Our findings extend the current understanding of GR biology, demonstrating that GR can regulate gene expression via non-canonical interactions with RUNX transcription factors, rather than exclusively through classical glucocorticoid response elements (GREs). Traditionally, GR-mediated transcriptional regulation has been attributed to direct DNA binding at GREs or through tethered interactions with factors such as NF-κB and AP-1[1,8]. Here, we provide evidence that, in cortisol-treated human CD8 T cells, GR preferentially occupies genomic regions enriched for RUNX motifs. Co-immunoprecipitation experiments confirm a

direct, ligand-dependent interaction between GR and RUNX3, positioning RUNX3 as a critical non-canonical partner that enables GR to access distinct regulatory regions beyond those governed by classical GRE-driven mechanisms.

RUNX transcription factors are established regulators of immune cell differentiation and function[14]. Our data uncover an unexpected role for RUNX3 in human CD8 T cells as a pivotal mediator of cortisol-induced immunomodulation. Genetic ablation of GR expression abolishes cortisol-induced gene expression changes, confirming direct GR dependency. Moreover, cortisol-induced genes with proximal GR-RUNX motif occupancy include key immunoregulatory factors such as IL7R, FKBP5, TSC22D3, PIK3IP1 and AREG, implicating RUNX as a crucial determinant of glucocorticoid-driven immunomodulation.

A particularly intriguing observation from our study is the paradoxical regulation of signalling pathways by cortisol in CD8 T cells. Cortisol robustly suppressed inflammatory pathways (e.g., TNF-α/NF-κB and IL-2/STAT5 signalling) while simultaneously activating pathways associated with cell survival and proliferation (mTORC1 signalling, G2/M checkpoint). This dual regulatory pattern suggests that cortisol enforces a metabolic checkpoint state, limiting inflammatory effector functions while promoting cellular survival pathways. Such metabolic reprogramming may facilitate prolonged persistence of dysfunctional or exhausted T cells within tumours, thus contributing to impaired antitumour immunity.

Our findings reveal that RUNX-GR dependent gene regulation is particularly prominent within the tumour microenvironment across multiple cancer types. This tumour-specific upregulation suggests that local factors within the tumour might be driving this regulatory programme. One potential mechanism could involve local steroid biosynthesis within the tumour microenvironment inducing the glucocorticoid signalling pathway, which subsequently activates RUNX-GR-mediated transcriptional programs that influence the fate and function of tumour-infiltrating CD8 T cells. The co-regulation by RUNX factors and GR in tumour-infiltrating T cells is particularly intriguing given that both transcription factor families have been implicated in immune cell differentiation and function. RUNX family transcription factors are known to function redundantly, with RUNX1, RUNX2, and RUNX3 often binding to the same genomic sites and collaboratively regulating similar target genes. Similarly, GR has been shown to collaborate with pioneer factors and other transcription factors to regulate cell type-specific transcriptional programs. The observed correlation between RUNX-GR co-regulated gene expression and T cell exhaustion markers suggests that this transcriptional program may contribute to the development of the exhausted phenotype that characterises many tumour-infiltrating T cells. This finding provides new insights into the molecular mechanisms that may

underlie T cell dysfunction within the tumour microenvironment and identifies potential
therapeutic targets for enhancing anti-tumour immunity. Indeed, recent studies have
highlighted local GC production within tumours as an immunosuppressive mechanism
exploited by cancer cells to evade immune surveillance[2,42,4,5-7,20-25]. Our results provide
mechanistic insights into how this local steroid signalling axis may operate at the
transcriptional level through GR-RUNX cooperation leading T cell dysfunction.

From a translational perspective, our findings suggest novel therapeutic opportunities aimed
at improving cancer immunotherapy efficacy by targeting context-specific GR interactions.
Systemic inhibition of glucocorticoid signalling is challenging due to adverse effects; thus,
selective disruption of GR-RUNX3 interactions or downstream targets identified herein may
represent promising therapeutic strategies to enhance antitumour immunity without broadly
compromising physiological GC functions. Future studies should explore small-molecule
inhibitors or targeted genetic approaches to specifically disrupt GR-RUNX3 interactions in
preclinical tumour models.

In conclusion, our study identifies RUNX3 as a novel non-canonical partner of GR in human
CD8 T cells and reveals a previously unrecognised mechanism by which endogenous
glucocorticoids modulate antitumour immunity. Targeting context-specific transcription factor
interactions represents an exciting new avenue for therapeutic intervention aimed at reversing
steroid-driven immunosuppression within tumours.

Limitations of the study
Nevertheless, several questions remain unanswered: First, although we demonstrated robust
ligand-dependent GR-RUNX3 interaction *in vitro* and identified direct target genes through
integrative genomic analyses, further studies are required to clarify the precise molecular
determinants mediating this interaction at endogenous loci and its functional outcome
changing chromatin structure at the endogenous locus. Second, additional research is needed
to delineate the relative contributions of RUNX family members in mediating GC responses in
distinct immunological contexts. Finally, future work should validate our findings using *in vivo*
tumour models to assess whether disrupting GR-RUNX interactions can effectively restore
CD8 T cell functionality and enhance antitumour responses.

Methods
Human CD8 T cell culture
Peripheral blood samples were obtained from healthy male adult donors by informed consent.
Peripheral blood mononuclear cells (PBMCs) were recovered through density gradient

centrifugation by Ficoll-Paque Plus (GE Healthcare). CD8 T cells were isolated from PBMCs

by immunomagnetic negative selection (Stem Cell Technologies). CD8 T cells were cultured

for a total of 8 days in Immunocult-XF T Cell Expansion media (Stem Cell Technologies) with

50 µg/mL of Gentamicin (Gibco). Isolated cells were stimulated with plate bound anti-CD3

(Biolegend, Clone: OKT3, 1 µg/mL) and anti-CD28 (Biolegend, Clone: CD28.2, 1 µg/mL) for

72 hours. Next, CD8 T Cells were rested and expanded in fresh Immunocult media in the

presence of 100 U/mL of human IL-2 (Peprotech) for a further 72 hours. Cells were then re-

stimulated with plate bound anti-CD3 (Biolegend, Clone: OKT3 1 µg/mL) and anti-CD28

(Biolegend Clone: CD28.2 1 µg/mL) in the presence of 100 nM of cortisol [hydrocortisone (HC)

– Sigma] or vehicle control (methanol, Sigma) for 48 hours.

CRISPR-Cas9 mediated *NR3C1* deletion

Seventy-two hours after T cell activation, cells were washed with PBS and resuspended in

complete P3 Primary Cell Nucleofection Solution (Lonza) according to the manufacturer's

instructions. For each nucleofection, 1×10^6 cells were used. Two distinct ribonucleoprotein

(RNP) complexes were prepared, each targeting a specific sgRNA designed for the deletion

of NR3C1. The sequences for the sgRNAs were as follows:

1. sgRNA 1: 5' – CUUUAAGUCUGUUUCCCCG – 3'

2. sgRNA 2: 5'– AACCUCCAACAGUGACACCA – 3'

Both sgRNAs were synthesised by Synthego, and the Alt-R S.p. High Fidelity Cas9 Nuclease

V3 (Integrated DNA Technologies) was used to form the RNP complexes. Each RNP complex

was assembled at a 3:1 molar ratio of sgRNA to Cas9 (2.48 µM Cas9 + 7.52 µM sgRNA). The

prepared Cas9 RNP complexes were mixed with the cells and transferred to a 20 µL

Nucleocuvette Vessel. Electroporation was performed using a 4D Nucleofector with the EO-

115 program (Lonza). For control experiments, NR3C1-sufficient CD8 T cells were processed

in parallel and electroporated with High Fidelity Cas9 protein only (Mock Control). Immediately

following nucleofection, 80 µL of pre-warmed ImmunoCult-XF media (without supplements)

was added directly to the Nucleocuvette Vessel, and the cells were allowed to recover at 37°C

for 20 minutes. The cells were then transferred to a 24-well plate and cultured for 72 hours in

fresh ImmunoCult-XF media supplemented with 100 U/mL of IL-2 (PeproTech) and 50 µg/mL

of Gentamicin (Gibco™).

Single-cell RNA sequencing (scRNA-seq), clustering and visualisation

The previously published and publicly available scRNA-seq datasets used in this study were

as follows GSE164690[26-28], GSE161529[29,30], GSE127465[31], GSE155698[32,33]. The data has

been downloaded from NCBI GEO (https://www.ncbi.nlm.nih.gov/geo/) and processed with

the standard scRNA-seq integration pipeline in Seurat. To cluster the cells, we first identified

the top 2,000 most variable genes using the FindVariableFeatures function in Seurat. For cell
visualization, we applied the Uniform Manifold Approximation and Projection (UMAP)
algorithm to project the cells' PCA space representation into two dimensions. Cell clustering
in PCA space was performed with the Shared Nearest Neighbor (SNN) algorithm, using
Seurat V3's FindNeighbors and FindClusters functions. We then visualised the resulting
clusters in UMAP space with the DimPlot function. The differentially expressed genes in each
cluster were identified with the FindAllMarkers function in Seurat. The clusters have been
identified either by previous annotation or based on the Cluster Identity Predictor (CIPR)
program and has been validated based on marker gene expression from Panglo DB. Gene
expression levels were visualised using heatmaps, violin plots, and bar plots, created with the
pheatmap   package   (version   1.0.12)   (available   at   https://cran.r-
project.org/web/packages/pheatmap) and the dittoSeq R package (version 1.2.4). Data
scaling during visualization was performed automatically using the default settings of these
packages. Cell type information was retrieved from the meta.data slot of the downloaded
object. Gene module scores for subpopulations, as defined above, were calculated using the
Seurat function AddModuleScore.
459
Single-cell trajectory analysis
Single-cell trajectory analysis was performed using Monocle (version 2.20.0). Briefly, size
factors and dispersions were estimated with default settings. Differentially expressed genes
were identified using Monocle's differentialGeneTest function (q-value < 0.01). Identified
genes were used as ordering genes. Dimensionality was reduced using the DDRTree
algorithm, and trajectories were constructed by ordering cells based on pseudotime.
466
Western blot analysis
72 hours post nucleofection, CD8 T cells were washed once with ice cold PBS and lysed in
radioimmunoprecipitation assay (RIPA) buffer (Merck) with proteinase and phosphatase
inhibitors (Roche). Protein concentrations were determined by Pierce™ BCA Protein Assay
Kit and performed according to the manufacturer's instructions (Thermo-Scientific). Protein
samples (15 µg) were separated on a NuPAGE 4-12% Bis-Tris polyacrylamide gels
(Invitrogen), transferred onto nitrocellulose membranes (Bio-Rad) and probed using rabbit
anti-GR (Clone: D6H2L, Cell Signaling Technology) and mouse beta-actin (Clone: 8H10,
Origene) antibodies. Detection of immunoreactive bands was performed using secondary goat
anti-rabbit IRDye™ 680CW and goat anti-mouse IRDye™ 800CW antibodies and images
acquired on the Odyssey DLx imaging system (Li-COR).
478
479

Bulk RNA sequencing analysis

Cell pellets were layered with 40 µL of RNAlater Stabilization solution (Invitrogen) and stored

at -80°C prior to RNA extraction. Samples were homogenised using the QIAshredder Kit

(Qiagen) according to manufacturer's instructions. Next, total RNA was extracted using the

RNeasy Plus Mini Kit (Qiagen). RNA sequencing was performed in paired-end on a NextSeq

500 (Illumina) (HiSeq1500) platform, yielding 40 million reads per sample (Novogene, UK).

Raw sequencing reads were quality-controlled with FastQC (version 0.11.9), followed by

alignment to the reference genome gencode version 42 (Homo sapiens genome

hg38/GRCh38) using the alignment tool hisat2 (version 2.2.1). The sam files were further

inspected, filtered, and processed using samtools (version 1.13). HTseq counts obtained from

bam files were annotated and subjected to DESeq2 (version 1.34.0) analysis in R (version

4.1.1).. For DEseq2 analysis, for each gene, a log2 fold change was computed and the

significance level was assessed using the Wald test statistic with a p value and an adjusted p

value. Differentially expressed genes across different conditions have been identified using

the cutoff of a p-adjusted value of 0.05. The volcano plot has been constructed in GraphPad

Prism 9.5.1 based on the differentially expressed genes found in bulk RNA sequencing. After

differential expression analysis, genes were ranked according to their fold change and by

using the fGSEA algorithm, we performed Gene Set Enrichment Analysis (GSEA). The human

collection H (hallmark gene sets) from the Molecular Signatures Database (MSigDB) were

used to identify positively and negatively enriched pathways. All steps were performed in R

version 4.1.1.

Chromatin immunoprecipitation (ChIP) followed by sequencing

Chromatin immunoprecipitation (ChIP) was performed using the SimpleChIP Enzymatic

Chromatin IP Kit (Cell Signaling Technology) following the manufacturer's protocol with minor

modifications. Briefly, cells were crosslinked with 1% formaldehyde (Merck) in serum-free

RPMI-1640 for 10 min at room temperature (RT) and quenched with 1X glycine for 5 min.

Nuclei were isolated and chromatin digested with 0.3 µL micrococcal nuclease in 200 µL ChIP

Buffer at 37 °C for 20 min, followed by brief sonication (Branson 450-D sonicator, amplitude

36%, 4 × 5 s pulses on ice). Chromatin was immunoprecipitated overnight at 4 °C with 1 µg

each of anti-GR antibodies (PA1-511A, Thermo-Scientific; D6H2L, Cell Signaling

Technology). Immune complexes were captured with ChIP-Grade Protein G Magnetic Beads

(2 h, 4 °C), washed sequentially with low-salt and high-salt buffers, and eluted with 1X ChIP

Elution Buffer at 65 °C for 30 min. Crosslinks were reversed at 65 °C for 2 hours (5 M NaCl,

Proteinase K), with DNA purified and submitted for Illumina sequencing (Novogene,

Cambridge, UK).

ChIP-seq data analysis

Raw reads from fastq files were first trimmed using fastp 0.20.0 with the setting '-l 25 -- detect_adapter_for_pe'. Trimmed reads were mapped to the human genome hg38 by hisat2 2.1.0 with default setting and sorted by samtools 1.9 with flag '-ShuF 4 -q 30 -f 2'. After mapping, duplicated reads were removed by the picard tool (v.2.20.8) (http://broadinstitute.github.io/picard/). Peaks were then called using MACS2 and annotated using HOMER.

Integration of bulk RNA-seq and ChIP-seq data

After analysing bulk RNA-Seq and ChIP-Seq data, we compared the differentially expressed genes in RNA-seq of cortisol treated vs. vehicle control and all annotated peaks in ChIP-seq data. Downstream analysis was performed based on the genes between the RNA-seq and HC-treated ChIP-seq peaks, which include a total of 35 genes. The venn diagram to visualise the common genes was constructed using InteractiVenn. A heatmap has been drawn in gplots (version 3.1.3) in R version 4.1.1 to show the normalised counts of genes (transformed into row z-score) common in differentially expressed genes in RNA-seq and annotated peaks from cortisol (HC) treated cells. The protein-protein interaction (PPI) network has been constructed in STRING (version 11.5) and visualised in cytoscape (version 3.91). All 35 differentially expressed genes have been used as inputs in STRING, but only 8 genes constitute an intact network. The pie chart demonstrating the genomic position diversity of peaks obtained from cortisol treated cells in ChIP-seq data has been constructed in GraphPad Prism 9.5.1.

Overexpression of NR3C1 and RUNX3 in HEK-293T

A custom polycistronic plasmid (EF1A > hRUNX3 [NM_001320672.1] : IRES : hNR3C1 [NM_001364185.1] : IRES : EGFP) was designed and generated by VectorBuilder. Driven by the EF1A promoter, this construct produces a single mRNA transcript translated into RUNX3, NR3C1 (glucocorticoid receptor), and EGFP via IRES elements. HEK-293T cells were transfected with 5 µg of plasmid using Lipofectamine 3000 (Invitrogen) in serum-free Opti-MEM, following the manufacturer's protocol. Transfection was performed at 70–80% confluence. After 6 hours, 100nM of cortisol (Sigma-Aldrich) was added directly to the culture medium, and cells were incubated for an additional 48 hours to allow for cortisol-mediated effects.

Co-Immunoprecipitation (Co-IP)

Cells were harvested by washing with ice-cold PBS and collected by centrifugation at 300 × g for 5 minutes at 4 °C. Cell lysis was performed using Pierce IP Lysis Buffer (Thermo Scientific) supplemented with protease and PhosSTOP phosphatase inhibitors (Roche). Lysates were

incubated on ice for 30 minutes, followed by mild sonication (three 5 s pulses at 36%
amplitude) using a Branson 4D Sonicator. Lysates were cleared by centrifugation at
16,000 × g for 20 minutes at 4 °C. The supernatant was pre-cleared by incubation with 10 µL
of Protein A/G Plus-Agarose Beads (Santa Cruz) for 1 hour at 4 °C with rotation, followed by
centrifugation at 3,000 × g for 10 minutes. Clarified lysates were incubated overnight at 4 °C
with either anti-NR3C1 (D6H2L, Cell Signaling Technology; 1:100) or control rabbit IgG (Cell
Signaling Technology), followed by addition of 30 µL Protein A/G Plus-Agarose Beads and
further incubation for 2 hours at 4 °C. Beads were washed three times with lysis buffer, and
bound proteins were eluted in Laemmli sample buffer. Eluates were analysed by western
blotting using anti-RUNX1/AML1, RUNX3, and RUNX2 antibodies (EPR3099, Abcam).
564

Resource availability

Lead contact

Further information and requests for resources and reagents should be directed to and will be

fulfilled by the Lead Contact, Bidesh Mahata (bm562@cam.ac.uk)

Material availability

This study did not generate unique reagents.

Data and code availability

○ Data: All large datasets (RNA-seq and ChIP-seq) generated in this study will be

submitted to public repository.

○ Code: This paper does not report original code.

○ Additional information: Any additional information required to reanalyse the data

reported in this paper is available from the lead contact upon request.

Author contribution

CJW, SC, SKS, CVV, QZ and XC, performed the experiments, analysed, interpreted &

visualised the data. BM conceptualised and supervised the study. All authors read and

approved the draft manuscript before submission.

Acknowledgements

We would like to thank Amit Mandal, University of Oxford, for critical suggestion and advice

on RNA-seq data analysis. We used BioRender.com to generate the graphical illustrations

presented in this manuscript.

Funding

The work is supported by CRUK Career Development Fellowship (RCCFEL\100095), NSF-BIO/UKRI-BBSRC project grant (BB/V006126/1) and MRC project grant (MR/V028995/1).

Competing interests

The authors declare that they have no competing interests.

Data and materials availability

All newly generated data such as RNA sequencing and ChIP sequencing data used in the study will be available in the public database during revised submission stage.

Supplementary tables

Supplementary Table S1: Cortisol-induced genes in human CD8 T cell

Supplementary Table S2: Cortisol-induced hallmark pathways in CD8 T cell

Supplementary Table S3: Glucocorticoid binding peaks in CD8 T cell

Supplementary Table S4: Glucocorticoid direct target genes in CD8 T cell

Supplementary Table S5: Cortisol induced genes, ChIP-seq peaks, RUNX regulated genes

References

1. Cain, D.W., and Cidlowski, J.A. (2017). Immune regulation by glucocorticoids. Nat Rev Immunol *17*, 233-247. 10.1038/nri.2017.1.
2. Vandewalle, J., Luypaert, A., De Bosscher, K., and Libert, C. (2018). Therapeutic Mechanisms of Glucocorticoids. Trends Endocrinol Metab *29*, 42-54. 10.1016/j.tem.2017.10.010.
3. Quatrini, L., and Ugolini, S. (2021). New insights into the cell- and tissue-specificity of glucocorticoid actions. Cell Mol Immunol *18*, 269-278. 10.1038/s41423-020-00526-2.
4. Obradovic, M.M.S., Hamelin, B., Manevski, N., Couto, J.P., Sethi, A., Coissieux, M.M., Munst, S., Okamoto, R., Kohler, H., Schmidt, A., and Bentires-Alj, M. (2019). Glucocorticoids promote breast cancer metastasis. Nature *567*, 540-544. 10.1038/s41586-019-1019-4.
5. Acharya, N., Madi, A., Zhang, H., Klapholz, M., Escobar, G., Dulberg, S., Christian, E., Ferreira, M., Dixon, K.O., Fell, G., et al. (2020). Endogenous Glucocorticoid Signaling Regulates CD8(+) T Cell Differentiation and Development of Dysfunction in the Tumor Microenvironment. Immunity *53*, 658-671 e656. 10.1016/j.immuni.2020.08.005.
6. Mahata, B., Pramanik, J., van der Weyden, L., Polanski, K., Kar, G., Riedel, A., Chen, X., Fonseca, N.A., Kundu, K., Campos, L.S., et al. (2020). Tumors induce de novo steroid biosynthesis in T cells to evade immunity. Nat Commun *11*, 3588. 10.1038/s41467-020-17339-6.

7.   Chakraborty, S., Pramanik, J., and Mahata, B. (2021). Revisiting steroidogenesis and its
role in immune regulation with the advanced tools and technologies. Genes Immun
22, 125-140. 10.1038/s41435-021-00139-3.
8.   Weikum, E.R., Knuesel, M.T., Ortlund, E.A., and Yamamoto, K.R. (2017). Glucocorticoid
receptor control of transcription: precision and plasticity via allostery. Nat Rev Mol
Cell Biol *18*, 159-174. 10.1038/nrm.2016.152.
9.   De Bosscher, K., Beck, I.M., Dejager, L., Bougarne, N., Gaigneaux, A., Chateauvieux, S.,
Ratman, D., Bracke, M., Tavernier, J., Vanden Berghe, W., et al. (2014). Selective
modulation of the glucocorticoid receptor can distinguish between transrepression of
NF-kappaB and AP-1. Cell Mol Life Sci *71*, 143-163. 10.1007/s00018-013-1367-4.
10. Escoter-Torres, L., Greulich, F., Quagliarini, F., Wierer, M., and Uhlenhaut, N.H. (2020).
Anti-inflammatory functions of the glucocorticoid receptor require DNA binding.
Nucleic Acids Res *48*, 8393-8407. 10.1093/nar/gkaa565.
11. Franco, L.M., Gadkari, M., Howe, K.N., Sun, J., Kardava, L., Kumar, P., Kumari, S., Hu,
Z., Fraser, I.D.C., Moir, S., et al. (2019). Immune regulation by glucocorticoids can be
linked to cell type-dependent transcriptional responses. J Exp Med *216*, 384-406.
10.1084/jem.20180595.
12. Wherry, E.J., and Kurachi, M. (2015). Molecular and cellular insights into T cell
exhaustion. Nat Rev Immunol *15*, 486-499. 10.1038/nri3862.
13. Philip, M., and Schietinger, A. (2022). CD8(+) T cell differentiation and dysfunction in
cancer. Nat Rev Immunol *22*, 209-223. 10.1038/s41577-021-00574-3.
14. Seo, W., and Taniuchi, I. (2020). The Roles of RUNX Family Proteins in Development of
Immune Cells. Mol Cells *43*, 107-113. 10.14348/molcells.2019.0291.
15. Mevel, R., Draper, J.E., Lie, A.L.M., Kouskoff, V., and Lacaud, G. (2019). RUNX
transcription   factors: orchestrators   of   development.   Development   *146*.
10.1242/dev.148296.
16. Korinfskaya, S., Parameswaran, S., Weirauch, M.T., and Barski, A. (2021). Runx
Transcription Factors in T Cells-What Is Beyond Thymic Development? Front Immunol
12, 701924. 10.3389/fimmu.2021.701924.
17. Woolf, E., Xiao, C., Fainaru, O., Lotem, J., Rosen, D., Negreanu, V., Bernstein, Y.,
Goldenberg, D., Brenner, O., Berke, G., et al. (2003). Runx3 and Runx1 are required for
CD8 T cell development during thymopoiesis. Proc Natl Acad Sci U S A *100*, 7731-7736.
10.1073/pnas.1232420100.
18. Milner, J.J., Toma, C., Yu, B., Zhang, K., Omilusik, K., Phan, A.T., Wang, D., Getzler, A.J.,
Nguyen, T., Crotty, S., et al. (2017). Runx3 programs CD8(+) T cell residency in non-
lymphoid tissues and tumours. Nature *552*, 253-257. 10.1038/nature24993.
19. Egawa, T., Tillman, R.E., Naoe, Y., Taniuchi, I., and Littman, D.R. (2007). The role of the
Runx transcription factors in thymocyte differentiation and in homeostasis of naive T
cells. J Exp Med *204*, 1945-1957. 10.1084/jem.20070133.
20. Taves, M.D., Otsuka, S., Taylor, M.A., Donahue, K.M., Meyer, T.J., Cam, M.C., and
Ashwell, J.D. (2023). Tumors produce glucocorticoids by metabolite recycling, not
synthesis, and activate Tregs to promote growth. J Clin Invest *133*. 10.1172/JCI164599.
21. Ahmed, A., Reinhold, C., Breunig, E., Phan, T.S., Dietrich, L., Kostadinova, F., Urwyler,
C., Merk, V.M., Noti, M., Toja da Silva, I., et al. (2023). Immune escape of colorectal
tumours via local LRH-1/Cyp11b1-mediated synthesis of immunosuppressive
glucocorticoids. Mol Oncol *17*, 1545-1566. 10.1002/1878-0261.13414.

22. Michalek, S., Goj, T., Plazzo, A.P., Marovca, B., Bornhauser, B., and Brunner, T. (2022).
LRH-1/NR5A2 interacts with the glucocorticoid receptor to regulate glucocorticoid
resistance. EMBO Rep *23*, e54195. 10.15252/embr.202154195.
23. Merk, V.M., Grob, L., Fleischmann, A., and Brunner, T. (2023). Human lung carcinomas
synthesize immunoregulatory glucocorticoids. Genes Immun *24*, 52-56.
10.1038/s41435-023-00194-y.
24. Sidler, D., Renzulli, P., Schnoz, C., Berger, B., Schneider-Jakob, S., Fluck, C., Inderbitzin,
D., Corazza, N., Candinas, D., and Brunner, T. (2011). Colon cancer cells produce
immunoregulatory glucocorticoids. Oncogene *30*, 2411-2419. 10.1038/onc.2010.629.
25. Pramanik, J., Shaji, S.K., Zaman, M., Brown, B., Zhang, B., Yamashita-Kanemaru, Y.,
Homer, N.Z.M., Hussein, H., Zhao, Q., Okkenhaug, K., et al. (2025). Drug repurposing
reveals Posaconazole as a CYP11A1 inhibitor enhancing anti-tumour immunity.
bioRxiv, 2025.2002.2021.639414. 10.1101/2025.02.21.639414.
26. Kurten, C.H.L., Kulkarni, A., Cillo, A.R., Santos, P.M., Roble, A.K., Onkar, S., Reeder, C.,
Lang, S., Chen, X., Duvvuri, U., et al. (2021). Investigating immune and non-immune
cell interactions in head and neck tumors by single-cell RNA sequencing. Nat Commun
12, 7338. 10.1038/s41467-021-27619-4.
27. Janjic, B.M., Kulkarni, A., Ferris, R.L., Vujanovic, L., and Vujanovic, N.L. (2022). Human
B Cells Mediate Innate Anti-Cancer Cytotoxicity Through Concurrent Engagement of
Multiple TNF Superfamily Ligands. Front Immunol *13*, 837842.
10.3389/fimmu.2022.837842.
28. Wang, J., Li, H., Kulkarni, A., Anderson, J.L., Upadhyay, P., Onyekachi, O.V., Arantes, L.,
Banerjee, H., Kane, L.P., Zhang, X., et al. (2025). Differential impact of TIM-3 ligands
on NK cell function. J Immunother Cancer *13*. 10.1136/jitc-2024-010618.
29. Pal, B., Chen, Y., Vaillant, F., Capaldo, B.D., Joyce, R., Song, X., Bryant, V.L., Penington,
J.S., Di Stefano, L., Tubau Ribera, N., et al. (2021). A single-cell RNA expression atlas of
normal, preneoplastic and tumorigenic states in the human breast. EMBO J *40*,
e107333. 10.15252/embj.2020107333.
30. Chen, Y., Pal, B., Lindeman, G.J., Visvader, J.E., and Smyth, G.K. (2022). R code and
downstream analysis objects for the scRNA-seq atlas of normal and tumorigenic
human breast tissue. Sci Data *9*, 96. 10.1038/s41597-022-01236-2.
31. Zilionis, R., Engblom, C., Pfirschke, C., Savova, V., Zemmour, D., Saatcioglu, H.D.,
Krishnan, I., Maroni, G., Meyerovitz, C.V., Kerwin, C.M., et al. (2019). Single-Cell
Transcriptomics of Human and Mouse Lung Cancers Reveals Conserved Myeloid
Populations across Individuals and Species. Immunity *50*, 1317-1334 e1310.
10.1016/j.immuni.2019.03.009.
32. Steele, N.G., Carpenter, E.S., Kemp, S.B., Sirihorachai, V.R., The, S., Delrosario, L.,
Lazarus, J., Amir, E.D., Gunchick, V., Espinoza, C., et al. (2020). Multimodal Mapping of
the Tumor and Peripheral Blood Immune Landscape in Human Pancreatic Cancer. Nat
Cancer *1*, 1097-1112. 10.1038/s43018-020-00121-4.
33. Halbrook, C.J., Thurston, G., Boyer, S., Anaraki, C., Jimenez, J.A., McCarthy, A., Steele,
N.G., Kerk, S.A., Hong, H.S., Lin, L., et al. (2022). Differential integrated stress response
and asparagine production drive symbiosis and therapy resistance of pancreatic
adenocarcinoma cells. Nat Cancer *3*, 1386-1403. 10.1038/s43018-022-00463-1.
718

Figure legends

Figure 1. Genome-wide effect of glucocorticoid signalling in activated CD8 T cell

a. Schematic flowchart of the experimental strategy. CD8 T cells from three healthy donors were purified, activated and propagated. Resting T cells were reactivated the presence or absence of physiological level (100 nM) cortisol to find out glucocorticoid-regulated genes in CD8 T cells by RNA-seq.
Specification of the samples (b-g): KO denotes GR knockout, Mock denotes wildtype, HC denotes hydrocortisone (cortisol) treated. Veh denotes vehicle.

b. Volcano plot depicting the differentially expressed genes in human primary activated CD8 T cells after cortisol treatment. The significantly upregulated genes have been shown in red and the significantly downregulated genes have been shown in blue. log2-Fold change versus –log10(adjusted p-value) has been plotted, n = 3 human samples in each group.

c. The hallmark biological processes associated with cortisol-treated CD8 T cells. Prognosis-ranked gene set enrichment analysis (GSEA) identified hallmark pathways that are differentially regulated. Normalised enrichment scores (NES) were calculated for all 50 hallmark pathways in GSEA. Upregulated (shown in red) and downregulated (shown in blue) gene sets are displayed, with their NES values represented on the x-axis.

d. Knockout (KO) of glucocorticoid receptor encoding gene *NR3C1* in CD8 T cells using CRISPR-Cas9 targeting exon 2 and exon 5 (top panel). The knockout efficiency has been validated by western blot (bottom panel).

e. Volcano plot showing the differentially expressed genes in *NR3C1* KO activated CD8 T cells compared to mock activated CD8 T cells, in the presence of cortisol.

f. The intersections in Venn diagrams showing the cortisol-induced GR-dependent upregulated and downregulated genes.

g. Expression heatmap showing the cortisol-induced transcription factors and receptors (cytokine receptor, transporter type receptor, cell surface and adhesion receptor, nuclear receptor) from differentially expressed genes (log2FC> 1, log2FC <-1 and p.adj > 0.05).

Figure 2. Genome-wide chromatin binding of GR identifies non-canonical transcription factor binding motif in CD8 T cells

a. Diagrammatic flowchart of the experimental strategy. CD8 T cells from healthy donors were purified, activated and propagated. Resting CD8 T cells were

reactivated the presence or absence of physiological level (100 nM) cortisol to find out genome-wide chromatin binding of GR by ChIP-seq.

   b. Homer motif analysis of cortisol-treated CD8 T cells. The top 4 enriched GR binding peaks and corresponding motifs have been shown for two donors (HC-A & HC-B) in cortisol-treated CD8 T Cells. HC-A & HC-B: Hydrocortisone (HC)/cortisol-treated donor-A & donor-B samples.

   c. Venn diagram showing the common number of peaks between hydrocortisone (cortisol) treated CD8 T cells from two donors ,

   d. Venn diagram showing glucocorticoid (GC)-induced peaks comparing vehicle-treated and cortisol (HC)-treated CD8 T cells.

   e. The Venn diagram shows the direct targets of glucocorticoids after intersecting GR binding peaks from ChIP-seq data and cortisol-induced upregulated genes.

   f. The Venn diagram shows all GR-RUNX co-regulated genes that are also direct targets of GR.

   g. Snapshot of glucocorticoid binding peaks around indicated representative genes from the UCSC genome browser. Glucocorticoid influenced and GR -RUNX regulated top five genes have been shown here.

   h. GR-binding site frequencies representing diversity of glucocorticoid receptor binding sites in CD8 T cells.

Figure 3. RUNX transcription factors cooperate with the glucocorticoid receptor to regulate gene expression

   a. Schematic illustration of the experimental strategy (left). HEK 293T cells were transfected with GR and RUNX3 expressing plasmid construct. Nuclear lysates were immunoprecipitated by using anti-GR antibody and immunoblotted by RUNX antibody. Representative immunoblot shown in the right panel.

   b. Venn diagram depicting the set of genes jointly regulated by the GR and RUNX1, identified through an integrative analysis of cortisol-induced upregulated genes, GR ChIP-seq peaks, GR peaks containing RUNX1 motifs, and known RUNX1 target genes. Expression levels of the overlapping gene set are visualised in a heatmap across multiple experimental conditions.

   c. Venn diagram illustrating the genes co-regulated by GR and RUNX2, identified by integrating cortisol-responsive genes (both upregulated and downregulated), GR ChIP-seq data, GR peaks with RUNX2 motifs, and annotated RUNX2 targets. Expression patterns of these shared genes under different conditions are shown in the accompanying heatmap. Specification of the samples: KO denotes glucocorticoid

receptor (GR) knockout, HC denotes hydrocortisone (cortisol) treated. Mock denotes wildtype. Veh denotes vehicle.

Figure 4. Single-cell transcriptomics study reveals the expression of cortisol responder genes in different tumour-infiltrating CD8 T cells

a-d. UMAP-based clustering of scRNA-seq data was used to identify CD8 T cell populations from tumour and matched blood or normal tissue samples in breast cancer (**a**), lung cancer (**b**), head and neck squamous cell carcinoma (HNSC) (**c**), and pancreatic ductal adenocarcinoma (PDAC) (**d**). Dot plots show the proportion of cells (dot size) and the average expression (colour intensity) of selected top cortisol responder genes, *NR3C1*, *TSC22D3*, *AREG*, *FKBP5*, and *PIK3IP1*, within tumour-infiltrating CD8 T cells and corresponding peripheral blood or matched tissue-resident CD8 T cells. The data highlight distinct patterns of steroid-responsive gene expression between tumour-infiltrating and non-tumour CD8 T cell populations, reflecting the influence of the tumour microenvironment on CD8 T cell transcriptional states and potential links to T cell dysfunction and immune evasion

Figure 5. RUNX and GR co-regulated gene expression in tumour-infiltrating CD8 T cells across cancer types

a-d. Single-cell transcriptomic profiling was performed on CD8 T cells isolated from tumours and matched adjacent normal tissue or blood in triple-negative breast cancer (TNBC) (**a**), lung cancer (**b**), head and neck squamous cell carcinoma (HNSC) (**c**), and pancreatic ductal adenocarcinoma (PDAC) (**d**). UMAP-based clustering delineated distinct CD8 T cell populations. Dot plots display both the proportion of cells (dot size) and the average expression (colour intensity) of RUNX (RUNX1, 2 and 3) and GR co-regulated genes within tumour-infiltrating and peripheral or tissue-resident CD8 T cells. This analysis reveals elevated expression of RUNX and GR combined target genes in tumour-infiltrating CD8 T cells compared to their non-tumour counterparts, suggesting a tumour microenvironment-driven transcriptional programme.

e-h. To assess the functional relevance of RUNX and GR co-regulated gene expression, module scores were calculated using the AddModuleScore method for each CD8 T cell. High module scores, indicative of elevated co-regulated gene expression, were associated with increased expression of exhaustion-related genes, highlighting a link between RUNX2/GR transcriptional activity and the exhausted phenotype in tumour-infiltrating CD8 T cells across TNBC **(e)**, lung cancer **(f)**, HNSC **(g)**, and PDAC **(h)**.

Supplementary Figure S1. Comparative *RUNX 1, 2* and *3* expression and identification of GR-RUNX 3 co-regulated genes

a. Comparative *RUNX 1, 2* and *3* expression in activated CD8 T cells in the RNAseq. Normalised expression of these genes from different donors has been plotted here.

b. Venn diagrams showing the GR-RUNX3 co-regulated genes by integrating cortisol induced genes (upregulated and downregulated), all GR binding peaks from ChIP-seq, GR binding site containing RUNX motif and RUNX3 target genes. The corresponding heatmap showing the expression of GR-RUNX3 co-regulated genes in vehicle-treated CD8 T cells, cortisol (hydrocortisone or HC) treated CD8 T cell and *NR3C1* KO CD8 T cells.

Supplementary Figure S2. Single-cell trajectory analysis of CD8 T cells from lung tumours and blood samples

a. Pseudo-temporal trajectories of CD8 T cells inferred using Monocle, coloured by sample origin and pseudotime.

b. Expression patterns of cortisol-responsive genes (*TSC22D3, IL7R, FKBP5*) and inhibitory receptors (*PDCD1, CTLA4, HAVCR2, TIGIT, LAYN, TOX*) along the inferred trajectories.

Figure 1

a

b

Mock HC vs Mock Veh

c

Hallmark pathways

d

e

KO HC vs Mock HC

f

Cortisol induced upregulated genes

Cortisol induced downregulated genes

g

Transcription factors

Receptors

Figure 2

a

PBMC → CD8 T cell → **Stimulation with anti-CD3, anti-CD28** Expansion/propagation → **100 nM HC or vehicle** Re-stimulation with anti-CD3, anti-CD28 → GR ChIP-seq

b

HC-A

	Best match	% peak
ATTCCATTCCAT	TEAD	5.92%
ATCATCGCATGG	SREBF?	4.23%
TTCTGAGAATGC	STAT	0.97%
AAACCACA	Runx	12.57%

HC-B

	Best match	% peak
TTGTGGTTTT	Runx	35.92%
ATTCCATTCG	TEAD	1.58%
CACTTCCTGT	ETS	27.34%
ASTGACGTCA	ATF	17.32%

HC-A and HC-B overlap

	Best match	% peak
ATTCCATTCGAT	TEAD	7.78%
CATTCGATGA	Hbp?	6.1%
GATGATTCCA	Hdx,Cux?	6.28%
AAACCACAGA	Runx	25.2%

c

HC-A / HC-B

524 | 1103 | 5523

d

Veh / HC

710 | 344 | 782

e

GR occupied genes from ChIP-seq — Cortisol-induced upregulated genes

4790 | 81 | 42

Direct targets of GR

f

GR and RUNX-regulated genes

31 | 50

Direct targets of GR

g

TSC22D3
chrX: 107,720,000
RUNX1 motif
RUNX2 motif
Donor A, Donor B, Donor C
Donor A + HC, Donor B + HC, Donor C + HC

IL7R
chr5: 35,855,000 35,860,000
RUNX1 motif
RUNX2 motif

FKBP5
chr6: 35,595,000 35,600,000 35,605,000 35,610,000
RUNX1 motif
RUNX2 motif

PIK3IP1
chr22: 31,280,000 31,290,000
RUNX1 motif
RUNX2 motif

FAM53B
chr10: 124,730,000 124,740,000
RUNX1 motif
RUNX2 motif

h

Genomic distribution of HC binding site on CD8 T cell

- 30.51% promoter-TSS
- 31.48% intron
- 23.78% Intergenic
- 9.11% exon
- 3.46% 5' UTR
- 1.66% TTS

Figure 3

a

b

GR and RUNX1 regulated genes

c

GR and RUNX2 regulated genes

Figure 4

Figure 5

Supplementary Figure S1

a

RUNX expression

b

All peaks from ChIP Seq — RUNX motifs from ChIP Seq — Cortisol induced downregulated genes — RUNX3 target genes

1542, 432, 293, 3, 2, 234, 941, 6460, 19, 5, 0, 0, 25, 1921, 31

All Peaks from ChIP Seq — RUNX motifs from ChIP Seq — Cortisol induced upregulated genes — RUNX3 Target genes

1535, 422, 289, 10, 12, 227, 905, 6478, 18, 41, 4, 7, 18, 1928, 13

GR and RUNX3 regulated genes

COTL1, MFSD11, PASK, SHF, PRDX1, BCL2, TGIF1, PARP8, S100A10, TSPAN14, JAML, CCND3, TSC22D3, IL7R, FKBP5, SYTL3, SCML4, APBB1, HPCAL1, RGCC, SMAP2, CCR7, ST3GAL1, PDE4A, TRERF1, AGO2, BIN1, CASP8, ISG20, HERPUD1, ADAMTS17, ESR2, SYNJ2, FAM53B, SELPLG, TNFAIP3, CXCR4, LDLRAD4, SMAD3, AKAP13, SNX29, PRDM1, CFLAR, SLCO4A1, FBXO32, SLA

Group
- Cortisol upregulated
- Cortisol downregulated

Z-score
2, 1, 0, -1, -2

Group, Mock_Veh, Mock_Veh.1, Mock_Veh.2, Mock_HC, Mock_HC.1, KO_HC, KO_HC.1, KO_HC.2

Supplementary Figure S2

a

b