Gut bacteria gatekeep host-specific colonization of a commensal fungus that boosts type 2 immunity

Geongoo Han[1], Rebecca Yunker[1], Mohammad H. Hasan[1], Alex Bruce[1], Kellie Baldaro[1], Jessica Pacia[1], Owen Lockwood[1], Dana Magana Miralles[1], Nicholas N. Jarjour[2,3], Karthikeyani Chellappa[1], Shipra Vaishnava[1]*, Lalit K. Beura[1]*

[1] Molecular Microbiology & Immunology Department, Brown University, Providence, RI 02912, USA.

[2] Center for Immunology, University of Minnesota, Minneapolis, MN 55455, USA.

[3] Department of Laboratory Medicine and Pathology, University of Minnesota, Minneapolis, MN 55455, USA.

* Corresponding author: lalit_beura@brown.edu, shipra_vaishnava@brown.edu

Abstract

Throughout evolution, organisms have developed exclusive and mutually beneficial relationships with microbes in their natural environments. While these species-specific host-microbe interactions are well-documented for in case of commensal bacteria, similar relationships for commensal fungi remain largely unexplored. In this study, we screened the gut mycobiome of non-laboratory rodents, to find host-specific fungi. We identified *Kazachstania pintolopesii* as a dominant fungus in pet store mice across various geographical locations. This fungus stably colonized laboratory mice, outcompeted non-murine fungi, and persisted independently of gut bacteria without triggering conventional antifungal responses. Colonization by *K. pintolopesii* significantly increased intestinal macrophage populations, which was crucial in preventing systemic immune recognition. *K. pintolopesii* also induced intestinal type 2 immune responses, providing cross-kingdom protection against helminths in free-living mammals exposed to parasitic protozoa. Notably, *K. pintolopesii* strains from non-murine hosts failed to colonize laboratory mice with intact gut bacteria despite high genomic similarity. In gnotobiotic mice, these non-murine strains disrupted bacterial communities and their associated metabolomes, unlike the mouse strains. Our findings establish host exclusivity as a hallmark of fungal commensalism and reveal commensal bacteria as gatekeepers of species-specific fungal colonization and their immune interactions with the mammalian host.

Keywords

Kazachstania pintolopesii, commensal fungi, host adaptation, microbiome, cross-kingdom protection, type 2 immunity

Introduction

Commensal microorganisms are crucial for host immunity and have coevolved with their natural hosts, often exhibiting high species restriction. This phenomenon is observed across various life forms. In invertebrates, the Hawaiian bobtail squid depends on *Vibrio fischeri* for tissue development [1] while tsetse flies rely on *Wigglesworthia glossinidia* for fitness and reproduction [2]. Similarly, in mammals, gut immune cell maturation is driven by host-specific bacteria. Experiments have shown that human or rat gut bacteria introduced into germ-free mice are less effective at stimulating immune cell development compared to mouse-specific bacteria, highlighting the importance of host-adapted microbes in shaping host physiology [3]. A well-studied example of such exclusive host-microbe interaction is segmented filamentous bacteria (SFB), which colonize their hosts with remarkable species specificity, providing significant immune benefits to their adapted host while failing to colonize non-specific hosts [4-6]. While these exclusive interactions have been extensively studied for symbiotic bacteria, similar examples are lacking for other members of the gut microbiome, such as fungi, despite their emerging role in immune modulation as gut colonizers.

The mammalian gut houses a complex ecosystem of microbes, including bacteria, viruses, and fungi. However, our understanding of whether and how commensal fungi interact exclusively with their hosts is limited due to the lack of true commensal models that can stably colonize the mouse gut without causing inflammation or requiring immune suppression. Most research efforts have focused on *Candida albicans,* a gut commensal that is the main cause of opportunistic infections in humans. However, studying *C. albicans* commensalism in humans is somewhat limited because of technical and ethical challenges. As such introducing *C. albicans* to murine gut has emerged as a dominant

model system. But this approach often requires antibiotic treatment to maintain stable

colonization and, can induce inflammatory responses compromising interpretation [7-9].

Therefore, alternative commensal fungi that have coevolved with their natural host, are

needed to better understand fungal-host symbiosis.

Laboratory mice are typically raised in highly controlled, specific pathogen-free (SPF)

environments. These conditions, while beneficial for reducing variability in experimental

outcomes, result in mice with limited exposure to diverse microbial communities, including

fungi. This lack of exposure leads to an immunologically naive phenotype, which is far

removed from the natural immune system of free-living mammals [10-12]. The differences in

microbial exposure between laboratory, wild, and pet store mice present a unique

opportunity to identify and characterize potential commensal fungi that have co-evolved

with their murine hosts [13,14]. By incorporating naturally occurring commensal fungi into

laboratory settings, researchers can create models that better reflect the complex

microbial environment of humans and reveal novel interactions between diverse fungal

commensals and their natural host.

In this study, we sought to isolate and characterize commensal fungi from pet store

mice to elucidate their unique interactions with host. We identified *Kazachstania*

pintolopesii as the dominant fungal species in the gut of pet store mice across different

geographical locations. Through a series of *in vivo* experiments and multi-omics analyses,

we investigated the colonization dynamics, host immune responses, and functional

consequences of *K. pintolopesii* colonization in laboratory mice. By comparing strains

isolated from murine and non-murine hosts, we uncovered the importance of a finely-

tuned host-specific metabolic adaptation for successful gut colonization by *K. pintolopesii*.

This discovery marks the first documented instance of host exclusivity for a fungal

commensal species, opening avenues for exploring how such interactions influence

immune responses, microbial competition, and overall host physiology.

Results

K. pintolopesii is the dominant gut fungus in pet store mice

Specific pathogen-free laboratory mice are maintained in controlled environments, shielded from potential pathogen exposure and typically harbor microorganisms with limited diversity. In contrast, pet store mice live in large breeding colonies and are routinely exposed to rich varieties of microbes, including pathogenic bacteria, viruses, parasites, and fungi [10]. As an example of this, electron microscopic imaging of the gut epithelial surface of laboratory mice revealed no significant epithelium-attached microbes. In contrast, the epithelial surfaces of pet store mice harbored a number of microbial-like structures of varying shapes and sizes (~2-15 μm) (Fig. 1A). Therefore, we used pet store mice as source to screen for commensal murine fungi.

Using ribosomal RNA-based probes, we were able to find presence of fungal species among these epithelial-associated microbes in pet store mice, which are largely absent in laboratory mice (Fig.1B). Fungal burden in the feces of pet store mice was notably high (~10^7 CFU/g of feces), while no fungal populations were detected in the feces of laboratory mice in our facility (Fig. 1C). Fungal colonization was observed throughout the gastrointestinal tract of the pet store mice, from the duodenum to the feces, with the fungal burden progressively increasing toward the caudal end of the intestine indicating that pet store mice are heavily colonized with fungi throughout the gastrointestinal tract (Fig. 1D).

To identify the commensal fungal species in the murine gut, we cultured the gut contents of the pet store mice for 5 days at 30 °C and randomly picked 26 colonies from the plates. Sequencing of the internal transcribed spacer (ITS) region of these colonies revealed that all colonies are the same fungal species, _Kazachstania pintolopesii_ (Fig.

S1A). The isolate appeared as a budding yeast and had an ellipsoidal shape with typical

yeast cell structures such as a thick cell wall, nucleus, vacuole, and mitochondria (Fig.

S1B and C). The isolate showed higher ITS sequence similarities with other *Kazachstania*

species and *Saccharomyces cerevisiae*, whereas lower similarities with *Candida* and

Aspergillus species (Fig. S1D). We also directly sequenced the ITS1 region of the gut

metagenome of pet store mice. Intriguingly, this culture-independent analysis also

showed that the intestines of pet store mice were predominantly colonized by a single

fungal species, *K. pintolopesii*, which accounted for over 99% of the total sequence reads

in both the luminal and mucosal regions of the small intestine and colon (Fig. 1E).

To determine whether this observation is specific to pet stores in a particular region

(Rhode Island), fecal samples were collected from pet store mice in other geographical

locations, including Midwest United States (Minnesota) and Southern United States

(Texas) (Fig. S1E). Higher fungal burden was observed across all regions (Fig. 1F), with

K. pintolopesii consistently identified as the predominant fungal species in the gut,

regardless of the region (Fig. 1G). These findings indicate that *K. pintolopesii* is not limited

to a specific geographical area but rather is a prevalent member of the mouse mycobiome

under non-laboratory setting. It colonizes the gastrointestinal tract extensively, achieving

high population levels both near the mucosal surface and within the intestinal lumen.

K. pintolopesii stably colonizes laboratory mice irrespective of gut bacteria and

provides colonization resistance against non-murine fungi

To test whether this fungal isolate from the pet store mice could colonize the intestines of

the laboratory mice, we orally administered the isolated *K. pintolopesii* to laboratory mice.

A high fungal burden (~10^7 CFU/g of feces) was observed in the feces following

colonization, and this level was maintained for 6 weeks (Fig. 2A). *K. pintolopesii* was

located closely to the gut epithelium of mice, co-localizing with bacteria, similar to the

colonization pattern observed in the pet store mice (Fig. 2B).

We performed 16S rRNA gene sequencing to assess the impact of *K. pintolopesii*

colonization on the bacterial community in the gut. *K. pintolopesii* colonization maintained

bacterial community diversity, composition, and load without significant alterations (Fig.

2C - F).

Germ-free (GF) mice showed *K. pintolopesii* fecal colonization levels (~10^7 CFU/g)

comparable to conventional mice (Fig. 2G), demonstrating that commensal bacteria do

not influence *K. pintolopesii*'s intestinal colonization. The ability of *K. pintolopesii* to

colonize gastrointestinal tract to high levels without impacting bacterial community is in

contrast to other fungal species that often require microbiome manipulation to establish

colonization [7,8,15-17].

We conducted a competition assay to examine whether *K. pintolopesii* affects the

colonization of non-murine-specific fungi. The assay included five fungal species: *K.*

pintolopesii and four previously used to colonize mice - *S. cerevisiae, C. albicans, C.*

parapsilosis, and *C. tropicalis* [16-19] (Fig. 2H). An equal number of viable fungal cells (~2 ×

10^6 CFU/species) were orally administered to both conventional mice and GF mice, and

fecal samples were collected on days 3, 7, 14, and 22, followed by ITS sequencing.

Notably, *K. pintolopesii* exhibited outstanding colonization ability, out-competing the other

fungal species in both conventional and GF mice (Fig. 2I). In conventional mice, *K.*

pintolopesii rapidly dominated the fungal community, accounting for over 95% of the total

fungal population within 3 days of treatment. In GF mice, *C. albicans* was initially the

dominant species, with *C. tropicalis* comprising over 10% of the population until day 7.

However, the proportion of *K. pintolopesii* increased over time, eventually representing

approximately 75% of the total fungal population by days 14 and 22 (Fig. 2I). This

suggests that *K. pintolopesii* is capable of stable colonization in laboratory mice,

independent of the presence of commensal gut bacteria, and provides colonization

resistance against non-murine fungi.

***K. pintolopesii* colonization does not induce key anti-fungal immune responses**

including expansion of neutrophils and the Th17 population

Fungal colonization in the gut has been reported to trigger immune responses including

the expansion of neutrophils and T helper 17 cells (Th17) populations. *C. albicans*, a well-

studied human commensal fungus, is known to induce neutrophils and Th17 populations

in humans and mice [20,21]. Rewilded mice, which harbor an increased fungal population in

the gut, exhibited increased circulating neutrophil levels [13,14]. In addition, mucosa-

associated fungi induced type 17 immune responses [22]. To evaluate whether *K.*

pintolopesii also induces key anti-fungal immune responses, we measured the levels of

neutrophils in blood. Interestingly, *K. pintolopesii* colonization did not induce circulating

neutrophil levels in the peripheral blood (Fig. 3A and S2A). To mimic the systemic fungal

dissemination, we introduced *K. pintolopesii* intravenously in mice, comparing it with *C.*

albicans systemic infection (Fig. 3B). As expected, *C. albicans*-infected mice exhibited a

poor survival rate and a high fungal burden in the kidneys, along with significant

upregulation of *Il17* gene expression in the kidney. Intriguingly, *K. pintolopesii*-infected

mice did not show any pathological symptoms. All *K. pintolopesii*-infected mice survived

and showed low to no detectable levels of fungal burden in the kidney. Additionally, *Il17*

gene expression in the kidney of *K. pintolopesii*-infected mice was significantly lower,

approximately 14-fold lower than that observed in *C. albicans*-infected mice. (Fig. 3C –

E). To assess neutrophils' role in *K. pintolopesii* clearance during systemic infection, we

depleted neutrophils using anti-Ly-6G (1A8 clone) antibody before *K. pintolopesii* infection

(Fig. 3F). Despite complete neutrophil depletion, all mice survived without detectable

kidney fungal burden (Fig. 3G-H and S2B).

We assessed *K. pintolopesii*'s impact on intestinal mucosa by measuring its distance

from the epithelium in both small intestine and colon. The fungi were found closer to the

epithelium on day 5 (small intestine: ~25 μm, colon: ~42 μm, mean distance) but

maintained a greater distance by day 13 (small intestine: ~41 μm, colon: ~55 μm, mean

distance) (Fig. S3A and B). We also measured the levels of Th17, another key anti-fungal

immune cell in mucosal immunity, following *K. pintolopesii* colonization. Interestingly, *K.*

pintolopesii colonization did not increase Th17 cells in the small intestinal lamina propria

(Fig. S3C). Furthermore, the gene expression levels of *S100a8* and *S100a9* in the small

intestine were comparable between control and *K. pintolopesii*-colonized mice (Fig. S3D).

Next, to test whether type-17 immune response can antagonize *K. pintolopesii*, we

provided IL-17A supplementation to colonized mice (Fig. S3E). Despite the IL-17A

injection upregulating the expression of type 17 immunity-related genes (*Il17a*, *Reg3b*,

Reg3g, *S100a8*, and *S100a9*), the fungal burden remained unchanged compared to the

control mice (Fig. S3F and G) indicating that IL-17 and associated downstream effectors

do not limit *K. pintolopesii* burden. To assess contribution of other components of adaptive

immune system in *K. pintolopesii* colonization, we used Rag1 knock-out (KO) mice (Fig.

S4A). Neither the fungal burden nor the body weight differed between wild-type (WT) and

Rag1 KO mice (Fig. S4B and C). To assess the potential for systemic fungal

dissemination and immune activation in the absence of B and T lymphocytes, we

measured the number of neutrophils in the spleen. No significant differences were

observed in neutrophil density between the two genotypes indicating that the adaptive

immune arms are neither engaged nor play a significant role in limiting *K. pintolopesii*

burden (Fig. S4D).

It has been reported that intestinal colonization by *C. albicans* can protect the host

against systemic *C. albicans* infection [21]. To assess whether this commensal fungus

provides a similar cross-species protection to the host, *C. albicans* was systemically

introduced into *K. pintolopesii*-colonized mice (Fig. S5A). However, no differences in

survival rates or fungal burden in the kidneys were observed between the control and *K.*

pintolopesii-colonized mice following systemic *C. albicans* infection (Fig. S5B and C). In

contrast, we observed that the intestinal colonization by *C. albicans* in antibiotics-treated

mice protected the host from systemic candidiasis (Fig. S5D – F).

These results indicate that *K. pintolopesii* does not trigger classical anti-fungal defenses

including neutrophilia and Th17 response and the complete absence of the adaptive arm

of the immune system does not enhance fungal pathogenicity. When *K. pintolopesii* was

introduced systemically, the host was able to control the fungus via neutrophil-

independent mechanisms suggesting a unique host-fungal relationship that differs from

typical anti-fungal immune responses.

***K. pintolopesii* increases the macrophage population in the small intestine**

Although we examined the effects of *K. pintolopesii* colonization on the host immune

system, we primarily focused on dominant anti-fungal immune responses, specifically

neutrophils and Th17 cells. However, to thoroughly investigate the impact of *K.*

pintolopesii colonization on the host's intestine in an unbiased manner, we collected ileum

samples 14 days post-colonization and performed bulk RNA sequencing. *K. pintolopesii*

colonization showed significant transcriptional changes with 602 up-regulated genes and

1,213 down-regulated genes compared with control mice (Fig. 4A and S6A). Notably,

genes associated with macrophage activation were remarkably upregulated following *K.*

pintolopesii colonization (Fig. 4B). GO terms related to macrophage activation, including

'regulation of macrophage activation', 'macrophage activation', and 'macrophage

cytokine production', were significantly enriched in the *K. pintolopesii*-colonized group

(Fig. 4C). We also observed that *K. pintolopesii*-colonized mice exhibited elevated

macrophage frequencies in the lamina propria of small intestine compared with control

mice (Fig. 4D and S6B).

To test the role of macrophages on *K. pintolopesii* colonization, we used the clodronate-

liposomes mediated macrophage depletion technique. Clodronate liposomes can also

target monocyte and monocyte-derived macrophages along with resident macrophages

[23,24]. Interestingly, in clodronate-treated mice, *K. pintolopesii* colonization significantly

increased the level of neutrophils in systemic circulation compared to non-colonized mice

(Fig. 4E-F and S6C). This indicates that in the absence of macrophages and monocytes,

K. pintolopesii can disseminate systemically and trigger neutrophilia. These findings

suggest that macrophages are a barrier to the systemic spread of *K. pintolopesii* and help

maintain this fungus as a commensal member within the gut.

K. pintolopesii colonization induces type 2 immune responses in the gut and

protects the host against roundworm infection

Our further analysis of the RNA-seq data revealed upregulation of tuft cell and goblet cell

marker genes in *K. pintolopesii*-colonized mice (Fig. 5A). To validate the RNA sequencing

data, we quantified the number of tuft cells and goblet cells in the small intestine. As we

saw in RNA sequencing data, the number of tuft cells and goblet cells was significantly

higher in *K. pintolopesii*-colonized mice than in control mice (Fig. 5B and C). Tuft cells are

a rare cell type in the intestinal epithelium, yet they play a crucial role in goblet cell

hyperplasia and initiating type 2 immune responses [25,26]. So, next examined *K.*

pintolopesii also modulates other type 2 associated cells. We observed significant

upregulation of T helper 2 (Th2) and group 2 innate lymphoid cells (ILC2s) by *K.*

pintolopesii colonization (Fig. 5D and S7A).

To check if this induction of type-2 immunity is not only limited to the C57BL/6 mouse

strain, we colonized *K. pintolopesii* in the prototypical Th2-type mice, BALB/c [27,28].

Colonized BALB/c mice exhibited a high fungal burden and significantly increased

frequencies of Th2 and ILC2 levels in the small intestine, similar to what was observed in

C57BL/6 mice (Fig. 5E-F and S7B). However, circulating neutrophil levels and the Th17

population in the small intestine were comparable to those in control mice (Fig. S7C and

D).

Next, to assess the functional relevance of this type-2 immunity, we challenged control

and *K. pintolopesii*-colonized mice with a murine intestinal nematode, *Heligmosomoides*

polygyrus, that is controlled by type-2 immune responses (Fig. 5G). The number of eggs

in the fecal samples was significantly lower in *K. pintolopesii*-colonized mice compared

with control mice at 21 days post-infection (Fig. 5H). This suggests that *K. pintolopesii*

colonization in the mouse intestine protects the host from a parasitic nematode.

273 *K. pintolopesii* **exhibits strict host specificity, with colonization of non-murine**

274 **strains causing microbiome and metabolic disruption in mice**

275 We described a mouse-originated strain of *K. pintolopesii* (Kp-M) that can stably colonize

276 the mouse intestine and remains non-virulent even when systemically exposed. Two

277 recent studies have identified other *Kazachstania* strains, from mice intestines [29,30].

278 Similarly, *Kazachstania* species have been found in pigs, primates, and humans [31-34]. The

279 presence in different hosts suggest that *Kazachstania* species are widely distributed

280 across mammals and might contribute similar symbiotic benefit to their host. However,

281 another feature of true commensal is the adaptation of the symbiont to the specific

282 environmental and physiological conditions of the natural host, borne out of long-period

283 of co-existence. There have been many examples of such host-restriction across the

284 microbiome literature [35-37]. To explore whether *K. pintolopesii* also displays such

285 colonization restriction, we isolated a different strain of *K. pintolopesii* from the gut of

286 another pet rodent, the guinea pig. The guinea pig-originated strain (Kp-GP) was isolated

287 from fecal samples based on the shape and size of colonies on the plates and was

288 identified as *K. pintolopesii* by their ITS sequences and clustered together with the Kp-M

289 strain (Fig. S8A). The genomes of Kp-M and Kp-GP were sequenced and assembled with

290 good completeness. The estimated genome sizes were 13.5 Mbp for Kp-M and 13.9 Mbp

291 for Kp-GP, with only minor differences observed in the aligned sequences between the

292 two strains (Fig. S8B - D). Average nucleotide identity (ANI) analysis revealed a 98%

293 identity between Kp-M and Kp-GP genome sequences. We obtained genome sequences

294 of additional *K. pintolopesii* strains isolated from cynomolgus macaque (KPIN-4417) and

295 mice (C89-2, ID1LA, IDNYC, QC1, W26G), as well as Kaz3, another murine commensal,

to conduct a comprehensive genomic analysis. The genomes of eight *K. pintolopesii*

strains and *K. weizmannii* showed a high level of similarity and shared a substantial

number of the function-known genes (Fig. S8E). We inferred the orthologs of the two *K.*

pintolopesii isolates with other fungal species and compared them using the phylogenetic

method, which showed that Kp-M, Kp-GP, and *K. pintolopesii* strains share more genomic

features with each other than with other species, such as *C. albicans* and *S. cerevisiae*

(Fig. S8F).

We first tested if Kp-GP can systemically spread through intravenous introduction.

Similar to Kp-M, Kp-GP showed no detectable fungal burden in the kidney and we

observed no differences between in body weight, circulating neutrophil levels, or kidney

Il17 gene expression (Fig. S9A – E). Next, we assessed if the the Kp-GP strain can

establish itself in gut. But unlike the mouse-derived strain, Kp-GP showed very poor

colonization in the intestine of the mouse host (Fig. 6A and B). In addition, continuous Kp-

GP exposure in mice did not improve the colonization (Fig. S10A and B).

However, the colonization level of Kp-GP in the GF mice was comparable to that of Kp-

M (Fig. 6C), indicating that commensal bacteria provide colonization resistance to Kp-GP.

To investigate how commensal bacteria influence fungal colonization in a non-native host,

we used the Altered Schaedler Flora (ASF) model system with the defined bacterial

species. A further advantage of the ASF-colonized mice is, it can support stable

colonization of non-murine commensal fungi like *Malassezia restricta* and *C. albicans*,

providing an ideal system to compare colonization patterns of fungi from different host

origins [38,39]. We gavaged Kp-M, Kp-GP, and *C. albicans* into ASF-colonized ex-GF mice

and confirmed stable colonization in the intestine, although the fungal burden of Kp-GP

and *C. albicans*-colonized mice was significantly lower compared with Kp-M-colonized

mice (Fig. 6D and E). 16S rRNA gene sequencing revealed that colonization of Kp-GP

and *C. albicans* altered the bacterial community, whereas Kp-M did not change the ASF

community in the gut (Fig. 6F and G).

We also analyzed cecal metabolites, and similar patterns were observed (Fig. 6H).

Specifically, mice colonized with non-murine fungi, Kp-GP or *C. albicans*, exhibited similar

metabolic alterations particularly among amino acids and organic compounds (Fig. 6I).

To infer the effects of bacterial alteration on the cecal metabolites, we performed a co-

occurrence analysis and found several significant concurrent patterns between cecal

metabolites and intestinal bacteria. Especially two *Lactobacillus* species (ASF360 and

361) showed positive associations with metabolites related to 'Alanine, aspartate and

glutamate metabolism', 'Glycine, serine and threonine metabolism', 'Amino sugar and

nucleotide sugar metabolism', and 'Pantothenate and CoA biosynthesis' pathways in Kp-

GP or *C. albicans*-colonized mice. In contrast, *Mucispirillum schaedleri* (ASF457) and

Parabacteroides sp. (ASF519) showed negative associations with these metabolites.

Conversely, level of these bacteria were negatively associated with metabolites like

acetoin, ethylmalonate, xanthurenate, diaminopimelate, hydroxypyruvate, anserine, and

diacetyl (Fig. 6I). These results suggest that *K. pintolopesii* colonization in non-native host

alters the host's intestinal bacterial communities, potentially affecting metabolic pathways

and overall gut homeostasis suggesting the host's commensal bacteria act as a barrier,

resisting colonization by non-native *K. pintolopesii* strains.

Discussion

Naturalized, wilded, and pet store cohoused mice have emerged as valuable models for studying the impact of microbes encountered outside the laboratory setting. These mice provide researchers with a more accurate representation of the complex microbial interactions that occur in natural environments, offering insights that conventional laboratory mice cannot. Transfer of gut microbiome from wild mice to the GF mice exhibited improved protection against viral infection and embryo transfer of laboratory mice into wild mice showed human-like immune phenotypes [40,41]. Exposing laboratory mice to wild environments, and then recapturing them, is another way to colonize the wild microbiome in laboratory mice. These rewilded mice exhibit increased circulating neutrophils and enhanced granulopoiesis [13,14]. Pet store mice are another valuable resource for studying host-microbe interactions in mice. Although they do not live in the natural environment like wild mice, they are exposed to the dirty environment in which humans live. When laboratory mice are exposed to pet store mice microbiome by co-housing, co-housed laboratory mice harbor various microbes such as viruses, fungi, and worms, and exhibit similar immune composition to those of human adults [10,42].

While studies have explored interactions between mice and their complex microbiomes in toto, the specific relationships between hosts and individual microbial kingdoms are less explored, especially in the presence of diverse pathogenic groups. Among various microbes, we focused on the role of commensal fungi in the murine gut. Fungi ubiquitously exist in environments including several body sites of mammals [43]. In general, the importance of commensal fungi is underestimated compared to commensal bacteria due to fewer available databases and lower cell numbers compared to bacteria in the gut.

However, fungi can contribute unique metabolic functions as eukaryotes and even minor

commensal member affects host phenotypes in the gut [6,44,45]. In our study, we aimed to

identify murine commensal fungi that will allow us to take advantage of the robust mouse

model to study fungal commensalism. We successfully isolated *K. pintolopesii* from the

intestines of pet store mice, which constituted the dominant fungal species in the pet store

mice regardless of geographic location. In laboratory settings, *K. pintolopesii* was capable

of stable colonization without any manipulation of the gut microbiome and was not harmful

to the host.

Two recent studies also identified *Kazachstania* species from murine hosts from

immunodeficient and wild mice that can colonize laboratory mice gut to high numbers.

While the study by Kralova et al. investigates *Kazachstania heterogenica var. weizmannii*

isolated from an immunodeficient mouse model [29], the study by Liao et al. examines *K.*

pintolopesii isolated from free-living mice [30]. The *Kazachastania* sp. in all three studies

demonstrate a remarkable ability to colonize the murine gastrointestinal tract and

outcompete or prevent colonization by non-murine fungi including *C. albicans*,

showcasing their potential as powerful modulators of the gut mycobiome.

K. weizmannii exhibits a stealth colonization strategy, inducing minimal immune

responses under normal conditions but mitigating *C. albicans*-induced pathology [29]. In

contrast, *K. pintolopesii* isolated in Liao et al. and our study triggers a robust type 2

immune response enhancing resistance to helminth infections while exacerbating

gastrointestinal allergy [30]. We additionally reveal the critical role of macrophages in

modulating the immune response during colonization by *K. pintolopesii*, particularly in

preventing excessive neutrophil activation. We show that upon colonization, *K.*

pintolopesii does not trigger the typical antifungal immune responses associated with

other gut-colonizing fungi, such as *C. albicans*, which often leads to increased neutrophil

levels. Instead, the presence of *K. pintolopesii* is associated with an increase in

macrophage populations within the small intestine. When macrophages/monocytes are

depleted using clodronate-liposomes, there is a significant rise in circulating neutrophils

following *K. pintolopesii* colonization. Macrophages play a crucial role in maintaining

immune homeostasis with commensal bacteria in the gut. These tissue-resident cells are

strategically positioned in the intestinal lamina propria, where they act as the first line of

defense against potential pathogens while simultaneously promoting tolerance to

beneficial microbes. Unlike neutrophils, which can cause bystander damage through their

potent inflammatory responses, macrophages employ a more measured approach to

microbial interactions [46-48].

Our study also provides compelling evidence for species-specific host adaptation of *K.*

pintolopesii. We observed that two *K. pintolopesii* strains isolated from different hosts,

mice and guinea pigs, act differently in mice despite their high genomic similarity. Only

the mouse-derived *K. pintolopesii* strain showed stable colonization in conventional mice

with colonization resistance against non-murine fungal strains. However, the Kp-GP strain

failed to colonize the mouse gut, and the colonization level of the Kp-GP was significantly

lower than that of the Kp-M, even in the presence of a simple bacterial community, the

ASF. Moreover, the Kp-M strain did not change the bacterial composition and metabolites

in the gut, whereas the Kp-GP strain altered the commensal bacteria and cecal metabolic

profiles in the ASF mouse model. Our results demonstrate that fungal species require

host adaptation to establish a commensal relationship with the host in the gut. Host

adaptation is crucial in becoming a colonizer of the host's microbial community [49]. For

example, pathogenic bacteria evolve genetically to colonize new host species, allowing

them better adhesion ability, host-specific nutrient acquisition, and immune evasion [49].

Tso et al. reported that the mouse gut-adapted *C. albicans* strain by experimental

evolution exhibits enhanced competitive fitness in the gut [50]. Although genome

sequences between Kp-M and Kp-GP are highly similar, small genomic differences during

host adaptation may result in different colonization abilities because even a single amino

acid difference can provide host specificity [51]. The presence of commensal bacteria plays

a crucial role in supporting fungal commensalism because we observed that the Kp-GP

strain colonized the gut of GF mice at levels similar to those of the Kp-M strain. One

possible explanation is that the host-adapted fungi do not directly compete with

commensal bacteria for nutrients within the gut environment because the Kp-M strain did

not alter the bacterial community and the cecal metabolic profiles. Another possibility is

that the host-adapted fungi can evade the immune system, which was trained by

commensal bacteria. Kralova et al.'s study identifies *K. heterogenica* or *K. weizmannii* in

human gut and vaginal microbiome samples, suggesting potential translational

implications [29]. Our findings underscore the importance of considering strain-level

differences and unique host-specific adaptation when studying fungal commensals.

The Liao et al. study shows that *K. pintolopesii* triggers a type 2 immune response,

characterized by eosinophilia and Th2 cell polarization but only when the mucosal barrier

is impaired (e.g., during fiber-free diet, antibiotic treatment, or in germ-free mice) [30]. In

contrast, we observed that *K. pintolopesii* strain from pet store mice increases type 2

immune cells in the small intestine and provides cross-kingdom protection against

helminth infection even without any need of barrier impairment suggesting strain-level

differences in *K. pintolopesii* mice strains between the two studies. Perhaps in the

microbial environment of pet store mice that have higher burden of parasitic worms and

pathogens has forced *K. pintolopesii* to induce type 2 immunity [52-54]. Cross-kingdom

interactions highlight the complexity and interconnectedness of microbial communities

and their host organisms. These interactions can have significant implications for

understanding disease resistance, developing new therapeutic strategies, and managing

ecosystems.

In conclusion, our study underscores the critical importance of strain-specific, host-

exclusive commensal fungal interactions in shaping the gut mycobiome and influencing

host physiology (Fig. 7). The discovery of *K. pintolopesii*'s species-level exclusivity in mice

points to the intricate evolutionary adaptations enabling specific fungi to establish stable,

host-specific relationships. Focusing on these naturally occurring, host-adapted fungi can

provide deeper insights into co-evolutionary processes, leading to a more nuanced

understanding of how commensal fungi contribute to host health, influence immune

system development, and potentially protect against pathogens or modulate disease

outcomes. This approach could reveal new therapeutic targets or probiotic strategies for

improving gut health and managing various gastrointestinal disorders.

Methods

Mice

Conventional wild-type C57BL/6 mice were purchased from Taconic Biosciences and

Jackson laboratory or were bred in the animal facility at Brown University. BALB/c mice

were purchased from Jackson laboratory. Rag1 KO mice were used as an adaptive

immunity-deficient mouse model. Germ-free (GF) mice were wild-type C57BL/6

background, and they were raised and bred in flexible film isolators in a gnotobiotic facility

at Brown University. Pet store mice were purchased from local pet stores. Fecal pellets

of pet store mice in Minnesota and Texas and pet guinea pigs were freshly delivered to

Brown University and used for further experiments. Altered schaedler flora (ASF) mice

were established by colonizing GF mice with ASF. Briefly, ASF fecal pellets (Taconic

Biosciences) were resuspended in pre-reduced sterile PBS (50 mg/1 mL) followed by

filtration using the 70 μm cell strainer. 100 uL filtrates were gavaged into GF mice.

Experiments were performed according to the protocols approved by the Institutional

Animal Care and Use Committees of Brown University.

Fungal isolation from the gut of pet store mice and guinea pigs

Cecal contents and feces, collected from pet store mice, were suspended in sterile PBS

and plated on Sabouraud Dextrose (SD) agar with streptomycin (40 mg/L) followed by

incubation at 30 °C for 5 days. A total of 26 single colonies were randomly picked from

the plates and the ITS region of fungal colonies was amplified by colony PCR using the

Phire Green Hot Start II PCR Master Mix (Thermo Scientific) with ITS1-F/ITS4 primer pair

(ITS1-F: 5'-CTTGGTCATTTAGAGGAAGTAA-3', ITS4: 5'-

TCCTCCGCTTATTGATATGC-3') [55,56]. The amplification program consisted of 94 °C for

10 min, followed by 35 cycles of 94 °C for 30 sec, 52 °C for 30 sec, and 72 °C for 1 min,

and 72 °C for 8 min. The amplicon was sequenced by Sanger sequencing (GENEWIZ)

and MOLE-BLAST (Basic Local Alignment Search Tool) was performed against the ITS

database of NCBI to identify the isolated colonies. The exported phylogenetic tree from

MOLE-BLAST was visualized using the R package *ggtree*.

To isolate *K. pintolopesii* from pet guinea pigs, feces were suspended in sterile PBS

and plated on SD agar with streptomycin (40 mg/L) followed by incubation at 37 °C for 2

days. Nine single colonies with morphological characteristics resembling those of mouse

K. pintolopesii colonies were selected from the plates. Identifying these isolates followed

the same procedures as we did for pet store mice samples.

Fungal culture

All fungal strains were cultured in SD or yeast extract-peptone-dextrose (YPD) media.

The isolate, *K. pintolopesii*, was cultured at 37 °C for 1 day with shaking. This culture was

inoculated in fresh media and cultured for 1 day in the same condition. *S. cerevisiae*

BY4717 was cultured at 30 °C for 2 days with shaking. *C. albicans* SC5314, *C.*

parapsilosis ATCC22019, and *C. tropicalis* MYA-3404 were cultured overnight at 30 °C

with shaking. For systemic infection, *C. albicans* overnight culture was inoculated in fresh

media and cultured for 4 h in the same condition.

Fungal infection

K. pintolopesii were treated in drinking water ($\sim 10^6$ CFU/mL) for intestinal colonization.

After three days, the fungal water changed to fresh regular drinking water.

For oral administration of various fungal species, each fungal culture was mixed with

the same CFU and gavaged to conventional or GF mice (~2 × 10^6 CFU/species in 200

μL per mouse).

For intestinal colonization of *C. albicans*, antibiotics water was prepared with penicillin

(750 U/mL), streptomycin (1 mg/mL), and 2.5% glucose. The antibiotic water was treated

four days before oral inoculation. *C. albicans* culture was resuspended in antibiotic water

(~10^6 CFU/mL) and was treated. After a day, fungal antibiotic water was changed to fresh

antibiotic water, and fresh antibiotic water was provided every three or four days during

the whole experimental period. The same antibiotics water was provided for mice in the

control group and the *K. pintolopesii* group.

For ASF mice colonization, each fungal strain was gavaged into mice after 2 weeks of

ASF treatment (2 × 10^6 CFU/mouse).

For systemic infections, *K. pintolopesii* (~10^5 CFU/200 μL) or *C. albicans* (~10^5

CFU/200 μL) was injected into mice via the retro-orbital venous sinus or lateral tail vein.

All fungal cultures were washed twice with sterile PBS and resuspended in the proper

solution for infection.

Fungal burden measurement

The fecal pellets and the contents of the intestine were resuspended in sterile PBS with

antibiotics (500 mg/L ampicillin, 250 mg/L streptomycin, 225 mg/L kanamycin). The other

tissues were homogenized in sterile PBS with antibiotics. All samples were serially diluted

in sterile PBS and plated on SD agar with streptomycin (40 mg/L) followed by incubation

at 30 °C or 37 °C for 1 or 2 days based on the optimal condition of each fungus.

Immune cell isolation and flow cytometry

To isolate peripheral blood lymphocytes, blood samples were collected from the

submandibular vein by cheek punch in heparin-containing polypropylene tubes. Red

blood cells were removed from blood samples by two rounds of ACK (Ammonium-

Chloride-Potassium) lysis buffer treatment. Lymphocytes from the spleen and the small

intestinal lamina propria were isolated as described in Beura et al. [57] and Grizotte-Lake

et al. [58].

Isolated cells were stained with the corresponding cocktail of fluorescently labeled

antibodies (CD3, CD4, CD8a, CD8b, CD11b, CD11c, CD44, CD45, CD62L, CD64,

CD90.2, B220, Ly6C, Ly6G, KLRG1, F4/80, ST2) and biotin-conjugated anti-mouse

lineage CD3e, CD5, Ly6G, CD19, CD11b, CD11c and streptavidin. Cell viability was

determined using Ghost Dye 780 or Fixable Viability Dye eFluor™ 450. After surface

staining, cells were fixed and permeabilized followed by intracellular staining with the

corresponding cocktail of fluorescently labeled antibodies (T-bet, Foxp3, GATA3, Rorγt,

IL-17). The stained samples were acquired using the Aurora flow cytometer (Cytek) and

were analyzed with FlowJo 10 software (BD). To visualize high-dimensional flow

cytometry data, 80,000 live $CD45^+$ cells per group were sampled using the DownSample

plugin and analyzed using the t-distributed stochastic neighbor embedding (t-SNE)

method in Flowjo 10 software.

IL-17 injection

Recombinant mouse IL-17A (BioLegend) was administered intraperitoneally to mice

every other day (0.5 µg/100 µL per mouse), beginning on the day of *K. pintolopesii*

treatment, to elevate the levels of IL-17. Control mice received intraperitoneal injections

of PBS (100 µL per mouse).

Neutrophil and macrophage/monocyte depletion

For neutrophil depletion, anti-Ly-6G antibody (1A8 clone; Biolegend) or isotype control (Biolegend) was administered intraperitoneally to mice every other day (200 μg in 200 μL per mouse) from the day before the *K. pintolopesii* challenge.

To deplete the macrophage/monocyte populations, Clodrosome (Encapsula Nano Science) was administered intraperitoneally to mice every other day (200 μL per mouse) from the day before *K. pintolopesii* treatment. Encapsome (Encapsula Nano Science), a control liposome formulation, was administered to control *K. pintolopesii* treatment mice (200 μL per mouse).

Genome sequencing of fungal isolate

DNA extraction and Sequencing. Fungal DNA was extracted from fungal culture as described in Jiang et al [16]. Briefly, fungal cells (~10^8 CFU) were collected by centrifugation at 4,000 rpm for 5 min. Cell pellets were suspended in 50 mM Tris buffer (pH 7.5) supplemented with 1 mM EDTA, 0.2% b-mercaptoethanol, and 1000 units/ml of lyticase (Sigma-Aldrich) followed by incubation at 37 °C for 30 min. After incubation, genomic DNA was extracted using a Quick-DNA™ Fungal/Bacterial Miniprep Kit (Zymo Research).

The extracted DNA sample was submitted to the MiGS or the SeqCenter for library construction and sequencing. For Illumina sequencing, sample libraries were prepared using the Illumina DNA prep kit and IDT UDI indices and then sequenced on Illumina NextSeq 2000 or NovaSeq X Plus 2 × 151 bp paired-end sequencing. Demultiplexing, quality control, and adaptor trimming were performed with bcl2fastq. For Oxford Nanopore Technology (ONT) sequencing, samples were prepared using the Genomic

DNA by Ligation kit (Oxford Nanopore) and then sequenced on Nanopore R9 flow cells.

Guppy was used for high-accuracy base calling.

Genome assembly and annotations. Adapter and low-quality sequences were trimmed

from Illumina reads using Trimmomatic [59]. ONT reads were trimmed and filtered using

Porechop [60] and Nanofilt [61]. The quality of reads before and after trimming was checked

using FastQC (https://www.bioinformatics.babraham.ac.uk/projects/fastqc/). The *k*-mer

was counted using jellyfish [62] with Illumina reads followed by visualization on

GenomeScope [63] to estimate genome size.

De novo assembly was performed using Flye [64] based on ONT data and the assembled

genome was polished with Illumina reads using medaka

(https://github.com/nanoporetech/medaka), Racon [65], and Pilon [66]. Assembly was

assessed using QUAST [67] and BUSCO [68]. Funannotate pipeline

(https://github.com/nextgenusfs/funannotate) was used for genome annotation.

Comparative genome analysis. Assembled *K. pintolopesii* genomes from mice and

guinea pigs were aligned using the Mauve alignment tool [69]. Anvi'o [70] was used to align

and visualize multiple genomes. Orthofinder [71] was used to infer orthologs between yeast

strains and compare their genes. All genome and protein sequences (*K. pintolopesii*

KPIN-4417, *K. pintolopesii* C89-2, *K. pintolopesii* ID1LA, *K. pintolopesii* IDNYC, *K.*

pintolopesii QC1, *K. pintolopesii* W26G, *K. weizmannii* Kaz3, *C. albicans* SC5314

ASM18296v3, and *S. cerevisiae* S288C R64) were downloaded from the NCBI database.

DNA extraction for microbiome and mycobiome analysis

The small intestine (ileum) and colon were flushed with sterile PBS to collect luminal

contents and intestinal material was obtained by centrifugation at 5,000 rpm for 10 min.

After the flush, the small intestine and colon were scraped and collected in sterile PBS

followed by centrifugation at 5,000 rpm for 10 min to get mucosal contents. Fecal samples

were directly collected from the rectum. Samples (~50 mg) were suspended in 50 mM

Tris buffer (pH 7.5) supplemented with 1 mM EDTA, 0.2% b-mercaptoethanol, and 1000

units/mL of lyticase (Sigma-Aldrich) followed by incubation at 37 °C for 30 min. After

incubation, genomic DNA was extracted from samples using a Quick-DNA Fecal/Soil

Microbe Microprep Kit (Zymo Research) according to the manufacturer's protocol and

was stored at −20°C until further processing.

Bacterial load measurement

Bacterial load in the feces was measured by quantitative real-time PCR (qPCR).

Reactions were prepared using Maxima SYBR Green/ROX qPCR Master Mix (Thermo

Scientific). Bacterial DNA contents in the intestine were determined by 16S rRNA gene

contents using the 340F/514R primer pair (340F: 5'-ACTCCTACGGGAGGCAGCAGT-3',

514R: 5'-ATTACCGCGGCTGCTGGC-3'). Bacterial load was expressed as a 2^{-Ct} value

and normalized to the weight of the starting fecal material.

16S rRNA gene sequencing and bacterial community analysis

The V4/V5 region of the bacterial 16S rRNA gene was amplified using the Phusion High-

Fidelity DNA polymerase (Thermo Scientific) with 518F/926R primer pair (518F: 5'-

CCAGCAGCYGCGGTAAN -3', 926R: 5'- CCGTCAATTCNTTTRAGT -3'), which were

conjugated with overhang adapter sequences for Illumina MiSeq. The amplification

program consisted of 95 °C for 3 min, followed by 35 cycles of 95 °C for 30 sec, 58 °C for

30 sec, and 72 °C for 30 sec, and 72 °C for 10 min. DNA libraries were constructed and

sequenced on Illumina MiSeq 2 × 300 bp paired-end sequencing (Rhode Island

Genomics and Sequencing Center).

The raw sequencing reads were processed using QIIME2 pipeline [72]. DADA2 [73] was

used to perform denoising and to construct the amplicon sequence variants (ASVs) table.

The pre-trained Naïve Bayes classifier on the SILVA 132 database [74] was used for

taxonomic assignment. Singletons and all features annotated as mitochondria or

chloroplast were removed from the table and the abundance of bacterial taxa was

expressed as a percentage of total 16S rRNA gene sequences. For alpha and beta

diversity, the feature table was rarefied to 10,434 reads. Shannon was used as an alpha

diversity index and principal coordinates analysis (PCoA) based on weighted UniFrac

distances was used for beta diversity. R package *qiime2R*

(https://github.com/jbisanz/qiime2R) was used for visualization.

ITS sequencing and fungal community analysis

The ITS1 region of the fungal DNA was amplified using the Phusion High-Fidelity DNA

polymerase (Thermo Scientific) with ITS1-F/ITS2 primer pair (ITS1-F: 5'-

CTTGGTCATTTAGAGGAAGTAA-3', ITS2: 5'-GCTGCGTTCTTCATCGATGC -3')[55,56],

which were conjugated with overhang adapter sequences for Illumina MiSeq. DNA library

construction and sequencing were conducted following the same procedures for 16S

rRNA gene sequencing.

The raw sequencing reads were processed using QIIME2 pipeline [72]. The reads were

trimmed using Q2-ITSxpress [75]. Trimmed sequence reads were denoised and sequence

variants were identified by DADA2 [73]. Taxonomy was assigned using a pre-trained Naïve

Bayes classifier on the UNITE v8.2 database [76]. Singletons were removed from the table

and the abundance of fungal taxa was expressed as a percentage of total sequence reads.

RNA extraction from tissues

Small pieces of the small intestine and the kidney were stored in RNAlater (Invitrogen).

Total RNA from tissues was extracted using PureLink™ RNA Mini Kit (Invitrogen) or

RNeasy Plus Mini Kit (Qiagen) according to the manufacturer's protocol and was stored

at −20°C until further processing.

Quantitative PCR (qPCR)

cDNA was synthesized from isolated RNA with M-MLV Reverse Transcriptase

(Invitrogen). qPCR reactions were prepared using Maxima SYBR Green/ROX qPCR

Master   Mix   (Thermo   Scientific)   with   primer   sets   for   *Il17*   (F:

ATCAGGACGCGCAAACATGA,   R: TTGGACACGCTGAGCTTTGA),   *Reg3b*   (F:

TACTGCCTTAGACCGTGCTTTCTG, R: GACATAGGGCAACTTCACCTCACA), *Reg3g*

(F: TTCCTGTCCTCCATGATCAAAA, R: CATCCACCTCTGTTGGGTTCA), *S100a8* (F:

TGTCCTCAGTTTGTGCAGAATATAAA,   R: TCACCATCGCAAGGAACTCC),   and

S100a9 (F: GGTGGAAGCACAGTTGGCA, R: GTGTCCAGGTCCTCCATGATG) genes.

Gene expression was normalized to *Gapdh* (F: TGGCAAAGTGGAGATTGTTGCC, R:

AAGATGGTGATGGGCTTCCCG) and relative expression was calculated using the $^{\Delta\Delta}$Ct

method.

RNA-Seq

Extracted RNA samples were submitted for library construction and sequencing

(Novogene). The constructed libraries were sequenced on the Illumina NovaSeq X Plus

platform (Paired-end 150 bp). Adapter and low-quality reads were trimmed from the raw

sequence reads using Trimmomatic [59]. Trimmed sequence reads were aligned to the

mm10 mouse genome using STAR aligner [77]. DESeq2 [78] was used to identify differentially

expressed genes (DEGs) between groups when $P < 0.05$. R packages Complex heatmap

[79] and EnhancedVolcano (https://github.com/kevinblighe/EnhancedVolcano) were used

to draw the heatmap and volcano plot. The cell-specific gene list was obtained from

PanglaoDB [80] and utilized as a cell marker gene set for gene set enrichment analysis

(GSEA). R package clusterProfiler [81] was used to perform GSEA and gene ontology (GO)

enrichment analysis.

Fluorescence *in situ* hybridization (FISH)

The small intestine and the colon were fixed in Methacarn (Bioworld) and embedded in

paraffin. Tissues were sectioned at 7 μm and deparaffinized in xylene followed by

rehydration in 95% EtOH, 90% EtOH, and H_2O. After deparaffinization and rehydration,

tissue sections were hybridized into universal bacterial probes (Alexa 488-

GCTGCCTCCCGTAGGAGT; Alexa 488-GCAGCCACCCGTAGGTGT) and a universal

fungal probe (Alexa 555-CTCTGGCTTCACCCTATTC) directed against the 16S rRNA

gene and 28S rRNA gene, respectively. DAPI was used to stain the nucleus.

Intestinal cell staining

The small intestine was fixed in Methacarn (Bioworld) and embedded in paraffin. Fixed

tissues were sectioned at 7 μm and deparaffinized in xylene followed by rehydration in

100% EtOH, 95% EtOH, 70% EtOH, and H_2O.

For tuft cells, slides were incubated in citrate buffer (10 mM Trisodium citrate, 0.05%

Tween 20, pH 6.0) for 20 min at 95 °C for antigen retrieval. Slides were blocked with 1%

bovine serum albumin (BSA) followed by overnight incubation with rabbit anti-mouse

DCAMKL1 (Abcam) antibodies at 4 °C. After incubation, slides were washed and

incubated with goat anti-rabbit Alexa Fluor 546 (Invitrogen) secondary antibodies for 1 h

at room temperature. Slides were counterstained with DAPI.

For goblet cells, slides were incubated with Alcian blue (StatLab) for 30 min followed

by washing in running tap water for 2 min. Slides were counterstained with hematoxylin

for 2 min followed by washing in running tap water for 30 sec. The slides were 10 dips in

the clarifier and bluing reagent with washing in tap water, respectively. Slides were

dehydrated in 95% EtOH, 100% EtOH, and xylene.

Scanning electron microscopy (SEM) & transmission electron microscopy (TEM)

For SEM, *K. pintolopesii* cultures were washed twice with sterile PBS, and a thin layer

was applied to the coverslips. The coverslips were fixed in fixation buffer (2.5%

glutaraldehyde, 0.15 M sodium cacodylate, 2% paraformaldehyde, 2 mM calcium chloride,

0.1 M sucrose, pH 7.4) followed by incubation for 2 h at room temperature. For the small

intestine, sections of the tissue were flushed with sterile PBS and then cut into 1 cm

sections. They were incubated in the same fixation buffer as previously described. After

incubation, the coverslips were washed three times with washing buffer (2 mM calcium

chloride, 0.15 M sodium cacodylate, 0.1 M sucrose). After washing, samples were

incubated overnight in 2% osmium tetroxide, 2.5% glutaraldehyde, 0.15 M sodium

cacodylate, and 2% paraformaldehyde. Samples were washed with DI water three times

and dehydrated in an EtOH gradient as follows: 20%, 50%, 70%, 90%, 95% and then

100%. Each EtOH step was incubated for 20 min at room temperature and then washed

twice with 100% EtOH. Samples were flash dehydrated using a critical point dryer and

then mounted on coverslips for imaging. Thermo Apreo SEM was used for imaging.

For TEM, *K. pintolopesii* cultures were pelleted in the 1.5 mL Eppendorf tubes. The fixation buffer was added to the tubes and incubated for 2 h at room temperature. The samples were then washed three times with the washing buffer. After washing, samples were incubated overnight in 2% osmium tetroxide, 2.5% glutaraldehyde, 0.15 M sodium cacodylate, and 2% paraformaldehyde. Samples were washed with DI water three times, followed by incubation overnight at 4 °C in saturated uranyl acetate solution. The next day, samples were dehydrated in the EtOH gradient as we described in SEM preparation and washed three times in 100% EtOH. Then, the samples were infiltrated with Hard Spurr's resin and rocked on a rotator at room temperature. The gradient started at 25% Hard Spurr's and 75% EtOH and gradually increased until 100% resin. Each step lasted 2 h, and the final 100% resin step was left overnight. The samples in resin were placed in resin molds and then let to harden for 18-24 h at 60 °C. The pellets in resin were cut with a knife into 100-200 nm sections and observed on PHILIPS 410 TEM.

Metabolomics

Metabolite extraction from serum and tissue samples. Frozen cecum lumen samples were weighed and ground on a cryomill (Retsch). Water-soluble polar metabolites from luminal contents were extracted with ice-cold 40:40:20 acetonitrile:methanol:water (25 mg tissue/mL extraction buffer), vortexed and incubated on ice for 10 min. Tissue samples were then centrifuged at 16,000 *xg* for 30 min at 4 °C. The tissue supernatants were transferred to new Eppendorf tubes and then centrifuged again at 16,000 *xg* for 30 min at 4 °C to remove residual debris before liquid chromatography coupled to a mass spectrometer (LC-MS) analysis. Extracts were immediately transferred to a mass spectrometry vial and analyzed on an LC-MS.

Metabolite Measurements. Metabolite separation and analysis were performed on a

Vanquish Horizon UHPLC system (Thermo Fisher Scientific) coupled to the Orbitrap

Exploris 240 Mass Spectrometer (Thermo Fisher Scientific) as reported previously [82]. The

hydrophilic interaction chromatography (HILIC) LC separation was performed on an

XBridge BEH Amide column (2.5 µm, 2.1 mm X 150 mm, Waters, Milford, MA). Solvent

A is 95% water in 5% acetonitrile with 40 mM ammonium hydroxide and 20 mM acetic

acid. Solvent B is 80% acetonitrile in 20% water with 40 mM ammonium hydroxide and

20 mM acetic acid. The gradient was 0 min, 100% B; 3 min, 100% B; 3.2 min, 90% B; 6.2

min, 90% B; 6.5 min, 80% B; 10.5 min, 80% B; 10.7 min, 70% B; 13.5 min, 70% B; 13.7

min, 45% B; 16 min, 45% B; 16.5 min, 100% B; and 22 min, 100% B. The flow rate was

300 µL/min. The column temperature was set to 30 °C. The autosampler temperature

was set to 4 °C, and the injection volume was 5 µL. MS scans were obtained in both

negative and positive ion modes with a resolution of 120,000 at m/z 200, in addition to an

automatic gain control target of 3×10^6 and m/z scan range of 72 to 1000.

Raw LC/MS data were converted to mzML format using the command line "msconvert"

utility [83]. Data were analyzed via MAVEN software and the annotations of the metabolites

were confirmed using our in-house library generated using the Mass Spectrometry

Metabolite Library (Millipore Sigma, supplied by IROA Technologies). MetaboAnalyst 6.0

[84] was used for statistical analyses and visualization.

***Heligmosomoides polygyrus* infection**

H. polygyrus L3 larvae were orally gavaged to mice (200 L3 larvae per mouse). After

infection, fecal samples were collected every week. Fecal samples were resuspended in

water and mixed with the same volume of super-saturated salt solution. The resuspended

fecal solution was immediately loaded on a McMaster counting chamber (Eggzamin) and

eggs were counted under the microscope. The counted eggs were normalized by fecal

weight.

Statistical analysis

All statistical analysis and plotting were performed on R (https://www.R-project.org) and

Prism (GraphPad). The normality of data was assessed by the Shapiro-Wilk test. If data

followed normal distribution, parametric tests (unpaired two-tailed t-test for two groups,

One-way ANOVA with Tukey's multiple comparisons test for more than two groups) were

performed. If data did not follow normal distribution, non-parametric tests (Mann-Whitney

test for two groups, Kruskal-Wallis test with Dunn's multiple comparisons test for more

than two groups) were performed to find significant differences between groups. For the

survival curve, the log-rank (Mantel-Cox) test was used to find significant differences

between groups. A two-way ANOVA test was used to find significant differences in body

weight changes at each time point between the two groups. $*p < 0.05$, $**p < 0.01$, $***p <$

0.001, $****p<0.0001$.

Data availability

Acknowledgments

Schematics were created with BioRender.com. We thank Dr. Richard Bennett (Brown University) for providing *S. cerevisiae* BY4717, *C. albicans* SC5314, *C. parapsilosis* ATCC22019, and *C. tropicalis* MYA-3404 strains. GH was supported by an American Association of Immunologists Careers in Immunology Fellowship. NNJ is supported by the Damon Runyon Cancer Research Foundation (DRG-2427-21) and NIAID 1K22AI177360-01. SV is supported by R01DK113265, P20GM109035 (pilot project) and Brown University Seed grant. LKB is supported by Searle Scholar Foundation and this work is also supported by Brown University Seed grant. The Thermo Apreo VS SEM was purchased with a high-end instrumentation grant from the Office of the Director at the National Institutes of Health (S10OD023461).

Author contributions

793   Conceptualization: SV, LKB, GH; Methodology: SV, LKB, GH, RY, MHH, NNJ, KC;

794   Investigation: SV, LKB, GH, RY, MHH, AB, KB, JP, OL, DMM, KC; Resources: SV, LKB,

795   NNJ; Funding acquisition: SV, LKB; Supervision: SV, LKB; Writing – original draft: SV,

796   LKB, GH; Writing – review & editing: SV, LKB, GH, KC

Competing interests

799   The authors declare no competing interests.

References

Koropatnick, T. A. *et al.* Microbial factor-mediated development in a host-bacterial mutualism. *Science* **306**, 1186-1188 (2004). https://doi.org:10.1126/science.1102218

Pais, R., Lohs, C., Wu, Y., Wang, J. & Aksoy, S. The obligate mutualist Wigglesworthia glossinidia influences reproduction, digestion, and immunity processes of its host, the tsetse fly. *Appl Environ Microbiol* **74**, 5965-5974 (2008). https://doi.org:10.1128/AEM.00741-08

Chung, H. *et al.* Gut immune maturation depends on colonization with a host-specific microbiota. *Cell* **149**, 1578-1593 (2012). https://doi.org:10.1016/j.cell.2012.04.037

Tannock, G. W., Miller, J. R. & Savage, D. C. Host specificity of filamentous, segmented microorganisms adherent to the small bowel epithelium in mice and rats. *Appl Environ Microbiol* **47**, 441-442 (1984). https://doi.org:10.1128/aem.47.2.441-442.1984

Gaboriau-Routhiau, V. *et al.* The key role of segmented filamentous bacteria in the coordinated maturation of gut helper T cell responses. *Immunity* **31**, 677-689 (2009). https://doi.org:10.1016/j.immuni.2009.08.020

Ivanov, II *et al.* Induction of intestinal Th17 cells by segmented filamentous bacteria. *Cell* **139**, 485-498 (2009). https://doi.org:10.1016/j.cell.2009.09.033

Koh, A. Y. Murine models of Candida gastrointestinal colonization and dissemination. *Eukaryot Cell* **12**, 1416-1422 (2013). https://doi.org:10.1128/EC.00196-13

8   Koh, A. Y., Kohler, J. R., Coggshall, K. T., Van Rooijen, N. & Pier, G. B. Mucosal

damage and neutropenia are required for Candida albicans dissemination. *PLoS*

Pathog **4**, e35 (2008). https://doi.org:10.1371/journal.ppat.0040035

9   Kumamoto, C. A. Inflammation and gastrointestinal Candida colonization. *Curr*

Opin Microbiol **14**, 386-391 (2011). https://doi.org:10.1016/j.mib.2011.07.015

10   Beura, L. K. *et al.* Normalizing the environment recapitulates adult human

immune traits in laboratory mice. *Nature* **532**, 512-516 (2016).

https://doi.org:10.1038/nature17655

11   Abolins, S. *et al.* The comparative immunology of wild and laboratory mice, Mus

musculus domesticus. *Nat Commun* **8**, 14811 (2017).

https://doi.org:10.1038/ncomms14811

12   Ma, J. *et al.* Laboratory mice with a wild microbiota generate strong allergic

immune responses. *Sci Immunol* **8**, eadf7702 (2023).

https://doi.org:10.1126/sciimmunol.adf7702

13   Yeung, F. *et al.* Altered Immunity of Laboratory Mice in the Natural Environment

Is Associated with Fungal Colonization. *Cell Host Microbe* **27**, 809-822 e806

(2020). https://doi.org:10.1016/j.chom.2020.02.015

14   Chen, Y. H. *et al.* Rewilding of laboratory mice enhances granulopoiesis and

immunity through intestinal fungal colonization. *Sci Immunol* **8**, eadd6910 (2023).

https://doi.org:10.1126/sciimmunol.add6910

15   Hedin, K. A. *et al.* Effects of broad-spectrum antibiotics on the colonisation of

probiotic yeast Saccharomyces boulardii in the murine gastrointestinal tract. *Sci*

Rep **12**, 8862 (2022). https://doi.org:10.1038/s41598-022-12806-0

16   Jiang, T. T. *et al.* Commensal Fungi Recapitulate the Protective Benefits of

Intestinal Bacteria. *Cell Host Microbe* **22**, 809-816 e804 (2017).

https://doi.org:10.1016/j.chom.2017.10.013

17   Mellado, E. *et al.* Sustained gastrointestinal colonization and systemic

dissemination by Candida albicans, Candida tropicalis and Candida parapsilosis

in adult mice. *Diagn Microbiol Infect Dis* **38**, 21-28 (2000).

https://doi.org:10.1016/s0732-8893(00)00165-6

18   Ost, K. S. & Round, J. L. Commensal fungi in intestinal health and disease. *Nat*

Rev Gastroenterol Hepatol **20**, 723-734 (2023). https://doi.org:10.1038/s41575-

023-00816-w

19   Roberts, K. *et al.* Candida tropicalis Affects Candida albicans Virulence by

Limiting Its Capacity to Adhere to the Host Intestinal Surface, Leading to

Decreased Susceptibility to Colitis in Mice. *J Fungi (Basel)* **10** (2024).

https://doi.org:10.3390/jof10040245

20   Bacher, P. *et al.* Human Anti-fungal Th17 Immunity and Pathology Rely on

Cross-Reactivity against Candida albicans. *Cell* **176**, 1340-1355 e1315 (2019).

https://doi.org:10.1016/j.cell.2019.01.041

21   Shao, T. Y. *et al.* Commensal Candida albicans Positively Calibrates Systemic

Th17 Immunological Responses. *Cell Host Microbe* **25**, 404-417 e406 (2019).

https://doi.org:10.1016/j.chom.2019.02.004

22   Leonardi, I. *et al.* Mucosal fungi promote gut barrier function and social behavior

via Type 17 immunity. *Cell* **185**, 831-846 e814 (2022).

https://doi.org:10.1016/j.cell.2022.01.017

23   Huang, L., Nazarova, E. V., Tan, S., Liu, Y. & Russell, D. G. Growth of

Mycobacterium tuberculosis in vivo segregates with host macrophage

metabolism and ontogeny. *J Exp Med* **215**, 1135-1152 (2018).

https://doi.org:10.1084/jem.20172020

24   Culemann, S. *et al.* Stunning of neutrophils accounts for the anti-inflammatory

effects of clodronate liposomes. *J Exp Med* **220** (2023).

https://doi.org:10.1084/jem.20220525

25   Gerbe, F. *et al.* Intestinal epithelial tuft cells initiate type 2 mucosal immunity to

helminth parasites. *Nature* **529**, 226-230 (2016).

https://doi.org:10.1038/nature16527

26   Howitt, M. R. *et al.* Tuft cells, taste-chemosensory cells, orchestrate parasite type

2 immunity in the gut. *Science* **351**, 1329-1333 (2016).

https://doi.org:10.1126/science.aaf1648

27   Sasaki, S., Ishida, Y., Nishio, N., Ito, S. & Isobe, K. Thymic involution correlates

with severe ulcerative colitis induced by oral administration of dextran sulphate

sodium in C57BL/6 mice but not in BALB/c mice. *Inflammation* **31**, 319-328

(2008). https://doi.org:10.1007/s10753-008-9081-3

28   Tsuchiya, T. *et al.* Role of gamma delta T cells in the inflammatory response of

experimental colitis mice. *J Immunol* **171**, 5507-5513 (2003).

https://doi.org:10.4049/jimmunol.171.10.5507

29   Sekeresova Kralova, J. *et al.* Competitive fungal commensalism mitigates

candidiasis pathology. *J Exp Med* **221** (2024).

https://doi.org:10.1084/jem.20231686

30 Liao, Y. *et al.* Fungal symbiont transmitted by free-living mice promotes type 2

immunity. *Nature* (2024). https://doi.org:10.1038/s41586-024-08213-2

31 Hu, J. *et al.* Core-predominant gut fungus Kazachstania slooffiae promotes

intestinal epithelial glycolysis via lysine desuccinylation in pigs. *Microbiome* **11**,

31 (2023). https://doi.org:10.1186/s40168-023-01468-3

32 Zhang, H. *et al.* Immune activation of characteristic gut mycobiota Kazachstania

pintolopesii on IL-23/IL-17R signaling in ankylosing spondylitis. *Front Cell Infect*

Microbiol **12**, 1035366 (2022). https://doi.org:10.3389/fcimb.2022.1035366

33 Gallotti, A. C. *et al.* Kazachstania slooffiae, an emerging pathogen to watch for in

humans? *Med Mycol Case Rep* **42**, 100604 (2023).

https://doi.org:10.1016/j.mmcr.2023.08.007

34 Brunet, K. *et al.* Invasive Candida bovina Infection, France. *Emerg Infect Dis* **26**,

626-627 (2020). https://doi.org:10.3201/eid2603.191371

35 Flannigan, K. L. & Denning, T. L. Segmented filamentous bacteria-induced

immune responses: a balancing act between host protection and autoimmunity.

Immunology **154**, 537-546 (2018). https://doi.org:10.1111/imm.12950

36 Baumler, A. & Fang, F. C. Host specificity of bacterial pathogens. *Cold Spring*

Harb Perspect Med **3**, a010041 (2013).

https://doi.org:10.1101/cshperspect.a010041

37 Gutierrez-Garcia, K. *et al.* A conserved bacterial genetic basis for commensal-

host specificity. *Science* **386**, 1117-1122 (2024).

https://doi.org:10.1126/science.adp7748

38   Limon, J. J. *et al.* Malassezia Is Associated with Crohn's Disease and

917   Exacerbates Colitis in Mouse Models. *Cell Host Microbe* **25**, 377-388 e376

918   (2019). https://doi.org:10.1016/j.chom.2019.01.007

919   39   Li, X. V. *et al.* Immune regulation by fungal strain diversity in inflammatory bowel

920   disease. *Nature* **603**, 672-678 (2022). https://doi.org:10.1038/s41586-022-04502-

921   w

922   40   Rosshart, S. P. *et al.* Wild Mouse Gut Microbiota Promotes Host Fitness and

923   Improves Disease Resistance. *Cell* **171**, 1015-1028 e1013 (2017).

924   https://doi.org:10.1016/j.cell.2017.09.016

925   41   Rosshart, S. P. *et al.* Laboratory mice born to wild mice have natural microbiota

926   and model human immune responses. *Science* **365** (2019).

927   https://doi.org:10.1126/science.aaw4361

928   42   Huggins, M. A. *et al.* Microbial Exposure Enhances Immunity to Pathogens

929   Recognized by TLR2 but Increases Susceptibility to Cytokine Storm through

930   TLR4 Sensitization. *Cell Rep* **28**, 1729-1743 e1725 (2019).

931   https://doi.org:10.1016/j.celrep.2019.07.028

932   43   Netea, M. G., Joosten, L. A., van der Meer, J. W., Kullberg, B. J. & van de

933   Veerdonk, F. L. Immune defence against Candida fungal infections. *Nat Rev*

934   *Immunol* **15**, 630-642 (2015). https://doi.org:10.1038/nri3897

935   44   Underhill, D. M. & Iliev, I. D. The mycobiota: interactions between commensal

936   fungi and the host immune system. *Nat Rev Immunol* **14**, 405-416 (2014).

937   https://doi.org:10.1038/nri3684

45   Han, G., Luong, H. & Vaishnava, S. Low abundance members of the gut

microbiome exhibit high immunogenicity. *Gut Microbes* **14**, 2104086 (2022).

https://doi.org:10.1080/19490976.2022.2104086

46   Muller, P. A., Matheis, F. & Mucida, D. Gut macrophages: key players in

intestinal immunity and tissue physiology. *Curr Opin Immunol* **62**, 54-61 (2020).

https://doi.org:10.1016/j.coi.2019.11.011

47   Mowat, A. M. & Bain, C. C. Mucosal macrophages in intestinal homeostasis and

inflammation. *J Innate Immun* **3**, 550-564 (2011).

https://doi.org:10.1159/000329099

48   Wang, S., Ye, Q. H., Zeng, X. F. & Qiao, S. Y. Functions of Macrophages in the

Maintenance of Intestinal Homeostasis. *J Immunol Res* **2019** (2019).

https://doi.org:Artn 1512969

10.1155/2019/1512969

49   Barber, M. F. & Fitzgerald, J. R. Mechanisms of host adaptation by bacterial

pathogens. *Fems Microbiol Rev* **48** (2024).

https://doi.org:10.1093/femsre/fuae019

50   Tso, G. H. W. *et al.* Experimental evolution of a fungal pathogen into a gut

symbiont. *Science* **362**, 589-595 (2018). https://doi.org:10.1126/science.aat0537

51   Lecuit, M. *et al.* A single amino acid in E-cadherin responsible for host specificity

towards the human pathogen Listeria monocytogenes. *EMBO J* **18**, 3956-3963

(1999). https://doi.org:10.1093/emboj/18.14.3956

52   Roble, G. S., Gillespie, V. & Lipman, N. S. Infectious disease survey of Mus

musculus from pet stores in New York City. *J Am Assoc Lab Anim Sci* **51**, 37-41

(2012).

53   Parker, S. E., Malone, S., Bunte, R. M. & Smith, A. L. Infectious diseases in wild

mice (Mus musculus) collected on and around the University of Pennsylvania

(Philadelphia) Campus. *Comp Med* **59**, 424-430 (2009).

54   Dammann, P. *et al.* Infectious microorganisms in mice (Mus musculus)

purchased from commercial pet shops in Germany. *Lab Anim* **45**, 271-275

(2011). https://doi.org:10.1258/la.2011.010183

55   Gardes, M. & Bruns, T. D. ITS primers with enhanced specificity for

basidiomycetes--application to the identification of mycorrhizae and rusts. *Mol*

Ecol **2**, 113-118 (1993). https://doi.org:10.1111/j.1365-294x.1993.tb00005.x

56   White, T. J., Bruns, T., Lee, S. & Taylor, J. in *PCR Protocols: A Guide to*

Methods and Applications   (eds M. A. Innis, D. H. Gelfand, J. J. Sninsky, & T. J.

White) 315-322 (Academic Press, 1990).

57   Beura, L. K. *et al.* Lymphocytic choriomeningitis virus persistence promotes

effector-like memory differentiation and enhances mucosal T cell distribution. *J*

Leukoc Biol **97**, 217-225 (2015). https://doi.org:10.1189/jlb.1HI0314-154R

58   Grizotte-Lake, M. *et al.* Commensals Suppress Intestinal Epithelial Cell Retinoic

Acid Synthesis to Regulate Interleukin-22 Activity and Prevent Microbial

Dysbiosis. *Immunity* **49**, 1103-1115 e1106 (2018).

https://doi.org:10.1016/j.immuni.2018.11.018

981 59 Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina

982   sequence data. *Bioinformatics* **30**, 2114-2120 (2014).

983   https://doi.org:10.1093/bioinformatics/btu170

984 60 Wick, R. R., Judd, L. M., Gorrie, C. L. & Holt, K. E. Completing bacterial genome

985   assemblies with multiplex MinION sequencing. *Microb Genom* **3**, e000132

986   (2017). https://doi.org:10.1099/mgen.0.000132

987 61 De Coster, W., D'Hert, S., Schultz, D. T., Cruts, M. & Van Broeckhoven, C.

988   NanoPack: visualizing and processing long-read sequencing data. *Bioinformatics*

989   **34**, 2666-2669 (2018). https://doi.org:10.1093/bioinformatics/bty149

990 62 Marcais, G. & Kingsford, C. A fast, lock-free approach for efficient parallel

991   counting of occurrences of k-mers. *Bioinformatics* **27**, 764-770 (2011).

992   https://doi.org:10.1093/bioinformatics/btr011

993 63 Vurture, G. W. *et al.* GenomeScope: fast reference-free genome profiling from

994   short reads. *Bioinformatics* **33**, 2202-2204 (2017).

995   https://doi.org:10.1093/bioinformatics/btx153

996 64 Kolmogorov, M., Yuan, J., Lin, Y. & Pevzner, P. A. Assembly of long, error-prone

997   reads using repeat graphs. *Nat Biotechnol* **37**, 540-546 (2019).

998   https://doi.org:10.1038/s41587-019-0072-8

999 65 Vaser, R., Sovic, I., Nagarajan, N. & Sikic, M. Fast and accurate de novo

1000   genome assembly from long uncorrected reads. *Genome Res* **27**, 737-746

1001   (2017). https://doi.org:10.1101/gr.214270.116

66   Walker, B. J. *et al.* Pilon: an integrated tool for comprehensive microbial variant

detection and genome assembly improvement. *PLoS One* **9**, e112963 (2014).

https://doi.org:10.1371/journal.pone.0112963

67   Gurevich, A., Saveliev, V., Vyahhi, N. & Tesler, G. QUAST: quality assessment

tool for genome assemblies. *Bioinformatics* **29**, 1072-1075 (2013).

https://doi.org:10.1093/bioinformatics/btt086

68   Manni, M., Berkeley, M. R., Seppey, M., Simao, F. A. & Zdobnov, E. M. BUSCO

Update: Novel and Streamlined Workflows along with Broader and Deeper

Phylogenetic Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral

Genomes. *Mol Biol Evol* **38**, 4647-4654 (2021).

https://doi.org:10.1093/molbev/msab199

69   Darling, A. C., Mau, B., Blattner, F. R. & Perna, N. T. Mauve: multiple alignment

of conserved genomic sequence with rearrangements. *Genome Res* **14**, 1394-

1403 (2004). https://doi.org:10.1101/gr.2289704

70   Eren, A. M. *et al.* Community-led, integrated, reproducible multi-omics with

anvi'o. *Nat Microbiol* **6**, 3-6 (2021). https://doi.org:10.1038/s41564-020-00834-3

71   Emms, D. M. & Kelly, S. OrthoFinder: phylogenetic orthology inference for

comparative genomics. *Genome Biol* **20**, 238 (2019).

https://doi.org:10.1186/s13059-019-1832-y

72   Bolyen, E. *et al.* Reproducible, interactive, scalable and extensible microbiome

data science using QIIME 2. *Nat Biotechnol* **37**, 852-857 (2019).

https://doi.org:10.1038/s41587-019-0209-9

73   Callahan, B. J. *et al.* DADA2: High-resolution sample inference from Illumina

amplicon data. *Nat Methods* **13**, 581-583 (2016).

https://doi.org:10.1038/nmeth.3869

74   Yilmaz, P. *et al.* The SILVA and "All-species Living Tree Project (LTP)"

taxonomic frameworks. *Nucleic Acids Res* **42**, D643-648 (2014).

https://doi.org:10.1093/nar/gkt1209

75   Rivers, A. R., Weber, K. C., Gardner, T. G., Liu, S. & Armstrong, S. D. ITSxpress:

Software to rapidly trim internally transcribed spacer sequences with quality

scores for marker gene analysis. *F1000Res* **7**, 1418 (2018).

https://doi.org:10.12688/f1000research.15704.1

76   Nilsson, R. H. *et al.* The UNITE database for molecular identification of fungi:

handling dark taxa and parallel taxonomic classifications. *Nucleic Acids Res* **47**,

D259-D264 (2019). https://doi.org:10.1093/nar/gky1022

77   Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-

21 (2013). https://doi.org:10.1093/bioinformatics/bts635

78   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

https://doi.org:10.1186/s13059-014-0550-8

79   Gu, Z. Complex heatmap visualization. *Imeta* **1**, e43 (2022).

https://doi.org:10.1002/imt2.43

80   Franzen, O., Gan, L. M. & Bjorkegren, J. L. M. PanglaoDB: a web server for

exploration of mouse and human single-cell RNA sequencing data. *Database*

(Oxford) **2019** (2019). https://doi.org:10.1093/database/baz046

81   Xu, S. *et al.* Using clusterProfiler to characterize multiomics data. *Nat Protoc* **19**,

3292-3320 (2024). https://doi.org:10.1038/s41596-024-01020-z

82   Wang, L. *et al.* Peak Annotation and Verification Engine for Untargeted LC-MS

Metabolomics. *Anal Chem* **91**, 1838-1846 (2019).

https://doi.org:10.1021/acs.analchem.8b03132

83   Adusumilli, R. & Mallick, P. Data Conversion with ProteoWizard msConvert.

Methods Mol Biol **1550**, 339-368 (2017). https://doi.org:10.1007/978-1-4939-

6747-6_23

84   Pang, Z. *et al.* MetaboAnalyst 6.0: towards a unified platform for metabolomics

data processing, analysis and interpretation. *Nucleic Acids Res* **52**, W398-W406

(2024). https://doi.org:10.1093/nar/gkae253

1058

1059

Figures

1061

Fig. 1. *K. pintolopesii* **is the dominant fungus in the gut of pet store mice.** (A) Representative scanning electron microscopy (SEM) image of the small intestine of laboratory mice and pet store mice. (B) Representative fluorescence *in situ* hybridization (FISH) images of fungal colonization in the small intestine and the colon of laboratory mice and pet store mice. Tissues were stained with DAPI (nuclei; blue), EUB338 probe (bacteria; green), and 28S probe (fungi; red). (C) Fungal burden in the feces of laboratory mice (*n* = 5) and pet store mice (*n* = 8). (D) Fungal burden in the intestinal tract in the pet

store mice (n = 3 for tissues and n = 4 for feces). (E) Bar plot of relative abundance of

fungal species from ITS-marker gene sequencing at different sections of the intestine –

luminal and mucosal part of the small intestine and the colon (n = 4 for SI-mucosal and n

= 5 for other samples). Fungal burden (F) (RI, n = 8; MN, n = 6; TX, n =3) and bar plot of

relative abundance of fungal species (G) in the feces of pet store mice (RI, n = 5; MN, n

= 6; TX, n =8) in Rhode Island (RI), Minnesota (MI), and Texas (TX) in the USA. Bars

indicate mean ± SEM. Significances were determined by the Kruskal-Wallis test with

Dunn's multiple comparisons test or the Mann-Whitney test. *p < 0.05 and **p < 0.01.

L.o.D., limit of detection.

1078

1079

Fig. 2. *K. pintolopesii* stably colonizes laboratory mice irrespective of gut bacteria and provides colonization resistance against non-murine fungi. (A) Fungal burden in the feces of mice after *K. pintolopesii* treatment (n = 5). (B) Representative images of *K. pintolopesii* colonization at 13 days post-colonization in the small intestine and colon. Tissues were stained with DAPI (nuclei; blue), EUB338 probe (bacteria; green), and 28S probe (fungi; red). (C-E) 16S rRNA gene sequencing was performed to explore fecal bacterial communities (Control, n = 4; Kp, n = 6). (C) Alpha diversity (Shannon) at 14 days post-colonization in the feces of control and *K. pintolopesii*-colonized mice. (D) PCoA plots based on the weighted-UniFrac distances at 14 days post-colonization in the feces. Significances were determined by PERMANOVA. (E) Bar plot of relative abundance of the bacterial genus from 16S rRNA gene sequencing of the feces. (F) Bacterial load in the feces of the control and *K. pintolopesii*-colonized mice (Control, n = 4; Kp, n = 6). Bacterial load was expressed as a 2^{-Ct} value and normalized to the weight of the starting fecal material. (G) Fungal burden in the feces of GF mice after *K. pintolopesii* treatment (n = 5). (H) Scheme of fungal competition experiment in the conventional and germ-free mice. (I) Line plot of relative abundance of fungal species from ITS-marker gene sequencing of the feces at different time points in conventional and germ-free mice (n = 5 per group). Dots indicate mean ± SEM. Significances were determined by unpaired two-tailed t-test. Kp, *K. pintolopesii*-colonized mice. CV, conventional mice. GF, germ-free mice

1101

Fig. 3. The systemic challenge with *K. pintolopesii* is efficiently resolved in a neutrophil-independent manner. (A) Frequency of circulating neutrophils in the peripheral blood at 5 (Control, *n* = 13; Kp, *n* = 17), 8 (Control, *n* = 30; Kp, *n* = 35), and 20 days (Control, *n* = 4; Kp, *n* = 5) post-colonization of *K. pintolopesii*. Neutrophil populations were gated as CD45+ B220- CD11b+ Ly6G+. (B) Scheme of the systemic challenge with *K. pintolopesii* or *C. albicans* (*n* = 5 per group). (C) The survival curve after *K. pintolopesii* or *C. albicans* systemic infection. Fungal burden (D) and *Il17* gene expression (E) in the

kidney after *C. albicans* or *K. pintolopesii* infection. (F) Experimental schedule of

neutrophils depletion and *K. pintolopesii* infection (*n* = 5 per group). (G) The survival curve

after *K. pintolopesii* systemic infection with or without neutrophils depletion. (H) Fungal

burden in the kidney after *K. pintolopesii* systemic infection with or without neutrophils

depletion. Bars indicate mean ± SEM. Significances were determined by unpaired two-

tailed t-test. For the survival curve, the log-rank (Mantel-Cox) test was used to find

significant differences between groups. *p < 0.05. L.o.D., limit of detection. Kp, *K.*

pintolopesii-colonized mice.

1117

1118

Fig. 4. *K. pintolopesii* **colonization increases the macrophage population in the**

small intestine. (A-C) Transcriptional profiles of the small intestines from control and *K.*

pintolopesii-colonized mice were generated using bulk RNA sequencing (*n* = 4 per

group). (A) Volcano plot of the transcriptional changes after *K. pintolopesii* colonization

in the small intestine. Down-regulated genes and up-regulated after *K. pintolopesii*

colonization were presented as blue and red dots, respectively. (B) The heat map of

macrophage activation-associated genes. (C) Bar plots from gene ontology (GO)

enrichment analysis. Macrophage activation-associated terms were chosen from the list

of significantly different GO terms. (D) Representative flow cytometry plots and

frequency of macrophages in the small intestinal lamina propria of control and *K.*

pintolopesii-colonized mice at 14 days post-colonization (Control, $n = 4$; Kp, $n = 5$).

Macrophage populations were gated as live $CD45^+$ $Ly6G^-$ $CD11b^+$ MHC II^+ $F4/80^+$

$CD64^+$. (E) Experimental schedule of clodronate injection and *K. pintolopesii*

colonization. (F) Representative flow cytometry plots and frequency of neutrophils in the

peripheral blood of clodronate-treated (Clod, $n = 4$), *K. pintolopesii*-colonized (Kp, $n =$

5), and *K. pintolopesii*-colonized clodronate-treated mice (Kp+Clod, $n = 4$) at 8 days

post-colonization. Neutrophil populations were gated as $CD45^+$ $B220^-$ $CD11b^+$ $Ly6G^+$.

Bars indicate mean ± SEM. Significances were determined by the unpaired two-tailed t-

test or One-way ANOVA with Tukey's multiple comparisons test. $*p < 0.05$ and $**p <$

0.01. Kp, *K. pintolopesii*-colonized mice. Clod, clodronate-treated mice.

1139

1140

Fig. 5. *K. pintolopesii* colonization increases the tuft cell population and induces type 2 immune responses in the gut and protects the host against *H. polygyrus* infection. (A) The heat map of expression of the tuft cell and the goblet cell marker genes in the small intestine (*n* = 4 per group). (B) Representative image and the number of tuft cells in the small intestine of control and *K. pintolopesii*-colonized mice. Tissues were

stained with DAPI (nuclei; blue) and DCLK1 (tuft cells; green). 1-7 images were used for

tuft cell counting per mouse with 4–6 mice per group. (C) Representative image and the

number of goblet cells in the small intestine of control and *K. pintolopesii*-colonized mice.

Tissues were stained with hematoxylin (nuclei) and alcian blue (goblet cells). 6-15 villi

were chosen for goblet cell counting per mouse with 4–5 mice per group. (D)

Representative flow cytometry plots and frequency of Th2 in the small intestinal lamina

propria of control and *K. pintolopesii*-colonized mice (n = 10 per group). Th2 populations

were gated as live $CD45^+$ Lin^+ $CD90.2^+$ $CD4^+$ $GATA3^+$ $Foxp3^-$ $ST2^+$. (E) Fungal burden in

the feces of BALB/c mice after *K. pintolopesii* colonization (n = 4 per group). (F)

Frequency of Th2 in the small intestinal lamina propria of control and *K. pintolopesii*-

colonized BALB/c mice (n = 4 per group). (G) Scheme of the *H. polygyrus* challenge. (H)

The quantification of worm eggs in the feces of the control and *K. pintolopesii*-colonized

mice (Control, n = 9; Kp, n = 10). Bars indicate mean ± SEM. Significances were

determined by the unpaired two-tailed t-test or Mann-Whitney test. *$p < 0.05$, **$p < 0.01$,

and ****$p < 0.0001$. Kp, *K. pintolopesii*-colonized mice.

1161

Fig. 6. *K. pintolopesii* **from guinea pigs shows different characteristics in colonization, microbiome, metabolome, and genome compared with mouse isolate.** (A) Scheme of the colonization of *K. pintolopesii* from the pet store mouse intestine (Kp-M) and *K. pintolopesii* strain from the pet store guinea pig intestine (Kp-GP) in the laboratory conventional mice. (B) Fungal burden of Kp-M and Kp-GP in the feces at different time points. Fecal samples were collected from 4-9 mice per group at each time point. (C) Fungal burden of Kp-M and Kp-GP in the feces of germ-free mice at 14 days post-colonization ($n = 5$ per group). (D) Scheme of the colonization of Kp-M, Kp-GP, and *C. albicans* in the altered Schaedler flora (ASF)-colonized ex-GF mice ($n = 4$ per group). (E) The fungal burden of Kp-M, Kp-GP, and *C. albicans* in the ASF-colonized ex-GF mice. (F) PCoA plots based on the weighted-UniFrac distances in the feces at 14 days post-colonization in the feces of ASF mice with different strains of fungal colonization. Significances were determined by PERMANOVA. (G) Bar plot of relative abundance of ASF species from 16S rRNA gene sequencing of the feces ASF mice with different strains of fungal colonization. (H) Sparse partial least squares discriminant analysis (sPLS-DA) plot of the cecal metabolome. (I) Heatmap of the top 33 different metabolites in the cecal contents of ASF mice with different strains of fungal colonization and co-occurrence results between metabolites and bacterial abundances. Different metabolites were chosen based on the one-way ANOVA test results. Bars indicate mean ± SEM. Significances were determined by the unpaired two-tailed t-test or Mann-Whitney test. *$p < 0.05$ and ***$p < 0.001$. L.o.D., limit of detection. Kp-M, *K. pintolopesii* from the mouse-colonized mice. Kp-GP, *K. pintolopesii* from the guinea pig-colonized mice. Ca, *C. albicans*-colonized mice.

1186

Fig. 7. Bacteria-dependent *K. pintolopesii* colonization in the mouse gut. *K. pintolopesii,* a dominant fungal species in pet store mice, stably colonizes the intestine of laboratory mice, outcompetes non-murine commensal fungi, and persists independently of gut bacteria. Its colonization expands intestinal macrophage populations, which is crucial in preventing systemic immune recognition. Additionally, *K. pintolopesii* induces type 2 immunity, conferring cross-kingdom protection against helminth infection. However,

1193   *K. pintolopesii* strains from non-murine hosts, such as guinea pigs (Kp-GP), fail to

1194   colonize laboratory mice with intact gut bacterial community despite high genomic

1195   similarity to mouse-derived strains (Kp-M). In gnotobiotic ASF mice, the Kp-GP strain

1196   disrupts bacterial communities and their associated metabolomes, a phenomenon not

1197   observed with the Kp-M strain.