Longitudinal and cross-sectional selection on flowering traits in a self-incompatible annual.

Laura Leventhal[1,2,3,8], Karen J. Bai[4,5], Madeline A. E. Peters[4,6], Emily J. Austen[4,5,7], Arthur E. Weis[4,5], Jennifer L. Ison[4,5,8]

[1]Department of Biology, Stanford University, Stanford, CA 94305, USA.
[2]Department of Integrative Biology, University of California Berkeley, Berkeley, CA 94720, USA
[3]Department of Plant Biology, Carnegie Institution for Science, Stanford, CA 94305, USA.
[4]Department of Ecology and Evolutionary Biology, University of Toronto, Toronto, ON, Canada
[5]Koffler Scientific Reserve, University of Toronto, 17000 Dufferin Street, King City, ON, Canada
[6]Michigan Institute for Data and AI in Society, University of Michigan, Ann Arbor, MI 48109, USA
[7]Department of Biology, Mount Allison University, Sackville, NB, Canada
[8]Department of Biology, College of Wooster, Wooster, OH 44691 USA

Corresponding author: Laura Leventhal; lauracleventhal@gmail.com

Abstract

Net selection on a trait reflects the association of phenotype to fitness, across an entire life cycle. This longitudinal estimate of selection can be viewed as the summation of selection episodes, each characterized by a cross-sectional estimate. Selection may be consistent in direction and strength across episodes for some traits, fluctuating in others, and for some, concentrated in a single intense event. Additionally, while selection on plant reproductive traits is predicted to be stronger through male fitness than female fitness, male fitness remains less studied. We investigated how selection on flowering traits in *Brassica rapa* varied temporally and spatially by measuring male reproductive fitness in four experimental populations with two spatial arrangements. To estimate longitudinal and cross-sectional selection, we introduced male-sterile plants at successive intervals. We genotyped over 3000 plants and calculated selection on flowering time, duration, and total flowers. Cross-sectional analyses revealed directional selection was common, but patterns were masked by longitudinal estimates. Comparing selection gradients and coefficients revealed conflicting selection directions, suggesting the importance of trait correlations in selection estimates. Spatial population arrangement significantly impacted pollen movement and the strength of selection, demonstrating how breeding timing, spatial aggregation, and male fitness interact to create complex evolutionary dynamics.

Keywords

gene traps, male fitness, selection, Brassica rapa, flowering time, trait variation

Introduction

Unlike many animal species with highly synchronized breeding periods lasting only days or hours (Fogarty et al., 2012; McDiarmid, 1994; Schreiber & Burger, 2001; Urquhart, 1960), most flowering plants deploy reproductive structures sequentially over weeks or even months. Pollen transfer from donor to recipient is largely passive, mediated by biotic or abiotic vectors whose effectiveness fluctuates with environmental conditions (Ackerman, 2000; Elzinga et al., 2007; Polatto et al., 2014). Consequently, a plant's reproductive success represents the sum of all day-to-day mating events within an entire breeding season. The association between trait values and realized mating (i.e., the direction and magnitude of selection on plant traits), could in theory vary across dates. However, many studies cannot evaluate reproductive success on such a fine scale. To address this limitation, we focus on the temporal scheduling of flower deployment and its impact on male reproductive success through pollen export over the entire breeding season. By dissecting the breeding season into discrete temporal windows, we can analyze how selection on male function varies day-to-day within a season, revealing how phenological traits shape reproductive outcomes across the flowering season.

Within-season variation in selection is most effectively demonstrated by cross-sectional selection analyses, which measure selection on a subset of the population at a particular point in time (Endler, 1986; Kingsolver et al., 2001). While longitudinal selection studies reveal the net direction and intensity of selection by measuring the relationship between phenotype and lifetime fitness proxies, the cross-sectional approach can illuminate the mechanistic underpinnings of differential survival, mating, and reproduction throughout a breeding season, i.e., the *drivers* of longitudinal selection. Comparing these approaches allows us to determine whether longitudinal selection arises through steady and consistent selection across an entire season or from a few intense selection events at specific timepoints. Understanding how selection varies across shorter, episodic temporal scales can provide critical insights into trait

evolution (Cockburn et al., 2008; Hoekstra et al., 2001; Losos et al., 2006; Walsh & Blows,

2009).

Natural fluctuations across a breeding season in population density of mates, competitors,

predators, and mutualists could give rise to within-season variation in the direction or magnitude

of cross-sectional selection. Such temporally variable spatial structures can influence selection

patterns, as documented in animal systems (Daupagne et al., 2023; Gyulavári et al., 2017;

Kasumovic et al., 2008; Punzalan et al., 2010; Wacker et al., 2014) and less frequently in plant

systems (Parachnowitsch et al., 2012; Parachnowitsch & Caruso, 2008). In flowering plants,

density-dependent selection can be mediated by multiple, non-exclusive mechanisms. For

instance, the density of competitors influences selection in flowering plants as competition for

pollinators can arise when there is temporal overlap with blooms of competing species

(Flanagan et al., 2011; Landry, 2013; Levin & Anderson, 1970; Richardson et al., 2021).

Besides competitors, within-season variation in the pollinator community itself (composition,

abundance, effectiveness) can influence selection (CaraDonna et al., 2017; Fisogni et al., 2016;

Ison et al., 2018). Finally, population-level flowering density varies as plants enter and exit the

flowering season and the individual-level floral display size rises and falls over time, both of

which affect both within-species and within-individual competition patterns for resources (e.g.

(Austen et al., 2015; Feldman, 2008; Grindeland et al., 2005; Makino et al., 2007; Thomson,

1980)). If male fitness (seed siring success) is limited by access to mates, we might expect

selection through male fitness to be particularly vulnerable to within-season changes in

conspecific density and pollinator service.

Despite the importance of male fitness components, our understanding of selection in plants is

heavily biased toward female fitness, as measuring seeds produced is more straightforward

than determining the paternity of seeds sired. This bias was illustrated in a meta-analysis of

selection flowering time, where only 5.7% (5/87) of studies evaluated selection via male fitness

(Munguía-Rosas et al., 2011). Because female fitness depends on resources and male fitness

depends on mate opportunity (Bateman & Innes, 1948), there may be a disagreement among

selection estimates depending on the fitness input (Conner et al., 1996; Delph & Ashman, 2006;

Hou et al., 2024; Tsuji et al., 2020). When male and female fitness optima for the same trait

differ, evolutionary outcomes could lead to sexually antagonistic selection, which may stifle the

opportunity for the population to reach the optimal trait value for either sex (H. Wang et al.,

2021).

Consider how components of the flowering schedule such as flowering onset, duration, and total

flower production shape male fitness outcomes. Variability in schedule can affect population-

level synchrony, influencing both mate availability and mating frequency (Augspurger, 1983;

Devaux & Lande, 2010; Fox, 2003; Weis et al., 2005). When flowering schedules vary among

individuals, plants enter and exit their reproductive phase asynchronously, creating a dynamic

environment in which selection acts. To illustrate our conceptual framework, we imagine two

hypothetical scenarios relating variation in these flowering traits and selection (**Fig. 1**). In

scenario 1 (**Fig. 1A**), flowering time (early, mid, or late) is uncorrelated with (or positively

covaries with) duration or total number of flowers. Thus, selection through male fitness favors

early flowering onset early in the season, and late flowering onset at the end. This shift in

selection is driven entirely by which plants are in flower. In scenario 2 (**Fig. 1B**), duration and

total flowers negatively covary with flowering time, thus early flowering plants have the longest

duration of flowering. In this case, early flowering plants maintain presence throughout the

season, but selection could weaken over time as more plants enter the mating pool. Note that

these scenarios assume no variation in per-flower pollinator service across the season. A

decline in per-flower transfer in any one of the three depicted intervals would diminish the

contribution of that episode to net selection. Temporal variation in factors affecting pollen

transfer and correlations between these factors can create heterogeneous selection pressures

across the flowering season. This raises the possibility of selection shifting in direction and

intensity on flowering traits within a flowering season.

One difficulty in estimating selection on the flowering schedule through the pollen-export fitness

component arises when female qualities vary across time. Pollen export is intrinsically a trait of

the donor but must be quantified through the number of seeds sired. This poses no problem if

all females are of equal and constant quality, i.e., invariant in the probability that pollen receipt

results in offspring production. However, flowers produced toward the end of the flowering

season typically have reduced fruit maturation rates (Austen et al., 2015; Weis & Kossler, 2004).

This decline can stem from priority effects: early flowers access the full pool of maternal

resources, which become progressively depleted for later blooming flowers. The "gene-trap"

method developed by Devaux & Lande (2010) and refined by Ison & Weis (2017) can help

circumvent the priority effect bias. This approach involves introducing successive batches of

genetically-uniform pollen-recipient plants– all at identical developmental stages– into the

experimental population, followed by paternity analysis of their seeds. By controlling for variation

among pollen recipients, this method reveals the strictly male component of selection through

analysis of covariance between donor traits and siring success.

Research goals

In this study, we investigate how selection via male fitness on flowering schedule components –

flowering time, flowering duration, and total flower production– varies throughout the breeding

season in *Brassica rapa* through the gene-trap method. Our primary research goal is to evaluate

if selection gradients measured across the entire season (longitudinal selection) emerge from

consistent selection or from episodic bursts of selection (cross-sectional selection). We extend

this goal by also examining if the longitudinal and cross-sectional selection estimates vary

based on the spatial arrangement of plants within a population. Another research goal is to

quantify if trait correlations create conflicts between direct and total selection. Lastly, to

contextualize effects of spatial arrangement, we examine how changes to plant spatial

arrangement affect the distances pollen travels to fertilize seeds. By comparing selection at

different temporal scales and spatial contexts, we aim to better understand the evolutionary

dynamics shaping flowering traits. This integrated approach provides insights into the basic

evolutionary processes shaping flowering traits and contributes to our knowledge of how

selection operates on reproductive traits in plant populations.

Figure 1. Conceptual diagram illustrating how selection on flowering traits (flowering time,
duration, and total flowers) varies temporally across a breeding season and how cross-sectional
and longitudinal selection estimates relate. Each bell curve represents an individual plant's
flowering distribution. **(A)** When flowering traits are uncorrelated, selection pressures shift
throughout the season. The population's flowering distribution (left) shows distinct patterns
across three time intervals (grey bars). Selection gradients (circles) show direction and
magnitude of selection with their position relative to zero indicating selection for increases or
decreases in the trait value. Selection gradients measured at these cross-sectional time points
reveal how discrete episodes of selection contribute to overall longitudinal selection pattern
(right). Since pollen from each individual dominates in their corresponding interval (e.g. pollen
from early-flowering individuals in interval 1) this results in weak longitudinal selection when
averaged across the season. **(B)** When first flower timing correlates negatively with duration and
total flowers, early-flowering individuals also flower longer and produce more flowers. Thus, the

first to flower dominate the pollen pool in interval 1 and 2, but early, intermediate, and late
flowering individuals are all equally represented in the pollen pool during interval 3. This
correlation creates strong selection patterns across intervals 1 and 2, which drive strong
longitudinal selection on all three flowering traits.

Methods

Study species: *Brassica rapa*

Brassica rapa is a non-clonal annual in the Brassicaceae. The species is native to Eurasia but is naturalized throughout much of North America (Gulden et al., 2008). Flowers are typically perfect (containing both pollen-producing, and ovule-producing, structures within the same flower), and a plant begins flowering 30-60 days after seed germination (Austen & Weis, 2016a; Ison & Weis, 2017). The inflorescence is an elongated raceme, and flowers will persist for one to three days but senesce more quickly once fertilized. Flowering can continue through the fruit set of the earliest flower (Austen & Weis, 2016a). *B. rapa* is an obligate out-crosser with a sporophytic self-incompatibility system (Bateman, 1955; Murase et al., 2020). Insect pollinators include generalist bee and fly pollinators in the Apidae, Halictidae, and Syrphidae, and butterflies in the Lepidoptera.

Experimental design

Field populations

We established four perfect-flowered experimental populations of *B. rapa* at the University of Toronto's Koffler Scientific Reserve (44°1′N, 079°32′W) during summer 2011 using seeds bulk-collected from Eastern Quebec in 2009 (Austen & Weis, 2016a). Two populations flowered from July 4 – Aug 11 ("early season"), and two flowered from Aug 16 – Oct 13 ("late season"). The inclusion of these two flowering periods allowed us to capture any possible changes in pollinator service (or other ecological factors) across a larger temporal range.

The two populations within each of the early- and late-season treatments differed in spatial arrangement (one "clumped" and one "even"). Both spatial arrangements consisted of a hexagonal grid with 4 m sides. In the even populations, plants were evenly spaced at 0.5 m apart (211 total plants; **Fig. 2A**). The clumped populations had six small hexagons within a

larger hexagon, with small hexagons 2.7 m apart and plants within the small hexagons 0.17 m

apart (216 total plants; **Fig. 2B**). In total, our design included four distinct experimental

populations (i.e. even early season, clumped early season, even late season, clumped late

season).

Gene-trap implementation

To assess male fitness and pollen movement, male sterile plants ("gene-traps") were deployed

as standardized pollen recipients. These male-sterile plants were produced by back-crossing

Wisconsin FastPlants line 1-107 (carrying male sterility) into the Eastern Quebec population

background (see Ison & Weis 2017: "constructing the gene trap line"). This breeding approach

yielded functionally female plants that were morphologically and developmentally similar to the

perfect-flowered plants while maintaining male sterility.

Each population contained six gene-trap positions where male-sterile plants were placed. These

positions were evenly spaced in the even populations and positioned at the center of each small

hexagon in the clumped populations (**Fig. 2**). Gene-trap plants were systematically replaced

approximately 3-5 days throughout the flowering season (**Fig. 2** and **Table S1**). The temporal

rotation of new male-sterile plants as pollen recipients at each interval eliminated variance in

mate quality available to prospective sires across the season, ensuring our selection estimates

reflected differences in the competitive abilities of the perfect-flowered pollen plants as pollen

donors. Seeds produced by these standardized maternal plants were later genotyped to

estimate siring success of the perfect-flowered pollen donors through paternity analysis.

Phenotypic measurements and genetic analysis

We recorded three flowering traits for each pollen-donor (854 perfect-flowered plants): day of

first flowering (flowering time), flowering duration, and the total number of fresh flowers

(summed across census days). Prior to analysis, we standardized all phenotypes by z-scoring

them within each of the four populations.

We collected tissue samples from all perfect-flowered plants (potential sires) and each gene-

trap plant (264 maternal plants; 42 per early-season populations and 48 per late-season

populations). During winter 2011-12, we germinated and grew gene-trap offspring in a

greenhouse (see Ison and Weis 2017 for methods). For this study, we aimed for twelve

offspring per gene-trap plant, but some gene-trap plants produced fewer than twelve viable

seeds and for a few gene-trap plants, we sampled more than twelve offspring (mean 10.9; 1970

total offspring; **Table S1**). Offspring were tissue sampled when at least two true leaves were

present. Leaf tissue from parents and offspring was dried in silica gel.

Genotyping methods

We extracted DNA from the dried tissue samples using isopropanol precipitation methods

(Rogers et al., 1996). We characterized the plants' DNA at ten *B. rapa* microsatellite loci (Lowe

et al., 2004; Suwabe et al., 2002) (for primer information, see **Table S2**). We conducted 10 µl

polymerase chain reactions with 1 µl of DNA template, 5 µl of GoTaq Colorless MasterMix

(Promega), 0.5 µl of BSA, 0.25 µl of additional $MgCl_2$, 0.16 µl M13-tagged fluorescent dye (ABI

dye set-33), 0.05 µl of forward primer with M13 tail, 0.6 µl reverse primer, and DNase free water

to volume. Each PCR consisted of a 3 min 94°C denaturing cycle followed by 30-32 cycles of 30

s at 94°C, 30 s at annealing temperature (see **Table S2**), and a 30 s extension step at 72°C.

Three additional cycles were run with an annealing temperature of 55°C followed by a final 10

min 72°C extension step. We sent the PCR products to the University of Toronto's Center for

Analysis of Genome Evolution and Function for capillary gel fragment analysis. We scored the

resulting genotypes using Peak Scanner version 1.0 (Biosystems, 2006). To calculate

genotyping error, we repeated the PCR, fragment analysis on 96 haphazardly selected

samples. We calculated both the allelic dropout error rate, ε1, and the stochastic error rates due

to typing error, ε2, which can be attributed to mutation, miscalling, polymerase error, and data

entry error (J. Wang, 2004).

Paternity Analysis/ MCMC Model

We conducted analyses using R 4.4.0 (R Development Core Team 2024). We assigned

proportional paternity, reconstructed pedigrees, and checked for convergence using

MasterBayes and coda for analysis (Hadfield, 2017; Plummer, 2024). MasterBayes uses

Bayesian statistics to find the maximum likelihood of paternity through the calculation of a

posterior distribution relying on both genetic and phenotypic data to find the highest probability

offspring-paternal pair (Hadfield, 2017; Jones et al., 2010). The statistical inference employs a

Bayesian probabilistic architecture to simultaneously derive posterior probability distributions

across 1) genealogical relationships represented in the pedigree structure *P, and* 2) the

coefficient vector β that parametrizes the log-linear model incorporating non-genetic covariates.

This joint estimation process integrates the uncertainty in both the familial relationships and the

magnitude of non-genetic effects within a unified probabilistic framework. The non-genetic

information that is an input of the model comes out as the multinomial log-linear model. The

coefficients from the multinomial log-linear model are analogous to the coefficient of a multiple

regression, thus the coefficient derived on non-genetic data from paternity assignment of

offspring outputs a selection gradient as conducted in (Austen & Weis, 2016a) with the R

package MasterBayes (Hadfield et al., 2006).

To parameterize our model, we initialized three independent MCMC chains with differing

starting conditions. The first two chains were initialized with fixed genotyping error rates (*e1* and

e2) and allele frequencies, while the third chain incorporated normally distributed random

starting values for the beta coefficients, with a mean vector of [-0.1,0.1] and a standard

deviation of 0.05. We optimized MCMC sampling using a tuning parameter of 0.325 to achieve

acceptance rates between 20% and 50%. Each chain was run for 110,000 iterations with a

burn-in period of 10,000 iterations. To reduce autocorrelation, we employed a thinning interval

of 100, resulting in 1,000 store iterations per chain. The chains were run with a maximum

mismatch tolerance of 5. This multi-chain approach with different initialization parameters

allowed us to assess convergence and mixing of the MCMC sampling while accounting for

potential genotyping errors in the pedigree reconstruction. We tested all chains for convergence

using the Gelman-Rubin Diagnostic using the coda package (Plummer et al., 2006), and

removed all models from the analysis that did not reach convergence (convergence is indicated

by values lying between 0.9-1.1).

Selection analysis framework

Measuring longitudinal and cross-sectional selection

We estimate selection gradients (β) on our phenotypes for each interval (cross-section) and for

the entire season (longitudinal). Longitudinal selection gradients simply evaluated selection

based on all offspring generated within each of the four populations. For the cross-sectional

approach, we estimated selection in discrete intervals across the flowering season by

implementing the same analytical framework as with longitudinal selection, but we restricted the

pollen pool to specific windows when a single batch of gene-trap plants were present in the field

(7 intervals for early season and 8 intervals for late season; **Table S1**). Here, potential sires

were limited to plants producing pollen during the active gene-trap interval and thus plants that

were non-flowering were removed from the interval-specific analysis. To account for temporal

variation in pollen availability, we randomly subsampled offspring proportionally to each time

period's contribution to the total seasonal pollen pool. This approach generated interval-specific

selection gradients that revealed how selection pressures shifted throughout the flowering

season.

Comparing multivariate and univariate models

Our multivariate selection framework had incorporated all three flowering traits (flowering time,

flowering duration, and total flower number) simultaneously into the log-linear model, providing

an estimate of direct selection on the focal trait. This multivariate approach disentangles the

network of trait relationships, allowing us to isolate the direct selection acting on each trait while

statistically controlling for the influence of correlated traits. Thus, the multivariate selection

model partitions the shared variance of traits to reveal which traits are directly under selection

versus those that change due to correlated responses. However, we also measured selection

using a univariate model where only the trait of interest was incorporated into the log-linear

model, providing a selection differential and a measure of total selection. Selection differentials

capture the total selective pressures acting on each trait, including both direct selection on the

trait itself and indirect selection arising from correlations with other traits because covariation

with other traits is not included in the univariate model. This univariate selection model reflects

the net selection on each trait in natural conditions.

We identified traits experiencing selective conflicts by comparing the directionality of direct (β)

and total (S) selection estimates. Traits exhibiting opposing signs between β and S were

classified as being in "conflict zones," indicating that their evolutionary trajectory is constrained

by antagonistic selection pressures mediated through trait correlations (Ruffley et al., 2023).

The comparison reveals how trait covariation can modify the adaptive landscape and potentially

constrain evolutionary responses.

Distance of pollen movement analysis

The log-linear model framework allows for the incorporation of spatial effects through pairwise

distances between maternal plants and potential sires. The spatial effect coefficient (β) from the

posterior distribution quantifies the decay in mating probability with increasing distance and

follows an exponential decay function (Exponential decay = e^{β^*x}, where β is the peak of the

posterior distribution and x is the pairwise distance).

The spatial component was incorporated directly into the multinomial log-linear model with

MasterBayes. The exponential decay model provides a mechanistic understanding of pollen

movement where negative β values indicate decreasing probability of successful mating with

increasing distance between pairs of plants.

388
389
390

Figure 2. Diagrams of the experimental population spatial arrangements and flowering
phenology. Populations were hexagonal in shape and each side was 4 m long. Each population
had six gene-trap (male-sterile plants) positions denoted by an open circle **(A&B)**. Gene-trap
plants were added and removed from populations in consecutive batches every 3-5 days
throughout the experiment (7 batches for July-Aug time period, and 8 batches for Aug-Oct time
period, see **Table S1** for placement schedule). For the even spatial arrangement **(A)** plants
were spaced 0.5 m apart. In the clumped arrangement **(B)** there were six smaller hexagons at
the corner of each of the population hexagons. Within each smaller hexagon plants were 0.17 m
apart and smaller hexagons were 2.7 m from each other. Flower counts were recorded for all
plants on census days corresponding to each data point shown in the flowering phenology curve
(C&D). The even populations are denoted by circles and the clumped populations are denoted
by triangles. Dark blue and orange colors indicate the experiment was conducted in the early
season (July-Aug) and the light blue and yellow colors indicate the experiment was conducted in
the late season (Aug-Oct).

Results

We successfully genotyped nearly all populations (n=854), gene-trap (n=264), and offspring (n=1970) plants at the 10 microsatellite loci. There was 3.7%, 3.8%, and 2.5% of the genotypic data missing across loci, for population, gene-trap, and offspring plants, respectively. For the paternity analysis, we tested convergence of our three chains per analysis. In total, across direct and total selection and cross-sectional and longitudinal estimates in the four populations, we ran 242 chains, and 238 of them converged (98.3%). Analyses that did not reach convergence with the Gelman-Rubin Diagnostic were removed from the results.

Longitudinal selection on flowering traits via male fitness

Selection varied among populations and traits when measured across the entire flowering season. One population (clumped, late season) showed significant directional selection favoring late flowering, while the remaining populations did not show significant evidence of selection on flowering time (**Fig. 2A**). We observed significant selection for longer flowering duration in both late season populations but no significant selection in early season populations (**Fig. 2B**). Selection favored more total flowers in three populations, with only the early season clumped population having no significant selection (**Fig. 2C**).

Cross-sectional selection on flowering traits via male fitness

Cross-sectional selection analysis using the gene-trap sampling periods revealed dynamic patterns of selection that varied both temporally and spatially. All populations showed temporal shifts in selection on flowering time, generally favoring early flowering at the beginning of the season and late flowering towards the end, with non-significant estimates during mid-season (**Fig 2A**). Notably, cross-sectional selection gradients on flowering time had the largest estimated magnitudes in early and late flowering plants in both experiment timings in clumped populations (flowering time $\beta = -0.37 \pm 0.098$ through 0.56 ± 0.105, duration $\beta = -0.28 \pm 0.048$

through 0.48 ±0.112, total flowers β = -0.03 ± 0.0166 through 0.03 ± 0.009; **Fig. 2**). There were

two trend-breaking anomalies in the final gene-trap sampling period: the late season even

population had directional selection for early flowering (β = -0.07 ± 0.040), while the early

season clumped population had no significant selection.

Cross-sectional selection on flowering duration and total flowers varied based on the population

spatial arrangements. Both spatial arrangements had modest selection for longer flowering

duration in many of the gene-trap intervals (13 intervals) with strong selection in late flowering

plants (clumped late season β =0.48 ±0.112, **Fig. 2B**). The even populations had relatively

consistent selection for more total flowers throughout the season (mean β =0.15, max β =

0.022, min β = 0.012, **Fig. 2C**). In contrast, the clumped populations exhibited selection for

shorter flowering duration (β = -0.28 ± 0.074 and β = -0.11 ± 0.048) and fewer total flowers (β =

-0.05 ± 0.0165) during early and mid-season (**Fig. 2B&C, Table S3**).

Conflicts in total and direct selection

While longitudinal analyses revealed no clear conflicts between direct and total selection (**Fig.**

S1 and Table S3), cross-sectional analyses uncovered temporal variation in selection conflicts

(**Fig. 4 and Table S3**). The late season clumped population showed discordance between direct

selection favoring late flowering and no total selection on flowering time (β = 0.02 ± 0.015, S =

0.001 ± 0.0154, **Fig. S1A**). Measurements from early flowering plants in the late season even

population revealed conflict in both duration (β = 0.04 ± 0.016 S = -0.023 ± 0.019, **Fig. 4B**) and

total flowers (β = 0.01 ± 0.010, S = -0.02 ± 0.009 (**Fig. 4C**), where direct and total selection

opposed each other in sign. Additionally, gene-trap sample periods showed varying degrees of

conflict between direct and total selection, though not always reaching statistical significance.

Pollen movement in differing spatial arrangements

Pollen movement patterns varied with both spatial arrangement and gene-trap sampling

periods. Clumped populations generally showed greater pollen dispersal distances than even

populations across the season (**Fig. 5A** and **Table S4**). The gene-trap sample periods revealed

that the even populations had more variance in pollen movement distances with the shortest

pollen movement distance being during peak flowering time (β = -1.74 ± 0.232 in early season

and β = -1.104 ± 0.181 in late season), while clumped populations had a relatively consistent

distance of pollen movement throughout the season (**Fig. 5B, Table S4**).

485
Figure 3. Longitudinal and cross-sectional estimates for selection gradients (±SE) on **(A)**
flowering time, **(B)** duration of flowering, and **(C)** total flowers in four experimental populations.
Longitudinal selection gradients are depicted in the dashed box, while selection gradients in
each time interval correspond with the days since being placed in the field axis. Blue circles
correspond to evenly spaced populations and yellow and orange triangles represent clumped
populations, and the dark blue and dark orange represent the early season populations while
the light blue and yellow represent the late season populations. A filled-in shape indicates the
estimate and ±SE do not cross zero while an empty shape indicates the estimate and ±SE cross
zero and are therefore non-significant estimates.

496
497 **Figure 4.** Direct and total cross-sectional selection plotted for the three traits: **(A)** flowering time,
498 **(B)** duration, **(C)** total flowers. Points in grey conflict zones indicate opposing significant signs of
499 selection estimates. Even populations are shown as circles and clumped populations are shown
500 as triangles, and the color of the points indicates the time within the season.

520
Figure 5. (A) Probability of paternity relative to distance (m) for the two population types in the
early and late season experiments plotted as an exponential decay of the distance decay
coefficient. Shaded regions represent 95% Bayesian Confidence Intervals. Blue colors
represent even populations and yellow and orange colors represent clumped populations. **(B)**
Coefficient of distance decays over cross-sections of time and plotted without exponential decay
function.

Discussion

Cross-sectional selection analyses reveal mechanisms of longitudinal selection

in a plant with a long-breeding season

While longitudinal analyses capture season-wide trends, examining selection at a finer temporal

scale uncovered nuanced patterns that varied in both strength and direction of selection

throughout the breeding season (**Fig. 3**). The overall selection on flowering time was

characterized by mostly slow and steady weak selection, punctuated by bursts favoring early

flowering (**Fig. 3A**). These bursts were often matched or countered by weaker selection for later

flowering onset throughout the season. Early in the season, we observed strong directional

selection for early flowering, as the mating pool was dominated by early-flowering individuals

(**Fig. 3A**). At the beginning of the season there is limited variance in flowering start time

providing less trait variation for selection to act on. As such, we expect a small magnitude of

selection on early flowering– this makes the strong selection we found for early flowering

surprising and suggests that there is a true advantage to being one of the first pollen producing

plants, even among the early flowering cohort. This aligns in the literature with findings that the

earliest flowering plants can experience reduced competition for pollinators, potentially

increasing reproductive success (Elzinga et al., 2007; Kehrberger & Holzschuh, 2019).

Late in the season, we found strong directional selection for late flowering in all four

experimental populations, apart from the final interval for the late-season-even population. We

might expect that if early flowering plants have a long duration and continue contributing to the

gene pool at the end of the season, later flowering plants may have an advantage of higher

quality pollen at that time in the plant's flowering curve (Ashman & Schoen, 1994) or generally

have a genetic advantage at any point in the season (Austen & Weis, 2016a). Because we

found selection for late flowering plants at the end of the season and flowering time was only

negatively correlated in ~35% of the intervals with duration of flowering (~7% of intervals had

positive correlations with duration and flowering time and the other ~57% of intervals had no

correlation with flowering time and duration, **Table S5**), we propose that selection at the end of

the season is partially due to simply the ratio of pollen available at that time but also, in the case

of plants that negatively covary duration and flowering time, decreased flower quality in earlier

flowering plants at the end of the season. Overall, the alternating advantages of early and late

flowering through male function, combined with the likelihood that female function favors early

flowering (although cross-sectional selection on flowering time via female function has to date

not been evaluated empirically), suggest that there may be no net directional selection from the

integration of these fitness components across the flowering season (McGlothlin, 2010;

Wadgymar, Daws, et al., 2017; Wadgymar, Lowry, et al., 2017).

Our cross-sectional analysis revealed that longitudinal selection on flowering duration (which

was non-significant or in the direction of longer duration, (**Fig. 3B)**) and on total flower

production (which was non-significant or in the direction of more total flowers (**Fig. 3C**)) arose

through steady selection throughout the season. However, in certain intervals of the early

season clumped population, we observed selection favoring shorter duration and fewer total

flowers, contrasting with theoretical predictions that longer flowering and more flowers would

increase siring success, and thus, dampening overall season estimates (**Fig. 3B** and **C**). These

patterns were temporally restricted to early and peak season days, suggesting that temporal

density dependent selection was at play. At peak flowering, when there is significant overlap in

flowering schedules, competition for pollinators may lead to selection pressures favoring

resource conservation (Barley et al., 2022; Eisen et al., 2020). The emergence of negative or

neutral selection estimates highlight that the evolution of flowering traits is contingent upon

temporal dynamics and a single event can undermine or cancel out season-wide trends, which

underscores the complexity of selection operating in natural populations.

Within the cross-sectional analysis, we observed differences based on the spatial arrangements

of the populations, though we do not have statistical power to test these observations.

Interestingly, the strongest selection estimates for early and late flowering, shorter and longer

duration, and fewer and more total flowers all occur in clumped populations. Although even

population had more frequent significant selection estimates, clumped populations showed

amplified directional shifts. This pattern of stronger selection in denser spatial arrangements

aligns with studies demonstrating that pollinator behavior mediates density-dependent selection:

floral abundance alters visitation patterns, which disproportionately rewards extreme trait values

in dense groups (Caruso, 2002; Cohen et al., 2021; Kunin, 1993). For example, in *B. rapa*, high

density populations exhibit fivefold stronger soft selection (rank based competition (e.g. driven

by social competition)) than hard selection (environmentally based and independent of

population composition) (Weis et al., 2015), with elevated Gini coefficients reflecting stark

reproductive inequality– consistent with our observation of intensified selection in clumped

populations. These findings suggest that spatial density amplifies pollinator-driven selection by

increasing both competition for visits and variance in mating success. Future work directly

linking pollinator foraging decisions to selection gradients across spatial arrangements could

clarify how density shapes floral trait evolution in wild and managed populations.

Measuring longitudinal selection on flowering traits via male fitness enhances

our understanding of their evolution

Flowering time

While the timing of flowering is affected by environmental conditions, studies have found a

strong genetic basis for variance in flowering time (e.g. (Fox, 1990; Dorn & Mitchell-Olds, 1991;

Geber & Griffen, 2003; Reed et al., 2022)) including the Quebec populations of *B. rapa* used in

this study (independent narrow-sense heritability estimates of 0.506, 0.568, and 0.610 (Austen

& Weis, 2016a; Ison & Weis, 2017). Despite widespread evidence for directional selection

favoring early flowering via female fitness (reviewed in (Austen et al., 2017; Munguía-Rosas et

al., 2011)), flowering time variation persists in natural populations. A key hypothesis posits that

opposing selection through male fitness counterbalances female-driven selection, weakening

net directional selection (Austen et al., 2017). If the counterbalance is present, then our previous

understanding of the nature of selection on flowering time based on female fitness alone is

misleading (Austen et al., 2017; Austen & Weis, 2016b). In short, variance in flowering time

could be maintained because selection is not as strong when evaluating total fitness (the

combination of male and female reproductive success) as when using female fitness alone.

By isolating male siring success using gene-trap plants and holding female qualities nearly

constant through time, we observed no directional selection for early flowering in any population

and even detected selection for later flowering in one clumped populations (**Fig. 3A**). This

contrasts starkly with the strong early-flowering bias reported in female fitness studies (Austen

et al., 2017; Austen & Weis, 2016b). However, it is important to note that non-gene-trap plants'

female fitness (seed production) was unmeasured, limiting our ability to directly test the male-

female counterbalance hypothesis. While total fitness integrates both components, our results

demonstrate that male fitness alone does not reinforce early flowering, challenging assumptions

based solely on female function.

Sex-specific selection can arise because male fitness is limited by mating opportunities (pollen

dispersal), while female fitness is resource limited (ovule maturation) (Anthes et al., 2010;

Ashman & Morgan, 2004; Bateman & Innes, 1948; Burd & Callahan, 2000; Campbell,

1989).This divergence may explain a reduction net directional selection. However, even studies

measuring both male and female fitness remain incomplete without accounting for unmeasured

components (e.g. the correlation between flowering time genotype and seed variability or early-

life mortality) (Wadgymar, Daws, et al., 2017; Wadgymar et al., 2024; Wadgymar, Lowry, et al.,

2017). Thus, future studies should aim to integrate measurements of both pre- and post-

reproductive fitness as well as try to measure total fitness through male and female fitness

components to fully understand selective forces in plant populations.

Duration of flowering

Duration of flowering theoretically increases mating opportunities through male fitness by

extending the period of potential pollen transfer (Austen & Weis, 2016a; Lloyd, 1980; Xu, 2024).

However, longer duration could incur resource costs that create trade-offs generally (e.g., longer

duration but lower quality flowers) (Ashman & Schoen, 1994) or on the male-female fitness axis

(Ashman & Morgan, 2004; Zhao et al., 2020). In our experiment, the relationship between

duration and fitness varied between experimental times and spatial contexts– both even spatial

populations in early and late-season experimental runs showed selection for longer duration

while the early-season clumped population displayed selection for shorter duration in certain

intervals. These results suggest that selection on duration of flowering is not as intuitive as

theoretically proposed, and that ecological context and fitness type can impact selection in

unexpected ways.

Flowering duration presents a complex trait for selection analysis because of its poorly

understood heritability. While duration is often correlated with heritable traits like flowering time,

understanding of its own heritability is limited because there are few direct measurements in

selection studies (Bhakta et al., 2017; Cheng et al., 2006; Hof et al., 1999; Williams et al., 2022).

These studies have found moderate to low heritability of duration. Because heritability is

influenced by both genetic and environmental factors, the often observed correlation of

flowering time with flowering duration may be environmental not genetically based (Austen &

Weis, 2016b). In our populations, duration was correlated with total flowers and flowering time

some but not all of the time (**Table S5**). This absence of pattern suggests that selection on

duration may operate independently from selection on flowering time in this experiment.

Total flowers

Three of our four populations showed longitudinal directional selection for more total flowers,

and one population showed no longitudinal selection for more total flowers (**Fig. 3C**; note the

magnitude of selection is lower on this trait than in duration or flowering time). Similar to

flowering duration, heritability estimates of total flowers are varied and much less common than

estimates for flowering time (Drennan et al., 1986; Fogaça et al., 2012; Hof et al., 1999). In our

study, total flowers are correlated with flowering time in two populations, indicating potential

genetic or developmental constraints on independent evolution of these traits (**Table S5**). We

also suspect that the relationship between total flowers and male fitness is not straight forward–

although producing more flowers theoretically increases opportunity for pollen dispersal, plants

face resource allocation trade-offs between flower quantity and quality (Lanuza et al., 2023), as

well as variation in the differences temporal distribution of flower production throughout the

season (Burd, 1999).

Cross-sectional selection analyses reveal periods of conflict between direct and total

selection

Our cross-sectional analysis uncovered conflicts between total (S) and direct (β) selection

across flowering traits. These discrepancies– where total selection favored one trait value while

direction selection was neutral or opposing– suggest indirect selection mediated by trait

correlations (e.g., pleiotropy or linkage disequilibrium). For instance, in the late-season clumped

population, direct selection favored late flowering, while total selection was non-significant (**Fig.**

S1), implying that indirect selection via correlated traits (e.g. duration or flower number) could

constrain flowering time evolution. Similarly, in early-season even populations, total selection

favored fewer flowers and shorter duration, but direct selection opposed this, favoring longer

duration and more flowers. Such conflicts highlight how trait covariance can mask or redirect

evolutionary trajectories, particularly when selection gradients interact with genetic architecture.

Evolutionary outcomes depend critically on the interplay between selection gradients and the G-

matrix (genetic variance-covariance matrix). For example, in *Lobelia siphilitica*, multivariate

models revealed opposing selection on flower size and number due to genetic constraints,

limiting floral display evolution despite univariate selection signals (Caruso, 2004). Similarly,

foundational studies by Campbell (1996) and Mitchell et al. (1998) explored genetic constraints

of floral traits, highlighting how trait correlations can limit evolutionary responses. Trade-offs can

be evident in life history strategies as well, as in the well-documented trade-off in *Arabidopsis*

thaliana where flowering time and water use efficiency (two traits linked to different drought

survival strategies) negatively covary (Kenney et al., 2014) and were found to be constrained

genetically (Ruffley et al., 2023). These studies align with our findings: if traits like flowering time

and duration share genetic covariance (as suggested by their correlated selection), the G-matrix

could amplify or dampen responses to selection. However, the heritability of these traits– and

thus the strength of constraint– remains unknown in *B. rapa* and many other plant systems.

Future work estimating G-matrices alongside selection gradients in wild populations will clarify

how genetic architecture mediates floral traits evolution.

Pollen moved farther in clumped populations than in even populations

We designed the population arrangements such that detected pollen movement via paternity

assignment is likely an accurate representation of how pollen moved in the populations. We

eliminated potential fine-scale spatial genetic structure by randomly planting pollen donor plants

which enabled nearly all population plants to be compatible with all maternal plants (i.e., gene-

traps) since they have different genetic backgrounds. Notably, we observed striking differences

in successful pollen movement between even and clumped spatial arrangements, where overall

pollen moved farther in clumped than even populations (**Fig. 5A**). When looking at the intervals

of time in even populations, successful pollen moved shorter distances at peak flowering time

and showed more seasonal variation in pollen movement distance (**Fig. 5B**). This is consistent

with other studies that also found longer-distance pollen movements early and late in the

season (Ison et al., 2014; Kitamoto et al., 2006). In contrast, pollen moved farther in clumped

populations, despite closer plant proximity, and maintained consistent movement distances

across the season (**Fig. 5B**). We observed these patterns in the distance pollen moves in space

in both early and late season experimental runs indicating that these patterns are robust to

seasonal changes in pollinator community composition, though the pollinator community stayed

relatively consistent between the two runs (**Fig. S2**).

While our design kept the overall number of plants constant between even and clumped

populations, the arrangement of the plants and density immediately around the gene traps

varied (50 cm in even populations and 17 cm in clumped populations). Other studies have found

that pollinator foraging behavior changes with plant density (Cresswell, 2000; Grindeland et al.,

2005; Kunin, 1997). However, how these changes in foraging behavior impacts pollen

movement distance is less understood. In one study, bumblebees were more likely to skip over

plants and their foraging had less directionality when plant density increased but that did not

lead to changes in pollen movement distances, based on foraging observations (Cresswell,

1997). In our study, we hypothesize that the dense clumped populations of individual plants

were hard to distinguish, and pollinations may have been making foraging decisions at a clump-

level, rather than an individual-plant level. In support of this, we see an elevated probability of

paternity at 4 m in clumped populations **(Fig. 5A**), which is of note as 4 m is the distance

between neighboring clumps (**Fig 2B**)–this heightened 4 m-travel in clumped populations

suggests that pollinators may be treating clumps as a "single plant".

Conclusion and future directions

Our study demonstrates how multiple approaches for estimating selection can reveal complex

evolutionary dynamics that might otherwise remain hidden. The cross-sectional approach

reveals nuanced patterns of selection throughout the flowering season that were masked in

longitudinal estimates, highlighting the importance of temporal scales in selection studies and

displaying the mechanisms behind longitudinal selection estimates. By examining selection

through male fitness in *B. rapa*, we uncovered selection for late flowering in one of four

populations, which may help explain the maintenance of flowering time variation, despite

consistent selection for early flowering through female fitness in previous studies. While we

found limited evidence of conflicting selection through direct and total selection, this analytical

framework provides a valuable tool for identifying potential evolutionary constraints in traits with

stronger trade-offs or genetic relationships. Finally, spatial arrangement emerged as a key

factor influencing both selection strength and pollen movement patterns, demonstrating how

local aggregation can modify selection pressures.

Future studies should integrate both male and female fitness components to provide a

comprehensive view of selection acting on plants. Additionally, incorporating temporal sampling

designs will better capture how selection varies throughout the season, allowing for deeper

insight into the drivers of selection. Finally, measuring suites of potentially correlated traits offers

opportunities to evaluate selection conflicts and understand how genetic architecture influences

evolutionary outcomes. This multifaceted approach to studying selection will enhance our

understanding of how natural populations respond to selection pressures across different spatial

and temporal scales.

Data accessibility statement: *The data underlying this article are available in: (doi for dryad if*
manuscript is accepted). If the reviewers would like to see the code, please have the editor
reach out to the corresponding author.
785

786   **Author contributions:** JLI and AEW conceived and designed the experimental populations and
787   gene-trap method with input from KJB and EJA. JLI and KJB collected field and genotype data.
788   LL led the analysis with help from EJA, JLI, MAEP and AEW. LL wrote the initial draft of the
789   manuscript with contributions from JLI and AEW. All authors contributed to manuscript
790   revisions.
791

Funding: Initial funding was provided by grants from the Canada Foundation for Innovation, the
Natural Sciences and Engineering Research Council of Canada (RGPIN-2016-06540), and the
University of Toronto. Additional funding was provided through the College of Wooster's
Copeland, Luce, and Wilson Awards. While working on the manuscript JLI was supported by the
College of Wooster's research leave program and LL was supported by the National Science
Foundation Graduate Research Fellowship (Fellow ID 2020304890).
798

799   **Conflict of interest statement:** The authors of this manuscript declare no conflict of interest.
800

801   **Acknowledgments:** The authors thank BA, DC, JEI, JJ, GRS, YS, and GT for field and
802   greenhouse assistance. JLI also thanks Raven for companionship in the field. CG, GRS, YS,
803   GT, and JT, helped extensively with lab work, and we are grateful for their efforts. We thank
804   SMW for feedback on research design and ML, JP, KSM, RE, TB, ARP, and EVS, for feedback
805   on early drafts of the manuscript.

References

Ackerman, J. D. (2000). Abiotic pollen and pollination: ecological, functional, and evolutionary perspectives. In *Pollen and Pollination* (pp. 167–185). Springer Vienna. https://doi.org/10.1007/978-3-7091-6306-1_9

Anthes, N., David, P., Auld, J. R., Hoffer, J. N. A., Jarne, P., Koene, J. M., Kokko, H., Lorenzi, M. C., Pélissié, B., Sprenger, D., Staikou, A., & Schärer, L. (2010). Bateman gradients in hermaphrodites: an extended approach to quantify sexual selection. *The American Naturalist*, *176*(3), 249–263. https://doi.org/10.1086/655218

Ashman, T. L., & Morgan, M. T. (2004). Explaining phenotypic selection on plant attractive characters: male function, gender balance or ecological context? *Proceedings. Biological Sciences*, *271*(1539), 553–559. https://doi.org/10.1098/rspb.2003.2642

Ashman, T. L., & Schoen, D. J. (1994). How long should flowers live? *Nature*, *371*(6500), 788–791. https://doi.org/10.1038/371788a0

Augspurger, C. K. (1983). Phenology, flowering synchrony, and fruit set of six neotropical shrubs. *Biotropica*, *15*(4), 257. https://doi.org/10.2307/2387650

Austen, E. J., Forrest, J. R. K., & Weis, A. E. (2015). Within-plant variation in reproductive investment: consequences for selection on flowering time. *Journal of Evolutionary Biology*, *28*(1), 65–79. https://doi.org/10.1111/jeb.12538

Austen, E. J., Rowe, L., Stinchcombe, J. R., & Forrest, J. R. K. (2017). Explaining the apparent paradox of persistent selection for early flowering. *The New Phytologist*, *215*(3), 929–934. https://doi.org/10.1111/nph.14580

Austen, E. J., & Weis, A. E. (2016a). Estimating selection through male fitness: three complementary methods illuminate the nature and causes of selection on flowering time. *Proceedings. Biological Sciences / The Royal Society*, *283*(1825), 20152635. https://doi.org/10.1098/rspb.2015.2635

Austen, E. J., & Weis, A. E. (2016b). The causes of selection on flowering time through male

fitness in a hermaphroditic annual plant. *Evolution*, *70*(1), 111–125.

https://doi.org/10.1111/evo.12823

Barley, T. A., Martinez Algarin, M. G., & Bauer, J. T. (2022). The effects of flower patch density

on pollinator visitation. *Environmental Entomology*, *51*(2), 482–491.

https://doi.org/10.1093/ee/nvab143

Bateman, A. J., & Innes, J. (1948). Intra-sexual selection in Drosophila. *Heredity*, *2*, 349–368.

https://doi.org/10.1038/hdy.1948.21

Bateman, A. J. (1955). Self-incompatibility systems in angiosperms: III. Cruciferae. *Heredity*,

9(1), 53–68. https://doi.org/10.1038/hdy.1955.2

Bhakta, M. S., Gezan, S. A., Clavijo Michelangeli, J. A., Carvalho, M., Zhang, L., Jones, J. W.,

Boote, K. J., Correll, M. J., Beaver, J., Osorno, J. M., Colbert, R., Rao, I., Beebe, S.,

Gonzalez, A., Ricaurte, J., & Vallejos, C. E. (2017). A predictive model for time-to-flowering

in the common bean based on QTL and environmental variables. *G3: Genes, Genomes,*

Genetics, *7*(12), 3901–3912. https://doi.org/10.1534/g3.117.300229

Biosystems, A. (2006). *Peak Scanner^{TM} software version 1.0 reference guide*. Retrieved from

https://www.umassmed.edu/globalassets/deep-sequencing-

core/mbcl/peak_scanner_manual.pdf

Burd, & Callahan. (2000). What does the male function hypothesis claim? *Journal of*

Evolutionary Biology, *13*(5), 735–742. https://doi.org/10.1046/j.1420-9101.2000.00220.x

Burd, M. (1999). Flower number and floral components in ten angiosperm species: an

examination of assumptions about trade-offs in reproductive evolution. *Biological Journal of*

the Linnean Society. Linnean Society of London, *68*(4), 579–592.

https://doi.org/10.1111/j.1095-8312.1999.tb01189.x

Campbell, D. R. (1989). Measurements of selection in a hermaphroditic plant: Variation in male

and female pollination success. *Evolution; International Journal of Organic Evolution*, *43*(2),

318–334. https://doi.org/10.1111/j.1558-5646.1989.tb04230.x

Campbell, D. R. (1996). Evolution of floral traits in a hermaphroditic plant: Field measurements of heritabilities and genetic correlations. *Evolution; International Journal of Organic Evolution*, *50*(4), 1442–1453. https://doi.org/10.1111/j.1558-5646.1996.tb03918.x

CaraDonna, P. J., Petry, W. K., Brennan, R. M., Cunningham, J. L., Bronstein, J. L., Waser, N. M., & Sanders, N. J. (2017). Interaction rewiring and the rapid turnover of plant-pollinator networks. *Ecology Letters*, *20*(3), 385–394. https://doi.org/10.1111/ele.12740

Caruso, C. M. (2002). Influence of plant abundance on pollination and selection on floral traits of *Ipomopsis aggregata*. *Ecology*, *83*(1), 241–254. https://doi.org/10.1890/0012-9658(2002)083[0241:iopaop]2.0.co;2

Caruso, C. M. (2004). The quantitative genetics of floral trait variation in *Lobelia*: potential constraints on adaptive evolution. *Evolution; International Journal of Organic Evolution*, *58*(4), 732–740. https://doi.org/10.1111/j.0014-3820.2004.tb00406.x

Cheng, C. H., Seal, A. G., Murphy, S. J., & Lowe, R. G. (2006). Variability and inheritance of flowering time and duration in *Actinidia chinensis* (kiwifruit). *Euphytica; Netherlands Journal of Plant Breeding*, *147*(3), 395–402. https://doi.org/10.1007/s10681-005-9036-9

Cockburn, A., Osmond, H. L., & Double, M. C. (2008). Swingin' in the rain: condition dependence and sexual selection in a capricious world. *Proceedings. Biological Sciences*, *275*(1635), 605–612. https://doi.org/10.1098/rspb.2007.0916

Cohen, H., Philpott, S. M., Liere, H., Lin, B. B., & Jha, S. (2021). The relationship between pollinator community and pollination services is mediated by floral abundance in urban landscapes. *Urban Ecosystems*, *24*(2), 275–290. https://doi.org/10.1007/s11252-020-01024-z

Conner, J. K., Rush, S., Kercher, S., & Jennetten, P. (1996). Measurements of natural selection on floral traits in wild radish (*Raphanus raphanistrum*). Ii. Selection through lifetime male and total fitness. *Evolution; International Journal of Organic Evolution*, *50*(3), 1137–1146.

https://doi.org/10.1111/j.1558-5646.1996.tb02354.x

Cresswell, J. E. (1997). Spatial heterogeneity, pollinator behaviour and pollinator-mediated gene

flow: Bumblebee movements in variously aggregated rows of oil-seed rape. *Oikos*, *78*(3),

546. https://doi.org/10.2307/3545616

Cresswell, J. E. (2000). Manipulation of female architecture in flowers reveals a narrow optimum

for pollen deposition. *Ecology*, *81*(11), 3244–3249. https://doi.org/10.1890/0012-

9658(2000)081[3244:MOFAIF]2.0.CO;2

Daupagne, L., Bouchard, C., Michaud, A., Dhamelincourt, M., Lasne, E., & Tentelier, C. (2023).

Considering temporal variations in mating opportunities: consequences for sexual selection

estimates. *Animal Behaviour*, *204*, 49–65. https://doi.org/10.1016/j.anbehav.2023.08.004

Delph, L. F., & Ashman, T. L. (2006). Trait selection in flowering plants: how does sexual

selection contribute? *Integrative and Comparative Biology*, *46*(4), 465–472.

https://doi.org/10.1093/icb/icj038

Devaux, C., & Lande, R. (2010). Selection on variance in flowering time within and among

individuals. *Evolution; International Journal of Organic Evolution*, *64*(5), 1311–1320.

https://doi.org/10.1111/j.1558-5646.2009.00895.x

Dorn, L. A., & Mitchell-Olds, T. (1991). Genetics of Brassica campestris. 1. Genetic constraints

on evolution of life-history characters. *Evolution; International Journal of Organic Evolution*,

45(2), 371–379. https://doi.org/10.1111/j.1558-5646.1991.tb04411.x

Drennan, D., Harding, J., & Byrne, T. G. (1986). Heritability of inflorescence and floret traits in

G*erbera*. *Euphytica; Netherlands Journal of Plant Breeding*, *35*(1), 319–330.

https://doi.org/10.1007/bf00028570

Eisen, K. E., Wruck, A. C., & Geber, M. A. (2020). Floral density and co-occurring congeners

alter patterns of selection in annual plant communities. *Evolution; International Journal of*

Organic Evolution, *74*(8), 1682–1698. https://doi.org/10.1111/evo.13960

Elzinga, J. A., Atlan, A., Biere, A., Gigord, L., Weis, A. E., & Bernasconi, G. (2007). Time after

time: flowering phenology and biotic interactions. *Trends in Ecology & Evolution*, 22(8),

432–439. https://doi.org/10.1016/j.tree.2007.05.006

Endler, J. A. (1986). *Natural selection in the wild* (Vol. 21). Princeton University Press.

Feldman, T. S. (2008). The plot thickens: does low density affect visitation and reproductive

success in a perennial herb, and are these effects altered in the presence of a co-flowering

species? *Oecologia*, 156(4), 807–817. https://doi.org/10.1007/s00442-008-1033-y

Fisogni, A., Rossi, M., Sgolastra, F., Bortolotti, L., Bogo, G., de Manincor, N., Quaranta, M., &

Galloni, M. (2016). Seasonal and annual variations in the pollination efficiency of a

pollinator community of *Dictamnus albus* L. *Plant Biology*, 18(3), 445–454.

https://doi.org/10.1111/plb.12417

Flanagan, R. J., Mitchell, R. J., & Karron, J. D. (2011). Effects of multiple competitors for

pollination on bumblebee foraging patterns and *Mimulus ringens* reproductive success.

Oikos, 120(2), 200–207. https://doi.org/10.1111/j.1600-0706.2010.18777.x

Fogaça, L. A., Oliveira, R. A., Cuquel, F. L., Filho, J. C. B., Vendrame, W. A., & Tombolato, A.

F. C. (2012). Heritability and genetic correlation in daylily selection. *Euphytica; Netherlands*

Journal of Plant Breeding, 184(3), 301–310. https://doi.org/10.1007/s10681-011-0478-y

Fogarty, N. D., Vollmer, S. V., & Levitan, D. R. (2012). Weak prezygotic isolating mechanisms in

threatened Caribbean *Acropora* corals. *PloS One*, 7(2), e30486.

https://doi.org/10.1371/journal.pone.0030486

Fox, G. (2003). Assortative mating and plant phenology: Evolutionary and practical

consequences. *Evolutionary Ecology Research*, 5, 1–18.

Fox, G. A. (1990). Components of flowering time variation in a desert annual. *Evolution;*

International Journal of Organic Evolution, 44(6), 1404–1423.

https://doi.org/10.1111/j.1558-5646.1990.tb03835.x

Geber, M. A., & Griffen, L. R. (2003). Inheritance and Natural Selection on Functional Traits.

International Journal of Plant Sciences, 164(S3), S21–S42. https://doi.org/10.1086/368233

Grindeland, J. M., Sletvold, N., & Ims, R. A. (2005). Effects of floral display size and plant

density on pollinator visitation rate in a natural population of *Digitalis purpurea*. *Functional*

Ecology, *19*(3), 383–390. https://doi.org/10.1111/j.1365-2435.2005.00988.x

Gulden, R. H., Warwick, S. I., & Thomas, A. G. (2008). The Biology of Canadian Weeds. 137.

Brassica napus L. and *B. rapa* L. *Canadian Journal of Plant Science*, *88*(5), 951–996.

https://doi.org/10.4141/cjps07203

Gyulavári, H. A., Tüzün, N., Arambourou, H., Therry, L., Dévai, G., & Stoks, R. (2017). Within-

season variation in sexual selection on flight performance and flight-related traits in a

damselfly. *Evolutionary Ecology*, *31*(1), 21–36. https://doi.org/10.1007/s10682-016-9882-z

Hadfield, J. D. (2017). *MasterBayes: ML and MCMC methods for pedigree reconstruction and*

analysis (Version 2.55).

Hadfield, J. D., Richardson, D. S., & Burke, T. (2006). Towards unbiased parentage

assignment: combining genetic, behavioural and spatial data in a Bayesian framework.

Molecular Ecology, *15*(12), 3715–3730. https://doi.org/10.1111/j.1365-294X.2006.03050.x

Hoekstra, H. E., Hoekstra, J. M., Berrigan, D., Vignieri, S. N., Hoang, A., Hill, C. E., Beerli, P., &

Kingsolver, J. G. (2001). Strength and tempo of directional selection in the wild.

Proceedings of the National Academy of Sciences of the United States of America, *98*(16),

9157–9160. https://doi.org/10.1073/pnas.161281098

Hof, L., Keizer, L. C. P., Elberse, I. A. M., & Dolstra, O. (1999). A model describing the flowering

of single plants, and the heritability of flowering traits of *Dimorphotheca pluvialis*. *Euphytica;*

Netherlands Journal of Plant Breeding, *110*(1), 35–44.

https://doi.org/10.1023/a:1003701700116

Hou, M., Opedal, Ø. H., & Zhao, Z.-G. (2024). Sexually concordant selection on floral traits

despite greater opportunity for selection through male fitness. *The New Phytologist*, *241*(2),

926–936. https://doi.org/10.1111/nph.19370

Ison, J. L., Prescott, L. J., Nordstrom, S. W., Waananen, A., & Wagenius, S. (2018). Pollinator-

973 mediated mechanisms for increased reproductive success in early flowering plants. *Oikos* ,

974  *127*(11), 1657–1669. https://doi.org/10.1111/oik.04882

975 Ison, J. L., Wagenius, S., Reitz, D., & Ashley, M. V. (2014). Mating between *Echinacea*

976  *angustifolia* (Asteraceae) individuals increases with their flowering synchrony and spatial

977  proximity. *American Journal of Botany*, *101*(1), 180–189.

978  https://doi.org/10.3732/ajb.1300065

979 Ison, J. L., & Weis, A. E. (2017). Temporal population genetic structure in the pollen pool for

980  flowering time: A field experiment with *Brassica rapa* (Brassicaceae). *American Journal of*

981  *Botany*, *104*(10), 1569–1580. https://doi.org/10.3732/ajb.1700210

982 Jones, A. G., Small, C. M., Paczolt, K. A., & Ratterman, N. L. (2010). A practical guide to

983  methods of parentage analysis. *Molecular Ecology Resources*, *10*(1), 6–30.

984  https://doi.org/10.1111/j.1755-0998.2009.02778.x

985 Kasumovic, M. M., Bruce, M. J., Andrade, M. C. B., & Herberstein, M. E. (2008). Spatial and

986  temporal demographic variation drives within-season fluctuations in sexual selection.

987  *Evolution; International Journal of Organic Evolution*, *62*(9), 2316–2325.

988  https://doi.org/10.1111/j.1558-5646.2008.00446.x

989 Kehrberger, S., & Holzschuh, A. (2019). How does timing of flowering affect competition for

990  pollinators, flower visitation and seed set in an early spring grassland plant? *Scientific*

991  *Reports*, *9*(1), 15593. https://doi.org/10.1038/s41598-019-51916-0

992 Kenney, A. M., McKay, J. K., Richards, J. H., & Juenger, T. E. (2014). Direct and indirect

993  selection on flowering time, water-use efficiency (WUE, δ (13)C), and WUE plasticity to

994  drought in Arabidopsis thaliana. *Ecology and Evolution*, *4*(23), 4505–4521.

995  https://doi.org/10.1002/ece3.1270

996 Kingsolver, J. G., Hoekstra, H. E., Hoekstra, J. M., Berrigan, D., Vignieri, S. N., Hill, C. E.,

997  Hoang, A., Gibert, P., & Beerli, P. (2001). The strength of phenotypic selection in natural

998  populations. *The American Naturalist*, *157*(3), 245–261. https://doi.org/10.1086/319193

Kitamoto, N., Ueno, S., Takenaka, A., Tsumura, Y., Washitani, I., & Ohsawa, R. (2006). Effect

of flowering phenology on pollen flow distance and the consequences for spatial genetic

structure within a population of *Primula sieboldii* (Primulaceae). *American Journal of*

Botany, *93*(2), 226–233. https://doi.org/10.3732/ajb.93.2.226

Kunin, W. E. (1993). Sex and the single mustard: Population density and pollinator behavior

effects on seed-set. *Ecology*, *74*(7), 2145–2160. https://doi.org/10.2307/1940859

Kunin, W. E. (1997). Population size and density effects in pollination: Pollinator foraging and

plant reproductive success in experimental arrays of *Brassica kaber*. *The Journal of*

Ecology, *85*(2), 225. https://doi.org/10.2307/2960653

Landry, C. L. (2013). Pollinator-mediated competition between two co-flowering Neotropical

mangrove species, *Avicennia germinans* (Avicenniaceae) and *Laguncularia racemosa*

(Combretaceae). *Annals of Botany*, *111*(2), 207–214. https://doi.org/10.1093/aob/mcs265

Lanuza, J. B., Rader, R., Stavert, J., Kendall, L. K., Saunders, M. E., & Bartomeus, I. (2023).

Covariation among reproductive traits in flowering plants shapes their interactions with

pollinators. *Functional Ecology*, *37*(7), 2072–2084. https://doi.org/10.1111/1365-

2435.14340

Levin, D. A., & Anderson, W. W. (1970). Competition for pollinators between simultaneously

flowering species. *The American Naturalist*, *104*(939), 455–467.

https://doi.org/10.1086/282680

Lloyd, D. G. (1980). Sexual strategies in plants. I. an hypothesis of serial adjustment of maternal

investment during one reproductive session. *The New Phytologist*, *86*(1), 69–79.

https://www.jstor.org/stable/2434416?seq=1

Losos, J. B., Schoener, T. W., Langerhans, R. B., & Spiller, D. A. (2006). Rapid temporal

reversal in predator-driven natural selection. *Science*, *314*(5802), 1111.

https://doi.org/10.1126/science.1133584

Lowe, A. J., Moule, C., Trick, M., & Edwards, K. J. (2004). Efficient large-scale development of

1025 microsatellites for marker and mapping applications in *Brassica* crop species. *Theoretical*

1026 *and Applied Genetics*, *108*(6), 1103–1112. https://doi.org/10.1007/s00122-003-1522-7

1027 Makino, T. T., Ohashi, K., & Sakai, S. (2007). How do floral display size and the density of

1028 surrounding flowers influence the likelihood of bumble bee revisitation to a plant?

1029 *Functional Ecology*, *21*(1), 87–95. https://doi.org/10.1111/j.1365-2435.2006.01211.x

1030 McDiarmid, R. W. (1994). Amphibian Diversity and Natural History: An Overview. In R. W.

1031 Heyer (Ed.), *Measuring and Monitoring Biological Diversity: Standard Methods for*

1032 *Amphibians* (p. 4726). Smithsonian Institution Press.

1033 McGlothlin, J. W. (2010). Combining selective episodes to estimate lifetime nonlinear selection.

1034 *Evolution; International Journal of Organic Evolution*, *64*(5), 1377–1385.

1035 https://doi.org/10.1111/j.1558-5646.2009.00914.x

1036 Mitchell, R. J., Shaw, R. G., & Waser, N. M. (1998). Pollinator selection, quantitative genetics,

1037 and predicted evolutionary responses of floral traits in *Penstemon centranthifolius*

1038 (Scrophulariaceae). *International Journal of Plant Sciences*, *159*(2), 331–337.

1039 https://doi.org/10.1086/297554

1040 Munguía-Rosas, M. A., Ollerton, J., Parra-Tabla, V., & De-Nova, J. A. (2011). Meta-analysis of

1041 phenotypic selection on flowering phenology suggests that early flowering plants are

1042 favoured. *Ecology Letters*, *14*(5), 511–521. https://doi.org/10.1111/j.1461-

1043 0248.2011.01601.x

1044 Murase, K., Moriwaki, Y., Mori, T., Liu, X., Masaka, C., Takada, Y., Maesaki, R., Mishima, M.,

1045 Fujii, S., Hirano, Y., Kawabe, Z., Nagata, K., Terada, T., Suzuki, G., Watanabe, M.,

1046 Shimizu, K., Hakoshima, T., & Takayama, S. (2020). Mechanism of self/nonself-

1047 discrimination in *Brassica* self-incompatibility. *Nature Communications*, *11*(1), 4916.

1048 https://doi.org/10.1038/s41467-020-18698-w

1049 Parachnowitsch, A. L., & Caruso, C. M. (2008). Predispersal seed herbivores, not pollinators,

1050 exert selection on floral traits via female fitness. *Ecology*, *89*(7), 1802–1810.

https://doi.org/10.1890/07-0555.1

Parachnowitsch, A. L., Caruso, C. M., Campbell, S. A., & Kessler, A. (2012). *Lobelia siphilitica*

plants that escape herbivory in time also have reduced latex production. *PloS One*, *7*(5),

e37745. https://doi.org/10.1371/journal.pone.0037745

Plummer, M. (2024). *Output Analysis and Diagnostics for MCMC*. https://CRAN.R-

project.org/package=coda

Plummer, M., Best, N., Cowles, K., & Vines, K. (2006). *CODA: convergence diagnosis and*

output analysis for MCMC (pp. 7–11). https://CRAN.R-project.org/package=coda

Polatto, L. P., Chaud-Netto, J., & Alves-Junior, V. V. (2014). Influence of abiotic factors and

floral resource availability on daily foraging activity of bees: influence of abiotic and biotic

factors on bees. *Journal of Insect Behavior*, *27*(5), 593–612.

https://doi.org/10.1007/s10905-014-9452-6

Punzalan, D., Rodd, F. H., & Rowe, L. (2010). Temporally variable multivariate sexual selection

on sexually dimorphic traits in a wild insect population. *The American Naturalist*, *175*(4),

401–414. https://doi.org/10.1086/650719

Reed, W. J., Ison, J. L., Waananen, A., Shaw, F. H., Wagenius, S., & Shaw, R. G. (2022).

Genetic variation in reproductive timing in a long-lived herbaceous perennial. *American*

Journal of Botany, *109*(11), 1861–1874. https://doi.org/10.1002/ajb2.16072

Richardson, L. K., Gallagher, M. K., Hayes, T. E., Gallinat, A. S., Kiefer, G., Manion, K., Jenkins,

M., Diersen, G., & Wagenius, S. (2021). Competition for pollination and isolation from

mates differentially impact four stages of pollination in a model grassland perennial. *The*

Journal of Ecology, *109*(3), 1356–1369. https://doi.org/10.1111/1365-2745.13562

Rogers, H. J., Burns, N. A., & Parkes, H. C. (1996). Comparison of small-scale methods for the

rapid extraction of plant DNA suitable for PCR analysis. *Plant Molecular Biology Reporter*,

14(2), 170–183. https://doi.org/10.1007/bf02684906

Ruffley, M., Lutz, U., Leventhal, L., Hateley, S., Yuan, W., Keck, J., Rhee, S. Y., Weigel, D., &

Exposito-Alonso, M. (2023). Selection constraints of plant adaptation can be relaxed by

gene editing. In *bioRxiv* (p. 2023.10. 16.562583).

https://doi.org/10.1101/2023.10.16.562583

Schreiber, E. A., & Burger, J. (2001). *Biology of Marine Birds*. CRC Press.

Suwabe, K., Iketani, H., Nunome, T., Kage, T., & Hirai, M. (2002). Isolation and characterization

of microsatellites in *Brassica rapa* L. *Theoretical and Applied Genetics*, *104*(6-7), 1092–

1098. https://doi.org/10.1007/s00122-002-0875-7

Thomson, J. D. (1980). Skewed flowering distributions and pollinator attraction. *Ecology*, *61*(3),

572–579. https://doi.org/10.2307/1937423

Tsuji, K., Kobayashi, K., Hasegawa, E., & Yoshimura, J. (2020). Dimorphic flowers modify the

visitation order of pollinators from male to female flowers. *Scientific Reports*, *10*(1), 9965.

https://doi.org/10.1038/s41598-020-66525-5

Urquhart, F. A. (1960). *The Monarch butterfly*. Heritage.

Wacker, S., Amundsen, T., Forsgren, E., & Mobley, K. B. (2014). Within-season variation in

sexual selection in a fish with dynamic sex roles. *Molecular Ecology*, *23*(14), 3587–3599.

https://doi.org/10.1111/mec.12826

Wadgymar, S. M., Daws, S. C., & Anderson, J. T. (2017). Integrating viability and fecundity

selection to illuminate the adaptive nature of genetic clines: stabilizing selection on

functional traits. *Evolution Letters*, *1*(1), 26–39. https://doi.org/10.1002/evl3.3

Wadgymar, S. M., Lowry, D. B., Gould, B. A., Byron, C. N., Mactavish, R. M., & Anderson, J. T.

(2017). Identifying targets and agents of selection: innovative methods to evaluate the

processes that contribute to local adaptation. *Methods in Ecology and Evolution*, *8*(6), 738–

749. https://doi.org/10.1111/2041-210X.12777

Wadgymar, S. M., Sheth, S., Josephs, E., DeMarche, M., & Anderson, J. (2024). Defining

Fitness in Evolutionary Ecology. *International Journal of Plant Sciences*, 000–000.

https://doi.org/10.1086/729360

Walsh, B., & Blows, M. W. (2009). Abundant genetic variation + strong selection = multivariate

genetic constraints: A geometric view of adaptation. *Annual Review of Ecology, Evolution,*

and Systematics, *40*(1), 41–59. https://doi.org/10.1146/annurev.ecolsys.110308.120232

Wang, H., Barrett, S. C. H., Li, X.-Y., Niu, Y., Duan, Y.-W., Zhang, Z.-Q., & Li, Q.-J. (2021).

Sexual conflict in protandrous flowers and the evolution of gynodioecy. *Evolution;*

International Journal of Organic Evolution, *75*(2), 278–293.

https://doi.org/10.1111/evo.14113

Wang, J. (2004). Sibship reconstruction from genetic data with typing errors. *Genetics*, *166*(4),

1963–1979. https://doi.org/10.1534/genetics.166.4.1963

Weis, A. E., & Kossler, T. M. (2004). Genetic variation in flowering time induces phenological

assortative mating: quantitative genetic methods applied to *Brassica rapa*. *American*

Journal of Botany, *91*(6), 825–836. https://doi.org/10.3732/ajb.91.6.825

Weis, A. E., Turner, K. M., Petro, B., Austen, E. J., & Wadgymar, S. M. (2015). Hard and soft

selection on phenology through seasonal shifts in the general and social environments: A

study on plant emergence time. *Evolution; International Journal of Organic Evolution*, *69*(6),

1361–1374. https://doi.org/10.1111/evo.12677

Weis, A. E., Winterer, J., Vacher, C., Kossler, T. M., Young, C., & LeBuhn, G. (2005).

Phenological assortative mating in flowering plants: the nature and consequences of its

frequency dependence. *Evolutionary Ecology Research*, *7*, 161–181.

Williams, O., Vander Schoor, J. K., Butler, J. B., Ridge, S., Sussmilch, F. C., Hecht, V. F. G., &

Weller, J. L. (2022). The genetic architecture of flowering time changes in pea from wild to

crop. *Journal of Experimental Botany*, *73*(12), 3978–3990.

https://doi.org/10.1093/jxb/erac132

Xu, K. (2024). Evolution of flowering time due to variation in the onset of pollen dispersal among

individuals. *Evolution; International Journal of Organic Evolution*, *78*(3), 401–412.

https://doi.org/10.1093/evolut/qpad215

Zhao, Z., Hou, M., Wang, Y., & Du, G. (2020). Phenological variation of flower longevity and

duration of sex phases in a protandrous alpine plant: potential causes and fitness

significance. *BMC Plant Biology*, *20*(1), 137. https://doi.org/10.1186/s12870-020-02356-w

Supplementary Figures

Figure S1. Overall selection conflict for three focal traits. Direct and total longitudinal selection plotted for the three traits (±SE): **(A)** flowering time, **(B)** duration, **(C)** total flowers. Points in grey conflict zones indicate opposing significant signs of selection estimates. Even populations are indicated by the blue points, and clumped populations are indicated by yellow and orange points. Dark blue and dark orange points are estimates from the early season run and light blue and yellow points are from the late season run.

1189
Figure S2. Pollinator counts by pollinator category for each of the four plots (BMB = bumblebee,
BSY = big syrphid fly, BUT = butterfly, HSF = house fly, HYB = honey bee, LBB = large black
bee, MGB = medium green bee, MSY = medium syrphid fly, SBB = small black bee, SGB =
small green bee, SSY = small syrphid fly, UBB = unknown black bee, USY = unknown syrphid
fly, WAS = wasp). Pollinator counts were obtained when we conducted six five-minute pollinator
observation periods every four to eight days at each of the four plots. Each observation period
included a gene-trap plant and the six closest plot plant positions. We recorded all insects that
visited a reproductive structure on one of the observed plants. We did not collect the pollinators,
so as to not impact the pollinator community.

Supplementary Tables

Table S1. Information about the gene-trap plant batches for early and late season populations. Batches are gene-trap plants that were placed into the experimental plots on the first day of the 'gene-trap time period' and removed on the last day. Approximately twelve offspring were genotyped from each of the six gene-trap plants from each batch.

Gene-trap batch	Gene-trap time period	Gene-trap offspring genotyped (even plot)	Gene-trap offspring genotyped (clumped plot)
Early season (1.1)	4-7 July	72	67
Early season (1.2)	11-14 July	71	39
Early season (1.3)	14-21 July	48	77
Early season (1.4)	21-25 July	55	51
Early season (1.5)	25-28 July	68	49
Early season (1.6)	28-July-1 August	69	60
Early season (1.7)	1-11 August	70	30
Late season (2.1)	16-20 August	72	71
Late season (2.2)	24-28 August	73	71
Late season (2.3)	1-5 September	72	72
Late season (2.4)	9-13 September	70	72
Late season (2.5)	17-21 September	71	72
Late season (2.6)	25-29 September	72	71
Late season (2.7)	3-7 October	71	72
Late season (2.8)	7-13 October	71	71

1222 **Table S2.** Microsatellite primer information.

Locus	F primer sequence	R primer sequence	Repeat motif	Annealing temperature °C	ε1	ε2	Primer source
BRMS-005	ACCTCCTGCAGATTCGTGTC	GCTGACCTTTCTTACCGCTC	$(GA)_{13}$	60	0.052449	0.013805	Suwabe et al., 2002
BRMS-007	AAATTGTTTCTCTTCCCCAT	GTGTTAGGGAGCTGGAGAAT	$(CT)_{24}$	45	0.022207	0.047789	Suwabe et al., 2002
BRMS-019	CCCAAACGCTTTTGACACAT	GGCACAATCCACTCAGCTTT	$(GT)_{10}$	60	0.063575	0.006244	Suwabe et al., 2002
BRMS-033	GCGGAAACGAACACTCCTCCCATGT	CCTCCTTGTGCTTTCCCTGGAGACG	$(CA)_{11}$	60	0.010682	0.006696	Suwabe et al., 2002
BRMS-037	CTGCTCGCATTTTTTATCATAC	TACGCTTGGGAGAGAAAACTAT	$(CA)_{10}$	56	0.024997	0.03898	Suwabe et al., 2002
BRMS-042-2	GGATCAGTTATCTGCACCACAA	TCGGAATTGGATAAGAATTCAA	$(GA)_4(CT)_{26}$	60	0.024431	0.004467	Suwabe et al., 2002
Na10-A08	CATGGTTAAAACAATGGCCC	CAAGAAACACCATCATTTCTCA	$(GA)_{21}$	58	0.013953	0.002916	Lowe et al., 2004

Locus	F primer sequence	R primer sequence	Repeat motif	Annealing temperature °C	ε1	ε2	Primer source
Na10-D09	AAGAACGTCAAGATCCTCTGC	ACCACCACGGTAGTAGAGCG	$(GA)_{24}$	61	0.036447	0.005979	Lowe et al., 2004
Na12-E02	TTGAAGTAGTTGGAGTAATTGGAGG	CAGCAGCCACAACCTTACG	$(TTG)_{13}$	55	0.007782	0.008447	Lowe et al., 2004
Ra2-A05	GCTAGTTTACGCGGCGG	AAACGACATCGGCAAAGAAG	$(GGC)_{6}$	50	0.009597	0.006421	Lowe et al., 2004

1223
1224 **Table SX**. Microsatellite primer information.

1250 **Table S3.** Tables providing mean, standard deviation, naive standard error, and batch and plot
1251 identifiers for both direct and total selection estimates in longitudinal and cross-sectional
1252 analyses for the three phenotypes flowering time, duration of flowering, and total flowers.
1253

Direct selection estimates (flowering time)				
Mean	**SD**	**Naive.SE**	**batch_id**	**Plot**
-0.006981909	0.028531554	0.000902247	Overall	Early season clumped
0.015903295	0.015035075	0.000475451	Overall	Late season clumped
-0.01608615	0.021168351	0.000669402	Overall	Early season even
0.00332643	0.017943455	0.000567422	Overall	Late season even
-0.375596626	0.098439972	0.003112945	1.1	Early season clumped
0.025871576	0.068861019	0.002177577	1.2	Early season clumped
-0.002016799	0.038472298	0.001216601	1.3	Early season clumped
-0.003970546	0.05747295	0.001817454	1.4	Early season clumped
0.000380185	0.047107278	0.001489663	1.5	Early season clumped
0.146935381	0.048386808	0.001530125	1.6	Early season clumped
0.089617636	0.111987284	0.003541349	1.7	Early season clumped
-8.151834252	6.138518812	0.194117009	2.1	Late season clumped
-0.158231295	0.03924648	0.001241083	2.2	Late season clumped
-0.046718881	0.024993799	0.000790373	2.3	Late season clumped
0.048687282	0.026372259	0.000833964	2.4	Late season clumped
0.102996043	0.026697738	0.000844257	2.5	Late season clumped
0.071973383	0.024355838	0.000770199	2.6	Late season clumped
0.17962825	0.039616412	0.001252781	2.7	Late season clumped
0.56566521	0.105095761	0.00332342	2.8	Late season clumped
-0.242616427	0.080214745	0.002536613	1.1	Early season even
-0.199546906	0.047097617	0.001489357	1.2	Early season even
-0.055462348	0.042598907	0.001347096	1.3	Early season even
0.008392488	0.037678651	0.001191504	1.4	Early season even
0.019336944	0.030348842	0.000959715	1.5	Early season even

0.079555724	0.030462853	0.00096332	1.6	Early season even
0.169496117	0.035322689	0.001117001	1.7	Early season even
-16.74500507	6.94164946	0.21951423	2.1	Late season even
-0.206475354	0.036369289	0.001150098	2.2	Late season even
0.014245518	0.026199672	0.000828506	2.3	Late season even
-0.002361999	0.027747698	0.000877459	2.4	Late season even
0.079928258	0.030323221	0.000958904	2.5	Late season even
0.229932993	0.043773556	0.001384241	2.6	Late season even
0.083331634	0.055573654	0.001757393	2.7	Late season even
-0.062178147	0.060055837	0.001899132	2.8	Late season even

Direct selection estimates (duration)				
Mean	SD	Naive.SE	batch_id	Plot
0.00882696	0.02500533	0.00079074	Overall	Early season clumped
0.02652461	0.01102161	0.00034853	Overall	Late season clumped
0.01132943	0.018089	0.00057202	Overall	Early season even
0.01154386	0.00985931	0.00031178	Overall	Late season even
-0.2815545	0.07362382	0.00232819	1.1	Early season clumped
0.00762596	0.0587306	0.00185722	1.2	Early season clumped
-0.099965	0.04848925	0.00153336	1.3	Early season clumped
-0.0343674	0.04663158	0.00147462	1.4	Early season clumped
-0.002068	0.04242457	0.00134158	1.5	Early season clumped
0.15472886	0.02680204	0.00084755	1.6	Early season clumped
0.2080262	0.07628213	0.00241225	1.7	Early season clumped
0.0628585	0.08120299	0.00256786	2.1	Late season clumped
0.03028139	0.01491263	0.00047158	2.2	Late season clumped
-0.0223375	0.01999688	0.00063236	2.3	Late season clumped
0.0227997	0.0184814	0.00058443	2.4	Late season clumped
0.04676684	0.01831795	0.00057926	2.5	Late season clumped
0.10846698	0.01892437	0.00059844	2.6	Late season clumped
0.16481722	0.0334763	0.00105861	2.7	Late season clumped
0.48793196	0.11279371	0.00356685	2.8	Late season clumped
0.05026352	0.03764932	0.00119058	1.1	Early season even
-0.0182263	0.03497106	0.00110588	1.2	Early season even
0.02385683	0.03355934	0.00106124	1.3	Early season even
0.00985617	0.03240137	0.00102462	1.4	Early season even
0.04074949	0.03014128	0.00095315	1.5	Early season even
0.09346784	0.02964106	0.00093733	1.6	Early season even
0.1798256	0.03475848	0.00109916	1.7	Early season even

0.04241331	0.01660166	0.00052499	2.1	Late season even
0.0113336	0.01784886	0.00056443	2.2	Late season even
0.00450894	0.01532011	0.00048446	2.3	Late season even
0.01477747	0.01447289	0.00045767	2.4	Late season even
-0.010332	0.01765665	0.00055835	2.5	Late season even
-0.01189	0.02315772	0.00073231	2.6	Late season even
0.01259347	0.02192172	0.00069323	2.7	Late season even
0.03622488	0.02324671	0.00073513	2.8	Late season even

Direct selection estimates (total flowers)				
Mean	SD	Naive.SE	batch_id	Plot
0.00510838	0.00605554	0.00019149	Overall	Early season clumped
0.00523285	0.00331455	0.00010482	Overall	Late season clumped
0.00680454	0.00286065	9.05E-05	Overall	Early season even
0.01631161	0.00294164	9.30E-05	Overall	Late season even
-0.0309894	0.01658643	0.00052451	1.1	Early season clumped
-0.0123147	0.01359962	0.00043006	1.2	Early season clumped
0.00475529	0.00986638	0.000312	1.3	Early season clumped
0.03381944	0.00944064	0.00029854	1.4	Early season clumped
0.01759863	0.00893975	0.0002827	1.5	Early season clumped
0.00182576	0.00931115	0.00029444	1.6	Early season clumped
0.02604876	0.01752567	0.00055421	1.7	Early season clumped
0.00953891	0.00984681	0.00031138	2.1	Late season clumped
0.00477074	0.00490168	0.00015501	2.2	Late season clumped
0.0094166	0.00581673	0.00018394	2.3	Late season clumped
0.00568217	0.00592814	0.00018746	2.4	Late season clumped
0.02042128	0.00482642	0.00015263	2.5	Late season clumped
0.01194299	0.00501812	0.00015869	2.6	Late season clumped
-0.0009671	0.00562092	0.00017775	2.7	Late season clumped
0.01230085	0.0047433	0.00015	2.8	Late season clumped
-0.0013418	0.0061301	0.00019385	1.1	Early season even
-0.003449	0.00570091	0.00018028	1.2	Early season even
0.00547881	0.0058345	0.0001845	1.3	Early season even
0.01221124	0.00466883	0.00014764	1.4	Early season even
0.01302366	0.00354156	0.00011199	1.5	Early season even
0.01435457	0.00303726	9.60E-05	1.6	Early season even
0.00379067	0.00459429	0.00014528	1.7	Early season even

0.01485659	0.01048946	0.00033171	2.1	Late season even
0.01799558	0.00803002	0.00025393	2.2	Late season even
-0.0004183	0.00578559	0.00018296	2.3	Late season even
0.01462476	0.00516053	0.00016319	2.4	Late season even
0.01467727	0.00472552	0.00014943	2.5	Late season even
0.01156951	0.00505552	0.00015987	2.6	Late season even
0.01390219	0.0041293	0.00013058	2.7	Late season even
0.02279289	0.00347257	0.00010981	2.8	Late season even

Total selection estimates (indirect + direct) (flowering time)				
Mean	SD	Naive.SE	batch_id	Plot
-0.006245153	0.027145249	0.000858408	Overall	Early season clumped
0.000502176	0.015385134	0.000486521	Overall	Late season clumped
-0.017027673	0.021456522	0.000678515	Overall	Early season even
0.000755943	0.017899217	0.000566023	Overall	Late season even
-0.116477424	0.075047471	0.002373209	1.1	Early season clumped
0.061961986	0.063664127	0.002013236	1.2	Early season clumped
0.016895209	0.037927546	0.001199374	1.3	Early season clumped
0.047557717	0.050022058	0.001581836	1.4	Early season clumped
-0.006949878	0.037874006	0.001197681	1.5	Early season clumped
0.052937902	0.035993165	0.001138204	1.6	Early season clumped
-0.118205263	0.060980513	0.001928373	1.7	Early season clumped
-1.206140182	0.316204997	0.00999928	2.1	Late season clumped
-0.13720504	0.036369516	0.001150105	2.2	Late season clumped
-0.056717202	0.024387774	0.000771209	2.3	Late season clumped
0.027278464	0.024642487	0.000779264	2.4	Late season clumped
0.079986311	0.029759593	0.000941081	2.5	Late season clumped
-0.006080636	0.02331375	0.000737246	2.6	Late season clumped
0.046692645	0.02598787	0.000821809	2.7	Late season clumped
0.212382724	0.049964516	0.001580017	2.8	Late season clumped
-0.237546966	0.080386225	0.002542036	1.1	Early season even
-0.136520557	0.046521306	0.001471133	1.2	Early season even
-0.041817789	0.044178334	0.001397042	1.3	Early season even
0.017976659	0.039324229	0.001243541	1.4	Early season even
-0.00075705	0.029611497	0.000936398	1.5	Early season even
0.030183431	0.026266082	0.000830606	1.6	Early season even
0.027862096	0.01488759	0.000470787	1.7	Early season even

-8.387812769	5.225045121	0.165230435	2.1	Late season even
-0.178649154	0.034457995	0.001089657	2.2	Late season even
0.019579195	0.026755947	0.000846097	2.3	Late season even
-0.004850016	0.026920949	0.000851315	2.4	Late season even
0.076617595	0.030610104	0.000967976	2.5	Late season even
0.231011633	0.043631702	0.001379756	2.6	Late season even
0.074558702	0.041999864	0.001328152	2.7	Late season even
-0.073420909	0.040465859	0.001279643	2.8	Late season even

	Total selection estimates (indirect + direct) (duration)			
Mean	**SD**	**Naive.SE**	**batch_id**	**Plot**
0.00888715	0.02414921	0.00076366	Overall	Early season clumped
0.03611464	0.00922804	0.00029182	Overall	Late season clumped
0.02701842	0.01631411	0.0005159	Overall	Early season even
0.0075893	0.00924095	0.00029222	Overall	Late season even
-0.1645653	0.05422724	0.00171482	1.1	Early season clumped
-0.0682685	0.06698963	0.0021184	1.2	Early season clumped
-0.1030512	0.04537273	0.00143481	1.3	Early season clumped
-0.0204566	0.04660923	0.00147391	1.4	Early season clumped
0.01016517	0.03782852	0.00119624	1.5	Early season clumped
0.12356042	0.02360052	0.00074631	1.6	Early season clumped
0.15860709	0.03610699	0.0011418	1.7	Early season clumped
-0.1768025	0.04110923	0.00129999	2.1	Late season clumped
0.04090976	0.01280845	0.00040504	2.2	Late season clumped
-0.0119696	0.01729475	0.00054691	2.3	Late season clumped
0.0342684	0.01641246	0.00051901	2.4	Late season clumped
0.08307472	0.01483972	0.00046927	2.5	Late season clumped
0.09911	0.01459202	0.00046144	2.6	Late season clumped
0.05598788	0.01618212	0.00051172	2.7	Late season clumped
0.01832311	0.0185168	0.00058555	2.8	Late season clumped
0.03103956	0.03212403	0.00101585	1.1	Early season even
-0.0243458	0.03432544	0.00108547	1.2	Early season even
0.02939477	0.03282804	0.00103811	1.3	Early season even
0.01956398	0.03261906	0.00103151	1.4	Early season even
0.04881801	0.02628494	0.0008312	1.5	Early season even
0.05201642	0.02455114	0.00077638	1.6	Early season even
0.05122485	0.02216105	0.00070079	1.7	Early season even

-0.0228835	0.01942158	0.00061416	2.1	Late season even
0.02265316	0.0178791	0.00056539	2.2	Late season even
0.00307507	0.01677021	0.00053032	2.3	Late season even
0.01056311	0.0145205	0.00045918	2.4	Late season even
-0.013963	0.01668804	0.00052772	2.5	Late season even
-0.0367376	0.02208735	0.00069846	2.6	Late season even
-0.0291383	0.0208311	0.00065874	2.7	Late season even
-0.0041043	0.02183622	0.00069052	2.8	Late season even

Total selection estimates (indirect + direct) (total flowers)				
Mean	SD	Naive.SE	batch_id	Plot
0.00591626	0.00594727	0.00018807	Overall	Early season clumped
0.00901461	0.00274265	8.67E-05	Overall	Late season clumped
0.00535489	0.00261246	8.26E-05	Overall	Early season even
0.01442986	0.00286848	9.07E-05	Overall	Late season even
-0.0257099	0.01352754	0.00042778	1.1	Early season clumped
-0.0135148	0.01398149	0.00044213	1.2	Early season clumped
-0.0050195	0.00973243	0.00030777	1.3	Early season clumped
0.02894547	0.00901032	0.00028493	1.4	Early season clumped
0.01443122	0.00875107	0.00027673	1.5	Early season clumped
0.00321208	0.00784956	0.00024823	1.6	Early season clumped
0.01296748	0.00929735	0.00029401	1.7	Early season clumped
-0.0300788	0.00686928	0.00021723	2.1	Late season clumped
0.01255315	0.00444451	0.00014055	2.2	Late season clumped
0.00624272	0.00506644	0.00016022	2.3	Late season clumped
0.00758897	0.0045251	0.0001431	2.4	Late season clumped
0.01863653	0.00397083	0.00012557	2.5	Late season clumped
0.01657257	0.00410618	0.00012985	2.6	Late season clumped
0.00404817	0.00493065	0.00015592	2.7	Late season clumped
0.00401624	0.0047312	0.00014961	2.8	Late season clumped
0.00478586	0.00600942	0.00019004	1.1	Early season even
-0.0038417	0.00619016	0.00019575	1.2	Early season even
0.00336287	0.00575715	0.00018206	1.3	Early season even
0.00909441	0.00458465	0.00014498	1.4	Early season even
0.01144983	0.00339332	0.00010731	1.5	Early season even
0.01060982	0.00290363	9.18E-05	1.6	Early season even
-0.0072674	0.00398486	0.00012601	1.7	Early season even

-0.0229094	0.00861401	0.0002724	2.1	Late season even
0.00479196	0.006677	0.00021115	2.2	Late season even
-0.0001432	0.00580432	0.00018355	2.3	Late season even
0.01322361	0.00520336	0.00016455	2.4	Late season even
0.01482225	0.00488533	0.00015449	2.5	Late season even
0.00748422	0.00538035	0.00017014	2.6	Late season even
0.01119549	0.00374181	0.00011833	2.7	Late season even
0.02101552	0.00337827	0.00010683	2.8	Late season even

1446 **Table S4.** Distance decay coefficients overall and by cross-section, standard deviation (SD), naive
1447 standard error (Naive.SE), batch unique identifier, and plot identification.
1448

Mean	SD	Naive.SE	Time.series.SE	batch_id	Plot
-0.317837465	0.044689822	0.001413216	0.001413216	Overall	Early season clumped
-0.510805082	0.074624019	0.002359819	0.002359819	1.1	Early season clumped
-0.322985353	0.101972742	0.003224661	0.003224661	1.2	Early season clumped
-0.268203257	0.066243369	0.002094799	0.002094799	1.3	Early season clumped
-0.507300335	0.108578235	0.003433545	0.003590788	1.4	Early season clumped
-0.263970904	0.06997566	0.002212825	0.002212825	1.5	Early season clumped
-0.096257626	0.074744094	0.002363616	0.002317706	1.6	Early season clumped
-0.427581789	0.116611641	0.003687584	0.003687584	1.7	Early season clumped

Mean	SD	Naive.SE	Time.series.SE	batch_id	Plot
-0.341095528	0.036489575	0.001153902	0.001210716	Overall	Late season clumped
-0.308492813	0.066002057	0.002087168	0.002087168	2.1	Late season clumped
-0.392679726	0.070532759	0.002230442	0.002230442	2.2	Late season clumped
-0.636592274	0.09309287	0.002943855	0.002943855	2.3	Late season clumped
-0.454286413	0.07547378	0.00238669	0.002252077	2.4	Late season clumped
-0.327838819	0.061945003	0.001958873	0.001958873	2.5	Late season clumped
-0.351887491	0.057437166	0.001816323	0.001816323	2.6	Late season clumped
-0.373146439	0.063896589	0.002020588	0.002020588	2.7	Late season clumped
-0.436147964	0.067142277	0.002123225	0.002268596	2.8	Late season clumped

Mean	SD	Naive.SE	Time.series.SE	batch_id	Plot
-0.797697697	0.085003163	0.002688036	0.002688036	Overall	Early season even
-0.552820264	0.13505356	0.004270769	0.004270769	1.1	Early season even
-1.744651908	0.232391077	0.007348851	0.008646116	1.2	Early season even
-1.161193606	0.247805254	0.00783629	0.00783629	1.3	Early season even

Mean	SD	Naive.SE	Time.series.SE	batch_id	Plot
-0.98966918	0.169600111	0.005363226	0.005363226	1.4	Early season even
-0.547981383	0.120386754	0.003806963	0.003461145	1.5	Early season even
-0.52810649	0.11458752	0.003623576	0.003623576	1.6	Early season even
-0.509307717	0.120664344	0.003815742	0.003815742	1.7	Early season even

Mean	SD	Naive.SE	Time.series.SE	batch_id	Plot
-0.618860274	0.071623246	0.002264926	0.002264926	Overall	Late season even
-0.413540403	0.099965021	0.003161172	0.003161172	2.1	Late season even
-0.357433907	0.118675583	0.003752851	0.003752851	2.2	Late season even
-1.103726098	0.181458818	0.005738232	0.005738232	2.3	Late season even
-0.638494874	0.116795807	0.003693408	0.003877345	2.4	Late season even
-0.391249216	0.114403781	0.003617765	0.003414186	2.5	Late season even
-0.607766055	0.121406375	0.003839207	0.003839207	2.6	Late season even
-0.453230443	0.115335253	0.003647221	0.003617791	2.7	Late season even
-0.23110786	0.119146619	0.003767747	0.003767747	2.8	Late season even

Table S5. Pearson correlation coefficients for all variables overall and by cross-section, p-value, batch id indicating the interval, and the population type. .

variable 1	variable 2	correlation coefficient (r)	p-value	batch_id	population
flowering time	duration	-0.108796	0.1168646	overall	early season even
flowering time	total flowers	-0.1694515	0.01417473	overall	early season even
duration	total flowers	0.30625275	6.49E-06	overall	early season even
flowering time	duration	-0.122673	0.07197802	overall	early season clumped
flowering time	total flowers	-0.0005354	0.99375777	overall	early season clumped
duration	total flowers	0.32026702	1.53E-06	overall	early season clumped
flowering time	duration	0.05469244	0.42933741	overall	late season even
flowering time	total flowers	0.24278174	0.00037219	overall	late season even
duration	total flowers	-0.0032296	0.96280442	overall	late season even
flowering time	duration	0.2238422	0.00092353	overall	late season clumped
flowering time	total flowers	-0.0882211	0.19650397	overall	late season clumped
duration	total flowers	0.52273278	0	overall	late season clumped
flowering time	duration	0.03124778	0.73914524	1.1	early season even
flowering time	total flowers	-0.1059658	0.25758493	1.1	early season even
duration	total flowers	0.16656088	0.07393866	1.1	early season even
flowering time	duration	0.02166525	0.7691269	1.2	early season even
flowering time	total flowers	-0.1097976	0.13573603	1.2	early season even
duration	total flowers	0.25179106	0.0005266	1.2	early season even
flowering time	duration	-0.0036244	9.59E-01	1.3	early season even
flowering time	total flowers	-0.0903504	0.19873559	1.3	early season even
duration	total flowers	0.28134582	4.57E-05	1.3	early season even
flowering time	duration	-0.087612	0.29811014	1.4	early season even
flowering time	total flowers	0.00420594	9.60E-01	1.4	early season even
duration	total flowers	0.20330619	0.01487775	1.4	early season even
flowering time	duration	-0.3014413	0.0029919	1.5	early season even
flowering time	total flowers	-0.1012531	0.32889555	1.5	early season even
duration	total flowers	0.14085603	0.17334982	1.5	early season even
flowering time	duration	-0.4644519	0.00024002	1.6	early season even

flowering time	total flowers		-0.0818464	0.54134498	1.6 early season even
duration	total flowers		0.03724077	0.78136586	1.6 early season even
flowering time	duration		-0.758817	8.19E-08	1.7 early season even
flowering time	total flowers		-0.2486999	0.14356848	1.7 early season even
duration	total flowers		0.08920431	0.60489775	1.7 early season even
flowering time	duration		-0.1789834	0.04407399	1.1 early season clumped
flowering time	total flowers		-0.1261353	0.15763571	1.1 early season clumped
duration	total flowers		0.33315217	0.00012959	1.1 early season clumped
flowering time	duration		-0.1899769	0.00586691	1.2 early season clumped
flowering time	total flowers		0.03126849	0.65311268	1.2 early season clumped
duration	total flowers		0.35542285	1.29E-07	1.2 early season clumped
flowering time	duration		-0.122673	0.07197802	1.3 early season clumped
flowering time	total flowers		-0.0005354	0.99375777	1.3 early season clumped
duration	total flowers		0.32026702	1.53E-06	1.3 early season clumped
flowering time	duration		-0.3259546	7.54E-05	1.4 early season clumped
flowering time	total flowers		-0.0281776	0.73923031	1.4 early season clumped
duration	total flowers		0.19870562	0.01775868	1.4 early season clumped
flowering time	duration		-0.4302741	0.00130086	1.5 early season clumped
flowering time	total flowers		-0.0836393	0.55155679	1.5 early season clumped
duration	total flowers		0.03612217	0.7973436	1.5 early season clumped
flowering time	duration		-0.2583463	0.13401365	1.6 early season clumped
flowering time	total flowers		-0.1303629	0.45541263	1.6 early season clumped
duration	total flowers		0.28688426	0.09473694	1.6 early season clumped
flowering time	duration		-0.4060562	0.06778209	1.7 early season clumped
flowering time	total flowers		-0.239039	0.2966868	1.7 early season clumped
duration	total flowers		0.30017967	0.18613492	1.7 early season clumped
flowering time	duration		0.386939	0.01135958	2.1 late season clumped
flowering time	total flowers		0.42047224	0.00556198	2.1 late season clumped
duration	total flowers		0.52932245	0.00031292	2.1 late season clumped
flowering time	duration		0.03237528	0.71248374	2.2 late season clumped
flowering time	total flowers		-0.0977349	0.26490164	2.2 late season clumped

duration	total flowers	0.6087358	9.77E-15	2.2	late season clumped
flowering time	duration	0.13939391	0.04957618	2.3	late season clumped
flowering time	total flowers	-0.0824127	0.24718247	2.3	late season clumped
duration	total flowers	0.53765172	2.22E-16	2.3	late season clumped
flowering time	duration	0.03546327	0.65011499	2.4	late season clumped
flowering time	total flowers	-0.3046599	6.58E-05	2.4	late season clumped
duration	total flowers	0.45083624	1.09E-09	2.4	late season clumped
flowering time	duration	-0.0519579	0.54355023	2.5	late season clumped
flowering time	total flowers	-0.3754992	5.24E-06	2.5	late season clumped
duration	total flowers	0.38947149	2.15E-06	2.5	late season clumped
flowering time	duration	-0.2656026	0.0134502	2.6	late season clumped
flowering time	total flowers	-0.3492361	0.00098172	2.6	late season clumped
duration	total flowers	0.39767881	0.00014969	2.6	late season clumped
flowering time	duration	-0.4220204	0.0092751	2.7	late season clumped
flowering time	total flowers	-0.2963079	0.07495489	2.7	late season clumped
duration	total flowers	0.43483869	0.00715556	2.7	late season clumped
flowering time	duration	-0.4763618	0.01038847	2.8	late season clumped
flowering time	total flowers	-0.3063095	0.11289407	2.8	late season clumped
duration	total flowers	0.43593564	0.02039783	2.8	late season clumped
flowering time	duration	0.16585614	0.30003689	2.1	late season even
flowering time	total flowers	0.29738475	0.05898921	2.1	late season even
duration	total flowers	0.21022224	0.1870857	2.1	late season even
flowering time	duration	0.01634887	0.85409787	2.2	late season even
flowering time	total flowers	0.2036342	0.02063286	2.2	late season even
duration	total flowers	0.06001309	0.49929938	2.2	late season even
flowering time	duration	0.0886604	0.22382051	2.3	late season even
flowering time	total flowers	0.23537003	0.00107931	2.3	late season even
duration	total flowers	0.01783815	0.80701488	2.3	late season even
flowering time	duration	0.02431246	0.76775072	2.4	late season even
flowering time	total flowers	0.05644605	0.49265999	2.4	late season even
duration	total flowers	-0.0307412	0.70882124	2.4	late season even

flowering time	duration	0.001234	0.98892637	2.5	late season even
flowering time	total flowers	-0.0085442	0.92344068	2.5	late season even
duration	total flowers	-0.0445466	0.61617654	2.5	late season even
flowering time	duration	0.09496412	0.37057185	2.6	late season even
flowering time	total flowers	-0.0242579	0.81945937	2.6	late season even
duration	total flowers	-0.0453514	0.6694783	2.6	late season even
flowering time	duration	-0.0632892	0.70979335	2.7	late season even
flowering time	total flowers	0.14199825	0.40183076	2.7	late season even
duration	total flowers	-0.1818482	0.28139682	2.7	late season even
flowering time	duration	0.04286	0.81892107	2.8	late season even
flowering time	total flowers	-0.0679444	0.71647574	2.8	late season even
duration	total flowers	-0.3198476	0.07941965	2.8	late season even

1475
1476
1477