An AI-Guided Framework Reveals Conserved Features Governing

microRNA Strand Selection

Author list

Dalton Meadows[1,2],

Hailee Hargis[1,2],

Amanda Ellis[1,2],

Heewook Lee[1],

Marco Mangone[1,*]

Author affiliations

[1)] The Biodesign Institute at Arizona State University, 1001 S McAllister Ave, Tempe, AZ

[2)] School of Life Sciences, Arizona State University, 427 E Tyler Mall, Tempe, AZ

Author list footnotes

[*)] Corresponding author

Corresponding author and e-mail address

Marco Mangone; mangone@asu.edu

Abstract

MicroRNAs (miRNAs) are central regulators of gene expression, yet how cells choose between the two strands (5p or 3p) of a miRNA duplex during biogenesis remains unresolved. Here, we present a comprehensive, experimentally grounded framework that decodes the logic of miRNA strand selection. Using *Caenorhabditis elegans* as a model system, we developed a high-throughput qPCR platform enabling precise quantification of strand usage across developmental stages and in specific somatic tissues. To uncover the molecular grammar guiding this process, we built a predictive machine learning model trained on experimentally validated strand usage data. This AI-driven model, integrating 77 biologically informed features, accurately predicts strand preference not only in nematodes but also across vertebrates, including humans, revealing compositional and structural biases that are conserved yet functionally repurposed. Our analysis shows that strand selection is not stochastic but follows conserved, context-dependent rules shaped by cellular and developmental cues. To support the research community, we provide open-access resources: a database of strand usage profiles, predictive scores across species, and code and protocols via GitHub. This work offers the first unified, generalizable model for miRNA strand selection, establishing a paradigm that combines large-scale experimentation with AI to reveal a conserved, programmable layer of gene regulation.

Introduction

miRNAs are ~22-nucleotide non-coding RNAs that guide the RNA-induced silencing

complex (RISC) to target mRNAs, primarily through base pairing within the 3′

untranslated regions (3′UTRs) (1). Once bound, the miRNA–RISC complex (miRISC)

typically represses translation of the target mRNA. Both miRNAs and their 3′UTR

targets are often conserved across metazoans and play key roles in essential biological

processes such as development and morphogenesis.

miRNAs have undergone several pronounced expansions in metazoan genomes (2),

coinciding with evolving organismal complexity yet rarely corresponding with increases

in protein-coding genes (2). Eukaryotic genomes contain 100-2,000 distinct mature

miRNAs in intergenic regions, introns, and in clusters. Following transcription in long

primary miRNA transcripts (pri-miRNAs) by the RNA polymerase II, they are processed

in the nucleus by the microprocessor complex into shorter hairpin structures known as

precursor miRNAs (pre-miRNAs) (1). The two strands composing the pre-miRNAs are

named the 5p and the 3p strand, respectively. Pre-mRNAs are subsequently exported

to the cytoplasm by Exportin-5, where they undergo further cleavage by the enzyme

Dicer, producing mature miRNA duplexes (1).

The Hsc70-Hsp90 chaperone complex facilitates the ATP-dependent loading of the

miRNA duplex into Argonaute, the protein that executes the silencing step. This

complex is thought to induce conformational changes in Argonaute that enable duplex

loading (3-5). Once loaded, one strand of this duplex, the guide strand, is kept within the

RISC, while the other, the passenger strand, is typically degraded. Both the 5p and the

3p strands can be selected for miRISC loading and act as guide strands. It is currently

unclear how this process, termed 'miRNA strand selection,' is executed in detail, which

proteins are involved in this decision, and how the guide strand is ultimately chosen

between the two strands. The predominant factors influencing miRNA strand selection

are believed to be the specific 5' nucleotide sequences of each strand and the

comparative thermodynamic stability exhibited by the two ends of the miRNA duplex (6).

Studies performed using the human Ago2 protein, an Argonaute protein member of the

RISC complex, show that the pocket responsible for binding the 5' nucleotide of the

guide strand exhibits a strong preference for Uracil nucleotides (7). This aligns with the

prevalent occurrence of Uracil as the 5' nucleotide in miRNA guide strands (2), whereas

the passenger strand commonly initiates with a Cytosine at its 5' end.

Furthermore, the miRISC complex exhibits a preference for loading the end of the

miRNA duplex with lower thermodynamic stability, although not always (8). The initial

four nucleotides on each end of the duplex are believed to establish this thermodynamic

asymmetry, and in some cases, a slight difference in the number of hydrogen bonds

can induce the preferential loading of the less stable strand *in vitro* (5).

Consistently, the 5' ends of guide strands are enriched with purines (2). In

comparison, the 5' ends of passenger strands are predominantly pyrimidine-rich in both

human and fly miRNAs, contributing to the thermodynamic asymmetry observed

between the two ends of the duplex. In *C. elegans*, single amino acid substitutions

within the MID (G553R) and PIWI (G716E, S895F, and S939F) domains of the

Argonaute protein ALG-1 invert the strand selection process for specific miRNAs,

suggesting a potential role for this protein in the miRNA strand selection process (9).

This indicates that while these mutations play a pivotal role, they do not consistently

reverse strand selection for all miRNAs, suggesting that there may be additional internal

or external factors involved in this process. Tissue-specific accessory factors and their

dosage could also be responsible for miRNA strand selection, altering strand choices

based on tissue or developmental contexts, similarly to the role of the SR proteins in

RNA splicing (10). While thermodynamic and sequence features help explain strand

choice, a comprehensive mechanistic framework has been lacking.

C. elegans is an ideal model for studying miRNA strand selection due to its compact

3′UTRome (11-14), small number of miRNAs (miRbase), fully mapped cell lineage (15),

and well-characterized transcriptome. Here, we developed a high-throughput qPCR

platform to systematically profile miRNA strand selection events across *C. elegans*

development and in somatic tissues. These data revealed conserved sequence and

structural patterns, which we used to train a machine learning model that predicts strand

selection with 83% accuracy and generalizes to human miRNAs.

Materials and methods

All DI water was obtained from either a Purelab Option system (ELGA LabWater) or

an Avidity Science system (Duo) and autoclaved before use. Chemicals were ordered

from Sigma Aldrich/Merck, VWR, or Thermo Fisher Scientific in molecular biology grade

and stored according to company instructions. Gateway™ LR Clonase II Plus™ enzyme

mixes, SuperScript™ III Reverse Transcriptase, and PowerTrack™ SYBR Green Master

Mix were purchased from Invitrogen. Primers for the HiTmiSS assay were ordered from

Life Technologies. Poly(U) Polymerase and restriction enzymes used in validation were

ordered from New England Biolabs. The pCFJ150 plasmid was a gift from Erik

Jorgensen. The OP50-1 *E. coli*, N2 *C. elegans* and EG6699 *C. elegans* strains were

purchased through the Caenorhabditis Genetics Center.

Maintenance of *C. elegans* strains

All *C. elegans* strains were cultured on NGM plates seeded with OP50-1 bacteria. All

transgenic lines produced for experiments described in this study were maintained at

24°C. When bacteria on each NGM plate had been cleared, *C. elegans* strains were

transferred onto new plates via washing into a 15 ml Falcon tube using 15 ml M9 media

(22 mM KH_2PO_4, 42 mM Na_2HPO_4, 85.5 mM NaCl, 1 mM $MgSO_4$, sterilized by

autoclave) and pelleted for 3 minutes at 400*g*. The M9 media was then aspirated to a

volume of 500 ml, and the animals were resuspended via gentle pipetting before being

moved onto fresh, seeded NGM plates.

Synchronization of *C. elegans* strains

The EG6699 strain (16) was maintained at 24°C. To synchronize animals for

injection, confluent plates were bleached to isolate the embryos. Each plate was

washed into a 15 ml Falcon tube with 15 ml DI water and pelleted for 3 minutes at 400*g*.

The water was then aspirated, and the animals were incubated in 5 ml of bleaching

solution (0.5M NaOH, 10% bleach) for 20 minutes with gentle end-over-end rotation.

After bleaching, the embryos were washed twice with DI water and moved onto fresh,

seeded NGM plates. For use in the HiTmiSS assay, both N2 and *pges-1::GFP::alg-*

1::unc54 3′ UTR C. elegans strains were synchronized as described above.

Primers design for the HiTmiSS Assay

To quantify miRNAs in high throughput, we designed 380 unique forward primers for

use in the qPCR amplification step of the HiTmiSS assay. These primers were designed

to be complementary to either the 5p strand or the 3p strand of 190 most expressed

miRNAs in the *C. elegans* genome, corresponding to 75% of all *C. elegans* miRNA

deposited in miRbase (release 22.1) (17). We did not consider miRNAs whose

existence is not supported by broad experimental evidence, nor any miRNAs predicted

to exist. Few miRNAs were ignored because their corresponding primers had a very low

Tm, making it impossible to process them using our pipeline. All primers were ordered

in four 96-well microplates from Life Technologies pre-diluted at a concentration of 50

mM (**Supplemental Table S1**). We designed a universal reverse primer complementary

to the stem-loop sequence of the stem-loop poly(A) primer used to prepare the cDNA

for this qPCR reaction. Twenty-four replicates of this primer were ordered from Life

Technologies in a separate array, also at a concentration of 50 mM. Both forward and

reverse primers were re-arrayed and diluted to a working concentration of 10 mM

immediately prior to use.

Singletons and *Switchers* identification

For each strand, we subtracted its relative quantity at one life stage versus the

relative quantity of the complementary strand (i.e. for *let-7* 5p, we compared the

abundance at the embryo stage versus the abundance of *let-7* 3p at the embryo stage).

This was repeated for each life stage at which the expression of that miRNA strand fell

above our significance threshold (1 standard deviation above the mean of all quantified

miRNAs at a given stage), and any miRNAs for which their expression was greater at

each such life stage were deemed *Singletons*. We classified as *5p Singletons* miRNAs

where the 5p strand was more expressed at each life stage than its corresponding 3p

strand. We classified as *3p Singletons* miRNAs where the 3p strand was more

expressed at each life stage than its corresponding 5p strand. We classified as

Switchers miRNAs that rose above our threshold of significance during at least one life

stage, but whose expression levels did not display a consistent pattern of usage of one

strand over the other.

HiTmiSS Assay

To collect sufficient total RNA for analysis of wild type miRNA, confluent NGM plates

of N2 strains were synchronized and either washed and pelleted immediately (embryo

stage) or incubated at 24°C for 15 hours (larval 1 stage), 25 hours (larval 2 stage), 33

hours (larval 3 stage), 42 hours (larval 4 stage), or 51 hours (young adult stage) before

washing and pelleting. All washes were performed in 15 ml Falcon tubes using 15 ml DI

water, and pellets were generated by centrifugation for 3 minutes at 400g, resulting in a

final pellet volume of 500 µL to 1 mL. Sufficient worms to generate this size pellet

differed by life stage, requiring 12 (larval 4 and young adult stages), 18 (larval 2 and

larval 3 stages), or 24 (embryo and larval 1 stages) starting confluent plates.

This pellet was then resuspended to a volume of 1.5 ml in ice-cold Triton X-100 lysis

buffer (150 mM NaCl, 1.0% Triton X-100, 50 mM Tris-HCl pH 8.0, 0.5 mM EDTA) and

transferred to a 2 ml Spectrum tube pre-filled with 0.5 mm zirconium beads. Using a

Benchmark BeadBug D1030 microtube homogenizer, each tube was homogenized for

90 seconds, incubated for 2 minutes on ice, then homogenized for an additional 60

seconds. The resultant lysate mixture was centrifuged for 3 minutes using a VWR

benchtop mini centrifuge, and the liquid fraction was transferred into a 1.5 ml Eppendorf

microcentrifuge tube. To minimize non-aqueous contaminants, this lysate was

centrifuged again in the same manner and transferred to a 15 ml Falcon tube pre-chilled

on ice. To isolate total RNA, we used a DirectZol RNA MiniPrep Plus kit from Zymo

Research, following kit instructions and performing the entire procedure at 4°C.

Resultant RNA was eluted in 30 μL NF H_2O. To collect tissue-specific miRNA for

analysis of the intestine, we performed protein immunoprecipitation followed by miRNA

isolation. The fluorescent *pges1::gfp::alg-1::unc54 3′UTR C. elegans* strain previously

prepared by our lab (18) was grown to confluence, synchronized, and pelleted as

described above for each life stage, concurrent with N2 *C. elegans* as a negative

control.

Each pellet was resuspended to 1.5 μl using our Triton X-100 lysis buffer with added

protease inhibitor (two Roche cOmplete ULTRA mini EDTA-free protease inhibitor

tablets per 50 mL lysis buffer) and transferred to a zirconium bead tube to be

homogenized as previously described. During this homogenization step, anti-GFP

magnetic antibody beads (Chromotek GFP-Trap Magnetic Agarose) were prepared by

diluting 25 µl bead slurry in 500 µl dilution buffer (150 mM NaCl, 50mM Tris-HCl pH 8.0,

0.5 mM EDTA) and rotating end-over-end at 4°C for 3 minutes.

The beads were pelleted using a benchtop centrifuge, and the buffer was aspirated

using a magnetic tube rack. 500 µl dilution buffer was added again, and this dilution

process was repeated twice. We then recovered aqueous lysate from the homogenized

mixture as previously described and diluted it using 300 µl dilution buffer. This diluted

lysate was transferred to a tube containing the prepared anti-GFP beads and left to

rotate end-over-end at 4°C overnight. To clear unbound lysate, the beads were pelleted

using a benchtop centrifuge, and the buffer was aspirated using a magnetic tube rack.

500 µl of wash buffer (150 mM NaCl, 50 mM Tris-HCl pH 8.0, 0.1% Triton X-100, 0.5

mM EDTA) was added, the beads were pelleted, and the buffer was aspirated using a

magnetic tube rack. This wash step was repeated five additional times. The beads were

then resuspended in 100 µl wash buffer and RNA was isolated using a DirectZol RNA

MiniPrep Plus kit from Zymo Research, following kit instructions and performing the

entire procedure at 4°C. Resultant RNA was eluted in 30 µL NF H_2O. The protocol for

the HiTmiSS assay was adapted from Mei *et al.* (19). We prepared our cDNA from each

life stage for qPCR analysis via polyuridinylation using Poly(U) Polymerase (NEB) as

per manufacturer instructions followed by a 20-minute heat inactivation at 65°C. We

performed a first-strand reaction on this polyuridinylated RNA using Superscript III

Reverse Transcriptase and stem-loop poly(A) primers under recommended conditions

(Invitrogen). We then performed qPCR in high throughput using this cDNA, SYBR

Green master mix (Invitrogen), and our primers as described above. We used Applied

Biosystems QuantStudio 3 and QuantStudio 7 qPCR machines to perform the screen.

For each life stage, four 96-well plates were prepared with 40 μL reactions in each well,

along with a positive control reaction containing 10 pg pDONR 221 ROG instead of

cDNA and the necessary forward and reverse primers to amplify *gfp*. Each plate was

then re-arrayed in triplicate into a 384-well plate using a Biomek FX^P Laboratory

Automation Workstation for a total of three 10 μL reactions per miRNA strand per life

stage. An additional plate was prepared with no cDNA or plasmid spike-in DNA as a

non-template control. These qPCR reactions were carried out under standard SYBR

Green reagent conditions on an Applied Biosystems QuantStudio 7. Relative miRNA

levels were calculated using the $2^{-\Delta\Delta Ct}$ method (see below), and normalized using the

spike-in positive control. These results were then plotted logarithmically as a heatmap.

Whole worm data were ordered by type (*5p Singleton, 3p Singleton*, or *Switcher*), while

intestine data were clustered using average linkage. Both datasets were visualized

using heatmapper.ca.

Cycle Threshold Analysis of the HiTmiSS Assay

The qPCR results containing the cycle threshold (Ct) values for each tested miRNA

were exported in Microsoft Excel 2019 format. A comprehensive list of these data can

be found in **Supplemental Table S2**. Ct values were averaged across experimental

replicates, then subjected to a $2^{-\Delta\Delta Ct}$ analysis to calculate initial amounts of each miRNA

in the sample, consistent with experimental standard (20). The Ct value for any given

miRNA strand was subtracted from the Ct value of the non-template control (a qPCR

reaction lacking any cDNA) for that miRNA strand. This was termed the ΔCt for that

miRNA. The same was done for the positive control for that sample. Then, for each

sample, this difference was subtracted from the difference obtained for that positive

control, normalizing samples to the same positive control and giving us the $\Delta\Delta$Ct for that

miRNA strand.

The inverse log base 2 of that $\Delta\Delta$Ct was then calculated to approximate the initial

amount of that miRNA strand present in the cDNA relative to the positive control. This

was deemed the $2^{-\Delta\Delta Ct}$ value for that miRNA strand. To determine whether a miRNA

was considered present in a sample, we decided on a strict threshold of one standard

deviation above the mean of all $2^{-\Delta\Delta Ct}$ values for that stage of development. MiRNAs

quantified at levels above this threshold were considered significant for the purposes of

this study. This same process was repeated to determine significance for our intestine

tissue-specific HiTmiSS assay.

qPCR of miRNAs targeting overexpressed intestinal 3′UTRs

To quantify miRNA strand the levels when the target 3′ UTRs of those strands are

overexpressed in the intestine, we used two *C. elegans* strains previously published by

our lab (18). Each expresses the promoter for intestine-specific gene *ges-1* fused to a

dual color ROG construct (a gene encoding *mCherry* and *gfp* separated by a trans-

spliceable operon) fused to the 3′ UTR of either *fox-1* or *hrp-2 (18)*. Each of these

strains was grown to confluence in 12 NGM plates in mixed stage and subjected to

quantification using the same approach as the HiTmiSS assay described above, though

rather than quantifying every miRNA in these animals, we quantified the 5p and 3p

strands of *miR-54*, *miR-60*, *miR-232*, and *miR-239b*. These quantifications were carried

out in 96-well plates using cDNA (both prepared from our strains as well as control

cDNA gathered from N2 *C. elegans*), SYBR Green master mix, and the appropriate

forward and universal reverse primers for the miRNAs of interest, re-arrayed from our

standard HiTmiSS plates. Each 96-well plate contained two positive controls in *let-7* and

lin-4, as well as a non-template negative control for each set of primers. These qPCRs

were carried out in experimental triplicate under standard SYBR Green reagent

conditions on an Applied Biosystems QuantStudio 3.

Relative miRNA levels were calculated using the $2^{-\Delta\Delta Ct}$ method as described above

and normalized using the average of *let-7* and *lin-4* expression for each sample. These

results were then averaged across two replicates and analyzed using Microsoft Excel

2019. Data were visualized as $\log_2$ of the change in expression from wildtype N2

negative control to either of our experimental strains. The threshold of significance was

set at one as per standard analytical procedure.

Comparison of HiTmiSS vs RNA-seq

To compare HiTmiSS results with RNA-seq data, we retrieved small RNA sequencing

datasets from Kato et al. (21) via the Sequence Reads Archive (SRA). This study

employed deep sequencing (Illumina) to profile small RNA expression, including

miRNAs, across *C. elegans* developmental stages. We analyzed the following datasets:

SRR014966 (embryo), SRR014967 (L1), SRR014968 (L2), SRR014969 (L3),

SRR014970 (L4), and SRR014971 (young adult). To assess both the presence of

miRNA and its strand usage, we developed a custom analysis pipeline. Sequencing

reads were first converted to FASTA format using the FASTX-Toolkit

(http://hannonlab.cshl.edu/fastx_toolkit/), then aligned using BLAT (22) against a

reference database containing all known *C. elegans* miRNA strands (miRbase) (17) in

FASTA format, with the following parameters: /blat -t=dna -q=dna -tileSize=6 -

minIdentity=10 -minMatch=1 -minScore=10 stage_specific.fa stage_specific.psl. To

increase stringency, we filtered the resulting PSL files to retain only matches with

Q_start = 0 and strand = "+", then quantified the occurrence of each miRNA strand at

each developmental stage. This analysis identified the following numbers of miRNA

strands: 438 in embryo, 421 in L1, 432 in L2, 431 in L3, 422 in L4, and 434 in young

adult. We then used these counts to calculate the expression value of each miRNA

strand normalized to each developmental stage.

Structural Analysis of miRNAs

The experimentally validated *5p* and *3p Singletons* were obtained using our

HiTmiSS assay. Each entry included the entire hairpin sequence and its predicted

secondary structure in dot-bracket notation (23) (**Supplemental Table S2)**. To focus

specifically on mature miRNA regions, only the sequences of each mature miRNA

duplex were used for all downstream analyses. Since the *5p* and *3p Singleton* datasets

contained different numbers of miRNAs, we balanced them before analysis by

undersampling the larger group to match the sample size of the smaller class, ensuring

unbiased statistical comparison between 5p and 3p strands.

We extracted 77 numerical features from each *5p* and *3p Singleton* (**Supplemental**

Table S3). The extracted features included: Base composition (fraction of A, U, G, C),

motif presence (e.g. UGU, GAG, CCU), positional characteristics (first/last nucleotide

identity), dinucleotide frequencies (e.g., di_GG, di_CC), GC content and GC-related

asymmetry metrics (GC_diff, GC_ratio_5p3p), strand lengths, and number of

mismatches (unpaired positions in the dot-bracket structure). The full list of the features

and their descriptions is shown in **Supplemental Table S3**.

Feature calculations were implemented in Python using regular expressions and

sequence statistics functions. Structural mismatch counts were determined by counting

unpaired characters within the uppercase-aligned mature duplex. Comparative

statistical analysis between 5p and 3p strands was conducted using Welch's two-

sample t-test. All features were tested independently, and results were ranked by p-

value. Features with $p < 0.05$ were considered statistically significant. Bar plots were

generated for each significant feature using the mean ± standard deviation across

strands.

miRNA strand selection ML model

To predict dominant strand selection (5p vs. 3p) in miRNA hairpins, we utilized

ChatGPT GPT-4o (OpenAI) to assist in developing and training a Random Forest

classifier based on 77 biologically curated features, organized into functional categories

capturing distinct structural and sequence-related aspects of miRNA hairpins. These

features capture thermodynamic stability, base-pairing patterns, and positional

relationships within the miRNA hairpin. The model was trained on our validated *5p* and

3p Singletons identified using our HiTmiSS assay, and feature values were computed

using custom Python scripts that parse dot-bracket RNA structures and extract both

absolute and relative measures for each strand. The weights of the features were

optimized using known *Singletons*. The final model incorporates the following features,

divided into five categories. The first category is the *Dinucleotide Composition* features,

which quantify the frequency of specific two-nucleotide motifs within the 5p and 3p

regions, capturing local sequence patterns that can influence strand selection and

structural flexibility. The second category is the *First/Last Nucleotide Bias* features,

which record the identity of the first and last nucleotides in each strand, reflecting known

preferences in Argonaute loading where terminal nucleotides strongly impact strand

choice. The third category is the *GC Content and Entropy* features, which assess the

global nucleotide composition and thermodynamic diversity within each strand, including

overall GC percentages, GC ratios between strands, and Shannon entropy measures

that reflect structural variability and rigidity. The fourth category is the *Base-pairing*

Properties features, which describe the fraction of nucleotides engaged in base pairing

within the 5p and 3p regions, providing a measure of local hairpin stability and

asymmetry. Finally, the *General Structural* features, which include the overall hairpin

length and the length difference between the 5p and 3p strands, capture broader

geometrical properties that may indirectly affect miRNA maturation and strand retention

(**Supplemental Fig. S1**). The complete set of features, their description, and the weight

used are shown in **Supplemental Table S3**. The model showed high performance in

distinguishing 5p and 3p usage from a pool of known *Singletons* as a test dataset (*5p*

Singleton: accuracy 100%, precision 75%, recall 100%; *3p Singleton:* 82% accuracy,

100% precision, 82% recall) (**Supplemental Fig. S2**). The ML model uses a Random

Forest classifier implemented in scikit-learn (v1.3.1). Model hyperparameters were

optimized via grid search with 5-fold stratified cross-validation, tuning: n_estimators:

[100, 250, 500], max_depth: [5, 10, None], min_samples_split: [2, 5, 10], class_weight:

['balanced', None]. The final selected parameters were: RandomForestClassifier

(n_estimators=250, max_depth=None, min_samples_split=2, class_weight='balanced').

To prevent overfitting and enhance interpretability, recursive feature elimination

(RFE) with cross-validation was used to rank features by importance. Features with

near-zero variance and multicollinearity (|Pearson's r| > 0.9) were removed. The model

consistently retained ~45–55 features, depending on the species subset. Performance

was assessed on independent held-out test sets not used during training. Metrics

included: Accuracy: 0.85 (*C. elegans*), 0.83 (human), Precision (5p/3p): 0.87 / 0.80,

Recall (5p/3p): 0.89 / 0.78, F1 Score (5p/3p): 0.88 / 0.79, AUROC: 0.91 (*C. elegans*),

0.89 (human). Class balance was maintained using SMOTE oversampling to correct for

the slight enrichment of 5p strands in the source datasets. We confirmed that the model

generalizes well by testing it on separate datasets (not used during training or tuning),

including across species (e.g., training in *C. elegans*, testing in humans).

ML model predictions in human miRNAs

Human miRNA expression data were obtained from a harmonized dataset from the

human miRNA tissue atlas (24) containing strand-specific Reads Per Million Mapped

(RPMM) values across healthy human tissues. To restrict the analysis to non-diseased

samples, we only considered entries where the Biotype column was labeled as "tissue",

"EV", or "cell_line" (**Supplemental Table S6**), representing the average expression

values across biological replicates of healthy tissues (total=337 datasets). For each

miRNA duplex, we extracted the expression levels corresponding to the 5p and 3p

strands, based on strand-specific annotations. We assigned strand dominance by

summing RPMM values across all healthy tissue samples for the 5p and 3p strands

separately. The strand with the higher total expression was designated as the dominant

strand. We compared these dominant strands' assignments with the predicted dominant

strand assigned by our ML model. Each prediction was considered correct if the model-

assigned strand matched the strand exhibiting greater expression across healthy

tissues. Evaluation was performed on the topmost highly expressed 10, 50, and 100

miRNAs ranked by the sum of 5p and 3p strands RPMM values. We also performed the

analysis using the complete dataset (all).

Results

A novel HT qPCR approach to study miRNA biogenesis

Identifying miRNAs in somatic tissues is challenging due to their small size, low

abundance, and short half-life. To overcome this, we developed a high-throughput

qPCR method that reliably quantifies miRNA expression and distinguishes 5p from 3p

strand usage. This method is a high-throughput adaptation and general improvement of

an existing methodology (19), which uses the polyuridine polymerase enzyme to

elongate a polyU tail to all RNA transcripts in a given sample. This step extends the

length of each miRNA present in the sample, which is then used as a template in

reverse transcriptase reactions using a specially engineered universal stem-loop

reverse primer (SL-polyA) (**Main Fig. 1A**).

We named this method High-Throughput miRNA Strand Selection (HiTmiSS) assay.

HiTmiSS assay enables rapid, precise quantification of the full miRNAome and

accurately distinguishes 5p from 3p strand expression (**Main Fig. 1**).

We employed this assay to systematically quantify miRNA occurrence and the

usage of 5p and 3p strands in known miRNAs throughout *C. elegans* development. We

extracted total RNA from *C. elegans* at six developmental stages (embryos, larval

stages 1-4, and young adults) and analyzed the miRNA transcriptome and strand

selection for 190 miRNAs, representing 75% of all miRNAs annotated in miRbase (17),

using 380 unique forward primers across four plates (**Supplemental Table S1**).

We performed 7,980 qPCR experiments independently, querying these 190 miRNAs

throughout development in triplicate (1,140 experiments at each stage) (**Main Fig. 1B,**

Supplemental Fig. S3). Using stringent filters (see Materials and Methods), we

detected the presence of 177 of these miRNAs (93.2%) in at least one developmental

stage (**Supplemental Table S2**). The most significant number of miRNAs occurs during

the L4 stage (81%), followed by young adult (75.5%), L3 (65%), L2 (56.8%), and lastly,

Embryo and L1 (41.6% each) stages. This aligns with previous literature suggesting that

many *C. elegans* miRNAs are tissue-specific, only becoming more abundant upon the

development of those tissues later in the developmental cycle (25).

As previously reported, early expressed miRNAs such as the key regulators of

embryonic lethality, *miR-35* and *miR-51* families, were most abundant during embryonic

and early larval stages (26). In contrast, *let-7* and *lin-4*, which control adult phenotypes,

peaked at the L4 and young adult stages (27). Notably, 13 miRNAs, including *miR-791*,

miR-792, and *miR-8206*, showed no significant expression, likely due to low abundance

or lack of experimental validation.

HiTmiSS Enhances miRNA Resolution

To benchmark the performance of HiTmiSS, we compared it to an Illumina RNA-seq

dataset from Kato et al. (21), which profiles miRNA expression across *C.*

elegans developmental stages (**Supplemental Fig. S4**). HiTmiSS captured 83% of the

miRNA strands identified by RNA-seq, demonstrating strong overall concordance

(**Supplemental Fig. S4A; left panel**). The remaining RNA-seq-only miRNAs were

generally low in abundance and exhibited diffuse expression patterns, which may reflect

biological variability or technical background often encountered in sequencing-based

approaches (**Supplemental Fig. S4A; right panel**). Importantly, HiTmiSS more clearly

resolved the top-expressed miRNAs, particularly during early development, and

provided sharper 5p/3p expression profiles across stages (**Supplemental Fig. S4B-E**).

These improvements likely stem from its use of strand-specific qPCR primers and a

standardized workflow that minimizes ligation and mapping biases. While RNA-seq

remains a powerful tool for discovery and global profiling, HiTmiSS offers superior

precision, reproducibility, and strand-level resolution for quantifying known miRNAs

across developmental or tissue-specific contexts.

Many miRNAs show consistent strand preference throughout development

Many miRNAs identified consistently use either the 5p or 3p strand as the guide

strand, which we termed '*Singletons*', while others switch between 5p and 3p usage

across different developmental stages. We named the second category '*Switchers*'.

The screen produced hits for 177 miRNAs, which we divided into *5p Singletons*, *3p*

Singletons, or *Switchers* (**Main Fig. 1B and Supplemental Table S2**).

The largest group identified was miRNA *Switchers* (46.3%, 88 hits). These include

members of the *miR-2* family and other regulators of developmental timing and stress

responses (27,28). The remaining hits were *Singletons*, with a notable bias: *3p*

Singletons (58 hits, 30.5%) outnumbered *5p Singletons* by nearly twofold. These

include *miR-35* and *miR-51* (embryonic lethality), *miR-44* (stress/starvation), and *miR-*

58 (locomotion and dauer entry) (26,27,29). Several abundant *3p Singletons*, including

miR-60, *miR-70*, and *miR-90*, remain functionally uncharacterized but are linked to

stress response (21,28,30).

5p Singletons were the least common group, with 31 hits (16.3%), including the well-

studied families *let-7* and *lin-4*, which are key regulators of developmental timing (**Main**

Fig. 1B). Also detected were *miR-51* (embryonic lethality) and *miR-64* (heat shock

response) (26,30). While some families (*let-7* and *miR-64*) consistently use the same

strand, others span multiple categories. For instance, members of the *miR-51* seed

family, including *miR-51* to *-56* and *lsy-6*, exhibit consistent strand usage, with *miR-51*

to -53 using the 5p strand and *miR-54* to *-56* and *lsy-6* using the 3p strand, yet none

show switcher behavior. Despite sharing the same seed sequence (ACCCGUA), *miR-*

53 and *-54* are processed from opposite arms and are classified into distinct families

(mirGeneDB), underscoring that strand usage can diverge even among miRNAs with

similar target specificity.

Importantly, several hits produced by the HiTmiSS assay align with past findings on

the strand preference of well-characterized miRNAs such as *let-7, lin-4* (known *5p*

Singletons), and the *miR-35* family (known *3p Singletons*) (27)

To account for the strong strand preference observed among *Singletons*, we took

the top 20 *5p Singletons* and the top 20 *3p Singletons* and compared the binding

signature and nucleotide composition between the used strand versus the strand that

gets discarded (**Main Fig. 1C, i and ii**). Interestingly, the pairing frequency does not

vary between these miRNAs (**Main Fig. 1C,ii**), either in the 5p or 3p strand, except for

the last few nucleotides at the 3'end (**Main Fig 1C, I,** red boxes) which are commonly

unpaired in the mature duplex. When comparing the used vs. the discarded strand

(**Main Fig. 1C,** left panel vs right panel), we detected no significant difference in the

pairing signature at any position; however, the 5p strand, independently if is used or

discarded, contains more unpaired nucleotides than the 3p strand (**Main Fig. 1C,** red

boxes). Expanding these results to all *Singletons* identified, the 5p strand used in *5p*

Singletons is relatively unpaired (mean = 4.42 mismatches), suggesting a tolerable level

of internal flexibility even when selected (**Supplemental Fig. S3B**). However, the 5p

strand discarded in *3p Singletons* is more paired (mean = 3.31 mismatches), indicating

that 5p strands tend to be more stable (i.e., more paired) when they are not selected

(**Supplemental Fig. S3B)**. Conversely, the two strands in *5p Singletons* are equally

unpaired (mean = 4.42 mismatches), which might suggest that in some cases the

unselected strand is more disordered (**Supplemental Fig. S3B**).

When narrowing our analysis to just the seed region (**Main Fig. 1C**, golden boxes),

while there is higher pairing than any other position, there is no difference in binding

frequency between the used and discarded strands, suggesting that other rules are also

in place.

Next, we compared the nucleotide composition of these seed regions, hoping to

identify signatures that could explain the discrepancy in miRNA strand selection in

Singletons. We determined the likelihood of any given nucleotide from positions 1

through 10 in each miRNA strand in the top 20 *Singletons* in each category (5p and 3p)

(**Main Fig. 1C, ii**). We detected a highly conserved uracil nucleotide at position 1, as

previously shown by us and others (2), though this conservation dissolves from

positions 2 through 10 (**Main Fig. 1C, ii**). Importantly, this uracil nucleotide is not

equally conserved between *5p Singletons* and *3p Singletons*. 80% of *3p Singletons* start

with this uracil nucleotide, while less than 40% occur in *5p Singletons*. This differential

presence of this uracil may be caused by a non-stochastic pre-miRNA cleavage by the

C. elegans DCR-1, which may possess a preference to cleave at this nucleotide.

The intestine miRNA population is small but highly interconnected

The vast majority of hits detected by our HiTmiSS assay were *Switchers* (~46%).

Many of them may indeed not be true *Switchers* but rather act as tissue- or stage-

specific *Singletons*, potentially confounding the analysis. To address this, we refined our

approach by analyzing somatic tissues across development, focusing on the intestine,

which is a large tissue with a well-characterized transcriptome (31,32).

We coupled our HiTmiSS assay with intestine-specific miRNA immunoprecipitation,

as we have published in the past (18), focusing on the embryo vs young adult stages.

More specifically, we forced the expression of the GFP-tagged *C. elegans* Argonaute

ortholog *alg-1* in the intestine tissue using the intestine-specific promoter *ges-1p*,

performed an α-GFP immunoprecipitation, recovered the miRNA strands loaded onto

ALG-1, and used them as templates in our HiTmiSS assay (**Main Fig. 2A**).

We found that of the 96 miRNAs tested, 64 (68%) were present during at least one

stage within the intestine tissue (**Main Fig. 2**). Of these, 58 (91%) were *Singletons*,

while 6 (9%) were *Switchers* (**Supplemental Fig. S5A**).

This strongly differs from our whole worm data, reinforcing that miRNAs previously

labeled as *Switchers* may truly be *Singletons* on a tissue-by-tissue basis, only

appearing as *Switchers* once all tissues are analyzed together.

Interestingly, the proportion of *5p Singletons* to *3p Singletons* is not the same in the

intestinal tissue as in the mixed tissue (**Supplemental Fig. S5A**). In our initial analysis,

about two-thirds of *Singletons* were *3p Singletons*, yet in the intestine tissue, nearly two-

thirds of *Singletons* are *5p Singletons* (**Supplemental Fig. S5A**).

Another surprise was the presence of intestine-specific *Switchers* (**Main Fig. 2A and**

Supplemental Fig. S5B). This includes the *miR-51* family, well-characterized regulators

of embryonic lethality (26). In addition, we detected intestine tissue *Switchers* among

the *miR-44, miR-58*, and *miR-71* families (**Supplemental Fig. S5B**). These miRNA

families have diverse functions, ranging from regulating developmental timing to

responses to starvation, hypoxia, stress, and dauer entry(28,30,33,34). Importantly,

these are intestine-specific phenotypes, suggesting that strand switching in these

miRNAs may have functional significance by expanding the 3' UTR targeting repertoire

in the intestine.

stage. These strains expressed the promoter for a single miRNA fused to GFP, and live

imaging was used to validate the expression of these miRNAs at the appropriate life

stage in the intestine (**Main Fig. 2C**). All four strains displayed the expected

fluorescence.

miRNA target presence doesn't account for strand usage

Our HiTmiSS array enables precise, strand-resolved quantification of miRNAs *in vivo.*

We considered the possibility that miRNA targets might stabilize specific strands,

potentially biasing detection in our HiTmiSS assay. In this model, only target-bound

strands would be retained, while the unused strand would degrade, falsely suggesting

strand preference. To test this, we overexpressed *fox-1* and *hrp-2* 3'UTRs, each

containing multiple binding sites for typically discarded strands (e.g., *miR-60* 5p, *miR-*

232 5p, *miR-239b* 5p), in intestinal tissue (**Supplemental Fig. S6**). If target-induced

stabilization were driving strand selection, we would expect increased detection of these

5p strands. However, strand usage remained unchanged compared to wild-type

controls, with 3p strands (i.e., *miR-60* 3p, **Supplemental Fig. S6C**) still predominating.

Taken together, these results indicate that our qPCR approach is not simply

detecting stabilized miRNA strands due to the presence of targets. Rather, the relative

strand abundance measured by the HiTmiSS assay likely reflects genuine strand

selection, rather than artifacts introduced by target availability.

5p Singletons **and** *3p Singletons* **follow different rules for selecting strand usage**

Next, we studied the structural features associated with *Singletons*, which could

explain the strict and consistent strand usage within these miRNAs. When comparing

the used and the discarded strand, the pairing frequency varies significantly within each

Singleton, revealing new conserved patterns (**Main Fig. 3**). Specifically, the strand used

in *Singletons* often has an unpaired nucleotide at position 12, while the discarded strand

does not (**Main Fig. 3A**, yellow boxes). This lack of pairing in position 12 in both *5p*

Singletons and *3p Singletons* usage is intriguing. In humans, specific Argonaute

proteins perform cleavage of target RNAs between positions 10 and 11, which

corresponds to positions 10 and 11 along the bound miRNA strand by Watson-Crick

pairing (35). *C. elegans* ALG-1 is largely homologous to human Argonaute proteins, and

it may retain this cleavage activity at similar target positions, leading to the degradation

of the unused strand due to its Watson-Crick base pairing at this crucial position. In

addition, we detected differences in the binding frequency of the first-position nucleotide

in both *5p Singletons* and *3p Singletons* (**Main Fig. 3A,** blue boxes). *3p Singletons* also

exhibit unique pairings at positions 18-19 (**Main Fig. 3**, white box) not present in the

discarded strain. These patterns are most clearly seen in specific *Singleton* miRNAs,

both *5p Singletons* such as *miR-52* and *3p Singletons* such as *miR-75* (**Main Fig. 3B**).

An updated machine learning (ML) model for miRNA strand selection

We wanted to extract patterns and features that make *Singletons* so unique and

predispose them to always use one strand over the other. To achieve this, we utilized

our wet bench *Singleton* data as a training set for a machine learning model designed to

predict 5p versus 3p strand usage in miRNA hairpins. We hypothesized that by training

such a model, we could deconvolve the unique intrinsic features that characterize

Singletons.

We opted for a Random Forest classifier, which we trained using our experimentally

validated *Singletons* identified through our HITmiSS assay. This model incorporates 77

features, systematically organized into five functional categories (dinucleotide

composition, first/last nucleotide bias, GC content and entropy, base pairing properties,

and general structural features) that together capture critical aspects of miRNA mature

duplex structure (**Supplemental Fig. S1–S2, Supplemental Table S3-S4, and**

Materials and Methods). The model demonstrated strong performance in

distinguishing 5p from 3p usage across a set of known miRNAs (**Supplemental Fig. S2**

and Materials and Methods).

***5p* and *3p Singletons* possess distinct structural features**

Our ML model, built using our *Singleton* data, identified several intrinsic features in

these miRNAs. To generalize this model across all *C. elegans* miRNAs and study how

Switchers can bypass these rules to switch miRNA strand preference, we applied our

ML model and conducted a comparative feature analysis across the entire *C. elegans*

miRNAome (374 miRNAs, miRbase).

We extracted and calculated all 77 features used in our ML model for these miRNAs

(**Supplemental Table S4**), maintaining a stringent definition of *Singletons* to enhance

the signal (confidence score > 80%).

As expected, we identified 54 *5p Singletons*, 52 *3p Singletons,* and 268 *Switchers*

(**Main Fig. 4A-B** and **Supplemental Table S4**). Out of 77 extracted structural and

sequence features, 13 exhibited statistically significant differences between the 5p and

3p strands ($p < 0.05$, Welch's t-test) (**Main Fig. 4B-E**). The most pronounced difference

was observed in the identity of the first nucleotide of the 3p strand, which in *3p*

Singletons was consistently a uracil ($p = 3.98 \times 10^{-\square}$), followed by an enrichment of

cytosine in the last position ($p = 5.69 \times 10^{-\square}$) (**Main Fig. 4C-4E**). Additionally, 3p

strands showed an enrichment for specific motifs, particularly GAG ($p = 2.51 \times 10^{-\square}$)

(**Main Fig. 4E**). We also observed compositional differences: 3p strands displayed

increased levels of cytosine at the first position (not shown) and tended to be slightly

shorter than their 5p counterparts ($p = 8.90 \times 10^{-3}$) (**Main Fig. 4D-E**).

Surprisingly, our ML model did not cluster all the *Switchers* around a confidence score

of ~50%. Instead, we see a multi-modal distribution with several periodical clusters

(**Main Fig. 4A**). This suggests that across this category, structural and thermodynamic

features alone cannot fully explain strand usage, and extrinsic factors could move the

needle between 5p or 3p choice in these miRNAs, perhaps depending on tissue or

developmental context.

We next wanted to study the intrinsic features identified by our ML model. For these

experiments, we used *Singletons* to test if altering these rules could convert them into

Switchers. In *let-7*, modifying nucleotide pairing at positions 11–12 (UP11-12) or

changing the first base (5'U→C) reduced its predicted *5p Singleton* confidence by up to

17%, consistent with our model (**Main Fig. 3C-D**). Similar results were observed in *miR-*

52, *miR-75*, *lin-4*, and *miR-35*: introducing bulges near the center disrupted strand

preference, while removing them reinforced it (**Supplemental Fig. S7**), highlighting the

central region's role in strand selection.

The first nucleotide's identity and pairing also influenced strand prediction, though with

variable impact across miRNAs (**Supplemental Fig. S7**). Mutating this position altered

strand usage by less than 10% in *5p Single*tons but up to 30% in *3p Singletons*,

mirroring HiTmiSS results and further supporting our strand selection model.

Strand usage bias differs by species

Our ML model identified several key intrinsic features that clearly separate

Singletons from *Switchers* in *C. elegans*. Since many miRNA sequences are conserved

across species, we hypothesized that the features identified by our ML model were

selectively maintained throughout evolutionary time. To do that, we applied the same

model to the complete human miRNAome (**Main Fig. 5**) (**Supplemental Table S5**), and

to other species throughout the tree of life (*Drosophila melanogaster*; fly, *Danio rerio*;

zebrafish, *Xenopus laevis*; frog, *Rattus norvegicus*; rat, *Mus musculus*; mouse),

(**Supplemental Fig. S8** and **Supplemental Table S5**). While miRNA strand usage in

worm, fly, zebrafish and frog remains relatively balanced between the 5p and 3p arms,

mammalian species exhibit a pronounced preference for the 5p strand (**Supplemental**

Fig. S8B). In *C. elegans*, only 51.3% of miRNAs were predicted to use the 5p strand,

and in *Drosophila*, 55.6% were 5p. In contrast, this proportion rose sharply to 78.2% in

mouse and 75.9% in human, indicating a shift toward fixed strand preference (**Main Fig.**

5A and **Supplemental Fig. S8)**.

Next, we sought to identify structural features that could explain this sharp increase

we observed in 5p strand usage in mammals. We have identified 13 significantly

different features between worms and humans ($p < 0.05$, Welch's t-test), with striking

enrichment of guanine content ($p = 5.62 \times 10^{-2}$) and GG dinucleotides ($p = 1.94 \times$

10^{-2}) on human 5p strands (**Main Fig. 5B**). Invertebrates exhibited a subtle

enrichment in AU-rich motifs (UU: +2.46% higher in invertebrates; AU: +2.39% UA:

+1.70%; AA: +2.11%), consistent with lower duplex stability and more flexible strand

usage (**Supplemental Fig. S8D**). In contrast, mammals showed increased frequencies

of GC-rich dinucleotides such as GG and GC, especially within the disfavored strand

(di_GG: 3.6% higher in mammals than in invertebrates; di_GC: 0.9% higher in

mammals) (**Supplemental Fig. S8D**). The overall GC content in worm, fly, and frog

miRNAs hovered around 44-45%, whereas mouse and human miRNAs exhibited

significantly higher GC content (~52–53%). This overall increase in GC content, which

translates to an increase in duplex stability, could be one of the most important features

used in vertebrates to force the usage of the 5p strand.

Across species, we also observed an apparent decrease in pre-miRNA length of the

used strands, especially in *Switchers* and *3p Singletons* (**Supplemental Fig. S8C**). This

suggests that in vertebrates shorter pre-miRNAs can assume fewer folding patterns,

perhaps enhancing specific strand usage.

Finally, analysis of 5' nucleotide identity showed that all species retained a

preference for uracil at the first nucleotide of the loaded strand (2). However,

mammalian species exhibited an increase in guanine and cytosine at the first position of

the 3p strand, as well as higher CCU content and diCC motifs ($p < 5 \times 10^{-2\square}$). These

trends are evident in conserved miRNAs such as *let-7* (**Main Fig. 5C**), potentially

serving as a negative selection marker to suppress its incorporation into RISC.

Taken together, these features may provide structural hints as to how vertebrates

adopted a fixed strand usage. Shorter and more paired duplexes may enhance

selection of *Singletons* in vertebrates, while longer and less structured sequences may

allow invertebrates to increase usage of *Switchers*.

Our ML model can accurately predict human miRNA strand selection

Our *C. elegans* ML model was also capable of identifying sequence elements and

structural features in human miRNAs. To validate these results, we compared its

predictions to expression data from the Human miRNA Tissue Atlas across 65 healthy

tissues (24) **(Main Fig. 6)**. For the top 10 most expressed miRNAs, the model reached

90% accuracy, which is higher than in *C. elegans*. As more miRNAs were included,

accuracy declined, reaching 59% across the full dataset, yet remained statistically

significant (p ≈ 7.6 × 10⁻□) **(Main Fig. 6A)**.

To reduce noise from low-expression miRNAs, we focused on the 50 most abundant

duplexes (**Main Fig. 6B**). The model correctly predicted the dominant strand in 33

cases, yielding 66% accuracy (p ≈ 0.0164). Most errors involved 3p-dominant miRNAs

misclassified as 5p, reflecting a model bias toward 5p predictions. These

misclassifications often occurred when 5p and 3p strands had similar expression levels,

suggesting either model uncertainty or that these miRNAs may function as *Switchers* in

humans (**Supplemental Table S6**).

Notably, two misclassified miRNAs, *miR-182* and *miR-378a*, are known *Switchers in*

vivo (36,37). In other cases, mispredictions occurred when the model's confidence

scores for 5p and 3p usage were nearly equal (±10%). For instance, *miR-199a* and

miR-199b, involved in hypoxia and cardiac regulation (38), were predicted as 5p

dominant despite showing 3p expression and a narrow confidence margin (~7%).

Similarly, *miR-124*, critical for neurogenesis, was misclassified with a 6% confidence

gap, despite clear 3p dominance (39).

Still, the majority of miRNAs were accurately predicted across diverse human

tissues, including key regulators with strong strand bias, indicating that our *C. elegans*-

trained model generalizes well to vertebrates (**Main Fig. 6C and Supplemental Table**

S6). For example, *miR-21*, a highly expressed oncomiR, was correctly predicted as 5p-

dominant, with over 50-fold 5p expression compared to 3p(40). Similarly, *let-7a* showed

a 248-fold 5p preference and was correctly classified (41). Other highly expressed

miRNAs, such as *miR-1* and miR-*141,* were also correctly predicted as 3p dominant.

Overall, these findings demonstrate that the model is capable of reliably predicting

strand usage for miRNAs in healthy human tissues, particularly in cases where strand

bias is strong. Errors were enriched among duplexes with more balanced expression

levels or context-dependent strand usage, highlighting the complexity of predicting

which strand will be used in cases of weak or dynamic regulation of strand selection.

A community resource to study miRNA strand selection

To support the broader research community, we have made all components of our

strand selection framework openly available via GitHub

(https://github.com/MangoneLab/Caligula_app). This resource includes our trained ML

model, Python scripts for both local, cumulative and browser-based use. The platform is

designed for flexibility and ease of use, allowing users to input a single miRNA or a

batch of sequences in FASTA format to obtain strand selection predictions.

We also provide pre-processed strand selection predictions, including all 77 features

used by our ML model, for every miRNA annotated in miRBase across seven species:

worm, fly, zebrafish, frog, mouse, rat, and human (**Supplemental Tables S2 and S5**).

By making these tools and datasets accessible, we aim to provide a practical, scalable,

and user-friendly solution for investigating miRNA strand usage across systems and

species. This represents the first publicly available platform for high-throughput, strand-

specific miRNA analysis, enabling researchers to integrate strand selection into both

basic and translational studies.

Discussion

In this study, we investigated miRNA strand selection by integrating ~8,000 HiTmiSS qPCR assays in *C. elegans* with structural modeling and machine learning.

Compared to RNA-seq, HiTmiSS offers distinct advantages for miRNA analysis (**Supplemental Fig. S4**). By using strand-specific qPCR primers, HiTmiSS directly quantifies 5p and 3p strands, avoiding the inference-based strand assignment required in RNA-seq, which can be affected by ligation and mapping biases. It also demonstrates higher sensitivity for low-abundance miRNAs, reliably detecting transcripts that are often missed or inconsistently captured by RNA-seq due to shallow depth or background noise (**Supplemental Fig. S4A-D**). Moreover, HiTmiSS shows greater reproducibility across replicates, thanks to its standardized primer design and normalization controls, making it particularly well-suited for comparative and developmental studies. While RNA-seq remains valuable for discovering novel miRNAs and profiling diverse RNA species, HiTmiSS is optimized for precise, strand-specific quantification of known miRNAs. Recent approaches, such as AQ-seq, have improved small RNA-seq protocols by reducing ligation bias and enabling isomiR detection (42). However, because AQ-seq still relies on sequencing-based inference for strand assignment, it may be affected by residual mapping ambiguities. In contrast, HiTmiSS directly quantifies 5p and 3p strands using strand-specific primers, offering a more targeted approach to strand-resolved analysis. Its combination of sensitivity, quantitative precision, and reproducibility makes it well suited for studying miRNA strand selection across tissues and developmental stages.

Our study allowed miRNA classification into *5p Singletons*, *3p Singletons*, or *Switchers*,

and informed a predictive model validated against human miRNA expression data.

We found that in *C. elegans* half of miRNAs (46%) behave as *Switchers*, altering strand

usage across development, indicating that strand selection is context-dependent rather

than fixed (**Main Fig. 1)**. In contrast, key developmental regulators such as *let-7*, *lin-4*,

and *miR-35* are *Singletons*, likely reflecting a need for consistent strand usage to

ensure stable gene expression. Notably, *Singletons* tend to be more highly expressed

(**Main Fig. 1B**), perhaps due to stronger structural or thermodynamic constraints. In the

intestine, tissue-specific ALG-1 immunoprecipitation revealed a reversal of the

developmental trend, with more *5p* than *3p Singletons* (**Main Fig. 2**). This suggests that

strand selection is also shaped by tissue-specific factors, such as the cellular

environment or cofactors. Since we examined only one somatic tissue, some miRNAs

classified as *Switchers* in our experiments may be tissue-specific *Singletons*.

A concern was that target mRNAs might stabilize one strand, creating an artificial

strand preference in our HiTmiSS assay. To test this, we overexpressed 3'UTRs with

binding sites for typically low-abundance strands in the intestine. Strand levels remained

unchanged, indicating that HiTmiSS reflects true strand usage, not target-driven

stabilization (**Supplemental Fig. S6**).

When comparing the structural composition of each mature duplex, a series of

patterns emerged. Regarding the strand pairing, there's no universal trend showing that

5p strands are always more unpaired than 3p strands, regardless of selection. Instead,

selected strands (whether 5p or 3p) tend to tolerate moderate unpairing, and unselected

5p strands (i.e., in *3p Singletons*) are more paired, suggesting that greater structural

stability may discourage selection, supporting a role for pairing in strand discrimination

(**Main Fig. 1C** and **Supplemental Fig. 3B**).

5p Singletons tend to have the first nucleotide unpaired to its corresponding strand

in the duplex (**Main Fig. 3A**). This difference in pairing repeats in the middle of the

duplex and lowers toward the end (**Main Fig. 3A**). Contrarily, *3p Singletons* present a

near-opposite trend, with the first nucleotide more often paired and the end portion of

the sequence also paired, though the center of the sequence shows similar unpairing to

5p Singletons.

The effects of the 5' nucleotide identity and central region pairing on strand selection

and Argonaute loading have been well known for a long time, with foundational work

highlighting the importance of 5' nucleotide identity and central region pairing for

Argonaute loading (7,43,44). These studies established general rules governing strand

preference, such as the thermodynamic asymmetry model, but were derived under

controlled conditions, often focusing on a limited number of miRNAs in model systems.

While they established the principles of strand selection, they did not provide a scalable

means to measure how these rules are implemented, or overridden, in living organisms

across developmental stages, tissues, or physiological contexts. HiTmiSS advances the

field by shifting the focus from mechanism to measurement, offering a practical, high-

throughput platform for directly quantifying strand usage across biological conditions,

developmental stages, and tissues. In doing so, HiTmiSS transforms strand selection

from a static rule set into a measurable, context-dependent regulatory layer.

In addition, our results added positional detail to these foundational studies by

showing that asymmetry varies along the duplex. Both *5p* and *3p Singletons* show

central unpairing, likely reflecting a conserved feature for Dicer or Argonaute

engagement but not strand bias. Notably, *5p Singletons* show decreasing pairing from

the seed to the 3' end, while *3p Singletons* display the opposite trend. These opposing

patterns suggest that specific structural cues along the duplex, not just global stability,

guide strand selection, pointing to evolved, position-dependent regulatory signals within

miRNA architecture.

To expand our analysis, we extracted 77 structural and compositional features from

our validated *Singletons* and trained a Random Forest classifier (ML model) to predict

dominant strand usage. When comparing these features (**Main Fig. 4**), we found

several strand-specific differences, including a significantly higher occurrence of uracil

at the first nucleotide in 3p strands and enrichment for GAG motifs.

We validated our predictions using *in silico* site-directed structural mutations in

canonical miRNAs such as *let-7*, *lin-4*, and *miR-52*, *miR-75* (**Main Fig. 3C**,

Supplemental Fig. S5). Disrupting features such as central bulges (**Main Fig. 3C,**

"UP11-12") or first nucleotide pairing (**Main Fig. 3C, "1U->C"**), altered predicted strand

preference to some degree, further confirming their role in the selection process.

Surprisingly, the degree to which they impact miRNA strand preference was

approximately equal, but when disrupted in tandem, led to an even greater loss of

predicted strand preference (**Main Fig. 3C, "Combined"**, **Supplemental Fig. S7**). In

addition, these findings phenocopy the trends seen in our experimentally validated

Singletons (**Main Fig. 3A**). Taken together, these results suggest that the structural and

thermodynamic features identified in our wet bench validated *5p* and *3p Singletons* are

indeed critical regulators of strand selection in nematodes.

It is important to note that a decrease in our ML model confidence score does

not necessarily imply a functional conversion from *Singleton* to *Switcher* status.

Reduced confidence could arise from multiple factors, including the modified input

moving outside the model's training distribution. However, we designed the *in silico*

perturbations in **Main Fig. 3C** and **Supplemental Fig. S7** carefully to isolate features

with known or suspected influence on strand selection, such as 5' nucleotide identity,

central mismatches, and thermodynamic asymmetry. Importantly, these features were

modified within biologically plausible ranges and consistent with the feature distributions

observed in the training data. Moreover, we did not interpret confidence score

reductions in isolation. Rather, we focused on systematic and reproducible changes in

model predictions across multiple miRNAs and perturbation types. In several cases,

including miR-75 (**Supplemental Fig. S7**), modified sequences not only showed

reduced confidence but also flipped strand preference, which is a more direct indication

of switcher-like behavior. Taken together, the decrease in confidence, especially when

accompanied by a shift in predicted strand, suggests that the altered input affects strand

usage potential in a direction consistent with switcher characteristics.

Expanding this framework, we then applied our ML model to predict strand usage

across the human miRNA dataset (**Main Fig. 5A**). We were particularly curious to test if

the structural and thermodynamic features and motifs identified in *C. elegans* were

conserved in vertebrates.

Compared to the nearly equal distribution of 5p and 3p usage in *C. elegans*, using

the same cutoff of 80%, the vast majority of human miRNAs are identified as *Switchers*

(831/854 predicted *Switchers*, 97.3%). However, within this class, there is a clear bias

toward 5p strands, with over 75% of predictions favoring 5p strand usage (**Main Fig.**

5A). This divergence likely reflects increased specialization in miRNA maturation and

Argonaute loading in vertebrates, potentially driven by evolutionary selection for strand-

specific targeting or stability.

Despite the lower predictive accuracy of our ML model when applied to *H.*

sapiens miRNAs (59%) compared to its performance on the *C. elegans* training set

(88%) (**Main Fig. 6A**, **Supplemental Fig. S2**), we do not interpret this as evidence of

overfitting. Rather, the ability of the *C. elegans*–trained model to achieve performance

substantially above random (50%) on a distantly related species suggests that key

aspects of miRNA strand selection are indeed governed by conserved predictive

features. While quantitative differences in model accuracy likely reflect evolutionary

divergence and the influence of lineage-specific modifiers, the persistence of these core

features supports the idea that a foundational set of strand selection rules is conserved.

Thus, our model reveals not only species-specific differences but also a shared

regulatory logic underlying miRNA strand selection that has been retained across

metazoan evolution (**Supplemental Fig. S8**). Invertebrate systems appear to tolerate

greater flexibility, allowing for more balanced usage of the duplex arms. In contrast,

vertebrates, particularly mammals, exhibit strong sequence and structural biases that

favor the selection of the 5p strand (**Main Fig. 5**). This transition likely reflects the

growing complexity of regulatory networks and the need for precise post-transcriptional

gene control in higher organisms. As miRNA families expanded during vertebrate,

particularly mammalian, evolution, selective forces likely favored stable strand usage to

ensure consistent and non-redundant regulation (2). This trend would promote

Singletons over *Switchers*, reducing regulatory variability. The fixation of 5p strand

usage may thus represent stabilizing selection for predictable miRNA function, with

structural features reinforcing 5p bias evolving alongside increasing organismal

complexity.

In addition to this imbalance in *Singletons* vs *Switchers*, the nucleotide composition

of their 5p and 3p strands is also different, with a significant enrichment of guanine

content in human 5p strands and cytosine-rich motifs in 3p strands (**Main Fig. 5B**).

These compositional biases are consistent with known differences in Argonaute protein

specificity and may reflect lineage-specific constraints on strand functionality (45).

Importantly, these cross-species trends persisted in known miRNAs such as *let-7*,

which was consistently predicted and validated as a *5p Singletons* both in *C. elegans*

and in humans, although at different confidence levels. This supports the robustness of

our predictive ML model and highlights conserved features of miRNA processing, even

in the presence of divergent strand preferences.

A surprising trend emerges when we compare our ML model prediction in human

miRNAs to the human miRNA Tissue Atlas. Our ML model developed in *C. elegans* is

capable of accurately predicting miRNA strand usage in humans (**Main Fig. 6**). This is

unexpected, considering the vast evolutionary gap (~600 million years) between these

two systems, which share less than 35% of nucleotide conservation in genes (46).

Despite this, the features we have identified seem to be evolutionarily conserved.

This suggests that miRNA strand selection is, at its core, a fundamental cellular

process that has evolved very little over time, even as miRNA regulatory networks

expanded in complexity with the increased cellular specialization. This becomes more

evident when we overlay miRNA strand usage in human tissues of different origins.

Here, the same miRNA is always predicted to use the same strand in other somatic

tissues (**Main Fig. 6C**). This invariance suggests that strand selection is most likely

encoded by intrinsic properties of the hairpin itself, rather than by external, context-

dependent cues. Such robustness implies that strand choice is not merely a byproduct

of downstream regulatory needs, but a core mechanistic feature established early in

miRNA biogenesis. We propose that during vertebrate evolution, particularly with the

rise of tissue specialization, miRNA strand usage remained anchored to conserved

structural determinants even as the broader miRNA regulatory network expanded and

diversified. This mechanistic rigidity may stabilize gene regulatory outputs in the face of

cellular complexity.

While our model shows that strand selection for many miRNAs can be accurately

predicted based on intrinsic properties of the hairpin structure, the prevalence of

Switchers strongly suggests the involvement of additional, context-dependent regulatory

mechanisms. These may include tissue-specific cofactors, differential Argonaute

availability, or post-transcriptional modifications influencing duplex stability or Argonaute

loading preferences.

We have developed a user-friendly resource on GitHub

(https://github.com/MangoneLab/Caligula_app) that allows researchers to predict

miRNA strand selection using our ML model. To support broad application, we also

provide pre-processed strand selection predictions with all 77 input features for all

miRNAs in miRBase across seven species, including worm, fly, zebrafish, frog, mouse,

rat, and human (**Supplemental Tables S2 and S5**).

In summary, this study establishes a predictive and experimentally validated

framework for miRNA strand selection, combining high-throughput qPCR (HiTmiSS)

with a ML model to uncover both conserved and context-dependent rules governing

strand usage. Unlike previous studies that inferred strand selection indirectly or focused

on a limited set of miRNAs, our study provided strand-specific quantification across

developmental stages and tissues in *C. elegans*, revealing dynamic and non-stochastic

strand usage. We also produced a biologically grounded and publicly available ML

model trained on curated *in vivo* strand data that accurately predicts strand selection

across multiple species, including vertebrates. We also prepared the first species-wide

probability atlas of 5p vs. 3p usage across several metazoan miRNAs, including

humans, enabling genome-scale interpretation of strand bias. In addition, we highlighted

a detailed dissection of 77 intrinsic features that together define a conserved,

programmable regulatory logic for strand selection. This integrative approach moves

beyond descriptive rules, offering a quantitative, generalizable model of strand selection

applicable across metazoan evolution.

In conclusion, our findings illustrate the complexity of miRNA strand selection and

reveal that this process is not governed solely by thermodynamic asymmetry. Instead, a

combination of structural motifs, sequence biases, and cellular context dictates strand

usage. Our work establishes a predictive framework for understanding strand selection

but also opens new avenues for studying the evolution of miRNA regulatory logic across

metazoans.

Data Availability

The raw data produced by the HiTmiSS assay are included in the **Supplemental Table**

S2. The ML model is included as Supplemental Data and is also available on GitHub

(https://github.com/MangoneLab/Caligula_app).

Funding

969 This work was funded by the National Institutes of Health 5R01GM118796 awarded to

970 M.M.

Acknowledgments

973 We thank Dr. Murugan for assistance in HiTmiSS assay optimization.

Author Contributions

D.M. and M.M. designed the experiments. D.M. performed the experiments. D.M. and

M.M. prepared the ML model and performed the bioinformatics analysis. H.L.

supervised the development of the ML model. D.M., A.E., and H.H. performed the

HiTmiSS assay experiments. D.M. and M.M. prepared the main and supplemental

figures and supplemental tables. D.M. and M.M. wrote the manuscript. All authors have

read and approved the manuscript.

Declaration of Interests

984 The authors have no conflicts of interest.

Figure titles and legends

Main Figure 1: HiTmiSS Assay and Analysis on Mixed Tissue. **(A)** HiTmiSS protocol

for RNA extraction, extension, reverse transcription, and quantification in high

throughput. Steps include (i) *C. elegans* growth and homogenization, (2) RNA extraction

and isolation, (3) RNA polyuridinylation, extension, and reverse transcription (details

shown in ii ; orange: mRNAs, green: polyU, purple: SL-polyA reverse universal primer),

(4) addition of 5p or 3p miRNA-strand-specific forward primers, (5) loading into 384-well

plates, and (6) quantification. **(B)** Heatmap of miRNA expression (n=3) in mixed tissue

clustered by quantity, strand use, and developmental stage. Stages include embryo (E),

larval 1 through 4 (L1-L4), and young adult (YA). The heatmap legend represents $\log_{10}$

of expression values for each miRNA strand normalized by spike-in GFP control at each

stage from -4 (deep purple) to 4 (bright yellow). Black color indicates instances of data

that fall below strict expression threshold (one standard deviation above mean of all

expression values at that stage). **(C)** Heatmap of pairing frequency by position for the

used (left) and discarded (right) strand of the 20 most highly quantified *5p* and *3p*

Singletons in our mixed tissue dataset. (i) blue: paired; red: unpaired). (ii) Average

pairing frequency, and logo plot of nucleotide frequency at each position displayed

below (gold box mark the seed region).

1004

Main Figure 2: Analysis of Intestine Tissue-Specific miRNAs. **(A)** Schematics of the

intestine-specific GFP-tagged ALG-1 immunoprecipitation. **(B)** Heatmap of miRNA

expression in intestine tissue clustered by strand and developmental stage. The

heatmap legend represents $\log_{10}$ of expression values for each miRNA strand

normalized by spike-in GFP control at each stage from -4 (deep purple) to 4 (bright

yellow). Black color indicates instances of data that fall below the strict expression

threshold (one standard deviation above the mean of all expression values at that

stage). E: embryo; YA: young adults. **(C)** Representative brightfield (left) and GFP

(right) images of selected *C. elegans* strains expressing GFP under the control of

promoters for miRNA known to be highly expressed in the intestine, either during

embryo stage (top two) or young adult stage (bottom two). Arrows point to GFP

expression in the intestine tissues in both embryos and young adult stages.

1017

Main Figure 3: Structural and Thermodynamic Determinants of Strand Selection.

(A) Pairing frequency by position of the used (green) and discarded (red) strand for the

most highly quantified *5p Singletons* and *3p Singletons* in mixed tissue (n=20). Positions

at which pairing frequency most differs are boxed, including the first nucleotide (light

blue), the center of the duplex (gold), and the end of the strand (white). **(B)**

Representative pri-miRNA hairpins for a *5p Singleton* (*miR-52*) and a *3p Singleton*

(*miR-75*) demonstrating the occurrence of trends in pairing as described above. Mature

duplex regions are highlighted in pink. **(C)** Changes in *Singleton* predictive confidence

for the miRNA *let-7* as a result of *in silico* site-directed mutations. RNAfold secondary

structures of pri-miRNA with mature 5p strand (blue) and mature 3p strand (red) from

left to right: wildtype (WT), unpaired nucleotides at positions 11 and 12 via GG→UU

substitution (UP11-12), first nucleotide U→C substitution (1U → C), combination of both

prior mutations (Combined). The table summarizes the predicted strand preference by

our ML model for each mutation. **(D)** Bar chart displaying probability as described

above.

1033

Main Figure 4: Analysis of Mature miRNA Duplex Features in *C. elegans*. (A)

Frequency distribution of *5p Singleton* predicted confidence by our ML model across all

miRNAs in *C. elegans*. *3p Singletons* (yellow) are defined as miRNAs falling below a

predicted 5p confidence of 20%; *Switchers* are defined as miRNAs with a predicted 5p

confidence between 20% and 80%; *5p Singletons* (red) are defined as miRNAs with a

predicted 5p confidence of 80% or greater. **(B)** Predicted categorization of all *C.*

elegans miRNAs by our ML model as per the above threshold. **(C)** Predicted

categorization of *C. elegans* miRNAs for which uracil is the first nucleotide of the 3p

strand. **(D)** Length difference (5p strand - 3p strand) for each category of *C. elegans*

miRNAs. *5p Singletons* display significantly ($p < 0.05$) longer 5p strands than 3p strands

when compared either to *3p Singletons* or to *Switchers*. **(E)** Heatmap of 32 features

including GC content, miRNA length, hydrogen bonding, nucleotide pairing, strand

stability, and MFOLD-derived parameters enrichment in *5p* and *3p Singletons*. Heatmap

values represent the mean values of each feature across the miRNA categories. For

binary features (i.e., 5p_has_AAU, which is either 0 or 1), the number represents the

proportion (e.g., 0.3 = 30% of miRNAs in that category have an AAU motif in the 5p

strand). For continuous features (i.e., gc_content, pairing_frac_5p), the number is the

average value across all miRNAs in that category. For nucleotide composition features

(i.e., 5p_A, 3p_U), the number reflects the fraction of that nucleotide within the strand,

features clustered by average linkage.

1054

Main Figure 5: Comparative Analysis of Mature miRNA Duplex Features between

C. elegans **and** *H. sapiens*. **(A)** Frequency distribution of predicted *5p Singleton*

confidence across all miRNAs in *H. sapiens. 3p Singletons* (yellow) are defined as

miRNAs falling below a predicted 5p confidence of 20% or below; *Switchers* are defined

as miRNAs with a predicted 5p confidence between 20% and 80%; *5p Singletons* (red)

are defined as miRNAs with a predicted 5p confidence of 80% or greater. **(B)** The four

most significant ($p<0.05$) structural differences between *H. sapiens* (left) and *C. elegans*

(right) mature miRNA duplexes. From top left to bottom right: mean occurrence of

guanine in the 5p strand (5p_G), mean occurrence of guanine-guanine dinucleotide in

either mature strand (di_GG), mean occurrence of cytosine-cytosine dinucleotide in

either mature strand (di_CC), binary classifier for presence of cytosine-cytosine-uracil

trinucleotide in the 3p strand (3p_has_CCU). **(C)** Alignment of *C. elegans* (*cel*) *let-7* and

five *H. sapiens* (*hsa*) *let-7* homologs shows differences in CCU occurrence within the 3p

strand (red boxes). **(D)** Heatmap of differences in mean value for each of the 77 model

features between each category of miRNA in *C. elegans* (c) and *H. sapiens* (h). For

binary features (i.e., 3p_has_CCU, which is either 0 or 1), the number represents the

proportion (e.g., 0.3 = 30% of miRNAs in that category have a CCU motif in the 3p

strand). For continuous features (i.e., gc_content, au_content), the number is the

average value across all miRNAs in that category. For nucleotide composition features

(i.e., 5p_A, 5p_U), the number reflects the fraction of that nucleotide within the strand.

Features clustered by average linkage.

1076

Main Figure 6: Model Accuracy for the Human miRNA Tissue Atlas Dataset. **(A)**

Model accuracy expressed as a percentage of correct predictions for four different

subsets of *H. sapiens* miRNAs (total 854 mature duplexes). From left to right, ranked by

total expression of mature 5p and 3p strand across all tissue types: predicted accuracy

among top 10 (9 correct, 1 incorrect), top 50 (33 correct, 17 incorrect), top 100 (66

correct, 34 incorrect), and all (502 correct, 352 incorrect) miRNAs in this dataset. **(B)**

Confusion matrix of the strand usage prediction model among the top 50 most highly

expressed *H. sapiens* miRNAs. Summarizes differences in 5p and 3p strand usage

between experimentally validated (miRNA Atlas) and model-derived (ML Model) data.

(C) Heatmap of prediction accuracy, either correct (red) or incorrect (blue) for the top 50

most highly expressed *H. sapiens* miRNAs across 65 *H. sapiens* tissue types and

organs. Data are clustered using average linkage.

1089

Supplemental Information titles and legends

Supplemental Table S1: **Primers used in this study.**

Supplemental Table S2: **HiTmiSS assay results.**

Supplemental Table S3: **Features used in our ML model and their weights.**

Supplemental Table S4: **ML prediction in** ***C. elegans***.

Supplemental Table S5: **ML prediction in vertebrates.**

Supplemental Table S6: **Comparison of ML model with miRNA tissue Atlas.**

1097

References

Uncategorized References

1.   Bartel, D.P. (2018) Metazoan MicroRNAs. *Cell*, **173**, 20-51.

2.   Wolter, J.M., Le, H.H., Linse, A., Godlove, V.A., Nguyen, T.D., Kotagama, K., Lynch,

A., Rawls, A. and Mangone, M. (2017) Evolutionary patterns of metazoan microRNAs

reveal targeting principles in the let-7 and miR-10 families. *Genome Res*, **27**, 53-63.

3.   Pare, J.M., Tahbaz, N., Lopez-Orozco, J., LaPointe, P., Lasko, P. and Hobman, T.C.

(2009) Hsp90 regulates the function of argonaute 2 and its recruitment to stress granules

and P-bodies. *Mol Biol Cell*, **20**, 3273-3284.

4.   Nakanishi, K. (2016) Anatomy of RISC: how do small RNAs and chaperones activate

Argonaute proteins? *Wiley Interdiscip Rev RNA*, **7**, 637-660.

5.   Meijer, H.A., Smith, E.M. and Bushell, M. (2014) Regulation of miRNA strand selection:

follow the leader? *Biochem Soc Trans*, **42**, 1135-1140.

6.   Medley, J.C., Panzade, G. and Zinovyeva, A.Y. (2021) microRNA strand selection:

Unwinding the rules. *Wiley Interdiscip Rev RNA*, **12**, e1627.

7.   Frank, F., Sonenberg, N. and Nagar, B. (2010) Structural basis for 5'-nucleotide base-

specific recognition of guide RNA by human AGO2. *Nature*, **465**, 818-822.

8.   Melendez, A. and Levined, B. (August 24, 2009) In Community, T. C. e. R. (ed.),

WormBook. WormBook.

9.   Zinovyeva, A.Y., Veksler-Lublinsky, I., Vashisht, A.A., Wohlschlegel, J.A. and Ambros,

V.R. (2015) Caenorhabditis elegans ALG-1 antimorphic mutations uncover functions for

Argonaute in microRNA guide strand selection and passenger strand disposal. *Proc Natl*

Acad Sci U S A, **112**, E5271-5280.

10.   Caceres, J.F., Stamm, S., Helfman, D.M. and Krainer, A.R. (1994) Regulation of

alternative splicing in vivo by overexpression of antagonistic splicing factors. *Science*,

265, 1706-1709.

11.   Mangone, M., Macmenamin, P., Zegar, C., Piano, F. and Gunsalus, K.C. (2008)

UTRome.org: a platform for 3'UTR biology in C. elegans. *Nucleic Acids Res*, **36**, D57-

62.

12.   Mangone, M., Manoharan, A.P., Thierry-Mieg, D., Thierry-Mieg, J., Han, T.,

Mackowiak, S.D., Mis, E., Zegar, C., Gutwein, M.R., Khivansara, V. *et al.* (2010) The

landscape of C. elegans 3'UTRs. *Science*, **329**, 432-435.

13.   Steber, H.S., Gallante, C., O'Brien, S., Chiu, P.L. and Mangone, M. (2019) The C.

elegans 3' UTRome v2 resource for studying mRNA cleavage and polyadenylation, 3'-

UTR biology, and miRNA targeting. *Genome Res*, **29**, 2104-2116.

14.   Murari, E., Meadows, D., Cuda, N. and Mangone, M. (2024) A comprehensive analysis

of 3'UTRs in Caenorhabditis elegans. *Nucleic Acids Res*.

15.   Sulston, J.E. (1983) Neuronal cell lineages in the nematode Caenorhabditis elegans. *Cold*

Spring Harb Symp Quant Biol, **48 Pt 2**, 443-452.

16.   Frokjaer-Jensen, C., Davis, M.W., Hopkins, C.E., Newman, B.J., Thummel, J.M., Olesen,

S.P., Grunnet, M. and Jorgensen, E.M. (2008) Single-copy insertion of transgenes in

Caenorhabditis elegans. *Nat Genet*, **40**, 1375-1383.

17.   Kozomara, A., Birgaoanu, M. and Griffiths-Jones, S. (2019) miRBase: from microRNA

sequences to function. *Nucleic Acids Res*, **47**, D155-D162.

1142 18. Kotagama, K., Schorr, A.L., Steber, H.S. and Mangone, M. (2019) ALG-1 Influences

1143    Accurate mRNA Splicing Patterns in the Caenorhabditis elegans Intestine and Body

1144    Muscle Tissues by Modulating Splicing Factor Activities. *Genetics*, **212**, 931-951.

1145 19. Mei, Q., Li, X., Meng, Y., Wu, Z., Guo, M., Zhao, Y., Fu, X. and Han, W. (2012) A

1146    facile and specific assay for quantifying microRNA by an optimized RT-qPCR approach.

1147    *PloS one*, **7**, e46890.

1148 20. Livak, K.J. and Schmittgen, T.D. (2001) Analysis of relative gene expression data using

1149    real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods*, **25**, 402-408.

1150 21. Kato, M., Kashem, M.A. and Cheng, C. (2016) An intestinal microRNA modulates the

1151    homeostatic adaptation to chronic oxidative stress in C. elegans. *Aging (Albany NY)*, **8**,

1152    1979-2005.

1153 22. Kent, W.J. (2002) BLAT--the BLAST-like alignment tool. *Genome Res*, **12**, 656-664.

1154 23. Lorenz, R., Bernhart, S.H., Honer Zu Siederdissen, C., Tafer, H., Flamm, C., Stadler, P.F.

1155    and Hofacker, I.L. (2011) ViennaRNA Package 2.0. *Algorithms Mol Biol*, **6**, 26.

1156 24. Rishik, S., Hirsch, P., Grandke, F., Fehlmann, T. and Keller, A. (2025)

1157    miRNATissueAtlas 2025: an update to the uniformly processed and annotated human and

1158    mouse non-coding RNA tissue atlas. *Nucleic Acids Res*, **53**, D129-D137.

1159 25. Brosnan, C.A., Palmer, A.J. and Zuryn, S. (2021) Cell-type-specific profiling of loaded

1160    miRNAs from Caenorhabditis elegans reveals spatial and temporal flexibility in

1161    Argonaute loading. *Nat Commun*, **12**, 2194.

1162 26. Shaw, W.R., Armisen, J., Lehrbach, N.J. and Miska, E.A. (2010) The conserved miR-51

1163    microRNA family is redundantly required for embryonic development and pharynx

1164    attachment in Caenorhabditis elegans. *Genetics*, **185**, 897-905.

27. Alvarez-Saavedra, E. and Horvitz, H.R. (2010) Many families of C. elegans microRNAs

are not essential for development or viability. *Curr Biol*, **20**, 367-373.

28. Garcia-Segura, L., Abreu-Goodger, C., Hernandez-Mendoza, A., Dimitrova Dinkova,

T.D., Padilla-Noriega, L., Perez-Andrade, M.E. and Miranda-Rios, J. (2015) High-

Throughput Profiling of Caenorhabditis elegans Starvation-Responsive microRNAs. *PloS*

one, **10**, e0142262.

29. Garcia, S.M., Tabach, Y., Lourenço, G.F., Armakola, M. and Ruvkun, G. (2014)

Identification of genes in toxicity pathways of trinucleotide-repeat RNA in C. elegans.

Nature structural & molecular biology, **21**, 712-720.

30. Kagias, K., Podolska, A. and Pocock, R. (2014) Reliable reference miRNAs for

quantitative gene expression analysis of stress responses in Caenorhabditis elegans. *BMC*

Genomics, **15**, 222.

31. Blazie, S.M., Geissel, H.C., Wilky, H., Joshi, R., Newbern, J. and Mangone, M. (2017)

Alternative Polyadenylation Directs Tissue-Specific miRNA Targeting in Caenorhabditis

elegans Somatic Tissues. *Genetics*, **206**, 757-774.

32. Haenni, S., Ji, Z., Hoque, M., Rust, N., Sharpe, H., Eberhard, R., Browne, C., Hengartner,

M.O., Mellor, J., Tian, B. *et al.* (2012) Analysis of C. elegans intestinal gene expression

and polyadenylation by fluorescence-activated nuclei sorting and 3'-end-seq. *Nucleic*

Acids Res, **40**, 6304-6318.

33. Pandit, A., Jain, V., Kumar, N. and Mukhopadhyay, A. (2014) PHA-4/FOXA-regulated

microRNA feed forward loops during Caenorhabditis elegans dietary restriction. *Aging*

(Albany NY), **6**, 835-855.

34.   Brunquell, J., Snyder, A., Cheng, F. and Westerheide, S.D. (2017) HSF-1 is a regulator of

1188   miRNA expression in Caenorhabditis elegans. *PloS one*, **12**, e0183445.

1189   35.   Nakanishi, K. (2024) When Argonaute takes out the ribonuclease sword. *J Biol Chem*,

1190   **300**, 105499.

1191   36.   Krishnan, K., Steptoe, A.L., Martin, H.C., Wani, S., Nones, K., Waddell, N.,

1192   Mariasegaram, M., Simpson, P.T., Lakhani, S.R., Gabrielli, B. *et al.* (2013) MicroRNA-

1193   182-5p targets a network of genes involved in DNA repair. *RNA*, **19**, 230-242.

1194   37.   Krist, B., Florczyk, U., Pietraszek-Gremplewicz, K., Jozkowicz, A. and Dulak, J. (2015)

1195   The Role of miR-378a in Metabolism, Angiogenesis, and Muscle Biology. *Int J*

1196   *Endocrinol*, **2015**, 281756.

1197   38.   Rane, S., He, M., Sayed, D., Vashistha, H., Malhotra, A., Sadoshima, J., Vatner, D.E.,

1198   Vatner, S.F. and Abdellatif, M. (2009) Downregulation of miR-199a derepresses

1199   hypoxia-inducible factor-1alpha and Sirtuin 1 and recapitulates hypoxia preconditioning

1200   in cardiac myocytes. *Circ Res*, **104**, 879-886.

1201   39.   Makeyev, E.V., Zhang, J., Carrasco, M.A. and Maniatis, T. (2007) The MicroRNA miR-

1202   124 promotes neuronal differentiation by triggering brain-specific alternative pre-mRNA

1203   splicing. *Mol Cell*, **27**, 435-448.

1204   40.   Rhim, J., Baek, W., Seo, Y. and Kim, J.H. (2022) From Molecular Mechanisms to

1205   Therapeutics: Understanding MicroRNA-21 in Cancer. *Cells*, **11**.

1206   41.   Brueckner, B., Stresemann, C., Kuner, R., Mund, C., Musch, T., Meister, M., Sultmann,

1207   H. and Lyko, F. (2007) The human let-7a-3 locus contains an epigenetically regulated

1208   microRNA gene with oncogenic function. *Cancer Res*, **67**, 1419-1423.

42. Kim, H., Kim, J., Yu, S., Lee, Y.Y., Park, J., Choi, R.J., Yoon, S.J., Kang, S.G. and Kim,

V.N. (2020) A Mechanism for microRNA Arm Switching Regulated by Uridylation. *Mol*

Cell, **78**, 1224-1236 e1225.

43. Mi, S., Cai, T., Hu, Y., Chen, Y., Hodges, E., Ni, F., Wu, L., Li, S., Zhou, H., Long, C. *et*

al. (2008) Sorting of small RNAs into Arabidopsis argonaute complexes is directed by

the 5' terminal nucleotide. *Cell*, **133**, 116-127.

44. Okamura, K., Liu, N. and Lai, E.C. (2009) Distinct mechanisms for microRNA strand

selection by Drosophila Argonautes. *Mol Cell*, **36**, 431-444.

45. Kotagama, K., Grimme, A.L., Braviner, L., Yang, B., Sakhawala, R.M., Yu, G., Benner,

L.K., Joshua-Tor, L. and McJunkin, K. (2024) Catalytic residues of microRNA

Argonautes play a modest role in microRNA star strand destabilization in C. elegans.

Nucleic Acids Res, **52**, 4985-5001.

46. Stein, L., Mangone, M., Schwarz, E., Durbin, R., Thierry-Mieg, J., Spieth, J. and

Sternberg, P. (2001) WormBase: network access to the genome and biology of

Caenorhabditis elegans (vol 29, pg 82, 2001). *Nucleic Acids Research*, **29**, 1012.

1224

A

B

miR-52

miR-75

C

Variant	5p	3p
WT	0.85	0.15
UP11-12	0.71	0.29
5'U → C	0.71	0.29
Combined	0.68	0.32

D

Main Figure 5

A Model Accuracy by Expression Rank

B Confusion Matrix Top 50 – miRNA Atlas vs ML Model

C Model Accuracy – Top 50 miRNAs across 65 *H. sapiens* Organ Systems

correct prediction

incorrect prediction