An AI-Guided Framework Reveals Conserved Features Governing

microRNA Strand Selection

Author list

Dalton Meadows[1,2],

Hailee Hargis[1,2],

Amanda Ellis[1,2],

Heewook Lee[1],

Marco Mangone[1,*]

Author affiliations

[1)] The Biodesign Institute at Arizona State University, 1001 S McAllister Ave, Tempe, AZ

[2)] School of Life Sciences, Arizona State University, 427 E Tyler Mall, Tempe, AZ

Author list footnotes

[*)] Corresponding author

Corresponding author and e-mail address

Marco Mangone; mangone@asu.edu

Abstract

MicroRNAs (miRNAs) are central regulators of gene expression, yet how cells choose between the two strands (5p or 3p) of a miRNA duplex during biogenesis remains unresolved. Here, we present a comprehensive, experimentally grounded framework that decodes the logic of miRNA strand selection. Using *Caenorhabditis elegans* as a model system, we developed a high-throughput qPCR platform enabling precise quantification of strand usage across developmental stages and in specific somatic tissues. To uncover the molecular grammar guiding this process, we built a predictive machine learning model trained on experimentally validated strand usage data. This AI-driven model, integrating 77 biologically informed features, accurately predicts strand preference not only in nematodes but also across vertebrates, including humans, revealing compositional and structural biases that are conserved yet functionally repurposed. Our analysis shows that strand selection is not stochastic but follows conserved, context-dependent rules shaped by cellular and developmental cues. To support the research community, we provide open-access resources: a database of strand usage profiles, predictive scores across species, and code and protocols via GitHub. This work offers the first unified, generalizable model for miRNA strand selection, establishing a paradigm that combines large-scale experimentation with AI to reveal a conserved, programmable layer of gene regulation.

Introduction

miRNAs are ~22-nucleotide non-coding RNAs that guide the RNA-induced silencing complex (RISC) to target mRNAs, primarily through base pairing within their 3′ untranslated regions (3′UTRs) (1). Once bound, the miRNA-RISC complex (miRISC) typically represses translation of the target mRNA. Both miRNAs and their 3′UTR targets are often conserved across metazoans and play key roles in essential biological processes such as development and morphogenesis.

miRNAs have undergone several pronounced expansions in metazoan genomes (2), coinciding with evolving organismal complexity yet rarely corresponding with increases in protein-coding genes (2). Eukaryotic genomes contain 100-2,000 distinct mature miRNAs in intergenic regions, introns, and in clusters. Following transcription in long primary miRNA transcripts (pri-miRNAs) by RNA polymerase II, they are processed in the nucleus by the microprocessor complex into shorter hairpin structures known as precursor miRNAs (pre-miRNAs) (1). The microprocessor is a multiprotein complex composed of the RNAse III endonuclease Drosha (3), DGCR8/Pasha (4-6) and auxiliary factors. The two strands composing the pre-miRNAs are called the 5p and the 3p strand, respectively. Pre-mRNAs are subsequently exported to the cytoplasm by Exportin-5, where they undergo further cleavage by the enzyme Dicer, producing mature miRNA duplexes (1). Beyond the canonical Drosha-Dicer processing pathway, some miRNAs are generated through non-canonical mechanisms such as mirtron biogenesis (7,8). Mirtrons are short introns that, after splicing and debranching, fold directly into pre-miRNA hairpins and enter the miRNA pathway downstream of Drosha.

The Hsc70-Hsp90 chaperone complex facilitates the ATP-dependent loading of the

miRNA duplex into an Argonaute protein, the class of proteins that execute the silencing

step (9). This complex is thought to induce conformational changes in Argonaute that

enable duplex loading (10-12). Once loaded, one strand of this duplex, the guide strand,

is kept within the RISC, while the other, the passenger strand, is typically degraded.

Both the 5p and the 3p strands can be selected for miRISC loading and act as guide

strands. It is currently unclear how this process, termed 'miRNA strand selection,' is

executed in detail, which proteins are involved in this decision, and how the guide

strand is ultimately chosen between the two strands.

The predominant factors influencing miRNA strand selection are believed to be the

specific 5' nucleotide sequences of each strand and the comparative thermodynamic

stability exhibited by the two ends of the miRNA duplex (13). Studies performed using

the human Ago2 protein, an Argonaute protein member of the RISC complex, show that

the pocket responsible for binding the 5' nucleotide of the guide strand exhibits a strong

preference for Uracil nucleotides (14). This aligns with the prevalent occurrence of uracil

as the 5' nucleotide in miRNA guide strands (2), whereas the passenger strand

commonly initiates with a cytosine at its 5' end.

Furthermore, the miRISC complex exhibits a preference for loading whichever end

of the miRNA duplex has lower thermodynamic stability, although not always (15). The

initial four nucleotides on each end of the duplex are believed to establish this

thermodynamic asymmetry, and in some cases, a slight difference in the number of

hydrogen bonds can induce the preferential loading of the less stably paired strand *in*

vitro (12).

Consistently, the 5' ends of guide strands are enriched with purines (2). In

comparison, the 5' ends of passenger strands are predominantly pyrimidine-rich in both

human and fly miRNAs, contributing to the thermodynamic asymmetry observed

between the two ends of the duplex. In *C. elegans*, single amino acid substitutions

within the MID (G553R) and PIWI (G716E, S895F, and S939F) domains of the

Argonaute protein ALG-1 invert the strand selection process for specific miRNAs,

suggesting a potential role for this protein in the miRNA strand selection process (16).

While these mutations play a pivotal role, they do not consistently reverse strand

selection for all miRNAs, suggesting that there may be additional internal or external

factors involved in this process. Tissue-specific accessory factors and their dosage

could also be responsible for miRNA strand selection, altering strand choices based on

tissue or developmental contexts, similarly to the role of SR proteins in RNA splicing

(17). While thermodynamic and sequence features help explain strand choice, a

comprehensive mechanistic framework has been lacking.

C. elegans is an ideal model for studying miRNA strand selection due to its compact

3′UTRome (18-21), small number of miRNAs (miRbase release 22.1) (22), fully mapped

cell lineage (23), and well-characterized transcriptome. Here, we developed a high-

throughput qPCR platform to systematically profile miRNA strand selection events

across *C. elegans* development and in somatic tissues. These data revealed conserved

sequence and structural patterns, which we used to train a machine learning model that

predicts strand selection with 83% accuracy and generalizes to human miRNAs.

Materials and methods

All deionized (DI) water was obtained from either a Purelab Option system (ELGA

LabWater) or an Avidity Science system (Duo) and autoclaved before use. Chemicals

were ordered from Sigma Aldrich/Merck, VWR, or Thermo Fisher Scientific in molecular

biology grade and stored according to company instructions. Gateway^{TM} LR Clonase II

Plus^{TM} enzyme mixes, SuperScript^{TM} III Reverse Transcriptase, and PowerTrack^{TM}

SYBR Green Master Mix were purchased from Invitrogen. Primers for the HiTmiSS

assay were ordered from Life Technologies. Poly(U) Polymerase and restriction

enzymes used in validation were ordered from New England Biolabs. The pCFJ150

plasmid was a gift from Erik Jorgensen. The OP50-1 *E. coli*, N2 *C. elegans,* and

EG6699 *C. elegans* strains were purchased through the Caenorhabditis Genetics

Center.

Maintenance of *C. elegans* strains

All *C. elegans* strains were cultured on NGM plates seeded with OP50-1 bacteria. All

transgenic lines produced for experiments described in this study were maintained at

24°C. When bacteria on each NGM plate had been cleared, *C. elegans* strains were

transferred onto new plates via washing into a 15 ml Falcon tube using 15 ml M9 media

(22 mM KH_2PO_4, 42 mM Na_2HPO_4, 85.5 mM NaCl, 1 mM $MgSO_4$, sterilized by

autoclave) and pelleted for 3 minutes at 400g. The M9 media was then aspirated to a

volume of 500 ml, and the animals were resuspended via gentle pipetting before being

moved onto fresh, seeded NGM plates.

Synchronization of *C. elegans* strains

The EG6699 strain (24) was maintained at 24°C. To synchronize animals for

injection, confluent plates were bleached to isolate the embryos. Each plate was

washed into a 15 ml Falcon tube with 15 ml DI water and pelleted for 3 minutes at 400g.

The water was then aspirated, and the animals were incubated in 5 ml of bleaching

solution (0.5 M NaOH, 10% bleach) for 20 minutes with gentle end-over-end rotation.

After bleaching, the embryos were washed twice with DI water and moved onto fresh,

seeded NGM plates. For use in the HiTmiSS assay, both N2 and *pges-1::GFP::alg-*

1::unc54 3ʹ UTR C. elegans strains were synchronized as described above.

Primers design for the HiTmiSS Assay

To quantify miRNAs in high throughput, we designed 380 unique forward primers for

use in the qPCR amplification step of the HiTmiSS assay. These primers were designed

to be complementary to either the 5p strand or the 3p strand of the 190 most expressed

miRNAs in the *C. elegans* genome, corresponding to 75% of all *C. elegans* miRNA

deposited in miRbase (release 22.1) (25). We did not consider miRNAs whose

existence is not supported by broad experimental evidence, nor any miRNAs predicted

to exist. A few miRNAs were ignored because their corresponding primers had a very

low Tm, making it impossible to process them using our pipeline. All primers were

ordered in four 96-well microplates from Life Technologies pre-diluted at a concentration

of 50 mM (**Supplemental Table S1**). We designed a universal reverse primer

(GCAGGGTCCGAGGTATTC) complementary to the stem-loop sequence of the stem-loop

poly(A) primer used to prepare the cDNA for this qPCR reaction

(GTCGTATCCAGTGCAGGGTCCGAGGTATTCGCACTGGATACGACAAAAAAAAAAAAA). Twenty-four

replicates of this primer were ordered from Life Technologies in a separate array, also

at a concentration of 50 mM. Both forward and reverse primers were re-arrayed and

diluted to a working concentration of 10 mM immediately prior to use.

Singletons **and** *Switchers* **identification**

For each strand, we subtracted its relative quantity at one life stage versus the

relative quantity of the complementary strand (i.e., for *let-7* 5p, we compared the

abundance at the embryo stage versus the abundance of *let-7* 3p at the embryo stage).

This was repeated for each life stage at which the expression of that miRNA strand fell

above our significance threshold (1 standard deviation above the mean of all quantified

miRNAs at a given stage), and any miRNAs for which their expression was greater at

each such life stage were deemed *Singletons*. We classified as *5p Singletons* miRNAs

where the 5p strand was more expressed at each life stage than its corresponding 3p

strand. We classified as *3p Singletons* miRNAs where the 3p strand was more

expressed at each life stage than its corresponding 5p strand. We classified as

Switchers miRNAs that rose above our threshold of significance during at least one life

stage, but whose expression levels did not display a consistent pattern of usage of one

strand over the other.

HiTmiSS Assay

To collect sufficient total RNA for analysis of wild type miRNA, confluent NGM plates

of N2 strains were synchronized and either washed and pelleted immediately (embryo

stage) or incubated at 24°C for 15 hours (larval 1 stage), 25 hours (larval 2 stage), 33

hours (larval 3 stage), 42 hours (larval 4 stage), or 51 hours (young adult stage) before

washing and pelleting. All washes were performed in 15 ml Falcon tubes using 15 ml DI

water, and pellets were generated by centrifugation for 3 minutes at 400g, resulting in a

final pellet volume of 500 µL to 1 mL. Sufficient worms to generate this size pellet

differed by life stage, requiring 12 (larval 4 and young adult stages), 18 (larval 2 and

larval 3 stages), or 24 (embryo and larval 1 stages) starting confluent plates.

This pellet was then resuspended to a volume of 1.5 ml in ice-cold Triton X-100 lysis

buffer (150 mM NaCl, 1.0% Triton X-100, 50 mM Tris-HCl pH 8.0, 0.5 mM EDTA) and

transferred to a 2 ml Spectrum tube pre-filled with 0.5 mm zirconium beads. Using a

Benchmark BeadBug D1030 microtube homogenizer, each tube was homogenized for

90 seconds, incubated for 2 minutes on ice, then homogenized for an additional 60

seconds. The resultant lysate mixture was centrifuged for 3 minutes using a VWR

benchtop mini centrifuge, and the liquid fraction was transferred into a 1.5 ml Eppendorf

microcentrifuge tube. To minimize non-aqueous contaminants, this lysate was

centrifuged again in the same manner and transferred to a 15 ml Falcon tube pre-chilled

on ice. To isolate total RNA, we used a DirectZol RNA MiniPrep Plus kit from Zymo

Research, following kit instructions and performing the entire procedure at 4°C.

Resultant RNA was eluted in 30 μL nuclease-free (NF) H_2O.

The protocol for the HiTmiSS assay was adapted from Mei *et al.* (26). We prepared

our cDNA from each life stage for qPCR analysis via polyuridinylation using Poly(U)

Polymerase (NEB) as per manufacturer instructions followed by a 20-minute heat

inactivation at 65°C. We performed a first-strand reaction on this polyuridinylated RNA

using Superscript III Reverse Transcriptase and stem-loop poly(A) primers under

recommended conditions (Invitrogen). We then performed qPCR in high throughput

using this cDNA, SYBR Green master mix (Invitrogen), and our primers as described

above. We used Applied Biosystems QuantStudio 3 and QuantStudio 7 qPCR

machines to perform the screen. For each life stage, four 96-well plates were prepared

with 40 μL reactions in each well, along with a positive control reaction containing 10 pg

pDONR 221 ROG instead of cDNA and the necessary forward and reverse primers to

amplify *gfp.* Each plate was then re-arrayed in triplicate into a 384-well plate using a

Biomek FX^P Laboratory Automation Workstation for a total of three 10 μL reactions per

miRNA strand per life stage. An additional plate was prepared with no cDNA or plasmid

spike-in DNA as a non-template control. These qPCR reactions were carried out under

standard SYBR Green reagent conditions on an Applied Biosystems QuantStudio 7.

Relative miRNA levels were calculated using the $2^{-\Delta\Delta Ct}$ method (see below), and

normalized using the spike-in positive control. These results were then plotted

logarithmically as a heatmap, with data below our analytical threshold visualized in

black. Data were ordered by type (*5p Singleton*, *3p Singleton*, or *Switcher*) and

visualized using the Heatmapper2 web application (27).

Additional MIQE-Compliance-related methods

RNA Integrity Reporting: RNA purity and integrity were assessed using a Nanodrop

spectrophotometer (Thermo Scientific). All samples had 260/280 ratios between 1.9-2.1

and 260/230 ratios >1.8.

Primer Validation: Primer specificity was inferred from the absence of amplification

in non-template controls and the presence of a single, well-defined amplification curve

for each reaction. All primer sets produced consistent Ct values across independent

replicates and developmental stages, and no secondary amplification products were

detected in NTC reactions. The HiTmiSS assay is a high-throughput adaptation of Mei

et al. (26), who demonstrated primer specificity using stem-loop qPCR with strand-

specific forward primers. Because our assay uses identical primer architecture,

specificity is expected from design and was verified by negative controls.

Biological Replicates: For each developmental stage and tissue, RNA was extracted

from independent biological populations of worms cultured on separate plates. Each

qPCR was run in technical triplicate.

NTC Controls: Non-template controls did not produce detectable amplification

across all primer sets.

Reference Stability: *let-7* and *lin-4* were chosen as reference miRNAs based on their

stable expression across developmental stages and tissues (confirmed by low Ct

variance across samples).

Cycle Threshold Analysis of the HiTmiSS Assay

HiTmiSS assay queries each miRNA strand separately, normalized by a per-assay

positive and negative control. The qPCR results containing the cycle threshold (Ct)

values for each tested miRNA were exported in Microsoft Excel 2019 format. A

comprehensive list of these data can be found in **Supplemental Table S2**. Ct values

were averaged across experimental replicates, then subjected to a $2^{-\Delta\Delta Ct}$ analysis to

calculate initial amounts of each miRNA in the sample, consistent with experimental

standard (28). The Ct value for any given miRNA strand was subtracted from the Ct

value of the non-template control (a qPCR reaction lacking any cDNA) for that miRNA

strand. This was termed the ΔCt for that miRNA. The same was done for the positive

control for that sample. Then, for each sample, this difference was subtracted from the

difference obtained for that positive control, normalizing samples to the same positive

control and giving us the $\Delta\Delta$Ct for that miRNA strand. 5p/3p comparisons for the same

pre-miRNA, use a universal reverse primer and strand-specific forward primers matched

for Tm, length, GC content, and amplicon size. The inverse log base 2 of that $\Delta\Delta$Ct was

then calculated to approximate the initial amount of that miRNA strand present in the

cDNA relative to the positive control. This was deemed the $2^{-\Delta\Delta Ct}$ value for that miRNA

strand. To determine whether a miRNA was considered present in a sample, we

decided on a strict threshold of one standard deviation above the mean of all $2^{-\Delta\Delta Ct}$

values for that stage of development. MiRNAs quantified at levels above this threshold

were considered significant for the purposes of this study.

HiTmiSS Assay in the Intestinal Tissue

To isolate tissue for use in an intestine-specific HiTmiSS assay, we first performed

protein immunoprecipitation followed by miRNA isolation. The fluorescent

pges1::gfp::alg-1::unc54 3′UTR C. elegans strain previously prepared by our lab (29)

was grown to confluence, synchronized, and pelleted as described above for each life

stage, concurrent with N2 *C. elegans* as a negative control.

Each pellet was resuspended to 1.5 µl using our Triton X-100 lysis buffer with added

protease inhibitor (two Roche cOmplete ULTRA mini EDTA-free protease inhibitor

tablets per 50 mL lysis buffer) and transferred to a zirconium bead tube to be

homogenized as previously described. During this homogenization step, anti-GFP

magnetic antibody beads (Chromotek GFP-Trap Magnetic Agarose) were prepared by

diluting 25 µl bead slurry in 500 µl dilution buffer (150 mM NaCl, 50mM Tris-HCl pH 8.0,

0.5 mM EDTA) and rotating end-over-end at 4°C for 3 minutes.

The beads were pelleted using a benchtop centrifuge, and the buffer was aspirated

using a magnetic tube rack. 500 µl dilution buffer was added again, and this dilution

process was repeated twice. We then recovered aqueous lysate from the homogenized

mixture as previously described and diluted it using 300 µl dilution buffer. This diluted

lysate was transferred to a tube containing the prepared anti-GFP beads and left to

rotate end-over-end at 4°C overnight. To clear unbound lysate, the beads were pelleted

using a benchtop centrifuge, and the buffer was aspirated using a magnetic tube rack.

500 μl of wash buffer (150 mM NaCl, 50 mM Tris-HCl pH 8.0, 0.1% Triton X-100, 0.5

mM EDTA) was added, the beads were pelleted, and the buffer was aspirated using a

magnetic tube rack. This wash step was repeated five additional times. The beads were

then resuspended in 100 μl wash buffer and RNA was isolated using a DirectZol RNA

MiniPrep Plus kit from Zymo Research, following kit instructions and performing the

entire procedure at 4°C. Resultant RNA was eluted in 30 μL NF H_2O.

The HiTmiSS assay and subsequent analysis was performed as described above,

with the following modifications: (1) Forward primers were re-arrayed from the 96-well

microplates described above, resulting in a screen of the 93 most highly quantified

miRNAs (93 5p and 93 3p strands) from our whole worm analysis. This was done to

avoid redundant quantification of miRNAs which in whole worm fell below our

quantification threshold and were therefore unlikely to display significant expression in a

tissue-specific screen performed on immunoprecipitated miRNAs. (2) Only embryo (E)

and young adult (YA) stages were included in this analysis. (3) The analytical threshold

per life stage was set at the mean expression of all miRNA strands at that life stage,

due to globally decreased cycle threshold values detected in a single tissue relative to

our spike-in positive control used for normalization. (4) Data was ordered by overall

strand expression across both life stages, leading to only two major clusters on the

resultant heatmap. Additionally, miRNAs which fell below our analytical threshold for

both strands at each life stage were excluded from the heatmap for clarity.

qPCR of miRNAs targeting overexpressed intestinal 3′UTRs

We used two *C. elegans* strains previously generated and published by our lab to measure miRNA strand levels upon intestinal overexpression of their target 3′UTRs,. (29). Each expressed the promoter for intestine-specific gene *ges-1* fused to a dual color ROG construct (a gene encoding *mCherry* and *gfp* separated by a trans-spliceable operon) fused to the 3′ UTR of either *fox-1* or *hrp-2 (29)*. Each of these strains was grown to confluence in 12 NGM plates in mixed stage and subjected to quantification using the same approach as the HiTmiSS assay described above, though rather than quantifying every miRNA in these animals, we quantified the 5p and 3p strands of *miR-54*, *miR-60*, *miR-232*, and *miR-239b*. These quantifications were carried out in 96-well plates using cDNA (both prepared from our strains as well as control cDNA gathered from N2 *C. elegans*), SYBR Green master mix, and the appropriate forward and universal reverse primers for the miRNAs of interest, re-arrayed from our standard HiTmiSS plates. Each 96-well plate contained two positive controls in *let-7* and *lin-4*, as well as a non-template negative control for each set of primers. These qPCRs were carried out in experimental triplicate under standard SYBR Green reagent conditions on an Applied Biosystems QuantStudio 3.

Relative miRNA levels were calculated using the $2^{-\Delta\Delta Ct}$ method as described above and normalized using the average of *let-7* and *lin-4* expression for each sample. These results were then averaged across two replicates and analyzed using Microsoft Excel 2019. Data were visualized as $\log_2$ of the change in expression from wild-type N2

negative control to either of our experimental strains. The threshold of significance was

set at one as per standard analytical procedure.

Comparison of HiTmiSS vs RNA-seq

To compare HiTmiSS results with RNA-seq data, we retrieved small RNA sequencing

datasets from Kato et al. (30) and Stubna et al (31) via the Sequence Reads Archive

(SRA). Both studies employed deep sequencing (Illumina) to profile the expression of

small RNA and miRNA across *C. elegans* developmental stages. For Kato et al., we

analyzed the following datasets: SRR014966 (embryo), SRR014967 (L1), SRR014968

(L2), SRR014969 (L3), SRR014970 (L4), and SRR014971 (young adult). For Stubna et

al., we analyzed the following datasets: L1: SRR29013898 (rep1), and SRR29013897

(rep2). L2: SRR29013894 (rep1), and SRR29013893 (rep2). L3: SRR29013890 (rep1),

and SRR29013889 (rep2). L4: SRR29013889 (rep1), and SRR29013885 (rep2). YA:

SRR29013882 (rep1), and SRR29013881 (rep2). Early Embryo: SRR29013881 (rep1),

and SRR29013905 (rep2). Late Embryo: SRR29013902 (rep1) and SRR29013901

(rep2). The early and late embryos were pooled together and averaged for the analysis

described below. To assess both the presence of miRNA and its strand usage, we

developed a custom analysis pipeline. Sequencing reads were first converted to FASTA

format using the FASTX-Toolkit (http://hannonlab.cshl.edu/fastx_toolkit/), then aligned

using BLAT (32) against a reference database containing all known *C. elegans* miRNA

strands (miRbase) (25) in FASTA format, with the following parameters: /blat -t=dna -

q=dna -tileSize=6 -minIdentity=10 -minMatch=1 -minScore=10 stage_specific.fa

stage_specific.psl. To increase stringency, we filtered the resulting PSL files to retain

only matches with strand = "+", and with Q_start = 0 (for Kato et al.) and 100% match

(For Stubna et al.,) then quantified the occurrence of each miRNA strand at each

developmental stage, with at least 50 reads. For the Kato et al. dataset, miRNA strands

with ≥50 reads were considered for analysis. The numbers of unique 5p and 3p strands,

together with their corresponding median read counts, were as follows: Embryo, 158

(5p) and 147 (3p), totaling 305 unique miRNA strands (median = 139.5 reads); L1, 81

(5p) and 87 (3p), totaling 168 unique miRNA strands (median = 359 reads); L2, 85 (5p)

and 82 (3p), totaling 167 unique miRNA strands (median = 373.5 reads); L3, 74 (5p)

and 74 (3p), totaling 148 unique miRNA strands (median = 483 reads); L4, 67 (5p) and

81 (3p), totaling 148 unique miRNA strands (median = 589.5 reads); and YA, 70 (5p)

and 82 (3p), totaling 152 unique miRNA strands (median = 507 reads). For the Stubna

et al. dataset, a consistent set of 192 (5p) and 190 (3p) strands was identified across all

stages. The median read counts per replicate were as follows: Early embryo, 6,501

(rep1) and 5,089 (rep2); Late embryo, 7,805 (rep1) and 7,630 (rep2); L1, 5,362 (rep1)

and 6,210 (rep2); L2, 5,310 (rep1) and 7,390 (rep2); L3, 9,187 (rep1) and 8,782 (rep2);

L4, 11,605 (rep1) and 11,313 (rep2); and YA, 16,194 (rep1) and 30,042 (rep2). We then

used these counts to calculate the expression value of each miRNA strand normalized

to each developmental stage.

Structural Analysis of miRNAs

The experimentally validated *5p* and *3p Singletons* were obtained using our

HiTmiSS assay. Each entry included the entire hairpin sequence and its predicted

secondary structure in dot-bracket notation (33) (**Supplemental Table S2)**. To focus

specifically on mature miRNA regions, only the sequences of each mature miRNA

duplex were used for all downstream analyses. Since the *5p* and *3p Singleton* datasets

contained different numbers of miRNAs, we balanced them before analysis by

undersampling the larger group to match the sample size of the smaller class, ensuring

unbiased statistical comparison between 5p and 3p strands.

We extracted 77 numerical features from each *5p* and *3p Singleton* (**Supplemental**

Table S3). The extracted features included: Base composition (fraction of A, U, G, C),

motif presence (e.g. UGU, GAG, CCU), positional characteristics (first/last nucleotide

identity), dinucleotide frequencies (e.g., di_GG, di_CC), GC content and GC-related

asymmetry metrics (GC_diff, GC_ratio_5p3p), strand lengths, and number of

mismatches (unpaired positions in the dot-bracket structure). The full list of the features

and their descriptions is shown in **Supplemental Table S3**.

Feature calculations were implemented in Python using regular expressions and

sequence statistics functions. Structural mismatch counts were determined by counting

unpaired characters within the uppercase-aligned mature duplex. Comparative

statistical analysis between 5p and 3p strands was conducted using Welch's two-

sample t-test. All features were tested independently, and results were ranked by p-

value. Features with $p < 0.05$ were considered statistically significant. Bar plots were

generated for each significant feature using the mean ± standard deviation across

strands.

miRNA strand selection ML model

    To predict dominant strand selection (5p vs. 3p) in miRNA hairpins, we utilized

ChatGPT GPT-4o (OpenAI) to assist in developing and training a Random Forest

classifier based on 77 biologically curated features, organized into functional categories

capturing distinct structural and sequence-related aspects of miRNA hairpins. These

features capture thermodynamic stability, base-pairing patterns, and positional

relationships within the miRNA hairpin. The model was trained on our validated *5p* and

3p Singletons identified using our HiTmiSS assay, and feature values were computed

using custom Python scripts that parse dot-bracket RNA structures and extract both

absolute and relative measures for each strand. The weights of the features were

optimized using known *Singletons*. The final model incorporates the following features,

divided into five categories. The first category is the *Dinucleotide Composition* features,

which quantify the frequency of specific two-nucleotide motifs within the 5p and 3p

regions, capturing local sequence patterns that can influence strand selection and

structural flexibility. The second category is the *First/Last Nucleotide Bias* features,

which record the identity of the first and last nucleotides in each strand, reflecting known

preferences in Argonaute loading where terminal nucleotides strongly impact strand

choice. The third category is the *GC Content and Entropy* features, which assess the

global nucleotide composition and thermodynamic diversity within each strand, including

overall GC percentages, GC ratios between strands, and Shannon entropy measures

that reflect structural variability and rigidity. The fourth category is the *Base-pairing*

Properties features, which describe the fraction of nucleotides engaged in base pairing

within the 5p and 3p regions, providing a measure of local hairpin stability and

asymmetry. Finally, the *General Structural* features, which include the overall hairpin

length and the length difference between the 5p and 3p strands, capture broader

geometrical properties that may indirectly affect miRNA maturation and strand retention.

The complete set of features, their description, and the weight used are shown in

Supplemental Table S3. The model showed high performance in distinguishing 5p and

3p usage from a pool of known *Singletons* as a test dataset (*5p Singleton*: accuracy

100%, precision 75%, recall 100%; *3p Singleton:* 82% accuracy, 100% precision, 82%

recall). The ML model uses a Random Forest classifier implemented in scikit-learn

(v1.3.1). Model hyperparameters were optimized via grid search with 5-fold stratified

cross-validation, tuning: n_estimators: [100, 250, 500], max_depth: [5, 10, None],

min_samples_split: [2, 5, 10], class_weight: ['balanced', None]. The final selected

parameters were: RandomForestClassifier (n_estimators=250, max_depth=None,

min_samples_split=2, class_weight='balanced'). To prevent overfitting and enhance

interpretability**,** recursive feature elimination (RFE) with cross-validation was used to

rank features by importance. Features with near-zero variance and multicollinearity

(|Pearson's r| > 0.9) were removed. The model consistently retained ~45–55 features,

depending on the species subset. Performance was assessed on independent held-out

test sets not used during training. Metrics included: Accuracy: 0.85 (*C. elegans*), 0.83

(human), Precision (5p/3p)**:** 0.87 / 0.80, Recall (5p/3p)**:** 0.89 / 0.78, F1 Score (5p/3p)**:**

0.88 / 0.79**,** AUROC**:** 0.91 (*C. elegans*), 0.89 (human). Class balance was maintained

using SMOTE oversampling to correct for the slight enrichment of 5p strands in the

source datasets. We confirmed that the model generalizes well by testing it on separate

datasets (not used during training or tuning), including across species (e.g., training

in *C. elegans*, testing in humans).

Unsupervised benchmark of strand-selection predictors.

We evaluated three predictors on the same set of human miRNA duplexes (n = 854): (i)

HiTmiSS (77-feature RF; outputs 5p/3p and P(5p)); (ii) a 5′-nucleotide rule (guide

predicted by first-base preference U > A > C > G; ties to 5p); and (iii) an end-stability

proxy (ΔΔG) that scores the first 5 nucleotides at each arm's 5′ end against the reverse-

complement of the opposite arm (GC=3, AU=2, GU=1, mismatch=0), predicting the arm

with the less-stable 5′ end. Because a comprehensive gold standard is unavailable, we

tested the three predictions using the Dawid-Skene label-aggregation model, which

jointly infers (a) a latent "true" label for each duplex from the pattern of agreements and

(b) a confusion matrix (reliability) for each predictor. We report each method's estimated

accuracy as the prior-weighted proportion of correct calls from its inferred confusion

matrix. All parameters were prespecified (no tuning), and the same items (n=854) were

used for all methods.

ML model predictions in human miRNAs

Human miRNA expression data were obtained from a harmonized dataset from the

human miRNA tissue atlas (34) containing strand-specific Reads Per Million Mapped

(RPMM) values across 65 healthy tissues and cell lines of human origins. Each sample

contained expression values for both the *5p* and *3p* arms of annotated miRNA hairpins,

together with metadata fields specifying Biotype, Organ_system, and Tissue. For every

miRNA-tissue pair, the total abundance was computed as the sum

of *5p* and *3p* expression values. To reduce noise from low-coverage observations,

entries with total expression below the 25th percentile of the global abundance

distribution were excluded prior to classification. Within each tissue, the relative

contribution of the *5p* and *3p* arms was determined as:

$$f_{5p} = \frac{E_{5p}}{E_{5p} + E_{3p}}, f_{3p} = 1 - f_{5p}.$$

486
A miRNA was designated as: *5p-Singleton* if $f_{5p} \geq 0.90$ and $f_{3p} < 0.90$, *3p-Singleton*

if $f_{3p} \geq 0.90$ and $f_{5p} < 0.90$, and *Switcher* otherwise. A 90% strand-dominance

threshold was applied uniformly to all tissues. Each tissue-specific matrix was converted

into a long-format table listing the dominant strand for every miRNA across organs and

cell types. A global classification for each miRNA was then obtained by majority vote

among all tissues and cell types that met the expression threshold. Concordance

between our ML predictions and atlas-derived classifications was assessed across all

miRNAs classified as *5p-* or *3p- Singletons* (excluding *Switchers*). Overall accuracy was

computed as:

$$\text{Accuracy} = \frac{\text{Number of matching 5p or 3p calls}}{\text{Total number of 5p/3p miRNAs compared}} \times 100.$$

496
At the 90% atlas cutoff, the observed global agreement between experimental and ML

model-predicted strand identity was 68.81% (median). All analyses were conducted

in Python 3.10 using pandas 2.2, numpy 1.26, and matplotlib 3.9 within a Google

Colab environment.

Results

A novel High-Throughput (HT) qPCR approach to study miRNA biogenesis

Identifying miRNAs in somatic tissues is challenging due to their small size, low

abundance, and short half-life. To overcome this, we developed a high-throughput

qPCR method that reliably quantifies miRNA expression and distinguishes 5p from 3p

strand usage. This method is a high-throughput adaptation and general improvement of

an existing methodology (26), which uses the polyuridine polymerase enzyme to

elongate a polyU tail to all RNA transcripts in a given sample. This step extends the

length of each miRNA present in the sample, which is then used as a template in

reverse transcriptase reactions using a specially engineered universal stem-loop

reverse primer (SL-polyA) (**Main Fig. 1A**).

We named this method High-Throughput miRNA Strand Selection (HiTmiSS) assay.

HiTmiSS assay enables rapid, precise quantification of the full miRNAome and

accurately distinguishes 5p from 3p strand expression (**Main Fig. 1**).

We employed this assay to systematically quantify miRNA occurrence and the

usage of 5p and 3p strands in known miRNAs throughout *C. elegans* development. We

extracted total RNA from *C. elegans* at six developmental stages (embryos, larval

stages 1-4, and young adults) and analyzed the miRNA transcriptome and strand

selection for 190 miRNAs, representing 75% of all miRNAs annotated in miRbase (25),

using 380 unique forward primers across four plates (**Supplemental Table S1**).

We performed 7,980 qPCR experiments independently, querying these 190 miRNAs

throughout development in triplicate (1,140 experiments at each stage) (**Main Fig. 1B,**

Supplemental Fig. S1). Using stringent filters (see Materials and Methods), we

detected the presence of 177 of these miRNAs (93.2%) in at least one developmental

stage (**Supplemental Table S2**). The most significant number of miRNAs occurred

during the L4 stage (81%), followed by young adult (75.5%), L3 (65%), L2 (56.8%), and

lastly, embryo and L1 (41.6% each) stages. This aligned with previous literature

suggesting that many *C. elegans* miRNAs are tissue-specific, only becoming more

abundant upon the development of those tissues later in the developmental cycle (35).

As previously reported, miRNAs such as the key regulators of embryonic lethality,

miR-35 and *miR-51* families, were most abundant during embryonic and early larval

stages (36). In contrast, *let-7* and *lin-4*, which control adult phenotypes, peaked at the

L4 and young adult stages (37). Notably, 13 miRNAs, including *miR-791*, *miR-792*, and

miR-8206, showed no significant expression, likely due to low abundance or being

mistakenly predicted to exist.

HiTmiSS Enhances miRNA Resolution

To benchmark the performance of HiTmiSS, we compared it to Illumina RNA-seq

datasets from Kato et al. (30) and Stubna et al. (31), which profile miRNA expression

across all *C. elegans* developmental stages (**Supplemental Fig. S2-S4**). HiTmiSS

captured 88% of the miRNA strands identified by Kato et al., and 87% identified by

Stubna et al., (overlap with both datasets = 81.6%), demonstrating strong overall

concordance (**Supplemental Fig. S2A)**. Importantly, HiTmiSS more clearly resolved the

top-expressed miRNAs, particularly during early development, and provided sharper

5p/3p expression profiles across stages (**Supplemental Fig. S2B**). These

improvements likely stem from the usage of strand-specific qPCR primers and a

standardized workflow that minimizes ligation and mapping biases. Kato et al., and

Stubna et al., display broader contrasts across developmental stages (**Supplemental**

Fig. S2C), reflecting the wider dynamic range that is typical of sequencing-based

quantification. In comparison, HiTmiSS values are more compressed, but they reveal

smoother and more gradual transitions between adjacent stages, which is characteristic

of targeted qPCR-based measurements. Despite these platform-related differences, the

stage-to-stage expression patterns are highly consistent across all three datasets. Many

embryonic miRNAs transition from low (blue) to high (red) levels as worms progress

through larval development, while a distinct subset instead peaks in young adults (YA).

The distributions of values overlap almost perfectly after normalization, and the

developmental trajectories of canonical miRNAs (*let-7*, *lin-4*, *miR-35* cluster, etc.) match

very closely between all three datasets (**Supplemental Fig. S3**). Density plots show

distinct global expression distributions for each dataset (**Supplemental Fig. S4**).

Stubna et al. is enriched for low-expressed miRNAs, Kato et al. displays a tighter peak

of moderate expression, and HiTmiSS covers the widest dynamic range with extended

low and high tails, reflecting methodological differences between RNA-seq and targeted

HiTmiSS quantification (**Supplemental Fig. S4A**). Boxplots analysis indicates

comparable medians across platforms, but HiTmiSS exhibits a broader interquartile

range and whiskers, consistent with increased sensitivity toward both low- and high-

abundance strands relative to RNA-seq (**Supplemental Figs. S4B-C**). Importantly,

most HiTmiSS-defined *5p-* and *3p-Singletons* are reproduced in Stubna et al. (73-88%

of 5p and 62-79% of 3p), and developmental profiles of representative miRNAs (e.g.,

let-7, *lin-4*, *miR-51*, *miR-42*, *lys-6*, *miR-82*) illustrate clear strand dominance and stage-

specific dynamics, highlighting the resolution of the HiTmiSS platform (**Supplemental**

Fig. S4D).

Together, these results show that HiTmiSS faithfully recapitulates the developmental

miRNA landscape captured by RNA-seq, but with enhanced clarity at the strand and

stage level. Rather than serving as a redundant measurement, HiTmiSS adds

resolution: it sharpens 5p/3p distinctions, smooths developmental transitions, and

identifies strand dominance in agreement with independent datasets. Thus, RNA-seq

provides breadth, but HiTmiSS delivers precision, making it an ideal platform for

quantitative, strand-specific miRNA profiling in defined biological contexts.

Many miRNAs show consistent strand preference throughout development

Many miRNAs identified consistently use either the 5p or 3p strand as the guide

strand, which we termed '*Singletons*', while others switched between 5p and 3p usage

across different developmental stages. We named the second category '*Switchers*'.

The screen produced hits for 177 miRNAs, which we divided into *5p Singletons*, *3p*

Singletons, or *Switchers* (**Main Fig. 1B and Supplemental Table S2**).

The largest group identified was miRNA *Switchers* (46.3%, 88 hits). These include

members of the *miR-2* family and other regulators of developmental timing and stress

responses (37,38). The remaining hits were *Singletons*, with a notable bias: *3p*

Singletons (58 hits, 30.5%) outnumbered *5p Singletons* by nearly twofold. These

include *miR-35* and *miR-51* (required for embryonic development), *miR-44*

(stress/starvation), and *miR-58* (locomotion and dauer entry) (36,37,39). Several

abundant *3p Singletons*, including *miR-60*, *miR-70*, and *miR-90*, remain functionally

uncharacterized but are linked to stress response (30,38,40).

5p Singletons were the least common group, with 31 hits (16.3%), including the well-

studied families *let-7* and *lin-4*, which are key regulators of developmental timing (**Main**

Fig. 1B). Also detected were *miR-51* (embryonic lethality) and *miR-64* (heat shock

response) (36,40). While some families (*let-7* and *miR-64*) consistently use the same

strand, others span multiple categories. For instance, members of the *miR-51* seed

family (*miR-51* to *miR-56*) and *lsy-6* exhibited consistent strand usage, with *miR-51 to*

miR-53 using the 5p strand and *miR-54* to *miR-56* and *lsy-6* using the 3p strand, yet

none showed *Switcher* behavior. Despite sharing the same seed sequence

(ACCCGUA), *miR-53* and *miR-54* are processed from opposite arms and are classified

into distinct families (mirGeneDB), underscoring that convergent evolution resulted in

miRNAs sharing a seed despite different evolutionary origins.

Importantly, several hits produced by the HiTmiSS assay aligned with past findings

on the strand preference of well-characterized miRNAs such as *let-7, lin-4* (known *5p*

Singletons), and the *miR-35* family (known *3p Singletons*) (37).

To account for the strong strand preference observed among *Singletons*, we took

the top 20 *5p Singletons* and the top 20 *3p Singletons*. We compared the binding

signature and nucleotide composition between the strand that is used versus the strand

that is discarded (**Main Fig. 1C, i and ii**). Interestingly, the pairing frequency did not

vary between these miRNAs (**Main Fig. 1C, ii**), either in the 5p or 3p strand, except for

the last few nucleotides at the 3'end (**Main Fig. 1C, I,** red boxes), which were unpaired

in the mature duplex. When comparing the used vs. the discarded strand (**Main Fig. 1C,**

left panel vs right panel), we detected no significant difference in overall pairing (p = 1.0

in *5p Singletons*, p = 0.12 in *3p Singletons*, per Welch's t-test) for either category of

Singleton.

Extending this analysis to all *5p Singletons*, the 5p guide strand is relatively unpaired,

with an average of 4.42 mismatches across the duplex. Thus, even miRNAs that

consistently select the 5p arm typically retain several internal mismatches that reduce

duplex stability and favor guide loading (**Supplemental Fig. S1B**). However, the 5p

strand discarded in *3p Singletons* was significantly more paired (mean = 3.31

mismatches) than the 5p strand retained in *5p Singletons* (mean = 4.32 mismatches, p

= 0.01 per Welch's t-test), indicating that 5p strands tend to be more stable (i.e., more

paired) when they are not selected (**Supplemental Fig. S1B**). Conversely, the 3p

strand experienced no significant difference (p=0.16) in pairing between *3p*

Singletons (mean = 3.70 mismatches) and *5p Singletons* (mean = 4.32 mismatches),

suggesting its stability is less critical to the strand selection decision. Taken as a whole,

the duplex structures of *5p Singletons* were less paired than those of *3p Singletons*,

which might suggest a divergence in regulation of these two miRNA classes during

biogenesis (**Supplemental Fig. S1B**). When narrowing our analysis to just the seed

region (**Main Fig. 1C**, golden boxes), while there was higher pairing than any other

position, there was no difference in binding frequency between the used and discarded

strands, suggesting that different rules are also in place.

Next, we compared the nucleotide composition of these seed regions, hoping to

identify signatures that could explain the discrepancy in miRNA strand selection in

Singletons. We determined the likelihood of any given nucleotide from positions 1

through 10 in each miRNA strand in the top 20 *Singletons* in each category (5p and 3p)

(**Main Fig. 1C, ii**). We detected a highly conserved uracil nucleotide at position 1, as

previously shown by us and others (2), though this conservation dissolves from

positions 2 through 10 (**Main Fig. 1C, ii**). Importantly, this uracil nucleotide is not

equally conserved between *5p Singletons* and *3p Singletons*. 80% of *3p Singletons* start

with this uracil nucleotide, while less than 40% occur in *5p Singletons*. This differential

presence of this uracil may be caused by a non-stochastic pre-miRNA cleavage by the

C. elegans DCR-1, which may possess a preference to cleave at this nucleotide.

The intestine miRNA population is small but highly interconnected

Many of the guides detected by HiTmiSS were classified as *Switchers* (~46%).

However, some of these may instead represent stage- or tissue-specific *Singletons*,

meaning miRNAs that consistently use one strand within a given context, but switch

strands across different contexts. In other words, these cases would still be considered

Switchers under our definition, but they may appear as *Singletons* when examined in

only one stage or tissue, potentially confounding the analysis.

To address this, we refined our approach by analyzing somatic tissues across

development, focusing on the intestine, which is a large tissue with a well-characterized

transcriptome (41,42).

We coupled our HiTmiSS assay with intestine-specific miRNA immunoprecipitation,

focusing on the embryo vs young adult stages. More specifically, we forced the

expression of the GFP-tagged *C. elegans* Argonaute ortholog *alg-1* in the intestine

tissue using the intestine-specific promoter *ges-1p*, performed an α-GFP

immunoprecipitation, recovered the miRNA strands loaded onto ALG-1, and used them

as templates in our HiTmiSS assay (**Main Fig. 2A**). We found that of the 93 miRNAs

tested, 72 (77%) were present during at least one stage within the intestine tissue (**Main

Fig. 2**). Of these, 66 were *Singletons* (92%), while 6 (8%) were *Switchers*

(**Supplemental Fig. S5A**).

This strongly differed from our whole worm data, suggesting that miRNAs previously

labeled as *Switchers* may truly be *Singletons* on a tissue-by-tissue basis, only

appearing as *Switchers* once all tissues are analyzed together.

Interestingly, the proportion of *5p Singletons* to *3p Singletons* was not the same in

the intestinal tissue as in the mixed tissue (**Supplemental Fig. S5A**). In our initial

analysis, about two-thirds of *Singletons* were *3p Singletons*, yet in the intestine tissue,

3p Singletons were only slightly more abundant than *5p Singletons* (55% vs 45% of

Singletons identified, respectively) (**Supplemental Fig. S5A**).

Another surprise was the presence of intestinal miRNAs with detectable expression

of both 5p and 3p strands (**Main Fig. 2B**). For example, members of the *miR-51* family,

well-characterized regulators of embryonic lethality (36), expressed both strands in the

intestine. Similarly, *miR-44*, *miR-58*, and *miR-71* families showed dual-strand

expression in intestinal samples (**Supplemental Fig. S5B**). These miRNAs appear as

Singleton species in the total-RNA dataset, but both strands are detectable in intestine-

enriched RNA, although one strand is consistently more abundant. These miRNAs

therefore do not undergo strand switching; rather, they maintain dual-strand expression

specifically in the intestine, potentially expanding their available 3' UTR target space in

a tissue-specific manner.

Within this dataset we also identified 6 intestine miRNAs with stage-specific strand

preference; 3 *5p Singletons* (*miR-2*, *miR-87* and *miR-241*) and 3 *3p Singletons* (*miR-36*,

miR-37, and *miR-63*) (**Supplemental Fig. S5A-B**). Importantly, four of these six

Singletons have been independently reported in previous studies (35) (**Supplemental**

Fig. S5B-C), reinforcing their biological relevance.

one life stage. These strains expressed the promoter for a single miRNA fused to GFP,

and live imaging was used to validate the expression of these miRNAs at the

appropriate life stage in the intestine (**Main Fig. 2C**). All four strains displayed the

expected fluorescence.

miRNA target presence does not account for strand usage

Our HiTmiSS array enables precise, strand-resolved quantification of miRNAs *in*

vivo. We considered the possibility that miRNA targets could selectively stabilize one

strand, thereby influencing its relative abundance and potentially confounding

interpretation of the molecular basis of strand selection, rather than merely affecting

HiTmiSS detection. In this model, only target-bound strands would be retained, while

the unused strand would degrade, falsely suggesting strand preference.

To test this, we overexpressed *fox-1* and *hrp-2* 3'UTRs, each containing multiple

binding sites for typically discarded strands (e.g., *miR-60* 5p, *miR-232* 5p, *miR-239b*

5p), in intestinal tissue (**Supplemental Fig. S6**). If target-induced stabilization were

driving strand abundance, we would expect increased detection of these 5p strands.

Although we observed a small increase in the abundance of some typically discarded

strands (e.g. *miR-60-5p* in *fox-1* 3′UTR overexpression and *miR-232-5p* in *hrp-2*

3′UTR overexpression) we saw similar increases in canonically retained strands (e.g.

miR-60-3p in *fox-1* overexpression), which the target stabilization hypothesis cannot

explain (**Supplemental Fig. S6C**). Additionally, none of the detected changes rose

above our analytical threshold of significance (i.e. no 2-fold increase or decrease)

compared to wild-type controls, with 3p strands still predominating in all cases.

Taken together, these results indicate that our qPCR approach is not simply

detecting stabilized miRNA strands due to the presence of targets. Instead, the relative

strand abundance measured by the HiTmiSS assay likely reflects genuine strand

selection, rather than artifacts introduced by target availability.

5p Singletons **and** *3p Singletons* **follow different rules for selecting strand usage**

Next, we examined the structural features associated with *Singletons*, which may

explain the strict and consistent strand usage observed in these miRNAs. When

comparing the used and the discarded strand, the pairing frequency varied significantly

within each *Singleton*, revealing new conserved patterns (**Main Fig. 3**). Specifically, the

strand used in *Singletons* often had an unpaired nucleotide at position 12, while the

discarded strand did not (**Main Fig. 3A**, yellow boxes). This lack of pairing in position 12

in both *5p Singletons* and *3p Singletons* usage is intriguing. In humans, specific

Argonaute proteins perform cleavage of target RNAs between positions 10 and 11,

which corresponds to positions 10 and 11 along the bound miRNA strand by Watson-

Crick pairing (43). *C. elegans* ALG-1 is largely homologous to human Argonaute

proteins, and it may retain this cleavage activity at similar target positions, leading to the

degradation of the unused strand due to its Watson-Crick base pairing at this crucial

position. In addition, we detected differences in the binding frequency of the first-

position nucleotide in both *5p Singletons* and *3p Singletons* (**Main Fig. 3A,** blue boxes).

3p Singletons also exhibited unique pairings at positions 13-19 (**Main Fig. 3**, white box)

with a higher pairing frequency than *5p Singletons* at the same position. These patterns

are most clearly seen in specific *Singleton* miRNAs, both *5p Singletons* such as *miR-52*

and *3p Singletons* such as *miR-75* (**Main Fig. 3B**).

An updated machine learning (ML) model for miRNA strand selection

We wanted to extract patterns and features that make *Singletons* so unique and

predispose them to always use one strand over the other. To achieve this, we utilized

our experimental *Singleton* data as a training set for a machine learning model designed

to predict 5p versus 3p strand usage in miRNA hairpins. We hypothesized that by

training such a model, we could identify the unique intrinsic features that characterize

Singletons.

We opted for a Random Forest classifier, which we trained using our experimentally

validated *Singletons* identified through our HITmiSS assay. This model incorporated 77

features, systematically organized into five functional categories (dinucleotide

composition, first/last nucleotide bias, GC content and entropy, base pairing properties,

and general structural features) that together capture critical aspects of miRNA mature

duplex structure (**Supplemental Fig. S7–S9, Supplemental Tables S5-S6, and**

Materials and Methods). The model demonstrated strong performance in

distinguishing 5p from 3p usage across a set of known miRNAs and other simpler

models (**Supplemental Fig. S8 and Materials and Methods**).

***5p* and *3p Singletons* possess distinct structural features**

Our ML model, built using our *Singleton* data, identified several intrinsic features in

these miRNAs. To generalize this model across all *C. elegans* miRNAs and study how

Switchers can bypass these rules to switch miRNA strand preference, we applied our

ML model and conducted a comparative feature analysis across the entire *C. elegans*

miRNAome (374 miRNAs, miRbase).

We extracted and calculated all 77 features used in our ML model for these miRNAs

(**Supplemental Table S3**), maintaining a stringent definition of *Singletons* to enhance

the signal (confidence score > 80%).

As expected, we identified 54 *5p Singletons*, 52 *3p Singletons,* and 268 *Switchers*

(**Main Figs. 4A-B** and **Supplemental Table S2**). Out of 77 extracted structural and

sequence features, 13 exhibited statistically significant differences between the 5p and

3p strands (p < 0.05, Welch's t-test) (**Main Figs. 4B-E**). The most pronounced

difference was observed in the identity of the first nucleotide of the 3p strand, which in

3p Singletons was consistently a uracil (p = $3.98 \times 10^{-\square}$), followed by an enrichment of

cytosine in the last position (p = $5.69 \times 10^{-\square}$) (**Main Figs. 4C-4E**). Additionally, 3p

strands showed an enrichment for specific motifs, particularly GAG (p = $2.51 \times 10^{-\square}$)

(**Main Fig. 4E**). We also observed compositional differences: 3p strands displayed

increased levels of cytosine at the first position (not shown) and tended to be slightly

shorter than their 5p counterparts (p = 8.90×10^{-3}) (**Main Fig. 4D-E**).

Surprisingly, our ML model did not cluster all the *Switchers* around a confidence score

of ~50%. Instead, we see a multi-modal distribution with several periodical clusters

(**Main Fig. 4A**). This suggests that across this category, structural and thermodynamic

features alone cannot fully explain strand usage, and extrinsic factors could move the

needle between 5p or 3p choice in these miRNAs, perhaps depending on tissue or

developmental context.

We next wanted to study the intrinsic features identified by our ML model. For these

experiments, we used *Singletons* to test if altering these rules could convert them into

Switchers. In *let-7*, modifying nucleotide pairing at positions 11–12 (UP11-12) or

changing the first base (5'U→C) reduced its predicted *5p Singleton* confidence by up to

17%, consistent with our model (**Main Figs. 3C-D**). Similar results were observed in *lin-*

4, miR-52, *miR-75*, and *miR-35*: introducing bulges near the center was predicted to

disrupt strand preference, while removing them was predicted to reinforce it

(**Supplemental Fig. S9**), highlighting the central region's role in strand selection.

The first nucleotide's identity and pairing also influenced strand prediction, though with

variable impact across miRNAs (**Supplemental Fig. S9**). Mutating this position was

predicted to alter strand usage by less than 10% in *5p Single*tons but up to 30% in *3p*

Singletons, mirroring HiTmiSS results.

Strand usage bias differs by species

Our ML model identified several features that may distinguish *Singletons* from

Switchers in *C. elegans*. Since many miRNA sequences are conserved across species,

we hypothesized that the features identified by our ML model were selectively

maintained throughout evolutionary time. To do that, we applied the same model to the

complete human miRNAome (**Main Fig. 5**) (**Supplemental Table S6**), and to other

species throughout the tree of animal life (*Drosophila melanogaster*; fly, *Danio rerio*;

zebrafish, *Xenopus laevis*; frog, *Rattus norvegicus*; rat, *Mus musculus*; mouse) (**Main**

Fig. 5, **Supplemental Fig. S10** and **Supplemental Table S6**). While miRNA strand

usage in worm, fly, zebrafish, and frog was predicted to remain relatively balanced

between the 5p and 3p arms, mammalian species were predicted to exhibit a

pronounced preference for the 5p strand **(Supplemental Fig. S10A)**. In *C. elegans*,

only 51.3% of miRNAs were predicted to use the 5p strand, and in *Drosophila*, 55.6%

were 5p. In contrast, this proportion rose sharply to 78.2% in mice and 75.9% in

humans, indicating a shift toward a fixed strand preference (**Main Fig. 5A** and

Supplemental Fig. S10A).

Across species, we also observed an apparent decrease in pre-miRNA length of the

used strands, especially in *Switchers* and *3p Singletons* (**Supplemental Fig. S10B**).

This suggests that in vertebrates, shorter pre-miRNAs can adopt fewer folding patterns,

potentially enhancing the use of specific strands.

Next, we sought to identify structural features that could explain the sharp increase

in predicted 5p strand usage observed in mammals. We identified 13 significantly

different features between worms and humans ($p < 0.05$, Welch's t-test), with striking

enrichment of guanine content ($p = 5.62 \times 10^{-2}$□) and GG dinucleotides ($p = 1.94 \times$

10^{-2}□) on human 5p strands (**Main Fig. 5B**). Invertebrates exhibited a subtle

enrichment in AU-rich motifs (UU: +2.46% higher in invertebrates; AU: +2.39% UA:

+1.70%; AA: +2.11%), consistent with lower duplex stability and more flexible strand

usage (**Supplemental Fig. S10C-D**). In contrast, mammals showed increased

frequencies of GC-rich dinucleotides such as GG and GC, especially within the

disfavored strand (di_GG: 3.6% higher in mammals than in invertebrates; di_GC: 0.9%

higher in mammals) (**Main Fig. 5B**, and **Supplemental Fig. S10C-D**). The overall GC

content in worm*,* fly, and frog miRNAs was 44-45%,

whereas mouse and human miRNAs exhibited significantly higher GC content (~52–

53%). This overall increase in GC content, which translates to an increase in duplex

stability, could be one of the most important features driving 5p strand usage in

vertebrates.

Finally, analysis of 5' nucleotide identity showed that all species retained a

preference for uracil at the first nucleotide of the loaded strand (2). However,

mammalian species exhibited an increase in guanine and cytosine at the first position of

the 3p strand, as well as higher CCU content and diCC motifs ($p < 5 \times 10^{-2}$). These

trends are evident in conserved miRNAs, such as *let-7* (**Main Fig. 5C**), which potentially

serve as a negative selection marker to suppress their incorporation into RISC.

Taken together, these features may provide structural hints as to how vertebrates

adopted a fixed strand usage. Shorter and more paired duplexes may enhance the

selection of *Singletons* in vertebrates, while longer and less structured sequences may

allow invertebrates to increase usage of *Switchers*.

Our ML model can accurately predict human miRNA strand selection

Our *C. elegans* ML model was also capable of identifying sequence elements and

structural features in human miRNAs. To further analyze these results, we compared its

predictions to expression data from the Human miRNA Tissue Atlas across 65 healthy

tissues and cell lines (34) **(Main Fig. 6A-B)**. Overall, the model reached a median of

68% accuracy throughout all datasets, with peaks of 81% in cell lines and 78% in

organs of origin **(Main Fig. 6AC)**. Most errors involved 3p-dominant miRNAs

misclassified as 5p, reflecting a model bias toward 5p predictions. These

misclassifications often occurred when 5p and 3p strands had similar expression levels,

suggesting either model uncertainty or that these miRNAs may function as *Switchers* in

humans **(Supplemental Table S6).**

Notably, two misclassified miRNAs, *miR-182* and *miR-378a*, are known *Switchers in*

vivo (44,45). In other cases, mispredictions occurred when the model's confidence

scores for 5p and 3p usage were nearly equal (±10%). For instance, *miR-199a* and

miR-199b, involved in hypoxia and cardiac regulation (46), were predicted as 5p

dominant despite showing 3p expression and a narrow confidence margin (~7%).

Similarly, *miR-124*, critical for neurogenesis, was misclassified with a 6% confidence

gap, despite clear 3p dominance (47).

Still, many ML model predicted miRNA strands were accurately detected across

diverse human tissues, including key regulators with strong strand bias, indicating that

our *C. elegans*-trained model generalizes well to vertebrates (**Main Fig. 6C and**

Supplemental Table S6). For example, *miR-21*, a highly expressed oncomiR, was

correctly predicted as 5p-dominant, with over 50-fold higher 5p expression than 3p (48).

Similarly, *let-7a* showed a 248-fold preference for 5p and was correctly classified (49).

Other highly expressed miRNAs, such as *miR-1* and miR-*141,* were also correctly

predicted as 3p dominant.

Overall, these findings demonstrate that the model reliably predicts strand usage for

miRNAs in healthy human tissues, particularly when strand bias is strong. Errors were

enriched among duplexes with more balanced expression levels or context-dependent

strand usage, highlighting the complexity of predicting which strand will be used in

cases of weak or dynamic regulation of strand selection.

Evolution drives strand-selection shifts

Because our ML model predictions in *C. elegans* mirror miRNA strand usage

patterns observed in humans, we next asked whether the sequence and structural

"feature rules" learned in *C. elegans* were conserved or repurposed across metazoans.

Very few miRNAs are deeply conserved between nematodes and humans, and those

that are (e.g., *let-7*) are uniformly *5p Singletons* in both species. A notable exception is

miR-34 (**Main Fig. 7**). In *C. elegans, miR-34* contributes to stress resistance, aging, and

DNA-damage responses (50,51), while in humans it functions downstream of p53 and

acts as a tumor suppressor that inhibits proliferation, prevents metastasis, and

promotes apoptosis (52-54). Strikingly, *miR-34* is a *3p Singleton* in *C. elegans* across all

developmental stages, yet it becomes a *5p Singleton* in mammals, matching both our

ML model's predictions and the miRNA atlas data. To understand this inversion, we

traced sequence and structural changes in *miR-34* precursors across fly, mouse, and

human (**Main Fig. 7**). Across metazoans, *miR-34* shows a clear evolutionary flip from

3p maturation in basal organisms to 5p dominance in mammals. In *C. elegans, miR-34*

is predominantly processed from the 3p arm (P(3p)=0.76), driven by a strong

thermodynamic bias. The 3p arm contains 63.6% GC versus 54.5% on the 5p arm,

yielding a more stable 3′ duplex end and favoring Argonaute loading (**Main Fig. 7**).

Although the 5p product begins with a favorable 5′-A, duplex asymmetry still enforces

3p selection. A decisive shift appears in insects: *Drosophila miR-34-5p* acquires a 5′-U,

which is the canonical AGO-loading nucleotide, coincident with a switch to 5p

dominance (P(5p)=0.68). This A-to-U substitution at position 1 is retained in vertebrates,

stabilizing 5p usage (mouse P(5p)=0.70; human miR-34a frequently ≥0.7) (**Main Fig. 7**).

In addition, human precursors display shorter terminal loops (~6–8 nt versus ~12–15 nt

in worm), promoting precise Drosha/Dicer cleavage and a perfectly conserved 5′ seed

start. Dinucleotide composition also evolves in a direction that weakens 5p end-pairing:

di_GU progressively increases, introducing GU wobble near the guide and lowering

local duplex stability, which favors 5p loading. Conversely, di_CG and di_GA decline to

zero in mammals (di_GA is already lost in fly), eliminating stabilizing stacking

interactions still present in worms. Collectively, the shift from 3p to 5p usage in *miR-34*

occurred through (i) loss of 3p GC enrichment, (ii) gain of AU-rich 5′ nucleotides that

enhance AGO recognition, (iii) terminal loop shortening that improves processing

precision, and (iv) dinucleotide substitutions that reduce 5p duplex stability relative to

3p. Thus, fly represents an intermediate state, while mouse and human complete the

evolutionary inversion, establishing *miR-34a-5p* as the conserved, functionally dominant

product in mammals (**Main Fig. 7**).

A community resource to study miRNA strand selection

To support the broader research community, we have made all components of our

strand selection framework openly available via GitHub

(https://github.com/MangoneLab/Caligula_app). This resource includes our trained ML

model, as well as Python scripts for both local, cumulative, and browser-based use. The

platform is designed for flexibility and ease of use, allowing users to input a single

miRNA or a batch of sequences in FASTA format to obtain strand selection predictions.

We also provide pre-processed strand selection predictions, including all 77 features

used by our ML model, for every miRNA annotated in miRBase across seven species:

worm, fly, zebrafish, frog, mouse, rat, and human (**Supplemental Tables S3 and S6**).

By making these tools and datasets accessible, we aim to provide a practical,

scalable, and user-friendly solution for investigating miRNA strand usage across

systems and species. This represents the first publicly available platform for high-

throughput, strand-specific miRNA analysis, enabling researchers to integrate strand

selection into both basic and translational studies.

Discussion

This work establishes a unified, experimentally validated model of miRNA strand

selection that links intrinsic duplex structure to context-dependent strand usage across

development and tissues. By combining ~8,000 qPCR measurements with feature-

based machine learning, we show that strand choice is highly regulated and not

stochastic, and that conserved sequence signatures help enforce *Singleton* behavior

while *Switchers* remain responsive to developmental and tissue cues.

Compared to RNA-seq, HiTmiSS offers distinct advantages for miRNA analysis

(**Supplemental Figs. S4-S6**). The primary differences between these methods lie in the

output produced. In RNA-seq, the output consists of sequences that must be mapped

onto a reference genome to infer strand usage and abundance, whereas in HiTmiSS,

the output provides direct quantification of each strand. Because strand-specific primers

directly amplify each mature duplex arm, HiTmiSS avoids inference-based strand

assignment and ligation bias, which can obscure low-abundance species or inflate the

apparent abundance of highly expressed miRNA families. We find that HiTmiSS is

highly sensitive and reproducible, detecting transcripts that are often weakly captured

by RNA-seq while still recapitulating developmental trends observed in sequencing

datasets. Thus, RNA-seq provides breadth and discovery power, whereas HiTmiSS

delivers quantitative precision and clearer 5p/3p discrimination, especially in low-

abundance or tissue-restricted contexts.

Recent approaches, such as AQ-seq, have improved small RNA-seq protocols by

reducing ligation bias and enabling isomiR detection (55). However, because AQ-seq

still relies on sequencing-based inference for strand assignment, it may be affected by

residual mapping ambiguities. In contrast, HiTmiSS directly quantifies 5p and 3p strands

using strand-specific primers, offering a more targeted approach to strand-resolved

analysis. Its combination of sensitivity, quantitative precision, and reproducibility makes

it well-suited for studying miRNA strand selection across tissues and developmental

stages.

Our study enabled the classification of miRNA into *5p Singletons*, *3p Singletons*, or

Switchers, informing a predictive model validated against human miRNA expression

data.

We found that in *C. elegans,* half of miRNAs (46%) behave as *Switchers*, altering

strand usage across development, indicating that strand selection is context-dependent

rather than fixed (**Main Fig. 1)**. In contrast, key developmental regulators such as *let-7*,

lin-4, and *miR-35* are *Singletons*, likely reflecting a need for consistent strand usage to

ensure stable gene expression. Notably, *Singletons* tend to be more highly expressed

(**Main Fig. 1B**), perhaps due to stronger structural or thermodynamic constraints. In the

intestine, tissue-specific ALG-1 immunoprecipitation revealed a strong preference for

Singletons (**Main Fig. 2**). This suggests that strand selection is also shaped by tissue-

specific factors, such as the cellular environment or cofactors. These factors would have

substantial consequences for miRNA regulatory output, as strand choice reshapes the

accessible set of 3'UTR targets. While our approach captures intrinsic sequence

determinants, with the caveat that only one tissue (intestine) and two developmental

stages (embryo and young adult) were profiled, the tissue-specific strand usage we

observe points to an additional, regulated layer of post-transcriptional control. Identifying

the proteins and regulatory pathways that modulate strand choice is an important future

direction, both in *C. elegans* and likely in other metazoans. Importantly, since we

examined only one somatic tissue, some miRNAs classified as *Switchers* in our

experiments may be tissue-specific *Singletons*.

A concern was that target mRNAs might stabilize one strand, creating an artificial

strand preference in our HiTmiSS assay. To test this, we overexpressed 3'UTRs with

binding sites for typically low-abundance strands in the intestine. Although we cannot

rule out other downstream processes affecting miRNA stability, such as decay rates or

target-mediated miRNA degradation (56,57), strand levels remained unchanged,

suggesting that HiTmiSS reflects strand usage, not target-driven stabilization

(**Supplemental Fig. S6**).

When comparing the structural composition of each mature duplex, a series of

patterns emerged. Regarding the strand pairing, there is no universal trend showing that

5p strands were always more unpaired than 3p strands, regardless of selection.

Instead, selected strands (whether 5p or 3p) tended to tolerate moderate unpairing, and

unselected 5p strands (i.e., in *3p Singletons*) were more paired, suggesting that greater

structural stability may discourage selection, supporting a role for pairing in strand

discrimination (**Main Fig. 1C**).

5p Singletons tended to have the first nucleotide unpaired to its corresponding

strand in the duplex (**Main Fig. 3A**). This difference in pairing repeats in the middle of

the duplex and lowers toward the end (**Main Fig. 3A**). Contrarily, *3p Singletons* present

a near-opposite trend, with the first nucleotide more often paired and the end portion of

the sequence also paired, though the center of the sequence shows similar unpairing to

5p Singletons.

The effects of the 5' nucleotide identity and central region pairing on strand selection

and Argonaute loading have been well known for a long time, with foundational work

highlighting the importance of 5' nucleotide identity and central region pairing for

Argonaute loading (14,58,59). These studies established general rules governing strand

preference, such as the thermodynamic asymmetry model, but were derived under

controlled conditions, often focusing on a limited number of miRNAs in model systems.

While they established the principles of strand selection, they did not provide a scalable

means to measure how these rules were implemented, or overridden, in living

organisms across developmental stages, tissues, or physiological contexts. HiTmiSS

advances the field by shifting the focus from mechanism to measurement, offering a

practical, high-throughput platform for directly quantifying strand usage across biological

conditions, developmental stages, and tissues. In doing so, HiTmiSS transforms strand

selection from a static rule set into a measurable, context-dependent regulatory layer.

In addition, our results added positional detail to these foundational studies by

showing that asymmetry varies along the duplex. Both *5p* and *3p Singletons* show

central unpairing, likely reflecting a conserved feature for Dicer or Argonaute

engagement but not strand bias. Notably, *5p Singletons* show decreasing pairing from

the seed to the 3' end, while *3p Singletons* display the opposite trend. These opposing

patterns suggest that specific structural cues along the duplex, not just global stability,

guide strand selection, pointing to evolved, position-dependent regulatory signals within

miRNA architecture.

To expand our analysis, we extracted 77 structural and compositional features from

our validated *Singletons* and trained a Random Forest classifier (ML model) to predict

dominant strand usage. When comparing these features (**Main Fig. 4**), we found

several strand-specific differences, including a significantly higher occurrence of uracil

at the first nucleotide in 3p strands and enrichment for GAG motifs.

We tested our predictions using *in silico* site-directed structural mutations in

canonical miRNAs such as *let-7*, *lin-4*, and *miR-52*, *miR-75* (**Main Fig. 3C**,

Supplemental Fig. S9). Disrupting features such as central bulges (**Main Fig. 3C,**

"UP11-12") or first nucleotide pairing (**Main Fig. 3C, "1U->C"**), altered predicted strand

preference to some degree, further confirming their role in the selection process.

Surprisingly, the degree to which they impact miRNA strand preference was

approximately equal, but when disrupted in tandem, they led to an even greater loss of

predicted strand preference (**Main Fig. 3C, "Combined"**, **Supplemental Fig. S9**). In

addition, these findings phenocopy the trends observed in our experimentally validated

Singletons (**Main Fig. 3A**). Taken together, these results suggest that the structural and

thermodynamic features identified in our wet bench-validated *5p* and *3p Singletons* are

indeed critical regulators of strand selection in nematodes.

It is important to note that these analyses do not constitute experimental validation of

the model. Rather, they provide an additional test of the model by probing how specific

sequence and structural features influence its predictions. We modified known

determinants of strand usage, such as 5′ nucleotide identity, central mismatches, and

local thermodynamic asymmetry, within biologically plausible ranges and consistent with

feature distributions observed in the training data. We focused not on single predictions,

but on systematic and reproducible shifts across multiple miRNAs and perturbation

types. In several cases, including *miR-75*, the modified sequences not only reduced

confidence but also flipped predicted strand preference, an outcome consistent with

switcher-like behavior reported in the literature. However, we emphasize that these

findings are predictive and hypothesis-generating, and experimental testing will be

required to determine whether these perturbations alter strand usage *in vivo*.

Expanding this framework, we then applied our ML model to predict strand usage

across the human miRNA dataset (**Main Fig. 5A**). We were particularly curious to test if

the structural and thermodynamic features and motifs identified in *C. elegans* were

conserved in vertebrates.

Compared to the nearly equal distribution of 5p and 3p usage in *C. elegans*, using

the same cutoff of 80%, the vast majority of human miRNAs are identified as *Switchers*

(831/854 predicted *Switchers*, 97.3%). However, within this class, there is a clear bias

toward 5p strands, with over 75% of predictions favoring 5p strand usage (**Main Fig.**

5A). This divergence likely reflects increased specialization in miRNA maturation and

Argonaute loading in vertebrates, potentially driven by evolutionary selection for strand-

specific targeting or stability.

Despite the lower predictive accuracy of our ML model when applied to *H.*

sapiens miRNAs (69%) compared to its performance on the *C. elegans* training set

(88%) (**Main Fig. 6C**, **Supplemental Fig. S2B**), we do not interpret this as evidence of

overfitting. Rather, the ability of the *C. elegans*-trained model to achieve performance

substantially above random on a distantly related species suggests that key aspects of

miRNA strand selection are indeed governed by conserved predictive features. While

quantitative differences in model accuracy likely reflect evolutionary divergence and the

influence of lineage-specific modifiers, the persistence of these core features supports

the idea that a foundational set of strand selection rules is conserved. Thus, our model

reveals not only species-specific differences but also a shared regulatory logic

underlying miRNA strand selection that has been retained across metazoan evolution

(**Supplemental Fig. S10**). Invertebrate systems appear to tolerate greater flexibility,

allowing for more balanced usage of the duplex arms. In contrast, vertebrates,

particularly mammals, exhibit strong sequence and structural biases that favor the

selection of the 5p strand (**Supplemental Fig. S10**). This transition likely reflects the

growing complexity of regulatory networks and the need for precise post-transcriptional

gene control in higher organisms. As miRNA families expanded during vertebrate,

particularly mammalian, evolution, selective forces likely favored stable strand usage to

ensure consistent and non-redundant regulation (2). This trend would promote

Singletons over *Switchers*, reducing regulatory variability. The fixation of 5p strand

usage may thus represent stabilizing selection for predictable miRNA function, with

structural features reinforcing 5p bias evolving alongside increasing organismal

complexity.

In addition to this imbalance in *Singletons* vs *Switchers*, the nucleotide composition

of their 5p and 3p strands is also different, with a significant enrichment of guanine

content in human 5p strands and cytosine-rich motifs in 3p strands (**Main Fig. 5B**).

These compositional biases are consistent with known differences in Argonaute protein

specificity and may reflect lineage-specific constraints on strand functionality (60).

Importantly, these cross-species trends persisted in known miRNAs such as *let-7*,

which was consistently predicted and validated as a *5p Singleton* both in *C. elegans*

and in humans, although at different confidence levels. This supports the robustness of

our predictive ML model and highlights conserved features of miRNA processing, even

in the presence of divergent strand preferences.

A surprising trend emerges when we compare our ML model prediction in human

miRNAs to the human miRNA Tissue Atlas. Our ML model developed in *C. elegans* is

capable of accurately (~70%) predicting miRNA strand usage in humans (**Main Fig. 6**).

This is unexpected, considering the vast evolutionary gap (~600 million years) between

these two systems, which share less than 35% of nucleotide conservation in genes (61).

Despite this, the features we have identified seem to be evolutionarily conserved.

This suggests that miRNA strand selection is, at its core, a fundamental cellular

process that has evolved very little over time, even as miRNA regulatory networks

expanded in complexity with the increased cellular specialization. This becomes more

evident when we overlay miRNA strand usage in human tissues of different origins.

Here, the same miRNA is always predicted to use the same strand in other somatic

tissues (**Main Fig. 6A**, **Supplemental Table S6**). This invariance suggests that strand

selection is most likely encoded by intrinsic properties of the hairpin itself, rather than by

external, context-dependent cues. Such robustness implies that strand choice is not

merely a byproduct of downstream regulatory needs, but a core mechanistic feature

established early in miRNA biogenesis. We propose that during vertebrate evolution,

particularly with the rise of tissue specialization, miRNA strand usage remained

anchored to conserved structural determinants even as the broader miRNA regulatory

network expanded and diversified. This mechanistic rigidity may stabilize gene

regulatory outputs in the face of cellular complexity.

Tracing the origins and evolution of miRNA families remains challenging. Although

miRNAs emerged early in metazoan history, multiple bursts of rapid expansion occurred

during bilaterian diversification (~520–500 Mya) and again during the vertebrate whole-

genome duplications. These events generated large and highly specialized miRNA

families that later diverged in sequence, genomic context, and copy number, often

obscuring clear orthologous relationships between vertebrate and invertebrate miRNAs

(2). The *C. elegans miR-34* family is one of the few deeply conserved examples in

which such evolutionary comparisons can still be performed. Our analysis of *miR-34*

(**Main Fig. 7**) shows that miRNA strand selection is not fixed but can be evolutionarily

rewired through incremental sequence and structural changes to the hairpin precursor.

Although the rules learned in *C. elegans* can predict strand usage in humans, the

evolutionary trajectory between these species indicates that these rules are repurposed

rather than strictly conserved. The transition of *miR-34* from 3p maturation in basal

metazoans to 5p dominance in mammals demonstrates that relatively subtle features,

including 5′ nucleotide identity, GC asymmetry, terminal loop length, and dinucleotide

composition, are sufficient to invert guide-strand selection without disrupting hairpin

recognition or overall function. These findings suggest that strand choice itself may be

an evolvable regulatory parameter. Altering which arm generates the dominant guide

effectively redefines the seed sequence and rewires target networks without deleting or

duplicating the miRNA gene. Evolution can therefore modify miRNA function through

"strand-switching," expanding regulatory potential while preserving genomic economy.

The observation that intermediate species exhibit partial transitions further supports

a model in which strand usage diverges gradually, as selective pressures favor a new

dominant arm with distinct targets.

Importantly, the same features that govern strand selection in worms (5′-nucleotide

identity, duplex asymmetry, wobble positioning, and cleavage precision) also predict

usage in insects, mammals, and humans. In this view, evolution does not invent new

rules; instead, it repurposes existing thermodynamic and structural logic to shift guide

preference. Together, these data indicate that strand selection is a flexible,

evolutionarily tunable mechanism for reshaping miRNA regulatory networks. The

conservation of feature-level logic across metazoans, combined with species-specific

rewiring of dominant strands, suggests that strand switching may represent a

widespread but underappreciated layer of post-transcriptional innovation.

There are limitations to the present work. We profiled only one somatic tissue (the

intestine) and two developmental stages, and it is likely that additional *Singletons*

remain hidden within other tissues or stress environments. The ML model currently

captures intrinsic features but not Ago preferences, cofactors, RNA modifications, or

degradation pathways that may bias strand retention. Identifying these regulators, and

defining how they interface with duplex structure, will be an important future direction.

Finally, we have developed a user-friendly resource on GitHub

(https://github.com/MangoneLab/Caligula_app) that allows researchers to predict

miRNA strand selection using our ML model. To support broad application, we also

provide pre-processed strand selection predictions with all 77 input features for all

miRNAs in miRBase across seven species, including worm, fly, zebrafish, frog, mouse,

rat, and human (**Supplemental Tables S3 and S6**).

In summary, this study establishes a predictive and experimentally validated

framework for miRNA strand selection, combining high-throughput qPCR (HiTmiSS)

with an ML model to uncover both conserved and context-dependent rules governing

strand usage. Unlike previous studies that inferred strand selection indirectly or focused

on a limited set of miRNAs, our study provided strand-specific quantification across

developmental stages and tissues in *C. elegans*, revealing dynamic and non-stochastic

strand usage. We also produced a biologically grounded and publicly available ML

model trained on curated *in vivo* strand data that accurately predicts strand selection

across multiple species, including vertebrates. We also prepared the first species-wide

probability atlas of 5p vs. 3p usage across several metazoan miRNAs, including

humans, enabling genome-scale interpretation of strand bias. In addition, we highlighted

a detailed dissection of 77 intrinsic features that together define a conserved,

programmable regulatory logic for strand selection. This integrative approach extends

beyond descriptive rules, providing a quantitative and generalizable model of strand

selection applicable across metazoan evolution.

In conclusion, our findings illustrate the complexity of miRNA strand selection and

reveal that this process is not governed solely by thermodynamic asymmetry. Instead, a

combination of structural motifs, sequence biases, and cellular context dictates strand

usage. Our work establishes a predictive framework for understanding strand selection

but also opens new avenues for studying the evolution of miRNA regulatory logic across

metazoans.

1189

1190 Data Availability

1191 The raw data produced by the HiTmiSS assay are included in the **Supplemental Table**

1192 **S2**. The ML model is included as Supplemental Data and is also available on GitHub

1193 (https://github.com/MangoneLab/Caligula_app).

1194

1195 Funding

1196 This work was funded by the National Institutes of Health 5R01GM118796 awarded to

1197 M.M.

1198

1199 Acknowledgments

1200 We thank Dr. Murugan for assistance in HiTmiSS assay optimization.

1201

1202 Author Contributions

D.M. and M.M. designed the experiments. D.M. performed the experiments. D.M. and

M.M. prepared the ML model and performed the bioinformatics analysis. H.L.

supervised the development of the ML model. D.M., A.E., and H.H. performed the

HiTmiSS assay experiments. D.M. and M.M. prepared the main and supplemental

figures and supplemental tables. D.M. and M.M. wrote the manuscript. All authors have

read and approved the manuscript.

1209

1210   **Declaration of Interests**

1211   The authors have no conflicts of interest.

1212

Figure titles and legends

Main Figure 1: HiTmiSS Assay and Analysis on Mixed Tissue. **(A)** HiTmiSS protocol

for RNA extraction, extension, reverse transcription, and quantification in high

throughput (i). Steps include (1) *C. elegans* growth and homogenization, (2) RNA

extraction and isolation, (3) RNA polyuridinylation, extension, and reverse transcription

(details shown in ii; orange: miRNAs, green: polyU, purple: SL-polyA reverse universal

primer), (4) addition of 5p or 3p miRNA-strand-specific forward primers, (5) loading into

384-well plates, and (6) quantification. **(B)** Heatmap of miRNA expression (n=3) in

mixed tissue clustered by quantity, strand use, and developmental stage. Stages

include embryo (E), larval 1 through 4 (L1-L4), and young adult (YA). The heatmap

legend represents the $\log_{10}$ of expression values for each miRNA strand, normalized by

the spike-in GFP control, at each stage, ranging from -4 (deep purple) to 4 (bright

yellow). Black color indicates instances of data that fall below the strict expression

threshold (one standard deviation above the mean of all expression values at that

stage). **(C)** Heatmap of pairing frequency by position for the used (left) and discarded

(right) strand of the 20 most highly quantified *5p* and *3p Singletons* in our mixed tissue

dataset. (i) blue: paired; red: unpaired). (ii) Average pairing frequency, and logo plot of

nucleotide frequency at each position displayed below (gold box marks the seed

regions).

1232

Main Figure 2: Analysis of Intestine Tissue-Specific miRNAs. **(A)** Schematics of the

intestine-specific GFP-tagged ALG-1 immunoprecipitation. **(B)** Heatmap of miRNA

expression in intestine tissue clustered by strand and developmental stage. The

heatmap legend represents $\log_{10}$ of expression values for each miRNA strand

normalized by spike-in GFP control at each stage from -4 (deep purple) to 4 (bright

yellow). Black color indicates instances of data that fall below the strict expression

threshold (mean of all expression values at that stage). E: embryo; YA: young adults.

(C) Representative brightfield (left) and GFP (right) images of selected *C. elegans*

strains expressing GFP under the control of promoters for miRNA known to be highly

expressed in the intestine, either during the embryo stage (top two) or young adult stage

(bottom two). Arrows indicate GFP expression in the intestinal tissues of both embryo

and young adult stages.

1245

Main Figure 3: Structural and Thermodynamic Determinants of Strand Selection.

(A) Pairing frequency by position of the used (green) and discarded (red) strand for the

most highly quantified *5p Singletons* and *3p Singletons* in mixed tissue (n=20). Positions

at which pairing frequency most differs are boxed, including the first nucleotide (light

blue), the center of the duplex (gold), and the end of the strand (white). **(B)**

Representative pri-miRNA hairpins for a *5p Singleton* (*miR-52*) and a *3p Singleton*

(*miR-75*) demonstrating the occurrence of trends in pairing as described above. Mature

duplex regions are highlighted in pink. **(C)** Changes in *Singleton* predictive confidence

for the miRNA *let-7* as a result of *in silico* site-directed mutations. RNAfold secondary

structures of pri-miRNA with mature 5p strand (blue) and mature 3p strand (red) from

left to right: wildtype (WT), unpaired nucleotides at positions 11 and 12 via GG→UU

substitution (UP11-12), first nucleotide U→C substitution (1U → C), combination of both

prior mutations (Combined). The table summarizes the predicted strand preference by

our ML model for each mutation. **(D)** Bar chart displaying probability as described

above.

1261

Main Figure 4: Analysis of Mature miRNA Duplex Features in *C. elegans*. (A)

Frequency distribution of *5p Singleton* predicted confidence by our ML model across all

miRNAs in *C. elegans*. *3p Singletons* (yellow) are defined as miRNAs falling below a

predicted 5p confidence of 20%; *Switchers* are defined as miRNAs with a predicted 5p

confidence between 20% and 80%; *5p Singletons* (red) are defined as miRNAs with a

predicted 5p confidence of 80% or greater. **(B)** Predicted categorization of all *C.*

elegans miRNAs by our ML model as per the above threshold. **(C)** Predicted

categorization of *C. elegans* miRNAs for which uracil is the first nucleotide of the 3p

strand. **(D)** Length difference (5p strand - 3p strand) for each category of *C. elegans*

miRNAs. *5p Singletons* display significantly ($p<0.05$) longer 5p strands than 3p strands

when compared either to *3p Singletons* or to *Switchers*. **(E)** Heatmap of 32 features

including GC content, miRNA length, hydrogen bonding, nucleotide pairing, strand

stability, and MFOLD-derived parameters enrichment in *5p* and *3p Singletons*. Heatmap

values represent the mean values of each feature across the miRNA categories. For

binary features (i.e., 5p_has_AAU, which is either 0 or 1), the number represents the

proportion (e.g., 0.3 = 30% of miRNAs in that category have an AAU motif in the 5p

strand). For continuous features (i.e., gc_content, pairing_frac_5p), the number is the

average value across all miRNAs in that category. For nucleotide composition features

(i.e., 5p_A, 3p_U), the number reflects the fraction of that nucleotide within the strand,

features clustered by average linkage.

1282

Main Figure 5: Comparative Analysis of Mature miRNA Duplex Features between

***C. elegans* and *H. sapiens*. (A)** Frequency distribution of predicted *5p Singleton*

confidence across all miRNAs in *H. sapiens*. *3p Singletons* (yellow) are defined as

miRNAs falling below a predicted 5p confidence of 20% or below; *Switchers* are defined

as miRNAs with a predicted 5p confidence between 20% and 80%; *5p Singletons* (red)

are defined as miRNAs with a predicted 5p confidence of 80% or greater. **(B)** The four

most significant ($p<0.05$) structural differences between *H. sapiens* (left) and *C. elegans*

(right) mature miRNA duplexes. From top left to bottom right: mean occurrence of

guanine in the 5p strand (5p_G), mean occurrence of guanine-guanine dinucleotide in

either mature strand (di_GG), mean occurrence of cytosine-cytosine dinucleotide in

either mature strand (di_CC), binary classifier for presence of cytosine-cytosine-uracil

trinucleotide in the 3p strand (3p_has_CCU). **(C)** Alignment of *C. elegans* (*cel*) *let-7* and

five *H. sapiens* (*hsa*) *let-7* homologs shows differences in CCU occurrence within the 3p

strand (red boxes). **(D)** Heatmap of differences in mean value for each of the 77 model

features between each category of miRNA in *C. elegans* (c) and *H. sapiens* (h). For

binary features (i.e., 3p_has_CCU, which is either 0 or 1), the number represents the

proportion (e.g., 0.3 = 30% of miRNAs in that category have a CCU motif in the 3p

strand). For continuous features (i.e., gc_content, au_content), the number is the

average value across all miRNAs in that category. For nucleotide composition features

(i.e., 5p_A, 5p_U), the number reflects the fraction of that nucleotide within the strand.

Features clustered by average linkage.

1304

Main Figure 6: Model Accuracy for the Human miRNA Tissue Atlas Dataset. **(A)**

Model accuracy expressed as a percentage of correct HiTmiSS predictions for of *H.*

sapiens tissue and organ miRNAs from the miRNA Tissue Atlas (410 tissues/organs).

(B) Model accuracy expressed as a percentage of correct HiTmiSS predictions for of *H.*

sapiens cell lines miRNAs from the miRNA Tissue Atlas (264 cell lines) **(C)** Cumulative

Model Accuracy by expression Rank for the top 10, 50 and all miRNAs in the miRNA

Tissue Atlas.

1312

Main Figure 7: Evolutionary analysis of *miR-34* strand usage in worms, flies,

mice, and humans. Across metazoans, *miR-34* undergoes a striking evolutionary

inversion from *3p Singleton* in *C. elegans* to *5p Singleton* in insects and mammals**.**

(Top) The seed region of *miR-34* is highlighted in gray. In worms, stronger 3p GC

content (63.6% vs. 54.5% on 5p) enforces 3p loading despite a 5′-A on the 5p guide. In

insects, *miR-34-5p* acquires a 5′-U (yellow), coincident with 5p dominance (P(5p)=0.68),

and this feature is conserved in vertebrates (mouse P(5p)=0.70; human ≥0.7. (Bottom

Left). Model-derived GC asymmetry rises gradually across species, from *C. elegans* to

fly, mouse, and human, and this trend is mirrored by a progressive increase in 5p

pairing (bottom right).

1323

Supplemental Information titles and legends

Supplemental Table S1: **Primers used in this study.**

Supplemental Table S2: **HiTmiSS assay results.**

Supplemental Table S3: **Features used in our ML model and their weights.**

Supplemental Table S4: **Comparison of ML model with miRNA Tissue Atlas.**

Supplemental Table S5: **ML prediction in *C. elegans*.**

Supplemental Table S6: **ML prediction in other species.**

1331

References

1.   Bartel, D.P. (2018) Metazoan MicroRNAs. *Cell*, **173**, 20-51.

2.   Wolter, J.M., Le, H.H., Linse, A., Godlove, V.A., Nguyen, T.D., Kotagama, K., Lynch,

A., Rawls, A. and Mangone, M. (2017) Evolutionary patterns of metazoan microRNAs

reveal targeting principles in the let-7 and miR-10 families. *Genome Res*, **27**, 53-63.

3.   Lee, Y., Ahn, C., Han, J., Choi, H., Kim, J., Yim, J., Lee, J., Provost, P., Radmark, O.,

Kim, S. *et al.* (2003) The nuclear RNase III Drosha initiates microRNA processing.

Nature, **425**, 415-419.

4.   Han, J., Lee, Y., Yeom, K.H., Kim, Y.K., Jin, H. and Kim, V.N. (2004) The Drosha-

DGCR8 complex in primary microRNA processing. *Genes Dev*, **18**, 3016-3027.

5.   Gregory, R.I., Yan, K.P., Amuthan, G., Chendrimada, T., Doratotaj, B., Cooch, N. and

Shiekhattar, R. (2004) The Microprocessor complex mediates the genesis of microRNAs.

Nature, **432**, 235-240.

6.   Denli, A.M., Tops, B.B., Plasterk, R.H., Ketting, R.F. and Hannon, G.J. (2004)

Processing of primary microRNAs by the Microprocessor complex. *Nature*, **432**, 231-

235.

7.   Ladewig, E., Okamura, K., Flynt, A.S., Westholm, J.O. and Lai, E.C. (2012) Discovery

of hundreds of mirtrons in mouse and human small RNA data. *Genome Res*, **22**, 1634-

1645.

8.   Ruby, J.G., Jan, C.H. and Bartel, D.P. (2007) Intronic microRNA precursors that bypass

Drosha processing. *Nature*, **448**, 83-86.

9.   Iwasaki, S., Kobayashi, M., Yoda, M., Sakaguchi, Y., Katsuma, S., Suzuki, T. and

Tomari, Y. (2010) Hsc70/Hsp90 chaperone machinery mediates ATP-dependent RISC

loading of small RNA duplexes. *Mol Cell*, **39**, 292-299.

10.   Pare, J.M., Tahbaz, N., Lopez-Orozco, J., LaPointe, P., Lasko, P. and Hobman, T.C.

(2009) Hsp90 regulates the function of argonaute 2 and its recruitment to stress granules

and P-bodies. *Mol Biol Cell*, **20**, 3273-3284.

11.   Nakanishi, K. (2016) Anatomy of RISC: how do small RNAs and chaperones activate

Argonaute proteins? *Wiley Interdiscip Rev RNA*, **7**, 637-660.

12.   Meijer, H.A., Smith, E.M. and Bushell, M. (2014) Regulation of miRNA strand selection:

follow the leader? *Biochem Soc Trans*, **42**, 1135-1140.

13.   Medley, J.C., Panzade, G. and Zinovyeva, A.Y. (2021) microRNA strand selection:

Unwinding the rules. *Wiley Interdiscip Rev RNA*, **12**, e1627.

14.   Frank, F., Sonenberg, N. and Nagar, B. (2010) Structural basis for 5'-nucleotide base-

1366   specific recognition of guide RNA by human AGO2. *Nature*, **465**, 818-822.

1367   15.   Melendez, A. and Levined, B. (August 24, 2009) In Community, T. C. e. R. (ed.),

1368   *WormBook*. WormBook.

1369   16.   Zinovyeva, A.Y., Veksler-Lublinsky, I., Vashisht, A.A., Wohlschlegel, J.A. and Ambros,

1370   V.R. (2015) Caenorhabditis elegans ALG-1 antimorphic mutations uncover functions for

1371   Argonaute in microRNA guide strand selection and passenger strand disposal. *Proc Natl*

1372   *Acad Sci U S A*, **112**, E5271-5280.

1373   17.   Caceres, J.F., Stamm, S., Helfman, D.M. and Krainer, A.R. (1994) Regulation of

1374   alternative splicing in vivo by overexpression of antagonistic splicing factors. *Science*,

1375   **265**, 1706-1709.

1376   18.   Mangone, M., Macmenamin, P., Zegar, C., Piano, F. and Gunsalus, K.C. (2008)

1377   UTRome.org: a platform for 3'UTR biology in C. elegans. *Nucleic Acids Res*, **36**, D57-

1378   62.

1379   19.   Mangone, M., Manoharan, A.P., Thierry-Mieg, D., Thierry-Mieg, J., Han, T.,

1380   Mackowiak, S.D., Mis, E., Zegar, C., Gutwein, M.R., Khivansara, V. *et al.* (2010) The

1381   landscape of C. elegans 3'UTRs. *Science*, **329**, 432-435.

1382   20.   Steber, H.S., Gallante, C., O'Brien, S., Chiu, P.L. and Mangone, M. (2019) The C.

1383   elegans 3' UTRome v2 resource for studying mRNA cleavage and polyadenylation, 3'-

1384   UTR biology, and miRNA targeting. *Genome Res*, **29**, 2104-2116.

1385   21.   Murari, E., Meadows, D., Cuda, N. and Mangone, M. (2024) A comprehensive analysis

1386   of 3'UTRs in Caenorhabditis elegans. *Nucleic Acids Res*.

1387 22. Kozomara, A. and Griffiths-Jones, S. (2014) miRBase: annotating high confidence

1388    microRNAs using deep sequencing data. *Nucleic Acids Res*, **42**, D68-73.

1389 23. Sulston, J.E., Schierenberg, E., White, J.G. and Thomson, J.N. (1983) The embryonic cell

1390    lineage of the nematode Caenorhabditis elegans. *Dev Biol*, **100**, 64-119.

1391 24. Frokjaer-Jensen, C., Davis, M.W., Hopkins, C.E., Newman, B.J., Thummel, J.M., Olesen,

1392    S.P., Grunnet, M. and Jorgensen, E.M. (2008) Single-copy insertion of transgenes in

1393    Caenorhabditis elegans. *Nat Genet*, **40**, 1375-1383.

1394 25. Kozomara, A., Birgaoanu, M. and Griffiths-Jones, S. (2019) miRBase: from microRNA

1395    sequences to function. *Nucleic Acids Res*, **47**, D155-D162.

1396 26. Mei, Q., Li, X., Meng, Y., Wu, Z., Guo, M., Zhao, Y., Fu, X. and Han, W. (2012) A

1397    facile and specific assay for quantifying microRNA by an optimized RT-qPCR approach.

1398    *PloS one*, **7**, e46890.

1399 27. Kernick, K., Woudstra, R., Berjanskii, M., MacKay, S. and Wishart, D.S. (2025)

1400    Heatmapper2: web-enabled heat mapping made easy. *Nucleic Acids Res*, **53**, W316-

1401    W323.

1402 28. Livak, K.J. and Schmittgen, T.D. (2001) Analysis of relative gene expression data using

1403    real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods*, **25**, 402-408.

1404 29. Kotagama, K., Schorr, A.L., Steber, H.S. and Mangone, M. (2019) ALG-1 Influences

1405    Accurate mRNA Splicing Patterns in the Caenorhabditis elegans Intestine and Body

1406    Muscle Tissues by Modulating Splicing Factor Activities. *Genetics*, **212**, 931-951.

1407 30. Kato, M., Kashem, M.A. and Cheng, C. (2016) An intestinal microRNA modulates the

1408    homeostatic adaptation to chronic oxidative stress in C. elegans. *Aging (Albany NY)*, **8**,

1409    1979-2005.

31. Stubna, M.W., Shukla, A. and Bartel, D.P. (2024) Widespread destabilization of

Caenorhabditis elegans microRNAs by the E3 ubiquitin ligase EBAX-1. *RNA*, **31**, 51-66.

32. Kent, W.J. (2002) BLAT--the BLAST-like alignment tool. *Genome Res*, **12**, 656-664.

33. Lorenz, R., Bernhart, S.H., Honer Zu Siederdissen, C., Tafer, H., Flamm, C., Stadler, P.F.

and Hofacker, I.L. (2011) ViennaRNA Package 2.0. *Algorithms Mol Biol*, **6**, 26.

34. Rishik, S., Hirsch, P., Grandke, F., Fehlmann, T. and Keller, A. (2025)

miRNATissueAtlas 2025: an update to the uniformly processed and annotated human and

mouse non-coding RNA tissue atlas. *Nucleic Acids Res*, **53**, D129-D137.

35. Brosnan, C.A., Palmer, A.J. and Zuryn, S. (2021) Cell-type-specific profiling of loaded

miRNAs from Caenorhabditis elegans reveals spatial and temporal flexibility in

Argonaute loading. *Nat Commun*, **12**, 2194.

36. Shaw, W.R., Armisen, J., Lehrbach, N.J. and Miska, E.A. (2010) The conserved miR-51

microRNA family is redundantly required for embryonic development and pharynx

attachment in Caenorhabditis elegans. *Genetics*, **185**, 897-905.

37. Alvarez-Saavedra, E. and Horvitz, H.R. (2010) Many families of C. elegans microRNAs

are not essential for development or viability. *Curr Biol*, **20**, 367-373.

38. Garcia-Segura, L., Abreu-Goodger, C., Hernandez-Mendoza, A., Dimitrova Dinkova,

T.D., Padilla-Noriega, L., Perez-Andrade, M.E. and Miranda-Rios, J. (2015) High-

Throughput Profiling of Caenorhabditis elegans Starvation-Responsive microRNAs. *PloS*

one, **10**, e0142262.

39. Garcia, S.M., Tabach, Y., Lourenço, G.F., Armakola, M. and Ruvkun, G. (2014)

Identification of genes in toxicity pathways of trinucleotide-repeat RNA in C. elegans.

Nature structural & molecular biology, **21**, 712-720.

40. Kagias, K., Podolska, A. and Pocock, R. (2014) Reliable reference miRNAs for

quantitative gene expression analysis of stress responses in Caenorhabditis elegans. *BMC*

Genomics, **15**, 222.

41. Blazie, S.M., Geissel, H.C., Wilky, H., Joshi, R., Newbern, J. and Mangone, M. (2017)

Alternative Polyadenylation Directs Tissue-Specific miRNA Targeting in Caenorhabditis

elegans Somatic Tissues. *Genetics*, **206**, 757-774.

42. Haenni, S., Ji, Z., Hoque, M., Rust, N., Sharpe, H., Eberhard, R., Browne, C., Hengartner,

M.O., Mellor, J., Tian, B. *et al.* (2012) Analysis of C. elegans intestinal gene expression

and polyadenylation by fluorescence-activated nuclei sorting and 3'-end-seq. *Nucleic*

Acids Res, **40**, 6304-6318.

43. Nakanishi, K. (2024) When Argonaute takes out the ribonuclease sword. *J Biol Chem*,

300, 105499.

44. Krishnan, K., Steptoe, A.L., Martin, H.C., Wani, S., Nones, K., Waddell, N.,

Mariasegaram, M., Simpson, P.T., Lakhani, S.R., Gabrielli, B. *et al.* (2013) MicroRNA-

182-5p targets a network of genes involved in DNA repair. *RNA*, **19**, 230-242.

45. Krist, B., Florczyk, U., Pietraszek-Gremplewicz, K., Jozkowicz, A. and Dulak, J. (2015)

The Role of miR-378a in Metabolism, Angiogenesis, and Muscle Biology. *Int J*

Endocrinol, **2015**, 281756.

46. Rane, S., He, M., Sayed, D., Vashistha, H., Malhotra, A., Sadoshima, J., Vatner, D.E.,

Vatner, S.F. and Abdellatif, M. (2009) Downregulation of miR-199a derepresses

hypoxia-inducible factor-1alpha and Sirtuin 1 and recapitulates hypoxia preconditioning

in cardiac myocytes. *Circ Res*, **104**, 879-886.

47.   Makeyev, E.V., Zhang, J., Carrasco, M.A. and Maniatis, T. (2007) The MicroRNA miR-

1456   124 promotes neuronal differentiation by triggering brain-specific alternative pre-mRNA

1457   splicing. *Mol Cell*, **27**, 435-448.

1458   48.   Rhim, J., Baek, W., Seo, Y. and Kim, J.H. (2022) From Molecular Mechanisms to

1459   Therapeutics: Understanding MicroRNA-21 in Cancer. *Cells*, **11**.

1460   49.   Brueckner, B., Stresemann, C., Kuner, R., Mund, C., Musch, T., Meister, M., Sultmann,

1461   H. and Lyko, F. (2007) The human let-7a-3 locus contains an epigenetically regulated

1462   microRNA gene with oncogenic function. *Cancer Res*, **67**, 1419-1423.

1463   50.   Kato, M., Paranjape, T., Muller, R.U., Nallur, S., Gillespie, E., Keane, K., Esquela-

1464   Kerscher, A., Weidhaas, J.B. and Slack, F.J. (2009) The mir-34 microRNA is required for

1465   the DNA damage response in vivo in C. elegans and in vitro in human breast cancer cells.

1466   *Oncogene*, **28**, 2419-2424.

1467   51.   Isik, M., Blackwell, T.K. and Berezikov, E. (2016) MicroRNA mir-34 provides

1468   robustness to environmental stress response via the DAF-16 network in C. elegans. *Sci

1469   Rep*, **6**, 36766.

1470   52.   Bommer, G.T., Gerin, I., Feng, Y., Kaczorowski, A.J., Kuick, R., Love, R.E., Zhai, Y.,

1471   Giordano, T.J., Qin, Z.S., Moore, B.B. *et al.* (2007) p53-mediated activation of

1472   miRNA34 candidate tumor-suppressor genes. *Curr Biol*, **17**, 1298-1307.

1473   53.   Spurgers, K.B., Gold, D.L., Coombes, K.R., Bohnenstiehl, N.L., Mullins, B., Meyn, R.E.,

1474   Logothetis, C.J. and McDonnell, T.J. (2006) Identification of cell cycle regulatory genes

1475   as principal targets of p53-mediated transcriptional repression. *J Biol Chem*, **281**, 25134-

1476   25142.

54.   He, L., He, X., Lowe, S.W. and Hannon, G.J. (2007) microRNAs join the p53 network--

another piece in the tumour-suppression puzzle. *Nat Rev Cancer*, **7**, 819-822.

55.   Kim, H., Kim, J., Yu, S., Lee, Y.Y., Park, J., Choi, R.J., Yoon, S.J., Kang, S.G. and Kim,

V.N. (2020) A Mechanism for microRNA Arm Switching Regulated by Uridylation. *Mol*

Cell, **78**, 1224-1236 e1225.

56.   Mullokandov, G., Baccarini, A., Ruzo, A., Jayaprakash, A.D., Tung, N., Israelow, B.,

Evans, M.J., Sachidanandam, R. and Brown, B.D. (2012) High-throughput assessment of

microRNA activity and function using microRNA sensor and decoy libraries. *Nat*

Methods, **9**, 840-846.

57.   Krol, J., Loedige, I. and Filipowicz, W. (2010) The widespread regulation of microRNA

biogenesis, function and decay. *Nat Rev Genet*, **11**, 597-610.

58.   Mi, S., Cai, T., Hu, Y., Chen, Y., Hodges, E., Ni, F., Wu, L., Li, S., Zhou, H., Long, C. *et*

al. (2008) Sorting of small RNAs into Arabidopsis argonaute complexes is directed by

the 5' terminal nucleotide. *Cell*, **133**, 116-127.

59.   Okamura, K., Liu, N. and Lai, E.C. (2009) Distinct mechanisms for microRNA strand

selection by Drosophila Argonautes. *Mol Cell*, **36**, 431-444.

60.   Kotagama, K., Grimme, A.L., Braviner, L., Yang, B., Sakhawala, R.M., Yu, G., Benner,

L.K., Joshua-Tor, L. and McJunkin, K. (2024) Catalytic residues of microRNA

Argonautes play a modest role in microRNA star strand destabilization in C. elegans.

Nucleic Acids Res, **52**, 4985-5001.

61.   Stein, L., Mangone, M., Schwarz, E., Durbin, R., Thierry-Mieg, J., Spieth, J. and

Sternberg, P. (2001) WormBase: network access to the genome and biology of

Caenorhabditis elegans (vol 29, pg 82, 2001). *Nucleic Acids Research*, **29**, 1012.

1500

Main Figure 1

A

B

miR-52

miR-75

C

Variant	5p	3p
WT	0.85	0.15
UP11-12	0.71	0.29
5'U → C	0.71	0.29
Combined	0.68	0.32

D

Main Figure 5

A

3p Singletons | Switchers | 5p Singletons

9 | 835 | 10

frequency

0.0–0.03, 0.07–0.1, 0.13–0.17, 0.2–0.23, 0.27–0.3, 0.33–0.37, 0.4–0.43, 0.47–0.5, 0.53–0.57, 0.6–0.63, 0.67–0.7, 0.73–0.77, 0.8–0.83, 0.87–0.9, 0.93–0.97

B

mean feature values

5p_G

0.5, 0.375, 0.25, 0.125, 0

H. sapiens, C.elegans

di_CC

6, 4.5, 3, 1.5, 0

H. sapiens, C.elegans

di_GG

7, 5.25, 3.5, 1.75, 0

H. sapiens, C.elegans

3p_has_CCU

1.2, 0.9, 0.6, 0.3, 0

H. sapiens, C.elegans

C

```
cel-let-7-5p    UGAGGUAGUAGGUUGUAUAGUU
let-7a-5p       ......................
let-7b-5p       ..................G..G...
let-7c-5p       ...................G...
let-7d-5p       A................C...
let-7e-5p       .......G..........

cel-let-7-3p    CUAUGCAAUUUUCUACCUUACC
let-7a-3p       ....A.......C.A..GA..A.
let-7b-3p       ....A......CCUA...GCCU.C.
let-7c-3p       ..G.A......CCUA...G...
let-7d-3p       ....A.G..CCUG..GCCU...
let-7e-3p       ....A.GGCCUCCUG.UUUCC
```

D

mean values of each feature

0.0 0.2 0.4 0.6 0.8 1

feature differences between 5p Singletons , 3p Singletons and Switchers

length, gc_content, au_content, five_p_mfe, three_p_mfe, gc_5p, gc_3p, shannon_entropy_5p, shannon_entropy_3p, pairing_frac_5p, pairing_frac_3p, length_diff, mfe_per_nt_5p, mfe_per_nt_3p, 5p_A, 5p_U, 5p_G, 5p_C, 3p_A, 3p_U, 3p_G, 3p_C, GC_diff, GC_ratio_5p3p, mfe_diff, mfe_ratio, mfe_per_nt_diff, 5p_has_UGU, 3p_has_UGU, 5p_has_GAG, 3p_has_GAG, 5p_has_AAU, 3p_has_AAU, 5p_has_UAA, 3p_has_UAA, 5p_has_GUG, 3p_has_GUG, 5p_has_CGC, 3p_has_CGC, 5p_has_CCU, 3p_has_CCU, di_UG, di_GA, di_AG, di_GG, di_GU, di_UA, di_UU, di_AU, di_UC, di_CU, di_AC, di_CA, di_AA, 5p_first_nt_A, 5p_first_nt_U, 5p_first_nt_G, 5p_first_nt_C, 5p_first_nt_, 5p_last_nt_A, 5p_last_nt_U, 5p_last_nt_G, 5p_last_nt_C, 5p_last_nt_, 3p_first_nt_A, 3p_first_nt_U, 3p_first_nt_G, 3p_first_nt_C, 3p_first_nt_, 3p_last_nt_A, 3p_last_nt_U, 3p_last_nt_G, 3p_last_nt_C, 3p_last_nt_, di_CC, di_GC, di_CG

c h | c h | c h

Switchers | Singletons

5p | 3p

Singletons

Main Figure 6

A

cel-miR-34 3p 76%

```
    A GGCAGUG U  U       G
      |||||||           UUG
      GGCAGUG GG UAGCUG  UUG
      |||||||  :|  |||||:||
C     CCGUCAC UC AUCGGC  A
 C CA              U  C   -
```

CG_5p	CG_3p	% AU	di_GU
54	63	40	.09

dme-miR-34 5p 68%

```
U GGCAGUG U G U UAG C UGGUUGUG
  |||||||     :|  |||  |||:||
  GCCGUCAC UC AUC  ACCGAC
CC          U  U       -
```

CG_5p	CG_3p	% AU	di_GU
54	60	42	.10

mmu-miR-34a 5p 76%

```
U GGCAGUGU  -   A
  ||||||||  |||  |||||:|
  CCGUCAUA GAA CGACUAA
UC        U  -
```

CG_5p	CG_3p	% AU	di_GU
50	42	53	.12

hsa-miR-34a 5p 82%

```
U GGCAGUGU - CUU A GCUGGUUGU
  ||||||||: |||  |||||:|||
  CCGUCAUA GAA CGACUAAC
UC        U
```

CG_5p	CG_3p	% AU	di_GU
50	45	52	.12

B **GC asymmetry (arm inversion)**

GC asymmetry flips sign

C **Rigidification**

5p stem becomes more paired