Exploring MicroRNA Strand Selection: Insights into Gene Regulation

and Development

Author list

Dalton Meadows[1,2],

Hailee Hargis[1,2],

Amanda Ellis[1,2],

Marco Mangone[1,3]

Author affiliations

[1)] The Biodesign Institute at Arizona State University, 1001 S McAllister Ave, Tempe, AZ

[2)] School of Life Sciences, Arizona State University, 427 E Tyler Mall, Tempe, AZ

Author list footnotes

[3)] Lead contact

Corresponding author and e-mail address

Marco Mangone; mangone@asu.edu

Abstract

MicroRNAs (miRNAs) are important post-transcriptional regulators that play key roles in development, cell identity, and disease. While significant progress has been made in identifying miRNA targets, the mechanisms governing miRNA strand selection during biogenesis remain poorly understood. Here we investigate the hypothesis that miRNA strand selection is not random but follows specific thermodynamic and regulatory rules that vary across tissues and developmental stages.

Using *Caenorhabditis elegans* as a model system, we employed a novel high-throughput qPCR-based approach (HiTmiSS) to quantify miRNA strand selection throughout development and within somatic tissues, coupled with a novel machine learning model trained with our validated wet bench data, that can predict miRNA strand selection with high accuracy. When applied to vertebrates, our ML model has identified specific conserved features with important evolutionary implications for miRNA biogenesis. Our findings indicate that miRNA strand preference is widespread but varies dynamically, with a subset of miRNAs switching strand usage depending on developmental stage or tissue types. Furthermore, our analysis suggests that thermodynamic asymmetry alone does not fully explain strand selection, implicating additional cellular factors. This work provides a comprehensive framework for studying miRNA biogenesis, offering insights into regulatory principles governing strand selection and its broader implications in gene regulation, development, and disease across metazoans.

Introduction

miRNAs are short non-coding RNAs, ~22 nucleotides in length, that are incorporated into a large protein complex called the RNA-induced silencing complex (RISC), where they guide the interaction between the RISC and target mRNA by complementary base pairing primarily within 3' Untranslated (3'UTRs) regions in mature mRNAs (Bartel 2018). mRNAs targeted by the miRNA/RISC complex (miRISC) are typically held in translational repression. miRNAs and their 3'UTR targets are frequently conserved and play various roles in regulating fundamental biological processes across metazoans, such as development and morphogenesis.

miRNAs have undergone several pronounced expansions in metazoan genomes (Wolter et al. 2017). These expansions coincide with evolving organismal complexity yet rarely correspond with increases in protein-coding genes (Wolter et al. 2017). Eukaryotic genomes contain 100-2,000 distinct mature miRNAs in intergenic regions, introns, and operon-like clusters. Following transcription in long primary miRNA transcripts (pri-miRNAs) by the RNA polymerase II, they are processed in the nucleus by the microprocessor complex, comprising the RNase III enzyme Drosha and its cofactor DGCR8, into shorter hairpin structures known as precursor miRNAs (pre-miRNAs) (Bartel 2018). The two strands composing the pre-miRNAs are named the 5p and the 3p strand, respectively.

Pre-mRNAs are subsequently exported to the cytoplasm by Exportin-5, where they undergo further cleavage by the enzyme Dicer, producing mature miRNA duplexes (Bartel 2018). The Hsc70-Hsp90 chaperone complex facilitates the ATP-dependent

loading of the miRNA duplex into Argonaute, the protein that execute the silencing step.

This complex is thought to induce conformational changes in Argonaute that enable

duplex loading (Pare et al. 2009; Meijer et al. 2014; Nakanishi 2016). Once loaded, one

strand of this duplex, the guide strand, is kept within the RISC, while the other, the

passenger strand, is typically degraded. Both the 5p and the 3p strands can be selected

for miRISC loading and act as guide strands. It is currently unclear how this process,

termed 'miRNA strand selection,' is executed in detail, which proteins are involved in

this decision, and how the guide strand is ultimately chosen between the two strands.

The mature miRNA within the RISC complex guides the complex to target mRNAs,

leading to either translational repression or mRNA degradation, thereby influencing a

wide array of cellular processes and contributing significantly to the regulation of gene

expression. MiRNA biogenesis is a fundamental mechanism with far-reaching

implications in development, homeostasis, and disease (Bartel 2018).

Recently, several groups (including ours) produced tissue-specific localization data

for many miRNAs in *C. elegans* (Blazie et al. 2015; Blazie et al. 2017; Kotagama et al.

2019; Zhang et al. 2022; Schorr et al. 2023), flies (Carthew et al. 2017), mice (Isakova

et al. 2020), rats (Smith et al. 2016), human tissues (Jima et al. 2010), and cancer

(Ilieva et al. 2022). These results show that there are distinct functional miRNA

populations in tissues, which can reshape transcriptomes and contribute to cell identity

acquisition and maintenance. Since most of their targets are only predicted, how these

events are initiated and maintained is still being determined.

miRNAs guide the RISC to complementary sites on target mRNAs by standard

Watson-Crick and G:U wobble pairing. This pairing has been shown to require as little

as six consecutive nucleotides in the 5' end, or seed region, of the mature miRNA.

Perfect complementarity between the miRNA seed region and its target is considered

the canonical indicator of miRNA targeting. However, many recent studies indicate that

miRNAs can target non-canonical elements in target mRNA (Wolter et al. 2017).

Because these recognition elements are degenerate and small, each miRNA can

potentially target hundreds of genes, dramatically reshaping cellular gene output. In this

context, loading the correct miRNA strand into a RISC complex is extremely important

since the 5p and the 3p target a completely different set of genes.

What mechanisms control Argonaute's choice in loading one strand of the miRNA

duplex over the other? The predominant factors influencing miRNA strand selection are

believed to be the specific 5' nucleotide sequences of each strand and the comparative

thermodynamic stability exhibited by the two ends of the miRNA duplex (Medley et al.

2021). Studies performed using the human Ago2 protein, an Argonaute protein member

of the RISC complex, show that the pocket responsible for binding the 5' nucleotide of

the guide strand exhibits a strong preference for Uracil nucleotides (Frank et al. 2010).

This aligns with the prevalent occurrence of Uracil as the 5' nucleotide in miRNA

guide strands (Wolter et al. 2017), whereas the passenger strand commonly initiates

with a Cytosine at its 5' end.

Furthermore, the miRISC complex exhibits a preference for loading the end of the

miRNA duplex with lower thermodynamic stability, although not all the time (Melendez

and Levined August 24, 2009). The initial four nucleotides on each end of the duplex

are believed to establish this thermodynamic asymmetry, and in some cases, a slight

difference in the number of hydrogen bonds can induce the preferential loading of the

less stable strand *in vitro* (Meijer et al. 2014).

Consistently, the 5' ends of guide strands are enriched with purines (REF). In

comparison, the 5' ends of passenger strands are predominantly pyrimidine-rich in both

human and fly miRNAs, contributing to the thermodynamic asymmetry observed

between the two ends of the duplex. In *C. elegans*, single amino acid substitutions

within the MID (G553R) and PIWI (G716E, S895F, and S939F) domains of the

Argonaute protein ALG-1 invert the strand selection process for specific miRNAs,

suggesting a potential role for this protein in miRNA strand selection process

(Zinovyeva et al. 2015). This indicates that while these mutations play a pivotal role,

they do not consistently reverse strand selection for all miRNAs, suggesting that there

may be additional internal or external factors that impact the selection of miRNA strand

selection in *C. elegans*, depending on the specific miRNA involved. Tissue specific

accessory factors could also be responsible of miRNA strand selection, altering strand

choices based on tissue or developmental contexts, similarly to the role of the SR

proteins in RNA splicing.

The nematode *C. elegans* represents an attractive system to study miRNA strand

selection. This organism was the first multicellular system with an annotated 3'UTR

reference dataset (3'UTRome), which was originally developed in 2011 within the

modENCODE Project (Mangone et al. 2008; Mangone et al. 2010) and refined later on

(Steber et al. 2019; Murari et al. 2024). The number of miRNAs is relatively small when

compared to humans (253 miRNAs versus 1,917 in humans (miRbase)), and the size of

3'UTRs is also small (average size in *C. elegans* 230nt, ~1,5k in humans). In addition,

C. elegans has a complete cell-lineage map (Sulston 1983), its development is well

studied at the physiological and molecular level (Chalfie et al. 1981), it has small,

relatively simple tissues, and its transcriptome and 3'UTRome have been extensively

characterized (Gerstein et al. 2010; Mangone et al. 2010; Jan et al. 2011; Ramani et al.

2011; Steber et al. 2019). *C. elegans* also possesses the best-annotated miRNA target

predictions available for downstream studies (Bartel 2018). It has homologs for all

members of the known human RISC complex (Grishok 2005) and is amenable to

genetic manipulation.

After years of research, we have begun to understand the role of thermodynamics

and nucleotide composition in this process, which can explain some but not all events.

However, we still lack a general mechanistic framework that explains how miRNA

strand selection is executed in eukaryotes.

Here, we developed a novel high-throughput approach to profile miRNA strand

selection through *C. elegans* development and in somatic tissues. We have identified,

pattern, motifs and trends that miRNA use in this process. These results were then used

to train a machine learning model capable of accurately predicting strand usage with

83% accuracy and use it to perform a comparative analysis of miRNA strand selection

between *C. elegans* and humans.

Materials and methods

All DI water was obtained from either a Purelab Option system (ELGA LabWater) or

an Avidity Science system (Duo) and autoclaved before use. Chemicals were ordered

from Sigma Aldrich/Merck, VWR, or Thermo Fisher Scientific in molecular biology grade

and stored according to company instructions. Gateway[TM] LR Clonase II Plus[TM] enzyme

mixes, SuperScript[TM] III Reverse Transcriptase, and PowerTrack[TM] SYBR Green Master

Mix were purchased from Invitrogen. Primers for the HiTmiSS assay were ordered from

Life Technologies. Poly(U) Polymerase and restriction enzymes used in validation were

ordered from New England Biolabs. The pCFJ150 plasmid was a gift from Erik

Jorgensen. The OP50-1 *E. coli*, N2 *C. elegans* and EG6699 *C. elegans* strains were

purchased through the Caenorhabditis Genetics Center.

Maintenance of *C. elegans* strains

All *C. elegans* strains were cultured on NGM plates seeded with OP50-1 bacteria. All

transgenic lines produced for experiments described in this study were maintained at

$24°$C. When bacteria on each NGM plate had been cleared, worms were transferred

onto new plates via washing into a 15 ml Falcon tube using 15 ml M9 media (22 mM

KH_2PO_4, 42 mM Na_2HPO_4, 85.5 mM NaCl, 1 mM $MgSO_4$, sterilized by autoclave) and

pelleted for 3 minutes at 400g. The M9 media was then aspirated to a volume of 500 ml

and the animals were resuspended via gentle pipetting before being dropleted onto

fresh, seeded NGM plates.

Synchronization of *C. elegans* strains

The EG6699 strain (Frokjaer-Jensen et al. 2008) was also maintained at 24°C. To

synchronize animals for injection, confluent plates were bleached to isolate the

embryos. Each plate was washed into a 15 ml Falcon tube with 15 ml DI water and

pelleted for 3 minutes at 400*g*. The water was then aspirated and the animals were

incubated in 5 ml of bleaching solution (0.5M NaOH, 10% bleach) for 20 minutes with

gentle end-over-end rotation. After bleaching, the embryos were washed twice with DI

water and dropleted onto fresh, seeded NGM plates.

For use in the HiTmiSS assay, both N2 and *pges-1::GFP::alg-1::unc54 3′ UTR C.*

elegans strains were synchronized as described above.

Primers design for the HiTmiSS Assay

To quantify microRNAs in high throughput, we designed 380 unique forward primers

for use in the qPCR amplification step of the HiTmiSS assay. These primers were

designed to be complimentary to either the 5p strand or the 3p strand of 190 most

expressed microRNAs in the *C. elegans* genome, corresponding to 75% of all *C.*

elegans miRNA deposited in miRbase (release 22.1). We did not consider microRNAs

whose existence is not supported by broad experimental evidence nor any microRNAs

predicted to exist. Few miRNAs were ignored because their corresponding primers had

a very low Tm, making it impossible to process using our pipeline. All primers were

ordered in four 96-well microplates from Life Technologies pre-diluted at a concentration

of 50 mM (**Supplemental Table S1**). We designed a universal reverse primer

complementary to the stem-loop sequence of the stem-loop poly(A) primer used to

prepare the cDNA for this qPCR reaction. 24 replicates of this primer were ordered from

Life Technologies in a separate array, also at a concentration of 50 mM. Both forward

and reverse primers were re-arrayed and diluted to a working concentration of 10 mM

immediately prior to use.

***Singletons* and *Switchers* identification**

For each strand, we subtracted its relative quantity at one life stage versus the

relative quantity of the complementary strand (i.e. for *let-7* 5p, we compared the

abundance at embryo stage versus the abundance of *let-7* 3p at the embryo stage).

This was repeated for each life stage at which the expression of that microRNA strand

fell above our significance threshold, and any microRNAs for which their expression

was greater at each such life stage were deemed *Singletons*. For microRNAs where the

5p strand was more highly quantified when significant at each life stage, these were

classified as *5p Singletons*. For microRNAs where the 3p strand was more highly

quantified when significant at each life stage, these were classified as *3p Singletons*.

microRNAs which rose above our threshold of significance during at least one life stage,

but whose expression levels did not display a consistent pattern of usage of one strand

over the other were designated *Switchers*.

HiTmiSS Assay

To collect sufficient RNA for analysis of wild type microRNA, confluent NGM plates

of N2 *C. elegans* were synchronized and either washed and pelleted immediately

(embryo stage) or incubated at 24°C for 15 hours (larval 1 stage), 25 hours (larval 2

stage), 33 hours (larval 3 stage), 42 hours (larval 4 stage), or 51 hours (young adult

stage) before washing and pelleting. All washes were performed in 15 ml Falcon tubes

using 15 ml DI water and pellets were generated by centrifugation for 3 minutes at

400*g*, resulting in a final pellet volume of 500 μl to 1 ml. Sufficient worms to generate

this size pellet differed by life stage, requiring 12 (larval 4 and young adult stages), 18

(larval 2 and larval 3 stages), or 24 (embryo and larval 1 stages) starting confluent

plates.

This pellet was then resuspended to a volume of 1.5 ml in ice cold Triton X-100 lysis

buffer (150 mM NaCl, 1.0% Triton X-100, 50 mM Tris-HCl pH 8.0, 0.5 mM EDTA) and

transferred to a 2 ml Spectrum tube pre-filled with 0.5 mm zirconium beads. Using a

Benchmark BeadBug D1030 microtube homogenizer, each tube was homogenized for

90 seconds, incubated for 2 minutes on ice, then homogenized for an additional 60

seconds. The resultant lysate mixture was centrifuged for 3 minutes using a VWR

benchtop mini centrifuge and the liquid fraction was transferred into a 1.5 ml Eppendorf

microcentrifuge tube. To minimize non-aqueous contaminants, this lysate was

centrifuged again in the same manner and transferred to a 15 ml Falcon tube pre-chilled

on ice.

To isolate total RNA, we used a DirectZol RNA MiniPrep Plus kit from Zymo

Research, following kit instructions and performing the entire procedure at 4°C.

Resultant RNA was eluted in 30 μL NF H_2O. To collect tissue-specific microRNA for

analysis of the intestine, we performed protein immunoprecipitation followed by

microRNA isolation. The fluorescent *pges1::gfp::alg-1::unc54 3′UTR* strain (PMID:

31073019) of *C. elegans* previously prepared by our lab was grown to confluence,

synchronized, and pelleted as described above for each life stage, concurrent with N2

C. elegans as a negative control.

Each pellet was resuspended to 1.5 µl using our Triton X-100 lysis buffer with added

protease inhibitor (two Roche cOmplete ULTRA mini EDTA-free protease inhibitor

tablets per 50 ml lysis buffer) and transferred to a zirconium bead tube to be

homogenized as previously described. During this homogenization step, anti-GFP

magnetic antibody beads (Chromotek GFP-Trap Magnetic Agarose) were prepared by

diluting 25 µl bead slurry in 500 µl dilution buffer (150 mM NaCl, 50mM Tris-HCl pH 8.0,

0.5 mM EDTA) and rotating end-over-end at 4°C for 3 minutes.

The beads were pelleted using a benchtop centrifuge and buffer was aspirated using

a magnetic tube rack. 500 µl dilution buffer were added once again, and this dilution

process was repeated two more times. We then recovered aqueous lysate from the

homogenized mixture as previously described and diluted it using 300 µl dilution buffer.

This diluted lysate was transferred to a tube containing the prepared anti-GFP beads

and left to rotate end-over-end at 4°C overnight. To clear unbound lysate, the beads

were pelleted using a benchtop centrifuge and buffer was aspirated using a magnetic

tube rack. 500 µl of wash buffer (150 mM NaCl, 50 mM Tris-HCl pH 8.0, 0.1% Triton X-

100, 0.5 mM EDTA) was added, the beads were pelleted, and the buffer was aspirated

using a magnetic tube rack. This wash step was repeated five additional times. The

beads were then resuspended in 100 µl wash buffer and RNA was isolated using a

DirectZol RNA MiniPrep Plus kit from Zymo Research, following kit instructions and

performing the entire procedure at 4°C. Resultant RNA was eluted in 30 μL NF H$_2$O.

For either whole worm or tissue-specific analysis, the remainder of the HiTmiSS

protocol was the same. We prepared our cDNA from each life stage for qPCR analysis

via polyuridinylation using Poly(U) Polymerase (NEB) as per manufacturer instructions

followed by a 20-minute heat inactivation at 65°C. We performed a first-strand reaction

on this polyuridinylated RNA using Superscript III Reverse Transcriptase and stem-loop

poly(A) primers under recommended conditions (Invitrogen).

We then performed qPCR in high throughput using this cDNA, SYBR Green master mix

(Invitrogen), and our primers as described above. We used Applied Biosystems

Quantstudio 3 and Quantstudio 7 qPCR machines to perform the screen. For each life

stage, four 96-well plates were prepared with 40 μL reactions in each well, along with a

positive control reaction containing 10 pg pDONR 221 ROG instead of cDNA and the

necessary forward and reverse primers to amplify *gfp*. Each plate was then re-arrayed

in triplicate into a 384-well plate using a Biomek FX^P Laboratory Automation

Workstation for a total of three 10 μL reactions per microRNA strand per life stage. An

additional plate was prepared with no cDNA or plasmid spike-in DNA as a non-template

control.

These qPCR reactions were carried out under standard SYBR Green reagent

conditions on an Applied Biosystems QuantStudio 7. Relative microRNA levels were

calculated using the 2$^{-\Delta\Delta Ct}$ method as described above and normalized using the spike-

in positive control. These results were then plotted logarithmically as a heatmap. Whole

worm data were ordered by type (*5p Singleton*, *3p Singleton*, or *Switcher*), while

intestine data were clustered using average linkage. Both data were visualized using

heatmapper.ca.

Cycle Threshold Analysis of the HiTmiSS Assay

The qPCR results containing the cycle threshold (Ct) values for each tested miRNA

was exported in Microsoft Excel 2019 format. A comprehensive list of these data can be

found in **Supplemental Table S2**. Ct values were averaged across experimental

replicates, then subjected to a $2^{-\Delta\Delta Ct}$ analysis to calculate initial amounts of each

microRNA in the sample, consistent with experimental standard (Livak and Schmittgen

2001). The Ct value for any given microRNA strand was subtracted from the Ct value of

the non-template control (a qPCR reaction lacking any cDNA) for that microRNA strand.

This was termed the ΔCt for that microRNA. The same was done for the positive control

for that sample. Then for each sample this difference was subtracted from the difference

obtained for that positive control, normalizing samples to the same positive control and

giving us the $\Delta\Delta Ct$ for that microRNA strand.

The inverse log base 2 of that $\Delta\Delta Ct$ was then calculated to approximate the initial

amount of that microRNA strand present in the cDNA relative to the positive control.

This was deemed the $2^{-\Delta\Delta Ct}$ value for that microRNA strand.To determine whether a

microRNA was considered present in a sample, we decided on a strict threshold of one

standard deviation above the mean of all $2^{-\Delta\Delta Ct}$ values for that stage of development.

MicroRNAs quantified at levels above this threshold were considered significant for the

purposes of this study, while microRNAs not quantified at levels above this threshold

were not considered significant for the purposes of this study. This same process was

repeated to determine significance for our intestine tissue-specific HiTmiSS assay.

qPCR of microRNAs targeting overexpressed intestinal 3′UTRs

To quantify the level of certain microRNA strands when the target 3′ UTRs of those

strands are overexpressed in the intestine, we used two *C. elegans* strains previously

published by our lab (Kotagama et al. 2019). Each expresses the promoter for intestine-

specific gene *ges-1* fused to our lab's dual color ROG construct (a gene encoding

mCherry and *gfp* separated by a trans-spliceable operon) fused to the 3′ UTR of either

fox-1 or *hrp-2*. Each of these strains was grown to confluence in 12 NGM plates in

mixed stage and subjected to quantification using the same approach as the HiTmiSS

assay described above, though rather than quantifying every microRNA in these

animals, we quantified the 5p and 3p strand of *miR-54*, *miR-60*, *miR-232*, and *miR-*

239b. These quantifications were carried out in 96-well plates using cDNA (both

prepared from our strains as well as control cDNA gathered from N2 *C. elegans*), SYBR

Green master mix, and the appropriate forward and universal reverse primers for the

miRNAs of interest, rearrayed from our standard HiTmiSS plates.

Each 96-well plate contained two positive controls in *let-7* and *lin-4*, both of which

are known highly expressed microRNAs in the intestine, as well as a non-template

negative control for each set of primers. These qPCRs were carried out in experimental

triplicate under standard SYBR Green reagent conditions on an Applied Biosystems

QuantStudio 3.

Relative microRNA levels were calculated using the $2^{-\Delta\Delta Ct}$ method as described

above and normalized using the average of *let-7* and *lin-4* expression for each sample.

These results were then averaged across two replicates and analyzed using Microsoft

Excel 2019. Data were visualized as $\log_2$ of the change in expression from wildtype N2

negative control to either of our experimental strains. The threshold of significance was

set at 1 as per standard analytical procedure.

Structural Analysis of microRNAs

The experimentally validated *5p* and *3p Singletons* were obtained using our

HiTmiSS assay. Each entry included the full hairpin sequence and its predicted

secondary structure in dot-bracket notation (**Supplemental Table S2)**. To focus

specifically on mature miRNA regions, only the sequences of each mature miRNA

duplex was used for all downstream analyses. Since the *5p* and *3p Singleton* datasets

contained different numbers of miRNAs, we balanced them prior to analysis by

undersampling the larger group to match the sample size of the smaller class, ensuring

unbiased statistical comparison between 5p and 3p strands.

We extracted 77 numerical features from each *5p* and *3p Singleton* (**Supplemental**

Table S3). The extracted features included: Base composition (fraction of A, U, G, C),

motif presence (e.g. UGU, GAG, CCU), positional characteristics (first/last nucleotide

identity), dinucleotide frequencies (e.g., di_GG, di_CC), GC content and GC-related

asymmetry metrics (GC_diff, GC_ratio_5p3p), strand lengths, and number of

mismatches (unpaired positions in the dot-bracket structure). The full list of the features

and their descriptions are shown in **Supplemental Table S3**.

Feature calculations were implemented in Python using regular expressions and

sequence statistics functions. Structural mismatch counts were determined by counting

unpaired characters within the uppercase-aligned mature duplex. Comparative

statistical analysis between 5p and 3p strands was conducted using Welch's two-

sample t-test. All features were tested independently, and results were ranked by p-

value. Features with $p < 0.05$ were considered statistically significant. Bar plots were

generated for each significant feature using the mean ± standard deviation across

strands.

miRNA strand selection ML model

To predict dominant strand selection (5p vs. 3p) in miRNA hairpins, we utilized

ChatGPT GPT-4o (OpenAI) to assist in developing and training a Random Forest

classifier based on 77 biologically curated features, organized into functional categories

capturing distinct structural and sequence-related aspects of miRNA hairpins. These

features capture thermodynamic stability, base-pairing patterns, and positional

relationships within the miRNA hairpin. The model was trained on our validated *5p* and

3p Singletons identified using our HiTmiSS assay, and feature values were computed

using custom Python scripts that parse dot-bracket RNA structures and extract both

absolute and relative measures for each strand. The final model incorporates the

following features divided into five categories. The first category is the *Dinucleotide*

Composition features, which quantify the frequency of specific two-nucleotide motifs

within the 5p and 3p regions, capturing local sequence patterns that can influence

strand selection and structural flexibility. The second category is the *First/Last*

Nucleotide Bias features, which record the identity of the first and last nucleotides in

each strand, reflecting known preferences in Argonaute loading where terminal

nucleotides strongly impact strand choice. The third category is the *GC Content and*

Entropy features, which assess the global nucleotide composition and thermodynamic

diversity within each strand, including overall GC percentages, GC ratios between

strands, and Shannon entropy measures that reflect structural variability and rigidity.

The fourth category is the *Base-pairing Properties* feature, which describe the fraction of

nucleotides engaged in base pairing within the 5p and 3p regions, providing a measure

of local hairpin stability and asymmetry. Finally, the *General Structural Features*, which

include the overall hairpin length and the length difference between the 5p and 3p

strands, capturing broader geometrical properties that may indirectly affect miRNA

maturation and strand retention (**Supplemental Figure S1**). The complete set of

features, their description and the weight used are shown in **Supplemental Table S3**.

The model showed high performance in distinguishing 5p and 3p usage from a pool of

known *Singletons* as test dataset, (*5p Singleton*: accuracy 100%, precision 75%, recall

100%; *3p Singleton:* 82% accuracy, 100% precision, 82% recall) (**Supplemental Figure**

S2).

ML model predictions in human miRNAs

Human microRNA (miRNA) expression data were obtained from a harmonized dataset

from the human miRNA tissue atlas (Rishik et al. 2025) containing strand-specific

Reads Per Million Mapped (RPMM) values across healthy human tissues. To restrict

the analysis to non-diseased samples, we only considered entries where the Biotype

column was labeled as "tissue" or "EV", "cell_line" (**Supplemental Table S6**),

representing the average expression values across biological replicates of healthy

tissues (total=337 datasets). For each miRNA duplex, we extracted the expression

levels corresponding to the 5p and 3p strands, based on strand-specific annotations.

We assigned strand dominance by summing RPMM values across all healthy tissue

samples for the 5p and 3p strands separately. The strand with the higher total

expression was designated as the dominant strand. We compared these dominat

strands assignments with the predicted dominant strand assigned by our ML model.

Each prediction was considered correct if the model-assigned strand matched the

strand exhibiting greater expression across healthy tissues. Evaluation was performed

on the topmost highly expressed 10, 50, 100 miRNAs ranked by the sum of 5p and 3p

strands RPMM values. We also performed the analysis using the complete dataset (all).

Results

A novel HT qPCR approach to study miRNA biogenesis

miRNA identification from somatic tissues is challenging because miRNAs are small,

not abundant, and have a short half-life. We tested several approaches and ultimately

developed a new, more robust HT qPCR-based approach capable of unambiguously

identifying and quantifying miRNA expression and discriminating between 5p or 3p

strand usage from total RNA isolated from somatic tissues. This method is a high

throughput adaptation and general improvement of an existing methodology (Mei et al.

2012), which uses the polyuridine polymerase enzyme to elongate a polyU tail to all

RNA transcripts in a given sample. This step extends the length of each miRNA present

in the sample, which is then used as a template in reverse transcriptase reactions using

a specially engineered universal stem-loop reverse primer (SL-polyA) (**Main Fig. 1A**).

To identify miRNA occurrence, measure their abundance, and pinpoint miRNA

strand usage for each miRNA in the *C. elegans* genome, we adapted this approach to

High-Throughput q-PCR experiments conducted in 96-well plate formats in triplicate. In

this step, each well in the microplate contains forward primers complementary to either

the 5p or 3p strand for mature *C. elegans* miRNAs, along with a universal reverse

primer (**Main Fig. 1A**). We named this updated assay High-Throughput miRNA strand

selection (HITmiSS) assay. HITmiSS enables the rapid and highly precise quantification

of the entire miRNAome. It also accurately determines the abundance of the 5p or 3p

strand for each miRNA present in the sample. (**Main Fig. 1A**).

miRNA expression is widespread throughout *C. elegans* development

We employed our HITmiSS methodology to systematically quantify miRNA

occurrence and the usage of 5p and 3p strands in known miRNAs throughout *C.*

elegans development. To achieve this, we extracted total RNA from *C. elegans* at six

developmental stages (embryos, larval stages 1–4, and young adults) and analyzed the

miRNA transcriptome and strand selection for 190 miRNAs, representing 75% of all

miRNAs annotated in miRbase, using a total of 380 unique forward primers across four

plates (**Supplemental Table S1**).

We have performed 7,980 qPCR experiments independently, querying these 190

miRNAs throughout development in triplicate (1,140 experiments at each stage) (**Main**

Fig. 1B, Supplemental Fig. S3). For each miRNA tested, we also repeated these

experiments without adding the template cDNA to the reaction (negative control). We

subtracted the signal obtained from our negative control from each miRNA tested,

averaging the results for each triplicate experiment (see Materials and Methods). Using

stringent filters, we detected the presence of 177 of these miRNAs (93.2%) in at least

one developmental stage (**Supplemental Table S2**). The greatest number of miRNAs

occur during L4 stage (81%), followed by young adult (75.5%), L3 (65%), L2 (56.8%),

and lastly, Embryo and L1 (41.6% each) stages. This aligns with previous literature

suggesting that many *C. elegans* miRNAs are tissue-specific, only becoming more

abundant upon the development of those tissues later in the developmental cycle

(Brosnan et al. 2021).

Consistent with past observations, most of the miRNAs detected in embryo and

early larval stages play a functional role in embryonic lethality and early development.

Namely, the *miR-35* and *miR-51* families, both important regulators of embryonic

lethality, are the most highly expressed miRNAs during these early stages (Shaw et al.

2010). Similarly, the *let-7* and *lin-4* families, miRNAs that regulate adult phenotypes,

were the most highly detected in L4 and young adult stages (Alvarez-Saavedra and

Horvitz 2010). Importantly, we did not detect significant expression of 13 miRNAs: *miR-*

791, *miR-792*, *miR-1823*, *miR-2208a*, *miR-2209a*, *miR-2213*, *miR-2220*, *miR-2953*,

miR-4808, *miR-4810b*, miR-*8201*, *miR-8203*, or *miR-8206*. Many of these miRNAs have

not been experimentally validated, and they fall below our threshold.

In conclusion, gathering data using the HiTmiSS assay is robust, sensitive,

reproducible, scalable and cost-effective. Although the filters we used in our analysis

were very stringent, HiTmiSS was sensitive enough to detect more than 93% of all

tested miRNAs above random background, including many which remain poorly

characterized despite their relatively high expression.

Many miRNAs show consistent strand preference throughout development

Our method accurately quantifies miRNA abundance and detects which strand (5p

or 3p) is present in each sample. Many miRNAs identified consistently use either the 5p

or 3p strand as the guide strand, which we termed '*Singletons*', while others switch

between 5p and 3p usage across different developmental stages. We named the

second category '*Switchers*'. Importantly, the hits our assay produced align with past

findings on the strand preference of well-characterized miRNAs such as *let-7, lin-4*

(known *5p Singletons*), and the *miR-35* family (known *3p Singletons*) (Alvarez-Saavedra

and Horvitz 2010). The screen produced hits for 177 miRNAs, which we divided into *5p*

Singletons, *3p Singletons*, or *Switchers* (**Main Fig. 1B and Supplemental Table S2**).

The largest category identified is miRNA *Switchers*, with 88 hits (46.3%). These

miRNAs do not consistently prefer one strand across all developmental stages and

instead switch strand usage between the 5p and the 3p strand throughout development.

These included members of the well-characterized *miR-2* families and others, regulators

of developmental timing, and heat shock (Alvarez-Saavedra and Horvitz 2010; Garcia-

Segura et al. 2015). *Singletons* collectively represent the other half of the total hits

obtained, and the strand usage is surprisingly not equally distributed, with *3p Singletons*

accounting for nearly double the number of *5p Singletons* detected and are the second

largest group detected, with 58 hits (30.5%) (**Main Fig. 1B**).

These included members of the *miR-35* and *miR-51* families, both regulators of

embryonic lethality, members of the *miR-44* family, which regulate stress and starvation

responses, as well as members of the *miR-58* family, responsible for locomotion, egg-

laying, dauer entry, and body size (Alvarez-Saavedra and Horvitz 2010; Shaw et al.

2010; Garcia et al. 2014). Interestingly, some of the most highly detected *3p-singletons*

are not well-defined functionally, such as *miR-60*, *miR-70*, and *miR-90*, all of which are

known to broadly affect the stress response, though their specific impacts and targets

remain unclear (Kagias et al. 2014; Garcia-Segura et al. 2015; Kato et al. 2016).

As mentioned, the least prevalent group detected is the *5p Singletons*, with 31 hits

(16.3% of all tested miRNAs) corresponding to several well-characterized miRNA

families (**Main Fig. 1B**). These included the *let-7* family and members of the *lin-4* family,

both of which are known to regulate developmental timing and progression into the later

stages of development (Alvarez-Saavedra and Horvitz 2010). We also detected

members of the *miR-51* family, which regulates embryonic lethality, and the *miR-64*

family, which regulates the heat shock response (Shaw et al. 2010; Kagias et al. 2014).

MiRNAs are often grouped by family containing similar seed regions and broadly

similar targets (Wolter et al. 2017). Some miRNA families use the same strand

throughout development (*let-7*, *miR-64* family). However, the majority of them are

spread between *5p Singletons*, *3p Singletons*, and *Switchers*, suggesting that even

within the same family, with similar structure, use different rules to identify the used

strand. For example, the *miR-51* family, a well-characterized developmental regulator

(Brenner et al. 2012), has its members equally split between *5p Singletons* (*miR-51*,

miR-52, *miR-53*) and *3p Singletons* (*miR-54*, *miR-55*, *miR-56*, *lsy-6*), with no *Switchers*.

To account for the strong strand preference observed among *Singletons*, we took

the top 20 *5p Singletons* and the top 20 *3p Singletons* and compared the binding

signature and nucleotide composition between the used strand versus the strand that

gets discarded (**Main Fig. 1C, i and ii**). Interestingly, the pairing frequency does not

vary between these miRNAs (**Main Fig. 1C,ii**), either in the 5p or 3p strand, with the

exception of the last few nucleotides at the 3'end (**Main Fig 1C, I, red boxes**) which are

commonly unpaired in the mature duplex.

When comparing the used vs. the discarded strand (**Main Fig. 1C, left panel vs**

right panel), we detected no significant difference in the pairing signature at any

position; however, the 5p strand, independently if is used or discarded, contains more

unpaired nucleotides (**Main Fig. 1C, red boxes**). When narrowing our analysis to just

the seed region (**Main Fig. 1C**, golden boxes), while we detect higher pairing than any

other position within pre-miRNAs, we still do not detect any difference in binding

frequency between the used and discarded strands, suggesting that other rules are also

in place.

Next, we compared the nucleotide composition of these seed regions, hoping to

identify signatures that could explain the discrepancy in miRNA strand selection in

Singletons. We determined the likelihood of any given nucleotide from positions 1

through 10 in each miRNA strand in the top 20 *Singletons* in each category (5p and 3p)

(**Main Fig. 1C, ii**). We detected a highly conserved uracil nucleotide at position 1, as

previously shown by us and others (Wolter et al. 2017), though this conservation

dissolves from positions 2 through 10 (**Main Fig. 1C, ii**). Importantly, this uracil

nucleotide is not equally conserved between *5p Singletons* and *3p Singletons.* 80% of

3p Singletons start with this uracil nucleotide, while less than 40% occur in *5p*

Singletons. This differential presence of this uracil may be caused by a non-stochastic

pre-miRNA cleavage by the *C. elegans* DCR-1, which may possess a preference to

cleave at this nucleotide. Unfortunately, the structure of DCR-1, including the cleavage

site and nucleotide-binding pocket, remains unsolved, and more experiments need to

be performed to validate if this cleavage preference occurs.

The intestine miRNA population is small but highly interconnected

In our experiments, we could not consistently detect 40.3% of the tested miRNAs

throughout development. We believe that this is caused by either our stringent filters

that exclude miRNAs with low signal in a specific developmental stage and/or because

these miRNAs are not widely expressed in all stages and may be restricted in specific

tissues. In addition, *Switchers*, which are the most abundant category in our whole

worm screen, may not be 'true' *Switchers*, as they could act as either *3p Singletons* or

5p Singletons in different somatic tissues and/or developmental contexts, skewing our

analysis.

Therefore, to better understand the rules behind miRNA strand selection, we

decided to expand and deconvolve our analysis to somatic tissues throughout

development. For these experiments, we focused on the intestine tissue because it is

large, and its transcriptome has been extensively characterized by us and others

(Haenni et al. 2012; Blazie et al. 2017).

We coupled our HiTmiSS assay with intestine-specific miRNA immunoprecipitation,

as we have published in the past (Kotagama et al. 2019). More specifically, we forced

the expression of the GFP-tagged *C. elegans* Argonaute ortholog *alg-1* in the intestine

tissue using the intestine-specific promoter *ges-1p*, performed an α-GFP

immunoprecipitation, recovered the miRNA strands loaded onto ALG-1, and used them

as templates in our HiTmiSS assay (**Main Fig. 2A**). We performed these experiments in

triplicates, focusing our analysis on the embryo and young adult stages for the most

highly expressed miRNAs from our whole mixed tissue analysis.

We found that of the 96 miRNAs tested, 64 (68%) were present during at least one

stage within the intestine tissue (**Main Fig. 2A-C**). Of these, 58 (91%) were *Singletons*,

while 6 (9%) were *Switchers* (**Supplemental Fig. S4A**).

This strongly differs from our whole worm data, reinforcing that miRNAs previously

labeled as *Switchers* may truly be *Singletons* on a tissue-by-tissue basis, only

appearing as *Switchers* once all tissues are analyzed at once.

Interestingly, the proportion of *5p Singletons* to *3p Singletons* is not the same in the

intestinal tissue as in the mixed tissue (**Supplemental Fig. S4A**). In our initial analysis,

about two-thirds of *Singletons* were *3p Singletons*, yet in the intestine tissue, nearly two-

thirds of *Singletons* are *5p Singletons* (**Supplemental Fig. S4A**). This change implies

additional factors influencing the strand selection decision specific to individual tissues,

though further analyses in more tissues will be needed to confirm these trends.

Another surprise was the presence of intestine-specific *Switchers* (**Main Fig. 2A and**

Supplemental Fig. S4B). This includes the miR-51 family, well-characterized regulators

of embryonic lethality (Shaw et al. 2010). For these miRNAs, usage of both strands may

be crucial to ensure proper development of intestine function during embryogenesis. In

addition, we detected intestine tissue Switchers among the *miR-44, miR-58*, and *miR-71*

families (**Supplemental Fig. S4B**). These miRNA families have diverse functions, from

regulating developmental timing to starvation, hypoxia, stress response, and dauer

entry (Kagias et al. 2014; Pandit et al. 2014; Garcia-Segura et al. 2015; Brunquell et al.

2017). Importantly, these are intestine-specific phenotypes, suggesting that strand

switching in these miRNAs may have functional significance by expanding 3'UTR

targeting repertoires in the intestine.

stage. These strains expressed the promoter for a single miRNA fused to GFP, and live

imaging was used to validate the expression of these miRNAs at the appropriate life

stage in the intestine (**Main Fig. 2C**). All four strains displayed the expected

fluorescence.

miRNA target presence doesn't account for strand usage

We were concerned that the presence of miRNA targets in cells could stabilize a

specific miRNA strand, thus influencing its detection by our HiTmiSS assay. In this

scenario, our approach might not accurately reflect genuine strand selection but rather

detect artifacts resulting from target-induced stabilization. According to this model, the

undetected strand lacks cellular targets, leading to its degradation and subsequent

absence in our qPCR results.

To test this possibility, we overexpressed in the intestine tissue two 3'UTRs

containing several miRNA binding sites for strands that are discarded as for our

HiTmiSS results (*fox-1* 3'UTR and *hrp-2* 3'UTR) (**Supplemental Figure S5**). *Fox-1*

3'UTR contains binding sites for *miR-60 5p* and *hrp-2* contains dinging sites for *miR-232*

5p and *miR-239b 5p*). If this model was correct, we should observe an increase

abundance of this typically discarded strand and a more equal ratio of 5p vs 3p usage.

Our test did not detect any difference in miRNA strand usage for these miRNAs, which

were always very similar to their wild type N2 controls, suggesting that the presence of

a given miRNA target is not per se responsible for miRNA strand selection

(**Supplemental Figure S5**). For example, *miR-60 3p*, which did not have binding sites

to *fox-1* 3'UTR, remained the dominant detected strand in each experiment.

(**Supplemental Figure S5**).

Taken together, these results indicate that our qPCR approach is not simply

detecting stabilized miRNA strands due to the presence of targets. Rather, the relative

strand abundance measured by the HiTmiSS assay likely reflects genuine strand

selection, rather than artifacts introduced by target availability.

***5p Singletons* and *3p Singletons* follow different rules for selecting strand usage**

We next studied the structural features associated with Singletons, which could

explain the strict and consistent strand usage within these miRNAs. When comparing

the used and the discarded strand, the pairing frequency vary significantly within each

Singleton, revealing new conserved patterns (**Main Fig. 3**). Specifically, the strand used

in *Singletons* often has an unpaired nucleotide at position 12, while the discarded strand

does not (**Main Fig. 3A**, yellow boxes). This lack of pairing in position 12 in both *5p*

Singletons and *3p Singletons* usage is intriguing. In humans, certain Argonaute proteins

perform cleavage of target RNAs between positions 10 and 11, which corresponds to

positions 10 and 11 along the bound miRNA strand by Watson-Crick pairing (Nakanishi

2024). *C. elegans alg-1* is largely homologous to human Argonaute proteins, and it may

retain this cleavage activity at similar target positions, leading to the degradation of the

unused strand due to its Watson-Crick base pairing at this crucial position. In addition,

we detected differences in the binding frequency of the first position nucleotide in both

5p Singletons and *3p Singletons* (**Main Fig. 3A, blue boxes**). *3p Singletons* also exhibit

unique pairings at positions 18-19 (**Main Fig. 3**, white box) not present in the discarded

strand. These patterns are most clearly seen in specific *singleton* miRNAs, both *5p*

Singletons like *miR-52* and *3p Singletons* such as *miR-75* (**Main Fig. 3B**).

An updated machine learning model for miRNA strand selection

We next sought to extend our findings by using our wet bench validated data as a

training set for a machine learning model designed to predict 5p versus 3p strand usage

in miRNA hairpins. We wanted to extract patterns and features that make *Singletons* so

unique and allow then to always use one strand over the other. We selected a Random

Forest classifier, trained on experimentally validated *Singletons* identified through our

HITmiSS assay. This model incorporates 77 features, systematically organized into five

functional categories (dinucleotide composition, first/last nucleotide bias, GC content

and entropy, base pairing properties, general structural features) that together capture

critical aspects of miRNA mature duplex structure (**Supplemental Fig. S1–S2,**

Supplemental Table S3-S4, and Materials and Methods). Dinucleotide Composition

features capture the frequency of specific dinucleotide motifs within the 5p and 3p

strands, which can influence local secondary structure and strand selection dynamics.

First/Last Nucleotide Bias features record the identity of the first and last nucleotides of

each strand, reflecting established preferences that impact Argonaute loading and

strand stability. GC Content and Entropy features summarize the overall nucleotide

composition and diversity of each strand, which assess rigidity, flexibility, and relative

stability. Base-pairing Properties include the fraction of paired nucleotides in each arm,

capturing local structural stability and asymmetry between the 5p and 3p strands.

Finally, General Structural Features encompass broader geometrical characteristics,

representing the overall size of the miRNA duplex and differences in strand lengths that

may influence processing.Together, these features allow for accurate modeling of the

structural and sequence-based factors governing strand selection. The model

demonstrated strong performance in distinguishing 5p from 3p usage across a set of

known miRNAs (**Supplemental Fig. S2, and Materials and Methods**).

5p and *3p Singletons* **possess distinct structural features**

To explore the molecular determinants that differentiate strand selection, we

conducted a comparative feature analysis of the mature *C. elegans* miRNAs.

We extracted and calculated all 77 features used in our ML model for the entire C,

elegans miRNAome (374 miRNAs, miRbase) (**Supplemental Table S4**). We kept our

definition of *Singletons* very stringent to increase the signal (confidence score >80%).

We identified 54 *5p Singletons*, 52 *3p Singletons* and 268 *Switchers* (**Main Fig. 4A-B**

and **Supplemental Table S4**). Importantly, many of the *Singletons* identified by the ML

model were previously described as using only one strand (*let-7*, *lin-4*, etc.), reinforcing

our confidence of the ML model.

Out of 77 extracted structural and sequence features, 13 exhibited statistically

significant differences between the 5p and 3p strands ($p < 0.05$, Welch's t-test) (**Main**

Fig. 4B-E). The most pronounced difference was observed in the identity of the first

nucleotide of the 3p strand, which in *3p Singletons* was consistently a uracil ($p = 3.98 \times$

$10^-\square$) (**Main Fig. 4C**). Additionally, 3p strands showed an enrichment for specific

motifs, particularly GAG ($p = 2.51 \times 10^-\square$), and were more likely to terminate

in cytosine ($p = 5.69 \times 10^-\square$) (**Main Fig. 4E**). We also observed compositional

differences: 3p strands displayed increased levels of cytosine at the first position and

tended to be slightly shorter than their 5p counterparts ($p = 8.90 \times 10^{-3}$) (**Main Fig. 4D-**

E).

These findings suggest intrinsic sequence and structural biases associated with

strand identity that may reflect evolutionary pressures or mechanistic preferences in

miRNA biogenesis. Together, these results highlight discrete, strand-specific features

that distinguish *5p* and *3p Singletons*.

Next, we decided to use our ML model and test if mutating these identified features,

we could alter the overall miRNA structure and strand usage. In the case of *let-7*,

altering the nucleotide pairing in position 11 and 12 (UP11-12) and/or altering the

identity of the first nucleotide (5'U→C) significantly disrupted the strand selection and

lowered its RNAfold predicted confidence as a *5p Singleton* up to 17%, supporting our

model (**Main Figure 3C-D**). To further validate these findings, we altered the structure

and composition of four other well-characterized *Singleton*: *miR-52, miR-75, lin-4, and*

miR-35 (**Supplemental Fig. S6**). In every case we saw a similar pattern that introducing

bulges near the center of the mature hairpin disrupted predicted strand usage, while

eliminating bulges in this region reinforced strand usage, further suggesting the

importance of this region in our model of strand selection.

Next, we decided to use our ML model and test whether mutating the features

identified in our *Singleton* data could alter overall miRNA structure and predicted strand

usage. In the case of *let-7*, altering the nucleotide pairing in position 11 and 12 (UP11-

12) and/or altering the identity of the first nucleotide (5'U→C) significantly disrupted the

strand selection and lowered its RNAfold predicted confidence as a *5p Singleton* up to

17%, supporting our model (**Main Figure 3C-D**). To further validate these findings, we

altered the structure and composition of four other well-characterized *Singletons*: *lin-4,*

miR-52, miR-75, and miR-60 (**Supplemental Fig. S6**). In every case we saw a similar

pattern that introducing bulges near the center of the mature hairpin disrupted predicted

strand usage, further reinforcing the importance of this region in our model of strand

selection.

Similarly, the identity and pairing of the first nucleotide remained an important, though

inconsistent indicator of predicted strand usage (**Supplemental Fig. S6**). Here we

witnessed a stark difference in the magnitude of outcomes predicted as a result of these

mutations between different miRNAs. While mutations of the first nucleotide in *5p*

Singletons did not alter predicted strand usage more than 10%, similar mutations in 3p

Singletons were predicted to change strand preference by as much as 30%

(**Supplemental Fig. S6**). This recapitulates the trends in our HiTmiSS assay data and

further reinforces our model of strand selection.

Strand Usage Bias Differs by Species

To investigate structural differences underlying miRNA biogenesis across species,

we compared predicted hairpin features between *C. elegans* and human miRNAs using

our trained model (**Main Fig. 5**) (**Supplemental Table S5**). In *C. elegans*, strand usage

was nearly balanced, with 51.3% of miRNAs predicted to favor the 5p strand and 48.7%

favoring the 3p strand (192 vs. 182 predictions, respectively). This symmetry suggests

that strand choice in *C. elegans* remains broadly permissive, possibly facilitating a more

dynamic or context-specific regulatory output. In contrast, human miRNAs exhibited a

marked asymmetry, with 75.9% of hairpins predicted to preferentially use the 5p strand

(648 vs. 206 predictions) (**Main Fig. 5A**). In addition, statistical comparison revealed 13

features with significant differences ($p < 0.05$, Welch's t-test) between human and *C.*

elegans miRNAs (**Main Fig. 5B**).

Among these, several features stood out with exceptionally strong statistical support.

The most significant divergence was observed in guanine content on the 5p strand ($p =$

5.62×10^{-2}), which was markedly higher in human miRNAs (**Main Fig. 5B**). Similarly,

GG dinucleotide frequency ($p = 1.94 \times 10^{-2}$) was enriched in humans suggesting

evolutionary divergence in sequence composition or processing preferences (**Main Fig.**

5B). Motif analysis further highlighted a strong enrichment of the CCU motif on the 3p

strand in humans ($p = 9.37 \times 10^{-22}$), alongside a general increase in cytosine frequency

in 3p regions (3p_C, $p = 1.56 \times 10^{-2}$). Additionally, di_CC ($p = 4.25 \times 10^{-2}$), another

cytosine-related motif, was significantly more common in human sequences. Importantly

these results are also evident in known miRNAs such as *let-7* (**Main Fig. 4B**). We have

also applied our ML model to predict miRNA strand selection in additional species

including mouse, rat, frog, zebrafish and fly (**Supplemental Table S5**).

Collectively, these findings suggest that human miRNAs exhibit cytosine- and

dinucleotide-enriched motifs**,** especially in 3p strands**.** These compositional signatures

likely reflect differences in strand selection, processing pathways, or functional targeting

between the two species.

Our ML model can accurately predict human miRNA strand selection

To validate the accuracy of our machine learning model in predicting human miRNA

strand selection, we compared model predictions to the comprehensive Human miRNA

Tissue Atlas expression dataset in 65 healthy human organ systems (Rishik et al. 2025)

(**Main Fig. 6**). When comparing model performance across subsets of highly expressed

miRNAs from this dataset, we observed a consistent trend: our model was very

accurate when predicting strand usage among subsets of highly expressed miRNAs

(**Main Fig. 6A**). Among the top 10 most highly expressed miRNAs, our model achieved

an accuracy of 90%, surpassing even its accuracy in *C. elegans*. However, as we

included more miRNAs from the atlas, the accuracy of the model, as expected, trended

downward, reaching a lower yet still statistically significant ($p \approx 7.6 \times 10^{-\square}$) accuracy of

59% when predicting strand usage among all human miRNAs in the dataset (**Main Fig.**

6A).

To refine our analysis and reduce potential noise from low-expression features, we

focused on the top 50 most abundantly expressed miRNA duplexes (**Main Fig. 6B**).

Among these, the model correctly predicted the dominant strand in 33 cases,

corresponding to an overall accuracy of 66% ($p \approx 0.0164$) (**Main Fig. 6B**). Notably, our

model displayed a strong bias toward the 5p strand, with several 3p dominant miRNAs

being misclassified as *5p Singletons* (**Main Fig. 6B**). The 17 incorrect predictions were

generally concentrated among miRNAs in the Atlas that had little difference in

expression between their 5p and 3p strands. This in turn suggests that errors in our ML

model predictions are more likely present when there is not a clear prevailing strand.

Alternatively, since the difference in the expression levels of the 5p and 3p strand is so

small, these miRNAs may be *Switchers* in Humans (**Supplemental Table S6**).

Of note, two of the mispredicted miRNAs, *miR-182* and *miR-378a,* have indeed been

documented as *Switchers in vivo.* (Krishnan et al. 2013; Krist et al. 2015).

On the other hand, this misprediction of strand dominance often aligns with miRNAs

where the confidence scores of the 3p and 5p strand are very close (+/- ~10%).

For example , *miR-199a* and *miR-199b*, both implicated in hypoxia responses and

cardiac regulation (Rane et al. 2009), exhibited dominant expression of the 3p strand in

healthy tissues but were predicted as 5p dominant by our ML model with a very small

confidence of +/- ~7%. *miR-124*, a neuron-specific miRNA essential for neurogenesis,

was similarly misclassified, despite showing strong 3p expression dominance in the

dataset with a confidence of +/- 6% (Makeyev et al. 2007).

Still, the majority of miRNAs were correctly predicted across a broad range of organ

systems, including several critical functional regulators known to exhibit strong strand

bias, suggesting that our ML model trained in *C. elegans* Singletons is capable of

accurately predict strand usage in vertebrates (**Main Fig. 6C and Supplemental Table**

S6). For example, *miR-21*, one of the most abundant miRNAs across multiple organ

systems and a well-established oncomiR (Rhim et al. 2022), was correctly predicted as

a 5p dominant, with a greater than 50-fold difference in expression between its 5p and

3p strands. Similarly, *let-7a*, a central regulator of developmental timing and cell

differentiation (Brueckner et al. 2007), was correctly classified as a 5p dominant with a

248-fold higher expression of the 5p strand relative to the 3p strand.

Other highly expressed miRNAs such as *miR-181a*, involved in T cell development

(Kim et al. 2021), and *miR-100*, implicated in stem cell regulation (Pakravan et al.

2017), also exhibited strong 5p dominance and were accurately predicted by our ML

model.

Overall, these findings demonstrate that the model is capable of reliably predicting

strand usage for miRNAs in healthy human tissues, particularly in cases where strand

bias is strong. Errors were enriched among duplexes with more balanced expression

levels or context-dependent strand usage, highlighting the complexity of predicting

which strand will be used in cases of weak or dynamic regulation of strand selection.

827 **Discussion**

In this study, we present a comprehensive exploration of miRNA strand selection

using a combination of high-throughput quantification, structural modeling, and

predictive machine learning. Our goal was to study the molecular features that govern

whether the 5p or 3p strand of a miRNA duplex is preferentially selected during

biogenesis.

Using an in-house developed HiTmiSS assay, we performed the largest qPCR

screen to date, performing ~8k qPCR experiments profiling miRNA strand usage

throughout *C. elegans* development, and in the intestine tissue.

Our analysis led to the identification of experimentally validated *5p Singletons*, *3p*

Singletons and *Switchers* throughout *C. elegans* developmental stages (**Main Fig. 1**).

We then used structural, thermodynamic and intrinsic properties from these

experimentally validated data to develop a ML model capable of predicting high quality

miRNA strand usage from a given pre-miRNA sequence and its corresponding hairpin

structure. Furthermore, we extended our findings to vertebrate systems, comparing our

ML model results to the Human miRNA Tissue Atlas.

We observed in our mixed tissue analysis in **Main Fig. 1** that a substantial portion of

miRNAs act as *Switchers*, shifting strand usage across time, suggesting that strand

selection is not static but can vary by context. However, many well-characterized

developmental regulators such as *let-7*, *lin-4* and *miR-35* are *Singletons*, highlighting

the need of strand specificity in development where steady-state gene expression

output is critical. In this context it is worth it to mention that *Singletons* in general are

more abundant than *Switchers*, with even the most highly expressed *Switchers* detected

at far lower levels than many *Singletons* (**Main Fig. 1B**). This is perhaps due to less

stringent structural and thermodynamic control in these miRNAs.

Strand usage in the intestine, examined using tissue-specific ALG-1

immunoprecipitation (**Main Fig. 2**), revealed striking differences from whole-organism

data. Here, *5p Singletons* outnumbered *3p Singletons*, inverting the developmental

trend in our mixed tissue data. This suggests tissue-specific regulation of strand

selection and supports the idea that miRNA strand choice is influenced not only by

intrinsic structural features but also by the cellular environment and possibly tissue-

specific protein cofactors.

Importantly, since we only tested one somatic tissue, we do not know whether any

Switchers identified from our mix stage analysis are indeed tissue-specific *Singletons,*

displaying opposing strand preference in different tissues and therefore appearing like

Switchers when quantified in mixed tissue. More experiments in different somatic

tissues need to be performed to address these questions.

One potential concern is that target presence may preferentially stabilize one strand

of a miRNA duplex, thereby influencing its detection in our HiTmiSS assay. In this

scenario, both 5p and 3p strands might initially be available for loading, but the

presence of a target could enhance the stability or retention of one strand, leading to an

apparent, but potentially artifactual strand preference.

To test this, we overexpressed specific 3′UTRs known to contain binding sites for

microRNA strands not typically abundant in the intestine (**Supplemental Figure S5**).

Importantly, this manipulation did not alter the abundance of the corresponding strands

detected by our HiTmiSS assay, indicating that our assay faithfully captures genuine

strand usage rather than target-driven stabilization effects (**Supplemental Figure S5**).

When comparing the structural composition of each mature duplex, a series of

patterns emerged. *5p Singletons* tend to have the first nucleotide unpaired to its

corresponding strand in the duplex. This difference in pairing repeats in the middle of

the duplex and drastically reverses toward the end (**Main Fig. 3A**). Contrarily, *3p*

Singletons present a near-opposite trend, with the first nucleotide more often paired and

the end portion of the sequence typically unpaired, though the center of the sequence

shows similar unpairing to *5p Singletons*. Perhaps this highlights structural requirements

that form the mechanistic basis to select for one strand over the other.

To address this question, we extracted 77 structural and compositional features from

our validated *Singletons* and trained a Random Forest classifier (ML Model) to predict

dominant strand usage. When comparing these features (**Main Fig 4**), we found several

strand-specific differences, including a significantly higher occurrence of uracil at the

first nucleotide in 3p strands and enrichment for GAG motifs.

We validated our predictions using *in silico* site-directed structural mutations in

canonical miRNAs such as *let-7*, *lin-4*, and *miR-52*, *miR-75* (**Main Fig. 3C**,

Supplemental Fig. S4). Disrupting features such as central bulges or first nucleotide

pairing, altered predicted strand preference to some degree, further confirming their role

in the selection process. Surprisingly, the degree to which they impact miRNA strand

preference was approximately equal, but when disrupted in tandem led to an even

greater loss of predicted strand preference (**Main Fig. 3C**, **Supplemental Fig. S6**). In

addition, these findings phenocopy the trends seen in our experimentally validated

Singletons (**Main Fig. 3A**). This suggest, that the structural and thermodynamic features

identified in our wet bench validated *5p* and *3p Singletons* are indeed critical regulator

of strand selection in nematodes.

Expanding this framework, we then applied our ML model to predict strand usage

across the human miRNA dataset (**Main Fig. 5A**). We were particularly curious to test if

the structural and thermodynamics features and motifs identified in *C. elegans* were

conserved in vertebrates.

Compared to the nearly equal distribution of 5p and 3p usage in *C. elegans*, human

miRNAs showed a clear bias toward 5p strands, with over 75% of predictions falling in

this category. This divergence likely reflects increased specialization in miRNA

maturation and Argonaute loading in vertebrates, potentially driven by evolutionary

selection for strand-specific targeting or stability. In this view, the reason for the 5p bias

observed among human miRNAs may be due to the large expansion of miRNA families

and miRNA hotspots seen across evolutionary times (Wolter et al. 2017), reinforcing

strict rules which favor *Singletons* over *Switchers*.

We also found significant enrichment of guanine content in human 5p strands and

cytosine-rich motifs in 3p strands (**Main Fig. 5B**). These compositional biases are

consistent with known differences in Argonaute protein specificity and may reflect

lineage-specific constraints on strand functionality (Kotagama et al. 2024).

Importantly, these cross-species trends persisted in canonical miRNAs such as *let-7*,

which was consistently predicted and validated as a *5p Singletons* across both systems,

although at different confidence levels. This supports the robustness of our predictive

ML model and highlights conserved features of miRNA processing, even in the

presence of divergent strand preferences.

When we compare our ML model prediction in humans miRNA to the human miRNA

Tissue Atlas, a public repository of known miRNA expression in their tissue of origin, an

surprising trend emerges. Our ML model developed in *C. elegans* is capable of

accurately predicting miRNA strand usage in humans (**Main Fig. 6**). This is unexpected,

considering the vast evolutionary gap (~600 million years) between these two systems,

which share less than 35% of nucleotide conservation in genes (Stein et al. 2001).

Despite this, the features we have identified seems to be evolutionary conserved.

This suggests that miRNA strand selection is a fundamental cellular process which has

evolved very little over time, even as miRNA regulatory networks expanded in

complexity with the increased cellular specialization. This becomes more evident when

we overlayed miRNA strand usage in human tissues of different origins. Here, the same

miRNA always tends to use the same strand in different somatic tissues.

In conclusion, our findings illustrate the complexity of miRNA strand selection and

reveal that this process is not governed solely by thermodynamic asymmetry. Instead, a

combination of structural motifs, sequence biases, and cellular context dictate strand

usage. Our work establishes a predictive framework for understanding strand selection

but also opens new avenues for studying the evolution of miRNA regulatory logic across

metazoans.

939 Data Availability

940 The raw data produced by the HiTmiSS assay are included in the **Supplemental Table**

941 **S2**. The ML model is included as Supplemental Data and is also available upon request.

943 Funding

944 This work was funded by the National Institutes of Health 5R01GM118796 awarded to

945 M.M.

947 Acknowledgments

948 We thank Dr. Murugan for assistance in HiTmiSS assay optimization.

950 Author Contributions

951 D.M. and M.M. designed the experiments. D.M. performed the experiments. D.M. and

952 M.M. prepared the ML model and performed the bioinformatics analysis. D.M., A.E. and

953 H.H. performed the HiTmiSS assay experiments. D.M. and M.M. prepared the main and

954 supplemental figures and supplemental tables. D.M. and M.M. wrote the manuscript. All

955 authors have read and approved the manuscript.

Declaration of Interests

The authors have no conflicts of interest.

Figure titles and legends

Main Figure 1: HiTmiSS Assay and Analysis on Mixed Tissue. **(A)** HiTmiSS protocol

for RNA extraction, extension, reverse transcription, and quantification in high

throughput. Steps include (1) *C. elegans* growth and homogenization, (2) RNA

extraction and isolation, (3) RNA polyuridinylation, extension, and reverse transcription

(details shown in ii), (4) addition of miRNA-strand-specific forward primers, (5) loading

into 384-well plates, and (6) quantification. **(B)** Heatmap of miRNA expression (n=3) in

mixed tissue clustered by quantity, strand use, and developmental stage. Stages

include embryo (E), larval 1 through 4 (L1-L4) and young adult (YA). The heatmap

legend represents $\log_{10}$ of expression values for each miRNA strand normalized by

spike-in GFP control at each stage from -4 (deep purple) to 4 (bright yellow). Black color

indicates instances of data which fall below strict expression threshold (one standard

deviation above mean of all expression values at that stage). **(C)** Heatmap of pairing

frequency by position for the used (left) and discarded (right) strand of the 20 most

highly quantified *5p* and *3p Singletons* in our mixed tissue dataset. Average pairing

frequency and logo plot of nucleotide frequency at each position displayed below.

Main Figure 2: Analysis of Intestine Tissue-Specific miRNAs. **(A)** Protein

immunoprecipitation of GFP-tagged ALG-1 under the control of the intestine-specific

promoter *ges-1p* for subsequent RNA isolation and quantification using the HiTmiSS

assay. **(B)** Heatmap of miRNA expression in intestine tissue clustered by strand and

developmental stage. The heatmap legend represents $\log_{10}$ of expression values for

each miRNA strand normalized by spike-in GFP control at each stage from -4 (deep

purple) to 4 (bright yellow). Black color indicates instances of data which fall below strict

expression threshold (one standard deviation above mean of all expression values at

that stage). **(C)** Representative brightfield (left) and GFP (right) images of *C. elegans*

strains expressing GFP under the control of promoters for miRNA known to be highly

expressed in the intestine, either during embryo stage (top two) or young adult stage

(bottom two).

Main Figure 3: Structural and Thermodynamic Determinants of Strand Selection.

(A) Pairing frequency by position of the used (green) and discarded (red) strand for the

most highly quantified *5p Singletons* and *3p Singletons* in mixed tissue (n=20). Positions

at which pairing frequency most differs are boxed, including the first nucleotide (light

blue), center of the duplex (gold), and end of the strand (white). **(B)** Representative pri-

miRNA hairpins for a *5p Singleton* (*miR-52*) and a *3p Singleton* (*miR-75*) demonstrating

occurrence of trends in pairing as described above. Mature duplex regions highlighted

in pink. **(C)** Changes in *Singleton* predictive confidence for the miRNA *let-7* as a result

of *in silico* site-directed mutations. RNAfold secondary structures of pri-miRNA with

mature 5p strand (blue) and mature 3p strand (red) from left to right: wildtype (WT),

unpaired nucleotides at positions 11 and 12 via GG→UU substitution (UP11-12), first

nucleotide U→C substitution (1U → C), combination of both prior mutations

(Combined). Table shown displays probability that each mutation will yield a *5p*

Singleton (5p) or *3p Singleton* (3p). **(D)** Bar chart displaying probability as described

above.

1006

Main Figure 4: Analysis of Mature miRNA Duplex Features in *C. elegans*. (A)

Frequency distribution of predicted *5p Singleton* confidence across all miRNAs in *C.*

elegans. 3p Singletons (yellow) are defined as miRNAs falling below a predicted 5p

confidence of 20% or below; *Switchers* are defined as miRNAs with a predicted 5p

confidence between 20% and 80%; *5p Singletons* (red) are defined as miRNAs with a

predicted 5p confidence of 80% or greater. **(B)** Predicted categorization of all *C.*

elegans miRNAs by our model as per the above threshold. **(C)** Predicted categorization

of *C. elegans* miRNAs for which uracil is the first nucleotide of the 3p strand. **(D)** Length

difference (5p strand - 3p strand) for each category of *C. elegans* miRNAs. *5p*

Singletons display significantly (p<0.05) longer 5p strands than 3p strands when

compared either to *3p Singletons* or to *Switchers*. **(E)** Heatmap of 32 features including

GC content, miRNA length, hydrogen bonding, nucleotide pairing, strand stability, and

MFOLD-derived parameters enrichment in *5p* and *3p Singletons*. Heatmap values

represent mean values of each feature across the miRNA categories. For binary

features (i.e. 5p_has_AAU, which is either 0 or 1) the number represents the proportion

(e.g. 0.3 = 30% of miRNAs in that category have an AAU motif in the 5p strand). For

continuous features (i.e. gc_content, pairing_frac_5p), the number is the average value

across all miRNAs in that category. For nucleotide composition features (i.e. 5p_A,

3p_U), the number reflects the fraction of that nucleotide within the strand. Features

clustered by average linkage.

1027

Main Figure 5: Comparative Analysis of Mature miRNA Duplex Features between

***C. elegans* and *H. sapiens*. (A)** Frequency distribution of predicted *5p Singleton*

confidence across all miRNAs in *H. sapiens*. *3p Singletons* (yellow) are defined as

miRNAs falling below a predicted 5p confidence of 20% or below; *Switchers* are defined

as miRNAs with a predicted 5p confidence between 20% and 80%; *5p Singletons* (red)

are defined as miRNAs with a predicted 5p confidence of 80% or greater. **(B)** Four most

significant (p<0.05) structural differences between *H. sapiens* (left) and *C. elegans*

(right) mature miRNA duplexes. From top left to bottom right: mean occurrence of

guanine in the 5p strand (5p_G), mean occurrence of guanine-guanine dinucleotide in

either mature strand (di_GG), mean occurrence of cytosine-cytosine dinucleotide in

either mature strand (di_CC), binary classifier for presence of cytosine-cytosine-uracil

trinucleotide in the 3p strand (3p_has_CCU). **(C)** Alignment of *C. elegans* (*cel*) *let-7* and

five *H. sapiens* (*hsa*) *let-7* homologs to demonstrate differences in CCU occurrence

within the 3p strand (red boxes). **(D)** Heatmap of differences in mean value for each of

the 77 model features between each category of miRNA in *C. elegans* (c) and *H.*

sapiens (h). For binary features (i.e. 3p_has_CCU, which is either 0 or 1) the number

represents the proportion (e.g. 0.3 = 30% of miRNAs in that category have an CCU

motif in the 3p strand). For continuous features (i.e. gc_content, au_content), the

number is the average value across all miRNAs in that category. For nucleotide

composition features (i.e. 5p_A, 5p_U), the number reflects the fraction of that

nucleotide within the strand. Features clustered by average linkage.

1049

Main Figure 6: **Model Accuracy for the Human miRNA Tissue Atlas Dataset**. **(A)**

Model accuracy expressed as a percentage of correct predictions for four different

subsets of *H. sapiens* miRNAs (total 854 mature duplexes). From left to right, ranked by

total expression of mature 5p and 3p strand across all tissue types: predicted accuracy

among top 10 (9 correct, 1 incorrect), top 50 (33 correct, 17 incorrect), top 100 (66

correct, 34 incorrect), and all (502 correct, 352 incorrect) miRNAs in this dataset. **(B)**

Confusion matrix of the strand usage prediction model among the top 50 most highly

expressed *H. sapiens* miRNAs. Summarizes differences in 5p and 3p strand usage

between experimentally validated (miRNA Atlas) and model-derived (ML Model) data.

(C) Heatmap of prediction accuracy, either correct (red) or incorrect (blue) for the top 50

most highly expressed *H. sapiens* miRNAs across 65 *H. sapiens* tissue types and

organs. Data are clustered using average linkage.

1062

1063   **Supplemental Information titles and legends**

1064

Supplemental Table S1: **Primers used in this study.**

Supplemental Table S2: **HiTmiSS assay results.**

Supplemental Table S3: **Features used in our ML model and their weights.**

Supplemental Table S4: **ML prediction in *C. elegans*.**

Supplemental Table S5: **ML prediction in vertebrates.**

Supplemental Table S6: **Comparison of ML model with miRNA tissue Atlas.**

1071

1072   **References**

1073

Alvarez-Saavedra E, Horvitz HR. 2010. Many families of C. elegans microRNAs are not

essential for development or viability. *Curr Biol* **20**: 367-373.

Bartel DP. 2018. Metazoan MicroRNAs. *Cell* **173**: 20-51.

Blazie SM, Babb C, Wilky H, Rawls A, Park JG, Mangone M. 2015. Comparative RNA-Seq

analysis reveals pervasive tissue-specific alternative polyadenylation in Caenorhabditis

elegans intestine and muscles. *BMC Biol* **13**: 4.

Blazie SM, Geissel HC, Wilky H, Joshi R, Newbern J, Mangone M. 2017. Alternative

Polyadenylation Directs Tissue-Specific miRNA Targeting in Caenorhabditis elegans

Somatic Tissues. *Genetics* **206**: 757-774.

Brenner JL, Kemp BJ, Abbott AL. 2012. The mir-51 family of microRNAs functions in diverse

regulatory pathways in Caenorhabditis elegans. *PloS one* **7**: e37185.

Brosnan CA, Palmer AJ, Zuryn S. 2021. Cell-type-specific profiling of loaded miRNAs from

Caenorhabditis elegans reveals spatial and temporal flexibility in Argonaute loading. *Nat*

Commun **12**: 2194.

Brueckner B, Stresemann C, Kuner R, Mund C, Musch T, Meister M, Sultmann H, Lyko F.

2007. The human let-7a-3 locus contains an epigenetically regulated microRNA gene

with oncogenic function. *Cancer Res* **67**: 1419-1423.

Brunquell J, Snyder A, Cheng F, Westerheide SD. 2017. HSF-1 is a regulator of miRNA

expression in Caenorhabditis elegans. *PloS one* **12**: e0183445.

Carthew RW, Agbu P, Giri R. 2017. MicroRNA function in Drosophila melanogaster. *Semin*

Cell Dev Biol **65**: 29-37.

Chalfie M, Horvitz HR, Sulston JE. 1981. Mutations that lead to reiterations in the cell lineages

of C. elegans. *Cell* **24**: 59-69.

Frank F, Sonenberg N, Nagar B. 2010. Structural basis for 5'-nucleotide base-specific

recognition of guide RNA by human AGO2. *Nature* **465**: 818-822.

Frokjaer-Jensen C, Davis MW, Hopkins CE, Newman BJ, Thummel JM, Olesen SP, Grunnet M,

Jorgensen EM. 2008. Single-copy insertion of transgenes in Caenorhabditis elegans. *Nat*

Genet **40**: 1375-1383.

Garcia SM, Tabach Y, Lourenço GF, Armakola M, Ruvkun G. 2014. Identification of genes in

toxicity pathways of trinucleotide-repeat RNA in C. elegans. *Nature structural &*

molecular biology **21**: 712-720.

Garcia-Segura L, Abreu-Goodger C, Hernandez-Mendoza A, Dimitrova Dinkova TD, Padilla-

Noriega L, Perez-Andrade ME, Miranda-Rios J. 2015. High-Throughput Profiling of

Caenorhabditis elegans Starvation-Responsive microRNAs. *PloS one* **10**: e0142262.

Gerstein MB Lu ZJ Van Nostrand EL Cheng C Arshinoff BI Liu T Yip KY Robilotto R

Rechtsteiner A Ikegami K et al. 2010. Integrative analysis of the Caenorhabditis elegans

genome by the modENCODE project. *Science* **330**: 1775-1787.

Grishok A. 2005. RNAi mechanisms in Caenorhabditis elegans. *FEBS Lett* **579**: 5932-5939.

Haenni S, Ji Z, Hoque M, Rust N, Sharpe H, Eberhard R, Browne C, Hengartner MO, Mellor J,

Tian B et al. 2012. Analysis of C. elegans intestinal gene expression and polyadenylation

by fluorescence-activated nuclei sorting and 3'-end-seq. *Nucleic Acids Res* **40**: 6304-

6318.

Ilieva M, Panella R, Uchida S. 2022. MicroRNAs in Cancer and Cardiovascular Disease. *Cells*

11.

Isakova A, Fehlmann T, Keller A, Quake SR. 2020. A mouse tissue atlas of small noncoding

RNA. *Proc Natl Acad Sci U S A* **117**: 25634-25645.

Jan CH, Friedman RC, Ruby JG, Bartel DP. 2011. Formation, regulation and evolution of

Caenorhabditis elegans 3'UTRs. *Nature* **469**: 97-101.

Jima DD, Zhang J, Jacobs C, Richards KL, Dunphy CH, Choi WW, Au WY, Srivastava G,

Czader MB, Rizzieri DA et al. 2010. Deep sequencing of the small RNA transcriptome of

normal and malignant human B cells identifies hundreds of novel microRNAs. *Blood*

116: e118-127.

Kagias K, Podolska A, Pocock R. 2014. Reliable reference miRNAs for quantitative gene

expression analysis of stress responses in Caenorhabditis elegans. *BMC Genomics* **15**:

222.

Kato M, Kashem MA, Cheng C. 2016. An intestinal microRNA modulates the homeostatic

adaptation to chronic oxidative stress in C. elegans. *Aging (Albany NY)* **8**: 1979-2005.

Kim C, Ye Z, Weyand CM, Goronzy JJ. 2021. miR-181a-regulated pathways in T-cell

differentiation and aging. *Immun Ageing* **18**: 28.

Kotagama K, Grimme AL, Braviner L, Yang B, Sakhawala RM, Yu G, Benner LK, Joshua-Tor

L, McJunkin K. 2024. Catalytic residues of microRNA Argonautes play a modest role in

microRNA star strand destabilization in C. elegans. *Nucleic Acids Res* **52**: 4985-5001.

Kotagama K, Schorr AL, Steber HS, Mangone M. 2019. ALG-1 Influences Accurate mRNA

Splicing Patterns in the Caenorhabditis elegans Intestine and Body Muscle Tissues by

Modulating Splicing Factor Activities. *Genetics* **212**: 931-951.

Krishnan K, Steptoe AL, Martin HC, Wani S, Nones K, Waddell N, Mariasegaram M, Simpson

PT, Lakhani SR, Gabrielli B et al. 2013. MicroRNA-182-5p targets a network of genes

involved in DNA repair. *RNA* **19**: 230-242.

Krist B, Florczyk U, Pietraszek-Gremplewicz K, Jozkowicz A, Dulak J. 2015. The Role of miR-

378a in Metabolism, Angiogenesis, and Muscle Biology. *Int J Endocrinol* **2015**: 281756.

Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using real-time

quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* **25**: 402-408.

Makeyev EV, Zhang J, Carrasco MA, Maniatis T. 2007. The MicroRNA miR-124 promotes

neuronal differentiation by triggering brain-specific alternative pre-mRNA splicing. *Mol*

Cell **27**: 435-448.

Mangone M, Macmenamin P, Zegar C, Piano F, Gunsalus KC. 2008. UTRome.org: a platform

for 3'UTR biology in C. elegans. *Nucleic Acids Res* **36**: D57-62.

Mangone M, Manoharan AP, Thierry-Mieg D, Thierry-Mieg J, Han T, Mackowiak SD, Mis E,

Zegar C, Gutwein MR, Khivansara V et al. 2010. The landscape of C. elegans 3'UTRs.

Science **329**: 432-435.

Medley JC, Panzade G, Zinovyeva AY. 2021. microRNA strand selection: Unwinding the rules.

Wiley Interdiscip Rev RNA **12**: e1627.

Mei Q, Li X, Meng Y, Wu Z, Guo M, Zhao Y, Fu X, Han W. 2012. A facile and specific assay

for quantifying microRNA by an optimized RT-qPCR approach. *PloS one* **7**: e46890.

Meijer HA, Smith EM, Bushell M. 2014. Regulation of miRNA strand selection: follow the

leader? *Biochem Soc Trans* **42**: 1135-1140.

Melendez A, Levined B. August 24, 2009. Autophagy in C. elegans. In *WormBook*, doi:1551-

8507 (ed. TCeR Community). WormBook.

Murari E, Meadows D, Cuda N, Mangone M. 2024. A comprehensive analysis of 3'UTRs in

Caenorhabditis elegans. *Nucleic Acids Res* doi:10.1093/nar/gkae543.

Nakanishi K. 2016. Anatomy of RISC: how do small RNAs and chaperones activate Argonaute

proteins? *Wiley Interdiscip Rev RNA* **7**: 637-660.

Nakanishi K. 2024. When Argonaute takes out the ribonuclease sword. *J Biol Chem* **300**:

105499.

Pakravan K, Babashah S, Sadeghizadeh M, Mowla SJ, Mossahebi-Mohammadi M, Ataei F,

Dana N, Javan M. 2017. MicroRNA-100 shuttled by mesenchymal stem cell-derived

exosomes suppresses in vitro angiogenesis through modulating the mTOR/HIF-

1alpha/VEGF signaling axis in breast cancer cells. *Cell Oncol (Dordr)* **40**: 457-470.

Pandit A, Jain V, Kumar N, Mukhopadhyay A. 2014. PHA-4/FOXA-regulated microRNA feed

forward loops during Caenorhabditis elegans dietary restriction. *Aging (Albany NY)* **6**:

835-855.

Pare JM, Tahbaz N, Lopez-Orozco J, LaPointe P, Lasko P, Hobman TC. 2009. Hsp90 regulates

the function of argonaute 2 and its recruitment to stress granules and P-bodies. *Mol Biol*

Cell **20**: 3273-3284.

Ramani AK, Calarco JA, Pan Q, Mavandadi S, Wang Y, Nelson AC, Lee LJ, Morris Q,

Blencowe BJ, Zhen M et al. 2011. Genome-wide analysis of alternative splicing in

Caenorhabditis elegans. *Genome Res* **21**: 342-348.

Rane S, He M, Sayed D, Vashistha H, Malhotra A, Sadoshima J, Vatner DE, Vatner SF,

Abdellatif M. 2009. Downregulation of miR-199a derepresses hypoxia-inducible factor-

1alpha and Sirtuin 1 and recapitulates hypoxia preconditioning in cardiac myocytes. *Circ*

Res **104**: 879-886.

Rhim J, Baek W, Seo Y, Kim JH. 2022. From Molecular Mechanisms to Therapeutics:

Understanding MicroRNA-21 in Cancer. *Cells* **11**.

Rishik S, Hirsch P, Grandke F, Fehlmann T, Keller A. 2025. miRNATissueAtlas 2025: an update

to the uniformly processed and annotated human and mouse non-coding RNA tissue

atlas. *Nucleic Acids Res* **53**: D129-D137.

Schorr AL, Mejia AF, Miranda MY, Mangone M. 2023. An updated C. elegans nuclear body

muscle transcriptome for studies in muscle formation and function. *Skelet Muscle* **13**: 4.

Shaw WR, Armisen J, Lehrbach NJ, Miska EA. 2010. The conserved miR-51 microRNA family

is redundantly required for embryonic development and pharynx attachment in

Caenorhabditis elegans. *Genetics* **185**: 897-905.

Smith A, Calley J, Mathur S, Qian HR, Wu H, Farmen M, Caiment F, Bushel PR, Li J, Fisher C

et al. 2016. The Rat microRNA body atlas; Evaluation of the microRNA content of rat

organs through deep sequencing and characterization of pancreas enriched miRNAs as

biomarkers of pancreatic toxicity in the rat and dog. *BMC Genomics* **17**: 694.

Steber HS, Gallante C, O'Brien S, Chiu PL, Mangone M. 2019. The C. elegans 3'-UTRome v2

resource for studying mRNA cleavage and polyadenylation, 3'-UTR biology, and

miRNA targeting. *Genome Research*.

Stein L, Mangone M, Schwarz E, Durbin R, Thierry-Mieg J, Spieth J, Sternberg P. 2001.

WormBase: network access to the genome and biology of Caenorhabditis elegans (vol

29, pg 82, 2001). *Nucleic Acids Research* **29**: 1012.

Sulston JE. 1983. Neuronal cell lineages in the nematode Caenorhabditis elegans. *Cold Spring*

Harb Symp Quant Biol **48 Pt 2**: 443-452.

Wolter JM, Le HH, Linse A, Godlove VA, Nguyen TD, Kotagama K, Lynch A, Rawls A,

Mangone M. 2017. Evolutionary patterns of metazoan microRNAs reveal targeting

principles in the let-7 and miR-10 families. *Genome Res* **27**: 53-63.

Zhang Q, Hrach H, Mangone M, Reiner DJ. 2022. Identifying the Caenorhabditis elegans vulval

transcriptome. *G3 (Bethesda)* **12**.

Zinovyeva AY, Veksler-Lublinsky I, Vashisht AA, Wohlschlegel JA, Ambros VR. 2015.

Caenorhabditis elegans ALG-1 antimorphic mutations uncover functions for Argonaute

in microRNA guide strand selection and passenger strand disposal. *Proc Natl Acad Sci U*

S A **112**: E5271-5280.

1216

Main Figure 1

A

B

miRNA Expression [log($2^{-\Delta\Delta Ct}$)]

5p strand   3p strand

E L1 L2 L3 L4 YA   E L1 L2 L3 L4 YA

5p Singletons
31 (16.3%)

3p Singletons
58 (30.5%)

Switchers
88 (46.3%)

C

5p Singletons

i)

5p strand (used)   3p strand (discarded)

position   position

ii)

pairing frequency by position   pairing frequency by position

3p Singletons

i)

3p strand (used)   5p strand (discarded)

position   position

ii)

pairing frequency by position   pairing frequency by position

Main Figure 2

A

5p Singletons

3p Singletons

C

Variant	5p	3p
WT	0.85	0.15
UP11-12	0.71	0.29
5'U → C	0.71	0.29
Combined	0.68	0.32

B

miR-52

miR-75

D

Main Figure 5

A Model Accuracy by Expression Rank

B Confusion Matrix Top 50 – miRNA Atlas vs ML Model

C Model Accuracy – Top 50 miRNAs across 65 *H. sapiens* Organ Systems

correct prediction
incorrect prediction

Additional File 1 for:

Exploring MicroRNA Strand Selection: Insights into Gene Regulation and Development

Dalton Meadows[1,2], Halee Hargis[1,2], Amanda Ellis[1,2] and Marco Mangone[2*]

[1] The Biodesign Institute at Arizona State University 1001 S McAllister Ave, Tempe, AZ

[2] School of Life Sciences, Arizona State University, 427 E Tyler Mall, Tempe, AZ

* To whom correspondence should be addressed. Tel: (480) 965-7957; Email: mangone@asu.edu

This PDF includes:

Supplemental Figs. S1-S6

Table of Contents

Additional File 1

A First/Last Nucleotide Bias
- 3p_first_nt_U
- 3p_first_nt_C
- 3p_first_nt_G
- etc.

GC Content and Entropy
- gc_3p
- gc_5p
- GC_ratio_5p3p
- Shannon_entropy_5p
- etc.

27.5% 24.5%

4.6%

4.3%

Base-pairing Properties
- pairing_frac_5p
- pairing_frac_3p
- etc.

Dinucleotide composition
- di_UU
- di_UG
- di_CA
- di_GA
- etc.

39.1%

General Structural Features
- length
- length_diff
- etc.

B

Input

Hairpin sequence

uacacuguggauccg
gUGAGGUAGUAGGUU
GUAUAGUUuggaaua
uuaccaccggugaaC
UAUGCAAUUUUCUA
CCUUACCggagacag
aacucuucga

Secondary structure

.....(.((..((((
(((((((((.(((((
(((((((.......
(((((...))))))))
))))))))))).))))
))))))))).).))..)
....

Features

77 features

- **Dinucleotide Composition** (frequency of specific two-nucleotide motifs within the 5p and 3p regions, etc.)
- **First/Last Nucleotide Bias** (identity of the first and last nucleotides in each strand, etc.)
- **GC Content and Entropy** (global nucleotide composition and thermodynamic diversity within each strand, etc.)
- **Base-pairing** (nucleotides engaged in base pairing within the 5p and 3p regions, etc.)
- **General Structural Features** (overall miRNA length and the length difference between the 5p and 3p strands, etc.)

Model
Random Forest Classifier

Output

Predicted strand: 5p Confidence score 5p: 0.85

Supplemental Figure S1: Overview of the miRNA Strand Usage Prediction Model. (A) Weight assigned to each feature of the model. **(B)** Flowchart illustrating the computational pipeline developed to predict 5p or 3p strand usage in miRNA precursors using machine learning. Sample *let-7* input and output shown.

A

B

C

Supplemental Figure S2: Overview of Model Performance. **(A)** Confusion matrix of the strand usage prediction model. Summarizes the classification performance of the Random Forest model trained on *C. elegans* miRNA data, showing the number of correctly (33 and 50) and incorrectly (0 and 11) predicted 5p and 3p strands. **(B)** Precision-recall curve illustrating the relationship between precision (positive predictive value) and recall (sensitivity) for classifying 5p versus 3p strand usage in miRNA precursors using the Random Forest model. Each point on the curve represents a different decision threshold. The shape of the curve reflects the model's ability to maintain high precision while recovering true positives across a range of thresholds. **(C)** Receiver Operating Characteristic (ROC) curve of the strand usage prediction model. The ROC curve shows the trade-off between true positive rate (sensitivity) and false positive rate (specificity) for predicting 5p versus 3p strand usage using a Random Forest model trained on *C. elegans* miRNA data. Each point represents a different classification threshold. The area under the curve (AUC) provides a summary measure of the model's overall performance, with higher values indicating stronger discrimination between 5p and 3p strands. The curve demonstrates reliable strand separation across a range of thresholds.

5p Singletons

3p Singletons

Switchers

Supplemental Figure S3: Strand Usage per Category in Representative miRNAs. Normalized mean expression values (n=3) for the 5p strand (red) and 3p strand (blue) of several representative miRNAs that fall into one of our categories, either *5p Singleton*, *3p Singleton*, or *Switcher*. Expression is demonstrated to fluctuate across developmental time for each of these miRNAs in mixed tissue, following their expression from embryo stage (E) through the four larval stages (L1-L4) and in young adult stage (YA). Within each *Singleton* category we find well characterized canonical regulators of development such as *let-7*, *lin-4*, *miR-35* and *miR-53*. Within the *Switcher* category the trends are more diverse, with some miRNAs like *miR-40* altering which strand is used after embryo stage, while others like *miR-48* display strand switching later in development. Still others like *miR-354* show relatively equal strand usage throughout all developmental stages. Perhaps this diversity in strand usage trend among *Switchers* in our whole organism data can be explained by tissue-specific regulation, with one strand preferred in some tissues and the opposite strand preferred in other tissues.

A miRNA Distribution by Type

Mixed Tissue

Intestine Tissue

■ 5p Singleton ■ 3p Singleton ■ Switcher

B

Family	Family Members	Function in the Intestine
miR-44	miR-44	heterochronic timing, development
miR-51	miR-54, miR-56, miR-65	polarity, hypoxia
miR-58	miR-58a	stress response, dauer entry
miR-71	miR-71	starvation, longevity

Supplemental Figure S4: HiTmiSS Comparison and Intestine Tissue *Switchers*. (A)
Comparison of the distribution of *5p Singletons*, *3p Singletons*, and *Switchers* between
HiTmiSS analysis results in mixed tissue (left) and intestine tissue (right). Results expressed
as a percentage of total detected miRNAs per analysis. *Singleton* strand preference shifted
to favor *5p Singletons* and overall abundance of *Switchers* steeply declined, both of which
suggest that strand selection is regulated at the level of individual tissues. **(B)** Breakdown of
Switchers detected in the intestine tissue by miRNA family and function. *Switcher* in this case
was defined as any miRNA whose strand preference changed from embryo to young adult
stage, either 5p to 3p or vice versa. Though few in number, each *Switcher* regulates target
3′UTRs of genes directly involved in the function of the intestine, further reinforcing the
importance of the tissue-specific regulation of strand selection.

A

pges1::ROG::fox-1 3′UTR

pges1::ROG::hrp-2 3′UTR

B

pges1::ROG::fox-1 3′UTR

pges1::ROG::hrp-2 3′UTR

C

Log$_2$ Fold Change in miRNA Abundance

Threshold of significance

■ *pges1::ROG::fox-1* 3′UTR   ■ *pges1::ROG::hrp-2* 3′UTR

Supplemental Figure S5: Testing the Target Stabilization Hypothesis. **(A)** Constructs used to overexpress GFP under the control of the *fox-1* 3′UTR (top) or *hrp-2* 3′UTR (bottom) using the promoter for the intestine-specific gene *ges-1.* miRNA strands which bind each target 3′UTR (shown above) are typically not expressed in wildtype N2 *C. elegans*. (B) Representative brightfield (left) and GFP (right) images of C. elegans strains expressing these constructs, demonstrating fluorescence in the intestine. (C) Change in abundance of each miRNA strand, expressed as log$_2$ of the difference in mean expression value as per our HiTmiSS assay. None of the tested miRNA strands were significantly (log$_2$ > 1) upregulated when overexpressing either test 3′UTR. These data suggest that overexpression of targets for a miRNA strand does not lead to retention of that strand, but rather strand selection occurs independent of the presence or absence of target 3′UTRs.

lin-4

| WT | ACC → UCG | UU → GUA | Combined |

Variant	5p	3p
WT	0.81	0.19
ACC → UCG	0.74	0.26
UU → GUA	0.77	0.23
Combined	0.76	0.24

miR-52

| WT | Del A | CC → U | Combined |

Variant	5p	3p
WT	0.89	0.11
Del A	0.83	0.17
CC → U	0.90	0.10
Combined	0.88	0.12

miR-75

| WT | Pair 3' 9-10 | Unpair 3' | Combined |

Variant	5p	3p
WT	0.36	0.64
Pair 3' 9-10	0.30	0.70
Unpair 3'	0.60	0.40
Combined	0.57	0.43

miR-60

| WT | Ins AA | 3p 5'U → A | Combined |

Variant	5p	3p
WT	0.41	0.59
Ins AA	0.46	0.54
3p 5'U → A	0.46	0.54
Combined	0.52	0.48

Supplemental Figure S6: Predicted Strand Usage in *5p* and *3p Singletons*. Representative, well-characterized *5p Singletons* (top) and *3p Singletons* (bottom) were selected and their strand preference predicted by our ML model (WT, far left). Each underwent two *in silico* site-directed mutations, one in the bulge region (middle left) and one in the pairing/identity of the first nucleotide (middle right). Finally, a combination mutation of both is shown (far right). The confidence of prediction is displayed as a decimal, both *5p Singleton* confidence (5p) or *3p Singleton* confidence (3p). These mutations consistently altered predicted strand confidence, with a more pronounced effect in *3p Singletons*, supporting the trends within our wet bench data.