An AI-Guided Framework Reveals Conserved Rules Governing

microRNA Strand Selection

Author list

Dalton Meadows[1,2],

Hailee Hargis[1,2],

Amanda Ellis[1,2],

Heewook Lee[1],

Marco Mangone[1,*]

Author affiliations

[1)] The Biodesign Institute at Arizona State University, 1001 S McAllister Ave, Tempe, AZ

[2)] School of Life Sciences, Arizona State University, 427 E Tyler Mall, Tempe, AZ

Author list footnotes

[*)] Corresponding author

Corresponding author and e-mail address

Marco Mangone; mangone@asu.edu

Abstract

MicroRNAs (miRNAs) are central regulators of gene expression, yet the fundamental question of how cells choose between the two strands (5p or 3p) of a miRNA duplex during biogenesis remains unresolved. Here, we present a comprehensive, experimentally grounded framework that decodes the logic of miRNA strand selection. Using *Caenorhabditis elegans* as a model system, we developed a high-throughput qPCR platform enabling precise quantification of strand usage across development and in specific somatic tissues. Our analysis reveals that strand selection is not stochastic but follows conserved, strand-specific rules integrated with cellular and developmental context. To uncover the molecular grammar underlying this decision, we used artificial intelligence to build a predictive machine learning model trained on experimentally validated data. This AI-driven model, integrating 77 biologically informed features, accurately predicts strand preference not only in nematodes but also across vertebrates, including humans, revealing compositional and structural biases that are conserved yet functionally repurposed. Our work establishes a new paradigm for understanding miRNA strand selection, combining large-scale experimentation with AI to reveal an ancient, conserved, and programmable layer of gene regulation.

Introduction

miRNAs are ~22-nucleotide non-coding RNAs that guide the RNA-induced silencing complex (RISC) to target mRNAs, primarily through base pairing within the 3′ untranslated regions (3′UTRs) [1]. Once bound, the miRNA–RISC complex (miRISC) typically represses translation of the target mRNA. Both miRNAs and their 3′UTR targets are often conserved across metazoans and play key roles in essential biological processes such as development and morphogenesis.

miRNAs have undergone several pronounced expansions in metazoan genomes [2], coinciding with evolving organismal complexity yet rarely corresponding with increases in protein-coding genes [2]. Eukaryotic genomes contain 100-2,000 distinct mature miRNAs in intergenic regions, introns, and in clusters. Following transcription in long primary miRNA transcripts (pri-miRNAs) by the RNA polymerase II, they are processed in the nucleus by the microprocessor complex into shorter hairpin structures known as precursor miRNAs (pre-miRNAs) [1]. The two strands composing the pre-miRNAs are named the 5p and the 3p strand, respectively. Pre-mRNAs are subsequently exported to the cytoplasm by Exportin-5, where they undergo further cleavage by the enzyme Dicer, producing mature miRNA duplexes [1].

The Hsc70-Hsp90 chaperone complex facilitates the ATP-dependent loading of the miRNA duplex into Argonaute, the protein that execute the silencing step. This complex is thought to induce conformational changes in Argonaute that enable duplex loading [3-5]. Once loaded, one strand of this duplex, the guide strand, is kept within the RISC, while the other, the passenger strand, is typically degraded. Both the 5p and the 3p strands

can be selected for miRISC loading and act as guide strands. It is currently unclear how

this process, termed 'miRNA strand selection,' is executed in detail, which proteins are

involved in this decision, and how the guide strand is ultimately chosen between the two

strands. The predominant factors influencing miRNA strand selection are believed to be

the specific 5' nucleotide sequences of each strand and the comparative

thermodynamic stability exhibited by the two ends of the miRNA duplex [6]. Studies

performed using the human Ago2 protein, an Argonaute protein member of the RISC

complex, show that the pocket responsible for binding the 5' nucleotide of the guide

strand exhibits a strong preference for Uracil nucleotides [7]. This aligns with the

prevalent occurrence of Uracil as the 5' nucleotide in miRNA guide strands [2], whereas

the passenger strand commonly initiates with a Cytosine at its 5' end.

Furthermore, the miRISC complex exhibits a preference for loading the end of the

miRNA duplex with lower thermodynamic stability, although not all the time [8]. The initial

four nucleotides on each end of the duplex are believed to establish this thermodynamic

asymmetry, and in some cases, a slight difference in the number of hydrogen bonds

can induce the preferential loading of the less stable strand *in vitro* [5].

Consistently, the 5' ends of guide strands are enriched with purines [2]. In comparison,

the 5' ends of passenger strands are predominantly pyrimidine-rich in both human and

fly miRNAs, contributing to the thermodynamic asymmetry observed between the two

ends of the duplex. In *C. elegans*, single amino acid substitutions within the MID

(G553R) and PIWI (G716E, S895F, and S939F) domains of the Argonaute protein ALG-

1 invert the strand selection process for specific miRNAs, suggesting a potential role for

this protein in the miRNA strand selection process [9]. This indicates that while these

mutations play a pivotal role, they do not consistently reverse strand selection for all

miRNAs, suggesting that there may be additional internal or external factors involved in

this process. Tissue specific accessory factors and their dosage could also be

responsible of miRNA strand selection, altering strand choices based on tissue or

developmental contexts, similarly to the role of the SR proteins in RNA splicing [10]. While

thermodynamic and sequence features help explain strand choice, a comprehensive

mechanistic framework has been lacking.

C. elegans is an ideal model for studying miRNA strand selection due to its compact

3′UTRome [11-14], small number of miRNAs (miRbase), fully mapped cell lineage [15], and

well-characterized transcriptome. Here we developed a high-throughput qPCR platform

to systematically profile miRNA strand selection events across *C. elegans* development

and in somatic tissues. These data revealed conserved sequence and structural

patterns, which we used to train a machine learning model that predicts strand selection

with 83% accuracy and generalizes to human miRNAs.

Materials and methods

All DI water was obtained from either a Purelab Option system (ELGA LabWater) or

an Avidity Science system (Duo) and autoclaved before use. Chemicals were ordered

from Sigma Aldrich/Merck, VWR, or Thermo Fisher Scientific in molecular biology grade

and stored according to company instructions. Gateway[TM] LR Clonase II Plus[TM] enzyme

mixes, SuperScript[TM] III Reverse Transcriptase, and PowerTrack[TM] SYBR Green Master

Mix were purchased from Invitrogen. Primers for the HiTmiSS assay were ordered from

Life Technologies. Poly(U) Polymerase and restriction enzymes used in validation were

ordered from New England Biolabs. The pCFJ150 plasmid was a gift from Erik

Jorgensen. The OP50-1 *E. coli*, N2 *C. elegans* and EG6699 *C. elegans* strains were

purchased through the Caenorhabditis Genetics Center.

Maintenance of *C. elegans* strains

All *C. elegans* strains were cultured on NGM plates seeded with OP50-1 bacteria. All

transgenic lines produced for experiments described in this study were maintained at

24°C. When bacteria on each NGM plate had been cleared, *C. elegans* strains were

transferred onto new plates via washing into a 15 ml Falcon tube using 15 ml M9 media

(22 mM KH_2PO_4, 42 mM Na_2HPO_4, 85.5 mM NaCl, 1 mM $MgSO_4$, sterilized by

autoclave) and pelleted for 3 minutes at 400*g*. The M9 media was then aspirated to a

volume of 500 ml and the animals were resuspended via gentle pipetting before being

moved onto fresh, seeded NGM plates.

Synchronization of *C. elegans* strains

The EG6699 strain [16] was maintained at 24°C. To synchronize animals for injection,

confluent plates were bleached to isolate the embryos. Each plate was washed into a

15 ml Falcon tube with 15 ml DI water and pelleted for 3 minutes at 400*g*. The water

was then aspirated, and the animals were incubated in 5 ml of bleaching solution (0.5M

NaOH, 10% bleach) for 20 minutes with gentle end-over-end rotation. After bleaching,

the embryos were washed twice with DI water and moved onto fresh, seeded NGM

plates. For use in the HiTmiSS assay, both N2 and *pges-1::GFP::alg-1::unc54 3′ UTR*

C. elegans strains were synchronized as described above.

Primers design for the HiTmiSS Assay

To quantify microRNAs in high throughput, we designed 380 unique forward primers

for use in the qPCR amplification step of the HiTmiSS assay. These primers were

designed to be complimentary to either the 5p strand or the 3p strand of 190 most

expressed microRNAs in the *C. elegans* genome, corresponding to 75% of all *C.*

elegans miRNA deposited in miRbase (release 22.1). We did not consider microRNAs

whose existence is not supported by broad experimental evidence nor any microRNAs

predicted to exist. Few miRNAs were ignored because their corresponding primers had

a very low Tm, making it impossible to process using our pipeline. All primers were

ordered in four 96-well microplates from Life Technologies pre-diluted at a concentration

of 50 mM (**Supplemental Table S1**). We designed a universal reverse primer

complementary to the stem-loop sequence of the stem-loop poly(A) primer used to

prepare the cDNA for this qPCR reaction. 24 replicates of this primer were ordered from

Life Technologies in a separate array, also at a concentration of 50 mM. Both forward

and reverse primers were re-arrayed and diluted to a working concentration of 10 mM

immediately prior to use.

Singletons **and** *Switchers* **identification**

For each strand, we subtracted its relative quantity at one life stage versus the

relative quantity of the complementary strand (i.e. for *let-7* 5p, we compared the

abundance at embryo stage versus the abundance of *let-7* 3p at the embryo stage).

This was repeated for each life stage at which the expression of that microRNA strand

fell above our significance threshold (1 standard deviation above the mean of all

quantified miRNAs at a given stage), and any microRNAs for which their expression

was greater at each such life stage were deemed *Singletons*. We classified as *5p*

Singletons microRNAs where the 5p strand was more expressed at each life stage than

its corresponding 3p strand. We classified as *3p Singletons* microRNAs where the 3p

strand was more expressed at each life stage than its corresponding 5p strand. were

We classified as *Switchers* microRNAs which rose above our threshold of significance

during at least one life stage, but whose expression levels did not display a consistent

pattern of usage of one strand over the other.

HiTmiSS Assay

To collect sufficient total RNA for analysis of wild type miRNA, confluent NGM plates

of N2 strains were synchronized and either washed and pelleted immediately (embryo

stage) or incubated at 24°C for 15 hours (larval 1 stage), 25 hours (larval 2 stage), 33

hours (larval 3 stage), 42 hours (larval 4 stage), or 51 hours (young adult stage) before

washing and pelleting. All washes were performed in 15 ml Falcon tubes using 15 ml DI

water, and pellets were generated by centrifugation for 3 minutes at 400g, resulting in a

final pellet volume of 500 µL to 1 mL. Sufficient worms to generate this size pellet

differed by life stage, requiring 12 (larval 4 and young adult stages), 18 (larval 2 and

larval 3 stages), or 24 (embryo and larval 1 stages) starting confluent plates.

This pellet was then resuspended to a volume of 1.5 ml in ice cold Triton X-100 lysis

buffer (150 mM NaCl, 1.0% Triton X-100, 50 mM Tris-HCl pH 8.0, 0.5 mM EDTA) and

transferred to a 2 ml Spectrum tube pre-filled with 0.5 mm zirconium beads. Using a

Benchmark BeadBug D1030 microtube homogenizer, each tube was homogenized for

90 seconds, incubated for 2 minutes on ice, then homogenized for an additional 60

seconds. The resultant lysate mixture was centrifuged for 3 minutes using a VWR

benchtop mini centrifuge and the liquid fraction was transferred into a 1.5 ml Eppendorf

microcentrifuge tube. To minimize non-aqueous contaminants, this lysate was

centrifuged again in the same manner and transferred to a 15 ml Falcon tube pre-chilled

on ice. To isolate total RNA, we used a DirectZol RNA MiniPrep Plus kit from Zymo

Research, following kit instructions and performing the entire procedure at 4°C.

Resultant RNA was eluted in 30 µL NF H_2O. To collect tissue-specific miRNA for

analysis of the intestine, we performed protein immunoprecipitation followed by miRNA

isolation. The fluorescent *pges1::gfp::alg-1::unc54 3′UTR C. elegans* strain previously

prepared by our lab [17] was grown to confluence, synchronized, and pelleted as

described above for each life stage, concurrent with N2 *C. elegans* as a negative

control.

Each pellet was resuspended to 1.5 µl using our Triton X-100 lysis buffer with added

protease inhibitor (two Roche cOmplete ULTRA mini EDTA-free protease inhibitor

tablets per 50 ml lysis buffer) and transferred to a zirconium bead tube to be

homogenized as previously described. During this homogenization step, anti-GFP

magnetic antibody beads (Chromotek GFP-Trap Magnetic Agarose) were prepared by

diluting 25 µl bead slurry in 500 µl dilution buffer (150 mM NaCl, 50mM Tris-HCl pH 8.0,

0.5 mM EDTA) and rotating end-over-end at 4°C for 3 minutes.

The beads were pelleted using a benchtop centrifuge, and the buffer was aspirated

using a magnetic tube rack. 500 µl dilution buffer was added again, and this dilution

process was repeated twice. We then recovered aqueous lysate from the homogenized

mixture as previously described and diluted it using 300 µl dilution buffer. This diluted

lysate was transferred to a tube containing the prepared anti-GFP beads and left to

rotate end-over-end at 4°C overnight. To clear unbound lysate, the beads were pelleted

using a benchtop centrifuge and buffer was aspirated using a magnetic tube rack. 500

µl of wash buffer (150 mM NaCl, 50 mM Tris-HCl pH 8.0, 0.1% Triton X-100, 0.5 mM

EDTA) was added, the beads were pelleted, and the buffer was aspirated using a

magnetic tube rack. This wash step was repeated five additional times. The beads were

then resuspended in 100 µl wash buffer and RNA was isolated using a DirectZol RNA

MiniPrep Plus kit from Zymo Research, following kit instructions and performing the

entire procedure at 4°C. Resultant RNA was eluted in 30 µL NF H$_2$O. The protocol for

the HiTmiSS assay was adapted from Mei *et al.* [18]. We prepared our cDNA from each

life stage for qPCR analysis via polyuridinylation using Poly(U) Polymerase (NEB) as

per manufacturer instructions followed by a 20-minute heat inactivation at 65°C. We

performed a first-strand reaction on this polyuridinylated RNA using Superscript III

Reverse Transcriptase and stem-loop poly(A) primers under recommended conditions

(Invitrogen). We then performed qPCR in high throughput using this cDNA, SYBR

Green master mix (Invitrogen), and our primers as described above. We used Applied

Biosystems Quantstudio 3 and Quantstudio 7 qPCR machines to perform the screen.

For each life stage, four 96-well plates were prepared with 40 μL reactions in each well,

along with a positive control reaction containing 10 pg pDONR 221 ROG instead of

cDNA and the necessary forward and reverse primers to amplify *gfp*. Each plate was

then re-arrayed in triplicate into a 384-well plate using a Biomek FX^P Laboratory

Automation Workstation for a total of three 10 μL reactions per microRNA strand per life

stage. An additional plate was prepared with no cDNA or plasmid spike-in DNA as a

non-template control. These qPCR reactions were carried out under standard SYBR

Green reagent conditions on an Applied Biosystems QuantStudio 7. Relative miRNA

levels were calculated using the $2^{-\Delta\Delta Ct}$ method (see below), and normalized using the

spike-in positive control. These results were then plotted logarithmically as a heatmap.

Whole worm data were ordered by type (*5p Singleton*, *3p Singleton*, or *Switcher*), while

intestine data were clustered using average linkage. Both data were visualized using

heatmapper.ca.

Cycle Threshold Analysis of the HiTmiSS Assay

The qPCR results containing the cycle threshold (Ct) values for each tested miRNA

was exported in Microsoft Excel 2019 format. A comprehensive list of these data can be

found in **Supplemental Table S2**. Ct values were averaged across experimental

replicates, then subjected to a $2^{-\Delta\Delta Ct}$ analysis to calculate initial amounts of each

microRNA in the sample, consistent with experimental standard [19]. The Ct value for any

given microRNA strand was subtracted from the Ct value of the non-template control (a

qPCR reaction lacking any cDNA) for that microRNA strand. This was termed the ΔCt

for that microRNA. The same was done for the positive control for that sample. Then,

for each sample, this difference was subtracted from the difference obtained for that

positive control, normalizing samples to the same positive control and giving us the

ΔΔCt for that microRNA strand.

The inverse log base 2 of that ΔΔCt was then calculated to approximate the initial

amount of that miRNA strand present in the cDNA relative to the positive control. This

was deemed the $2^{-\Delta\Delta Ct}$ value for that miRNA strand. To determine whether a miRNA

was considered present in a sample, we decided on a strict threshold of one standard

deviation above the mean of all $2^{-\Delta\Delta Ct}$ values for that stage of development. MiRNAs

quantified at levels above this threshold were considered significant for the purposes of

this study. This same process was repeated to determine significance for our intestine

tissue-specific HiTmiSS assay.

qPCR of microRNAs targeting overexpressed intestinal 3′UTRs

To quantify miRNA strand the levels when the target 3′ UTRs of those strands are

overexpressed in the intestine, we used two *C. elegans* strains previously published by

our lab [17]. Each expresses the promoter for intestine-specific gene *ges-1* fused to a

dual color ROG construct (a gene encoding *mCherry* and *gfp* separated by a trans-

spliceable operon) fused to the 3′ UTR of either *fox-1* or *hrp-2* [17]. Each of these strains

was grown to confluence in 12 NGM plates in mixed stage and subjected to

quantification using the same approach as the HiTmiSS assay described above, though

rather than quantifying every microRNA in these animals, we quantified the 5p and 3p

strand of *miR-54*, *miR-60*, *miR-232*, and *miR-239b*. These quantifications were carried

out in 96-well plates using cDNA (both prepared from our strains as well as control

cDNA gathered from N2 *C. elegans*), SYBR Green master mix, and the appropriate

forward and universal reverse primers for the miRNAs of interest, re arrayed from our

standard HiTmiSS plates. Each 96-well plate contained two positive controls in *let-7* and

lin-4, as well as a non-template negative control for each set of primers. These qPCRs

were carried out in experimental triplicate under standard SYBR Green reagent

conditions on an Applied Biosystems QuantStudio 3.

Relative miRNA levels were calculated using the $2^{-\Delta\Delta Ct}$ method as described above

and normalized using the average of *let-7* and *lin-4* expression for each sample. These

results were then averaged across two replicates and analyzed using Microsoft Excel

2019. Data were visualized as $\log_2$ of the change in expression from wildtype N2

negative control to either of our experimental strains. The threshold of significance was

set at 1 as per standard analytical procedure.

Structural Analysis of microRNAs

The experimentally validated *5p* and *3p Singletons* were obtained using our

HiTmiSS assay. Each entry included the entire hairpin sequence and its predicted

secondary structure in dot-bracket notation [20] (**Supplemental Table S2)**. To focus

specifically on mature miRNA regions, only the sequences of each mature miRNA

duplex were used for all downstream analyses. Since the *5p* and *3p Singleton* datasets

contained different numbers of miRNAs, we balanced them before analysis by

undersampling the larger group to match the sample size of the smaller class, ensuring

unbiased statistical comparison between 5p and 3p strands.

We extracted 77 numerical features from each *5p* and *3p Singleton* (**Supplemental**

Table S3). The extracted features included: Base composition (fraction of A, U, G, C),

motif presence (e.g. UGU, GAG, CCU), positional characteristics (first/last nucleotide

identity), dinucleotide frequencies (e.g., di_GG, di_CC), GC content and GC-related

asymmetry metrics (GC_diff, GC_ratio_5p3p), strand lengths, and number of

mismatches (unpaired positions in the dot-bracket structure). The full list of the features

and their descriptions are shown in **Supplemental Table S3**.

Feature calculations were implemented in Python using regular expressions and

sequence statistics functions. Structural mismatch counts were determined by counting

unpaired characters within the uppercase-aligned mature duplex. Comparative

statistical analysis between 5p and 3p strands was conducted using Welch's two-

sample t-test. All features were tested independently, and results were ranked by p-

value. Features with $p < 0.05$ were considered statistically significant. Bar plots were

generated for each significant feature using the mean ± standard deviation across

strands.

miRNA strand selection ML model

To predict dominant strand selection (5p vs. 3p) in miRNA hairpins, we utilized

ChatGPT GPT-4o (OpenAI) to assist in developing and training a Random Forest

classifier based on 77 biologically curated features, organized into functional categories

capturing distinct structural and sequence-related aspects of miRNA hairpins. These

features capture thermodynamic stability, base-pairing patterns, and positional

relationships within the miRNA hairpin. The model was trained on our validated *5p* and

3p Singletons identified using our HiTmiSS assay, and feature values were computed

using custom Python scripts that parse dot-bracket RNA structures and extract both

absolute and relative measures for each strand. The weights of the features were

optimized using known *Singletons*. The final model incorporates the following features,

divided into five categories. The first category is the *Dinucleotide Composition* features,

which quantify the frequency of specific two-nucleotide motifs within the 5p and 3p

regions, capturing local sequence patterns that can influence strand selection and

structural flexibility. The second category is the *First/Last Nucleotide Bias* features,

which record the identity of the first and last nucleotides in each strand, reflecting known

preferences in Argonaute loading where terminal nucleotides strongly impact strand

choice. The third category is the *GC Content and Entropy* features, which assess the

global nucleotide composition and thermodynamic diversity within each strand, including

overall GC percentages, GC ratios between strands, and Shannon entropy measures

that reflect structural variability and rigidity. The fourth category is the *Base-pairing*

Properties features, which describes the fraction of nucleotides engaged in base pairing

within the 5p and 3p regions, providing a measure of local hairpin stability and

asymmetry. Finally, the *General Structural* features, which include the overall hairpin

length and the length difference between the 5p and 3p strands, capture broader

geometrical properties that may indirectly affect miRNA maturation and strand retention

(**Supplemental Figure S1**). The complete set of features, their description, and the

weight used are shown in **Supplemental Table S3**. The model showed high

performance in distinguishing 5p and 3p usage from a pool of known *Singletons* as a

test dataset (*5p Singleton*: accuracy 100%, precision 75%, recall 100%; *3p Singleton:*

82% accuracy, 100% precision, 82% recall) (**Supplemental Figure S2**). The ML model

uses a Random Forest classifier implemented in scikit-learn (v1.3.1). Model

hyperparameters were optimized via grid search with 5-fold stratified cross-validation,

tuning: n_estimators: [100, 250, 500], max_depth: [5, 10, None], min_samples_split: [2,

5, 10], class_weight: ['balanced', None]. The final selected parameters were:

RandomForestClassifier (n_estimators=250, max_depth=None, min_samples_split=2,

class_weight='balanced'). To prevent overfitting and enhance interpretability**,** recursive

feature elimination (RFE) with cross-validation was used to rank features by importance.

Features with near-zero variance and multicollinearity (|Pearson's r| > 0.9) were

removed. The model consistently retained ~45–55 features, depending on the species

subset. Performance was assessed on independent held-out test sets not used during

training. Metrics included: Accuracy: 0.85 (*C. elegans*), 0.83 (human), Precision (5p/3p)**:**

0.87 / 0.80, Recall (5p/3p)**:** 0.89 / 0.78, F1 Score (5p/3p)**:** 0.88 / 0.79**,** AUROC**:** 0.91 (*C.*

elegans), 0.89 (human). Class balance was maintained using SMOTE oversampling to

correct for the slight enrichment of 5p strands in the source datasets. We confirmed that

the model generalizes well by testing it on separate datasets (not used during training or

tuning), including across species (e.g., training in *C. elegans*, testing in humans).

ML model predictions in human miRNAs

Human miRNA expression data were obtained from a harmonized dataset from the

human miRNA tissue atlas [21] containing strand-specific Reads Per Million Mapped

(RPMM) values across healthy human tissues. To restrict the analysis to non-diseased

samples, we only considered entries where the Biotype column was labeled as "tissue"

"EV", or "cell_line" (**Supplemental Table S6**), representing the average expression

values across biological replicates of healthy tissues (total=337 datasets). For each

miRNA duplex, we extracted the expression levels corresponding to the 5p and 3p

strands, based on strand-specific annotations. We assigned strand dominance by

summing RPMM values across all healthy tissue samples for the 5p and 3p strands

separately. The strand with the higher total expression was designated as the dominant

strand. We compared these dominant strands assignments with the predicted dominant

strand assigned by our ML model. Each prediction was considered correct if the model-

assigned strand matched the strand exhibiting greater expression across healthy

tissues. Evaluation was performed on the topmost highly expressed 10, 50, and 100

miRNAs ranked by the sum of 5p and 3p strands RPMM values. We also performed the

analysis using the complete dataset (all).

Results

A novel HT qPCR approach to study miRNA biogenesis

Identifying miRNAs in somatic tissues is challenging due to their small size, low

abundance, and short half-life. To overcome this, we developed a high-throughput

qPCR method that reliably quantifies miRNA expression and distinguishes 5p from 3p

strand usage.

This method is a high throughput adaptation and general improvement of an existing

methodology [18], which uses the polyuridine polymerase enzyme to elongate a polyU tail

to all RNA transcripts in a given sample. This step extends the length of each miRNA

present in the sample, which is then used as a template in reverse transcriptase

reactions using a specially engineered universal stem-loop reverse primer (SL-polyA)

(**Main Fig. 1A**).

We named this method High-Throughput miRNA Strand Selection (HiTmiSS) assay.

HiTmiSS assay enables rapid, precise quantification of the full miRNAome and

accurately distinguishes 5p from 3p strand expression (**Main Fig. 1**).

We employed this assay to systematically quantify miRNA occurrence and the

usage of 5p and 3p strands in known miRNAs throughout *C. elegans* development. We

extracted total RNA from *C. elegans* at six developmental stages (embryos, larval

stages 1–4, and young adults) and analyzed the miRNA transcriptome and strand

selection for 190 miRNAs, representing 75% of all miRNAs annotated in miRbase, using

380 unique forward primers across four plates (**Supplemental Table S1**).

We performed 7,980 qPCR experiments independently, querying these 190 miRNAs

throughout development in triplicate (1,140 experiments at each stage) (**Main Fig. 1B,**

Supplemental Fig. S3). Using stringent filters (see Materials and Methods), we

detected the presence of 177 of these miRNAs (93.2%) in at least one developmental

stage (**Supplemental Table S2**). The greatest number of miRNAs occur during L4

stage (81%), followed by young adult (75.5%), L3 (65%), L2 (56.8%), and lastly,

Embryo and L1 (41.6% each) stages. This aligns with previous literature suggesting that

many *C. elegans* miRNAs are tissue-specific, only becoming more abundant upon the

development of those tissues later in the developmental cycle [22].

As previously reported, early expressed miRNAs such as the key regulators of

embryonic lethality *miR-35* and *miR-51* families, were most abundant during embryonic

and early larval stages [23]. In contrast, *let-7* and *lin-4*, which control adult phenotypes,

peaked at the L4 and young adult stages [24]. Notably, 13 miRNAs, including *miR-791*,

miR-792, and *miR-8206*, showed no significant expression, likely due to low abundance

or lack of experimental validation.

Many miRNAs show consistent strand preference throughout development

Many miRNAs identified consistently use either the 5p or 3p strand as the guide

strand, which we termed '*Singletons*', while others switch between 5p and 3p usage

across different developmental stages. We named the second category '*Switchers*'.

The screen produced hits for 177 miRNAs, which we divided into *5p Singletons*, *3p*

Singletons, or *Switchers* (**Main Fig. 1B and Supplemental Table S2**).

The largest group identified was miRNA *Switchers* (46.3%, 88 hits). These include

members of the *miR-2* family and other regulators of developmental timing and stress

responses [24,25]. The remaining hits were *Singletons*, with a notable bias: *3p Singletons*

(58 hits, 30.5%) outnumbered *5p Singletons* by nearly twofold. These include *miR-35*

and *miR-51* (embryonic lethality), *miR-44* (stress/starvation), and *miR-58* (locomotion

and dauer entry) [23,24,26]. Several abundant *3p Singletons*, including *miR-60*, *miR-70*, and

miR-90, remain functionally uncharacterized but are linked to stress response [25,27,28].

5p Singletons were the least common group, with 31 hits (16.3%), including the well-

studied families *let-7* and *lin-4*, which are key regulators of developmental timing (**Main**

Fig. 1B). Also detected were *miR-51* (embryonic lethality) and *miR-64* (heat shock

response) [23,28]. While some families (*let-7* and *miR-64*), consistently use the same

strand, others span multiple categories. For instance, the *miR-51* family splits evenly

between *5p* (*miR-51* to *-53*) and *3p Singletons* (*miR-54* to *-56*, *lsy-6*), showing no

Switchers, highlighting that strand usage can diverge even within structurally similar

family members.

Importantly, several hits produced by the HiTmiSS assay align with past findings on

the strand preference of well-characterized miRNAs such as *let-7, lin-4* (known *5p*

Singletons), and the *miR-35* family (known *3p Singletons*) [24]

To account for the strong strand preference observed among *Singletons*, we took

the top 20 *5p Singletons* and the top 20 *3p Singletons* and compared the binding

signature and nucleotide composition between the used strand versus the strand that

gets discarded (**Main Fig. 1C, i and ii**). Interestingly, the pairing frequency does not

vary between these miRNAs (**Main Fig. 1C,ii**), either in the 5p or 3p strand, except for

the last few nucleotides at the 3'end (**Main Fig 1C, I, red boxes**) which are commonly

unpaired in the mature duplex.

When comparing the used vs. the discarded strand (**Main Fig. 1C,** left panel vs right

panel), we detected no significant difference in the pairing signature at any position;

however, the 5p strand, independently if is used or discarded, contains more unpaired

nucleotides (**Main Fig. 1C,** red boxes). When narrowing our analysis to just the seed

region (**Main Fig. 1C**, golden boxes), while there is higher pairing than any other

position, there is no difference in binding frequency between the used and discarded

strands, suggesting that other rules are also in place.

Next, we compared the nucleotide composition of these seed regions, hoping to

identify signatures that could explain the discrepancy in miRNA strand selection in

Singletons. We determined the likelihood of any given nucleotide from positions 1

through 10 in each miRNA strand in the top 20 *Singletons* in each category (5p and 3p)

(**Main Fig. 1C, ii**). We detected a highly conserved uracil nucleotide at position 1, as

previously shown by us and others [2], though this conservation dissolves from positions

2 through 10 (**Main Fig. 1C, ii**). Importantly, this uracil nucleotide is not equally

conserved between *5p Singletons* and *3p Singletons*. 80% of *3p Singletons* start with

this uracil nucleotide, while less than 40% occur in *5p Singletons*. This differential

presence of this uracil may be caused by a non-stochastic pre-miRNA cleavage by the

C. elegans DCR-1, which may possess a preference to cleave at this nucleotide.

The intestine miRNA population is small but highly interconnected

The vast majority hits detected by our HiTmiSS assay were *Switchers* (~46%). Many

of them may indeed not be true *Switchers* but rather act as tissue- or stage-specific

Singletons, potentially confounding the analysis. To address this, we refined our

approach by analyzing somatic tissues across development, focusing on the intestine,

which is a large tissue with a well-characterized transcriptome [29,30].

We coupled our HiTmiSS assay with intestine-specific miRNA immunoprecipitation,

as we have published in the past [17], focusing on the embryo vs young adult stages.

More specifically, we forced the expression of the GFP-tagged *C. elegans* Argonaute

ortholog *alg-1* in the intestine tissue using the intestine-specific promoter *ges-1p*,

performed an α-GFP immunoprecipitation, recovered the miRNA strands loaded onto

ALG-1, and used them as templates in our HiTmiSS assay (**Main Fig. 2A**).

We found that of the 96 miRNAs tested, 64 (68%) were present during at least one

stage within the intestine tissue (**Main Fig. 2**). Of these, 58 (91%) were *Singletons*,

while 6 (9%) were *Switchers* (**Supplemental Fig. S4A**).

This strongly differs from our whole worm data, reinforcing that miRNAs previously

labeled as *Switchers* may truly be *Singletons* on a tissue-by-tissue basis, only

appearing as *Switchers* once all tissues are analyzed together.

Interestingly, the proportion of *5p Singletons* to *3p Singletons* is not the same in the

intestinal tissue as in the mixed tissue (**Supplemental Fig. S4A**). In our initial analysis,

about two-thirds of *Singletons* were *3p Singletons*, yet in the intestine tissue, nearly two-

thirds of *Singletons* are *5p Singletons* (**Supplemental Fig. S4A**).

Another surprise was the presence of intestine-specific *Switchers* (**Main Fig. 2A and**

Supplemental Fig. S4B). This includes the *miR-51* family, well-characterized regulators

of embryonic lethality [23]. In addition, we detected intestine tissue *Switchers* among the

miR-44, miR-58, and *miR-71* families (**Supplemental Fig. S4B**). These miRNA families

have diverse functions, from regulating developmental timing to starvation, hypoxia,

stress response, and dauer entry [25,28,31,32]. Importantly, these are intestine-specific

phenotypes, suggesting that strand switching in these miRNAs may have functional

significance by expanding 3'UTR targeting repertoires in the intestine.

stage. These strains expressed the promoter for a single miRNA fused to GFP, and live

imaging was used to validate the expression of these miRNAs at the appropriate life

stage in the intestine (**Main Fig. 2C**). All four strains displayed the expected

fluorescence.

miRNA target presence doesn't account for strand usage

Our HiTmiSS array enables precise, strand-resolved quantification of miRNAs *in vivo.*

We considered the possibility that miRNA targets might stabilize specific strands,

potentially biasing detection in our HiTmiSS assay. In this model, only target-bound

strands would be retained, while the unused strand would degrade, falsely suggesting

strand preference. To test this, we overexpressed *fox-1* and *hrp-2* 3'UTRs, each

containing multiple binding sites for typically discarded strands (e.g., *miR-60* 5p, *miR-*

232 5p, *miR-239b* 5p), in intestinal tissue (**Supplemental Figure S5**). If target-induced

stabilization were driving strand selection, we would expect increased detection of these

5p strands. However, strand usage remained unchanged compared to wild-type

controls, with 3p strands (i.e. *miR-60* 3p, **Supplemental Figure S5C**) still

predominating.

Taken together, these results indicate that our qPCR approach is not simply

detecting stabilized miRNA strands due to the presence of targets. Rather, the relative

strand abundance measured by the HiTmiSS assay likely reflects genuine strand

selection, rather than artifacts introduced by target availability.

***5p Singletons* and *3p Singletons* follow different rules for selecting strand usage**

Next, we studied the structural features associated with *Singletons*, which could

explain the strict and consistent strand usage within these miRNAs. When comparing

the used and the discarded strand, the pairing frequency varies significantly within each

Singleton, revealing new conserved patterns (**Main Fig. 3**). Specifically, the strand used

in *Singletons* often has an unpaired nucleotide at position 12, while the discarded strand

does not (**Main Fig. 3A**, yellow boxes). This lack of pairing in position 12 in both *5p*

Singletons and *3p Singletons* usage is intriguing. In humans, certain Argonaute proteins

perform cleavage of target RNAs between positions 10 and 11, which corresponds to

positions 10 and 11 along the bound miRNA strand by Watson-Crick pairing [33]. *C.*

elegans ALG-1 is largely homologous to human Argonaute proteins, and it may retain

this cleavage activity at similar target positions, leading to the degradation of the unused

strand due to its Watson-Crick base pairing at this crucial position. In addition, we

detected differences in the binding frequency of the first-position nucleotide in both *5p*

Singletons and *3p Singletons* (**Main Fig. 3A,** blue boxes). *3p Singletons* also exhibit

unique pairings at positions 18-19 (**Main Fig. 3**, white box) not present in the discarded

strain. These patterns are most clearly seen in specific *Singleton* miRNAs, both *5p*

Singletons such as *miR-52* and *3p Singletons* such as *miR-75* (**Main Fig. 3B**).

An updated machine learning (ML) model for miRNA strand selection

We wanted to extract patterns and features that make *Singletons* so unique and predispose them to always use one strand over the other. To do that, we used our wet bench *Singleton* data as a training set for a machine learning model designed to predict 5p versus 3p strand usage in miRNA hairpins. We hypothesized that by training such model we could deconvolve the unique intrinsic features that characterize *Singletons*. We opted for a Random Forest classifier, which we trained using our experimentally validated *Singletons* identified through our HITmiSS assay. This model incorporates 77 features, systematically organized into five functional categories (dinucleotide composition, first/last nucleotide bias, GC content and entropy, base pairing properties, and general structural features) that together capture critical aspects of miRNA mature duplex structure (**Supplemental Fig. S1–S2, Supplemental Table S3-S4, and Materials and Methods**). The model demonstrated strong performance in distinguishing 5p from 3p usage across a set of known miRNAs (**Supplemental Fig. S2, and Materials and Methods**).

***5p* and *3p Singletons* possess distinct structural features**

Our ML model built using our *Singleton* data identified several intrinsic features in these miRNAs. To generalize this model across all *C. elegans* miRNAs, and study how *Switchers* can instead bypass these rules and switch miRNA strand preference, we applied our ML model and conducted a comparative feature analysis across the entire

C. elegans miRNAome (374 miRNAs, miRbase).

We extracted and calculated all 77 features used in our ML model for these miRNAs

(**Supplemental Table S4**), keeping our definition of *Singletons* very stringent to

increase the signal (confidence score >80%).

As expected, we identified 54 *5p Singletons*, 52 *3p Singletons,* and 268 *Switchers*

(**Main Fig. 4A-B** and **Supplemental Table S4**). Out of 77 extracted structural and

sequence features, 13 exhibited statistically significant differences between the 5p and

3p strands (p < 0.05, Welch's t-test) (**Main Fig. 4B-E**). The most pronounced difference

was observed in the identity of the first nucleotide of the 3p strand, which in *3p*

Singletons was consistently a uracil (p = 3.98 × 10⁻□), followed by an enrichment of

cytosine in the last position (p = 5.69 × 10⁻□) (**Main Fig. 4C-4E**). Additionally, 3p

strands showed an enrichment for specific motifs, particularly GAG (p = 2.51 × 10⁻□)

(**Main Fig. 4E**). We also observed compositional differences: 3p strands displayed

increased levels of cytosine at the first position (not shown) and tended to be slightly

shorter than their 5p counterparts (p = 8.90 × 10⁻³) (**Main Fig. 4D-E**).

Surprisingly, our ML model did not cluster all the *Switchers* around a confidence score

of ~50%. Instead, we see a multi-modal distribution with several periodical clusters

(**Main Fig. 4A)**. This suggests that across this category, structural and thermodynamic

features alone cannot fully explain strand usage, and extrinsic factors could move the

needle between 5p or 3p choice in these miRNAs, perhaps depending on tissue or

developmental context.

We next wanted to study the intrinsic features identified by our ML model. For these

experiments we used *Singletons* to test if altering these rules could convert them into

Switchers. In *let-7*, modifying nucleotide pairing at positions 11–12 (UP11-12) or

changing the first base (5'U→C) reduced its predicted *5p Singleton* confidence by up to

17%, consistent with our model (**Main Figure 3C-D**). Similar results were observed in

miR-52, *miR-75*, *lin-4*, and *miR-35*: introducing bulges near the center disrupted strand

preference, while removing them reinforced it (**Supplemental Fig. S6**), highlighting the

central region's role in strand selection.

The first nucleotide's identity and pairing also influenced strand prediction, though with

variable impact across miRNAs (**Supplemental Fig. S6**). Mutating this position altered

strand usage by less than 10% in *5p Single*tons but up to 30% in *3p Singletons*,

mirroring HiTmiSS results and further supporting our strand selection model.

Strand usage bias differs by species

Our ML model identified several key intrinsic features that clearly separate

Singletons from *Switchers* in *C. elegans*. Since many miRNA sequences are conserved

across species, we hypothesized that the features identified by our ML model were

selectively maintained throughout evolutionary time. To do that, we applied the same

model to the complete human miRNAome (**Main Fig. 5**) (**Supplemental Table S5**).

Strand usage in *C. elegans* was nearly balanced (51.3% 5p vs. 48.7% 3p), suggesting

flexible regulatory potential. In contrast, human miRNAs showed a strong 5p bias

(75.9% 5p), indicating a shift toward fixed strand preference (**Main Fig. 5A**).

We identified 13 significantly different features between species ($p < 0.05$, Welch's t-

test), with striking enrichment of guanine content ($p = 5.62 \times 10^{-2}$□) and GG

dinucleotides (p = 1.94 × 10⁻²▯) on human 5p strands (**Main Fig. 5B**). Humans also

showed higher CCU motif presence and cytosine content in 3p regions, including di_CC

(p < 5 × 10⁻²▯), suggesting evolved sequence or processing preferences. These trends

are evident in conserved miRNAs such as *let-7* (**Main Fig. 4C**). We also extended

predictions to other species (mouse, rat, frog, zebrafish, fly) to explore evolutionary

patterns (**Supplemental Table S5**).

Our ML model can accurately predict human miRNA strand selection

Our *C. elegans* ML model was also capable to identify sequence elements and

structural features in human miRNAs. To validate these results, we compared its

predictions to expression data from the Human miRNA Tissue Atlas across 65 healthy

tissues [21] (**Main Fig. 6**). For the top 10 most expressed miRNAs, the model reached

90% accuracy, which is higher than in *C. elegans*. As more miRNAs were included,

accuracy declined, reaching 59% across the full dataset, yet remained statistically

significant (p ≈ 7.6 × 10⁻▯) (**Main Fig. 6A**).

To reduce noise from low-expression miRNAs, we focused on the 50 most abundant

duplexes (**Main Fig. 6B**). The model correctly predicted the dominant strand in 33

cases, yielding 66% accuracy (p ≈ 0.0164). Most errors involved 3p-dominant miRNAs

misclassified as 5p, reflecting a model bias toward 5p predictions. These

misclassifications often occurred when 5p and 3p strands had similar expression levels,

suggesting either model uncertainty or that these miRNAs may function as *Switchers* in

humans (**Supplemental Table S6**).

Notably, two misclassified miRNAs, *miR-182* and *miR-378a*, are known *Switchers in*

vivo [34,35]. In other cases, mispredictions occurred when the model's confidence scores

for 5p and 3p usage were nearly equal (±10%). For instance, *miR-199a* and *miR-199b*,

involved in hypoxia and cardiac regulation [36], were predicted as 5p dominant despite

showing 3p expression and a narrow confidence margin (~7%). Similarly, *miR-124*,

critical for neurogenesis, was misclassified with a 6% confidence gap, despite clear 3p

dominance [37].

Still the majority of miRNAs were accurately predicted across diverse human

tissues, including key regulators with strong strand bias, indicating that our *C. elegans*-

trained model generalizes well to vertebrates (**Main Fig. 6C and Supplemental Table**

S6). For example, *miR-21*, which is a highly expressed oncomiR, was correctly

predicted as 5p-dominant, with over 50-fold 5p vs. 3p expression [38]. Similarly, *let-7a*

showed a 248-fold 5p preference and was correctly classified [39]. Other highly expressed

miRNAs such *miR-1* and miR-*141* were also correctly predicted as 3p dominant.

Overall, these findings demonstrate that the model is capable of reliably predicting

strand usage for miRNAs in healthy human tissues, particularly in cases where strand

bias is strong. Errors were enriched among duplexes with more balanced expression

levels or context-dependent strand usage, highlighting the complexity of predicting

which strand will be used in cases of weak or dynamic regulation of strand selection.

Discussion

In this study, we investigated miRNA strand selection by integrating ~8,000 HiTmiSS

qPCR assays in *C. elegans* with structural modeling and machine learning. This allowed

classification into *5p Singletons*, *3p Singletons*, or *Switchers*, and informed a predictive

model validated against human miRNA expression data.

Nearly half of miRNAs (46%) behave as *Switchers*, altering strand usage across

development, indicating that strand selection is context-dependent rather than fixed

(**Main Fig. 1)**. In contrast, key developmental regulators such as *let-7*, *lin-4*, and *miR-35*

are *Singletons*, likely reflecting a need for consistent strand usage to ensure stable

gene expression. Notably, *Singletons* tend to be more highly expressed (**Main Fig. 1B**),

perhaps due to stronger structural or thermodynamic constraints. In the intestine, tissue-

specific ALG-1 immunoprecipitation revealed a reversal of the developmental trend, with

more *5p* than *3p Singletons* (**Main Fig. 2**). This suggests that strand selection is also

shaped by tissue-specific factors, such as the cellular environment or cofactors. Since

we examined only one somatic tissue, some miRNAs classified as *Switchers* in our

experiments may in fact be tissue-specific *Singletons*.

A concern was that target mRNAs might stabilize one strand, creating an artificial

strand preference in our HiTmiSS assay. To test this, we overexpressed 3'UTRs with

binding sites for typically low-abundance strands in the intestine. Strand levels remained

unchanged, indicating that HiTmiSS reflects true strand usage, not target-driven

stabilization (**Supplemental Figure S5**).

When comparing the structural composition of each mature duplex, a series of

patterns emerged. *5p Singletons* tend to have the first nucleotide unpaired to its

corresponding strand in the duplex (**Main Fig. 3A**). This difference in pairing repeats in

the middle of the duplex and lowers toward the end (**Main Fig. 3A**). Contrarily, *3p*

Singletons present a near-opposite trend, with the first nucleotide more often paired and

the end portion of the sequence also paired, though the center of the sequence shows

similar unpairing to *5p Singletons*. These results support earlier models linking 5' end

thermodynamic instability to strand selection [42,43], but add positional detail by showing

that asymmetry varies along the duplex. Both *5p* and *3p Singletons* show central

unpairing, likely reflecting a conserved feature for Dicer or Argonaute engagement but

not strand bias. Notably, *5p Singletons* show decreasing pairing from the seed to the 3'

end, while *3p Singletons* display the opposite trend. These opposing patterns suggest

that specific structural cues along the duplex, not just global stability, guide strand

selection, pointing to evolved, position-dependent regulatory signals within miRNA

architecture.

To expand this analysis, we extracted 77 structural and compositional features from

our validated *Singletons* and trained a Random Forest classifier (ML model) to predict

dominant strand usage. When comparing these features (**Main Fig 4**), we found several

strand-specific differences, including a significantly higher occurrence of uracil at the

first nucleotide in 3p strands and enrichment for GAG motifs.

We validated our predictions using *in silico* site-directed structural mutations in

canonical miRNAs such as *let-7*, *lin-4*, and *miR-52*, *miR-75* (**Main Fig. 3C**,

Supplemental Fig. S4). Disrupting features such as central bulges (**Main Fig. 3C,**

"UP11-12") or first nucleotide pairing (**Main Fig. 3C, "1U->C"**), altered predicted strand

preference to some degree, further confirming their role in the selection process.

Surprisingly, the degree to which they impact miRNA strand preference was

approximately equal, but when disrupted in tandem led to an even greater loss of

predicted strand preference (**Main Fig. 3C, "Combined"**, **Supplemental Fig. S6**). In

addition, these findings phenocopy the trends seen in our experimentally validated

Singletons (**Main Fig. 3A**). Taken together, these results suggest that the structural and

thermodynamic features identified in our wet bench validated *5p* and *3p Singletons* are

indeed critical regulators of strand selection in nematodes.

Expanding this framework, we then applied our ML model to predict strand usage

across the human miRNA dataset (**Main Fig. 5A**). We were particularly curious to test if

the structural and thermodynamic features and motifs identified in *C. elegans* were

conserved in vertebrates.

Compared to the nearly equal distribution of 5p and 3p usage in *C. elegans*, using

the same cutoff of 80%, the vast majority of human miRNAs are identified as *Switchers*

(831/854 predicted *Switchers*, 97.3%). However, within this class there is a clear bias

toward 5p strands, with over 75% of predictions favoring 5p strand usage (**Main Fig.**

5A). This divergence likely reflects increased specialization in miRNA maturation and

Argonaute loading in vertebrates, potentially driven by evolutionary selection for strand-

specific targeting or stability.

The prevalent 5p bias in human miRNAs may reflect evolutionary pressures rather

than just biochemical preference. As miRNA families expanded during vertebrate,

particularly mammalian, evolution, selective forces likely favored stable strand usage to

ensure consistent and non-redundant regulation [2]. This trend would promote *Singletons*

over *Switchers*, reducing regulatory variability. The fixation of 5p strand usage may thus

represent stabilizing selection for predictable miRNA function, with structural features

reinforcing 5p bias evolving alongside increasing organismal complexity.

In addition to this imbalance in *Singletons* vs *Switchers*, the nucleotide composition

of their 5p and 3p strands is also different, with a significant enrichment of guanine

content in human 5p strands and cytosine-rich motifs in 3p strands (**Main Fig. 5B**).

These compositional biases are consistent with known differences in Argonaute protein

specificity and may reflect lineage-specific constraints on strand functionality [44].

Importantly, these cross-species trends persisted in known miRNAs such as *let-7*,

which was consistently predicted and validated as a *5p Singletons* both in *C. elegans*

and in human, although at different confidence levels. This supports the robustness of

our predictive ML model and highlights conserved features of miRNA processing, even

in the presence of divergent strand preferences.

A surprising trend emerges when we compare our ML model prediction in human

miRNAs to the human miRNA Tissue Atlas. Our ML model developed in *C. elegans* is

capable of accurately predicting miRNA strand usage in humans (**Main Fig. 6**). This is

unexpected, considering the vast evolutionary gap (~600 million years) between these

two systems, which share less than 35% of nucleotide conservation in genes [45].

Despite this, the features we have identified seem to be evolutionarily conserved.

This suggests that miRNA strand selection is, at its core, a fundamental cellular

process that has evolved very little over time, even as miRNA regulatory networks

expanded in complexity with the increased cellular specialization. This becomes more

evident when we overlay miRNA strand usage in human tissues of different origins.

Here, the same miRNA always is predicted to use the same strand in other somatic

tissues (**Main Fig. 6C**). This invariance suggests that strand selection is most likely

encoded by intrinsic properties of the hairpin itself, rather than by external, context-

dependent cues. Such robustness implies that strand choice is not merely a byproduct

of downstream regulatory needs, but a core mechanistic feature established early in

miRNA biogenesis. We propose that during vertebrate evolution, particularly with the

rise of tissue specialization, miRNA strand usage remained anchored to conserved

structural determinants even as the broader miRNA regulatory network expanded and

diversified. This mechanistic rigidity may stabilize gene regulatory outputs in the face of

cellular complexity.

While our model shows that strand selection for many miRNAs can be accurately

predicted based on intrinsic properties of the hairpin structure, the prevalence of

Switchers strongly suggests the involvement of additional, context-dependent regulatory

mechanisms. These may include tissue-specific cofactors, differential Argonaute

availability, or post-transcriptional modifications influencing duplex stability or Argonaute

loading preferences.

In conclusion, our findings illustrate the complexity of miRNA strand selection and

reveal that this process is not governed solely by thermodynamic asymmetry. Instead, a

combination of structural motifs, sequence biases, and cellular context dictate strand

usage. Our work establishes a predictive framework for understanding strand selection

but also opens new avenues for studying the evolution of miRNA regulatory logic across

metazoans.

Data Availability

The raw data produced by the HiTmiSS assay are included in the **Supplemental Table**

S2. The ML model is included as Supplemental Data and is also available upon request.

Funding

This work was funded by the National Institutes of Health 5R01GM118796 awarded to

M.M.

Acknowledgments

We thank Dr. Murugan for assistance in HiTmiSS assay optimization.

Author Contributions

D.M. and M.M. designed the experiments. D.M. performed the experiments. D.M. and

M.M. prepared the ML model and performed the bioinformatics analysis. H.L.

supervised the development of the ML model. D.M., A.E. and H.H. performed the

HiTmiSS assay experiments. D.M. and M.M. prepared the main and supplemental

figures and supplemental tables. D.M. and M.M. wrote the manuscript. All authors have

read and approved the manuscript.

Declaration of Interests

The authors have no conflicts of interest.

Figure titles and legends

Main Figure 1: HiTmiSS Assay and Analysis on Mixed Tissue. **(A)** HiTmiSS protocol for RNA extraction, extension, reverse transcription, and quantification in high throughput. Steps include (i) *C. elegans* growth and homogenization, (2) RNA extraction and isolation, (3) RNA polyuridinylation, extension, and reverse transcription (details shown in ii ; orange: mRNAs, green: polyU, purple: SL-polyA reverse universal primer), (4) addition of 5p or 3p miRNA-strand-specific forward primers, (5) loading into 384-well plates, and (6) quantification. **(B)** Heatmap of miRNA expression (n=3) in mixed tissue clustered by quantity, strand use, and developmental stage. Stages include embryo (E), larval 1 through 4 (L1-L4) and young adult (YA). The heatmap legend represents $\log_{10}$ of expression values for each miRNA strand normalized by spike-in GFP control at each stage from -4 (deep purple) to 4 (bright yellow). Black color indicates instances of data which fall below strict expression threshold (one standard deviation above mean of all expression values at that stage). **(C)** Heatmap of pairing frequency by position for the used (left) and discarded (right) strand of the 20 most highly quantified *5p* and *3p* *Singletons* in our mixed tissue dataset. (i) blue: paired; red: unpaired). (ii) Average pairing frequency, and logo plot of nucleotide frequency at each position displayed below (gold box mark the seed region).

Main Figure 2: Analysis of Intestine Tissue-Specific miRNAs. **(A)** Schematics of the intestine-specific GFP-tagged ALG-1 immunoprecipitation. **(B)** Heatmap of miRNA expression in intestine tissue clustered by strand and developmental stage. The heatmap legend represents $\log_{10}$ of expression values for each miRNA strand

normalized by spike-in GFP control at each stage from -4 (deep purple) to 4 (bright

yellow). Black color indicates instances of data which fall below strict expression

threshold (one standard deviation above mean of all expression values at that stage). E:

embryo; YA: young adults. **(C)** Representative brightfield (left) and GFP (right) images

of selected *C. elegans* strains expressing GFP under the control of promoters for

miRNA known to be highly expressed in the intestine, either during embryo stage (top

two) or young adult stage (bottom two). Arrows point to GFP expression in the intestine

tissues in both embryos and young adult stages.

Main Figure 3: Structural and Thermodynamic Determinants of Strand Selection.

(A) Pairing frequency by position of the used (green) and discarded (red) strand for the

most highly quantified *5p Singletons* and *3p Singletons* in mixed tissue (n=20). Positions

at which pairing frequency most differs are boxed, including the first nucleotide (light

blue), center of the duplex (gold), and end of the strand (white). **(B)** Representative pri-

miRNA hairpins for a *5p Singleton* (*miR-52*) and a *3p Singleton* (*miR-75*) demonstrating

occurrence of trends in pairing as described above. Mature duplex regions highlighted

in pink. **(C)** Changes in *Singleton* predictive confidence for the miRNA *let-7* as a result

of *in silico* site-directed mutations. RNAfold secondary structures of pri-miRNA with

mature 5p strand (blue) and mature 3p strand (red) from left to right: wildtype (WT),

unpaired nucleotides at positions 11 and 12 via GG→UU substitution (UP11-12), first

nucleotide U→C substitution (1U → C), combination of both prior mutations

(Combined). Table summarizes the predicted strand preference by our ML model for

each mutation. **(D)** Bar chart displaying probability as described above.

Main Figure 4: Analysis of Mature miRNA Duplex Features in *C. elegans*. (A)

Frequency distribution of *5p Singleton* predicted confidence by our ML model across all

miRNAs in *C. elegans*. *3p Singletons* (yellow) are defined as miRNAs falling below a

predicted 5p confidence of 20%; *Switchers* are defined as miRNAs with a predicted 5p

confidence between 20% and 80%; *5p Singletons* (red) are defined as miRNAs with a

predicted 5p confidence of 80% or greater. **(B)** Predicted categorization of all *C.*

elegans miRNAs by our ML model as per the above threshold. **(C)** Predicted

categorization of *C. elegans* miRNAs for which uracil is the first nucleotide of the 3p

strand. **(D)** Length difference (5p strand - 3p strand) for each category of *C. elegans*

miRNAs. *5p Singletons* display significantly ($p < 0.05$) longer 5p strands than 3p strands

when compared either to *3p Singletons* or to *Switchers*. **(E)** Heatmap of 32 features

including GC content, miRNA length, hydrogen bonding, nucleotide pairing, strand

stability, and MFOLD-derived parameters enrichment in *5p* and *3p Singletons*. Heatmap

values represent mean values of each feature across the miRNA categories. For binary

features (i.e. 5p_has_AAU, which is either 0 or 1) the number represents the proportion

(e.g. 0.3 = 30% of miRNAs in that category have an AAU motif in the 5p strand). For

continuous features (i.e. gc_content, pairing_frac_5p), the number is the average value

across all miRNAs in that category. For nucleotide composition features (i.e. 5p_A,

3p_U), the number reflects the fraction of that nucleotide within the strand. Features

clustered by average linkage.

Main Figure 5: Comparative Analysis of Mature miRNA Duplex Features between

C. elegans **and** *H. sapiens*. **(A)** Frequency distribution of predicted *5p Singleton* confidence across all miRNAs in *H. sapiens*. *3p Singletons* (yellow) are defined as miRNAs falling below a predicted 5p confidence of 20% or below; *Switchers* are defined as miRNAs with a predicted 5p confidence between 20% and 80%; *5p Singletons* (red) are defined as miRNAs with a predicted 5p confidence of 80% or greater. **(B)** Four most significant ($p < 0.05$) structural differences between *H. sapiens* (left) and *C. elegans* (right) mature miRNA duplexes. From top left to bottom right: mean occurrence of guanine in the 5p strand (5p_G), mean occurrence of guanine-guanine dinucleotide in either mature strand (di_GG), mean occurrence of cytosine-cytosine dinucleotide in either mature strand (di_CC), binary classifier for presence of cytosine-cytosine-uracil trinucleotide in the 3p strand (3p_has_CCU). **(C)** Alignment of *C. elegans* (*cel*) *let-7* and five *H. sapiens* (*hsa*) *let-7* homologs shows differences in CCU occurrence within the 3p strand (red boxes). **(D)** Heatmap of differences in mean value for each of the 77 model features between each category of miRNA in *C. elegans* (c) and *H. sapiens* (h). For binary features (i.e. 3p_has_CCU, which is either 0 or 1) the number represents the proportion (e.g. 0.3 = 30% of miRNAs in that category have an CCU motif in the 3p strand). For continuous features (i.e. gc_content, au_content), the number is the average value across all miRNAs in that category. For nucleotide composition features (i.e. 5p_A, 5p_U), the number reflects the fraction of that nucleotide within the strand. Features clustered by average linkage.

Main Figure 6: Model Accuracy for the Human miRNA Tissue Atlas Dataset. **(A)**

Model accuracy expressed as a percentage of correct predictions for four different

subsets of *H. sapiens* miRNAs (total 854 mature duplexes). From left to right, ranked by

total expression of mature 5p and 3p strand across all tissue types: predicted accuracy

among top 10 (9 correct, 1 incorrect), top 50 (33 correct, 17 incorrect), top 100 (66

correct, 34 incorrect), and all (502 correct, 352 incorrect) miRNAs in this dataset. **(B)**

Confusion matrix of the strand usage prediction model among the top 50 most highly

expressed *H. sapiens* miRNAs. Summarizes differences in 5p and 3p strand usage

between experimentally validated (miRNA Atlas) and model-derived (ML Model) data.

(C) Heatmap of prediction accuracy, either correct (red) or incorrect (blue) for the top 50

most highly expressed *H. sapiens* miRNAs across 65 *H. sapiens* tissue types and

organs. Data are clustered using average linkage.

Supplemental Information titles and legends

Supplemental Table S1: **Primers used in this study.**

Supplemental Table S2: **HiTmiSS assay results.**

Supplemental Table S3: **Features used in our ML model and their weights.**

Supplemental Table S4: **ML prediction in *C. elegans*.**

Supplemental Table S5: **ML prediction in vertebrates.**

Supplemental Table S6: **Comparison of ML model with miRNA tissue Atlas.**

References

1   Bartel, D. P. Metazoan MicroRNAs. *Cell* **173**, 20-51 (2018).

https://doi.org:10.1016/j.cell.2018.03.006

2   Wolter, J. M. *et al.* Evolutionary patterns of metazoan microRNAs reveal targeting

principles in the let-7 and miR-10 families. *Genome Res* **27**, 53-63 (2017).

https://doi.org:10.1101/gr.209361.116

3   Pare, J. M. *et al.* Hsp90 regulates the function of argonaute 2 and its recruitment to stress

granules and P-bodies. *Mol Biol Cell* **20**, 3273-3284 (2009).

https://doi.org:10.1091/mbc.e09-01-0082

4   Nakanishi, K. Anatomy of RISC: how do small RNAs and chaperones activate Argonaute

proteins? *Wiley Interdiscip Rev RNA* **7**, 637-660 (2016).

https://doi.org:10.1002/wrna.1356

5   Meijer, H. A., Smith, E. M. & Bushell, M. Regulation of miRNA strand selection: follow

the leader? *Biochem Soc Trans* **42**, 1135-1140 (2014).

https://doi.org:10.1042/BST20140142

6   Medley, J. C., Panzade, G. & Zinovyeva, A. Y. microRNA strand selection: Unwinding

the rules. *Wiley Interdiscip Rev RNA* **12**, e1627 (2021). https://doi.org:10.1002/wrna.1627

7   Frank, F., Sonenberg, N. & Nagar, B. Structural basis for 5'-nucleotide base-specific

recognition of guide RNA by human AGO2. *Nature* **465**, 818-822 (2010).

https://doi.org:10.1038/nature09039

8   Melendez, A. & Levined, B. in *WormBook*   (ed The C. elegans Research Community)

(WormBook, August 24, 2009).

9   Zinovyeva, A. Y., Veksler-Lublinsky, I., Vashisht, A. A., Wohlschlegel, J. A. & Ambros,

V. R. Caenorhabditis elegans ALG-1 antimorphic mutations uncover functions for

937   Argonaute in microRNA guide strand selection and passenger strand disposal. *Proc Natl*

938   *Acad Sci U S A* **112**, E5271-5280 (2015). https://doi.org:10.1073/pnas.1506576112

939 10 Caceres, J. F., Stamm, S., Helfman, D. M. & Krainer, A. R. Regulation of alternative

940   splicing in vivo by overexpression of antagonistic splicing factors. *Science* **265**, 1706-

941   1709 (1994). https://doi.org:10.1126/science.8085156

942 11 Mangone, M., Macmenamin, P., Zegar, C., Piano, F. & Gunsalus, K. C. UTRome.org: a

943   platform for 3'UTR biology in C. elegans. *Nucleic Acids Res* **36**, D57-62 (2008).

944   https://doi.org:10.1093/nar/gkm946

945 12 Mangone, M. *et al.* The landscape of C. elegans 3'UTRs. *Science* **329**, 432-435 (2010).

946   https://doi.org:10.1126/science.1191244

947 13 Steber, H. S., Gallante, C., O'Brien, S., Chiu, P. L. & Mangone, M. The C. elegans 3'

948   UTRome v2 resource for studying mRNA cleavage and polyadenylation, 3'-UTR

949   biology, and miRNA targeting. *Genome Res* **29**, 2104-2116 (2019).

950   https://doi.org:10.1101/gr.254839.119

951 14 Murari, E., Meadows, D., Cuda, N. & Mangone, M. A comprehensive analysis of 3'UTRs

952   in Caenorhabditis elegans. *Nucleic Acids Res* (2024). https://doi.org:10.1093/nar/gkae543

953 15 Sulston, J. E. Neuronal cell lineages in the nematode Caenorhabditis elegans. *Cold Spring*

954   *Harb Symp Quant Biol* **48 Pt 2**, 443-452 (1983).

955   https://doi.org:10.1101/sqb.1983.048.01.049

956 16 Frokjaer-Jensen, C. *et al.* Single-copy insertion of transgenes in Caenorhabditis elegans.

957   *Nat Genet* **40**, 1375-1383 (2008). https://doi.org:10.1038/ng.248

958 17 Kotagama, K., Schorr, A. L., Steber, H. S. & Mangone, M. ALG-1 Influences Accurate

959   mRNA Splicing Patterns in the Caenorhabditis elegans Intestine and Body Muscle

Tissues by Modulating Splicing Factor Activities. *Genetics* **212**, 931-951 (2019). https://doi.org:10.1534/genetics.119.302223

Mei, Q. *et al.* A facile and specific assay for quantifying microRNA by an optimized RT-qPCR approach. *PloS one* **7**, e46890 (2012). https://doi.org:10.1371/journal.pone.0046890

Livak, K. J. & Schmittgen, T. D. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* **25**, 402-408 (2001). https://doi.org:10.1006/meth.2001.1262

Lorenz, R. *et al.* ViennaRNA Package 2.0. *Algorithms Mol Biol* **6**, 26 (2011). https://doi.org:10.1186/1748-7188-6-26

Rishik, S., Hirsch, P., Grandke, F., Fehlmann, T. & Keller, A. miRNATissueAtlas 2025: an update to the uniformly processed and annotated human and mouse non-coding RNA tissue atlas. *Nucleic Acids Res* **53**, D129-D137 (2025). https://doi.org:10.1093/nar/gkae1036

Brosnan, C. A., Palmer, A. J. & Zuryn, S. Cell-type-specific profiling of loaded miRNAs from Caenorhabditis elegans reveals spatial and temporal flexibility in Argonaute loading. *Nat Commun* **12**, 2194 (2021). https://doi.org:10.1038/s41467-021-22503-7

Shaw, W. R., Armisen, J., Lehrbach, N. J. & Miska, E. A. The conserved miR-51 microRNA family is redundantly required for embryonic development and pharynx attachment in Caenorhabditis elegans. *Genetics* **185**, 897-905 (2010). https://doi.org:10.1534/genetics.110.117515

24 Alvarez-Saavedra, E. & Horvitz, H. R. Many families of C. elegans microRNAs are not

essential for development or viability. *Curr Biol* **20**, 367-373 (2010).

https://doi.org:10.1016/j.cub.2009.12.051

25 Garcia-Segura, L. *et al.* High-Throughput Profiling of Caenorhabditis elegans Starvation-

Responsive microRNAs. *PloS one* **10**, e0142262 (2015).

https://doi.org:10.1371/journal.pone.0142262

26 Garcia, S. M., Tabach, Y., Lourenço, G. F., Armakola, M. & Ruvkun, G. Identification of

genes in toxicity pathways of trinucleotide-repeat RNA in C. elegans. *Nature structural*

& molecular biology **21**, 712-720 (2014).

27 Kato, M., Kashem, M. A. & Cheng, C. An intestinal microRNA modulates the

homeostatic adaptation to chronic oxidative stress in C. elegans. *Aging (Albany NY)* **8**,

1979-2005 (2016). https://doi.org:10.18632/aging.101029

28 Kagias, K., Podolska, A. & Pocock, R. Reliable reference miRNAs for quantitative gene

expression analysis of stress responses in Caenorhabditis elegans. *BMC Genomics* **15**,

222 (2014). https://doi.org:10.1186/1471-2164-15-222

29 Blazie, S. M. *et al.* Alternative Polyadenylation Directs Tissue-Specific miRNA

Targeting in Caenorhabditis elegans Somatic Tissues. *Genetics* **206**, 757-774 (2017).

https://doi.org:10.1534/genetics.116.196774

30 Haenni, S. *et al.* Analysis of C. elegans intestinal gene expression and polyadenylation by

fluorescence-activated nuclei sorting and 3'-end-seq. *Nucleic Acids Res* **40**, 6304-6318

(2012). https://doi.org:10.1093/nar/gks282

31   Pandit, A., Jain, V., Kumar, N. & Mukhopadhyay, A. PHA-4/FOXA-regulated

microRNA feed forward loops during Caenorhabditis elegans dietary restriction. *Aging*

(Albany NY) **6**, 835-855 (2014). https://doi.org:10.18632/aging.100697

32   Brunquell, J., Snyder, A., Cheng, F. & Westerheide, S. D. HSF-1 is a regulator of

miRNA expression in Caenorhabditis elegans. *PloS one* **12**, e0183445 (2017).

https://doi.org:10.1371/journal.pone.0183445

33   Nakanishi, K. When Argonaute takes out the ribonuclease sword. *J Biol Chem* **300**,

105499 (2024). https://doi.org:10.1016/j.jbc.2023.105499

34   Krishnan, K. *et al.* MicroRNA-182-5p targets a network of genes involved in DNA

repair. *RNA* **19**, 230-242 (2013). https://doi.org:10.1261/rna.034926.112

35   Krist, B., Florczyk, U., Pietraszek-Gremplewicz, K., Jozkowicz, A. & Dulak, J. The Role

of miR-378a in Metabolism, Angiogenesis, and Muscle Biology. *Int J Endocrinol* **2015**,

281756 (2015). https://doi.org:10.1155/2015/281756

36   Rane, S. *et al.* Downregulation of miR-199a derepresses hypoxia-inducible factor-1alpha

and Sirtuin 1 and recapitulates hypoxia preconditioning in cardiac myocytes. *Circ Res*

104, 879-886 (2009). https://doi.org:10.1161/CIRCRESAHA.108.193102

37   Makeyev, E. V., Zhang, J., Carrasco, M. A. & Maniatis, T. The MicroRNA miR-124

promotes neuronal differentiation by triggering brain-specific alternative pre-mRNA

splicing. *Mol Cell* **27**, 435-448 (2007). https://doi.org:10.1016/j.molcel.2007.07.015

38   Rhim, J., Baek, W., Seo, Y. & Kim, J. H. From Molecular Mechanisms to Therapeutics:

Understanding MicroRNA-21 in Cancer. *Cells* **11** (2022).

https://doi.org:10.3390/cells11182791

39   Brueckner, B. *et al.* The human let-7a-3 locus contains an epigenetically regulated

microRNA gene with oncogenic function. *Cancer Res* **67**, 1419-1423 (2007).

https://doi.org:10.1158/0008-5472.CAN-06-4074

40   Kim, C., Ye, Z., Weyand, C. M. & Goronzy, J. J. miR-181a-regulated pathways in T-cell

differentiation and aging. *Immun Ageing* **18**, 28 (2021). https://doi.org:10.1186/s12979-

021-00240-1

41   Pakravan, K. *et al.* MicroRNA-100 shuttled by mesenchymal stem cell-derived exosomes

suppresses in vitro angiogenesis through modulating the mTOR/HIF-1alpha/VEGF

signaling axis in breast cancer cells. *Cell Oncol (Dordr)* **40**, 457-470 (2017).

https://doi.org:10.1007/s13402-017-0335-7

42   Khvorova, A., Reynolds, A. & Jayasena, S. D. Functional siRNAs and miRNAs exhibit

strand bias. *Cell* **115**, 209-216 (2003). https://doi.org:10.1016/s0092-8674(03)00801-8

43   Schwarz, D. S. *et al.* Asymmetry in the assembly of the RNAi enzyme complex. *Cell*

115, 199-208 (2003). https://doi.org:10.1016/s0092-8674(03)00759-1

44   Kotagama, K. *et al.* Catalytic residues of microRNA Argonautes play a modest role in

microRNA star strand destabilization in C. elegans. *Nucleic Acids Res* **52**, 4985-5001

(2024). https://doi.org:10.1093/nar/gkae170

45   Stein, L. *et al.* WormBase: network access to the genome and biology of Caenorhabditis

elegans (vol 29, pg 82, 2001). *Nucleic Acids Research* **29**, 1012 (2001).

1043

Main Figure 1

Main Figure 2

A

5p Singletons

strand
- used
- discarded

pairing frequency

position

3p Singletons

pairing frequency

position

C

WT   UP11-12   1U → C   Combined

Variant	5p	3p
WT	0.85	0.15
UP11-12	0.71	0.29
5'U → C	0.71	0.29
Combined	0.68	0.32

B

miR-52

miR-75

D

probability (%)

5p 3p

WT   UP11-12   5' U → C   Unpaired 5'

A Model Accuracy by Expression Rank

B Confusion Matrix Top 50 – miRNA Atlas vs ML Model

C Model Accuracy – Top 50 miRNAs across 65 *H. sapiens* Organ Systems

correct prediction
incorrect prediction