Non-Destructive Larval Genotyping of *Danio rerio* for Mitochondrial and Nuclear DNA Genetics

Kyler S. Mitra[1,2], Shannon R. Holmberg[3], Mireya Mota[2,4], Ankit Sabharwal[2,4], and Stephen C. Ekker[1,2,4,#]

[1]Department of Molecular Biosciences, The University of Texas at Austin, Texas, USA;

[2]Center for Rare Disease, Dell Medical School, The University of Texas at Austin, Texas, USA;

[3]Department of Biochemistry and Molecular Biology, Mayo Clinic, Rochester, Minnesota, USA;

[4]Department of Pediatrics, Dell Pediatric Research Institute, Dell Medical School, The University of Texas at Austin, Texas, USA

[#]Corresponding author: Stephen C. Ekker, Ph.D., stephen.ekker@austin.utexas.edu

Keywords: Zebrafish, Mitochondrial DNA, Nuclear DNA Non-destructive genotyping

Abstract

The rapid advancement of nuclear and mitochondrial genomic editing tools has created an urgent need for efficient, non-lethal larval genotyping methods in zebrafish (*Danio rerio*) research. This study optimizes and validates a non-destructive proteinase K digestion method for mitochondrial and nuclear DNA genotyping while characterizing its impact on larval survival and gene expression. Using optimized protocol parameters, we demonstrate successful amplification of different mitochondrial and nuclear genetic loci with consistently high sensitivity. Molecular validation through PCR, restriction fragment length polymorphism analysis, and Sanger sequencing confirmed the specificity and reliability of the extracted DNA. The method successfully detected C-to-T base edits in the *mt-tl1* gene introduced using the FusX TALE Base editor system, demonstrating its applicability to gene editing studies. Both 48-well and optimized 96-well formats were used, enabling this approach to be deployed at scale. This optimized method enables researchers to correlate genotypes with phenotypes in longitudinal studies while maintaining specimen viability particularly valuable for investigating early-onset mitochondrial diseases and utilizes standard laboratory equipment and reagents, facilitating widespread adoption in zebrafish research while adhering to ethical principles in reducing animal mortality.

Introduction

Zebrafish (*Danio rerio*) are the most widely used aquatic model system in scientific research, transcending their initial role in developmental biology to become instrumental in unraveling the genetic and biochemical intricacies of various cellular processes. This small vertebrate boasts an impressive array of advantages as a model organism, including rapid reproductive cycles, high offspring yield, external embryonic development, transparent embryos, swift organ formation, and straightforward husbandry requirements.[1–5] The versatility of zebrafish has led to groundbreaking insights across a spectrum of biological processes and human diseases. From hematological and cardiovascular disorders to neurological conditions and cancer, zebrafish models have significantly contributed to our understanding of these complex systems.[6–8] Zebrafish shine particularly bright in the realm of mitochondrial research. Their mitochondrial genome shares remarkable similarities with humans, with 100% gene synteny and approximately 65% sequence identity at the nucleotide level. This genetic kinship extends to codon usage, strand-specific nucleotide bias, and gene order, making zebrafish an ideal vertebrate model for studying human mitochondrial disorders.[9–11] Since mitochondrial DNA (mtDNA) is extra-nuclearly segregated, it has different requirements for genotype to phenotype correlation. The complex genetics underlying mtDNA disorders, particularly their variable penetrance and diverse clinical presentations, can be attributed to the presence of either homoplasmy or heteroplasmy of mtDNA type within cells. The severity and progression of mitochondrial diseases are directly influenced by the ratio of mutated to normal (wild type) mtDNA within a cell.

Recent years have witnessed significant advancement in the mitochondrial DNA (mtDNA) editing toolbox such as DddA-derived cytosine base editors (DdCBEs) and TALE-linked deaminase (TALEDs) to engineer precise edits across different species enabling to model pathogenic mutations.[12–17] Notably, DddA-derived cytosine base editors (DdCBEs) have achieved editing efficiencies at targeted mtDNA loci, enabling accurate modeling of pathogenic mutations. Additionally, engineered variants such as DddA6 and DddA11 have expanded the editing scope by targeting previously inaccessible sequence contexts, while monomeric DdCBE (mDdCBE) constructs facilitate broader applicability and simplified delivery.[13,18] These developments collectively enhance the utility of zebrafish as a robust vertebrate model for mitochondrial genetic research. The advent of these mitochondrial genomic editing tools underscores the critical need for rapid and efficient larval genotyping in zebrafish research. Traditional approaches to live larval genotyping have relied on microsurgical fin biopsy performed in water droplets,[19] while alternative methods involve the complete lysis of embryos for cohort genotype determination[20] or an automated genotyping device.[21] However, these conventional techniques present significant limitations: fin microsurgery is labor-intensive and frequently lethal, while embryonic lysis precludes the survival of genotyped larvae for subsequent analysis or breeding. Non-destructive enzyme genotyping presents a scalable solution that overcomes these constraints by employing standard laboratory reagents and materials. This

technique facilitates DNA extraction through controlled proteinase K (ProK) digestion of
epithelial cells, yielding genetic material that can be PCR amplified while preserving larval
viability.[22] This protocol enables greater efficiency in husbandry and high-throughput assays,
enabling genotype determination within days of fertilization. Notably, this non-lethal approach
proves particularly valuable for investigating early-onset lethal phenotypes, where rapid
genotyping of living specimens is essential for understanding disease progression.

In this study, we employed and refined a non-destructive genotyping protocol for the
mitochondrial and nuclear genome and characterized the impact of these assays on larvae
through survival and transcriptomics data. During testing and troubleshooting of this non-
destructive assay, we identified both advantages and disadvantages of the proteinase K method.
Additionally, we demonstrated successful amplification of mitochondrially encoded genes
alongside nuclear genes. Finally, modifications to the previously published enzyme protocol
have enabled stronger survival.

Materials and Methods

Zebrafish handling and husbandry

82 All adult zebrafish and embryos were maintained according to the guidelines established by
83 Mayo Clinic Institutional Animal Care and Use Committee (IACUC number: A34513-13-R16)
84 and The University of Texas at Austin Institutional Animal Care and Use Committee (IACUC
85 number: AUP-2023-00200).

Proteinase K-based larval genotyping

Wild-type (WT) Mayo Recessive Free (MRF) zebrafish and mutant transgenic lines, *mt-tl1*[14]
were used to generate larvae for non-destructive genotyping.[22] Three days post-fertilization (dpf)
larvae were rinsed three times in embryo media followed by three washes in tricaine buffer (30
mM Tris-HCl in 0.4% tricaine solution). Using a p200 pipette tip trimmed to accommodate larval
width, individual larvae were aspirated in 40 µL of ProK genotyping buffer (25 µg/mL
Proteinase K (catalog no.: EO0491, Thermo Fisher Scientific, USA) in tricaine buffer) and
placed in 96-well (catalog no.: 3455, Thermo Fisher Scientific, USA) and 48-well plates
(catalog no.: CC7672-7548, CytoOne). For 48-well plates, droplets were dispensed centrally
within each well to minimize contact between larvae and well walls during agitation. Proteinase
K concentration and incubation time can be adjusted based on larval age and condition to
optimize survival rates. The plate was incubated in a heated shaker (catalog no.: VORTEMP 56,
Labnet International, Inc., USA) at 37°C with 500 rpm shaking for 20 minutes. After incubation,
30 µL of the liquid around the fish was carefully pipetted (avoiding larval aspiration) into 15 µL
of lysis buffer (10 mM Tris-HCl, pH 8.0, 50 mM KCl, 0.3% Tween-20, 0.3% NP-40, 1 mM
EDTA) in 0.2 mL strip tubes and pipetted up and down ~10 times. The solution was then
incubated at 98°C for 5 minutes to extract DNA. Following media collection, wells were quickly

replenished with embryo media using a Pasteur pipette, and larvae were transferred to 24-well
plates containing fresh embryo media.

To prevent tail damage during the shaking procedure, plates can be pre-treated with a protein-
rich blocking solution. If extensive tail damage is observed within shaken larvae, plates can be
incubated with 5% non-fat dry milk (NFDM) solution prepared in embryo media for 5 minutes,
followed by three washes with embryo media prior to genotyping (Fig. 1A).

Survival assay

Larvae collected after enzymatic genotyping were transferred into individual wells of a 24-well
plate. Unshaken age-matched zebrafish larvae obtained from the same clutch were used as
controls. Live counts for larvae from each group were recorded at 24 hours post-genotyping, and
morphological defects were examined pre- and post-procedure.

PCR amplification and Restriction Fragment Length Polymorphism for mitochondrial
genetic loci

To investigate the sensitivity of the protocol for mitochondrial DNA genotyping, we prioritized
protein-coding genes from each of the respiratory chain complex, transfer RNA and ribosomal
RNA genes. We targeted six mitochondrial genetic loci with distinct functional roles. Four of
them were protein coding genes: mitochondrially encoded NADH:ubiquinone oxidoreductase
core subunit 1 (*mt-nd1*), mitochondrially encoded cytochrome b (*mt-cytb*), mitochondrially
encoded ATP synthase membrane subunit 6 (*mt-atp6*) and mitochondrially encoded cytochrome c
oxidase subunit 1 (*mt-co1*). Remaining two were non-coding genes: mitochondrially encoded
tRNA-Leu (*mt-tl1*) and mitochondrially encoded 12S rRNA (*mt-rnr1*) (Fig. 1B). PCR
amplification was performed for each of the six genes using their respective primers mentioned
in Supplementary Table 1. PCR consisted of 10 μL of 2X OneTaq master mix with standard
buffer (NEB, USA), 1 μL of 10 μM forward primer, 1 μL of 10 μM reverse primer and
8 μL of template. Samples were run in a thermal cycler with the following PCR conditions: (1)
95°C, 4 min; (2) 95°C, 30 sec; (3) 60°C, 30 sec; (4) 72°C, 30 s; (5) repeat steps 2-4 for
30 cycles ; (6) 72°C, 4 min; and (7) 12°C, hold. PCR amplicons were run on 1% agarose gel
and analyzed.

Five to eight microliters of the PCR amplicon (depending on intensity of the amplicon on the
gel) were used as the substrate for restriction fragment length polymorphism (RFLP) in 20 µL
reaction solution (2 µL of the 10x corresponding buffer and 5 U of restriction enzyme and
remaining volume with nuclease-free water). PCR amplicons for genetic loci *mt-nd1*, *mt-cytb*,
mt-tl1, and *mt-rnr1* were surveyed for restriction digestion by SpeI, AvaI, SacI and HindIII,
respectively. A subset of the PCR amplicons were then submitted to Sanger sequencing at

GENEWIZ (Azenta, USA) to confirm the sequence specificity. Chromatograms files (.ab1) were
then uploaded to the EditR[23] web server to estimate the heteroplasmy in the mtDNA mutants.

PCR amplification of nuclear genetic loci

To investigate the sensitivity of the protocol for nuclear gene genotyping, we prioritized protein-
coding nuclear genes with roles in mitochondrial function and signaling. The three nuclear loci
were: mitochondrial antiviral signaling (*mavs*), cytochrome c oxidase subunit 7B (*cox7b*), and
leucine-rich pentatricopeptide repeat-containing (*lrpprc*). PCR amplification was performed for
each of these genes using their respective primers listed in Supplementary Table 1. PCR
reactions consisted of 10 μL of 2X OneTaq master mix with standard buffer (NEB, USA),
1 μL of 10 μM forward primer, 1 μL of 10 μM reverse primer, and 8 μL of template DNA.
PCR amplification was carried out under the following conditions: (1) 95°C, 4 min; (2) 95°C,
30 s; (3) annealing at 58°C for *cox7b* and *lrpprc*, or 54°C for *mavs,* 30 s; (4) 72°C, 30 s for
mavs and *lrpprc*; and 2 min for *cox7b* (5) repeat steps 2–4 for 34 cycles; (6) 72°C, 4 min; (7)
hold at 12°C. PCR amplicons were run on 1% agarose gel and analyzed. A subset of the PCR
products was submitted to Sanger sequencing at GENEWIZ (Azenta, USA) to confirm sequence
specificity.

RNA extraction and sample preparation

To assess for any differential transcriptional signatures in the proteinase K-shaken larvae to those
unshaken, we performed whole body RNA sequencing for the two groups. Adult zebrafish were
incrossed and non-destructive genotyping was performed on larvae aged 3 dpf. Larvae were
subjected to the protocol for the non-destructive enzymatic proteinase K conditions and were
then allowed to recover in embryo water until 6 dpf. Six larvae were pooled, and RNA isolation
was performed using the Qiagen RNeasy Mini kit as per manufacturer's instructions. An
additional DNAse treatment step was performed before the final wash and elution. RNA
concentration was measured using a spectrophotometer.

RNA sequencing and analyses

RNA library preparations and sequencing reactions were performed at GENEWIZ (Azenta
USA). For transcriptome sequencing, RNA was isolated by pooling six individual larvae from
each condition. RNA quantification was performed using the Qubit 2.0 Fluorometer (Thermo
Fisher Scientific, USA), and RNA integrity was assessed using the Agilent TapeStation (Agilent
Technologies, USA). RNA sequencing libraries were constructed following the NEBNext Ultra
II RNA Library Prep Kit for Illumina protocol (NEB, USA). Briefly, mRNA was enriched using
oligo(dT) beads, fragmented for 15 minutes at 94°C, and then reverse-transcribed into cDNA.
The cDNA was end-repaired, adenylated at the 3' ends, and ligated with universal adapters,
followed by index addition and PCR-based library amplification. The sequencing library was
validated on the Agilent TapeStation (Agilent Technologies, USA) and quantified using both the

Qubit 2.0 Fluorometer and quantitative PCR (KAPA Biosystems, USA). The sequencing libraries were then multiplexed, clustered onto a flow cell, and loaded onto the Illumina HiSeqX instrument according to the manufacturer's instructions. Sequencing was performed using a 2x150 bp Paired-End (PE) configuration. Image analysis and base calling were carried out with HiSeq Control Software (HCS). The raw sequence data (.bcl files) generated from the Illumina HiSeqX platform were converted into fastq files and demultiplexed using Illumina's bcl2fastq 2.20 software with one mismatch allowed for index sequence identification. The raw fastq data was uploaded to the Basepair cloud bioinformatics platform (https://www.basepairtech.com/). Filtered reads were aligned to the zebrafish reference genome (Zv11 version) using STAR. Raw and normalized gene and transcript counts were calculated. Differential expression analysis was performed using DESeq2[24], with a threshold of p-adjusted <0.05 for identifying differentially expressed genes between the genotyped and unshaken control larvae.

Results

Survival rates of enzyme-genotyped zebrafish larvae support high-throughput screening

Non-destructive enzyme genotyping of zebrafish larvae demonstrates high survival rates and compatibility with high-throughput screening. Larvae shaken in 96-well plates without milk pretreatment had a median survival rate of 75%, while 48-well larvae had a median of 94%, compared to 100% in unshaken controls (Fig. 1C). Most larval deaths occurred within 24 hours post-shaking, primarily due to tail fin damage, with no observed differences in survival between undamaged shaken and unshaken fish after this period. Key survival-enhancing steps included rapidly replacing embryo water after DNA collection and promptly removing larvae from the heated shaker. These parameters can be further adapted to accommodate different larval ages and sensitive phenotypes. Importantly, no morphological changes were observed in surviving genotyped larvae compared to unshaken controls, suggesting the method's minimal physiological impact.

Mitigating characteristic tail fin damage among ProK-genotyped larvae via 48-well plates and milk solution pretreatment

Our initial attempts using 96-well plates for ProK genotyping resulted in a subset of larvae exhibiting fatal damage, including partial to complete tail fin tearing. We attributed this damage pattern to tail fin adhesion to the well walls during heated shaking, exacerbated by ProK-induced weakening of fin structural integrity. We determined this damage pattern likely resulted from tail fin adhesion to well walls during heated shaking, exacerbated by ProK-induced weakening of fin structural integrity. Two effective mitigation strategies were thus developed. First, utilization of 48-well plates enabled placement of larvae in centered droplets without wall contact, significantly increasing survival and reducing tail fin damage. However, the efficacy of this approach may vary depending on the strength of the hydrophobic coating applied by different manufacturers to maintain protective droplet formation. Alternatively, pretreating 96-well plates

210 with 5% non-fat dry milk (NFDM) solution created a protective proteinaceous film that reduced
211 larval tail fin adhesion during shaking (Fig. 1A). We observed decreased tail fin damage and
212 improved survival rates among genotyped larvae in the 96-well plate treated with 5% NFDM in
213 embryo media (Supplementary Fig. 1).

High sensitivity of amplification across diverse mitochondrial genetic loci

Non-destructive enzyme genotyping demonstrates high amplification sensitivity across diverse
mitochondrial genes encoding critical cellular machinery. Amplification sensitivity was assessed
for *mt-nd1*, *mt-cytb*, *mt-rnr1*, and *mt-tl1* by calculating the percentage of successfully amplified
samples across these genes. For each gene-well combination, we conducted three experimental
trials, with each trial containing 20 larvae. Thus, we analyzed a total of 60 larvae for each gene-
well combination, resulting in 120 total samples per gene across both formats. Consistently high
amplification rates were observed across all four genes in both formats. All experimental sets
exhibited over 80% sensitivity (Fig. 1D), demonstrating robust performance of this non-
destructive genotyping method across diverse mitochondrial genes.

Specificity of mitochondrial DNA amplification verified by PCR and RFLP analysis

We validated the specificity of our non-destructive enzyme genotyping method through
comprehensive molecular analysis of four mitochondrial genes. PCR amplification and
restriction fragment length polymorphism (RFLP) analysis were performed on wild-type
zebrafish larval DNA samples from 96-well and 48-well plates, with all samples referenced
against Quick-Load Purple 50 bp DNA Ladder (NEB, USA). For *mt-nd1*, we observed a 350 bp
PCR amplicon with SpeI restriction digest yielding 236 bp and 114 bp fragments (Fig. 2A, B).
The *mt-cytb* gene generated a 460 bp PCR amplicon with AvaI restriction digest yielding 317 bp
and 143 bp fragments (Fig. 2D, E). Amplification of *mt-rnr1* showed a 395 bp PCR amplicon
with HindIII restriction digest producing 253 bp and 142 bp fragments (Fig. 2G, H). The *mt-tl1*
gene produced a 363 bp PCR product (Fig. 3A, B) with SacI restriction digest creating 186 bp
and 177 bp fragments. PCR amplification of *mt-atp6* produced a 369 bp amplicon
(Supplementary Fig. 2A), while *mt-co1* generated a 494 bp PCR product (Supplementary Fig.
2B). Sanger sequencing confirmed the presence of restriction enzyme sites for the first four
genes (Fig. 2C, F, I, 3E), and all samples selected for gel visualization represented the median
amplification range within their respective cohorts.

Genotyping of mtDNA mutant

We demonstrated the utility of our non-destructive genotyping approach for gene editing by
introducing a C-to-T base substitution in zebrafish *mt□tl1* mutants.[14] Three dpf *mt-tl1* zebrafish
mutants were genotyped non-destructively, enabling subsequent longitudinal studies (Fig. 3C).
The engineered C-to-T transition at position m.C3744 (C10 on the forward strand) resulted in the
loss of the SacI restriction site, thereby facilitating the identification of larvae with mtDNA

variations. RFLP analysis demonstrated the loss of the SacI restriction site in mutant larvae, with
digested products of 186 bp and 177 bp, and an undigested product of 363 bp. Wild-type controls
from both 96-well and 48-well cohorts showed only digested products of 186 bp and 177 bp
(Fig. 3D). RFLP analysis was confirmed by Sanger sequencing where sequencing files (.ab1)
were assessed using EditR[23] software to quantify the level of editing in the mutant larvae. Sanger
sequencing confirmed the C-to-T edit, revealing over 40% mtDNA mutant heteroplasmy levels at
the target locus (Fig. 3E).

Amplification across different nuclear genetic loci

To expand the utility of the optimized protocol, we adopted non-destructive genotyping to
amplify different nuclear genetic loci. We observed successful amplification of the nuclear genes
mavs (Fig. 4A, B), *cox7b* (Fig. 4C, D) and *lrpprc* (Fig. 4E, F) by agarose gel electrophoresis
(Fig. 4). PCR amplicons for *mavs* and *lrpprc* were referenced against Quick-Load Purple 50 bp
DNA Ladder (NEB, USA), and for *cox7b*, amplicons were referenced against HyperLadder 1kb
(Meridian Bioscience, USA) Sanger sequencing confirmed that PCR products corresponded to
the intended nuclear gene loci. This demonstrated that these optimizations can be employed for
nuclear DNA genotyping as well.

Transcriptomic signature in non-destructively genotyped zebrafish larvae

To examine the genome-wide transcriptional changes in zebrafish larvae following non-
destructive genotyping, we performed strand-specific paired-end RNA sequencing to identify a
candidate differential expression profile. Sequence data were processed using the DESeq2[24]
pipeline on the Basepair cloud bioinformatics platform (https://www.basepairtech.com/). For
both Proteinase K-shaken larvae and wild-type controls, we generated between 34 and 39 million
reads. Reads below a Phred score of Q30 were discarded, and the remaining high-quality reads
were aligned to the Zv11 zebrafish reference genome, with approximately 78% mapping
uniquely. These uniquely mapped reads were then assembled to quantify transcript levels across
the genome. Upon applying the criteria of a log□ fold change of ≥1 or ≤–1 with an adjusted p-
value < 0.05 in the dataset (Supplementary Fig. 3), zero downregulated genes were observed. We
identified seven annotated genes and six uncharacterized genes to be significantly upregulated in
the 96-well genotyping treatment as compared to unshaken controls (Supplementary Fig. 3 and
Supplementary File 1).

Discussion

In this study, we optimized and validated a non-destructive larval genotyping method for
efficient molecular analysis of zebrafish mitochondrial and nuclear DNA while maintaining high
survival rates. Building upon the non-destructive genotyping protocol originally established by
Zhang et al.,[22] we demonstrated its utility specifically for mitochondrial DNA genetics and
adapted it for high-throughput applications, allowing simultaneous processing of up to 96 larvae.

Our approach addresses a significant challenge in zebrafish research—the ability to collect
sufficient DNA for reliable genotyping from larvae without sacrificing the animal or
compromising its development, which is inherently important for correlating genotype to
phenotype in mitochondrial genetics. The optimized protocol achieves high amplification
sensitivity across diverse mitochondrial and nuclear genes while preserving larval viability.

The protocol provides a balanced approach for mtDNA genotyping that minimizes stress to the
larvae. Notably, the high survival rates observed, particularly in 48-well plates and 96-well with
5% NFDM, demonstrated the method's suitability for high-throughput screening applications.
Our technical improvements, particularly the implementation of 48-well plates and milk
pretreatment steps, significantly mitigated tail fin damage—a principal cause of larval mortality
in our earlier approaches. These modifications addressed the critical issue of tail fins adhering to
well walls during heated shaking, which when combined with ProK-induced weakening of fin
structural integrity, frequently resulted in fatal damage including torn tail fins. The 48-well plates
allowed for centered droplet placement that prevented wall contact, while the milk pretreatment
created a proteinaceous coating that reduced the adhesive properties of the well surfaces,
preventing tail fins from sticking to the walls during agitation. Together, these adjustments
substantially improved survival rates, enabling effective non-destructive genotyping without
compromising experimental viability. The absence of morphological changes in surviving larvae,
supported by our mRNA transcriptomics data showing no significant changes in the
mitochondrial gene expression, further confirmed the minimal physiological impact of the
protocol on mitochondrial homeostasis. These findings are particularly valuable for longitudinal
studies and subsequent phenotypic analyses that require the same individual to be monitored
over time. Several technical considerations emerged during out optimization process. We found
that rapid replenishment of embryo water post-DNA collection and prompt removal from the
heated shaker significantly improved larvae survival. These insights highlight the importance of
minimizing exposure to Proteinase K solutions and elevated temperatures. Additionally, our
observation that most larval mortality occurred within 24 hours of shaking due to tail fin damage
suggests areas for further protocol refinement. The DNA collected through this non-destructive
method was found to be stable for long periods comparable to conventionally collected fin clip
DNA, with successful amplification and sequencing more than 6 months post-collection.

Our results demonstrated consistent amplification success (>80% sensitivity) across different
mitochondrial DNA and nuclear DNA genetic loci. The robust performance across these diverse
genetic loci suggested that our method could reliably detect a range of mitochondrial and nuclear
sequences, which is particularly important for studies examining mitochondrial and nuclear
genetics. The molecular validation through PCR, RFLP analysis, and Sanger sequencing
confirmed the specificity and reliability of the genotyping method. The precise restriction
enzyme digestion patterns obtained for each mitochondrial gene demonstrated the specificity of
the extracted DNA. This methodological validation is crucial for ensuring the accuracy of
downstream genetic analyses and interpretations. Additionally, we successfully amplified two

other important mitochondrial genes—*mt-atp6* and *mt-co1*—conducting only PCR amplification
for these loci with strong amplification throughout. Significantly, we successfully applied our
non-destructive genotyping method to detect gene editing changes in the *mt-tl1* mutants
established earlier using the FusX TALE Base Editor system.[14] The detection of C-to-T base
edits with over 40% mtDNA mutant heteroplasmy levels underscored the protocol's sensitivity
for identifying genetic modifications. This capacity is particularly pertinent for mitochondrial
disorders, where early detection of heteroplasmic variations is essential for elucidating
pathogenic mechanisms.

The RNA-seq analysis revealed a small number of significantly upregulated genes, several of
which are known regulators of circadian rhythm and transcriptional control. Notably, period
circadian clock 1a (*per1a*), nuclear receptor subfamily 1 group d member 1 (*nr1d1*), and aryl
hydrocarbon receptor nuclear translocator-like protein 2 (*arntl2*) were among the most strongly
induced. Their coordinated overexpression suggests that the experimental condition may directly
influence circadian rhythm regulation.[25] In addition to these well-characterized regulators, the
dataset highlights several uncharacterized or poorly annotated transcripts, including
LOC101886224, LOC100535095, and LOC101883914, which showed moderate to strong
upregulation. Interestingly, granulin 2 (*grn2*) and granulin antisense (*grnas*) was also
significantly upregulated. Grn2 protein belong to family of proteins known as granulins that play
a role in cell proliferation, growth and inflammation.[26] However, we didn't find any
mitochondrial resident nuclear encoded genes or mitochondrial DNA encoded genes to be
differentially expressed in the genotyped larvae. These results highlight an important point to
consider that while the method enables efficient genotyping of live zebrafish for longitudinal
research, it can transiently alter gene expression profiles. This effect has the potential to
confound downstream results, especially if the genes or pathways altered by genotyping overlap
with those under investigation. Laboratories should consider whether genotyping could influence
cellular homeostasis or other key processes central to their studies and adopt controls
accordingly during experimental analysis.

Another important consideration for downstream experiments following genotyping is the
incubation temperature. Incubation at 37 °C may transiently activate heat shock response
pathways, particularly in transgenic lines carrying heat shock–inducible promoters. Although our
RNA-seq analysis did not detect heat shock–related gene expression changes, we cannot exclude
the possibility of transient stress-related transcriptional responses immediately after treatment.

By preserving larval integrity, this method enables longitudinal analyses that are crucial for
determining the germline inheritance of mitochondrial variations, an aspect that remains
underexplored in current zebrafish models. Additionally, the ability to correlate specific mtDNA
mutations with metabolic dysfunction and disease progression opens avenues for more detailed
phenotypic and metabolic assays. This continuous tracking across developmental stages not only
enhances the robustness of genotype–phenotype correlations but also paves the way for

translational applications in modeling mitochondrial diseases with early onset lethality. Moreover, the high-throughput nature of the protocol facilitates the screening of large sample sizes, enhancing statistical power when investigating rare mitochondrial variants and their associated phenotypes.

In conclusion, the optimization of the non-destructive enzyme genotyping method provides a valuable tool for zebrafish researchers, enabling efficient molecular characterization of both nuclear and mitochondrial genomes while maintaining animal viability. This approach not only enhances the ethical aspects of zebrafish research by reducing animal mortality but also expands the experimental possibilities for developmental, genetic, and disease modeling studies, particularly for conditions with early onset lethality.

Acknowledgments

The authors would like to thank the staff of the Mayo Clinic Zebrafish Facility and the University of Texas at Austin Zebrafish Facility. We would also like to acknowledge Dr. Karl J. Clark, Texas A&M University, Texas, USA for the valuable suggestions and feedback.

Authors' Contributions

The idea was conceived by A.S and S.C.E. The article was written by K.S.M, S.R.H, A.S., and S.C.E. Experiments were executed by K.S.M., S.R.H., M.M., and A.S. with experimental guidance from S.C.E. Data analysis was completed by K.S.M, S.R.H., and A.S. The article was reviewed and edited by K.S.M, A.S, and S.C.E.

Data Availability

Raw sequencing data has been uploaded on NCBI SRA (SRA ID: PRJNA1243465).

Author Disclosure Statement

The authors declare that there are no conflicts of interest related to the work reported in this manuscript.

Funding Information

This work was funded by grants from the NIH R01-GM063904, Dell Medical School at The University of Texas at Austin and Mayo Foundation for Medical Education and Research.

References

1. Driever W, Solnica-Krezel L, Schier AF, et al. A genetic screen for mutations affecting embryogenesis in zebrafish. Development 1996;123:37–46; doi: 10.1242/dev.123.1.37.

2. Amsterdam A, Burgess S, Golling G, et al. A large-scale insertional mutagenesis screen in
zebrafish. Genes Dev 1999;13(20):2713–2724; doi: 10.1101/gad.13.20.2713.

3. Lieschke GJ, Currie PD. Animal models of human disease: zebrafish swim into view. Nat Rev
Genet 2007;8(5):353–367; doi: 10.1038/nrg2091.

4. Nasevicius A, Ekker SC. Effective targeted gene "knockdown" in zebrafish. Nat Genet
2000;26(2):216–220; doi: 10.1038/79951.

5. Haffter P, Granato M, Brand M, et al. The identification of genes with unique and essential
functions in the development of the zebrafish, Danio rerio. Development 1996;123:1–36; doi:
10.1242/dev.123.1.1.

6. Xu C, Zon LI. The Zebrafish as a Model for Human Disease. In: Fish Physiology Elsevier;
2010; pp. 345–365; doi: 10.1016/S1546-5098(10)02909-2.

7. Patton EE, Zon LI, Langenau DM. Zebrafish disease models in drug discovery: from
preclinical modelling to clinical trials. Nat Rev Drug Discov 2021;20(8):611–628; doi:
10.1038/s41573-021-00210-8.

8. Best JD, Alderton WK. Zebrafish: An in vivo model for the study of neurological diseases.
Neuropsychiatr Dis Treat 2008;4(3):567–576; doi: 10.2147/ndt.s2056.

9. Broughton RE, Milam JE, Roe BA. The complete sequence of the zebrafish (Danio rerio)
mitochondrial genome and evolutionary patterns in vertebrate mitochondrial DNA. Genome
Res 2001;11(11):1958–1967; doi: 10.1101/gr.156801.

10.   Steele SL, Prykhozhij SV, Berman JN. Zebrafish as a model system for mitochondrial
biology and diseases. Transl Res 2014;163(2):79–98; doi: 10.1016/j.trsl.2013.08.008.

11.   Sabharwal A, Campbell JM, Schwab TL, et al. A Primer Genetic Toolkit for Exploring
Mitochondrial Biology and Disease Using Zebrafish. Genes (Basel) 2022;13(8):1317; doi:
10.3390/genes13081317.

12.   Mok BY, de Moraes MH, Zeng J, et al. A bacterial cytidine deaminase toxin enables
CRISPR-free mitochondrial base editing. Nature 2020;583(7817):631–637; doi:
10.1038/s41586-020-2477-4.

13.   Cho S-I, Lee S, Mok YG, et al. Targeted A-to-G base editing in human mitochondrial
DNA with programmable deaminases. Cell 2022;185(10):1764-1776.e12; doi:
10.1016/j.cell.2022.03.039.

14.   Sabharwal A, Kar B, Restrepo-Castillo S, et al. The FusX TALE Base Editor (FusXTBE)
for Rapid Mitochondrial DNA Programming of Human Cells In Vitro and Zebrafish Disease
Models In Vivo. CRISPR J 2021;4(6):799–821; doi: 10.1089/crispr.2021.0061.

15.   Guo J, Zhang X, Chen X, et al. Precision modeling of mitochondrial diseases in zebrafish
via DdCBE-mediated mtDNA base editing. Cell Discov 2021;7(1):78; doi: 10.1038/s41421-
021-00307-9.

16.   Qi X, Chen X, Guo J, et al. Precision modeling of mitochondrial disease in rats via
DdCBE-mediated mtDNA editing. Cell Discov 2021;7(1):95; doi: 10.1038/s41421-021-
00325-7.

17.   Lee H, Lee S, Baek G, et al. Mitochondrial DNA editing in mice with DddA-TALE
fusion deaminases. Nat Commun 2021;12(1):1190; doi: 10.1038/s41467-021-21464-1.

18.   Mok BY, Kotrys AV, Raguram A, et al. CRISPR-free base editors with enhanced activity
and expanded targeting scope in mitochondrial and nuclear DNA. Nat Biotechnol
2022;40(9):1378–1387; doi: 10.1038/s41587-022-01256-8.

19.   Wilkinson RN, Elworthy S, Ingham PW, et al. A method for high-throughput PCR-based
genotyping of larval zebrafish tail biopsies. Biotechniques 2013;55(6):314–316; doi:
10.2144/000114116.

20.   Meeker ND, Hutchinson SA, Ho L, et al. Method for isolation of PCR-ready genomic
DNA from zebrafish tissues. Biotechniques 2007;43(5):610, 612, 614; doi:
10.2144/000112619.

21.   Lambert CJ, Freshner BC, Chung A, et al. An automated system for rapid cellular
extraction from live zebrafish embryos and larvae: Development and application to
genotyping. Wen Z. ed. PLoS ONE 2018;13(3):e0193180; doi:
10.1371/journal.pone.0193180.

22.   Zhang X, Zhang Z, Zhao Q, et al. Rapid and Efficient Live Zebrafish Embryo
Genotyping. Zebrafish 2020;17(1):56–58; doi: 10.1089/zeb.2019.1796.

23.   Kluesner MG, Nedveck DA, Lahr WS, et al. EditR: A Method to Quantify Base Editing
from Sanger Sequencing. CRISPR J 2018;1(3):239–250; doi: 10.1089/crispr.2018.0014.

24.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome Biol 2014;15(12):550; doi: 10.1186/s13059-014-0550-
8.

25.   Sacksteder RE, Kimmey JM. Immunity, Infection, and the Zebrafish Clock. Infect Immun
2022;90(9):e0058821; doi: 10.1128/iai.00588-21.

26.   Cadieux B, Chitramuthu BP, Baranowski D, et al. The zebrafish progranulin gene family
and antisense transcripts. BMC Genomics 2005;6:156; doi: 10.1186/1471-2164-6-156.

Figure Legends

457
458
Figure 1: Non-Destructive Enzyme Genotyping Workflow and Validation of Survival and
Sensitivity. (A) Schematic of non-destructive enzyme genotyping protocol. **(B)** Schematic of the
zebrafish mitochondrial genome showing genes targeted for non-destructive genotyping
(highlighted in red boxes) **(C)** Box-and-whisker plot comparing survival percentages of larvae

24-hour post genotyping in 96-well, 48-well, and unshaken control groups. Whiskers represent
maximum and minimum values. p-Values were determined by Student's t-test (*p$<$0.05;
p$<$0.01; ***p$<$0.001). (N=100 technical replicates across 5 experiments). **(D) Sensitivity
of the protocol is represented by the percentage of successfully amplified samples across four
mitochondrial genes (N=60 technical replicates across 3 experiments). No statistically significant
difference was observed between the well formats with respect to the amplicon sensitivity. Error
bars are represented as standard error of mean. Figure 1 A and B was created using
Biorender.com.
471

Figure 2: Specificity of Mitochondrial DNA Amplification. **(A)** Gel electrophoresis of the mitochondrial gene *mt-nd1*, showing amplification of the 350 bp band. **(B)** Screening of *mt-nd1* amplicon by RFLP. PCR amplicons from wild type larvae show the presence of uncut product (350 bp) and expected 236 and 114 bp digested bands. UD = undigested PCR product. **(C)**

Representative chromatogram of the *mt-nd1* amplicon depicting the restriction enzyme recognition site. **(D)** Agarose gel depicting amplification of the mitochondrial gene *mt-cytb*, with the expected 460 bp fragment. **(E)** RFLP analysis of the *mt-cytb* PCR product, showing the intact 460 bp band and the predicted 317 and 143 bp digestion fragments. UD = undigested PCR product. **(F)** Representative electropherogram of the *mt-cytb* amplicon, with the restriction enzyme recognition site indicated. **(G)** Visualization of *mt-rnr1* amplification, showing a discrete 395 bp band. **(H)** Restriction digest of the *mt-rnr1* PCR product, displaying both the undigested 395 bp fragment and the expected 253 and 142 bp cleavage products. UD = undigested PCR product. **(I)** Sanger sequencing trace of the *mt-rnr1* amplicon, highlighting the enzyme cut site.

Figure 3: Screening for mtDNA mutants using non-destructive genotyping in *mt-tl1* locus. **(A)** PCR amplification of the *mt-tl1* loci from wild-type larvae resolved on a 1% agarose gel to confirm product integrity. **(B)** Schematic of the SacI recognition site within the amplicon (363 bp) and the expected fragment sizes (186 and 177 bp) for both unedited and edited alleles. **(C)** PCR products from *mt-tl1* mutants are shown on an agarose gel, demonstrating successful

amplification. **(D)** RFLP analysis reveals that C☐to☐T conversion at the target cytosine leading
to loss of SacI recognition site: control larvae yield the expected 186☐bp and 177☐bp fragments,
whereas edited samples display a single undigested band (each lane represents individual sample
DNA; UD = undigested PCR product). **(E)** Sanger sequencing chromatograms from control and
mutant larvae. Asterisk (*) denotes the site of edit with corresponding editing percentage (C-to-
T). Chromatograms and editing table plot were obtained using EditR.

498
Figure 4: Specificity of Nuclear DNA Amplification. (A–F) Gel electrophoresis demonstrates
successful PCR amplification for each nuclear locus: *mavs* **(A)** yields a distinct 367 bp product;
cox7b **(C)** a 1367 bp product; and *lrpprc* **(E)** a 380 bp product, corresponding to expected
amplicon size for each locus. **(B, D, F)** Representative Sanger sequencing chromatograms for the
mavs **(B)**, *cox7b* **(D)**, and *lrpprc* **(F)** amplicons show high-quality, clearly resolved peaks,
confirming the fidelity of PCR amplification and enabling downstream sequence analysis.
505
506

Supplementary Figure Legends

Supplementary Figure 1: Effect of 5% non-fat dry milk (NFDM) solution on larval survival
post genotyping. (A) Survival percentage of wild type larvae at 24 hours post genotyping in 96-
well plate treated with NFDM, 96-well plate without 5% NFDM and unshaken control group.
Bars represent survival percentage from three independent genotyping experiments. p-Values
were determined by Student's t-test (*p < 0.05; **p < 0.01; ***p<0.001). (N=60 technical
replicates across 3 experiments). **(B-D)** Representative bright-field images of 3-dpf wild type
larvae: **(B)** unshaken control, **(C)** 96-well plate without 5% NFDM showing tail damage in
subset of larvae, and **(D)** 96-well plate with 5% NFDM showing undamaged, intact larvae.

(Scale bar: 500 µm). **(E-F)** Gel electrophoresis demonstrating that 5% NFDM pre-treatment to
96-well plates does not interfere with the mitochondrial DNA genotyping, showing successful
amplification of mitochondrial genes from larvae in 5% NFDM-treated 96-well plates: **(E)** *mt-*
cytb (460 bp) and **(F)** *mt-tl1* (363 bp).

Supplementary Figure 2: Mitochondrial DNA Amplification for zebrafish *mt-atp6* and *mt-*
523 ***co1*. (A)** Gel electrophoresis of the mitochondrial gene *mt-atp6*, showing amplification of the
524 369 bp band. **(B)** Gel electrophoresis of the mitochondrial gene *mt-co1*, showing amplification of
525 the 494 bp band.

Supplementary Figure 3: Volcano plot illustrating differentially expressed genes. The x-axis
528 represents the log_2 fold change in gene expression, with positive values indicating upregulation

529 and negative values indicating downregulation. The y-axis shows the minus $\log_{10}$ of the adjusted
530 p-value, representing statistical significance. Each dot represents a gene, with black dots
531 indicating genes not significantly changed and green dots indicating significantly upregulated
532 genes above the threshold (dotted line at adjusted p-value = 0.05).

Supplementary Figure Legend

534 **Supplementary Table 1:** List of oligonucleotides used in this study

Target gene	Forward primer (5′ -3′)	Reverse primer (5′-3′)
mt-nd1	CAGGAAGACACATGACTTCTA	AGTTGGTCGTATCGGAATCG
mt-cytb	GCAATACACTACACCTCAGACAT	ATCCGGTTTCGTGGAGAAATAG
mt-rnr1	GGTTAAACGAGAGGCTCTAGT	ATTCCTTCACAGGGTAAGCTG
mt-tl1	GTTCAGACCGGAGTAATCCA	GGAGTTGTATATAGCCTAACAC
mt-atp6	ACCGTCCAAACATGATTGACT	TGCTGTGAGGTTTGCAGTAAG
mt-co1	GGAAACTGACTTGTGCCACT	ACTTCTGGGTGGCCAAAGAA
mavs	GGCCGTATATGGGTCGTTTT	TTGTGATCCCCCTGATCCTG
cox7b	CGTAAAACCACGGAACAGTTT	GGAAAGTCCATTTGATAGTTGACC
lrpprc	CAAGTCCTCACATTTGTTACC	CTTTCACTGCAGACATGTGG

Supplementary File Legend

537 **Supplementary File 1: Fold change for all the genes from RNA sequencing experiment**