Efficient packaging of HIV-1 genomes via recognition of its adenosine-rich content by a

heterologous RNA-binding domain

Hung R. Vuong[1], Qianzi Zhou[1], Sydney Lesko[2], Kasyap Tenneti[1], Keanu Davis[1], Shanyqua

Scott[2], Daphne Boodwa-Ko[1], Jenna E. Eschbach[1], Kamya Gopal[3], Jessica M. Porter[1], Yiqing

Wang[1], Shawn Mohammed[1], Nakyung Lee[1], Alice Telesnitsky[3, 4], Nathan Sherer[2], Sebla B.

Kutluay[1*]

[1]Department of Molecular Microbiology, Washington University School of Medicine, Saint Louis,

Missouri, USA

[2]McArdle Laboratory for Cancer Research and Institute for Molecular Virology, University of

Wisconsin, Madison, USA.

[3]Cell and Molecular Biology Program, University of Michigan Medical School, Ann Arbor, MI

USA.

[4]Department of Microbiology and Immunology, University of Michigan Medical School, Ann

Arbor, MI USA.

*Corresponding author: kutluay@wustl.edu

ABSTRACT

Selective packaging of a dimeric HIV-1 genome (gRNA) is thought to be driven by specific

binding of the nucleocapsid (NC) domain of viral Gag protein to the packaging signal (Ψ) in the

host cell cytosol. Through replacement of NC with heterologous RNA-binding domains (RBDs)

with distinct RNA-binding properties, we show that the biased adenosine-rich nucleotide content

of the gRNA facilitates its selective packaging. Despite disparate RNA binding specificities, all

Gag-RBD chimeras successfully recruited the gRNA to the plasma membrane, but many were

arrested at the assembly stage. Only the Gag-SRSF5 chimera, which multimerized efficiently on

adenosine-rich sequences on the gRNA, packaged gRNA at near wild-type levels. Many Gag

chimeras displayed potent dominant negative activities, highlighting NC functions as a

targetable step in virus replication. Together, our findings reveal an unexpected aspect of the

HIV-1 gRNA, its biased nucleotide content, as a key driver of selective genome packaging.

INTRODUCTION

Selective packaging of viral genomes is integral for viral replication and contributes to rapid viral evolution. For HIV-1, a non-covalent dimer of the unspliced viral mRNA serves as the packageable viral genome (gRNA)[1-3]. During reverse transcription in a target cell, the gRNA dimer allows the virus to undergo recombination at a high rate, which maintains viral genome integrity and helps HIV-1 overcome the selective pressures imposed by host immune responses and antiviral drugs[4-7]. Similar to HIV-1, RNA viruses with segmented genomes (e.g. members of *Orthomyxoviridae*, *Sedoreoviridae*) can acquire novel antigenic properties through genetic reassortment, where genome segments from co-infecting viruses are exchanged and packaged by viral progeny[8-12].

HIV-1 gRNA packaging is a highly efficient process with nearly all HIV-1 particles (>90%) containing a dimer of gRNA[1]. How such a high level of enrichment is achieved despite gRNA competing with cellular mRNAs in infected cells is a long-standing question. It is well established that the interaction of the nucleocapsid (NC) domain of the major HIV-1 structural protein, Gag, with gRNA is required for genome encapsidation[13,14]. Efficient gRNA packaging is widely thought to be initiated in the cytosol by high affinity binding of NC to distinct sites within the cis-acting packaging signal, Ψ, located near the 5' end of gRNA[15,16]. Ψ folds into three conserved stem loops known as SL1, SL2, and SL3[3]. Multiple *in vitro* studies have shown that unpaired guanosine residues within these stem loops are preferentially bound by Gag/NC and act synergistically to mediate gRNA packaging[17-19]. In agreement, crosslinking immunoprecipitation (CLIP) studies in cells have also shown a clear propensity of cytosolic Gag to bind to multiple regions within Ψ[20]. However, deletions of the Ψ stem loops do not result in a complete loss of packaging[21-23]. Furthermore, Gag binds with high affinity to both Ψ-containing and Ψ-null RNAs at physiological ionic strength *in vitro*[24] and packages cellular mRNAs into

virions, particularly in the absence of HIV-1 gRNA[25,26]. Hence, selective binding of Gag to Ψ

appears to be insufficient to support selective gRNA packaging.

While Gag-Ψ interactions likely initiate in the cytosol[20,27], whether this early genome recognition

event is the most critical step for selective gRNA packaging is unclear. Based on live-cell

imaging studies, a small number of Gag molecules appear to be required for trafficking and

stable anchoring of the gRNA at the plasma membrane (PM)[28-30]. This nucleation complex

subsequently recruits additional Gag molecules from the cytosol to drive the formation of high-

order Gag multimers, which ultimately results in encapsidation of the gRNA in budding

virions[3,31-33]. In addition, RNA plays a structural component in virions[34] and capsid (CA) domain

mutations that block Gag lattice assembly can impair specific binding to Ψ and genome

packaging[35,36]. Based on these findings, gRNA was proposed to be selectively packaged due to

having multiple Gag binding sites that help increase local Gag concentration and consequently

making HIV-1 RNAs more efficient at promoting Gag-Gag interactions that drive virus

assembly[3,16]. However, the specific features of HIV-1 gRNA that promote efficient virus

assembly and the mechanism by which Gag multimerizes more readily on viral RNAs are not

understood.

The HIV-1 gRNA has a highly biased nucleotide content: rich in adenosines (~36%) and poor in

cytosines (~18%) with purines overrepresented in all codon positions of all HIV-1 genes[37,38]. It is

plausible that the unique nucleotide content of the HIV-1 gRNA can facilitate high avidity Gag-

gRNA interactions at the PM, which in turn enhances Gag multimerization on the HIV-1 gRNA

during virion assembly. While prior CLIP studies have revealed the propensity of Gag to bind to

A/G-rich motifs in cellular mRNAs recruited to the PM[20], direct evidence that supports the role of

purine-rich elements in gRNA packaging is lacking.

In this study, we generated Gag-RBD chimeras with altered RNA binding specificities by

replacing its NC domain with RNA binding domains (RBDs) derived from the heterogeneous

nuclear ribonucleoprotein (hnRNP) and serine-arginine-rich (SRSF) protein families with the

hypothesis that like NC, Gag-RBD chimeras with purine-rich binding preference could package

gRNA efficiently. Remarkably while all chimeras were capable of binding to Ψ in the cytosol and

recruiting gRNA to the PM, we found that Gag-SRSF5, which preferentially binds to A/G-rich

sequences, was the sole chimera that packaged gRNA at near WT level. Gag-SRSF5 chimera

efficiently multimerized on the HIV-1 gRNA at the PM and displayed a remarkably similar

purine-rich RNA binding specificity with WT Gag in immature virions. Notably, mutations that

reduced the specificity of Gag-SRSF5 for A/G-rich motifs also reduced its gRNA packaging

efficiency. While each of the Gag-RBD chimeras we generated yielded non-infectious virions

and several displayed particle release defects, they were nonetheless able to co-assemble with

WT Gag and potently inhibited virion infectivity. Taken together, our findings indicate that Gag

binding to gRNA in the cytosol and trafficking it to PM assembly sites are insufficient to ensure

selective genome packaging. Rather, our findings provide strong support for the notion that

efficient Gag multimerization on the unusually A-rich HIV-1 gRNA at the PM favors virus

assembly on—and thus selective encapsidation of—gRNA over competing cellular mRNAs.

RESULTS

Gag-SRSF5 is unique in its ability amongst Gag-RBD chimeras to efficiently package the

HIV-1 gRNA

To alter the RNA binding specificity of Gag, we replaced its NC domain with tandem RBDs (e.g.

RNA-recognition motifs (RRMs) and K homology domains (KH)) from the hnRNP and SRSF

protein families within the context of a fully infectious HIV-1$_{NL4-3}$ proviral plasmid bearing a 3xHA

tag between capsid (CA) and matrix (MA) domains of Gag (**Fig. 1A**). We reasoned that RBDs

that preferentially bind to and multimerize on the A/G-rich sequences of HIV-1 gRNA would be

able to replace the genome packaging function of NC. Internal controls included replacement of

NC with leucine zipper domain from GCN4 (bZIP) that facilitates Gag multimerization in the

absence of NC [39] and complementation by NC itself. The set of hnRNP and SRSF proteins (**Fig.**

1A) were chosen because their RNA binding properties are well defined and prior studies have

shown that they directly bind to HIV-1 RNAs to regulate their alternative splicing [40-45]. For

example, hnRNP A1 recognizes "AGG" motifs, hnRNP F binds to stretches of oligonucleotide G

motifs, hnRNP K binds to short C-rich tracks, and SRSF1, 4 and 5 all recognize "AAG" motifs.

Analogous to NC, which binds to cognate RNA via two zinc finger motifs, hnRNP and SR

proteins possess tandem RBDs that create a larger binding interface enabling recognition of

longer sequence motifs with enhanced affinity compared to individual RBDs.

As expected, replacing the NC domain, which has multiple functions in HIV-1 replication beyond

packaging[46-49], rendered all chimeric viruses non-infectious **(Fig. 1B)**. Notwithstanding, all

chimeras expressed similar levels of Gag and Gag-Pol in cells, irrespective of the presence or

absence of a functional PR **(Fig. 1C, S1A).** The expression of Env and Nef proteins, generated

through translation of partially and completely spliced HIV-1 transcripts, respectively, were

similar across WT Gag and Gag-RBD chimeric viruses, suggesting that the isolated RBDs of

hnRNPs and SR proteins did not have unintended consequences on viral mRNA splicing (**Fig.**

1C, S1A). In the case of PR$^+$ viruses, Gag processing intermediates and fully processed CA

(p24) were present (**Fig. 1C**). Notably, particle release was largely blocked for Gag-hnRNP A1, -

hnRNP F and hnRNP K chimeras **(Fig. 1C, S1B)**. Gag-SRSF1 and Gag-SRSF4 displayed a

more modest release defect whereas Gag-SRSF5 release was largely unaffected **(Fig. 1C,**

S1B). Gag-SRSF5 particles contained equivalent levels of Gag-Pol and integrase,

demonstrating that virion incorporation and processing of Pol was also unaffected **(Fig. 1C,**

S1B).

As intended, all Gag-RBD chimeras could bind RNAs in cells, with a much higher propensity to

form Gag-RNA adducts than WT as revealed by autoradiography analysis of

immunoprecipitated Gag-RNA complexes (**Fig. 1D, 1E**). However, only the Gag-SRSF5

packaged gRNA nearly as efficiently as WT Gag based on RT-qPCR analysis of purified mature

and immature virions **(Fig. 1F, S1C, S1D)**. On the other hand, Gag-SRSF5 appeared to be less

selective for gRNA compared to WT Gag and packaged a higher proportion of multiply spliced

HIV-1 RNAs **(Fig. 1F, S1C, S1D)**. In addition to the viral gRNA, certain host RNAs including 7SL

RNA are markedly enriched in HIV-1 virions[20,50-52]. Surprisingly, Gag-SRSF5 also encapsidated

host 7SL RNA at levels comparable to that of WT Gag (**Fig. 1F, S1C, S1D**). Northern blotting in

Gag-SRSF5 virions confirmed these results and additionally demonstrated that both gRNA and

7SL RNA in Gag-SRSF5 particles were intact (**Fig. S1E**).

To gain further insight into the extent with which Gag-SRSF5 selectively packages gRNA over

cellular RNAs, we next conducted total RNA-seq experiments in purified virions. Expectedly, for

WT Gag and ΔNC-NC virions HIV-1 gRNA mapping reads constituted ~80% of the total reads

(**Fig. 1G**). Roughly half of the remaining reads were derived from rRNAs (**Fig. S1F**) and other

host-derived RNAs primarily composed of mRNAs and 7SL RNA (**Fig.1H, S1G**). In comparison,

only about 25% of RNA-seq-derived reads were of HIV-1 origin in Gag-SRSF5 virions, whereas

the majority of the remaining reads mapped to host mRNAs as well as 7SL RNA (**Fig. 1G, 1H,**

S1F, S1G), supporting the above finding that Gag-SRSF5 is less selective towards gRNAs. With

the exception of hnRNP A1, viral RNAs constituted only a small proportion of RNA-seq derived

reads for the closely related Gag-SRSF1 and Gag-SRSF4 as well as other Gag-RBD chimeras,

and the remaining reads mapped to mRNAs and host rRNAs (**Fig. 1G, 1H, S1F**). Note that the

observed high abundance of rRNA reads in Gag-RBD chimeras with low virus yields as well as

Gag-bZIP and Gag-ΔNC chimeras is likely due to PCR overamplification (per RNA-seq protocol)

of otherwise low abundance RNAs in virus preps. Nevertheless, comparison of mRNA-mapping

reads revealed that the host mRNAs packaged by ΔNC-NC virions were most similar to those

packaged by Gag-SRSF1, 4 and 5 as indicated by the highest Jaccard index values with > 0.2

generally considered significant correlation (**Fig. 1I**).

Single virion analysis of Gag-SRSF5 reveals its ability to package the HIV-1 gRNA at near

WT Gag efficiency

We next sought to gain more granular insight into gRNA packaging and virion assembly

properties of Gag-SRSF5 at a single virion level. To do so, we used a modified, visible virus

encoding YFP tagged Gag (Gag-YFP) and gRNA bearing 24 copies of an RNA stem loop

derived from the MS2 bacteriophage (MSL) that can be tracked in proxy by a virally-encoded

MS2 coat protein fused to mCherry (**Fig. 2A**)[28,53-55]. No differences were observed in virion-

incorporated Gag-YFP levels for WT compared to Gag-SRSF5 particles (**Fig. S2A**), suggesting

that Gag incorporation per virion and particle size were not affected. In contrast, ΔNC virions

incorporated considerably lower levels of Gag-YFP per virion (**Fig. 2B, S2A**). In agreement with

previous reports[1], ~90% of WT and ΔNC-NC particles contained gRNA, whereas the majority of

ΔNC virions lacked detectable genomes, also as expected (**Fig. 2B**). While ~70% of Gag-

SRSF5 chimeras encapsidated gRNA **(Fig. 2B),** we noted a trend towards lower gRNA signal

intensity per virion (**Fig. 2C, S2A**) suggesting that not all Gag-SRSF5 particles may contain a

gRNA dimer.

To interrogate the virion morphology of Gag-SRSF5 particles further, we conducted thin section

electron microscopy (TEM) analyses of purified virions. Similar to WT and ΔNC-NC particles,

Gag-SRSF5 particles were ~100nm in diameter but contained eccentrically localized

condensates and a higher fraction of particles with immature morphology **(Fig. 2D).** Biochemical

fractionation of viral cores isolated from Gag-SRSF5 virions indicated the presence of unstable

or improperly assembled cores similar to those observed for ΔNC and bZIP virions **(Fig. 2E, 2F)**

and possibly due to the presence of aberrant Gag processing intermediates (**Fig. S2B**). These

aberrations combined with the inability to execute various NC functions likely explain the early

reverse transcription defects of Gag-SRSF5 particles in target cells (**Fig. S2C**) and the

accompanying decrease in virion infectivity (**Fig. 1B**).

Gag-RBD chimeras efficiently recruit HIV-1 gRNA to the PM but many display virion

assembly defects

HIV-1 virion assembly involves stepwise localization of a gRNA dimer at the PM, Gag

multimerization on the PM-associated gRNA forming an immature Gag lattice, and recruitment

of host ESCRT-III machinery for budding[56]. The particle release defects of the several Gag-RBD

chimeras prompted us to evaluate at what step of virion assembly they were blocked at, which

we reasoned would help explain what makes WT Gag and Gag-SRSF5 successful at gRNA

packaging.

First, we focused on Gag and gRNA localization at the PM, given that gRNA recruitment and its

stable tethering at the PM are mediated by NC[28,29]. Like WT Gag, all Gag-RBD chimeras

localized in the cytosol and at the PM at steady state, except for Gag myristoylation mutant

(G2A) which is defective in PM anchoring and was enriched in the cytosol **(Fig. 3A, S3A, S3B)**.

Surprisingly, even though only Gag-SRSF5 packaged gRNA, all Gag-RBD chimeras recruited

gRNA to the PM as efficiently as WT Gag **(Fig. 3B)**. Of note, while Gag-hnRNP A1 chimera

preferentially retained gRNA over multiply spliced viral RNAs at the PM, other Gag chimeras

were comparably less selective **(Fig. 3B)**. In contrast to viral RNAs, the abundance of the

control host GAPDH mRNA at the PM was not dependent on the presence of Gag with a

functional NC domain or RBDs (**Fig. 3B**). To corroborate these findings and interrogate the

influence of heterologous RRMs on the trafficking properties of Gag chimeras with gRNA at a

single-cell level, we next conducted live-cell imaging studies using the dual labeling system

described in Fig. 2A. In agreement with biochemical experiments, all Gag chimeras were able to

recruit the gRNA to the PM **(Fig. 3C, S3C)**. However, only Gag-SRSF5 formed WT-like foci with

gRNA at the PM whereas the other Gag chimeras appeared to form sheet-like assemblies **(Fig.**

3C, S3D-E).

These data suggested that the particle release defect of select Gag-RBD chimeras is not due to

impaired trafficking with gRNA to the PM. To interrogate the next steps in virion assembly,

namely whether Gag-RBD chimeras can multimerize at the PM and form membrane curvature

necessary for recruitment of (ESCRT-III) machinery[57], we conducted TEM experiments in virus

producer cells. In agreement with live-cell imaging experiments, we found that Gag-RBD

chimeras with release defects formed extended electron dense sheet-like structures at the PM

(Fig. 3D), suggesting that they can all multimerize and form Gag lattices. However, unlike WT

and ΔNC-NC which formed well-defined curved buds, the electron dense sheet-like structures

from hnRNP A1, F, K and Gag-SRSF1 chimeras had little to no membrane curvature. Gag-

SRSF4, and SRSF5, on the other hand, formed buds that were closer in shapes and sizes to

WT and ΔNC-NC control **(Fig. 3D)**. Altogether, these data suggest that all Gag-RBD chimeras

possessing diverse RNA-binding modules can find and retain gRNA at the PM. However, events

post-Gag:gRNA localization, possibly the inability to multimerize on the gRNA at the PM,

rendered most of the chimeras unable to bud and package gRNA efficiently.

Gag-RBD chimeras maintain the known RNA-binding specificities of the isolated RBDs

but bind similarly to HIV-1 gRNA

To determine which feature of Gag-SRSF5 makes it uniquely capable of gRNA packaging, we

interrogated the RNA-binding specificities of Gag-RBD chimeras using CLIP in cells, at the PM

and in immature virions. In line with a prior report[20], a relatively small percentage of CLIP-

derived reads mapped to viral RNAs for WT Gag and Gag-ΔNC-NC **(Fig. 4A)**. Similarly, despite

co-immunoprecipitating with higher levels of total RNA (**Fig. 1D, 1E**), a relatively small

percentage of CLIP reads for Gag-RBD chimeras were derived from viral RNAs (**Fig. 4A**).

Analysis of reads that map to cellular RNAs revealed that similar to WT Gag and Gag-ΔNC-NC,

all Gag chimeras, with the exception of Gag-hnRNP F, bound most frequently to cellular tRNAs,

mediated by the matrix domain[20], and mRNAs (**Fig. 4B**). mRNAs constituted comparably a

greater fraction of cell-derived RNAs for Gag-hnRNP A1, -SRSF4 and -SRSF5 chimeras **(Fig.**

4B). Of note, for Gag-hnRNP F, tRNAs made up a considerably smaller proportion of reads for

compared to intron-mapping reads (**Fig. 4B**). This can be ascribed to the nuclear trafficking of

this chimera **(Fig. 3C)**, due to the presence of glycine-tyrosine-arginine-rich (GYR) domain

between RRM2 and RRM3 combined with its known preference to bind to intronic regions in

cognate RNAs[43,45,58]. Motif analysis within the bound mRNAs revealed that sequence

preferences of Gag-RBD chimeras are in alignment with the known RNA binding specificity of

the hnRNP/SR proteins that the RBDs were derived from **(Fig. 4C)**[43,45,59-62]: WT Gag as well as

Gag-SRSF1, 4 and 5 chimeras bound preferentially to sequences containing A/G-rich

sequences (**Fig. 4C**), Gag-hnRNP K bound to C-rich oligonucleotides, Gag-hnRNP A1 and

Gag-hnRNP F preferentially bound to sequences containing AGGGG motif and short G repeats,

respectively (**Fig. 4C**).

On the HIV-1 genome, WT Gag and Gag-ΔNC-NC bound to discrete sites within the 5' leader,

including SL1 and SL3 (**Fig. 4D, S4A**), in agreement with prior reports[20]. Surprisingly, all Gag-

RBD chimeras also frequently bound to these sites despite their diverse RNA-binding

specificities, though the extent with which they were bound to SL1 and SL3 varied **(Fig. 4D,**

S4A). Since Gag-SRSF5 also bound to the 5' leader, we tested whether progressive deletions

of structured RNA elements within this region would affect gRNA packaging **(Fig. S4B)**. None of

the 5'UTR deletion mutations impaired gRNA packaging for Gag-SRSF5 (**Fig. S4C, S4D**),

suggesting that sequences or features outside of the 5' leader contributes to its ability to

mediate selective genome packaging.

Gag-SRSF5 multimerizes on the A-rich HIV-1 gRNA at the plasma membrane and in

immature virions

Gag was proposed to undergo a multimerization-driven change in its RNA-binding specificity

towards more frequent binding to A-rich motifs at the PM, which is further enhanced in immature

virions[20]. We next examined whether a similar switch accounts for the unique ability of Gag-

SRSF5 to package gRNA through CLIP experiments in PM fractions and immature virions.

Comparison of membrane-associated Gag-RBD:RNA adducts revealed that unlike cytosolic

Gag (**Fig. 1D, E**), WT Gag and Gag-RBD chimeras bound to comparable levels of RNA **(Fig.**

5A, S5A). The increased level of RNA binding for WT Gag demonstrates its ability to more

extensively engage with cognate RNAs upon localizing to the PM through multimerization. While

this pattern continued in virions for WT Gag, which had the highest level of immunoprecipitated

Gag:RNA adducts, only the Gag-SRSF5 yielded comparable level of protein:RNA complexes

whereas Gag-SRSF1 and Gag-SRSF4 were largely impaired in RNA-binding **(Fig. 5B, S5B)**.

Expectedly, the percentages of viral RNA-derived CLIP reads were enriched in immature virions

compared to cells for WT Gag and Gag-ΔNC-NC **(Fig. 5C)**. However, the level of enrichment

was lower for Gag-SRSF5 **(Fig. 5C)**, in agreement with it packaging gRNA less selectively **(Fig.**

1G).

Strikingly, in immature virions the binding profile of Gag-SRSF5 on gRNA mirrored that of WT

Gag and Gag-ΔNC-NC (**Fig. 5D**) and exhibited a statistically significant correlation nearing that

observed between WT Gag vs. Gag-ΔNC-NC (**Fig. 5E**). In fact, Gag-SRSF5, along with Gag-

SRSF1 and Gag-SRSF4, which bind to A-rich motifs in cells **(Fig. 4C)**, exhibited a notable

switch in their RNA binding profile upon association with the PM. Similar to WT Gag, each of

these chimeras bound more frequently to a larger number of sites throughout the HIV-1 gRNA
at the PM and less prominently to the 5' UTR **(Fig. S5C)**. In contrast, Gag-CAΔCTD mutant that
is significantly impaired in forming high-order multimers bound to more discrete sites on the
gRNA (**Fig. S5C**). Gag-hnRNP A1 and Gag-hnRNP F, which bind to G-rich sequences in cells,
showed little to no increase in binding sites, with their cytosol and PM RNA binding profiles
resembling each other (**Fig. S5C vs. Fig. S4A**).

In addition to the marked similarity between WT Gag and Gag-SRSF5 binding to the HIV-1
gRNA in immature virions, Gag-SRSF5-bound host mRNAs also resembled those bound by WT
Gag and Gag-ΔNC-NC, with mRNAs and 7SL RNA representing the most frequently bound
RNAs in virions **(Fig. 5F)** Lastly, motif analyses on bound cellular mRNAs showed that virion-
associated Gag-SRSF5, WT Gag, and Gag-ΔNC-NC were all preferentially bound to A/G-rich
sequences **(Fig. 5G)**. Taken together, these results provide strong support for the notion that
the ability of Gag to efficiently multimerize on the A-rich gRNA may underlie its capacity to
efficiently package the HIV-1 genome.

Changing the RNA-binding specificity of Gag-SRSF5 specifically impairs gRNA
packaging

SRSF1, SRSF4, and SRSF5 bear tandem RRMs bridged by a flexible linker. A previous
structural study showed that the highly conserved $S_{133}WQDLKD_{139}$ sequence residing within
RRM2 of SRSF1 was responsible for its binding specificity for a minimal GGA motif [63].
Specifically, residues D136, K138 and D139 were engaged in an extensive hydrogen binding
network within the purine-rich recognition site. Based on this structural prediction, we designed
two Gag-SRSF5 mutants, mGGA-1 and mGGA-2, carrying substitutions in the homologous
residues (e.g. D123, K125, D126) to weaken site specific RNA recognition (**Fig. S6A**). In

addition, we engineered two sets of mutations to shift the RNA-binding specificity of SRSF5

towards a CCG motif (mCCG-1 and mCCG-2) or a GGG motif (mGGG) **(Fig. S6B)**.

Reassuringly, Gag-SRSF5 mutations did not impair its overall ability to bind RNA in cells, and all

Gag-SRSF5 variants immunoprecipitated with RNA at higher than WT level **(Fig. 6A, 6C)**. In

contrast, in virions, the parental Gag-SRSF5 variant bound to RNA at moderately higher levels

than the mutants **(Fig. 6B, 6C)**. Gag-SRSF5 mGGA-1 and mGGA-2 mutants packaged gRNA at

levels approximately two-fold lower than the parental Gag-SRSF5 variant **(Fig. 6D)**. The mCCG-

1, mCCG-2 and mGGG mutants displayed stronger defects, with gRNA, multiply spliced viral

RNAs, and 7SL RNA packaged at significantly lower levels **(Fig. 6D-F)**. Importantly, there was

no significant difference for any of these mutants in the packaging efficiency of cellular RPL36A,

RPS6 and RPS16 mRNAs that we and others have found to be abundantly incorporated into

HIV-1 VLPs [25] **(Fig. 6G-I)**. These findings strongly suggest that the shifting the RNA-binding

specificity of Gag-SRSF5 towards A-poor sequences specifically impair packaging of the A-rich

gRNA without affecting its overall ability to bind and package non-viral RNAs.

Gag-RBD chimeras potently interfere with virion infectivity

Since Gag-RBD chimeras are non-infectious, with some having severe release defects, we next

tested whether they can inhibit the release of WT Gag and/or co-assemble with it and

subsequently interfere with HIV-1 replication. Presence of WT Gag in producer cells enhanced

virion incorporation of Gag-hnRNP A1, F and K chimeras with negligible effects on Gag-SRSF1-

5 chimeras **(Fig. 7A, 7B)**. On the other hand, Gag-hnRNP F and Gag-hnRNP K chimeras were

present in virions at comparably lower levels than WT Gag and their presence didn't impair

virion release by WT Gag levels **(Fig. 7A, 7B)**. On the other hand, co-transfection with Gag-

hnRNP A1, Gag-hnRNP K, Gag-SRSF1, Gag-SRSF4, and Gag-SRSF5 substantially reduced

viral titers (≥100 fold), again with negligible effects on particle release **(Fig. 7C, 7D)**. With the

exception of the Gag-hnRNPK chimera which reduced gRNA packaging by ~3-fold, other Gag-

RBD chimeras did not interfere with gRNA packaging (**Fig. 7E**). Hence, the potent antiviral

activities of these chimeras are unlikely due to interference with gRNA packaging by NC. It is

noteworthy that Gag-hnRNP K exerted the most substantial dominant negative effect even

though it was incorporated into particles at a much lower level than WT Gag **(Fig. 7A)**. In

agreement, dose-response experiments corroborated these findings and demonstrated that

Gag-hnRNP K is the most potent antiviral chimera **(Fig. 7F, 7G)**. Interestingly, co-transfection

with Gag-ΔNC and Gag-bZIP also led to a significant decrease in virion infectivity (**Fig. 7C, 7D**),

suggesting that the functional amount of NC copies in virions is finely gauged.

These findings prompted us to test whether the antiviral activity of Gag-RBD chimeras can be

mapped to individual RBDs, and conversely, whether other hnRNP/SR proteins that naturally

bear a single RBDs (e.g. hnRNP C, SRSF2, SRSF3, SRSF7) can also exert antiviral effects. All

Gag chimeras bearing a single RBD were expressed in cells and virion release largely

correlated with their cognate expression level in cells **(Fig. S7A, S7B)**. Similar to Gag chimeras

with tandem RBDs, Gag-single RRM viruses were non-infectious **(Fig. S7C)**. While all of these

Gag chimeras bound RNA in cells **(Fig. S7A, S7B),** none of them, including the Gag-SRSF5-

RRM1 and Gag-SRSF5-RRM2 chimeras, were able to package gRNA **(Fig. S7D)**. Numerous

Gag-RBD chimeras reduced virion infectivity, with unique patterns emerging for each chimera.

For example, individual RRMs of hnRNP A1 each reduced virion infectivity by 10-fold as

compared to 100-fold reduction with both RRMs, suggesting cooperativity between the RRMs

(**Fig. 7C, 7H, 7I**). The potent antiviral activities of Gag-hnRNP K and Gag-SRSF4 largely

mapped to the KH2 and RRM1 domains, respectively (**Fig. 7C, 7H, 7I**). Interestingly, while the

Gag-hnRNP F chimera did not impair virion infectivity, Gag-hnRNP F-RRM1 chimera was

potently antiviral (**Fig. 7C, 7H, 7I**). Of the proteins with single RBDs, Gag-SRSF2 and Gag-

SRSF3 were the most potently antiviral, whereas Gag-SRSF7 did not inhibit virion infectivity and

Gag-hnRNP C1 modestly inhibited infectivity (**Fig. 7H, 7I**). Like with the full-length chimeras,

despite their potent antiviral activities, none of the chimeras interfered with gRNA packaging by

WT Gag (**Fig. 7J**).Lasty, dose-response experiments demonstrated that Gag-hnRNP K-KH2

has the most potent antiviral activity **(Fig. 7K-L)**.While it is unclear how the Gag-RBD chimeras

inhibit virion infectivity and whether they all inhibit the same step in HIV-1 replication, these

findings highlight that they can be repurposed as potent antiviral modules to target NC-

dependent functions post gRNA packaging.

DISCUSSION

HIV-1 genome packaging is widely thought to depend on the selective binding of Gag to gRNA

through recognition of a *cis*-acting packaging signal, Ψ, within the 5'-UTR. Gag-gRNA

interaction initiates in the cytosol in the excess of cellular mRNAs and this initial "finding" step is

thought to be the major driver of gRNA selection. Our findings challenge this notion and

demonstrate that heterologous RBDs with diverse RNA-binding specificities, without exception,

can successfully recruit gRNA to the host cell PM. This suggests that NC in and on its own is

not required for gRNA selection in the cytosol and rather its properties post-PM recruitment is

more critical.

Our findings highlight that the high adenosine content of HIV-1 gRNA serves as a key molecular

signature that mediates selective gRNA packaging following its recruitment to the PM. Hence an

important feature of NC that emerges from our findings is its ability to multimerize on A-rich

sequences on the HIV-1 gRNA. This idea is strengthened by the key finding in our study that

Gag-SRSF5 chimera efficiently packages gRNA through mimicking the ability of NC to

multimerize on A-rich sequences on the viral gRNA both at the PM and in immature virions.

Strong support for this notion is further provided by targeted mutations that altered the RNA-

binding specificity of Gag-SRSF5 resulting in specific inhibition of gRNA packaging.

An unexpected finding from our study is that Gag-RBD chimeras with diverse RNA-binding

specificities appeared to bind regions within Ψ in cells, while maintaining their known RNA-

binding specificities for cellular RNA targets.   Multiple studies have shown that WT Gag

recognizes Ψ via high affinity interactions with unpaired guanosines within SL1 and SL3[17,64-66].

Since the Gag-RBD chimeras tested here, with the exception of Gag-hnRNP K, preferentially

bind to single-stranded G-rich or A/G-rich sequences, it is possible that they bound to Ψ through

similar interactions with unpaired guanosines. Moving forward, it will be important to determine

whether RNA-binding modules that recognize dsRNAs can similarly recruit the HIV-1 gRNA to

the PM and successfully replace NC. In addition, understanding whether Ψ and HIV-1 gRNA

themselves bear unique structural features that destine the gRNA more readily for PM trafficking

by diverse RBDs will provide crucial insight into selective gRNA packaging.

While our findings suggest that multimerization-driven A-rich sequence content recognition

underlies the ability of Gag-SRSF5 to package gRNA, why the closely related Gag-SRSF1 and

Gag-SRSF4 chimeras with similar RNA-binding properties were unable to package gRNA is

unclear.   Together, these findings suggest that simply switching to A/G-rich binding at the PM

may be insufficient for virion genesis. It is possible that the higher RNA binding affinity of these

Gag-RBD chimeras could lead to an excessive amount of RNA being brought to virus assembly

sites, sterically hindering formation of Gag high-order multimers around a single gRNA dimer

and instead forming multimers on other non-productive RNA molecules that is not conducive to

forming membrane curvature. In line with this, Gag-hnRNP and Gag-SRSF1 chimeras exhibited

defects in membrane curvature while effectively multimerizing on the PM. Notably, the release

defects of these chimeras could be rescued at least in part by WT Gag, presumably by reducing

the levels of RNA retained at the PM and suggesting that these chimeras are not inherently

defective in making it into budding virions. Future studies in defined in vitro systems will be

informative to more directly test these models.

The unusual nucleotide composition of HIV-1 gRNA has remarkably remained stable through

multiple decades since the beginning of the HIV-1 pandemic despite the HIV-1 genome being

highly variable. While recent evidence suggest that this property of HIV-1 gRNA is partly shaped

by selective pressure imposed by host antiviral factors and higher G-to-A mutation rates during

reverse transcription, our study raises the possibility that selective packaging of the HIV-1

genome also contribute to the maintenance of A-richness. Specifically, viral RNAs with higher

adenosine content will be preferentially packaged because they are more efficient at driving

Gag multimerization. Furthermore, because Gag binds to unpaired nucleotides, A-richness

could also contribute to genome packaging by decreasing the prevalence of structured regions

throughout the HIV-1 gRNA, owing to adenosine only forming a weak base pair with uracil.

Indeed, an analysis of the secondary structure model of a complete HIV-1 genome found that

single-stranded regions have a high degree of adenosine accumulation, but regions where

secondary RNA structures are formed[67].

In addition to providing a unique tool for studying gRNA packaging, Gag-RBD chimeras will be

instrumental in investigating functions of NC beyond gRNA packaging and a multitude of other

RNA-binding proteins essential for virus replication. NC has been implicated in various steps of

HIV-1 replication, including budding, reverse transcription and integration[47,49,68]. Deeper

dissection of what makes a Gag-RBD chimera successful in executing these steps will provide

key insights into NC functions critical for virus replication. Considering that a single dimeric

gRNA in a virion is bound by a magnitude of viral (e.g. integrase and reverse transcriptase) and

host RNA-binding proteins (e.g. APOBEC3 family), the ability to package gRNAs independent of

NC by Gag-SRSF5 will also help determine the interdependency of how these diverse RNA-

binding proteins interact with the same gRNA.

Given the potent antiviral activities of Gag-RBD chimeras, understanding how they inhibit HIV-1

replication can reveal new vulnerabilities of HIV-1 and inform the development of new virus-

targeted therapeutics. Another relevant application of our findings is the design of gene therapy

vectors. While altering the nucleotide content of a transgene in this setting may enhance vector

production, whether protein expression from mRNAs bearing suboptimal codons will suffice to

obtain the desired phenotype remains to be determined. Hence, fine-tuning of the transgene

codon usage may help enhance packaging while still allowing sufficient protein production.

Because HIV-1 has a high mutation rate, a less restrictive mechanism of selecting viral

genomes for packaging — through efficient virion assembly on viral RNAs with high adenosine

bias rather than specific binding of viral proteins to a packaging signal made up of conserved

RNA sequences — could allow Gag to package the constantly diversifying genomes. Besides

HIV-1, a large number of other viruses with RNA genomes (e.g. HTLV-1 and Rubella) also have

a biased nucleotide composition including [69]. For many RNA viruses, in particular for those with

segmented genomes, molecular features that allow genome selection are poorly defined. While

our study suggests that nucleotide composition may serve as a molecular signature for HIV-1

genome packaging, similar principles can in principle be applicable to a broad range of RNA

viruses.

 FIGURE LEGENDS

Figure 1. Replacement of HIV-1 NC with heterologous RNA-binding modules reveals the

unique ability of Gag-SRSF5 chimera to package gRNA. A. Cartoon diagram of HIV-1 Gag-

RBD chimeras in which the NC domain was replaced by RNA-binding domains from

heterologous RNA-binding proteins of host hnRNP and SRSF proteins. **B.** The indicated HIV-1

virions were titered on TZM-bl reporter cells, with 100 µg/mL dextran-sulfate added at 6-16 hpi

to limit infection to a single cycle. The titers were subsequently normalized based on particle

numbers as quantified by a reverse transcriptase (RT) activity assay and presented relative WT

(set to 1). Titers are mean ± SEM from n=3 independent biological replicates. **C.** HEK293T cells

were transfected with PR^+ proviral plasmids encoding the indicated Gag-RBD chimeric proteins.

Cell lysates and purified virions were harvested 2 days post-transfection and analyzed with

immunoblotting for Gag-Pol (GP), Gag, Envelope (gp120), Nef, Capsid (p24) and Integrase (IN).

D. Representative autoradiogram of Gag:RNA adducts (top) and corresponding anti-HA

immunoblot of Gag proteins (bottom) immunoprecipitated from cells that were transfected with

PR^- proviral plasmids encoding the indicated Gag-RBD chimeras following PAR-CLIP protocol.

E. Densitometry analysis of Gag:RNA adducts from autoradiograms (e.g. panel D) of

independent biological replicates. Data are normalized relative to immunoprecipitated protein

levels and graphed as fold differences relative to WT (set to 1). Graph shows the average of 3-5

independent experiments denoted by symbols, error bars show the SEM. **F.** RT-qPCR

quantification of relative copies of gRNA, multiply spliced HIV-1 RNA (multiply spl.), and 7SL

RNA (7SL) from purified PR^+ virions collected from HEK293T cells expressing the indicated Gag

chimeras. Data show the average of 3 independent experiments and the error bars represent

standard error of the mean (SEM). Indicated statistical analyses (gRNA, multiply spliced RNA

and 7SL RNA) were done by one-way ANOVA with Dunnet's multiple comparison test for

correction, whereby n.s. denotes non-significant comparisons. Other group comparisons are not

shown due to clarity. **G-H.** Proportion of reads that map to viral RNA (G) and cellular non-

ribosomal RNAs (H) obtained from total RNA-seq experiments performed with purified virions.

Data show the average of 2-3 independent experiments, error bars represent the SEM. **I.** Bar

plot showing Jaccard index of reads that map to host mRNAs derived from virion total RNA-seq

to quantify the similarity between the indicated Gag chimeras and WT Gag.

Figure 2. Single virion analysis of gRNA packaging by and virion morphology of the Gag-

SRSF5 chimera. A. Cartoon depiction of strategy used for simultaneous fluorescence imaging

of Gag protein and HIV-1 gRNA. YFP, yellow fluorescent protein; MSL, 24 copies of MS2 RNA

stem-loop; MS2-mChe, bacteriophage MS2 coat protein fused with the fluorescent protein

mCherry. **B.** Representative fluorescence microscopy images of HIV-1 particles harvested from

293T cells transfected with MS2-mCherry plasmid and MSL HIV-1 plasmids expressing the

indicated Gag chimera-YFP fusion protein. Quantification of the percentage of total virions that

contains gRNA (mCherry$^+$YFP$^+$) based on single-virion analysis. **C.** Violin plot showing MS2-

mCherry/Gag-YFP mean fluorescence intensity (MFI) ratios measured for WT, Gag-ΔNC, and

Gag-SRSF5 particles shown in panel B. 60 particles were quantified for each condition. **D.**

Representative thin section electron microscopy (TEM) images showing WT Gag, Gag-ΔNC,

Gag-ΔNC-NC,Gag-bZIP, and Gag-SRSF5 virions. **E-F.** The indicated virions were subjected to

equilibrium density sedimentation following treatment with 0.5% Triton as detailed in Materials

and Methods. Ten fractions harvested from the top of sucrose gradients were analyzed for RT

activity via a qPCR assay (E) or CA content via immunoblotting using a mouse monoclonal anti-

CA antibody (F). Results in E and F show the mean values ± SEM of three independent

biological replicates.

Figure 3. Gag-RBD chimeras efficiently recruit and retain gRNA at the PM. A. Immunoblot

analysis of Gag from membrane (M) and cytoplasmic (C) fractions isolated from 293T cells

transfected with the indicated PR$^-$ proviral plasmids using a mouse monoclonal anti-HA

antibody. **B.** RT-qPCR quantitation of the relative PM/cytosol ratios of gRNA, multiply spliced

HIV-1 mRNA, and GAPDH mRNA copies. Data show the average of three independent

experiments and the error bars represent SEM. Indicated statistical analyses for gRNA were

done by one-way ANOVA with Dunnet's multiple comparison test for correction (**** $p<0.0001$).

Non-significant comparisons are not shown for clarity. **C.** Fluorescence imaging of HeLa cells

transfected with MS2-mCherry and HIV-1 constructs. Scale bars represent 20 µm in full images

and 2.5 µm in regions of interest (ROI) at the PM. ROIs are indicated by yellow dotted boxes.

Yellow arrowheads indicate where Gag and gRNA colocalize. **D.** Representative TEM images

from two independent experiments showing virion assembly sites at the PM of 293T cells

transfected with the indicated Gag plasmids. Scale bar represents 500 nm.

Figure 4. CLIP-seq analysis of Gag-RBD chimeras reveals their RNA-binding properties

in cells. A. Proportion of reads that map to cellular and viral genomes obtained from Gag-CLIP

experiments performed on cell lysates. Each point represents an independent experiment. **B.**

Classification of read clusters generated from cell Gag-CLIP experiments that map to cellular

RNAs; total number of reads is indicated below each pie chart. **C.** Representative sequence

motifs most frequently occurring in Gag-bound cellular mRNA clusters generated from CLIP

experiments. **D.** Read density distribution within the 5' leader (1-400 nt) from cell Gag-CLIP

experiments.

Figure 5. Gag-SRSF5 mimics NC in its ability to multimerize on the gRNA in immature

virions. A-B Densitometry analysis of Gag:RNA adducts from autoradiograms (See Figure

S5A, S5B) of independent CLIP replicates done on PM fractions (A) or immature virions (B).

Data are normalized relative to immunoprecipitated protein levels and graphed as fold

differences relative to WT (set to 1). Graph shows the average of 3 independent experiments

and the error bars represent SEM. **C.** Proportion of reads that map to cellular and viral genomes

obtained from virion Gag-CLIP experiments. Each point represents an independent experiment.

D. Read density distribution on full-length HIV-1 gRNA from virion CLIP-seq experiments. **E.**

Correlation analysis of WT, ΔNC-NC, and Gag-SRSF5 binding to viral RNA in virions from

independent CLIP experiments. **F.** Classification of read clusters generated from virion Gag-

CLIP experiments that map to cellular RNAs; total number of reads is indicated below each pie

chart**.** Representative sequence motifs most frequently occurring in Gag-bound cellular mRNA

clusters generated from virion Gag-CLIP experiments.

Figure 6. Altering the RNA-binding specificity of Gag-SRSF5 specifically impairs gRNA

packaging. A-B. Representative autoradiograms of Gag-RNA complexes (top) and immunoblot

analysis of Gag-SRSF5 mutants (bottom) immunoprecipitated from cells expressing the

indicated PR$^-$ Gag chimeras (A) and progeny virions (B). (*) marks the IgG-heavy chain from the

immunoprecipitating antibody. Eluates in virion IPs in (B) were normalized to load similar

amounts of protein for Gag and Gag-SRSF5 variants to more clearly illustrate RNA binding

ability per protein. **C.** Densitometry analysis of Gag:RNA adducts from autoradiograms (e.g.

panels A and B) of independent replicates. Data are normalized relative to immunoprecipitated

protein levels and graphed as fold differences relative to WT (set to 1). Graph shows the

average of 3 technical replicates, representative of two independent experiments, and the error

bars represent SEM. **D-I.** RT-qPCR quantification of relative copies of gRNA, multiply spliced

HIV-1 RNA, 7SL, RPL36A, RPS6, and RPS16 RNA from HIV-1 particles collected from 293T

cells expressing the indicated Gag. Data show the average of 3 independent experiments and

the error bars represent SEM. Indicated statistical analyses (gRNA, multiply spliced RNA and

7SL RNA) were done by one-way ANOVA with Dunnet's multiple comparison test for correction,

whereby n.s. denotes non-significant comparisons (*$P < 0.05$; **$P < 0.01$; ***$P < 0.001$; ****$P <$

0.0001). Other group comparisons are not shown due to clarity.

Figure 7. Gag-RBD chimeras are potently antiviral. A. Proviral plasmids encoding ΔGag or

WT Gag was co-transfected with the indicated Gag-RBD proviral plasmids in 293T cells. 2 days

post-transfection, cell lysates and purified virions were harvested for immunoblot analysis of

Gag using a mouse monoclonal antibody against the HA-tag. **B.** Quantitation of virion

immunoblot results from panel A. Fold increase of the virion-incorporated Gag-RBD chimeric

proteins in the presence of WTGag over ΔGag co-transfection control was graphed. Data show

the mean fold increase from n=2 independent experiments. **C-L.** 293T cells were transfected

with equal ratio of tandem RRM or single RRM Gag-RBD chimeric HIV-1 proviral plasmids and

a WT Gag HIV-1 proviral plasmid. Progeny virions were titered on TZM-bl reporter cells, with

dextran-sulfate added to limit infection to one cycle. The titers were presented as relative to WT

which was set to 1 (C, H). Particle release from (C) and (H) was quantified by RT activity assays

(D, I). Data are mean ± SEM from three-four independent experiments. RT-qPCR quantitation of

relative HIV-1 gRNA and multiply spliced HIV-1 RNA copies from purified virions co-assembled

from tandem RRM (E) or single RRM (J) with WT Gag. Purified virions harvested from 293T co-

expressing tandem RRM (F) or single RRM (K) Gag chimera with WT Gag at the indicated

ratios were titered on Tzm-bl reporter cells. RT activity from purified virions were quantified to

assess particle release (G, L). Data are mean ± SEM of 2-3 independent experiments.

SUPPLEMENTAL FIGURE LEGENDS

Figure S1. Properties of Gag-RBD chimeras with heterologous RNA binding modules in place of NC. A-B. HEK293T cells were transfected with PR⁻ pNL4-3 proviral plasmids encoding the indicated Gag-RBD chimeras. Cell lysates (A) and purified virions (B) were harvested 2 days post-transfection and analyzed by immunoblotting for Gag, Gag-Pol (GP), Nef and Env (gp120). **C-D.** RT-qPCR quantification of relative copies of gRNA, multiply spliced viral RNAs, and 7SL RNA (7SL) from purified virions with (C, PR⁻) or without (D, PR⁺) an inactivation mutation in viral protease collected from 293T cells expressing the indicated Gag chimeras. For PR⁺ virions, copies of packaged RNA was normalized relative to virus particle numbers quantified based on RT activity. The columns represent the average of 3 independent experiments and the error bars represent SEM. Indicated statistical analyses (gRNA) were done by one-way ANOVA with Dunnet's multiple comparison test for correction, whereby (*) denotes p<0.05 comparisons. **E.** Northern blot analysis of unspliced (US), partially spliced (PS), multiply spliced (MS) HIV-1 RNAs and host 7SL RNA isolated from transfected cells and purified HIV-1 particles. **F.** Proportion of cellular reads, following removal of viral RNA-derived reads, that map to rRNA from total RNA-seq experiments using purified virions. Data show the average of 2-3 independent experiments and the error bars represent SEM. **G.** Classification of non-rRNA cellular RNA reads from virion total RNA-seq experiments as 7SL, mRNA, and Other.

Figure S2. Assembly and gRNA packaging properties of Gag-SRSF5 virions. A. Violin plots of MFI of Gag-YFP (left) and MS2-mCherry (right) measured for 60 WT, Gag-ΔNC, Gag-SRSF5 particles imaged in Figure 2B. **B.** WT and Gag-SRSF5 virions were purified from culture supernatants of transfected HEK293T cells and subjected to equilibrium density sedimentation following treatment with 0.5% Triton as detailed in Materials and Methods. Ten fractions (top-to-bottom) were analyzed by immunoblotting using a mouse monoclonal antibody against CA. **C.** MT4-GFP cells were infected with VSV-G pseudotyped HIV-1 at MOI of 0.5 or 1 in the presence

and absence nevirapine (NVP, 25 µM). At 24 hpi, cells were collected for DNA extraction as

detailed in Materials and Methods. Extracted DNA was analyzed by qPCR for accumulation of

early RT products (ERT), late RT products (LRT), 2-LTR circles (2-LTR), and integrated viral

DNA (Alu-PCR).

Figure S3. Trafficking and assembly properties of Gag-RBD chimeras. A. Representative

(n=3) immunoblot analysis of Na^+K^+ ATPase and β-tubulin from membrane (M) and cytoplasmic

(C) fractions isolated from 293T cells transfected with the indicated PR⁻ proviral plasmids.

Samples are the same as those shown in Fig. 3A. **B.** Representative confocal microscopy

images of HeLa cells transfected with PR⁻ proviral plasmids encoding the indicated Gag. Cells

were stained for Gag (red) and nuclei by DAPI (blue). **C.** Quantification of Gag-gRNA

colocalization rate from fluorescence imaging experiments shown in Figure 3C. **D.** Confocal

imaging of the bottom surface of cells transfected with two-color Gag-YFP-SRSF5/MS2-

mCherry or Gag-YFP-hnRNPA1/MS2-mCherry viruses. **E.** Images from live cell imaging of cells

expressing two-color Gag-YFP-hnRNPA1/MS2-mCherry virus.

Figure S4. Gag-SRSF5 gRNA packaging is not affected by deletions of structured

elements within the HIV-1 5' leader. A. Read density distribution on full-length HIV-1 gRNA (x-

axis) from CLIP-seq experiments in which PR⁻ WT Gag and Gag-RBD chimeras were

immunoprecipitated from transfected 293T cell lysates. **B.** Depiction of secondary structures

within the 5'-UTR of HIV-1 RNA (top) and table (bottom) showing sequences deleted in ΔΨ1-6

mutants used for this study. TAR, trans-activation response element; PBS, primer-binding site;

DIS, dimerization initiation signal; SD, major splice donor. **C-D.** RT-qPCR analysis of relative

copies of virion-incorporated HIV-1 gRNA and multiply spliced HIV-1 mRNA (C) and

immmunoblot analysis using a mouse monoclonal anti-HA antibody (D) of purified PR⁻ ΔΨ

virions. The columns in panel C represent the average of two independent experiments (each

done in two technical replicates representing independent transfections), and the error bars

represent SEM.

Figure S5. Analysis of the RNA binding properties of Gag-RBD chimeras at the PM and

immature virions. A-B. Autoradiogram of Gag-RNA complexes (top) and immunoblot analysis

of Gag proteins (bottom) immunoprecipitated from the PM fraction (A) or immature virions (B) in

CLIP experiments. (*) marks the IgG-heavy chain from the immunoprecipitating anti-HA

antibody. **C.** Frequency distribution of nucleotide occurrence (read density) in reads mapping to

the HIV-1 genome derived from CLIP experiments done in PM fractions.

Figure S6. Rational design of Gag-SRSF5 variants with reduced affinity for GGA and

enhanced affinity for CCG or GGG. A-B. Structural depiction of key residues involved in GGA

motif recognition in Gag-SRSF5 variants with reduced GGA affinity (A, mGGA1- and mGGA-2).

Residues of Gag-SRSF5 variants with enhanced affinity for CCG or GGG motif (mCCG-1,

mCCG-2, and mGGG) are shown in panel B. The amino acid positions shown are based on full-

length SRSF5.

Figure S7. Single RRM Gag chimeras bound RNA in cells, but do not package gRNA

efficiently. A-B. 293T cells were transfected with PR⁻ pNL4-3 proviral plasmids encoding single

RRM (e.g. M1, M2) Gag chimeras. Cell lysates and purified virions were harvested 2 days post-

transfection and analyzed with immunoblot for Gag and Gag-Pol (GP). Single RRM Gag-RNA

complexes were immunoprecipitated from cell lysates and analyzed using autoradiography and

immunoblotting. **C.** The relative titers of purified virions were determined via single-cycle

infection of TZM-bl reporter cells. **D.** RT-qPCR quantification of relative copies of gRNA, multiply

spliced HIV-1 RNA, and 7SL RNA from purified virions. Data are mean ± SEM from 4

independent experiments.

MATERIALS AND METHODS

Plasmids and cloning.

pNL4-3-derived plasmids containing three copies of a HA-tag in the stalk region of MA [20] and with (PR-) or without (PR+) an inactivating mutation in the protease (PR) gene were modified to insert the heterologous RNA-binding domains as follows. First, the NC coding sequence was replaced by a NotI restriction endonuclease site through conventional cloning using outward BssHII and SphI sites (see list below). Sequences coding for the first four and last three amino acids of NC were left intact to facilitate the proper proteolytic cleavage of the inserted heterologous RNA-binding domains. PCR amplicons containing individual or the complete set of RNA-binding domains from hnRNP and SR proteins were inserted into the Not I site (see list below). Chimeras bearing a subset of RRMs were also subcloned into a previously described two-color virus backbone [70] using SphI and SacII cut sites for microscopy analyses.

Deletions within the 5' leader were generated by overlap-extension PCR mutagenesis. The overlap PCR fragments were restriction digested and subsequently cloned into pNL4-3/MA-3xHA backbone via conventional cloning. ΔΨ5 was built on the ΔΨ1 backbone. ΔΨ3 and ΔΨ6 was built on the ΔΨ2 backbone. AatII/SphI fragment derived from these ΔΨ plasmids were then cloned into Gag-SRSF5 backbone. Gag-SRSF5 mutants (mGGA-1, mGGA-2, mCCG-1, mCCG-2, and mGGG) were generated using overlap-extension PCR mutagenesis, with PCR amplicons being cloned into the Not I site via ligation or Gibson Assembly. mGGA-1 and mGGA-2 overlap PCR amplicons were generated with 2 fragments; mCCG-1 and mCCG-2 were generated with 4 fragments; mGGG was generated with 3 fragments.

Primer pairs used in cloning are listed in the below table:

Primer	Sequence
ΔNC-NotI-F	ACTGAGCGGCCGCAATTTTAGGAACCAAAGAAAG
ΔNC-NotI-R	ACTGAGCGGCCGCCTCTCTCAGTACAATCTTTCATTTG
BssHII-F	ATGGCGCCCGAACAGGGACT
AatII-F	CCTGACGTCTAAGAAACCATTATTATC
SphI-R	GCCCTGCATGCACTGGATGCAATCTATCCCA
hnRNP A1-RRM1-F	ACTGAGCGGCCGCCAGCTGAGGAAGCTCTTCATTG
hnRNP A1-RRM1-R	ACTGAGCGGCCGCCTCTGGAGACAGCTCTCTTTGG
hnRNP A1-RRM2-F	ACTGAGCGGCCGCAAAAAGATATTTGTTGGTGGC
hnRNP A1-RRM2-R	ACTGAGCGGCCGCCCAGGGCTTTTCTAACTTCACAG
hnRNP F-RRM1-F	ACTGAGCGGCCGCGGCTTTGTGGTCAAGCTCCG
hnRNP F-RRM1-R	ACTGAGCGGCCGCCGGACTTGAACACCTCAATG
hnRNP F-RRM2-F	ACTGAGCGGCCGCATGGATTGGGTGTTGAAGC
hnRNP F-RRM2-R	ACTGAGCGGCCGCCCTGGCTGCTCTTAAACACCTC
hnRNP F-RRM3-F	ACTGAGCGGCCGCCACTGTGTCCACATGAGGG
hnRNP F-RRM3-R	ACTGAGCGGCCGCCTGAATTCAAGAAGAGTTCTAT
hnRNP K-KH1-F	ACTGAGCGGCCGCATGGTTGAATTACGCATTCTG
hnRNP K-KH1-R	ACTGAGCGGCCGCCGATTTTCTTCAGAATTTCTC
hnRNP K-KH2-F	ACTGAGCGGCCGCGACTGCGAGTTGAGGCTGTTG
hnRNP K-KH2-R	ACTGAGCGGCCGCCGATGATCTTTATGCACTCTAC
hnRNP K-KH3-F	ACTGAGCGGCCGCATTATTACTACACAAGTAAC
hnRNP K-KH3-R	ACTGAGCGGCCGCCCTGCAGCAAATACTGTGCAT
SRSF1-RRM1-F	ACTGAGCGGCCGCTGCCGCATCTACGTGGGTA
SRSF1-RRM1-R	ACTGAGCGGCCGCCGCTTCGAGGAAACTCCACCC

SRSF1-RRM2-F	ACTGAGCGGCCGCAACAGAGTGGTTGTCTCTGG
SRSF1-RRM2-R	ACTGAGCGGCCGCCATCAACTTTAACCCGGATGTA
SRSF3-RRM1-F	ACTGAGCGGCCGCTGTAAGGTTTATGTAGGCAATC
SRSF3-RRM1-R	ACTGAGCGGCCGCCACCATTCGACAGTTCCACTCT
SRSF4-RRM1-F	ACTGAGCGGCCGCCCGCGGGTGTACATCGGCC
SRSF4-RRM1-R	ACTGAGCGGCCGCCGCCGCGGGCATGCTCAACAA
SRSF4-RRM2-F	ACTGAGCGGCCGCTACAGACTTATTGTGGAGAA
SRSF4-RRM2-R	ACTGAGCGGCCGCCTGGCTTGTCTTCAACTAATCT
SRSF5-RRM1-F	ACTGAGCGGCCGCTGTCGGGTATTCATCGGGA
SRSF5-RRM1-R	ACTGAGCGGCCGCCAGCCCTAGCATGTTCAATAGT
SRSF5-RRM2-F	ACTGAGCGGCCGCAATCGTCTTATAGTTGAGA
SRSF5-RRM2-R	ACTGAGCGGCCGCCTTTGCTGCCTTCAATTAATTTTA
Fragment 1-F (mGGA-1, mGGA-2, mCCG-1, mCCG-2, mGGG)	TAATGATACAGAAAGGCAGCGGCCGCTGTCGGGTATTCATCGGG
Fragment 1-R (mGGA	GGCGGCGAGGGCCTGCCAGCTGACTCTTGAGGATAAATTCTC
Fragment 2-F (mGGA-1)	CAGCTGGCAGGCCCTCGCCGCCTTCATGAGACAAGCTGGGGAAG
Fragment 1-R (mGGA-2)	GTTCTGGAGGTTCTGCCAGCTGACTCTTGAGGATAAATTCTC
Fragment 2-F (mGGA-2)	CAGCTGGCAGAACCTCCAGAACTTCATGAGACAAGCTGGGGAAG
Fragment 1-R (mCCG-1, mCCG-2)	CTCTACAGGTGGAGCATTTCGTCTATCATTTCGAGGTCTG
Fragment 2-F (mCCG-1, mCCG-2)	ACGAAATGCTCCACCTGTAGAGAGAACAGAAAATCGTCTTATAG
Fragment 2- R (mCCG-1)	GGCGGCGAGCTTCTGCCAGCTGACTCTTGAGGATAAATTCTC
Fragment 3- F (mCCG-1)	CAGCTGGCAGAAGCTCGCCGCCTTCATGAGACAAGCTGGGGAAG

Fragment 3-R (mCCG-1, mCCG-2) & Fragment 2-R (mGGG)	CTCCTCCCCATTTATTTCCTTTCCAGAAAGTTTTTCAATAGC
Fragment 4-F (mCCG-1, mCCG-2) & Fragment 3-F (mGGG)	AAAGGAAATAAATGGGGAGGAGATAAAATTAATTGAAGGC
Fragment 2-R (mCCG-2) & Fragment 1-R(mGGG)	ATCTTTGAGCTTCTGCCAGCTGACTCTTGAGGATAAATTCTC
Fragment 3-F (mCGG-2) & Fragment 2-F (mGGG)	CAGCTGGCAGAAGCTCAAAGATTTCATGAGACAAGCTGGGGAAG
Fragment 2 (mGGA-1, mGGA-2) & Fragment 4 (mCCG-1, mCCG-2) & Fragment 3 (mGGG) (R)	CCCTAAAAAATTAGCCTGGCGGCCGCCTTTGCT

Cells, transfections, and viruses.

HEK293T (ATCC CRL-11268), HeLa (ATCC CCL-2) and HeLa-derived TZM-bL (NIH AIDS

Reagent Program) cells were maintained in Dulbecco's Modified Eagle Medium (DMEM)

supplemented with 10% fetal bovine serum (FBS) at 37°C. MT4-GFP indicator cells containing

an HIV-1 LTR-driven GFP reporter gene were described before[71] and grown in RPMI-10% FBS

supplemented with puromycin. For virus production, HEK293T cells were transfected with the

indicated proviral plasmid using polyethylenimine (PEI) and viral particles were harvested at 48

hours post transfection from filtered cell culture supernatants. HeLa cell transfections were

performed with either PEI or Lipofectamine 3000 (Thermo Fisher Scientific) reagent according

to the manufacturer's instructions.

Immunoblotting.

Cells were lysed in 1X radioimmunoprecipitation assay (RIPA) buffer and cleared lysates were

diluted in 1X final sodium dodecyl sulfate (SDS) sample buffer. Viruses were pelleted on a 20%

sucrose cushion prepared in 1X PBS and similarly resuspended in 1X sodium dodecyl sulfate

(SDS) sample buffer. Samples were separated by electrophoresis on Bolt 4–12% Bis-Tris Plus

gels (Life Technologies), transferred to nitrocellulose membranes and probed with the following

antibodies in Odyssey/Intercept Blocking Buffer (LI-COR): mouse monoclonal anti-CA (183-

H12-5C, NIH AIDS reagents), mouse monoclonal anti-HA (Biolegend, clone 16B12), goat

polyclonal anti-Env (ARP, 12-6205-1), rabbit polyclonal anti-Nef (NIH AIDS Reagents,

ARP2949), mouse monoclonal anti-β-tubulin (Santa Cruz, SC-5274), mouse monoclonal anti-

actin (Santa Cruz, SC-8432), mouse monoclonal anti-Na$^+$-K$^+$ ATPase (Santa Cruz, SC-21712).

Membranes were probed with fluorophore-conjugated secondary antibodies and visualized

using the LI-COR Odyssey system. Where indicated, Gag protein in cells and virions were

quantified using the LI-COR Image Studio software to normalize for CLIP protein:RNA adduct

signal.

RNA extraction and RT-qPCR.

Cell and virion associated RNA was harvested with TRIzol (Thermo Fisher Scientific) and

precipitated overnight in isopropanol. RNA was resuspended and reverse transcribed with the

High-Capacity cDNA Reverse Transcription Kit (Applied Biosystems) according to the

manufacturer's instructions. qPCR was performed using PowerUp 2X SYBR Green Master Mix

(Applied Biosystems). Following primers were used to quantify their respective targets:

Primer	Sequence
Unspliced/Gag-Pol-F	TTCTTCAGAGCAGACCAGAGC

Unspliced/Gag-Pol-R	GCTGCCAAAGAGTGATCTGA
Multiply spliced-F	TCTATCAAAGCAACCCACCTC
Multiply spliced-R	CGTCCCAGATAAGTGCTAAGG
7SL RNA-F	CACCAGGTTGCCTAAGGAGGGG
7SL RNA-R	TCCCACTACTGATCAGCACGGG
GAPDH-F	AGGTGAAGGTCGGAGTCAACG
GAPDH-R	GGTCATTGATGGCAACAATATCCACTTTAC
RPL36A-F	GCACCAACCCCATAAAGTGAC
RPL36A-R	CTGGGCGTACAGAGAATCCT
RPS6-F	TGGACGATGAACGCAAACTTC
RPS6-R	TTCGGACCACATAACCCTTCC
RPS16-F	TCGGACGCAAGAAGACAGC
RPS16-R	AGCAGCTTGTACTGTAGCGTG

Quantification of reverse transcription and integration

MT4-GFP cells were grown in 24-well plates and infected with VSV-G pseudotyped WT or Gag

chimera viruses at an MOI of 0.5 or 1 in the presence of polybrene. Cells were collected at 24

hpi, washed with PBS, and resuspended in 200 μL lysis buffer (100 mM NaCl, 10 mM Tris-HCl

pH 8, 25 mM EDTA pH 8, 0.5% SDS, 0.1 mg/mL Proteinase K). Samples were incubated at

50°C by constant agitation (2,000 rpm) in a thermal mixer (Eppendorf) for 2 h. DNA was

extracted by phenol-chloroform extraction. Extracted DNA was analyzed for accumulation of RT

products, 2-LTR circles, and integration as described before [72,73].

Northern blotting

For northern blotting experiments, 293T cells were transiently transfected with 5 μg of each

plasmid using a 4:1 ratio of PEI and 150 mM NaCl. Media was replaced 24- and 44-hours post-

infection. 4 hours after the 44-hour media change, extracellular viral particles were harvested by

passing culture supernatants through a 0.22 um filter. Particles were then concentrated via

ultracentrifugation through a 20% sucrose cushion. RNA was isolated from transfected cells and

concentrated virions using Trizol Reagent. Northern blot analysis was performed using a

modified version of a previously described protocol. Briefly, RNA samples were incubated in 1x

MOPS electrophoresis buffer, 8% formaldehyde, 50% formamide, and gel loading buffer and

then heated at 85°C for 10 minutes. The denatured samples were loaded into the wells of a 1%

agarose gel containing 2.2M formaldehyde, and electrophoresis proceeded for approximately 7

hours at 50V. RNA species were then vertically transferred to a nylon membrane. Following

transfer, the membrane was exposed to UV to induce crosslinking. The membrane was then

incubated in prehybridization buffer (50 mM Tris, pH 7.5, 120 µg/mL salmon sperm DNA) at

55°C. After 4 hours, a ^{32}P-labelled probe complementary to the viral genome or 7SL RNA was

added to the prehybridization buffer and allowed to hybridize with the membrane at 55°C

overnight. The membrane was washed the next day and then exposed overnight in a

phosphorimaging cassette. The screen was imaged using an Amersham™ Typhoon™ scanner.

Membrane Flotation Assays.

Membrane flotation assays were performed as previously described [33]. HEK293T cells grown in

10 cm plates were transfected with 10µg of the indicated proviral plasmids using PEI and cell

culture media was replaced 6-8 hours post-transfection. 24 hours post transfection, cells were

collected, washed once with 1X STE buffer (100 mM NaCl, 10 mM Tris-Cl pH 8.0, 1 mM EDTA).

Afterwards, cells were resuspended in 500µl of hypotonic solution (10mM Tris [pH 7.4], 10mM

KCl, 1 mM EDTA) supplemented with Complete Protease Inhibitor Cocktail (Roche), and placed

on ice for 15 minutes. Cells were then lysed by dounce homogenization and centrifuged at 1000

x g for 5 minutes, 4°C. 350µl of post-nuclear supernatant was collected and mixed with 1.65 mL

of 90% sucrose solution. 6.5 mL of 65% sucrose solution followed by 2.5 mL of 10% sucrose

solution were layered on top of the mixture. Centrifugation was then performed with a Beckman

SW41 Ti rotor at 35,000 rpm for 4-18 hours, 4°C. 1 mL fractions were collected, with the 3rd

fraction and 11th fraction from top corresponding to the membrane and cytosol, respectively.

100 µL of each fraction was kept for RNA analysis while the remaining 900 µL was used for

protein precipitation. Aliquots for RNA analysis were treated with Proteinase K for 10 min at

37°C and RNA was extracted with an equal volume of phenol:chloroform:isoamyl alcohol

(125:24:1) (Sigma-Aldrich) and precipitated with the addition of 1 µL Glycoblue (Ambion), 0.25

volume of 3M NaOAc pH 5.2, and 2.5 volumes of 1:1 ethanol:isopropanol. The resulting RNA

samples were treated with DNase, reextracted using one more round of phenol:chloroform

extraction and used for RT-qPCR. Protein was precipitated in 10% trichloroacetic acid (TCA) for

2-4 hours and pelleted by centrifugation. Protein pellets were washed twice with 10% TCA and

once with ice cold acetone before being resuspended in SDS-PAGE sample buffer and used for

western blot analyses.

Equilibrium density sedimentation of virion core components in vitro

Equilibrium density sedimentation of virion core components was performed as previously

described [74]. Briefly, HEK293T cells grown on 10-cm dishes were transfected with the above-

mentioned plasmids. 48 hours post-transfection, cell-free virions collected from cell culture

supernatants were pelleted through a 20% sucrose cushion. Pelleted virions were resuspended

in 1X PBS and treated with 0.5% Triton X-100 for 2 minutes at room temperature. Immediately

after, samples were layered on top of a 30-70% linear sucrose gradients in 1X STE buffer and

centrifuged for 16 hours at 4°C and 28,500 rpm using an SW55Ti rotor. 500 µL fractions

collected from the top were analyzed by immunoblotting using a mouse monoclonal anti-HIV

p24 antibody (183-H12-5C, NIH AIDS reagents) and Q-PCR-based assays for RT activity [75].

Transmission electron microscopy.

To visualize virus assembly sites in cells, HEK293T cells were grown in 10 cm plates and

transfected with 10µg of proviral plasmid as previously described. 24 h post-transfection, cells

were harvested and fixed in 2% paraformaldehyde/2.5% glutaraldehyde (Ted Pella Inc.,

Redding, CA) in 100mM cacodylate buffer, pH 7.2 for 2 h at room temperature. Samples were

washed in cacodylate buffer and postfixed in 1% osmium tetroxide (Ted Pella Inc.)/1.5%

potassium ferricyanide (Sigma, St. Louis, MO) for 1 h. Samples were then rinsed extensively in

dH20 prior to en bloc staining with 1% aqueous uranyl acetate (Ted Pella Inc.) for 1 h. Following

several rinses in water, samples were dehydrated in a graded series of ethanol and embedded

in Eponate 12 resin (Ted Pella Inc.). Ultrathin sections of 95nm were cut with a Leica Ultracut

UCT ultramicrotome (Leica Microsystems Inc., Bannockburn, IL), stained with uranyl acetate

and lead citrate. To visualize viral particles, HEK293T cells grown in 15-cm plates were

transfected with 30 µg of the indicated plasmid. Cell free virions were harvested from the

supernatant 48 h post transfection, filtered through 0.22 µm filters and pelleted through a 20%

sucrose cushion via centrifugation using a Beckman SW32-Ti rotor at 28,000 rpm for 1.5 h, 4°C.

Virion pellets were fixed and processed for EM as described above. All samples were viewed on

a JEOL 1200 EX transmission electron microscope (JEOL USA Inc., Peabody, MA) equipped

with an AMT 8-megapixel digital camera and AMT Image Capture Engine V602 software

(Advanced Microscopy Techniques, Woburn, MA).

Immunofluorescence microscopy.

HeLa cells were plated in 24-well glass-bottom dishes (Mattek Corporation, Ashland, MA, USA)

coated with poly-L-lysine (50,000 cells/well) and transfected with 500ng of the indicated proviral

plasmid using either PEI or Lipofectamine 3000 transfection reagent. At 24 h post-transfection

cells were fixed with 4% paraformaldehyde in PBS, for 10 min and, for immunofluorescence-

based experiments, permeabilized with 0.1% Triton X-100 in PBS for 10 minutes. Permeabilized

cells were blocked with 5% goat serum in PBS-T (PBS, 0.1% Tween 20) blocking solution for 30

min at room temperature. Cells were probed with a mouse monoclonal anti-HA (Biolegend,

clone 16B12) in blocking solution for 1-2 h at room temperature, followed by goat anti-mouse

Alexa Fluor 594 secondary antibody (1:1000, Thermo Fisher Scientific) for 1 h at room

temperature. Nuclei were stained with DAPI for 10 min. Cells were visualized with a Zeiss

LSM880 Confocal Microscope with Airyscan, oil immersion x63 or A1R Confocal Microscope, oil

immersion x60.

Single virion analysis

To generate labeled, two-colored virus-like particles for single virion analysis, approximately

500,000 HEK293T cells were plated in each well of a six-well dish and transfected with plasmids

encoding wild-type, SRSF5 RBD, or ΔNC two-color self-tagging viruses using PEI. The media

was exchanged at 24 h post-transfection. Virus particle-containing supernatants were harvested

at 48 h post-transfection, filtered through a 0.22-µm filter, and centrifuged through 20% sucrose

for 2 h at 18,213 rpm. The medium was discarded after centrifugation, and concentrated viral

particles were resuspended in 1× PBS, plated in a 24-well glass-bottom dish (Cellvis, Mountain

View, CA), and left overnight at 4°C to allow virus particles to settle down on the glass wells.

Microscopy was performed using a Nikon Ti-Eclipse inverted wide-field microscope (Nikon

Corp, Minato, Tokyo, Japan) using a 100× Plan Apo oil objective lens (numerical aperture [NA]

1.45). Cell and virion images were captured using an ORCA—Flash4.0 CMOS camera

(Hamamatsu Photonics, Skokie, IL, USA) and the following excitation/emission filter sets:

510/535 nm (YFP) and 585/610 nm (mCherry). All images were processed and analyzed using

FIJI/ImageJ2. FIJI/ImageJ2's built-in tools were used to measure the mean fluorescent

intensities (MFIs) of both the virions and the background by selecting regions of interest in the

Gag-YFP channel and the corresponding areas in the MS2–mCherry (unspliced RNA) channel.

Virion background subtracted MFIs for MS2–mCherry and Gag–YFP channels were plotted

using GraphPad Prism (version 10.1.2). The data were plotted for MS2–mCherry MFI, Gag–

YFP MFI, and MS2–mCherry/Gag–YFP MFI.

RNA-seq.

HEK293T cells were grown in 24-well plates and transfected with 400ng of the indicated proviral

plasmid using PEI. 48 h post transfection, cell culture media was collected and filtered through

0.22 μm filters, before being pelleted through a 20% sucrose cushion at 15,000 rpm for 1.5 h,

4°C. RNA was harvested from purified cell free virions with TRIzol as previously described. RNA

samples were prepared for sequencing using the TruSeq Stranded Total RNA Sample Prep Kit

(Illumina) according to the manufacturer's instructions but without rRNA depletion and as

detailed previously [76]. Barcoded samples were pooled and sequenced using an Illumina

NextSeq platform (1x75) at the Center for Genome Sciences at Washington University.

Photoactivatable ribonucleoside-enhanced cross-linking and immunoprecipitation (PAR-CLIP).

PAR-CLIP experiments were conducted as previously described with some modifications [20,77,78].

HEK293T cells were grown in 10-cm plates and transfected with 10 μg of the indicated proviral

plasmid. Cell culture media was replaced 6-8 h post-transfection, and 4-thiouridine (4SU) was

added to the cell culture medium at a concentration of 100 μM 16 h before harvesting cells and

virions. 48 hours post-transfection, virions and cells were collected. Virions in the cell culture

supernatant were filtered through 0.22 μm filters and pelleted through a 20% sucrose cushion at

28,000 rpm for 1.5 hr, 4°C. Transfected cells were harvested with a cell scraper. Both cells and

purified virions were resuspended in PBS and protein-RNA complexes UV-crosslinked (365nm)

before being lysed in 1X RIPA lysis buffer. Cell and virus lysates were then treated with RNase

A (Thermo Fisher Scientific) and DNase I (Roche) at a final concentration of 20and 60 U/mL,

respectively. HA-tagged Gag proteins were immunoprecipitated using mouse monoclonal anti-

HA antibody conjugated Protein G-coupled Dynabeads (Thermo Fisher Scientific), as previously

described. Gag associated RNAs were radiolabeled with γ-^{32}P-ATP using T4 Polynucleotide

Kinase (New England Biolabs). The protein-RNA adducts were separated via SDS-PAGE and

transferred onto nitrocellulose membranes before being visualized with autoradiography film.

Protein-RNA adducts were excised from nitrocellulose membranes and digested with

Proteinase K, and RNA was extracted with phenol:chloroform:isoamyl alcohol (125:24:1)

(Sigma-Aldrich). Resultant RNA was pelleted with the addition of 1μL Glycoblue (Ambion), 0.25

volume of 3M NaOAc pH 5.2, and 2.5 volumes of 1:1 ethanol:isopropanol. The RNA pellet was

washed with 80% ethanol, dried and dissolved in 13 μL RNase-free H2O. RNA was ligated with

barcoded adapters and ligation products were separated and purified on Novex 15% TBE-Urea

gel (Invitrogen) as previously described with some exceptions [77]. Here we used barcoded 3'

adapters and thus samples were pooled into groups of two prior to 5' adapter ligation.

Furthermore, the amount of 3' adapter added to each reaction was reduced to 5pmole. An end-

labeled 15-nucleotide RNA oligonucleotide was prepared in parallel as positive control for

adapter ligations. The resulting ligation products were reverse transcribed using Superscript III

First Strand Synthesis Kit (Invitrogen), and cDNA was amplified using Phusion Polymerase

(New England Biolabs). PCR products were gel purified and sequenced on a NextSeq platform

(1x75bp) at the Center for Genome Sciences at Washington University.

3' barcoded adapters	Sequence
CLIP1	5'adenylated/NNTGACTG TGG AAT TCT CGG GTG CCA AGG-3'dideoxyC
CLIP2	5'adenylated/NNACACTCTGG AAT TCT CGG GTG CCA AGG-3'dideoxyC
CLIP3	5'adenylated/NNACAGAG TGG AAT TCT CGG GTG CCA AGG-3'dideoxyC
CLIP4	5'adenylated/NNGCGATA TGG AAT TCT CGG GTG CCA AGG-3'dideoxyC

CLIP31	5′adenylated/NN<u>TAGCGA</u> TGG AAT TCT CGG GTG CCA AGG-3′dideoxyC
CLIP32	5′adenylated/NN<u>CTGTAG</u> TGG AAT TCT CGG GTG CCA AGG-3′dideoxyC
CLIP33	5′adenylated/NN<u>TAGTCG</u> TGG AAT TCT CGG GTG CCA AGG-3′dideoxyC
CLIP34	5′adenylated/NN<u>AGTGTC</u> TGG AAT TCT CGG GTG CCA AGG-3′dideoxyC
CLIP35	5′adenylated/NN<u>ATCGAC</u> TGG AAT TCT CGG GTG CCA AGG-3′dideoxyC
CLIP36	5′adenylated/NN<u>GACATG</u> TGG AAT TCT CGG GTG CCA AGG-3′dideoxyC
CLIP37	5′adenylated/NN<u>GACTAC</u> TGG AAT TCT CGG GTG CCA AGG-3′dideoxyC
CLIP38	5′adenylated/NN<u>ATCTCG</u> TGG AAT TCT CGG GTG CCA AGG-3′dideoxyC
5' adapter	rGUU CAG AGU UCU ACA GUC CGA CGA UC
RT primer	GCCTTGGCACCCGAGAATTCCA
cDNA PCR (F)	AATGATACGGCGACCACCGAGATCTACACGTTCAGAGTTCTACAGTCCGA
cDNA PCR (R)	CAAGCAGAAGACGGCATACGAGATCGTGATGTGACTGGAGTTCCTTGGCACCCGAGAATTCCA
Control oligo	rAUAGCUACGAUUGCA

Bioinformatic analysis of NGS data

Clip-seq:

<u>Mapping:</u>

Adapter trimming was performed using BBDuk from BBTools version 39.06. The 3′ adapter

sequences were removed with parameters k=8 and mink=7 to define the k-mer size and

minimum k-mer size. Additional parameters included ml=18 and maxlength=52 to filter reads

based on length. The rcomp function was disabled to only consider kmers in the forward

orientation.

Trimmed reads were further processed with BBDuk for separation based on their 3′ barcode

sequences. Parameters included a k-mer size of 8, maskmiddle=f to ensure only exact matches

were trimmed, and copyundefined was set to handle undefined bases. The resulting FASTQ

files are the trimmed reads specific to each barcode.

Next, the trimmed reads were mapped to their corresponding Gag chimera viral genomes and

WT viral genome (HIV-1$_{NL4-3}$/PR⁻) with Bowtie version 1.3.1. Uniquely mapped reads were

reported, allowing one mismatch (-v 1 -m 1). Mapped reads were processed with the mpileup

function form SAMtools to generate pileups for HIV-1$_{NL4-3}$/PR⁻, and these pileups were then

analyzed using customized in-house scripts and visualized as viral density maps with GraphPad

Prism (v6). Reads that failed to map to their corresponding viral genomes were extracted using

SAMtools (-b -f 4) and then aligned to the hg19 human genome with Bowtie, allowing up to 10

multi-mapped reads with one mismatch. Various file formats of mapped reads were generated,

including BED and SAM files, for further analysis.

<u>Cluster Generation by PARalyzer and Cluster Annotations</u>

PARalyzer was used for peak calling using SAM or BOWTIE outputs to generate clusters, which

represent binding sites derived from overlapping mapped reads. Clusters were annotated using

publicly available databases, ENSEMBL v72 and tRNASCANse [79].

For a cluster or a read that mapped to a location with multiple gene features (i.e., intron-exon or

coding-sequence-3′ UTR junctions), we required greater than half the length of the read/cluster

to map to the assigned class. In other cases where a cluster/read could be assigned to multiple

gene features due to the presence of alternatively spliced isoforms in the ENSEMBL database,

the following order of preference was chosen: coding sequence, 5′UTR, 3′UTR, intron. In some

experiments, the number of reads associated with each cluster was taken as a measurement of

how frequently a given cluster (i.e., binding site) was bound by Gag. To identify sequence motifs

within mRNAs, PARalyzer parameters were adjusted to generate clusters with a mean length of

~10 nucleotides surrounding the crosslinking site and which overlapped > 95% with the clusters

generated above. Perl scripts were written for comparison of clusters in two separate data sets,

determining the nucleotide composition of clusters and identifying features of mRNA molecules

(i.e., 5'UTR, 3'UTR, CDS length) bound by Gag.

<u>Correlation Analysis</u>

Cross-correlation analysis was performed to measure the similarity of Gag binding frequencies

between two datasets (Oppenheim & Schafer, 1975). The correlation function (CF(s)) compared

the signals by shifting one dataset relative to the other and calculating their normalized

correlation. The search window for correlation analysis was set to 400 nucleotides.

To assess the statistical significance of observed correlations, Monte Carlo simulations were

used to generate 10,000 randomized datasets with the same structure as the original (i.e.,

identical numbers of binding sites and reads per site). Confidence intervals were calculated for

p-values of 0.05 based on the cross-correlation results of these randomized samples.

<u>Motif analysis</u>

The bed files were annotated and reads in the mRNA regions were extracted using in-house

scripts. These extracted reads were then converted to FASTA format using BEDTools for the

subsequent analysis.

Motif analysis was conducted using MEME Suite with the following parameters: -nmotifs 10 -

minw 5 -maxw 20 -dna. The parameters specified the identification of up to 10 motifs with a

minimum width of 5 and a maximum width of 20 base pairs, applied to DNA sequences.

RNA-seq:

The STAR aligner was used to map the RNA-seq data to human genome indices (hg19) with a

mismatch rate threshold of 4%. Mapped reads were than annotated with featureCounts.

All CLIP-seq and RNA-seq data were analyzed using publicly available programs (BBDuk

toolkit, Bowtie, and SAMtools) and in-house scripts [20,77]. Briefly, reads that contained ambiguous

nucleotides, did not contain the 3′ adapter sequence or were shorter than 10nt were discarded.

The remaining reads were then separated based on their 3′ barcode sequences and collapsed

to generate a set of unique sequences. Reads were then aligned to the human genome (hg19)

or to corresponding viral genomes, with up to 1 mismatch allowed using the Bowtie algorithm [80].

In the case of reads that mapped to the human genome, locations were only reported for those

with the minimum number of observed mismatches for each read (Bowtie criteria: -v 1 -m 10 for

mapping to hg19, and - v 1 -m 1 for mapping to the viral genomes). SAMtools was used to

generate pileups of mapped reads, which were subsequently processed using in-house scripts

and visualized by Prism GraphPad software (v6) [81].

ACKNOWLEDGEMENTS

We thank Dr. Wandy Beatty and the Molecular Microbiology Imaging Facility at the Washington

University School of Medicine for their assistance with confocal microscopy and transmission

electron microscopy experiments. H.R.V. was supported by the National Science Foundation

Graduate Research Fellowship under Grant No. DGE-1745038, DGE-2139839 and the National

Institute of Allergy and Infectious Diseases of the National Institutes of Health under Award

Number T32AI007172. This work was supported by NIH grants U54 AI170660 (A.T., S.B.K.,

N.S.), R01 AI179691 (S.B.K., N.S.) and R56 AI110221 (N.S.).

CLIP-seq and RNA-seq data were deposited to the Gene Expression Omnibus Database with

accession numbers GSE281422, GSE281423, and GSE281424.

REFERENCES

1.   Chen, J., Nikolaitchik, O., Singh, J., Wright, A., Bencsics, C.E., Coffin, J.M., Ni, N., Lockett,
S., Pathak, V.K., and Hu, W.S. (2009). High efficiency of HIV-1 genomic RNA packaging
and heterozygote formation revealed by single virion analysis. Proc Natl Acad Sci U S A
106, 13535-13540. 10.1073/pnas.0906822106.
2.   Moore, M.D., and Hu, W.S. (2009). HIV-1 RNA dimerization: It takes two to tango. AIDS
Rev *11*, 91-102.
3.   Duchon, A., and Hu, W.S. (2024). HIV-1 RNA genome packaging: it's G-rated. mBio *15*,
e0086123. 10.1128/mbio.00861-23.
4.   Hu, W.S., and Temin, H.M. (1990). Genetic consequences of packaging two RNA
genomes in one retroviral particle: pseudodiploidy and high rate of genetic
recombination. Proc Natl Acad Sci U S A *87*, 1556-1560. 10.1073/pnas.87.4.1556.
5.   Rawson, J.M.O., Nikolaitchik, O.A., Keele, B.F., Pathak, V.K., and Hu, W.S. (2018).
Recombination is required for efficient HIV-1 replication and the maintenance of viral
genome integrity. Nucleic Acids Res *46*, 10535-10545. 10.1093/nar/gky910.
6.   Streeck, H., Li, B., Poon, A.F., Schneidewind, A., Gladden, A.D., Power, K.A., Daskalakis,
D., Bazner, S., Zuniga, R., Brander, C., et al. (2008). Immune-driven recombination and
loss of control after HIV superinfection. J Exp Med *205*, 1789-1796.
10.1084/jem.20080281.
7.   Kellam, P., and Larder, B.A. (1995). Retroviral recombination can lead to linkage of
reverse transcriptase mutations that confer increased zidovudine resistance. J Virol *69*,
669-674. 10.1128/jvi.69.2.669-674.1995.
8.   Vijaykrishna, D., Mukerji, R., and Smith, G.J. (2015). RNA Virus Reassortment: An
Evolutionary Mechanism for Host Jumps and Immune Evasion. PLoS Pathog *11*,
e1004902. 10.1371/journal.ppat.1004902.
9.   McDonald, S.M., Matthijnssens, J., McAllen, J.K., Hine, E., Overton, L., Wang, S., Lemey,
P., Zeller, M., Van Ranst, M., Spiro, D.J., and Patton, J.T. (2009). Evolutionary dynamics
of human rotaviruses: balancing reassortment with preferred genome constellations.
PLoS Pathog *5*, e1000634. 10.1371/journal.ppat.1000634.
10. Nelson, M.I., Viboud, C., Simonsen, L., Bennett, R.T., Griesemer, S.B., St George, K.,
Taylor, J., Spiro, D.J., Sengamalay, N.A., Ghedin, E., et al. (2008). Multiple reassortment
events in the evolutionary history of H1N1 influenza A virus since 1918. PLoS Pathog *4*,
e1000012. 10.1371/journal.ppat.1000012.
11. Ghedin, E., Fitch, A., Boyne, A., Griesemer, S., DePasse, J., Bera, J., Zhang, X., Halpin,
R.A., Smit, M., Jennings, L., et al. (2009). Mixed infection and the genesis of influenza
virus diversity. J Virol *83*, 8832-8841. 10.1128/jvi.00773-09.
12. Komoto, S., Tacharoenmuang, R., Guntapong, R., Ide, T., Tsuji, T., Yoshikawa, T.,
Tharmaphornpilas, P., Sangkitporn, S., and Taniguchi, K. (2016). Reassortment of Human
and Animal Rotavirus Gene Segments in Emerging DS-1-Like G1P[8] Rotavirus Strains.
PLOS ONE *11*, e0148416. 10.1371/journal.pone.0148416.
13. Aldovini, A., and Young, R.A. (1990). Mutations of RNA and protein sequences involved
in human immunodeficiency virus type 1 packaging result in production of noninfectious
virus. J Virol *64*, 1920-1926. 10.1128/jvi.64.5.1920-1926.1990.

14. Gorelick, R.J., Nigida, S.M., Jr., Bess, J.W., Jr., Arthur, L.O., Henderson, L.E., and Rein, A.
(1990). Noninfectious human immunodeficiency virus type 1 mutants deficient in
genomic RNA. J Virol *64*, 3207-3211. 10.1128/jvi.64.7.3207-3211.1990.
15. Kuzembayeva, M., Dilley, K., Sardo, L., and Hu, W.S. (2014). Life of psi: how full-length
HIV-1 RNAs become packaged genomes in the viral particles. Virology *454-455*, 362-370.
10.1016/j.virol.2014.01.019.
16. Rein, A. (2019). RNA Packaging in HIV. Trends Microbiol *27*, 715-723.
10.1016/j.tim.2019.04.003.
17. Nikolaitchik, O.A., Somoulay, X., Rawson, J.M.O., Yoo, J.A., Pathak, V.K., and Hu, W.S.
(2020). Unpaired Guanosines in the 5' Untranslated Region of HIV-1 RNA Act
Synergistically To Mediate Genome Packaging. J Virol *94*. 10.1128/jvi.00439-20.
18. De Guzman, R.N., Wu, Z.R., Stalling, C.C., Pappalardo, L., Borer, P.N., and Summers, M.F.
(1998). Structure of the HIV-1 nucleocapsid protein bound to the SL3 psi-RNA
recognition element. Science *279*, 384-388. 10.1126/science.279.5349.384.
19. Ding, P., Kharytonchyk, S., Waller, A., Mbaekwe, U., Basappa, S., Kuo, N., Frank, H.M.,
Quasney, C., Kidane, A., Swanson, C., et al. (2020). Identification of the initial
nucleocapsid recognition element in the HIV-1 RNA packaging signal. Proc Natl Acad Sci
U S A *117*, 17737-17746. 10.1073/pnas.2008519117.
20. Kutluay, S.B., Zang, T., Blanco-Melo, D., Powell, C., Jannain, D., Errando, M., and
Bieniasz, P.D. (2014). Global changes in the RNA binding specificity of HIV-1 gag regulate
virion genesis. Cell *159*, 1096-1109. 10.1016/j.cell.2014.09.057.
21. Clever, J.L., and Parslow, T.G. (1997). Mutant human immunodeficiency virus type 1
genomes with defects in RNA dimerization or encapsidation. J Virol *71*, 3407-3414.
10.1128/JVI.71.5.3407-3414.1997.
22. Laham-Karam, N., and Bacharach, E. (2007). Transduction of human immunodeficiency
virus type 1 vectors lacking encapsidation and dimerization signals. J Virol *81*, 10687-
10698. 10.1128/JVI.00653-07.
23. McBride, M.S., and Panganiban, A.T. (1997). Position dependence of functional hairpins
important for human immunodeficiency virus type 1 RNA encapsidation in vivo. J Virol
71, 2050-2058. 10.1128/JVI.71.3.2050-2058.1997.
24. Comas-Garcia, M., Datta, S.A., Baker, L., Varma, R., Gudla, P.R., and Rein, A. (2017).
Dissection of specific binding of HIV-1 Gag to the 'packaging signal' in viral RNA. Elife *6*.
10.7554/eLife.27055.
25. Rulli, S.J., Hibbert, C.S., Mirro, J., Pederson, T., Biswal, S., and Rein, A. (2007). Selective
and Nonselective Packaging of Cellular RNAs in Retrovirus Particles. Journal of Virology
81, 6623-6631. doi:10.1128/jvi.02833-06.
26. Muriaux, D., Mirro, J., Harvin, D., and Rein, A. (2001). RNA is a structural element in
retrovirus particles. Proc Natl Acad Sci U S A *98*, 5246-5251. 10.1073/pnas.091000398.
27. Kutluay, S.B., and Bieniasz, P.D. (2010). Analysis of the initiating events in HIV-1 particle
assembly and genome packaging. PLoS Pathog *6*, e1001200.
10.1371/journal.ppat.1001200.
28. Jouvenet, N., Simon, S.M., and Bieniasz, P.D. (2009). Imaging the interaction of HIV-1
genomes and Gag during assembly of individual viral particles. Proc Natl Acad Sci U S A
106, 19114-19119. 10.1073/pnas.0907364106.

29.   Sardo, L., Hatch, S.C., Chen, J., Nikolaitchik, O., Burdick, R.C., Chen, D., Westlake, C.J.,
Lockett, S., Pathak, V.K., and Hu, W.S. (2015). Dynamics of HIV-1 RNA Near the Plasma
Membrane during Virus Assembly. J Virol *89*, 10832-10840. 10.1128/jvi.01146-15.
30.   Chen, J., Rahman, S.A., Nikolaitchik, O.A., Grunwald, D., Sardo, L., Burdick, R.C., Plisov,
S., Liang, E., Tai, S., Pathak, V.K., and Hu, W.S. (2016). HIV-1 RNA genome dimerizes on
the plasma membrane in the presence of Gag protein. Proc Natl Acad Sci U S A *113*,
E201-208. 10.1073/pnas.1518572113.
31.   Freed, E.O. (2015). HIV-1 assembly, release and maturation. Nat Rev Microbiol *13*, 484-
496. 10.1038/nrmicro3490.
32.   Ivanchenko, S., Godinez, W.J., Lampe, M., Kräusslich, H.-G., Eils, R., Rohr, K., Bräuchle,
C., Müller, B., and Lamb, D.C. (2009). Dynamics of HIV-1 Assembly and Release. PLOS
Pathogens *5*, e1000652. 10.1371/journal.ppat.1000652.
33.   Kutluay, S.B., and Bieniasz, P.D. (2010). Analysis of the Initiating Events in HIV-1 Particle
Assembly and Genome Packaging. PLOS Pathogens *6*, e1001200.
10.1371/journal.ppat.1001200.
34.   Muriaux, D., Mirro, J., Harvin, D., and Rein, A. (2001). RNA is a structural element in
retrovirus particles. Proceedings of the National Academy of Sciences *98*, 5246-5251.
doi:10.1073/pnas.091000398.
35.   Duchon, A., Santos, S., Chen, J., Brown, M., Nikolaitchik, O.A., Tai, S., Chao, J.A., Freed,
E.O., Pathak, V.K., and Hu, W.-S. (2021). Plasma Membrane Anchoring and Gag:Gag
Multimerization on Viral RNA Are Critical Properties of HIV-1 Gag Required To Mediate
Efficient Genome Packaging. mBio *12*, e03254-03221. doi:10.1128/mbio.03254-21.
36.   Lei, X., Gonçalves-Carneiro, D., Zang, T.M., and Bieniasz, P.D. (2023). Initiation of HIV-1
Gag lattice assembly is required for recognition of the viral genome packaging signal.
eLife *12*, e83548. 10.7554/eLife.83548.
37.   Kypr, J., and Mrazek, J. (1987). Unusual codon usage of HIV. Nature *327*, 20.
10.1038/327020a0.
38.   van der Kuyl, A.C., and Berkhout, B. (2012). The biased nucleotide composition of the
HIV genome: a constant factor in a highly variable virus. Retrovirology *9*, 92.
10.1186/1742-4690-9-92.
39.   Accola, M.A., Strack, B., and Göttlinger, H.G. (2000). Efficient particle production by
minimal Gag constructs which retain the carboxy-terminal domain of human
immunodeficiency virus type 1 capsid-p2 and a late assembly domain. J Virol *74*, 5395-
5402. 10.1128/jvi.74.12.5395-5402.2000.
40.   Han, Siew P., Tang, Yue H., and Smith, R. (2010). Functional diversity of the hnRNPs:
past, present and perspectives. Biochemical Journal *430*, 379-392. 10.1042/bj20100396.
41.   Jeong, S. (2017). SR Proteins: Binders, Regulators, and Connectors of RNA. Mol Cells *40*,
1-9. 10.14348/molcells.2017.2319.
42.   Sertznig, H., Hillebrand, F., Erkelenz, S., Schaal, H., and Widera, M. (2018). Behind the
scenes of HIV-1 replication: Alternative splicing as the dependency factor on the quiet.
Virology *516*, 176-188. 10.1016/j.virol.2018.01.011.
43.   Huelga, S.C., Vu, A.Q., Arnold, J.D., Liang, T.Y., Liu, P.P., Yan, B.Y., Donohue, J.P., Shiue,
L., Hoon, S., Brenner, S., et al. (2012). Integrative genome-wide analysis reveals

cooperative regulation of alternative splicing by hnRNP proteins. Cell Rep *1*, 167-178.
10.1016/j.celrep.2012.02.001.

44. Pandit, S., Zhou, Y., Shiue, L., Coutinho-Mansfield, G., Li, H., Qiu, J., Huang, J., Yeo, G.W.,
Ares, M., Jr., and Fu, X.D. (2013). Genome-wide analysis reveals SR protein cooperation
and competition in regulated splicing. Mol Cell *50*, 223-235.
10.1016/j.molcel.2013.03.001.

45. Kutluay, S.B., Emery, A., Penumutchu, S.R., Townsend, D., Tenneti, K., Madison, M.K.,
Stukenbroeker, A.M., Powell, C., Jannain, D., Tolbert, B.S., et al. (2019). Genome-Wide
Analysis of Heterogeneous Nuclear Ribonucleoprotein (hnRNP) Binding to HIV-1 RNA
Reveals a Key Role for hnRNP H1 in Alternative Viral mRNA Splicing. J Virol *93*.
10.1128/jvi.01048-19.

46. Derdowski, A., Ding, L., and Spearman, P. (2004). A novel fluorescence resonance energy
transfer assay demonstrates that the human immunodeficiency virus type 1 Pr55Gag I
domain mediates Gag-Gag interactions. J Virol *78*, 1230-1242. 10.1128/jvi.78.3.1230-
1242.2004.

47. Levin, J.G., Mitra, M., Mascarenhas, A., and Musier-Forsyth, K. (2010). Role of HIV-1
nucleocapsid protein in HIV-1 reverse transcription. RNA Biol *7*, 754-774.
10.4161/rna.7.6.14115.

48. Darlix, J.L., Gabus, C., Nugeyre, M.T., Clavel, F., and Barré-Sinoussi, F. (1990). Cis
elements and trans-acting factors involved in the RNA dimerization of the human
immunodeficiency virus HIV-1. J Mol Biol *216*, 689-699. 10.1016/0022-2836(90)90392-y.

49. Gao, K., Gorelick, R.J., Johnson, D.G., and Bushman, F. (2003). Cofactors for human
immunodeficiency virus type 1 cDNA integration in vitro. J Virol *77*, 1598-1603.
10.1128/jvi.77.2.1598-1603.2003.

50. Onafuwa-Nuga, A.A., Telesnitsky, A., and King, S.R. (2006). 7SL RNA, but not the 54-kd
signal recognition particle protein, is an abundant component of both infectious HIV-1
and minimal virus-like particles. Rna *12*, 542-546. 10.1261/rna.2306306.

51. Itano, M.S., Arnion, H., Wolin, S.L., and Simon, S.M. (2018). Recruitment of 7SL RNA to
assembling HIV-1 virus-like particles. Traffic *19*, 36-43. 10.1111/tra.12536.

52. Eckwahl, M.J., Arnion, H., Kharytonchyk, S., Zang, T., Bieniasz, P.D., Telesnitsky, A., and
Wolin, S.L. (2016). Analysis of the human immunodeficiency virus-1 RNA packageome.
Rna *22*, 1228-1238. 10.1261/rna.057299.116.

53. Becker, J.T., and Sherer, N.M. (2017). Subcellular Localization of HIV-1 gag-pol mRNAs
Regulates Sites of Virion Assembly. J Virol *91*. 10.1128/jvi.02315-16.

54. Guerrero, J.F., Lesko, S.L., Evans, E.L., and Sherer, N.M. (2024). Studying Retroviral Life
Cycles Using Visible Viruses and Live Cell Imaging. Annu Rev Virol *11*, 125-146.
10.1146/annurev-virology-100422-012608.

55. GC, K., Lesko, S., Emery, A., Burnett, C., Gopal, K., Clark, S., Swanstrom, R., Sherer, N.M.,
Telesnitsky, A., and Kharytonchyk, S. (2025). HIV-1 single-transcription start-site mutants
display complementary replication functions that are restored by reversion. Journal of
Virology *0*, e02139-02124. doi:10.1128/jvi.02139-24.

56. Sundquist, W.I., and Kräusslich, H.G. (2012). HIV-1 assembly, budding, and maturation.
Cold Spring Harb Perspect Med *2*, a006924. 10.1101/cshperspect.a006924.

57.   Lee, I.H., Kai, H., Carlson, L.A., Groves, J.T., and Hurley, J.H. (2015). Negative membrane
1086   curvature catalyzes nucleation of endosomal sorting complex required for transport
1087   (ESCRT)-III assembly. Proc Natl Acad Sci U S A *112*, 15892-15897.
1088   10.1073/pnas.1518765113.
1089   58.   Van Dusen, C.M., Yee, L., McNally, L.M., and McNally, M.T. (2010). A glycine-rich domain
1090   of hnRNP H/F promotes nucleocytoplasmic shuttling and nuclear import through an
1091   interaction with transportin 1. Mol Cell Biol *30*, 2552-2562. 10.1128/mcb.00230-09.
1092   59.   Sanford, J.R., Wang, X., Mort, M., Vanduyn, N., Cooper, D.N., Mooney, S.D., Edenberg,
1093   H.J., and Liu, Y. (2009). Splicing factor SFRS1 recognizes a functionally diverse landscape
1094   of RNA transcripts. Genome Res *19*, 381-394. 10.1101/gr.082503.108.
1095   60.   Änkö, M.L., Müller-McNicoll, M., Brandl, H., Curk, T., Gorup, C., Henry, I., Ule, J., and
1096   Neugebauer, K.M. (2012). The RNA-binding landscapes of two SR proteins reveal unique
1097   functions and binding to diverse RNA classes. Genome Biol *13*, R17. 10.1186/gb-2012-
1098   13-3-r17.
1099   61.   Li, J., Li, G., Qi, Y., Lu, Y., Wang, H., Shi, K., Li, D., Shi, J., Stovall, D.B., and Sui, G. (2021).
1100   SRSF5 regulates alternative splicing of DMTF1 pre-mRNA through modulating SF1
1101   binding. RNA Biol *18*, 318-336. 10.1080/15476286.2021.1947644.
1102   62.   Buratti, E., Muro, A.F., Giombi, M., Gherbassi, D., Iaconcig, A., and Baralle, F.E. (2004).
1103   RNA folding affects the recruitment of SR proteins by mouse and human polypurinic
1104   enhancer elements in the fibronectin EDA exon. Mol Cell Biol *24*, 1387-1400.
1105   10.1128/mcb.24.3.1387-1400.2004.
1106   63.   Cléry, A., Sinha, R., Anczuków, O., Corrionero, A., Moursy, A., Daubner, G.M., Valcárcel,
1107   J., Krainer, A.R., and Allain, F.H.-T. (2013). Isolated pseudo–RNA-recognition motifs of SR
1108   proteins can regulate splicing using a noncanonical mode of RNA recognition.
1109   Proceedings of the National Academy of Sciences *110*, E2802-E2811.
1110   doi:10.1073/pnas.1303445110.
1111   64.   Keane, S.C., Heng, X., Lu, K., Kharytonchyk, S., Ramakrishnan, V., Carter, G., Barton, S.,
1112   Hosic, A., Florwick, A., Santos, J., et al. (2015). Structure of the HIV-1 RNA packaging
1113   signal. Science *348*, 917-921. 10.1126/science.aaa9266.
1114   65.   Wilkinson, K.A., Gorelick, R.J., Vasa, S.M., Guex, N., Rein, A., Mathews, D.H., Giddings,
1115   M.C., and Weeks, K.M. (2008). High-Throughput SHAPE Analysis Reveals Structures in
1116   HIV-1 Genomic RNA Strongly Conserved across Distinct Biological States. PLOS Biology *6*,
1117   e96. 10.1371/journal.pbio.0060096.
1118   66.   Abd El-Wahab, E.W., Smyth, R.P., Mailler, E., Bernacchi, S., Vivet-Boudou, V., Hijnen, M.,
1119   Jossinet, F., Mak, J., Paillart, J.-C., and Marquet, R. (2014). Specific recognition of the
1120   HIV-1 genomic RNA by the Gag precursor. Nature Communications *5*, 4304.
1121   10.1038/ncomms5304.
1122   67.   van Hemert, F.J., van der Kuyl, A.C., and Berkhout, B. (2013). The A-nucleotide
1123   preference of HIV-1 in the context of its structured RNA genome. RNA Biol *10*, 211-215.
1124   10.4161/rna.22896.
1125   68.   Sette, P., O'Connor, S.K., Yerramilli, V.S., Dussupt, V., Nagashima, K., Chutiraka, K.,
1126   Lingappa, J., Scarlata, S., and Bouamr, F. (2016). HIV-1 Nucleocapsid Mimics the
1127   Membrane Adaptor Syntenin PDZ to Gain Access to ESCRTs and Promote Virus Budding.
1128   Cell Host Microbe *19*, 336-348. 10.1016/j.chom.2016.02.004.

1129   69.   Gaunt, E.R., and Digard, P. (2022). Compositional biases in RNA viruses: Causes,
1130   consequences and applications. Wiley Interdiscip Rev RNA *13*, e1679.
1131   10.1002/wrna.1679.
1132   70.   Gc, K., Lesko, S., Emery, A., Burnett, C., Gopal, K., Clark, S., Swanstrom, R., Sherer, N.M.,
1133   Telesnitsky, A., and Kharytonchyk, S. (2024). HIV-1 single transcription start site mutants
1134   display complementary replication functions that are restored by reversion. bioRxiv.
1135   10.1101/2024.12.04.626847.
1136   71.   Kane, M., Zang, T.M., Rihn, S.J., Zhang, F., Kueck, T., Alim, M., Schoggins, J., Rice, C.M.,
1137   Wilson, S.J., and Bieniasz, P.D. (2016). Identification of Interferon-Stimulated Genes with
1138   Antiretroviral Activity. Cell Host Microbe *20*, 392-405. 10.1016/j.chom.2016.08.005.
1139   72.   Mohammed, K.D., Topper, M.B., and Muesing, M.A. (2011). Sequential deletion of the
1140   integrase (Gag-Pol) carboxyl terminus reveals distinct phenotypic classes of defective
1141   HIV-1. J Virol *85*, 4654-4666. 10.1128/jvi.02374-10.
1142   73.   Shema Mugisha, C., Dinh, T., Kumar, A., Tenneti, K., Eschbach, J.E., Davis, K., Gifford, R.,
1143   Kvaratskhelia, M., and Kutluay, S.B. (2022). Emergence of Compensatory Mutations
1144   Reveals the Importance of Electrostatic Interactions between HIV-1 Integrase and
1145   Genomic RNA. mBio *13*, e0043122. 10.1128/mbio.00431-22.
1146   74.   Madison, M.K., Lawson, D.Q., Elliott, J., Ozanturk, A.N., Koneru, P.C., Townsend, D.,
1147   Errando, M., Kvaratskhelia, M., and Kutluay, S.B. (2017). Allosteric HIV-1 Integrase
1148   Inhibitors Lead to Premature Degradation of the Viral RNA Genome and Integrase in
1149   Target Cells. J Virol *91*. 10.1128/JVI.00821-17.
1150   75.   Pizzato, M., Erlwein, O., Bonsall, D., Kaye, S., Muir, D., and McClure, M.O. (2009). A one-
1151   step SYBR Green I-based product-enhanced reverse transcriptase assay for the
1152   quantitation of retroviruses in cell culture supernatants. J Virol Methods *156*, 1-7.
1153   10.1016/j.jviromet.2008.10.012.
1154   76.   Mugisha, C.S., Dinh, T., Kumar, A., Tenneti, K., Eschbach, J.E., Davis, K., Gifford, R.,
1155   Kvaratskhelia, M., and Kutluay, S.B. (2022). Emergence of Compensatory Mutations
1156   Reveals the Importance of Electrostatic Interactions between HIV-1 Integrase and
1157   Genomic RNA. mBio *13*, e00431-00422. doi:10.1128/mbio.00431-22.
1158   77.   Shema Mugisha, C., Tenneti, K., and Kutluay, S.B. (2020). Clip for studying protein-RNA
1159   interactions that regulate virus replication. Methods *183*, 84-92.
1160   https://doi.org/10.1016/j.ymeth.2019.11.011.
1161   78.   Kutluay, S.B., and Bieniasz, P.D. (2016). Analysis of HIV-1 Gag-RNA Interactions in Cells
1162   and Virions by CLIP-seq. Methods Mol Biol *1354*, 119-131. 10.1007/978-1-4939-3046-
1163   3_8.
1164   79.   Chan, Patricia P., Lin, Brian Y., Mak, Allysia J., and Lowe, Todd M. (2021). tRNAscan-SE
1165   2.0: improved detection and functional classification of transfer RNA genes. Nucleic
1166   Acids Research *49*, 9077-9096. 10.1093/nar/gkab688.
1167   80.   Langmead, B., Trapnell, C., Pop, M., and Salzberg, S.L. (2009). Ultrafast and memory-
1168   efficient alignment of short DNA sequences to the human genome. Genome Biol *10*,
1169   R25. 10.1186/gb-2009-10-3-r25.
1170   81.   Danecek, P., Bonfield, J.K., Liddle, J., Marshall, J., Ohan, V., Pollard, M.O., Whitwham, A.,
1171   Keane, T., McCarthy, S.A., Davies, R.M., and Li, H. (2021). Twelve years of SAMtools and
1172   BCFtools. GigaScience *10*. 10.1093/gigascience/giab008.

1173

A

Gag (M)

64 –
51 –

Gag (C)

64 –
51 –

WT ΔNC G2A hnRNP A1 hnRNP F hnRNP K SRSF1 SRSF4 SRSF5

B

gRNA multiply spl. GAPDH

relative copies

1×10¹
1×10⁰
1×10⁻¹
1×10⁻²

****

WT ΔNC G2A hnRNP A1 hnRNP F hnRNP K SRSF1 SRSF4 SRSF5

C

WT ΔNC ΔNC-NC hnRNP A1 hnRNP F hnRNP K SRSF5

MS2-mCherry

Gag-YFP

Merge with DAPI

Scale bar = 20µm

ROI

Scale bar = 2.5µm

D

WT ΔNC-NC hnRNP A1 hnRNP F

hnRNP K SRSF1 SRSF4 SRSF5

Scale bar = 0.5µm