A physiology-based mathematical model of the renin-angiotensin system, bone remodeling, and calcium homeostasis: Effects of estrogen and renin-angiotensin system inhibitors

Melissa M. Stadt[1,*] and Anita T. Layton[1,2]

[1]Department of Applied Mathematics, University of Waterloo, Waterloo, ON, Canada
[2]Cheriton School of Computer Science, Department of Biology, School of Pharmacology, University of Waterloo, Waterloo, ON, Canada
[*]corresponding author, mstadt@uwaterloo.ca

Abstract

During menopause, estrogen levels decline significantly, leading to substantial physiological changes due to estrogen's regulatory role in various systems. In particular, estrogen helps prevent excessive bone resorption by its impact on bone remodeling. When estrogen levels decrease, bone resorption increases, often resulting in weakened bones and osteoporosis in post-menopausal women. Experimental studies have also shown that estrogen regulates the renin-angiotensin system (RAS), a hormone system involved in many physiological processes, including blood pressure regulation. Additionally, the RAS has an interconnected relationship with calcium regulatory and bone remodeling systems. Given these dynamic interplays, how would perturbations in one system affect the others? To answer that question, we developed a physiology-based mathematical model that simulates the interactions of estrogen, key RAS components, calcium regulation, and bone remodeling. Through sensitivity analysis and model simulations, we investigated how declining estrogen levels affect the RAS and bone mineral density. Furthermore, we quantified how RAS inhibitors, specifically angiotensin-converting enzyme inhibitors and angiotensin receptor blockers, may increase bone density during post-menopausal estrogen decline.

1 Introduction

Bone remodeling is a complex physiological process that maintains the balance between bone formation and resorption. This process is mediated by bone cells: osteoblasts, which are responsible for bone formation, and osteoclasts, which are responsible for bone resorption. The activity and proliferation of osteoblasts and osteoclasts is regulated by a system of various receptors, their corresponding ligands, and hormonal signals [1, 2]. Estrogen, in particular, impacts bone remodeling by inhibiting osteoclast activity, thereby protecting bones from excessive resorption. When estrogen levels in women decline post-menopause, osteoclast activity increases, resulting in increased

resorption, and ultimately weakening bones. This often leads to post-menopausal osteoporosis. Post-menopausal osteoporosis is the most common form of osteoporosis, which is characterized by increased bone fragility, making fractures more likely from minor trauma or in severe cases, even spontaneous fracture. It is estimated that about 1 in 3 women over 50 years of age will suffer a fracture related to post-menopausal osteoporosis [3, 4].

The main component of bone structure is hydroxyapatite, a compound made up of phosphate and calcium. Given their role in developing the key structure of bones, maintaining balanced levels of calcium and phosphate is critical for bone health. The regulation of calcium and phosphate levels is driven by calciotropic hormones: calcitriol and parathyroid hormone (PTH) [5]. Researchers have reported a bidirectional relationship between the calciotropic hormones and the renin-angiotensin system (RAS) [6], a hormone system that plays a key role in many physiological processes, including blood pressure regulation, in addition to a direct impact of the RAS on bone remodeling [7]. The RAS is a hormone cascade that produces the bioactive peptide angiotensin II (Ang II). Ang II binds to angiotensin type 1 receptors (AT1R) and angiotensin type 2 receptors (AT2R) to trigger various physiological effects. The RAS is also modulated by estrogen levels, and therefore is impacted by the rapidly declining estrogen levels in post-menopausal women [8]. This dysregulation contributes to the higher susceptibility of post-menopausal women to the development of hypertension, cardiovascular disease, and osteoporosis [8].

Inhibition of RAS components, particularly angiotensin-converting enzyme (ACE) and AT1R binding, has shown to be effective targets for antihypertensive treatment [9]. These drugs are classified as RAS inhibitors. Experimental data has shown that the use of RAS inhibitors improves bone quality independent of their effect on blood pressure [10, 11]. Given that RAS inhibitors are relatively well tolerated, safe, and are reasonably priced drugs, there has been interest in using RAS inhibitors to prevent osteoporosis in post-menopausal women [10–12]. Additionally, since hypertension is highly prevalent in post-menopausal women, understanding the impacts of antihypertensive treatments, such as RAS inhibitors, on bone remodeling is important for achieving blood pressure control and bone health in post-menopausal women [10–12].

Many mathematical models have been developed to study bone remodeling as recently reviewed in Ref. [2]. Peterson and Riggs [13] pioneered a mathematical model of calcium homeostasis coupled to a detailed bone biology model. Their model consists of bone, renal, and blood compartments to predict levels of PTH, calcitriol, calcium, phosphate, and bone remodeling activity. This model has been applied to investigate applications related to bone remodeling and calcium homeostasis, including the impact of estrogen [14], mineral bone disorder in patients with chronic kidney disease [15], and the impact of fibroblast growth factor 23 and vascular calcification [16]. Mathematical models have also been developed for the RAS to understand its role in blood pressure regulation. Specifically, Lo et al. [17] developed a model of the RAS in humans to investigate the differences between normo-tensive and hypertensive patients. This model was later included in a full body blood pressure model by Hallow et al. [18]. Leete et al. [19] further developed this model to investigate the effects of sex-based differences in the RAS on blood pressure regulation. However, no known models have been developed that include the direct impact of estrogen on the RAS or have coupled the RAS impacts with calcium homeostasis and bone remodeling in humans.

In this study, we (i) investigate the impact of age-related estrogen decline by including the impact of estrogen on the RAS and (ii) couple the calcium homeostasis and bone remodeling model (Refs. [13,14]) with the RAS model (Refs. [17–19]) to predict the impact of RAS inhibitors on this system.

2 Methods

In this study, we develop a coupled ordinary differential equation model that represents (i) the impact of estrogen on the RAS (Section 2.1.1) and (ii) couple this estrogen-RAS model with a bone biology and calcium homeostasis model (Section 2.3). We then use this model to simulate the effects of post-menopausal estrogen decline on the RAS, bone remodeling, and calcium homeostasis. Additionally, we simulate the impact of RAS inhibitors. To quantify the impact of parameter variation on model output, we conduct a global sensitivity analysis. A model schematic is shown in Fig. 2.1. A list of the model equations is given in Appendix D.

2.1 Renin-angiotensin system model

In this study, we use the RAS model with the female-specific parameter set from Ref. [19]. The model variables consist of the systemic concentrations of angiotensinogen (AGT), Ang I, Ang IV, Ang (1-7), AT1R-bound Ang II, and AT2R-bound Ang II. AT1R-bound Ang II inhibits renin secretion through a feedback loop. For a detailed description of the underlying RAS model used in this study, see Refs. [18, 19].

2.1.1 Impact of estrogen decline in aging women on the RAS

Estrogen levels impact the RAS by activating and inhibiting various key processes. Specifically, estrogen activates AGT production and AT2R binding, while inhibiting renin secretion, ACE activity, and AT1R binding [8]. These processes are illustrated in Fig. 2.1.

After the onset of menopause, estrogen levels decrease dramatically. To model this decline in estrogen levels, we use the estrogen model for post-menopausal women from Ref. [20]. The normalized level of estrogen, denoted by E, is modeled through explicit time dependence so that

$$E(t) = \begin{cases} 1 & t < t_e \\ \frac{1}{1+(t-t_e)/\tau_e} & t \geq t_e \end{cases} \tag{2.1}$$

where t is the time in years (i.e., age), $t_e = 50$ years is the age of onset of estrogen decline, and $\tau_e = 2.6$ years is a parameter fit to normalized estrogen data from Ref. [21] that determines the slope of the decline in estrogen (see Ref. [20] for details). Note that before t_e, $E = 1$ so that estrogen is assumed to remain at baseline level pre-menopause. Indeed, estrogen levels vary during the menstrual cycle [22]. In this study, we do not consider variations in estrogen over the menstrual cycle in pre-menopausal women and consider baseline estrogen level pre-menopause as the average level through the full cycle.

To include the impact of estrogen on the RAS, we use functions of the form:

$$g_i^+(E) = V_{\max,i}^+ \times \frac{(E/e_i)}{1 + (E/e_i)} \tag{2.2}$$

$$g_i^-(E) = V_{\max,i}^- \times \frac{1}{1 + (E/e_i)} \tag{2.3}$$

to model activation (g^+) or inhibition (g^-) with

$$V_{\max,i}^+ = e_i + 1, \qquad V_{\max,i}^- = \frac{e_i + 1}{e_i}$$

Figure 2.1: Schematic model of calcium regulation, bone remodeling, the renin-angiotensin system (RAS), and the impact of estrogen on these systems. Purple nodes represent phosphate and calcium levels, orange nodes represent phosphate and calcium renal handling. Yellow nodes represent the RAS, red nodes represent parathyroid hormone, pink nodes represent calcitriol, blue nodes represent bone cells and bone mineral density, green nodes represent signaling mechanisms in the bone. Nodes with a white background represent fixed values in the model. The impacts of estrogen are indicated by the red and green symbols for inhibition and activation, respectively. The solid arrows represent flux, secretion, synthesis, or expression, while the dashed lines represent impacts on other biological components. The pointed arrows represent activation and the rounded arrows represent inhibition. ACE: angiotensin-converting enzyme, AGT: angiotensinogen, Ang: angiotensin, AOH: 1α-hydroxylase, AT1R: angiotensin type 1 receptor, AT2R: angiotensin type 2 receptor, BMD: bone mineral density, Ca^{2+}: calcium, GFR: glomerular filtration rate, OB: osteoblast, OC: osteoclast, OPG: osteoprotegerin, PO$_4$: phosphate, PRC: plasma renin concentration, PT: parathyroid, PTH: parathyroid hormone, RANK: Receptor Activator of Nuclear Factor Kappa-B, RANKL: RANK ligand, TGFβ: transforming growth factor β

where i denotes the target mechanism, e_i is a parameter that is determined for each relevant target mechanism, and E is given by Eq. 2.1. Note that when $E = 1$ we have that $g_i^+(E) = g_i^-(E) = 1$ for any value of e_i, so that at baseline estrogen levels, there are no changes.

The enzyme renin is secreted by the juxtaglomerular cells of the kidneys to convert AGT to Ang I. Estrogen inhibits renin levels [8, 23, 24]. We let g_{renin}^- be the impact of estrogen on renin secretion so that renin secretion is modeled by

$$R_{\mathrm{sec}} = N_{\mathrm{rs}} \times \nu_{\mathrm{AT1R}} \times g_{\mathrm{renin}}^-(E) \tag{2.4}$$

where N_{rs} is baseline renin secretion and ν_{AT1R} is the feedback of [AT1R-bound Ang II] on renin secretion from Ref. [18]. Plasma renin concentration (PRC) is modeled by

$$\frac{d[\mathrm{PRC}]}{dt} = R_{\mathrm{sec}} - \frac{\ln(2)}{h_{\mathrm{renin}}} \times [\mathrm{PRC}] \tag{2.5}$$

where h_{renin} denotes the half life of renin as in Ref. [18].

AGT is synthesized in the liver and is stimulated by estrogen [8, 23, 25]. To model the impact of estrogen on AGT, we multiply the production of AGT, denoted by k_{AGT}, by $g_{\mathrm{AGT}}^+(E)$ so that [AGT] is modeled by

$$\frac{d[\mathrm{AGT}]}{dt} = g_{\mathrm{AGT}}^+(E) \times k_{\mathrm{AGT}} - \mathrm{PRA} - \frac{\ln(2)}{h_{\mathrm{AGT}}} \times [\mathrm{AGT}] \tag{2.6}$$

where

$$\mathrm{PRA} = X_{\mathrm{PRCPRA}} \times [\mathrm{PRC}] \times \frac{[\mathrm{AGT}]}{K_{\mathrm{AGT0}} + [\mathrm{AGT}]} \tag{2.7}$$

is plasma renin activity (PRA) with parameter X_{PRCPRA} and h_{AGT} is the half-life for AGT as in Ref. [18]. The equation for PRA is a slight modification from the equation used in Refs. [18, 19] to include the impact of AGT on PRA as well. The K_{AGT0} parameter was fit to predict the same model output in the original blood pressure model from Ref. [19]. In post-menopausal women, AGT levels decline from pre-menopause levels [23].

Angiotensin-converting enzyme (ACE) converts Ang I into Ang II. Researchers have reported that both ACE expression and activity are inhibited by estrogen in rodent models [26–29]. We let g_{ACE}^- represent the inhibition of estrogen levels on ACE activity so that the concentration of Ang I and Ang II are modeled by

$$\frac{d[\mathrm{AngI}]}{dt} = \mathrm{PRA} - (g_{\mathrm{ACE}}^-(E) \times c_{\mathrm{ACE}} + c_{\mathrm{chym}} + c_{\mathrm{nep}})[\mathrm{AngI}] - \frac{\ln(2)}{h_{\mathrm{AngI}}}[\mathrm{AngI}] \tag{2.8}$$

$$\begin{aligned}
\frac{d[\mathrm{AngII}]}{dt} = {} & (g_{\mathrm{ACE}}^-(E) \times c_{\mathrm{ACE}} + c_{\mathrm{chym}})[\mathrm{AngI}] \\
& - (c_{\mathrm{ACE2}} + c_{\mathrm{II,IV}} + g_{\mathrm{AT1}}^-(E) \times c_{\mathrm{AT1}} + g_{\mathrm{AT2}}^+(E) \times c_{\mathrm{AT2}})[\mathrm{AngII}] \\
& - \frac{\ln(2)}{h_{\mathrm{AngII}}}[\mathrm{AngII}] + k_{\mathrm{inf}}
\end{aligned} \tag{2.9}$$

where c_i denotes the corresponding enzyme activities and h_{AngI} and h_{AngII} are the half lives of Ang I and Ang II, respectively, as in Ref. [18]. k_{inf} is an input used for Ang II infusion (see Section 2.3.4) and is set to 0 in simulations with no Ang II infusion.

Ang II is reported to remain around the same in aging humans [30]. Ovariectomized rats showed a small decrease in plasma Ang II levels, which was reversed by estrogen replacement at pre-ovariectomy levels [31, 32]. We note that Ang II levels are often reported in post-menopausal women

on estrogen replacement therapy, but researchers have found significant differences in the impact of estrogen replacement therapy on the RAS when taking via oral or transdermal therapy [33]. This is likely due to the high exposure to estrogen in the liver when ingesting oral estrogen.

Ang II binds to AT1R and AT2R, triggering different physiological responses. Estrogen inhibits Ang II binding to AT1R, but activates Ang II binding to AT2R [8, 29, 34]. Angiotensin type 1 receptor density is increased over 2-fold in ovariectomized rats [35]. We based our parameter value for e_{AT1} on these results. AT1R-bound Ang II ([AT1R-AngII]) and AT2R-bound Ang II ([AT2R-AngII]) concentrations are modeled by

$$\frac{d[\text{AT1R} - \text{AngII}]}{dt} = g_{\text{AT1}}^{-}(E) \times c_{\text{AT1}}[\text{AngII}] - \frac{\ln(2)}{h_{\text{AT1}}}[\text{AT1R} - \text{AngII}] \tag{2.10}$$

$$\frac{d[\text{AT2R} - \text{AngII}]}{dt} = g_{\text{AT2}}^{+}(E) \times c_{\text{AT2}}[\text{AngII}] - \frac{\ln(2)}{h_{\text{AT2}}}[\text{AT2R} - \text{AngII}] \tag{2.11}$$

with h_{AT1} and h_{AT2} denoting the half-lives of AT1R-bound Ang II and AT2R-bound Ang II from Ref. [18]. c_{AT1} and c_{AT2} denote the binding rates of Ang II to AT1R and AT2R, respectively, as in Ref. [18]. g_{AT1}^{-} and g_{AT2}^{+} represent the impact of estrogen on the respective binding rate.

We chose parameter values for e_i based on reported changes in the respective RAS levels in the literature as described above. The values for e_i for each RAS component impacted by estrogen levels are listed in Table 2.1. Model simulations of the estrogen-RAS model in isolation are shown in Appendix A.

Name	Value	Equation(s)
e_{AGT}	0.15	g_{AGT}^{+}, Eq. 2.6
e_{renin}	0.15	g_{renin}^{-}, Eq. 2.4
e_{ACE}	5.0	g_{ACE}^{-}, Eqs. 2.8,2.9
e_{AT1}	0.5	g_{AT1}^{-}, Eqs. 2.9,2.10
e_{AT2}	0.05	g_{AT2}^{+}, Eqs. 2.9,2.11

Table 2.1: *Values of the parameters e_i that determine the impact of estrogen on renin-angiotensin system components. The e_i parameters are unitless.*

2.1.2   RAS inhibitor simulations

To simulate the RAS inhibitors, we use the same approach as in Ref. [18]. Specifically, to simulate treatment with an angiotensin receptor blocker (ARB), we let

$$c_{\text{AT1}} = c_{\text{AT1,0}} \times (1 - \sigma_{\text{ARB}}) \tag{2.12}$$

where $c_{\text{AT1,0}}$ is the baseline value for c_{AT1} and σ_{ARB} is the percent inhibition of AT1R binding by the ARB. Similarly, to simulate treatment with an ACE inhibitor (ACEi), we let

$$c_{\text{ACE}} = c_{\text{ACE,0}} \times (1 - \sigma_{\text{ACEi}}) \tag{2.13}$$

where $c_{\text{ACE,0}}$ is the baseline value for c_{ACE} and σ_{ACEi} is the percent inhibition of ACE by the ACEi.

2.2 Calcium homeostasis and bone biology model

The mathematical model developed by Peterson and Riggs [13] consists of detailed bone, renal, and blood compartments, and predicts levels of PTH, calcitriol, bone cell activity, calcium, and phosphate. In this study, we primarily used the original model from Peterson and Riggs [13] for the calcium homeostasis and bone remodeling components with the following modifications from published model extensions:

- Gaweda et al. [16] extended this model to include other known regulatory mechanisms on calcium and phosphate homeostasis. In this study, we also included the impact of phosphate levels on parathyroid gland capacity as added by Gaweda et al. [16] (equation for $H_{5,11}$ in Ref. [16]). Note that Gaweda et al. [16] used the Peterson and Riggs [13] model as is with their extensions so all other parameters and biological components are the same as in Ref. [13].

- A compartment for bone mineral density was added by Riggs et al. [14] and is included in the model in this study.

- Riggs et al. [14] added the impact of estrogen on bone remodeling and calcium homeostasis to the model in Ref. [13]. The authors included the impact of estrogen on latent and active transforming growth factor-β (TGF-β), osteoblast signaling, and renal calcium handling. TGF-β is a key cytokine in bone remodeling that regulates osteoblast and osteoclast proliferation and activity. Estrogen inhibits latent TGF-β formation, while stimulating its activation. TGF-β impacts bone remodeling activity activity through stimulating the expression of RANK on osteoclasts, stimulating osteoclast apoptosis, stimulating pre-osteoblast formation, and inhibiting osteoblast proliferation. Estrogen directly inhibits both the formation of pre-osteoblasts as well as the apoptosis rate of osteoblasts. Riggs et al. [14] also included the impact of estrogen on renal calcium reabsorption by decreasing renal calcium reabsorption with declining estrogen levels post-menopause. All of these estrogen impacts were included in the model use in this study.

A model schematic is shown in Fig. 2.1. The model equations are listed in Appendix D.

2.3 Composite calcium homeostasis, bone remodeling, and RAS model

To formulate a composite model, we connected the estrogen-RAS model (Section 2.1) with the calcium homeostasis and bone biology model (Section 2.2) through known regulatory mechanisms. Specifically, the composite model includes the impact of AT1R-bound Ang II on PTH secretion (Section 2.3.1), the impacts of extracellular calcium levels and the calciotropic hormones on renin secretion (Section 2.3.2), and the impact of AT1R-bound Ang II on bone remodeling (Section 2.3.3).

To model those interactions, we follow the approach inRef. [13] and characterize the changes in responses influenced by a given stimulant using hyperbolic functions:

$$H_x^- = \alpha - (\alpha - \rho) \times \frac{x^\gamma}{\delta^\gamma + x^\gamma} \tag{2.14}$$

$$H_x^+ = \rho + (\alpha - \rho) \times \frac{x^\gamma}{\delta^\gamma + x^\gamma} \tag{2.15}$$

where x denotes the stimulus variable, α is the maximum anticipated response, γ is the steepness of the response (note when $\gamma = 1$ the functions are sigmoidal), δ is the value of x that produces

the half-maximal response, and ρ is the minimum anticipated response. The minus or plus superscript in H denotes whether the response is a decrease or increase from the baseline steady state, respectively. Note that when $\rho = 0$, the response is a classical $E_{\max}$ response often used in modeling in pharmacology (see Ref. [36]). Parameter values for the equations added in this study are listed in Table 2.2. Additional parameter values can be found in Refs. [13, 14, 16] or using our publicly available model code (see Section 2.5).

Name	Units	Value	Equation
$\alpha_{\text{AT1R,PTH}}$	–	3.2	Eq. 2.16
$\rho_{\text{AT1R,PTH}}$	–	0.7	Eq. 2.16
$\delta_{\text{AT1R,PTH}}$	pmol/L	10.0	Eq. 2.16
$\gamma_{\text{AT1R,PTH}}$	–	1.5	Eq. 2.16
$\alpha_{\text{CaSR,renin}}$	–	1.2	Eq. 2.18
$\rho_{\text{CaSR,renin}}$	–	0.25	Eq. 2.18
$\delta_{\text{CaSR,renin}}$	mmol/L	2.6	Eq. 2.18
$\gamma_{\text{CaSR,renin}}$	–	5.0	Eq. 2.18
$\alpha_{\text{PTH,renin}}$	–	2.5	Eq. 2.19
$\rho_{\text{PTH,renin}}$	–	0.5	Eq. 2.19
$\delta_{\text{PTH,renin}}$	pmol/L	4.0	Eq. 2.19
$\gamma_{\text{PTH,renin}}$	–	3.0	Eq. 2.19
$\alpha_{\text{Ctriol,renin}}$	–	1.1	Eq. 2.20
$\rho_{\text{Ctriol,renin}}$	–	0.8	Eq. 2.20
$\delta_{\text{Ctriol,renin}}$	pmol/L	180	Eq. 2.20
$\gamma_{\text{Ctriol,renin}}$	–	1.0	Eq. 2.20
$\alpha_{\text{AT1R,RANKL}}$	–	5.0	Eq. 2.22
$\rho_{\text{AT1R,RANKL}}$	–	0.95	Eq. 2.22
$\delta_{\text{AT1R,RANKL}}$	pmol/L	16.0	Eq. 2.22
$\gamma_{\text{AT1R,RANKL}}$	–	3.0	Eq. 2.22
$\alpha_{\text{AT1R,OPG}}$	–	2.5	Eq. 2.23
$\rho_{\text{AT1R,OPG}}$	–	0.9	Eq. 2.23
$\delta_{\text{AT1R,OPG}}$	pmol/L	14	Eq. 2.23
$\gamma_{\text{AT1R,OPG}}$	–	2.0	Eq. 2.23

Table 2.2: Parameter values for coupling the RAS components to the calcium homeostasis and bone remodeling model. Additional parameters can be found in Refs. [13, 16].

2.3.1 AT1R-bound Ang II impact on PTH secretion

AT1R-bound Ang II concentration, denoted by [AT1R-AngII], increases PTH secretion from the parathyroid gland [6]. To model this effect, we let $H^{+}_{\text{AT1R,PTH}}$ denote the impact of AT1R-bound Ang II on PTH secretion:

$$H^{+}_{\text{AT1R,PTH}} = \rho_{\text{AT1R,PTH}} + (\alpha_{\text{AT1R,PTH}} - \rho_{\text{AT1R,PTH}}) \times \frac{[\text{AT1R} - \text{AngII}]^{\gamma_{\text{AT1R,PTH}}}}{\delta^{\gamma_{\text{AT1R,PTH}}}_{\text{AT1R,PTH}} + [\text{AT1R} - \text{AngII}]^{\gamma_{\text{AT1R,PTH}}}}. \quad (2.16)$$

where the parameters correspond to the function form of type Eq. 2.15. PTH concentration ([PTH]) in the blood is modeled by

$$\frac{d[\text{PTH}]}{dt} = S_{\text{PTH}} \times H^{+}_{\text{AT1R,PTH}} - k_{\text{deg−PTH}} \times [\text{PTH}] \quad (2.17)$$

where S_{PTH} is the secretion of PTH from the parathyroid gland and $k_{\text{deg-PTH}}$ is degradation as in
Ref. [13]. The parameters for Eq. 2.16 were fit using the Ang II only experiment data from Grant
et al. [37] (Section 2.3.4). The parameter values are listed in Table 2.2.

2.3.2 Calciotropic hormones, calcium, and renin secretion

Renin is secreted from specialized cells in the kidneys called juxtaglomerular cells. Renin secretion,
denoted by R_{sec}, is impacted by [PTH], calcitriol ([Ctriol]), and extracellular calcium levels. Specifically, increased extracellular calcium and [Ctriol] inhibit renin secretion, while [PTH] stimulates
renin secretion [6].

Extracellular calcium levels are detected by the calcium sensing receptor (CaSR) in the juxtaglomerular cells [38]. To model the impact of CaSR on R_{sec} we let:

$$H^-_{\text{CaSR,renin}} = \alpha_{\text{CaSR,renin}} - (\alpha_{\text{CaSR,renin}} - \rho_{\text{CaSR,renin}}) \times \frac{[\text{Ca}]^{\gamma_{\text{CaSR,renin}}}}{\delta^{\gamma_{\text{CaSR,renin}}}_{\text{CaSR,renin}} + [\text{Ca}]^{\gamma_{\text{CaSR,renin}}}}. \tag{2.18}$$

Ortiz-Capisano et al. [39,40] found that renin release was about 40% lower when using a calcimimetic
or at elevated calcium levels in the juxtaglomerular cells of the kidneys. We used this result to
estimate parameters so that at high calcium levels, renin release is decreased by about 40%.

To model stimulation of R_{sec} by [PTH] we let:

$$H^+_{\text{PTH,renin}} = \rho_{\text{PTH,renin}} + (\alpha_{\text{PTH,renin}} - \rho_{\text{PTH,renin}}) \times \frac{[\text{PTH}]^{\gamma_{\text{PTH,renin}}}}{\delta^{\gamma_{\text{PTH,renin}}}_{\text{PTH,renin}} + [\text{PTH}]^{\gamma_{\text{PTH,renin}}}}. \tag{2.19}$$

The parameters (listed in Table 2.2) were fit using the PTH only infusion experiment from Grant
et al. [37] (Section 2.3.4).

To model inhibition of R_{sec} by calcitriol we let:

$$H^-_{\text{Ctriol,renin}} = \alpha_{\text{Ctriol,renin}} - (\alpha_{\text{Ctriol,renin}} - \rho_{\text{Ctriol,renin}}) \times \frac{[\text{Ctriol}]^{\gamma_{\text{Ctriol,renin}}}}{\delta^{\gamma_{\text{Ctriol,renin}}}_{\text{Ctriol,renin}} + [\text{Ctriol}]^{\gamma_{\text{Ctriol,renin}}}}. \tag{2.20}$$

The parameters were chosen (listed in Table 2.2) based on the data from Tomaschitz et al. [41]
that related calcitriol levels with renin concentration. Taken together, we incorporate the combined
impact of calcium, PTH, and calcitriol on renin secretion by modifying Eq. 2.4 to

$$R_{\text{sec}} = N_{\text{rs}} \times \nu_{\text{AT1R}} \times g^-_{\text{renin}}(E) \times H^-_{\text{CaSR,renin}} \times H^+_{\text{PTH,renin}} \times H^-_{\text{Ctriol,renin}}. \tag{2.21}$$

2.3.3 AT1R-bound Ang II activation of RANKL and OPG

Bone remodeling is primarily regulated by osteoblasts and osteoclasts. Osteoblasts develop the
bone matrix, while osteoclasts break down bone tissue. RANKL (receptor activator of nuclear factor kappa-B ligand) is expressed by osteoblasts and binds to its receptor RANK (receptor activator
of nuclear factor kappa-B) on osteoclasts to promote osteoclast differentiation and activation. Osteoprotegerin (OPG) is a decoy receptor that binds to RANKL, preventing its binding with RANK
and thereby inhibiting osteoclast activity. The balance between RANKL and OPG is essential for
maintaining bone homeostasis as an increase in RANKL or decrease in OPG can lead to excessive or
reduced bone resorption, respectively. This system is often referred to as the RANK-RANKL-OPG
system and is illustrated in Fig. 2.1.

Shimizu et al. [7] reported that in ovariectomized rats, infusion of Ang II increased RANKL
expression and led to a significant decrease in bone volume, but this impact was abolished by

treatment with an AT1R blocker and not an AT2R blocker. Their results showed that RANKL production is activated by AT1R-bound Ang II levels. Additionally, Shimizu et al. [7] found that OPG expression is also increased in osteoblast cells due to infusion of Ang II, but to a much lesser extent than RANKL.

The impact of AT1R-bound Ang II on RANKL and OPG expression is modeled using the form in Eq. 2.15 for activation so that:

$$H^+_{\text{AT1R,RANKL}} = \rho_{\text{AT1R,RANKL}} + (\alpha_{\text{AT1R,RANKL}} - \rho_{\text{AT1R,RANKL}}) \times \frac{[\text{AT1R} - \text{AngII}]^{\gamma_{\text{AT1R,RANKL}}}}{\delta^{\gamma_{\text{AT1R,RANKL}}}_{\text{AT1R,RANKL}} + [\text{AT1R} - \text{AngII}]^{\gamma_{\text{AT1R,RANKL}}}},$$
(2.22)

$$H^+_{\text{AT1R,OPG}} = \rho_{\text{AT1R,OPG}} + (\alpha_{\text{AT1R,OPG}} - \rho_{\text{AT1R,OPG}}) \times \frac{[\text{AT1R} - \text{AngII}]^{\gamma_{\text{AT1R,OPG}}}}{\delta^{\gamma_{\text{AT1R,OPG}}}_{\text{AT1R,OPG}} + [\text{AT1R} - \text{AngII}]^{\gamma_{\text{AT1R,OPG}}}}.$$
(2.23)

These are included in the equations for the production of RANKL and OPG, respectively (Eqs. D.38 and D.40 in Appendix D). The values for $\alpha_{\text{AT1R,RANKL}}$ and $\alpha_{\text{AT1R,OPG}}$ represent the highest possible amount of activation of RANKL and OPG by AT1R binding, respectively. These values were set to the fold-change in RANKL and OPG expression after Ang II infusion in Shimizu et al. [7]. The parameter values are listed in Table 2.2.

2.3.4 Parameter fitting to PTH and Ang II infusion experiments

Grant et al. [37] conducted infusion experiments in healthy volunteers to determine the relationship between the RAS and PTH in humans. To test this, the authors gave healthy volunteers 1 hour infusions on different days of (i) Ang II only, (ii) PTH (i.e., teriparatide) only, and (iii) Ang II and PTH together. We simulated these experiments and used the data from this study to determine parameters values (listed in Table 2.2) for Eqs. 2.16, 2.18, and 2.19.

In their experiments, Ang II infusions were given in graded doses of 1, 3, and 10 ng/kg·min for consecutive infusions of 20 minutes each. We simulated Ang II infusion by letting $k_{\text{inf}} = 1$ ng/kg·min (Eq. 2.9) for the first 20 minutes, 3 ng/kg·min for the next 20 minutes, and 10 ng/kg·min for the final 20 minutes of the 1 hour infusion to simulate this infusion type. For PTH infusions we used the pharmacokinetic parameters from Peterson and Riggs [13] with a dose of 200 U of teriaparatide as reported by Grant et al. [37]. To fit the parameters, we used the `fmincon` function from `MATLAB` using the `interior-point` algorithm to minimize the difference between the predicted values and the data for PRA, intact PTH, and [Ang II] reported in Grant et al. [37]. In all of these simulations we let $E=1$ so that there are no estrogen effects.

2.4 Sobol sensitivity analysis

Mathematical models of complex biological systems, like the one used in this study, involve a large number of parameters with unknown or uncertain values. To investigate the impact of variations in parameter values on model output, we can use methods such as sensitivity analysis [42, 43]. Global sensitivity analysis methods evaluate the model at many points in the parameter space, providing an assessment of how individual parameters contribute to variations in model output across the parameter space, including interactions with other parameter changes [42–44].

The Sobol method is a variance-based global sensitivity analysis technique that decomposes the variance in the model output into contributions from each individual parameter as well as its interactions with other parameters [42, 45]. The results of a Sobol sensitivity analysis are called

"Sobol sensitivity indices" (or "Sobol indices" for short) and are computed for each parameter included in the analysis. First-order Sobol indices (denoted by S_{first}) are the fraction of the variance in the model output the individual parameter contributes to. Total order Sobol indices (denoted by S_{total}) quantify the fraction of output variance the parameter contributes to, including interactions with other parameter changes. By evaluating Sobol indices, we can identify which parameters are the most influential on model output. For more information about Sobol sensitivity analysis and its use in applications in biology, we refer readers to Refs. [42–45].

We conducted a Sobol sensitivity analysis on the full composite model (described in Section 2.3). The model outputs we considered are bone mineral density, AT1R-bound Ang II, PTH, calcitriol, extracellular calcium, and renin concentration over a simulation time from 20 to 80 years with age-related estrogen decline (i.e., same as the "no RASi" simulations in Figs. 3.2, 3.3). We computed Sobol indices at every 10 years of simulation time. The full composite model involves 181 parameters, so running a Sobol sensitivity analysis over all the parameters can lead to large numerical errors and computation time. To reduce the number of parameters used in the full Sobol sensitivity analysis, we applied a two-step approach:

1. We first conducted a Sobol sensitivity analysis varying all parameters only in the calcium homeostasis and bone biology model described in Section 2.2 by $\pm10\%$. This includes the baseline biological components of bone remodeling (including the detailed impacts of RANKL, RANK, OPG, and TGFβ on osteoblast and osteoclast activity), PTH secretion and degradation, calcitriol synthesis and degradation, renal calcium and phosphate reabsorption, intestinal calcium and phosphate absorption, and extracellular calcium and phosphate levels. We screened the 121 parameters included in this analysis for parameters that had an S_{first} value of greater than 1% in at least one of the outputs of interest (bone mineral density, AT1R-bound Ang II, PTH, calcitriol, or calcium) at some point in the simulation. This resulted in 30 sensitive parameters.

2. We then included all the new parameters from this study (listed in Tables 2.1 and 2.2) with the 30 parameters selected from the initial screening in Step 1, resulting in 59 total parameters. We conducted a Sobol sensitivity analysis for the full composite model with parameters varied by $\pm25\%$ of their baseline value.

The results of the Sobol sensitivity analysis are described in Section 3.3.

2.5   Code availability

All model code used in this study is available at `https://github.com/Layton-Lab/est-ras-bone`. Model simulations and code were conducted using `MATLAB R2022a`. The Sobol sensitivity analysis was conducted using `Simbiology`. Data originally presented in graphs from published papers (e.g., the data from Grant et al. [37] shown in Fig. 3.1) were extracted using Web Plot Digitizer (version 4.7).

3   Results

3.1   PTH and Ang II infusion experiments simulations

We conducted simulations of infusion of (i) Ang II, (ii) PTH, and (iii) PTH and Ang II together (described in Section 2.3.4). Model results, together with the experimental data reported for the clinical studies done by Grant et al. [37], are shown in Fig. 3.1.

Figure 3.1: Simulation results for infusion of Ang II ("Ang II only"), PTH ("PTH only"), and Ang II and PTH together ("Ang II + PTH"). Results for (A) plasma renin activity, (B) angiotensin II plasma concentration, (C), intact parathyroid hormone, and (D) angiotensin receptor type 1-bound Ang II are shown for during and after 1 hour infusions. Lines indicate simulations, while the dots and error bars in panels A–C are from the relevant experimental data presented in Grant et al. [37]. Ang II: angiotensin II, AT1R: angiotensin receptor type 1, iPTH: intact parathyroid hormone, PRA: plasma renin activity

When Ang II is infused in isolation, Ang II levels rise resulting in an increase in AT1R-bound Ang II levels (Fig. 3.1B, D), which stimulates PTH secretion so that endogenous PTH (intact PTH, denoted by [iPTH]) increases (Fig. 3.1C). Grant et al. [37] reported that during an Ang II infusion, [iPTH] increased, while PRA decreased slightly likely due to the negative feedback on renin secretion by AT1R-bound Ang II (Fig. 3.1 "Ang II only" data). Model simulations capture both the rise in [iPTH] (Fig. 3.1C "Ang II only") and slight decrease in PRA (Fig. 3.1A "Ang II only") during infusion.

When infusing PTH in healthy volunteers, PRA nearly doubled on average (Fig. 3.1A "PTH only" data) [37]. Model simulations predicted a 69% increase in PRA (Fig. 3.1A "PTH only"). While total PTH levels increased substantially during the PTH infusion experiment, endogenous PTH production steadily decreased (Fig. 3.1C "PTH only" data). Model simulations did not result in a significant decrease [iPTH]. That discrepancy may be attributed to a more detailed impact of PTH on renal calcium reabsorption, that is not represented in the model, which would increase calcium levels and thus decrease PTH secretion via the CaSR [46].

When infusing both Ang II and PTH together in healthy volunteers, the authors reported that

Ang II levels were similar to those reported in the Ang II only experiment, likely due to PTH having only a small impact on Ang II levels [37]. In our model simulations, Ang II levels were slightly higher in the Ang II and PTH infusion experiment than the Ang II only experiment, but were still within the range reported by Grant et al. [37] at the end of the 1 hour infusion time (Fig. 3.1B "Ang II + PTH"). Grant et al. [37] reported that PRA levels remained near baseline for the Ang II and PTH infusion experiment. In our model simulations, the "Ang II + PTH" simulations resulted in a transient increase in PRA but declines to near the levels prior to infusion by the end of the simulation time (Fig. 3.1A).

To summarize, model parameters were chosen to fit the observed responses of key RAS components and PTH following Ang II and PTH infusions with baseline estrogen levels.

3.2 Impact of estrogen decline and RAS inhibitors

How do declining estrogen levels in post-menopausal women impact the RAS? How do changes in the RAS and declining estrogen levels contribute to altered bone remodeling? Does bone mineral density improve with RAS inhibition? To investigate these questions, we conducted simulations of the composite calcium homeostasis, bone remodeling, and estrogen-RAS model with estrogen decline observed in post-menopausal women with and without drug interventions.

In each simulation, estrogen decline starts at 50 years, which is about the average age of menopause onset in women. We conducted simulations with key RAS inhibitors (ACEi and ARB) starting at age 60 years, unless otherwise indicated. For ARB simulations, we let σ_{ARB} (Eq. 2.12) vary from 88–98.3% as reported for different types and typical dose levels of ARBs by Hallow et al. [18]. For ACEi, we conducted simulations with σ_{ACEi} (Eq. 2.13) varying from 94–97.2% as reported for common types of ACEi and doses by Hallow et al. [18]. The results are shown in Figs. 3.2 and 3.3.

ACEi and ARB inhibit ACE activity and AT1R binding, respectively. While these are different components of the RAS, both drugs result in reduced levels of AT1R-bound Ang II (Fig. 3.2E). The impact of these drugs on the various components of the RAS are drug and dose-dependent. The varied range in values of σ_{ACEi} had a negligible effect on our model simulation results (Fig. 3.2, "ACEi"). In the ARB simulations, the impact on the RAS has a wider variation based on the value of σ_{ARB} (Fig. 3.2, "ARB"). Note that we simulated a wider range for σ_{ARB} than σ_{ACEi}.

We first consider the impact of ACEi and ARB on the RAS components. ARB directly inhibit the binding of Ang II to AT1R. As a result, AT1R-bound Ang II levels decreased by about 66–89%, depending on efficacy (Fig. 3.2E, "ARB" vs. "no RASi"). Recall that AT1R-bound Ang II inhibits renin secretion (Fig. 2.1). Hence, decreased AT1R-bound Ang II levels result in increased renin levels (Fig. 3.2B) as well as increased downstream Ang I (Fig. 3.2C) and Ang II levels (Fig. 3.2D). The lower AT1R binding along with increased Ang II levels results in more Ang II binding to AT2R, so that AT2R-bound Ang II levels increase by about 4-fold on average (Fig. 3.2F "ARB" vs. "no RASi"'). ACEi impact the RAS by inhibiting the activity of ACE, which is the enzyme that converts Ang I into Ang II (Fig. 2.1). Ang II levels decreased by an average of 52% in the ACEi simulations despite increased Ang I levels (Fig. 3.2D, C "ACEi" vs. "no RASi"). As a result, both AT2R-bound Ang II and AT1R-bound Ang II decreased by about 52% from the no RAS inhibitor case (Fig. 3.2E–F, "ACEi" vs. "no RASi").

Next, we analyze the age-related estrogen decline effect (Appendix A) on bone mineral density and its response to ACEi and ARB. Estrogen maintains bone density by regulating factors involved in bone remodeling that promote the differentiation and activation of osteoblasts, while inhibiting osteoclast activity (Fig. 2.1). When estrogen levels decline, osteoclast activity increases, which

Figure 3.2: Impact of post-menopausal estrogen decline and renin-angiotensin system inhibitors on the renin-angiotensin system. For the angiotensin receptor blocker simulations (labeled "ARB" in the legend), σ_{ARB} (Eq. 2.12) was varied between 88–98.3% based on the ranges reported in Ref. [18]. For the angiotensin-converting enzyme inhibitor simulations (labeled "ACEi" in the legend), σ_{ACEi} (Eq. 2.13) was varied between 94–97.2% as reported in Ref. [18]. Lines represent the mean values over the varied σ_i values for the respective treatment with shaded area representing the range. AGT: angiotensinogen, Ang I: angiotensin I, Ang II: angiotensin II, AT1R: angiotensin type 1 receptor, AT2R: angiotensin type 2 receptor, RAS: renin angiotensin system, RASi: RAS inhibitor, ARB: angiotensin receptor blocker, ACEi: angiotensin-converting enzyme inhibitor

increases bone resorption, ultimately resulting in a decrease in bone mineral density (Fig. 3.3A–B "no RASi"). The bone mineral density simulations results with the decline in estrogen levels is consistent with the data reported for aging women in Looker et al. [47] (Fig. 3.3A "no RASi" data). This decrease is a result of a combination of the direct impacts of estrogen on bone remodeling and increased AT1R-bound Ang II levels, which stimulates RANKL production to bind with RANK (Fig. 3.3C; Fig. 3.2E "no RASi"). RANK-RANKL stimulates osteoclast differentiation and activation. Throughout the estrogen decline simulations, the calciotropic hormones, PTH and calcitriol, as well as key electrolytes, calcium and phosphate, remained within normal range (Fig. B.2).

Age-related decreases in estrogen levels result in 4.5-fold increase in osteoclast activity (Fig. 3.3B "no RASi"). However, with ARB, AT1R-bound Ang II levels decrease (Fig. 3.2E), resulting in reduced RANKL production, so that osteoclast levels return to about 40% above baseline levels (Fig. 3.3B,C). As a result, by the end of the simulation time (i.e, $t = 80$ years), bone mineral density is at about 90% of baseline for ARB users versus 75% before estrogen decline with no treatment (Fig. 3.3A "ARB" vs. "no RASi"). ACEi also decreased AT1R-bound Ang II levels by a lesser amount (Fig. 3.2E) so that osteoclast levels are lowered to about 2-fold baseline levels. This results in a slight improvement in bone mineral density to about 85% of the level before estrogen decline

Figure 3.3: Impact of post-menopausal estrogen decline and RAS inhibitors on (A) bone mineral density, (B) osteoclast levels, (C) RANK-RANKL levels, and (D) RANK-OPG levels. The values are normalized to their baseline levels and shown as a percent of the baseline value. For the angiotensin receptor blocker simulations (labeled "ARB" in the legend), σ_{ARB} (Eq. 2.12) was varied between 88–98.3% based on the ranges reported in Ref. [18]. For the angiotensin-converting enzyme inhibitor simulations (labeled "ACEi" in the legend), σ_{ACEi} (Eq. 2.13) was varied between 94–97.2% as reported in Ref. [18]. Lines represent the mean values over the varied σ_i values for the respective treatment with shaded area representing the range. Data in panel (A) are used from the female data for femoral neck bone mineral density in Looker et al. [47]. The simulations and data are for femoral neck bone mineral density. BMD: bone mineral density, RANK: receptor activator of NF-kappaB, RANKL: RANK ligand, OPG: osteoprotegerin, RASi: RAS inhibitor, ARB: angiotensin receptor blocker, ACEi: angiotensin-converting enzyme inhibitor

(Fig. 3.3A "ACEi"). We observed that the range in the amount of AT1R-bound Ang II inhibition due to varied ARB efficacy did not have a large impact the change in bone levels on these drugs (Fig. 3.3; Fig. 3.2). Thus, the improvement to bone mineral density may not depend significantly on the dose of the RAS inhibitors.

We conducted simulations of RAS inhibitor treatment starting at different ages with age-related estrogen decline as in Eq. 2.1. In these simulations, we let $\sigma_{ARB} = 93\%$ and $\sigma_{ACEi} = 95\%$ based

on the average value for the respective drugs from Ref. [18]. We did this because varying σ_i did not have much impact on bone mineral density (Fig. 3.3A). The results are shown in Fig. 3.4. In all simulations, bone mineral density improved on both the ARB and ACEi drugs with a greater improvement with ARB treatment. When the administration of ARB or ACEi was started before the onset of estrogen decline (i.e., starting a RAS inhibitor before 50 years old), bone mineral density increased from baseline by about 5% (Fig. 3.4A1–A2). When estrogen decline started at 50 years, the bone mineral density decreased, but stays above 95% for patients on ARB (Fig. 3.4A1). When the administration of ACEi began before estrogen decline, bone mineral density declined below 90% of baseline during age-related estrogen decline (Fig. 3.4A2). Our model simulations suggest that early RAS inhibition prevents bone loss driven by age-related estrogen decline.

Figure 3.4: Model simulation results for varied starting age of angiotensin receptor blocker (A1–B1) and angiotensin-converting enzyme inhibitor (A2–B2). $\sigma_{\mathrm{ARB}} = 93\%$ in all simulations for ARB and $\sigma_{\mathrm{ACEi}} = 95.6\%$ in all ACEi simulations. The bone mineral density is femoral neck bone mineral density. AT1R: angiotensin type 1 receptor, ARB: angiotensin receptor blocker, ACEi: angiotensin-converting enzyme inhibitor, RASi: RAS inhibitor

In summary, the RAS, calcium, and bone remodeling model developed in this study represents the expected decrease in bone mineral density due to age-related estrogen decline. This is driven

by both the direct impacts of estrogen on bone remodeling and calcium regulation, as well as RAS activation by estrogen decline. Model simulations suggest that ARBs offer stronger protection for bone health than ACEi, and that result is consistent when starting treatment at different ages.

3.3 Sensitivity analysis

We conducted a Sobol sensitivity analysis for the full composite model during age-related estrogen decline in women (see Section 2.4 for details). The parameters with the largest average S_{total} values in Fig. 3.5A are the parameters that have the largest impact on variation in bone mineral density. The top five parameters, based on the average S_{total} value, are all related to the regulation of RANK and RANKL (Fig. 3.5A). The RANK-RANKL complex directly impacts bone remodeling the stimulating osteoclasts (Fig. 2.1). Specifically, the parameters $k_{\text{in,L}}$ and $k_{\text{in,RNK}}$ are baseline production rates of RANKL and RANK, respectively. The parameter $E_{\text{max,L−PTH}}$ relates to the impact of PTH on RANKL production. The parameters k_3 and k_4 are the binding and unbinding rate of the RANK-RANKL complex, respectively. The next parameter with the largest impact on bone mineral density is $\rho_{\text{AT1R,L}}$, which represents the minimum impact of AT1R on RANKL production (Eq. 2.22). $\delta_{\text{CaSR,renin}}$ represents the sensitivity of renin secretion to calcium levels. This is the parameter with the largest impact on [AT1R-bound Ang II] (Fig. 3.5B). In summary, the top parameters that impact bone mineral density all primarily relate to RANK, RANKL and RANK-RANKL binding.

AT1R-bound Ang II is the key biological component that connects the RAS with the calcium homeostasis and bone remodeling model (Fig. 2.1). The parameters with the 7 largest average S_{first} and S_{total} values for the model output of [AT1R-bound Ang II] are $\delta_{\text{CaSR,renin}}$, $\Phi_{\text{Plas2BoneCa0}}$, $K_{\text{Ca,PTH}}$, $\Phi_{\text{Bone2PlasCa0}}$, N_{rs}, c_{AT1R}, and X_{PRCPRA} (Fig. 3.5B). $\delta_{\text{CaSR,renin}}$ represents half-maximal response of renin secretion to calcium levels. $\Phi_{\text{Plas2BoneCa0}}$ and $\Phi_{\text{Bone2PlasCa0}}$ represent the baseline fluxes of calcium between the plasma and bone and both impact variation in calcium and renin levels (Fig. 3.6). Calcium level changes affect the RAS via the CaSR in the renin-secreting juxtaglomerular cells of the kidneys. $K_{\text{Ca,PTH}}$ is the half-maximal response of PTH secretion to calcium levels as mediated by the CaSR. This parameter plays a key role in variations in model output for [PTH], [Calcitriol], [Calcium], and [Renin] (Fig. 3.6 and 3.7), which all regulate the RAS. N_{rs}, c_{AT1R}, and X_{PRCPRA} are all directly related to the RAS (Section 2.1.1). Specifically, N_{rs} determines baseline renin secretion, c_{AT1R} is the binding rate for AT1R, and X_{PRCPRA} is baseline conversion rate from renin concentration to activity. Our sensitivity analysis shows that variations in [AT1R-bound Ang II] are driven by the CaSR impact on renin secretion and the RAS cascade.

The calciotropic hormones are the primary regulators of calcium and bone remodeling. Additionally, these hormones have a bidirectional relationship with the RAS (Fig. 2.1). In our Sobol sensitivity analysis, the parameters with the largest impact on PTH had high S_{total} values relative to the respective S_{first} values, likely due to the large number of regulators on PTH; these parameters are $V_{\text{max,PTH−reabs}}$, $K_{\text{Ca,PTH}}$, and $V_{\text{PTH−AOH}}$ (Fig. 3.7A). $K_{\text{Ca,PTH}}$ denotes the half-maximal response of PTH secretion driven by calcium sensing via the CaSR in the parathyroid glands. $V_{\text{max,PTH−reabs}}$ relates to the impact of PTH on calcium reabsorption in the kidneys. This parameter has a high S_{total} value for [PTH] as well as [Calcium] (Fig. 3.6 and 3.7), likely due to the interactions between renal calcium reabsorption, calcium levels, and PTH secretion. $V_{\text{PTH−AOH}}$ represents the maximal impact of PTH on 1α-hydroxylase, the enzyme that synthesizes calcitriol. PTH and calcitriol regulate one another, as calcitriol inhibits PTH secretion and PTH stimulates calcitriol production (Fig. 2.1). Both PTH and calcitriol are sensitive to the parameter $k_{\text{base,AOH}}$ which is the baseline production rate of 1α-hydroxylase, and to the parameters $\Phi_{\text{Bone2PlasCa0}}$ and $\Phi_{\text{Plas2BoneCa0}}$, which de-

Figure 3.5: Mean (bar) and standard deviation (error bar) of first and total order Sobol indices (denoted by S_{first} and S_{total} in legend, respectively) of full model ran from age 20 to 80 years with estrogen decline for output variables (A) femoral neck bone mineral density and (B) AT1R-bound Ang II. Parameters shown had a maximal value of $S_{total} > 0.05$ for at least one output variable. A description of the parameters is given in Table C.1. AT1R: angiotensin type 1 receptor, Ang II: angiotensin II

note the baseline rate of calcium exchange between the bone and plasma and vice versa (Fig. 3.7). The latter is likely due to the sensitivity of PTH to calcium levels, which are impacted by these two parameters as well (Fig. 3.6A).

The Sobol sensitivity analysis results shown here are the average (bar) and standard deviation (error bar) of S_{first} and S_{total} values over the full simulation time from age 20 to 80 years. We also analyzed the results over the simulation time from age 20 to 50, i.e., before estrogen decline. The parameters with the largest impact on model output were the same (results not shown) as those for the full simulation time. This suggests that the model output is most sensitive to parameters that represent the overall biological components of the composite model rather than parameters that represent the impact of estrogen on calcium regulation and bone remodeling.

In summary, our global sensitivity analysis shows that, in our model, bone remodeling is most sensitive to RANK-RANKL binding and that the RAS is sensitive to both calcium levels and the calciotropic hormones. These are the key biological components that impact RAS levels and bone

Figure 3.6: Mean (bar) and standard deviation (error bar) of first and total order Sobol indices (denoted by S_{first} and S_{total} in legend, respectively) of full model simulation from age 20 to 80 years with estrogen decline. Parameters shown had a maximal value of $S_{total} > 0.05$ for at least one output variable. A description of the parameters is given in Table C.1.

468 remodeling.

Figure 3.7: Mean (bar) and standard deviation (error bar) of first and total order Sobol indices (denoted by S_{first} and S_{total} in legend, respectively) of full model ran from age 20 to 80 years with estrogen decline for output variables (A) parathyroid hormone and (B) calcitriol. Parameters shown had a maximal value of $S_{total} > 0.05$ for at least one output variable. A description of the parameters is given in Table C.1. PTH: parathyroid hormone

4 Discussion

In this study we developed a mathematical model that includes the interconnected systems of estrogen, the RAS, PTH, calcitriol, calcium regulation, and bone remodeling. We used this model to:

- Investigate the impact of estrogen decline, as experienced in post-menopausal women, on the RAS, bone remodeling, and calcium homeostasis.

- Simulate potential therapeutic effects of ACEi and ARB on bone remodeling and calcium regulation.

- Assess the sensitivity of key model outputs to variations in the model's many unknown parameters using a Sobol sensitivity analysis.

To the best of our knowledge, this is the first model to incorporate the direct effects of estrogen on the RAS, as well as the integrated effects of the RAS on calcium regulatory and bone remodeling systems. As such, our model development and analysis have brought major advances to the active area of using mathematical modeling to investigate bone remodeling and calcium regulation.

Given that the prevalence of hypertension increases markedly with aging, hypertension and osteoporosis often coexist. RAS inhibitors are widely used, safe, reasonably priced long term treatments for hypertension. RAS inhibitors have also been proposed as a treatment to improve bone density due to the relationship between the RAS and bone remodeling [10, 11]. Since RAS inhibitors are already widely used for the treatment of hypertension in aging populations, including post-menopausal women, it may be beneficial to consider which RAS inhibitor may prevent or improve bone loss.

We conducted model simulations to investigate the impacts of RAS inhibitors, specifically ACEi and ARB, on bone remodeling for various stages of age-related estrogen decline. Since treatment with an ARB results in a larger reduction in AT1R-bound Ang II for all dose levels, the improvement in bone mineral density was greater in simulations of ARB than ACEi (Fig. 3.2E, Fig. 3.3A). It seems that the targeted impact of ARB on AT1R binding results in a better improvement in bone mineral density during post-menopausal estrogen decline than ACEi, which inhibit Ang II production via ACE. These results were consistent when we varied the starting age for ACEi and ARB before and after estrogen decline (Fig. 3.4). In summary, our model simulations showed that reductions in AT1R-bound Ang II abundance results in reduced osteoclast activity, which prevents and improves declines in bone mineral density driven by low estrogen levels.

In a clinical study, Kim et al. [10] reported that patients that have used ARB only experienced fewer fractures than patients who have never used a RAS inhibitor. Similarly, Solomon et al. [48] reported reduced fracture risk in users of ARB, while ACEi users had no change in fracture risk. The systemic review by Wu et al. [12] reported that lower fracture risk is consistently reported in users of ARB, but not ACEi. In contrast, our mathematical model predicted that bone mineral density is increased not only by ARB treatment but ACEi as well (Fig 3.3A). That discrepancy may be due to mechanisms not captured by our model. It is possible that the improvement in bone mineral density predicted by our model for ACEi is not enough to have a notable impact on fracture risk. We also note that bone mineral density is not the only factor that impacts fracture risk; the architecture of the skeleton, alcohol usage, smoking, physical activity, and previous fractures are also major factors in fracture risk that are not represented in the model in this study [49].

4.1 Clinical perspectives

Although estrogen replacement therapy (or hormone replacement therapy) can effectively manage menopause symptoms and post-menopausal osteoporosis, its benefits must be weighed against the heightened risks of blood clots, stroke, and breast cancer, especially with long-term use [50, 51]. Furthermore, side effects such as weight gain, diarrhea, and mood changes can affect patient well-being and adherence [51]. The balance of risks and benefits associated with hormone replacement therapy varies considerably based on individual physiology, time since menopause onset, and the specific type of hormone replacement therapy [50]. Other current osteoporosis treatments, including antiresorptives and osteoanabolic therapies, have limitations on their duration of use, typically ranging from one to five years [52]. Therefore, the development of alternative long-term osteoporosis treatments are needed to broaden the range of effective options for patients. Our mathematical modeling analysis presented in this study supports clinical evidence suggesting that ARBs may contribute to maintaining bone health during post-menopausal estrogen decline by reducing AT1R-bound Ang II levels.

Indeed, there are a number of clinical applications that this model could be extended to study. The RAS is a key hormone system in blood pressure regulation. The RAS model used in this study is part of a larger model that involves whole-body blood pressure regulation [18, 19]. Using a similar approach to Ref. [53], the model presented in this study may be incorporated into the blood pressure model (Ref. [19]) through the RAS to investigate the interactions of hypertension and bone remodeling. Chronic kidney disease (CKD) is a disorder of the loss of kidney function that affects over 10% of the global population [54]. Many patients with CKD experience CKD-metabolic bone disorder, where their impaired renal function results in loss of bone mineral density. Previous mathematical modeling studies have used the underlying calcium homeostasis and bone remodeling model to study this disorder [15, 16, 55]. However, none have considered the RAS. Many CKD patients take RAS inhibitors to protect their kidney function, prevent hypertension, and have been shown to improve bone health [56]. The model presented in this study may be used to investigate how RAS inhibitors may impact bone density and electrolyte balance in patients with CKD.

4.2 Limitations and future directions

In this study, we developed a mathematical model that describes the interactions between the RAS, calciotropic hormones, calcium regulation, and bone remodeling. One major limitation of this model is the large number of unknown parameters, resulting in uncertainty in the model outcomes. In this study, we analyzed the impacts of variations in parameter values on model output by conducting a global sensitivity analysis using the Sobol method. However, we do recognize that there may be parameters and mechanisms that are not included or could be changed in this model.

A worthwhile future direction is to accurately model an aging population. In this study, we conducted simulations that represent the estrogen decline that is seen in post-menopausal women. However, aging populations, including post-menopausal women, have shown other physiological changes related to the systems in the model presented in this study. For example, aging populations have shown a blunted response to ACE inhibitors, have a lower GFR, decreased intestinal calcium absorption, lower vitamin D levels, and higher PTH levels [57–59]. These effects are not included in the model developed in this study at this time. Indeed, future work may consider the specific physiological impacts of aging for a more comprehensive description of post-menopausal physiology.

Given the importance of estrogen and using a female-specific RAS, the model presented in this study only represents a female population. However, sex differences have been found in the RAS [60] and renal handling of calcium [61,62]. Men also can form osteoporosis as they age, which

is not driven by the declining estrogen levels found in post-menopausal women. Rianon et al. [63] found that there are sex- and race-specific effects of ACEi on the prevention of age-related bone loss. Specifically, the authors reported that ACEi were effective at protecting against bone loss in African-American elderly men [63]. This is different than reports that did not consider sex and race differences and only reported significant changes in fracture risk in patients using ARB. The model presented in this study may be extended to consider how underlying sex and racial differences may explain altered treatment outcomes [64].

In this study, we do not include the hormone aldosterone, which is connected with the RAS as a part of the renin angiotensin aldosterone system. Researchers have shown that aldosterone also has a bidirectional relationship with the calciotropic hormones [6]. Clinical reports have shown that the drug spironolactone (an aldosterone inhibitor), has a significant impact on the bone health of hypertensive patients [65]. Indeed this hormone may be added to this model to study the impacts of hypertension, spironolactone, hyperaldosteronism, or other relevant conditions and medications on bone density and calcium regulation. Additionally, in this study, we assume that the RAS activity in the bone is the same as the systemic RAS, though this may not always be the case. Experimental evidence has shown that there is a bone-specific RAS [66]. Future work may involve modeling a bone-specific RAS to consider the impacts on bone remodeling similar to the renal-specific RAS models in Refs. [17, 67].

Estrogen levels vary during the different phases of the menstrual cycle in menstruating women [22]. Women who take oral contraceptives also have different estrogen levels [68]. The mathematical model presented here may be adapted to consider how changes in estrogen impacts the RAS, calcium homeostasis, and bone remodeling during the different phases of the menstrual cycle or on oral contraceptives [69]. During pregnancy, estrogen levels rise dramatically, while post-parturition, the estrogen levels drop below pre-pregnancy levels to support lactation [70]. Each of these changes results in major impacts on physiology, including the RAS, calcium homeostasis, and bone remodeling [70, 71]. Previously, we developed mathematical models that represent calcium homeostasis in rats that consider sex differences and maternal adaptations of pregnancy and lactation in Ref. [72], but these models do not include a detailed bone compartment or the impact of the RAS. The model presented in this study may be similarly adapted to consider the major changes that women undergo during pregnancy and lactation.

Grants

This work is supported by the Canada 150 Research Chair program and by the National Science and Engineering Research Council of Canada via a Discovery award (RGPIN-2019-03916) to A.T.L. and a Canada Graduate Scholarship (CGS-D) to M.M.S. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Disclosures

No conflicts of interest, financial or otherwise, are declared by the authors.

Author Contributions

A.T.L. and M.M.S. conceived and designed research; M.M.S. wrote model code; M.M.S. conducted simulations; M.M.S. and A.T.L. analyzed data; M.M.S. and A.T.L. interpreted results of simulations; M.M.S. prepared figures; M.M.S. and A.T.L. drafted manuscript; M.M.S. and A.T.L. edited and revised manuscript; M.M.S. and A.T.L. approved final version of manuscript.

Supplemental Materials

A Impact of declining estrogen levels on the RAS

How do the various impacts of estrogen on the RAS change RAS levels during post-menopausal estrogen decline? To investigate this question, we conducted a simulations of the estrogen-RAS model presented in this study. The results are shown in Fig. A.1. We assume that estrogen decline starts at age 50 years (Fig. A.1A; Eq. 2.1). The decline in estrogen levels, decreases AGT production so that by age 80 years [AGT] is 60% below pre-menopausal levels (Fig. A.1B). Plasma renin secretion is inhibited by estrogen levels, which are rapidly decreasing. Renin secretion is inhibited by estrogen, so the lowered estrogen levels result in driving increased renin secretion so that ultimately plasma renin levels rise (Fig. A.1C).

Figure A.1: Simulation results of the impact of decreasing (A) estrogen levels on the plasma concentration of (B) AGT, (C) renin, (D) Ang I, (E) Ang II, (F) AT1R-bound Ang II, and (G) AT2R-bound Ang II. AGT: angiotensinogen, Ang: angiotensin, AT1R: angiotensin type 1 receptor, AT2R: angiotensin type 2 receptor

While renin concentration increases, the decreased AGT levels result in decreased PRA, which drives Ang I production. This results in stable Ang I levels (Fig. A.1D). ACE activity is increased so we see a slight increase in Ang II levels (Fig. A.1E). Ang II then binds to AT1R and AT2R. Estrogen inhibits AT1R binding and activates AT2R binding (Fig. 2.1). When estrogen levels decrease, AT1R receptors are inhibited less, thus leading to increased AT1R-bound Ang II levels (Fig A.1F). For AT2R receptors, initially, AT2R-bound Ang II levels slightly increase from baseline due to the increased Ang II levels, but later AT2R-bound Ang II levels decrease from baseline due to the loss of activation by estrogen levels (Fig. A.1G).

We conducted a Sobol sensitivity analysis on the estrogen-RAS model to investigate which parameters impact the estrogen deficiency model simulation the most for AGT, Ang II, AT1R-

₆₁₈ bound Ang II, and AT2R-bound Ang II. We varied the parameters that represent the rate of
₆₁₉ enzyme impacts on the RAS (i.e., parameters c_i) as well as the parameters that represent estrogen
₆₂₀ impact on the RAS (i.e., parameters e_i listed in Table 2.1). Each parameter was varied by a 2-fold
₆₂₁ increase and decrease from the baseline value. We then conducted a Sobol analysis at the time
₆₂₂ point of every 10 years for the estrogen deficiency simulation.

₆₂₃ The results for the total order Sobol indices (S_{total}) over the estrogen decline RAS simulation
₆₂₄ are shown in Fig. A.2. Before the onset of estrogen decline due to menopause (i.e., $t < 50$), the
₆₂₅ parameters e_i ($i=$ renin, AGT, ACE, etc), which represent the impact of estrogen on the RAS
₆₂₆ (Eqs. 2.2–2.3), do not have any impact on the model variation as expected because $E = 1$.

Figure A.2: Sobol total indices (S_{total}) for estrogen impact on (A) AGT, (B) Ang II, (C) AT1R-bound Ang II, and (D) AT2R-bound Ang II at different time points in the simulation. Parameters not shown did not have a S_{total} over 0.01 for any variable. AGT: angiotensinogen, Ang: angiotensin, AT1R, angiotensin type 1 receptor, AT2R: angiotensin type 2 receptor

₆₂₇ For renin, e_{renin} represents the impact of estrogen on renin secretion and leads to about 10% of
₆₂₈ the model output variation by $t = 70$ years (Fig. A.2A). The parameters with the largest output
₆₂₉ on model output of Ang II are c_{ACE} and c_{AT1R} which correspond to ACE activity and AT1R
₆₃₀ binding, respectively (Fig. A.2B). After the decline in estrogen, the impact of estrogen on renin
₆₃₁ secretion, e_{renin} contributes to the variation in the model output of Ang II as well (Fig. A.2B).
₆₃₂ AT1R-bound Ang II is also impacted by c_{ACE} and c_{AT1R} the most due to the dependence on Ang II

levels (Fig. A.2C). After the onset of estrogen decline, the impact of estrogen on renin and AT1R binding contribute to variation in the model output. By the end of the simulation time, about 10.2% and 3.2% of model output variation of AT1R-bound Ang II can be attributed to e_{renin} and e_{AT1R}, respectively (Fig. A.2C). For AT2R-bound Ang II, about 11% of the total variation can be attributed to the impact of estrogen on AT2R binding, e_{AT2R}, by the end of the simulation time (Fig. A.2D).

 # B   Additional Figures

Figure B.1: Replication of Fig. 3.1 for a longer simulation time. Simulation results for replication of the experiments in Grant et al. [37]. Results for (A) plasma renin activity (PRA), (B) angiotensin II (Ang II) plasma concentration, (C), parathyroid hormone (PTH) concentration, and (D) angiotensin receptor type II bound Ang II are shown for during and after 1 hour of infusions of Ang II only, PTH only, and Ang II and PTH infusions together. Lines indicate simulations, while the dots and error bars in panels A–C are from the relevant experimental data presented in Grant et al. [37]. AT1R: angiotensin receptor type 1, iPTH: intact parathyroid hormone

Figure B.2: Impact of estrogen deficiency and RAS inhibitors on extracellular levels of calciotropic hormones (A) parathyroid hormone and (B) calcitriol as well as electrolytes (C) calcium and (D) phosphate. For the angiotensin receptor blocker simulations (labeled "ARB" in the legend), σ_{ARB} was varied between 92–98.3% based on the ranges reported in Ref. [18]. For the angiotensin-converting enzyme inhibitor simulations (labeled "ACEi" in the legend), σ_{ACEi} was varied between 94–97.2% as reported in Ref. [18]. Lines represent the mean values over the varied σ_i values for the respective treatment with shaded area representing the range. PTH: parathyroid hormone, RASi: RAS inhibitor, ARB: angiotensin receptor blocker, ACEi: angiotensin-converting enzyme inhibitor

C Key parameter descriptions

Name	Description	Reference
$E_{\mathrm{max,L-PTH}}$	half-maximal response of effect of PTH on RANKL production	[13]
k_3	binding rate of RANK and RANKL	[13]
k_4	unbinding rate of RANK-RANKL	[13]
$k_{\mathrm{in,L}}$	baseline RANKL production	[13]
$k_{\mathrm{out,L}}$	degradation rate of RANKL	[13]
$k_{\mathrm{in,RNK}}$	baseline RANK production	[13]
$k_{\mathrm{out,RNK}}$	degradation rate of RANK	[13]
$\rho_{\mathrm{AT1R,L}}$	minimum impact of AT1R on RANKL	Eq. 2.22
$\delta_{\mathrm{AT1R,L}}$	half-maximal response of AT1R impact on RANKL	Eq. 2.22
$\alpha_{\mathrm{CaSR,renin}}$	maximum impact of CaSR on renin	Eq. 2.18
$\delta_{\mathrm{CaSR,renin}}$	half-maximal response of CaSR impact on renin	Eq. 2.18
$\Phi_{\mathrm{Plas2BoneCa0}}$	baseline calcium exchange from extracellular fluid to bone	[13]
$\Phi_{\mathrm{Bone2PlasCa0}}$	baseline calcium exchange from bone to extracellular fluid	[13]
c_{AT1R}	binding rate of Ang II to AT1R	[18, 19], Eq. 2.10
N_{rs}	baseline renin secretion	[18, 19], Eq. 2.4
X_{PRCPRA}	equilibrium ratio of PRA to PRA	[18]
$\delta_{\mathrm{PTH,renin}}$	half-maximal response of PTH impact on renin	Eq. 2.19
$K_{\mathrm{Ca,PTH}}$	half-maximal response of PTH secretion to CaSR	[13]
$k_{\mathrm{base,AOH}}$	baseline AOH production	[13]
$k_{\mathrm{deg,AOH}}$	degradation rate of AOH	[13]
$V_{\mathrm{max,PTH-AOH}}$	maximum effect of PTH on AOH production	[13]
$\delta_{\mathrm{AOH,Phos}}$	half-maximal response of impact of phosphate on AOH	[13]
$V_{\mathrm{max,Phos}}$	maximum impact of phosphate on AOH production	[13]
$V_{\mathrm{max,PTH-reabs}}$	maximum impact of PTH on renal reabsorption	[13]

Table C.1: Descriptions of parameters shown in Figs. 3.5, 3.6, and 3.7. The reference is the original source of the parameter value. Ang II: angiotensin II, AOH: 1α-hydroxylase, AT1R: angiotensin type 1 receptor, CaSR: calcium sensing receptor, OC: osteoclast, PTH: parathyroid hormone

D Model equations

A list of the ordinary differential equations in the model is given below.

D.1 Renin-angiotensin system

$$\frac{d[\text{PRC}]}{dt} = R_{\text{sec}} - \frac{\ln(2)}{h_{\text{renin}}} \times [\text{PRC}] \tag{D.1}$$

where

$$R_{\text{sec}} = N_{\text{rs}} \times \nu_{\text{AT1R}} \times g_{\text{renin}}^{-}(E) \times H_{\text{CaSR,renin}}^{-} \times H_{\text{PTH,renin}}^{+} \times H_{\text{Ctriol,renin}}^{-}. \tag{D.2}$$

$$\frac{d[\text{AGT}]}{dt} = g_{\text{AGT}}^{+}(E) \times k_{\text{AGT}} - \text{PRA} - \frac{\ln(2)}{h_{\text{AGT}}} \times [\text{AGT}] \tag{D.3}$$

where

$$\text{PRA} = X_{\text{PRCPRA}} \times [\text{PRC}] \tag{D.4}$$

$$\frac{d[\text{AngI}]}{dt} = \text{PRA} - (g_{\text{ACE}}^{-}(E) \times c_{\text{ACE}} + c_{\text{chym}} + c_{\text{nep}})[\text{AngI}] - \frac{\ln(2)}{h_{\text{AngI}}}[\text{AngI}] \tag{D.5}$$

$$\begin{aligned}
\frac{d[\text{AngII}]}{dt} &= (g_{\text{ACE}}^{-}(E) \times c_{\text{ACE}} + c_{\text{chym}})[\text{AngI}] \\
&\quad - (c_{\text{ACE2}} + c_{\text{II,IV}} + g_{\text{AT1}}^{-}(E) \times c_{\text{AT1}} + g_{\text{AT2}}^{+}(E) \times c_{\text{AT2}})[\text{AngII}] \\
&\quad - \frac{\ln(2)}{h_{\text{AngII}}}[\text{AngII}] + k_{\text{inf}}
\end{aligned} \tag{D.6}$$

$$\frac{d[\text{Ang}(1-7)]}{dt} = c_{\text{NEP}} \times [\text{AngI}] + c_{\text{ACE2}} \times [\text{AngII}] - \frac{\ln(2)}{h_{\text{Ang}(1-7)}}[\text{Ang}(1-7)] \tag{D.7}$$

$$\frac{d[\text{AngIV}]}{dt} = c_{\text{AngII=AngIV}} \times [\text{AngII}] - \frac{\ln(2)}{h_{\text{AngIV}}} \times [\text{AngIV}] \tag{D.8}$$

$$\frac{d[\text{AT1R} - \text{AngII}]}{dt} = g_{\text{AT1}}^{-}(E) \times c_{\text{AT1}}[\text{AngII}] - \frac{\ln(2)}{h_{\text{AT1}}}[\text{AT1R} - \text{AngII}] \tag{D.9}$$

$$\frac{d[\text{AT2R} - \text{AngII}]}{dt} = g_{\text{AT2}}^{+}(E) \times c_{\text{AT2}}[\text{AngII}] - \frac{\ln(2)}{h_{\text{AT2}}}[\text{AT2R} - \text{AngII}] \tag{D.10}$$

D.2 Calcium regulation

$$\frac{d[\text{Ca}]}{dt} = \Phi_{\text{Bone2Plas}-\text{Ca}} - \Phi_{\text{Plas2Bone}-\text{Ca}} - \Phi_{\text{Urine}-\text{Ca}} + \Phi_{\text{Gut2Plas}-\text{Ca}} \tag{D.11}$$

 with

$$\Phi_{\text{Gut2Plas}-\text{Ca}} = I_{\text{Ca}} \times \left(\Phi_{\text{Gut2Plas}-\text{Ca0}} + V_{\text{max}-\text{CtriolGut}} \times \frac{[\text{Ctriol}]^{\gamma_{\text{CtriolGut}}}}{(K_{\text{CtriolGut}-\text{Ca}})^{\gamma_{\text{CtriolGut}}} + [\text{Ctriol}]^{\gamma_{\text{CtriolGut}}}} \right) \tag{D.12}$$

$$\Phi_{\text{Plas2Bone}-\text{Ca}} = \xi_{15} \times [\text{Ca}] \times (1 - \nu_{\text{J15}} + \nu_{\text{J15}} \times \text{HAp}) \tag{D.13}$$

$$\Phi_{\text{Bone2Plas}-\text{Ca}} = \text{HAp} \times \left(\Phi_{\text{Bone2Plas}-\text{Ca0}} + V_{\text{max,OC}-\text{Ca}} \times \frac{\text{OC}^{\gamma_{\text{OCCa}}}}{K_{\text{OC}}^{\gamma_{\text{OCCa}}} + \text{OC}^{\gamma_{\text{OCCa}}}} \right) \tag{D.14}$$

$$\Phi_{\text{Urine}-\text{Ca}} = \kappa_{Ca} \times \Phi_{\text{GFR}} \times [\text{Ca}] - \Phi_{\text{Reabs}-\text{Ca}} \tag{D.15}$$

$$\tag{D.16}$$

 where

$$\Phi_{\text{Reabs}-\text{Ca}}^{\text{max}} = \Phi_{\text{GFR}} \times [\text{Ca}] \times (1 + m_{\text{EST}}(E - 1)) \tag{D.17}$$

$$\Phi_{\text{Reabs}-\text{Ca}} = \Phi_{\text{Reabs}-\text{Ca}}^{\text{max}} \times \frac{V_{\text{max}-\text{Ca}} \times [\text{Ca}]}{K_{\text{reabs}-\text{Ca}} + [\text{Ca}]} \times \frac{V_{\text{max}-\text{PTHreabs}} \times [\text{PTH}]}{K_{\text{PTH}-\text{reabs}} + [\text{PTH}]} \tag{D.18}$$

 ## D.3   Parathyroid hormone

$$\frac{d[\text{PTH}]}{dt} = S_{\text{PTH}} \times H_{\text{AT1R,PTH}}^{+} - k_{\text{deg,PTH}} \times [\text{PTH}] \tag{D.19}$$

$$S_{\text{PTH}} = H_{\text{CaSR,PTH}}^{-} \times \frac{\text{PTG}}{0.5} \times \text{PT}_{\text{max}} \tag{D.20}$$

$$\frac{d\text{PTG}}{dt} = k_{\text{prod}-\text{PTG}} - k_{\text{deg}-\text{PTG}} \times \text{PTG} \tag{D.21}$$

$$\frac{d\text{PT}_{\text{max}}}{dt} = \text{PT}_{\text{max}-\text{base}} \times H_{\text{Ctriol,PTmax}}^{-} \times H_{\text{Phos,PTmax}}^{+} - k_{\text{deg}-\text{PTmax}} \times \text{PT}_{\text{max}} \tag{D.22}$$

 ## D.4   Calcitriol

$$\frac{d[\text{AOH}]}{dt} = k_{\text{prod}-\text{AOH}} - k_{\text{deg}} \times [\text{AOH}] \tag{D.23}$$

 with

$$k_{\text{prod}-\text{AOH}} = k_{\text{base}-\text{AOH}} \times H_{\text{PTH,AOH}}^{+} \times H_{\text{Phos,AOH}}^{-} \tag{D.24}$$

$$\frac{d[\text{Ctriol}]}{dt} = \text{AOH} - k_{\text{deg}-\text{Ctriol}} \times [\text{Ctriol}] \tag{D.25}$$

 ## D.5   Phosphate regulation

$$\frac{d[\text{PhosECC}]}{dt} = \Phi_{\text{Bone2Plas}-\text{Phos}} - \Phi_{\text{Plas2Bone}-\text{Phos}} - \Phi_{\text{Urine}-\text{Phos}}$$
$$+ \Phi_{\text{Gut2Plas}-\text{Phos}} - \Phi_{\text{ECC2Int}-\text{Phos}} + \Phi_{\text{Int2ECC}-\text{Phos}} \tag{D.26}$$

$$\frac{d[\text{PhosInt}]}{dt} = \Phi_{\text{ECC2Int}-\text{Phos}} - \Phi_{\text{Int2ECC}-\text{Phos}} \tag{D.27}$$

654 with

$$\Phi_{\text{Gut2Plas-Phos}} = I_{\text{Phos}} \times \left(\Phi_{\text{Gut2Plas-Phos0}} + V_{\text{max-CtriolGutPhos}} \times \frac{[\text{Ctriol}]^{\gamma_{\text{CtriolGut}}}}{(K_{\text{CtriolGut-Ca}})^{\gamma_{\text{CtriolGut}}} + [\text{Ctriol}]^{\gamma_{\text{CtriolGut}}}} \right) \tag{D.28}$$

$$\Phi_{\text{Urine-Phos}} = \kappa_{\text{phos}} \times \Phi_{\text{GFR}} \times [\text{ECCPhos}] - \Phi_{\text{Phos-reabs}} \tag{D.29}$$

$$\Phi_{\text{Plas2Bone-Phos}} = \eta_{\text{bone-phos}} \times \Phi_{\text{Plas2Bone-Ca}} \tag{D.30}$$

$$\Phi_{\text{Bone2Plas-Phos}} = \eta_{\text{bone-phos}} \times \Phi_{\text{Bone2Plas-Ca}} \tag{D.31}$$

655 where

$$\Phi_{\text{Phos-reabs}} = \Phi_{\text{Phos-reabs0}} \times \Phi_{\text{GFR}} \tag{D.32}$$

D.6 Bone remodeling

$$\frac{d\text{OC}}{dt} = k_{\text{prod-OC}} \times H^{+}_{\text{RANK-RANKL}} \times \text{OC}_{\text{base}}$$
$$- k_{\text{deg-OC}} \times H^{+}_{\text{TGFBact,OC}} \times H^{-}_{\text{RNK-RNKL}} \times \text{OC} \tag{D.33}$$

$$\frac{d\text{pOB}}{dt} = k_{\text{prod-pOB}} \times \left(\frac{1}{E}\right)^{\gamma_{\text{pOB,EST}}} - k_{\text{deg,pOB}} \times H^{+}_{\text{TGFBact,pOB}} \times \text{pOB} \tag{D.34}$$

$$\frac{d\text{OB}}{dt} = k_{\text{prod-OB}} \times H^{-}_{\text{TGFBact,OB}} \times \text{pOB} - k_{\text{deg-OB}} \times \text{OB} \tag{D.35}$$

$$\frac{d\text{RNK}}{dt} = -k_3 \times \text{RNK} \times \text{L} + k_4 \times \text{RNK} - \text{RNKL}$$
$$+ k_{\text{prod-RNK}} \times H^{+}_{\text{TBFBact,RNK}} - k_{\text{deg-RNK}} \times \text{RNK} \tag{D.36}$$

$$\frac{d\text{RNKL}}{dt} = -k_3 \times \text{RNK} \times \text{L} + k_4 \times \text{RNK} - \text{RNKL} \tag{D.37}$$

$$+ k_{\text{prod-L}} \times \left(\frac{\text{OB}}{\text{OB}_0}\right)^{\gamma_{\text{OBeff}}} \times H^{+}_{\text{PTH,L}} \times H^{+}_{\text{AT1R,RANKL}}$$
$$- k_{\text{deg-L}} \times \text{RNKL} - k_1 \times \text{OPG} \times \text{L} + k_2 \times \text{RNK} - \text{OPG} \tag{D.38}$$

$$\frac{d\text{RNK} - \text{RNKL}}{dt} = k_3 \times \text{RNK} \times \text{L} - k_4 \times \text{RNK} - \text{RNKL} \tag{D.39}$$

$$\frac{d\text{OPG}}{dt} = -k_1 \times \text{OPG} \times \text{RNKL} + k_2 \times \text{RNK} - \text{OPG} +$$
$$k_{\text{prod-OPG}} \times H^{+}_{\text{PTH,OPG}} \times H^{+}_{\text{AT1R,OPG}} - k_{\text{deg-OPG}} \times \text{OPG} \tag{D.40}$$

$$\frac{d\text{RNK} - \text{OPG}}{dt} = k_1 \times \text{OPG} \times RNKL - k_2 \times \text{RNK} - \text{OPG} \tag{D.41}$$

$$\frac{d\text{TGFB}}{dt} = k_{\text{prod-TGF}} \times \left(\frac{\text{OB}}{\text{OB}_{\text{base}}}\right)^{\gamma_{\text{OBtgf}}} \times \left(\frac{1}{E}\right)^{\gamma_{\text{TGFB,E}}} \tag{D.42}$$

$$- k_{\text{act,TGFB}} \times \text{TGFB} \times \left(\frac{\text{OC}}{\text{OC}_{\text{base}}}\right)^{\gamma_{\text{OCtgf}}} \times E^{\gamma_{\text{tgfbactE}}} \tag{D.43}$$

$$\frac{d\text{TGFB}_{\text{active}}}{dt} = k_{\text{act,TGFB}} \times \text{TGFB} \times \left(\frac{\text{OC}}{\text{OC}_{\text{base}}}\right)^{\gamma_{\text{OCtgf}}} \times E^{\gamma_{\text{tgfbactE}}}$$
$$- k_{\text{deg-TGFact}} \times \text{TGFB}_{\text{active}} \tag{D.44}$$

$$\frac{d\text{BMD}}{dt} = k_{\text{BMD-form}} \times \left(\frac{\text{OB}}{\text{OB}_{\text{base}}}\right)^{\gamma_{\text{OBform}}} - k_{\text{BMD-res}} \times \left(\frac{\text{OC}}{\text{OC}_{\text{base}}}\right)^{\gamma_{\text{OBres}}} \times \text{BMD} \tag{D.45}$$

D.7 List of variable names

- AGT: angiotensinogen

- Ang I: angiotensin I

- Ang II: angiotensin II

- Ang (1-7): angiotensin (1-7)

- Ang IV: angiotensin IV

- AOH: 1α-hydroxylase

- AT1R-AngII: angiotensin type 1 receptor bound Ang II

- AT2R-AngII: angiotensin type 2 receptor bound Ang II

- BMD: bone mineral density

- Ca: plasma calcium

- Ctriol: calcitriol in plasma

- E: normalized estrogen level

- OB: osteoblasts

- OC: osteoclasts

- OPG: osteoprotegerin

- PhosECC: extracellular phosphate

- PhosInt: intracellular phosphate

- pOB: pre-osteoblasts

- PRA: plasma renin activity

- PRC: plasma renin concentration

- $\mathrm{PT_{max}}$: maximal parathyroid gland secretion

- PTG: parathyroid hormone in the parathyroid glands

- PTH: parathyroid hormone

- RNK: RANK

- RNKL: RANKL

- TGFB: latent $\mathrm{TGF}\beta$

- $\mathrm{TGFB_{active}}$: active $\mathrm{TGF}\beta$

References

[1] Allen MR, Burr DB. Chapter 4 - Bone Modeling and Remodeling. In: Burr DB, Allen MR, editors. Basic and Applied Bone Biology. San Diego: Academic Press; 2014. p. 75-90.

[2] Cook CV, Lighty AM, Smith BJ, Ford Versypt AN. A Review of Mathematical Modeling of Bone Remodeling from a Systems Biology Perspective. Frontiers in Systems Biology. 2024 Apr;4.

[3] Melton LJ III, Chrischilles EA, Cooper C, Lane AW, Riggs BL. Perspective How Many Women Have Osteoporosis? Journal of Bone and Mineral Research. 1992 Sep;7(9):1005-10.

[4] Holloway-Kew KL, Betson AG, Anderson KB, Gaston J, Kotowicz MA, Liao WH, et al. Association between Bone Measures and Use of Angiotensin-Converting Enzyme Inhibitors or Angiotensin II Receptor Blockers. Archives of Osteoporosis. 2021 Sep;16(1):137.

[5] Renkema KY, Alexander RT, Bindels RJ, Hoenderop JG. Calcium and Phosphate Homeostasis: Concerted Interplay of New Regulators. Annals of Medicine. 2008 Jan;40(2):82-91.

[6] Vaidya A, Brown JM, Williams JS. The Renin–Angiotensin–Aldosterone System and Calcium-Regulatory Hormones. Journal of Human Hypertension. 2015 Sep;29(9):515-21.

[7] Shimizu H, Nakagami H, Osako MK, Hanayama R, Kunugiza Y, Kizawa T, et al. Angiotensin II Accelerates Osteoporosis by Activating Osteoclasts. The FASEB Journal. 2008;22(7):2465-75.

[8] Gersh FL, O'Keefe JH, Lavie CJ, Henry BM. The Renin-Angiotensin-Aldosterone System in Postmenopausal Women: The Promise of Hormone Therapy. Mayo Clinic Proceedings. 2021 Dec;96(12):3130-41.

[9] Ibrahim MM. RAS Inhibition in Hypertension. Journal of Human Hypertension. 2006 Feb;20(2):101-8.

[10] Kim KM, Hwang EJ, Lee S, Yoon JH. The Impact of Renin-Angiotensin System Inhibitors on Bone Fracture Risk: A Nationwide Nested Case-Control Study. BMC Musculoskeletal Disorders. 2024 Jan;25(1):3.

[11] Saavedra JM. Angiotensin Receptor Blockers Are Not Just for Hypertension Anymore. Physiology. 2021 May;36(3):160-73.

[12] Wu J, Wang M, Guo M, Du XY, Tan XZ, Teng FY, et al. Angiotensin Receptor Blocker Is Associated with a Lower Fracture Risk: An Updated Systematic Review and Meta-Analysis. International Journal of Clinical Practice. 2022;2022:7581110.

[13] Peterson MC, Riggs MM. A Physiologically Based Mathematical Model of Integrated Calcium Homeostasis and Bone Remodeling. Bone. 2010 Jan;46(1):49-63.

[14] Riggs M, Bennetts M, van der Graaf P, Martin S. Integrated Pharmacometrics and Systems Pharmacology Model-Based Analyses to Guide GnRH Receptor Modulator Development for Management of Endometriosis. CPT: Pharmacometrics & Systems Pharmacology. 2012;1(10):11.

[15] Riggs MM, Peterson MC, Gastonguay MR. Multiscale Physiology-Based Modeling of Mineral Bone Disorder in Patients With Impaired Kidney Function. The Journal of Clinical Pharmacology. 2012;52(S1):45S-53S.

[16] Gaweda AE, McBride DE, Lederer ED, Brier ME. Development of a Quantitative Systems Pharmacology Model of Chronic Kidney Disease: Metabolic Bone Disorder. American Journal of Physiology-Renal Physiology. 2021 Feb;320(2):F203-11.

[17] Lo A, Beh J, De Leon H, Hallow MK, Ramakrishna R, Rodrigo M, et al. Using a Systems Biology Approach to Explore Hypotheses Underlying Clinical Diversity of the Renin Angiotensin System and the Response to Antihypertensive Therapies. In: Kimko HHC, Peck CC, editors. Clinical Trial Simulations: Applications and Trends. AAPS Advances in the Pharmaceutical Sciences Series. New York, NY: Springer; 2011. p. 457-82.

[18] Hallow KM, Lo A, Beh J, Rodrigo M, Ermakov S, Friedman S, et al. A Model-Based Approach to Investigating the Pathophysiological Mechanisms of Hypertension and Response to Antihypertensive Therapies: Extending the Guyton Model. American Journal of Physiology-Regulatory, Integrative and Comparative Physiology. 2014 May;306(9):R647-62.

[19] Leete J, Layton AT. Sex-Specific Long-Term Blood Pressure Regulation: Modeling and Analysis. Computers in biology and medicine. 2019;104:139-48.

[20] Jörg DJ, Fuertinger DH, Cherif A, Bushinsky DA, Mermelstein A, Raimann JG, et al. Modeling Osteoporosis to Design and Optimize Pharmacological Therapies Comprising Multiple Drug Types. eLife. 2022 Aug;11:e76228.

[21] Sowers MR, Zheng H, McConnell D, Nan B, Harlow SD, Randolph JF Jr. Estradiol Rates of Change in Relation to the Final Menstrual Period in a Population-Based Cohort of Women. The Journal of Clinical Endocrinology & Metabolism. 2008 Oct;93(10):3847-52.

[22] Prior JC. Women's Reproductive System as Balanced Estradiol and Progesterone Actions—A Revolutionary, Paradigm-Shifting Concept in Women's Health. Drug Discovery Today: Disease Models. 2020 Dec;32:31-40.

[23] Fischer M. Renin Angiotensin System and Gender Differences in the Cardiovascular System. Cardiovascular Research. 2002 Feb;53(3):672-7.

[24] Schunkert H, Danser AHJ, Hense HW, Derkx FHM, Ku¨rzinger S, Riegger GAJ. Effects of Estrogen Replacement Therapy on the Renin-Angiotensin System in Postmenopausal Women. Circulation. 1997 Jan;95(1):39-45.

[25] Gordon MS, Chin WW, Shupnik MA. Regulation of Angiotensinogen Gene Expression by Estrogen. Journal of Hypertension. 1992 Apr;10(4):361.

[26] Gallagher PE, Li P, Lenhart JR, Chappell MC, Brosnihan KB. Estrogen Regulation of Angiotensin-Converting Enzyme mRNA. Hypertension. 1999 Jan;33(1):323-8.

[27] Brosnihan KB, Li P, Ganten D, Ferrario CM. Estrogen Protects Transgenic Hypertensive Rats by Shifting the Vasoconstrictor-Vasodilator Balance of RAS. American Journal of Physiology-Regulatory, Integrative and Comparative Physiology. 1997 Dec;273(6):R1908-15.

[28] Brosnihan KB, Weddle D, Anthony MS, Heise C, Li P, Ferrario CM. Effects of Chronic Hormone Replacement on the Renin–Angiotensin System in Cynomolgus Monkeys. Journal of Hypertension. 1997 Jul;15(7):719.

[29] Hussien NI, El-kerdasy HI, Sorour SM, Shoman AA. Chronic Oestrogen Deficiency Induced by Ovariectomy May Cause Lung Fibrosis through Activation of the Renin-Angiotensin System in Rats. Archives of Physiology and Biochemistry. 2022 Mar;128(2):290-9.

[30] Joseph D, Kilfeather S, O'Brien E, O'Malley K, D'Amore A, Nussberger J. Effects of Aging and Hypertension on Plasma Angiotensin II and Platelet Angiotensin II Receptor Density. American Journal of Hypertension. 1992 Oct;5(10):687-93.

[31] Xu X, Xiao JC, Luo LF, Wang S, Zhang JP, Huang JJ, et al. Effects of Ovariectomy and 17β-Estradiol Treatment on the Renin–Angiotensin System, Blood Pressure, and Endothelial Ultrastructure. International Journal of Cardiology. 2008 Nov;130(2):196-204.

[32] Sullivan JC, Semprun-Prieto L, Boesen EI, Pollock DM, Pollock JS. Sex and Sex Hormones Influence the Development of Albuminuria and Renal Macrophage Infiltration in Spontaneously Hypertensive Rats. American Journal of Physiology-Regulatory, Integrative and Comparative Physiology. 2007 Oct;293(4):R1573-9.

[33] Ichikawa J, Sumino H, Ichikawa S, Ozaki M. Different Effects of Transdermal and Oral Hormone Replacement Therapy on the Renin-Angiotensin System, Plasma Bradykinin Level, and Blood Pressure of Normotensive Postmenopausal Women. American Journal of Hypertension. 2006 Jul;19(7):744-9.

[34] Sampson AK, Hilliard LM, Moritz KM, Thomas MC, Tikellis C, Widdop RE, et al. The Arterial Depressor Response to Chronic Low-Dose Angiotensin II Infusion in Female Rats Is Estrogen Dependent. American Journal of Physiology-Regulatory, Integrative and Comparative Physiology. 2012 Jan;302(1):R159-65.

[35] Nickenig G, Bäumer AT, Grohè C, Kahlert S, Strehlow K, Rosenkranz S, et al. Estrogen Modulates AT1 Receptor Gene Expression In Vitro and In Vivo. Circulation. 1998 Jun;97(22):2197-201.

[36] Bonate PL. Pharmacokinetic-Pharmacodynamic Modeling and Simulation. Boston, MA: Springer US; 2011.

[37] Grant FD, Mandel SJ, Brown EM, Williams GH, Seely EW. Interrelationships between the Renin-Angiotensin-Aldosterone and Calcium Homeostatic Systems. The Journal of Clinical Endocrinology & Metabolism. 1992 Oct;75(4):988-92.

[38] Hannan FM, Kallay E, Chang W, Brandi ML, Thakker RV. The Calcium-Sensing Receptor in Physiology and in Calcitropic and Noncalcitropic Diseases. Nature Reviews Endocrinology. 2019 Jan;15(1):33-51.

[39] Ortiz-Capisano MC, Reddy M, Mendez M, Garvin JL, Beierwaltes WH. Juxtaglomerular Cell CaSR Stimulation Decreases Renin Release via Activation of the PLC/IP3 Pathway and the Ryanodine Receptor. American Journal of Physiology-Renal Physiology. 2013 Feb;304(3):F248-56.

[40] Ortiz-Capisano MC, Ortiz PA, Garvin JL, Harding P, Beierwaltes WH. Expression and Function of the Calcium-Sensing Receptor in Juxtaglomerular Cells. Hypertension. 2007 Oct;50(4):737-43.

[41] Tomaschitz A, Pilz S, Ritz E, Grammer T, Drechsler C, Boehm BO, et al. Independent Association between 1,25-Dihydroxyvitamin D, 25-Hydroxyvitamin D and the Renin–Angiotensin System: The Ludwigshafen Risk and Cardiovascular Health (LURIC) Study. Clinica Chimica Acta. 2010 Sep;411(17):1354-60.

[42] Qian G, Mahdi A. Sensitivity Analysis Methods in the Biomedical Sciences. Mathematical Biosciences. 2020 May;323:108306.

[43] Saltelli A, Ratto M, Andres T, Campolongo F, Cariboni J, Gatelli D, et al. Global Sensitivity Analysis: The Primer. John Wiley & Sons; 2008.

[44] Renardy M, Hult C, Evans S, Linderman JJ, Kirschner DE. Global Sensitivity Analysis of Biological Multiscale Models. Current Opinion in Biomedical Engineering. 2019 Sep;11:109-16.

[45] Sobolʹ IM. Global Sensitivity Indices for Nonlinear Mathematical Models and Their Monte Carlo Estimates. Mathematics and Computers in Simulation. 2001 Feb;55(1):271-80.

[46] Alexander RT, Dimke H. Effects of Parathyroid Hormone on Renal Tubular Calcium and Phosphate Handling. Acta Physiologica. 2023;238(1):e13959.

[47] Looker AC, Wahner HW, Dunn WL, Calvo MS, Harris TB, Heyse SP, et al. Updated Data on Proximal Femur Bone Mineral Levels of US Adults. Osteoporosis International. 1998 Aug;8(5):468-90.

[48] Solomon DH, Mogun H, Garneau K, Fischer MA. Risk of Fractures in Older Adults Using Antihypertensive Medications. Journal of Bone and Mineral Research. 2011 Jul;26(7):1561-7.

[49] McClung MR. The Relationship between Bone Mineral Density and Fracture Risk. Current Osteoporosis Reports. 2005 Jun;3(2):57-63.

[50] Flores VA, Pal L, Manson JE. Hormone Therapy in Menopause: Concepts, Controversies, and Approach to Treatment. Endocrine Reviews. 2021 Dec;42(6):720-52.

[51] Vigneswaran K, Hamoda H. Hormone Replacement Therapy – Current Recommendations. Best Practice & Research Clinical Obstetrics & Gynaecology. 2022 May;81:8-21.

[52] Foessl I, Dimai HP, Obermayer-Pietsch B. Long-Term and Sequential Treatment for Osteoporosis. Nature Reviews Endocrinology. 2023 Jul:1-14.

[53] Stadt M, Layton AT. Modulation of Blood Pressure by Dietary Potassium and Sodium: Sex Differences and Modeling Analysis. American Journal of Physiology-Renal Physiology. 2025 Mar;328(3):F406-17.

[54] Francis A, Harhay MN, Ong ACM, Tummalapalli SL, Ortiz A, Fogo AB, et al. Chronic Kidney Disease and the Global Public Health Agenda: An International Consensus. Nature Reviews Nephrology. 2024 Jul;20(7):473-85.

[55] Gaweda AE, Lederer ED, Brier ME. Artificial Intelligence–Guided Precision Treatment of Chronic Kidney Disease–Mineral Bone Disorder. CPT: Pharmacometrics & Systems Pharmacology. 2022;11(10):1305-15.

[56] Gebru Y, Diao TY, Pan H, Mukwaya E, Zhang Y. Potential of RAS Inhibition to Improve Metabolic Bone Disorders. BioMed Research International. 2013 Jan;2013(1):932691.

[57] Chapuy MC, Durr F, Chapuy P. Age-Related Changes in Parathyroid Hormone and 25 Hydroxycholecalciferol Levels. Journal of Gerontology. 1983 Jan;38(1):19-22.

[58] Bhattarai HK, Shrestha S, Rokka K, Shakya R. Vitamin D, Calcium, Parathyroid Hormone, and Sex Steroids in Bone Health and Effects of Aging. Journal of Osteoporosis. 2020 Jun;2020:e9324505.

[59] Yoon HE, Choi BS. The Renin-Angiotensin System and Aging in the Kidney. The Korean Journal of Internal Medicine. 2014 May;29(3):291-5.

[60] Hilliard LM, Sampson AK, Brown RD, Denton KM. The "His and Hers" of the Renin-Angiotensin System. Current Hypertension Reports. 2013 Feb;15(1):71-9.

[61] Khalil R, Kim NR, Jardi F, Vanderschueren D, Claessens F, Decallonne B. Sex Steroids and the Kidney: Role in Renal Calcium and Phosphate Handling. Molecular and Cellular Endocrinology. 2018 Apr;465:61-72.

[62] Hakimi S, Dutta P, Layton AT. Coupling of Renal Sodium and Calcium Transport: A Modeling Analysis of Transporter Inhibition and Sex Differences. American Journal of Physiology-Renal Physiology. 2023 Nov;325(5):F536-51.

[63] Rianon N, Ambrose CG, Pervin H, Garcia M, Mama SK, Schwartz AV, et al. Long-Term Use of Angiotensin-Converting Enzyme Inhibitors Protects against Bone Loss in African-American Elderly Men. Archives of Osteoporosis. 2017 Oct;12(1):94.

[64] Layton AT. We Are All Different: Modeling Key Individual Differences in Physiological Systems. Mathematical Biosciences. 2024 Dec;378:109338.

[65] Song S, Cai X, Hu J, Zhu Q, Shen D, Ma H, et al. Effectiveness of Spironolactone in Reducing Osteoporosis and Future Fracture Risk in Middle-Aged and Elderly Hypertensive Patients. Drug Design, Development and Therapy. 2024 Dec;18:2215-25.

[66] Sernia C, Huang H, Nguyuen K, Li YH, Hsu S, Chen M, et al. Bone Homeostasis: An Emerging Role for the Renin-Angiotensin System. In: Leung PS, editor. Frontiers in Research of the Renin-Angiotensin System on Human Disease. Dordrecht: Springer Netherlands; 2007. p. 179-95.

[67] Smith D, Layton A. The Intrarenal Renin-Angiotensin System in Hypertension: Insights from Mathematical Modelling. Journal of Mathematical Biology. 2023 Mar;86(4):58.

[68] Sims ST, Heather AK. Myths and Methodologies: Reducing Scientific Design Ambiguity in Studies Comparing Sexes and/or Menstrual Cycle Phases. Experimental Physiology. 2018;103(10):1309-17.

[69] Abo S, Smith D, Stadt M, Layton A. Modelling Female Physiology from Head to Toe: Impact of Sex Hormones, Menstrual Cycle, and Pregnancy. Journal of Theoretical Biology. 2022 May;540:111074.

[70] Napso T, Yong HEJ, Lopez-Tello J, Sferruzzi-Perri AN. The Role of Placental Hormones in Mediating Maternal Adaptations to Support Pregnancy and Lactation. Frontiers in Physiology. 2018 Aug;9:1091.

[71] Kovacs CS. Maternal Mineral and Bone Metabolism During Pregnancy, Lactation, and Post-Weaning Recovery. Physiological Reviews. 2016 Apr;96(2):449-547.

[72] Stadt MM, Layton AT. Mathematical Modeling of Calcium Homeostasis in Female Rats: An Analysis of Sex Differences and Maternal Adaptations. Journal of Theoretical Biology. 2023 Sep;572:111583.