The reticulocyte restriction: invasion ligand RBP1a of *Plasmodium vivax* targets human TfR1, prohibitin-2, and basigin

Jessica Molina-Franky[1-4], Daniel Röth[1], Monica Ararat-Sarria[3,5], Manuel Alfonso Patarroyo[3,6*], Markus Kalkum[1*]

[1]Department of Immunology and Theranostics, Arthur Riggs Diabetes and Metabolism Research Institute, Beckman Research Institute of the City of Hope, Duarte, California, United States of America

[2]Global Scholars Program, Beckman Research Institute of the City of Hope, Duarte, California, United States of America

[3]Grupo de Investigación Básica en Biología Molecular e Inmunología (GIBBMI), Fundación Instituto de Inmunología de Colombia (FIDIC), Bogotá, Colombia

[4]PhD Program in Biotechnology, Universidad Nacional de Colombia, Bogotá, Colombia

[5]Health Sciences Faculty, Universidad de Ciencias Aplicadas y Ambientales (U.D.C.A), Bogotá, Colombia

[6]Microbiology Department, Faculty of Medicine, Universidad Nacional de Colombia, Bogotá, Colombia

[*] Corresponding authors:

E-mail: mkalkum@coh.org (MK)

E-mail: mapatarr.fidic@gmail.com (MAP)

Abstract

Plasmodium vivax is the most widespread malaria species with predominance in Latin America and Southeast Asia. Devising effective controls against *P. vivax* infections presents unique challenges due to the parasite's exclusive invasion of reticulocytes, a transient developmental stage of red blood cells. The ephemeral nature and limited availability of reticulocytes complicate efforts to establish continuous *in vitro* cultures to study the parasite's biology and receptor-ligand interactions. Here, the potential of the erythroid cell lines JK-1 and BEL-A as alternatives to reticulocytes was confirmed through successful parasite invasion, marking the first report of *P. vivax* entry to BEL-A cells. Furthermore, LC-MS was used to quantitatively compare the membrane proteomes of these cell lines with those of reticulocytes and mature erythrocytes. This analysis revealed significant similarities between the membrane proteomes of JK-1, BEL-A, and reticulocytes. In addition to the known *Plasmodium* receptors, transferrin receptor protein 1 (TfR1 or CD71), CD98hc, and basigin (BSG), potential receptor candidates involved in the parasite's invasion pathway were identified, including prohibitin-2 (PHB2), CAT-1 (SLC7A1), ATB(0) (SLC1A5), CD36, integrin beta-1 (ITGB1), and Metal transporter CNNM3. Proximity labeling using TurboID enabled the identification of specific interactions between PvRBP1a$_{158\text{-}650}$ and TfR1, BSG, and prohibitin-2 in the erythroid cell lines. Notably, this is the first report of prohibitin-2 as a receptor for *P. vivax*. These findings advance the understanding of *P. vivax* receptor-ligand interactions and underscore the potential of JK-1 and BEL-A cell lines as alternative models for the study of parasite biology.

Author Summary

Author Summary

Plasmodium vivax is a major cause of malaria, particularly in Latin America and Southeast

Asia. Unlike other malaria parasites, it infects only immature red blood cells

(reticulocytes), which are rare and short-lived, making laboratory studies challenging. No

continuous *in vitro* culture system currently exists for *P. vivax*, limiting research into its

invasion mechanisms. In this study, we evaluated two erythroid cell lines, JK-1 and BEL-

A, as potential reticulocyte surrogates. We demonstrate that *P. vivax* can invade both cell

types, including BEL-A cells, which had not previously been tested. Comparative analysis

revealed key differences in surface protein expression between these cell lines and

reticulocytes, compared to mature red blood cells, identifying seven potential parasite-

receptor proteins. We further applied a proximity-based protein tagging approach to

identify host-parasite interactions mediated by a specific *P. vivax* protein ligand, RBP1a.

This technique revealed a potential role for prohibitin-2, TfR1, and basigin in mediating *P.*

vivax entry into red blood cells by RBP1a binding. The role of prohibitin-2 as a

Plasmodium receptor is a novel Discovery. Our results highlight JK-1 and BEL-A cells as

valuable tools for *P. vivax* research and suggest new avenues for investigating host-

pathogen interactions, with implications for malaria treatment and vaccine development.

Introduction

Malaria remains a major global health issue, with 263 million cases reported in 2023.

Plasmodium vivax, one of the six human-infecting *Plasmodium* species, is noteworthy

for its global distribution. In the Americas, most malaria cases are due to *P. vivax*

(72.1% in 2023) [1]. Its blood stage form, the merozoites, interact with red blood cell

(RBC) receptors leading to RBC invasion [2–4]. However, *P. vivax* exclusively infects

reticulocytes [5], immature, short-lived precursors of erythrocytes, posing a significant

challenge to research progress. In contrast, *P. falciparum* has been extensively

researched due to the availability of a well-established *in vitro* culture system for over 40

years [6]. This critical disparity highlights the urgent need to develop alternative

research models for *P. vivax*.

The mechanism of *P. vivax* reticulocyte invasion remains unclear. It was previously

believed that *P. vivax* exclusively targeted reticulocytes through the Duffy antigen

receptor for chemokines (DARC) and Duffy binding protein (PvDBP) interaction [7], as

individuals with the Fy(a-b-) mutation in West Africa were resistant to the infection.

However, DARC is present on both reticulocytes and erythrocytes, and *P. vivax*

infections have been documented in Duffy-negative populations [8–10]. Current

evidence suggests that the reticulocyte-binding protein (RBP) family may play a role,

interacting with transferrin receptor 1 (TfR1) and CD98 heavy chain (SLC3A2), which

are lost during the maturation of reticulocytes to erythrocytes [2,3,11]. This implies that

*P. vivax (*Pv*)*RBP family proteins may specifically target receptors unique to the

reticulocyte membrane. Our previous studies on PvRBP1 of the *P. vivax* strain Belem

(GenBank AAA29743.3) identified eleven high-affinity reticulocyte binding peptides

(HABPs) corresponding to residues 158–653 of PvRBP1a in the *P. vivax* Salvador I

strain (GenBank AAS85749.1). Among these, HABP 3742 (KLLGEEISEVSHLYV) and

HABP 3459 (KEILDKMAKKVHYLK) exhibited dissociation constants (*K*d) of 131 nM

and 155 nM, respectively [12]. Additionally, an extracellular portion of PvRBP1a,

residues 157-650, binds strongly (~50%) to reticulocytes and moderately (~20%) to

erythrocytes [13]. The identity of PvRBP1a$_{157-650}$ binding-receptors within the

reticulocyte membrane has been unclear. Therefore, this study evaluated the

erythroleukemic cell line JK-1 [14] and the Bristol Erythroid Line Adult (BEL-A) [15] as

surrogates for reticulocytes, examining their susceptibility to *P. vivax* invasion. A

comparison of the cell lines' membrane proteomes revealed similarities with those of

reticulocytes, and dissimilarities with erythrocyte membrane proteomes, thereby

identifying potential *P. vivax* receptors. Furthermore, TurboID proximity labelling implied

specific interactions of PvRBP1a$_{158-650}$ with prohibitin-2 (PHB2), TfR1, and basigin

(BSG). These interactions were confirmed by ELISA, highlighting key molecular

determinants of *P. vivax*'s reticulocyte tropism.

Results

P. vivax can invade the erythroid cell lines BEL-A and JK-1

Cultured BEL-A and JK-1 cells were successfully invaded by *P. vivax* from a validated

mono-infected malaria patient's blood sample. The experiment required forgoing

enrichment of the RBCs to 4.0% parasitemia (Fig 1 A-E). Parasite invasion was

confirmed by immunofluorescence microscopy, detecting the intracellular presence of *P.*

vivax lactate dehydrogenase (PvLDH), which all blood stages of the parasite are known

to express [16]. PvLDH was detected in the positive control of infected reticulocytes,

and in BEL-A and JK-1 cells, incubated with infected reticulocytes (Fig 2, FITC). The

PvLDH signal was absent from non-infected control cells. Moreover, the presence of

hemozoin (Hz) pigment, characteristic for hemoglobin consumption by *Plasmodia* within

infected RBCs [17], was observed (Fig 2, bright field, and merge). The dark Hz pigment

was visible inside the parasite-infected nucleated erythroid cells as well as in infected

reticulocytes that originated from the donor. Hz is an insoluble, crystallized digestion

product of heme derived from the digestion of hemoglobin by malaria parasites,

containing heme-derived β-hematin, which neutralizes the toxicity of free heme released

after parasite invasion through a digestive process that involves the digestive vacuole

structure [18]. On days 3 to 6 post-infection, no parasite-infected erythroid cells were

observed, and cell mortality had substantially increased. Therefore, the experiment was

stopped on day 6.

Fig 1. *P. vivax* **mono-infected reticulocytes.** (A) Three representative micrographs of

P. vivax-infected enriched reticulocytes (red arrows) from a malaria patient. Scale bars

are 3 µm. (B-D) Species-specific nested PCR for the small subunit 18S ribosomal RNA.

Lane M, 100 bp marker; Lane *PvSx*, sample used in the infection of erythroid cells;

Lanes *Pv*, *Pf*, and *Pm* belong to the positive controls for each species; lane NCh,

negative control using genomic DNA from a healthy human; lane NC, negative control

using water instead of genomic DNA. (B) *P. vivax* (~120 bp amplicon). (C) *P. falciparum*

(~205 bp amplicon). (D) *P. malariae* (~144 bp amplicon). (E) DNA sequence alignment

of the positive *PvSx* amplicon with the corresponding gene segment of the *P. vivax*

Salvador-1 reference strain (GenBank No. XR_003001206.1) [19], showing 100%

identity.

Fig 2. *P. vivax* invades BEL-A and JK-1 cells. The micrographs show *P. vivax*-

infected reticulocytes as the positive Control (+); infected BEL-A and JK-1 cells at 24

and 48 h, and their non-infected negative Control (-); *P. vivax* lactate dehydrogenase

(FITC green); DNA (DAPI blue); hemozoin crystals (black dots marked with red arrows);

BF (bright field); Scale bars are 5 µm.

Overlapping membrane proteomes reveal potential *P. vivax* invasion receptors

Because *P. vivax* was able to invade the erythroid BEL-A and JK-1 cells, their membranes must contain the same essential receptor molecules as reticulocytes that enable parasite invasion. Furthermore, the membranes of mature erythrocytes are expected to lack these receptors or to express them only at insufficient abundances. Consequently, a quantitative comparison of the membrane proteome of these cells with those of human reticulocytes and erythrocytes identified potential receptors for *P. vivax* merozoite ligands that are most likely responsible for its reticulocyte-restricted invasion.

Stringent isolation procedures were necessary to obtain membrane proteins of pure reticulocytes. The isolated reticulocytes (CD71$^+$, CD45$^-$) used in this proteomic comparison had a purity of 98.4% (S2 Fig). Expression of CD71 is diminished during maturation into fully functional erythrocytes [20]. Simultaneous determination of CD45 negativity was necessary, as CD45$^+$ leukocytes also express CD71, to ensure purity of the isolated reticulocytes.

The BEL-A and JK-1 cells used in the membrane proteomic comparisons were harvested from *in vitro* cultures and exhibited distinct nucleated erythroid maturation stages, including proerythroblasts, basophilic erythroblasts, polychromatic erythroblasts, and orthochromatic erythroblasts (S3 Fig), with slight dominance of the basophilic and polychromatic stages.

In total, 2,100 proteins were identified in BEL-A cells and 2,178 in JK-1 cells. The number of proteins was lower in reticulocytes (1,234) and in erythrocytes (1,347). After filtering this data for membrane proteins (see S1 Text & S4 Fig), 1,530 and 1,595 such

proteins were obtained from BEL-A and JK-1 cell ghosts, respectively, while 846 and

974 proteins were identified for reticulocyte and erythrocyte ghosts.

Changes in membrane protein abundance were assessed by comparing erythroid cell

lines and reticulocytes to mature erythrocytes. The protein abundancies of reticulocytes

clustered better with those of BEL-A and JK-1 cells than with those of erythrocytes (Fig

3A). It was found that compared to erythrocytes 256 proteins were more abundant in

reticulocytes, 1,179 in JK-1, and 1,554 in BEL-A. Of these, 144 identical proteins were

increased in reticulocytes, JK-1, and BEL-A. However, reticulocytes and JK-1 cells

share 18 proteins that are less abundant in erythrocytes, while BEL-A and reticulocytes

have 40 proteins in common, that are less abundant in erythrocytes. Only 54 membrane

proteins with higher abundance than in erythrocytes were identified exclusively in

reticulocytes, 237 in JK-1, and 590 in BEL-A cells (Fig 3B).

Fig 3. Plasma membrane proteomes of reticulocytes resemble those of erythroid

cell lines. (A) Clustering of plasma membrane proteins in JK-1, BEL-A, reticulocytes,

and erythrocytes, measured in triplicate. Log2 intensities. (B) Abundance of intersecting

membrane proteins in reticulocytes, JK-1, and BEL-A cells compared to erythrocytes,

represented in an UpSet plot [21]. 144 proteins share higher abundance among cell

lines and reticulocytes but are reduced in erythrocytes (mauve bar). (C-E) Putative

receptors (blue) and characterized receptors (orange) for *P. vivax* merozoite invasion.

Gene names are displayed instead of protein names for simplicity. The x-axis

represents the $\log_2$ fold change, and the y-axis shows the P-value, indicating statistical

significance.

When comparing the membrane protein abundance in reticulocytes, cell lines, and

erythrocytes, known *P. vivax* receptors such as TfR1 (CD71), CD98hc, ACKR1/DARC,

BSG, CR1, and band 3 (SLC4A1) were identified. TfR1 and CD98hc, which are lost

during reticulocyte maturation to erythrocytes, were significantly more abundant in

reticulocytes and cell lines. In contrast, the other receptors showed higher levels in

erythrocytes (Fig 3C-E).

In silico topological analysis of membrane proteins enriched in reticulocytes and

erythroid cell lines identified several candidates — CD98lc (SLC7A5), high-affinity

cationic amino acid transporter 1 (CAT-1, SLC7A1), neutral amino acid transporter B^0

(ATB(0), SLC1A5), CD36, Integrin β-1 (ITGB1), prohibitin-2 (PHB2), and the metal

transporter CNNM3 — as possessing sizable extracellular regions that are potentially

accessible for interaction with *P. vivax* merozoite ligands (S1 Table & S5 Fig).

The significantly higher abundance of these proteins in reticulocytes, BEL-A, and JK-1

cells compared to erythrocytes (Fig 3C-E & S1 Table) highlights them as potential

candidates for *P. vivax* merozoite protein receptors, which may explain the parasite's

exclusivity for reticulocyte invasion.

Receptors for PvRBP1a$_{158-650}$LTID identified via proximity labeling

To identify potential receptors of the PvRBP1a$_{158-650}$LTID protein, the TurboID proximity

labeling technique was used. This technique enables the biotinylation of proteins that

come into close contact with the fused protein (within 10 nm), facilitating the

identification of their interactions [22,23]. Therefore, the fusion protein PvRBP1a$_{158-}$

$_{650}$LTID and the LTID control were obtained in soluble form, with PvRBP1a$_{158-650}$LTID

having a molecular weight of ~94 kDa and LTID ~36 kDa (Fig 4A). Both proteins

exhibited enzymatic activity and self-biotinylation at 1, 2, and 3 µM (Fig 4B-C).

Fig 4. Proximity labeling with PvRBP1a$_{158-650}$LTID in JK-1, BEL-A, reticulocytes,

and erythrocytes. Recombinantly expressed soluble PvRBP1a$_{158-650}$LTID and LTID by

(A) SDS PAGE (B, C) auto-biotinylation assay with increasing concentrations of each

protein at 1, 2, and 3 µM, detected with IRDye 800 Streptavidin (green bands),

LICORbio molecular weight markers (MW, red bands); (D) biotin labeling of JK-1

(orange) and BEL-A (fuchsia) cells, in presence of PvRBP1a$_{158-650}$LTID or LTID

(negative control, JK-1 - black, and BEL-A - gray) at 1, 2, and 3 µM. (E-G) Flow

cytometry (E) histograms of erythroid cells and human RBCs in presence of

PvRBP1a$_{158-650}$LTID (blue) or LTID (grey), both at 3 µM, demonstrating (F) significant

degrees of cell surface labeling by PvRBP1a$_{158-650}$LTID in erythroid cells and

reticulocytes compared to erythrocytes (**** $p \leq 0.0001$). (G) Receptors to PvRBP1a$_{158-}$

$_{650}$ are sensitive to cell surface treatment with trypsin and chymotrypsin but resistant to

neuraminidase (labeling % as normalized to untreated cells).

PvRBP1a$_{158-650}$LTID exhibits comparable binding to reticulocytes and

erythroid cell lines via a proteinaceous receptor

Proximity biotinylation mediated by TurboID facilitated binding evaluation through the

biotin-streptavidin interaction. The assays showed that PvRBP1a$_{158-650}$LTID binding is

concentration-dependent (Fig 4D). Since the highest percentage of biotin labeling on

the cell surface was obtained at 3 µM, this concentration was selected to analyze

PvRBP1a$_{158-650}$LTID binding to enriched reticulocytes (87.5% purity) (S6 Fig),

erythrocytes, and cell lines. The results showed that PvRBP1a$_{158-650}$ had 80%

biotinylation on the surface of reticulocytes, 4.7% on erythrocytes, and 79.85% and

83.2% on the surface of JK-1 and BEL-A cells, respectively. A statistically significant

difference was found between the cell lines and reticulocytes compared to erythrocytes

(P ≤ 0.0001). However, no significant difference was observed between the cell lines

and reticulocytes (Fig 4E-F). These data suggest that erythroid cell lines exhibit

PvRBP1a$_{158-650}$ binding activity comparable to reticulocytes, indicating that they may

express the receptor for this specific *P. vivax* ligand on their surface. Additionally, no

labelling was detected with LTID, confirming the specificity of the PvRBP1a$_{158-650}$

receptor interaction.

Cell surface labelling with PvRBP1a$_{158-650}$LTID was sensitive to trypsin and

chymotrypsin treatment but resistant to neuraminidase, which removes sialic acid from

glycans that modify proteins in vertebrates (Fig 4G). Therefore, the cell surface receptor

function for PvRBP1a$_{158-650}$ is proteinaceous and not dependent on sialic acid-

terminated glycans.

Enrichment of PvRBP1a$_{158-650}$LTID biotinylated cell surface proteins,

identifies TfR1 and prohibitin-2 as the likely reticulocyte-restricting

receptors

PvRBP1a$_{158-650}$LTID biotinylated membrane proteins from erythroid cells were captured

via streptavidin-affinity and subjected to proteomics analysis, revealing a total of 278

proteins, of which 12 were localized to the plasma membrane. However, the LTID

control contained five of these proteins, leaving seven unique to enrichment after

biotinylation with PvRBP1a$_{158-650}$LTID. Four of these membrane proteins do not possess

extracellular regions, while TfR1, prohibitin-2, and BSG do. Such extracellular regions

should be required to facilitate an interaction with *P. vivax* merozoite ligands (Fig 5A).

Interaction of PvRBP1a$_{158-650}$ was successfully validated by targeted (PRM) LC-MS

analysis for TfR1, BSG, and prohibitin-2 (Fig 5B). It clearly demonstrated that only

PvRBP1a$_{158-650}$LTID biotinylated these three proteins, while LTID did not.

BSG is more abundant in erythrocytes than in reticulocytes and in the erythroid cell

lines. In contrast, TfR1 and prohibitin-2 were significantly less abundant in erythrocytes

(Fig 3C-E). In fact, TfR1 and prohibitin-2 were among the most abundant membrane

proteins in reticulocytes and in the erythroid cell lines JK-1 and BEL-A. These findings

suggest that PvRBP1a$_{158-650}$ likely facilitate the recognition and invasion of reticulocytes

through interaction with TfR1 and prohibitin-2.

Fig 5. Proximity biotin-labeled TfR1 (CD71), Basigin, and Prohibitin-2, are

identified as *P. vivax* receptor candidates after interaction of JK-1 cells with

PvRBP1a$_{158-650}$LTID and validated by targeted LC-MS. (A) Heatmap of labeled

plasma membrane protein intensities in the PvRBP1a$_{158-650}$LITD (fuchsia) and LTID

(yellow) control samples, analyzed in duplicate by DDA LC-MS proteomics. Intensities

are represented on a Z-score scale, where each value was transformed by the number

of standard deviations (SD) from the mean. Topology categorizes proteins by their

cellular localization: intracellular (Inside, dark gray) and those with extracellular domains

(Outside, cyan). They include Shroom3, GPR132 – probable G-protein coupled receptor

132, EEF1A1 – Elongation factor 1-alpha 1, protein S100-A8, protein S100-A6, TfR1,

PHB2 – prohibitin-2, BSG, protein S100-A9, JUP – junction plakoglobin, DSC3 –

Desmocollin-3, and VIM – vimentin. (B) Validated interactions of transferrin receptor 1,

basigin, and prohibitin-2 with PvRBP1a$_{158-650}$LITD, but not with LITD, during the TurboID

procedure, followed by PRM LC-MS. (1) (2) – duplicates. Stacked bars are the sum of

the LC-MS ion chromatographic peak areas of the trypsin digested peptides (colored

boxes) of each protein, indicating the contribution of each peptide to the individual

protein abundance.

PvRBP1a$_{158\text{-}650}$LTID interacts with high-affinity binding to

TfR1, BSG, and prohibitin-2

The titration curves fit well to a single-site binding saturation model. In contrast, the

negative control LTID displayed a nonspecific binding pattern "unstable", corroborating

the specificity of the interactions (Fig 6A). The Kd obtained from the ELISA titration

indicated that PvRBP1a$_{158\text{-}650}$LTID had high affinity for TfR1 (Kd: 1.15 nM), followed by

BSG (Kd: 2.16 nM) and prohibitin-2 (Kd: 2.77 nM) (Fig 6B).

Fig 6. PvRBP1a$_{158\text{-}650}$ binds to TfR1, BSG, and PHB2 at nanomolar affinities. (A)

Titration ELISAs of protein-protein interactions between soluble ligand PvRBP1a$_{158\text{-}}$

$_{650}$LTID, control LTID, and the immobilized receptor candidates TfR1, BSG, and PHB2.

OD$_{450}$ absorbance values represent the binding of a TurboID-specific HRP-labeled

antibody for the biotin-free quantification of ligand and control in triplicate, fitted by a

single-site binding model; (B) average dissociation constants (kd) summarized as

determined from the fitted titration curves above.

Discussion

This study demonstrates the utility of erythroid cell lines JK-1 and BEL-A as suitable

surrogates for reticulocytes for studying the invasion process of the *P. vivax* malaria

parasite. While these cell lines and their culture conditions did not support a continuous

P. vivax culture *in vitro*, the formation of the Hz pigment and immuno detection of

PvLDH strongly supported parasite invasion. In contrast to human reticulocytes, both

erythroid cell lines are nucleated, which might permit them to initiate a cell-death

program upon parasite invasion. Consistently, JK-1 cells were previously reported to

support cell entry by both *P. vivax* and *P. falciparum* [24,25], while BEL-A cells have so

far only been studied with *P. falciparum* [26]. This study is the first to report *P. vivax*

invasion of BEL-A cells, confirming their susceptibility alongside JK-1 cells.

The quantitative membrane proteome comparison of reticulocytes and erythroid cell

lines with erythrocytes revealed Prohibitin-2 (PHB2), TfR1 (CD71), the CD98 heavy

chain (4F2hc, gene SLC3A2), the CD98 light chain (LAT-1, gene SLC7A5), ATB(0)

(gene SLC1A5), CAT-1 (SLC7A1), CD36, Integrin β-1 (gene ITGB1), and Metal

transporter CNNM3 to be of significantly increased abundance in those cell lines and

reticulocytes. Whereas they are strongly decreased (practically absent) in fully matured

erythrocytes. The increased abundance of TfR1 and CD98 in reticulocytes over

erythrocytes is consistent with previous studies [2,20]. However, genetic manipulation

[3] or antibody blockade [2] of these proteins only partially reduced *P. vivax* invasion,

suggesting the involvement of additional receptors. The heavy chain of CD98 (SLC3A2)

was reported to be bound by *P. vivax* in immature red blood cells via PvRBP2a [2].

Therefore, other potential *P. vivax* receptor candidates with extracellular regions,

namely Prohibitin-2, the CD98 light chain (LAT-1), ATB(0), CAT-1, CD36, Integrin β-1,

and CNNM3 should be considered. These membrane proteins participate in various

protein-protein interactions that facilitate the entry of microorganisms into host cells [27–

39]. Interestingly, LAT-1, that together with its heavy chain 4F2hc forms the heteromeric

CD98 [40,41], plays a role in hepatitis C virus entry [42], raising the question of whether

P. vivax may also interact with LAT-1.

Proximity labeling of erythroid cells with PvRBP1a$_{158-650}$LTID was largely consistent with

previous observations, in which ~50% of reticulocytes binding and ~20% of erythrocytes

bound to PvRBP1a$_{157-650}$ [13]. Furthermore, 20 of reticulocytes and only 1% of

erythrocytes bound to PvRBP1a$_{351-599}$ [43], while 31.5% of reticulocytes were reported to

bind to PvRBP1a$_{30-778}$ [44]. Additionally, the trypsin and chymotrypsin sensitivity of these

recombinant protein [13,43,44], as well as the PvRBP1a$_{157-653}$ HAPBs [12] align with our

results.

PvRBP1a$_{158-650}$ was found to interact with Prohibitin-2, TfR1, and BSG. Prohibitin-2 and

TfR1 are more abundant in reticulocyte membranes and cell lines compared to

erythrocytes, while BSG is more abundant in erythrocytes. These findings suggest that

PvRBP1a$_{158-650}$ may facilitate reticulocyte recognition and invasion through interaction

with Prohibitin-2 and TfR1. Additionally, interaction with BSG may contribute to binding

activity to erythrocytes, but not their restricted invasion, consistent with previous studies

on PvRBP1a binding [12,13,43,44].

This study demonstrated strong binding of PvRBP1a$_{158-650}$ with the 89-760 domain of

TfR1, contrasting with prior work that did not detect this interaction, possibly due to the

crucial role of TfR1's 89-120 region, not included in the previous protein construct [25].

TfR1 is a known receptor for PvRBP2b [3], as well as for various New World

arenaviruses [45,46]. BSG, a known receptor for *P. falciparum* RH5 [47,48], a member

of the PfRH family homologous to the PvRBP proteins of *P. vivax* [49,50], also serves

as a receptor for *P. vivax* TRAg38 [51]. Similarly, prohibitin-2, facilitates viral entry into

host cells, including enteroviruses, coronaviruses, HIV-1, and flaviviruses such as

dengue [39,52–54].

This research demonstrated the binding versatile of $PvRBP1a_{158-650}$, which interacts

with three different membrane proteins. In biological systems, ligands often bind

multiple receptors, as seen with *Plasmodium* interactions; PvTRAg38 binds both BSG

and band 3 [51,55], and PfEMP1binds to several receptors [56–60].

This study demonstrates that the BEL-A and JK-1 cells are suitable models for studying

P. vivax receptor-ligand interactions, providing viable alternatives to reticulocytes. The

similarity in the abundance of potential receptor candidates between cell lines and

reticulocytes, and their dissimilarity with erythrocytes, validates the use of JK-1 and

BEL-A cell lines as surrogate models for the study of *P. vivax* merozoite ligand-receptor

interactions, and suggests the existence of other potential *P. vivax* receptors. prohibitin-

2 and TfR1 may contribute to a redundant reticulocyte-restricted invasion pathway

because they exhibit high binding affinities to $PvRBP1a_{158-650}$. These findings lay the

foundation for the comprehensive study of all *P. vivax* invasion mechanisms and for the

development of targeted therapies against malaria.

363   Materials and methods

364   Ethics statement

The study was conducted in accordance with the Declaration of Helsinki. Use of

anonymized discarded blood from therapeutic phlebotomy was approved by the

Institutional Review Board of City of Hope, Duarte, California, USA, as exempt category

4, under 45CFR46.104 (d). Blood samples from malaria patients were obtained under

informed consent with the approval of the Bioethics central committee of the

Universidad de Córdoba, Monteria, Colombia, and imported into the United States

under CDC permit No.: 20210830-3188A0.

Collection, processing, and enrichment of *P. vivax* parasites from blood samples

P. vivax infected blood samples were collected from malaria patients in Tierralta—

Córdoba, Colombia, into 5-mL sodium citrate tubes. After transportation to Bogotá,

RBCs were enriched by centrifugation, mixed with an equal volume of Glycerolyte 57,

cryopreserved, shipped to the U.S. lab, thawed using the NaCl method [61], and

resuspended in 3 mL of Iscove's Modified Dulbecco's Medium (IMDM). These RBCs

were then enriched from 0.2% to 4.0% parasitemia by concentrating *P. vivax*-infected

reticulocytes through a KCl-Percoll gradient [62]. Enrichment was evaluated by

microscopy with Giemsa staining (Fig 1A).

Identification of the *Plasmodium* species

383   Genomic DNA was extracted from infected cells, and nested PCR was performed using

384   this DNA to identify the *Plasmodium* species. Genus- and species-specific primers

385   targeting the parasite's 18S ribosomal small subunit RNA were used as previously

386   described [63] (see S1 Text).

P. vivax entry into BEL-A and JK-1 cells

The enriched *P. vivax* mono-infected RBC sample (~40 μL) was divided into two equal

aliquots, one for co-incubation with BEL-A cells and the other with JK-1 cells. Each cell

line (1.5×10^5 cells) was cultured in 500 μL of medium, for BEL-A in StemSpan serum-

free expansion medium (SFEM, StemCell Technologies), containing 25% human serum

(Type AB, Sigma-Aldrich), 50 ng/mL stem cell factor (SCF), 3 U/mL erythropoietin

(EPO), 1 µM dexamethasone, 1 µg/mL doxycycline, 1 µg/mL chemically defined lipid

concentrate (CDLC), and 100 µM hypoxanthine; and JK-1 in IMDM supplemented with

GlutaMAX, with the same components, except for SCF, EPO, dexamethasone, and

doxycycline. Both cultures were incubated at 37°C with 5% CO_2 and 5% O_2. Fresh

medium and 2×10^5 erythroid cells were added every two days, and cultures were

evaluated by immunofluorescence assay (IFA) every 24 hours for 6 days (see S1 Text).

Quantitative comparison of membrane proteomes by (DIA) - LC-MS/MS

Reticulocytes, erythrocytes, JK-1 and BEL-A cells were collected, cytoplasmic content

was removed by osmolytic lysis, and membrane proteins of the remaining ghosts were

extracted. From each sample, 23 µg of proteins were processed for proteomics using S-

Trap columns (ProtiFi) according to the manufacturer instructions [64,65]. The resulting

trypsin/LysC-digested peptides were analyzed by LC-MS/MS in data-independent

acquisition (DIA) mode, as detailed in the S1 Text.

To enrich the plasma membrane proteins from the data set, proteins were filtered based

on at least one of the following annotations from the UniProt subcellular localization

database: "Cell membrane", "Apical cell membrane", "Basolateral cell membrane",

"Peripheral membrane protein" and "Plasma membrane", and the Gene Ontology (GO)

annotation term: "plasma membrane". The topology of the membrane proteins abundant

in reticulocytes compared to erythrocytes was evaluated using several predictors.

Protein sequences were analyzed with Protter for overall visualization of proteoforms

[66], TMHMM 2.0 for transmembrane region prediction [67], SignalP 6.0 for signal

peptide identification [68], and PredGPI for GPI anchor site prediction [69].

PvRBP1A$_{158-650}$ proximity labeling for the identification of

likely receptor candidates

Cloning, expression, purification, and activity of TurboID fusion

proteins

The DNA sequence encoding PvRBP1a$_{158-650}$ (GenBank AAS85749.1) was derived

from the *P. vivax* Salvador I reference strain (txid126793). This sequence was fused to

an acidic linker (L), GDEVDEDEG, to improve solubility, and the TurboID protein (TID)

[22], followed by a C-terminal 6xHis tag for purification, resulting in the PvRBP1a$_{158-650}$LTID fusion protein. An equivalent gene encoding L with TurboID alone (LTID) was

designed as a negative control. Both gene constructs were obtained as customized

synthetic genes, optimized for expression in *E. coli,* and cloned into a pET-28a(+)

expression vector between its *Nco*I and *Xho*I sites. The constructs were expressed in

soluble form in *E. coli* BL21 cells and purified by affinity chromatography, as detailed in

the S1 Text. Protein purity and expression were verified by polyacrylamide gel

electrophoresis. The biotinylation activity of both recombinant TurboID fusion proteins

was confirmed by evaluating their autobiotinylation activity through incubation in the

presence or absence of biotinylation reaction buffer Brxn (20 mM Tris-HCl, 500 µM

biotin, 2.5 mM ATP, pH 7.5) at 37°C for 15 minutes, quenching on ice and Western blot

analysis with streptavidin-IRDye 800CW conjugate (1:1,000) and an Odyssey DLx

imaging system (LICORbio).

PvRBP1a$_{158-650}$LTID proximity labeling

To evaluate PvRBP1a$_{158-650}$LTID's interaction with JK-1, BEL-A, reticulocytes, and erythrocytes, proximity labeling assays were performed in duplicate, and repeated up to three times. Cells were washed twice with PBS supplemented with 2% human serum (HS 2%) and incubated with either PvRBP1a$_{158-650}$LTID or LTID (negative control) for 3 hours at room temperature with constant mild agitation at 10 rpm. Following incubation, cells were washed three times with HS 2% to remove unbound proteins, then incubated with Brxn for 15 minutes at 37°C. The reaction was stopped by cooling on ice for 5 minutes, and the samples were washed with cold HS 2% before labeling with Alexa Fluor 488-conjugated streptavidin (10 µg/mL, Invitrogen) for 1 hour at room temperature. Biotinylation was quantified by cytometry, acquiring 100,000 events per sample on a FACSAria Fusion (BD). Data were analyzed with FlowJo v10.8.1 [70], calculating the percentage of biotinylated cells relative to total cells. LTID-treated and unlabeled cells served as negative controls.

The biochemical nature of PvRBP1a$_{158-650}$ receptors

JK-1 and BEL-A cells were treated with trypsin (1 mg/mL, Sigma-Aldrich), chymotrypsin (1 mg/mL, Sigma-Aldrich), or neuraminidase (50 mU, Roche) for 1 hour. After enzymatic treatment, proteolytic enzymes were inactivated with soy trypsin inhibitor (0.5 mg/mL, Sigma-Gibco) [71]. Proximity labeling assays were then performed as described above, using PvRBP1a$_{158-650}$LTID or LTID (3 µM).

Affinity enrichment and LC-MS identification of PvRBP1a$_{158-650}$ proximity-labeled receptor candidates

JK-1 cells were incubated with either PvRBP1a$_{158-650}$LTID or LTID (negative control),

each at 3 μM for 3 hours. After incubation, cells were washed three times with HS 2%,

then incubated with 100 μL of Brxn for 15 minutes at 37°C. The reaction was stopped as

described above, and cells were resuspended in 500 μL of IP-MS lysis buffer (MS-

compatible Magnetic IP kit, streptavidin, Pierce, Thermo Scientific), incubated on ice for

30 minutes with intermittent vortexing every 5 minutes. After centrifugation, the lysate's

supernatant was collected and combined with Streptavidin magnetic beads (50 μL,

Thermo Scientific), incubated for 1 hour at 21°C, and then overnight at 4°C, to enrich

biotinylated membrane proteins. The beads were washed, and biotinylated proteins

were eluted sequentially with 100 μL of 50 mM biotin, 100 μL of elution buffer, and 100

μL of 5% SDS at 95°C.

The eluted proteins were reduced, alkylated, and processed for proteomics using S-

Trap spin columns (ProtiFi) according to the manufacturer's instructions [64,65]. The

resulting trypsin/LysC digested peptides were analyzed by LC-MS in data-dependent

acquisition mode. Data analysis was performed using FragPipe v22.0 [72]. Candidate

receptor proteins for PvRBP1a$_{158-650}$ were selected based on the presence of

extracellular regions that are favorable for ligand interaction, evaluated using UniProt

GO annotations and the TMHMM 2.0 predictor [67]. Receptor candidates in the

PvRBP1a$_{158-650}$LTID sample that were detected in both duplicates and of significantly

higher abundance (≥ 2 fold) compared to the negative control (LTID) were also

considered. Subsequently, a parallel reaction monitoring (PRM) method was applied to

validate and quantify the peptides of interest (as detailed in the S1 Text).

Binding affinities of *Pv*RBP1a$_{158-650}$ to select receptor candidates by ELISA

The recombinant extracellular protein domains of receptor candidates TfR1 (Cys^{89}-Phe^{760}, SinoBiological), BSG (Met^{1}-His^{205}, SinoBiological), and full-length PHB2 (Origene) were used to evaluate the interaction between the PvRBP1a$_{158-650}$ and its binding membrane proteins (S1 Fig). Maxisorp plates were coated in triplicates with 5 µg/mL of each protein for 2 hours at room temperature and blocked with SuperBlock Buffer (Thermo Scientific). Serial dilutions of PvRBP1a$_{158-650}$LTID and LTID were prepared in blocking solution, ranging from 48,000 pM to 187.5 pM (1:2 dilution) and 4,000 pM to 1.28 pM (1:5 dilution), and incubated for 16 hours at 4°C. Bound PvRBP1a$_{158-650}$LTID and LTID were detected with a TurboID-specific polyclonal rabbit antibody (anti-BirA mutated/TurboID, Agrisera, 1:10,000). After five washes with PBST, 100 µL of 3,3',5,5',-Tetramethylbenzidine (TMB) substrate was added, and the reaction was stopped with 50 µL of 1 M phosphoric acid. Absorbance was measured at 450 nm. Dissociation constants (Kd) were determined using GraphPad Prism v10.3.1 with non-linear regression and a one-site binding saturation model.

Acknowledgments

The authors are grateful to NHS Blood and Transplant, especially Professor Jan Frayne, for providing the BEL-A cell line. *"BEL-A cell lines were created by Professor Jan Frayne, Professor David Anstee and Dr Kongtana Trakarnsanga with funding from the Wellcome Trust (grant numbers 087430/Z/08 and 102610), NHS Blood and Transplant and Department of Health (England)"*. Furthermore, they wish to extend their gratitude to the

References

1. World Health Organization. World malaria report 2024: addressing inequity in the global malaria response. 2024.

2. Malleret B, El Sahili A, Tay MZ, Carissimo G, Ong ASM, Novera W, et al. *Plasmodium vivax* binds host CD98hc (SLC3A2) to enter immature red blood cells. Nat Microbiol. 2021;6: 991–999. doi:10.1038/s41564-021-00939-3

3. Gruszczyk J, Kanjee U, Chan L-J, Menant S, Malleret B, Lim NTY, et al. Transferrin receptor 1 is a reticulocyte-specific receptor for *Plasmodium vivax*. Science. 2018;359: 48–55. doi:10.1126/science.aan1078

4. Molina-Franky J, Patarroyo ME, Kalkum M, Patarroyo MA. The Cellular and Molecular Interaction Between Erythrocytes and *Plasmodium falciparum* Merozoites. Front Cell Infect Microbiol. 2022;12: 816574. doi:10.3389/fcimb.2022.816574

5. Malleret B, Li A, Zhang R, Tan KSW, Suwanarusk R, Claser C, et al. Plasmodium vivax: restricted tropism and rapid remodeling of CD71-positive reticulocytes. Blood. 2015;125: 1314–1324. doi:10.1182/blood-2014-08-596015

6. Trager W, Jensen JB. Human Malaria Parasites in Continuous Culture. Science. 1976;193: 673–675. doi:10.1126/science.781840

7. Horuk R, Chitnis CE, Darbonne WC, Colby TJ, Rybicki A, Hadley TJ, et al. A Receptor for the Malarial Parasite *Plasmodium vivax*: the Erythrocyte Chemokine Receptor. Science. 1993;261: 1182–1184. doi:10.1126/science.7689250

8. Ryan JR, Stoute JA, Amon J, Dunton RF, Mtalib R, Koros J, et al. Evidence for transmission of *Plasmodium vivax* among a duffy antigen negative population in Western Kenya. Am J Trop Med Hyg. 2006;75: 575–581.

9.   Reyes C, Picón Jaimes YA, Kalkum M, Patarroyo MA. The Black Box of Cellular and
Molecular Events of *Plasmodium vivax* Merozoite Invasion into Reticulocytes.
IJMS. 2022;23: 14528. doi:10.3390/ijms232314528

10. Picón-Jaimes YA, Lozada-Martinez ID, Orozco-Chinome JE, Molina-Franky J,
Acevedo-Lopez D, Acevedo-Lopez N, et al. Relationship between Duffy
Genotype/Phenotype and Prevalence of *Plasmodium vivax* Infection: A Systematic
Review. TropicalMed. 2023;8: 463. doi:10.3390/tropicalmed8100463

11. Galinski MR, Medina CC, Ingravallo P, Barnwell JW. A reticulocyte-binding protein
complex of *Plasmodium vivax* merozoites. Cell. 1992;69: 1213–1226.
doi:10.1016/0092-8674(92)90642-P

12. Urquiza M, Patarroyo M, Marí V, Ocampo M, Suarez J, Lopez R, et al. Identification
and polymorphism of *Plasmodium vivax* RBP-1 peptides which bind specifically to
reticulocytes. Peptides. 2002;23: 2265–2277. doi:10.1016/S0196-9781(02)00267-X

13. Ntumngia FB, Thomson-Luque R, Galusic S, Frato G, Frischmann S, Peabody DS,
et al. Identification and Immunological Characterization of the Ligand Domain of
Plasmodium vivax Reticulocyte Binding Protein 1a. The Journal of Infectious
Diseases. 2018;218: 1110–1118. doi:10.1093/infdis/jiy273

14. Okuno Y, Suzuki A, Ichiba S, Takahashi T, Nakamura K, Hitomi K, et al.
Establishment of an Erythroid Cell Line (JK-1) that Spontaneously Diferentiates to
Red Cells. 66: 8.

15. Trakarnsanga K, Griffiths RE, Wilson MC, Blair A, Satchwell TJ, Meinders M, et al.
An immortalized adult human erythroid line facilitates sustainable and scalable
generation of functional red cells. Nat Commun. 2017;8: 14750.
doi:10.1038/ncomms14750

16. Cao P, Kho S, Grigg MJ, Barber BE, Piera KA, William T, et al. Characterisation of
Plasmodium vivax lactate dehydrogenase dynamics in *Plasmodium vivax*
infections. Commun Biol. 2024;7: 355. doi:10.1038/s42003-024-05956-6

17. Pandey AV, Tekwani BL. Formation of haemozoin/β-haematin under physiological
conditions is not spontaneous. FEBS Letters. 1996;393: 189–192.
doi:10.1016/0014-5793(96)00881-2

18. Coronado LM, Nadovich CT, Spadafora C. Malarial hemozoin: From target to tool.
Biochimica et Biophysica Acta (BBA) - General Subjects. 2014;1840: 2032–2041.
doi:10.1016/j.bbagen.2014.02.009

19. Carlton JM, Adams JH, Silva JC, Bidwell SL, Lorenzi H, Caler E, et al. Comparative
genomics of the neglected human malaria parasite *Plasmodium vivax*. Nature.
2008;455: 757–763. doi:10.1038/nature07327

20. Malleret B, Li A, Zhang R, Tan KSW, Suwanarusk R, Claser C, et al. *Plasmodium*
vivax: restricted tropism and rapid remodeling of CD71-positive reticulocytes.
Blood. 2015;125: 1314–1324. doi:10.1182/blood-2014-08-596015

21. Lex A, Gehlenborg N, Strobelt H, Vuillemot R, Pfister H. UpSet: Visualization of
Intersecting Sets. IEEE Trans Visual Comput Graphics. 2014;20: 1983–1992.
doi:10.1109/TVCG.2014.2346248

22. Branon TC, Bosch JA, Sanchez AD, Udeshi ND, Svinkina T, Carr SA, et al.
Efficient proximity labeling in living cells and organisms with TurboID. Nat
Biotechnol. 2018;36: 880–887. doi:10.1038/nbt.4201

23. Cho KF, Branon TC, Udeshi ND, Myers SA, Carr SA, Ting AY. Proximity labeling in
mammalian cells with TurboID and split-TurboID. Nat Protoc. 2020;15: 3971–3999.
doi:10.1038/s41596-020-0399-0

24. Kanjee U, Grüring C, Chaand M, Lin K-M, Egan E, Manzo J, et al. CRISPR/Cas9
knockouts reveal genetic interaction between strain-transcendent erythrocyte
determinants of *Plasmodium falciparum* invasion. Proc Natl Acad Sci U S A.
2017;114: E9356–E9365. doi:10.1073/pnas.1711310114

25. Gruszczyk J, Kanjee U, Chan L-J, Menant S, Malleret B, Lim NTY, et al. Transferrin
receptor 1 is a reticulocyte-specific receptor for *Plasmodium vivax*. Science.
2018;359: 48–55. doi:10.1126/science.aan1078

26. Satchwell TJ, Wright KE, Haydn-Smith KL, Sánchez-Román Terán F, Moura PL,
Hawksworth J, et al. Genetic manipulation of cell line derived reticulocytes enables
dissection of host malaria invasion requirements. Nat Commun. 2019;10: 3806.
doi:10.1038/s41467-019-11790-w

27. Smith JosephD, Kyes S, Craig AlisterG, Fagan T, Hudson-Taylor D, Miller LouisH,
et al. Analysis of adhesive domains from the A4VAR Plasmodium falciparum
erythrocyte membrane protein-1 identifies a CD36 binding domain. Molecular and
Biochemical Parasitology. 1998;97: 133–148. doi:10.1016/S0166-6851(98)00145-5

28. Albritton LM, Kim JW, Tseng L, Cunningham JM. Envelope-binding domain in the
cationic amino acid transporter determines the host range of ecotropic murine
retroviruses. J Virol. 1993;67: 2091–2096. doi:10.1128/jvi.67.4.2091-2096.1993

29. Yoshimoto T, Yoshimoto E, Meruelo D. Identification of amino acid residues critical
for infection with ecotropic murine leukemia retrovirus. J Virol. 1993;67: 1310–
1314. doi:10.1128/jvi.67.3.1310-1314.1993

30. Olaya-Galán NN, Salas-Cárdenas SP, Rodriguez-Sarmiento JL, Ibáñez-Pinilla M,
Monroy R, Corredor-Figueroa AP, et al. Risk factor for breast cancer development
under exposure to bovine leukemia virus in Colombian women: A case-control
study. Jaworski JP, editor. PLoS ONE. 2021;16: e0257492.
doi:10.1371/journal.pone.0257492

31. Tailor CS, Nouri A, Zhao Y, Takeuchi Y, Kabat D. A sodium-dependent neutral-
amino-acid transporter mediates infections of feline and baboon endogenous
retroviruses and simian type D retroviruses. J Virol. 1999;73: 4470–4474.
doi:10.1128/JVI.73.5.4470-4474.1999

32. Feire AL, Roy RM, Manley K, Compton T. The Glycoprotein B Disintegrin-Like
Domain Binds Beta 1 Integrin To Mediate Cytomegalovirus Entry. J Virol. 2010;84:
10026–10037. doi:10.1128/JVI.00710-10

33. Xiao J, Palefsky JM, Herrera R, Berline J, Tugizov SM. The Epstein–Barr virus
BMRF-2 protein facilitates virus attachment to oral epithelial cells. Virology.
2008;370: 430–442. doi:10.1016/j.virol.2007.09.012

34. Weigel-Kelley KA, Yoder MC, Srivastava A. α5β1 integrin as a cellular coreceptor
for human parvovirus B19: requirement of functional activation of β1 integrin for
viral entry. Blood. 2003;102: 3927–3933. doi:10.1182/blood-2003-05-1522

35. Graham KL, Halasz P, Tan Y, Hewish MJ, Takada Y, Mackow ER, et al. Integrin-
Using Rotaviruses Bind α2β1 Integrin α2 I Domain via VP4 DGE Sequence and
Recognize αXβ2 and αVβ3 by Using VP7 during Cell Entry. J Virol. 2003;77: 9969–
9978. doi:10.1128/JVI.77.18.9969-9978.2003

36. Maginnis MS, Forrest JC, Kopecky-Bromberg SA, Dickeson SK, Santoro SA, Zutter
MM, et al. Beta1 integrin mediates internalization of mammalian reovirus. J Virol.
2006;80: 2760–2770. doi:10.1128/JVI.80.6.2760-2770.2006

37. Nägele V, Heesemann J, Schielke S, Jiménez-Soto LF, Kurzai O, Ackermann N.
Neisseria meningitidis adhesin NadA targets beta1 integrins: functional similarity to
Yersinia invasin. J Biol Chem. 2011;286: 20536–20546.
doi:10.1074/jbc.M110.188326

38. Wintachai P, Wikan N, Kuadkitkan A, Jaimipuk T, Ubol S, Pulmanausahakul R, et
al. Identification of prohibitin as a Chikungunya virus receptor protein. J Med Virol.
2012;84: 1757–1770. doi:10.1002/jmv.23403

39. Su W, Huang S, Zhu H, Zhang B, Wu X. Interaction between PHB2 and
Enterovirus A71 VP1 Induces Autophagy and Affects EV-A71 Infection. Viruses.
2020;12: 414. doi:10.3390/v12040414

40. Lee Y, Wiriyasermkul P, Jin C, Quan L, Ohgaki R, Okuda S, et al. Cryo-EM
structure of the human L-type amino acid transporter 1 in complex with
glycoprotein CD98hc. Nat Struct Mol Biol. 2019;26: 510–517. doi:10.1038/s41594-
019-0237-7

41. Yan R, Zhao X, Lei J, Zhou Q. Structure of the human LAT1–4F2hc heteromeric
amino acid transporter complex. Nature. 2019;568: 127–130. doi:10.1038/s41586-
019-1011-z

42. Nguyen NNT, Lim Y-S, Nguyen LP, Tran SC, Luong TTD, Nguyen TTT, et al.
Hepatitis C Virus Modulates Solute carrier family 3 member 2 for Viral Propagation.
Sci Rep. 2018;8: 15486. doi:10.1038/s41598-018-33861-6

43. Han J-H, Lee S-K, Wang B, Muh F, Nyunt MH, Na S, et al. Identification of a
reticulocyte-specific binding domain of *Plasmodium vivax* reticulocyte-binding
protein 1 that is homologous to the PfRh4 erythrocyte-binding domain. Sci Rep.
2016;6: 26993. doi:10.1038/srep26993

44. Gupta ED, Anand G, Singh H, Chaddha K, Bharti PK, Singh N, et al. Naturally
Acquired Human Antibodies Against Reticulocyte-Binding Domains of *Plasmodium*
vivax Proteins, PvRBP2c and PvRBP1a, Exhibit Binding-Inhibitory Activity. J Infect
Dis. 2017;215: 1558–1568. doi:10.1093/infdis/jix170

45. Radoshitzky SR, Kuhn JH, Spiropoulou CF, Albariño CG, Nguyen DP, Salazar-
Bravo J, et al. Receptor determinants of zoonotic transmission of New World
hemorrhagic fever arenaviruses. Proc Natl Acad Sci USA. 2008;105: 2664–2669.
doi:10.1073/pnas.0709254105

46. Radoshitzky SR, Abraham J, Spiropoulou CF, Kuhn JH, Nguyen D, Li W, et al.
Transferrin receptor 1 is a cellular receptor for New World haemorrhagic fever
arenaviruses. Nature. 2007;446: 92–96. doi:10.1038/nature05539

47. Crosnier C, Bustamante LY, Bartholdson SJ, Bei AK, Theron M, Uchikawa M, et al.
Basigin is a receptor essential for erythrocyte invasion by *Plasmodium falciparum*.
Nature. 2011;480: 534–537. doi:10.1038/nature10606

48. Chen L, Xu Y, Healer J, Thompson JK, Smith BJ, Lawrence MC, et al. Crystal
structure of PfRh5, an essential *Plasmodium falciparum* ligand for invasion of
human erythrocytes. eLife. 2014;3: e04187. doi:10.7554/eLife.04187

49. Rayner JC, Galinski MR, Ingravallo P, Barnwell JW. Two Plasmodium falciparum
genes express merozoite proteins that are related to Plasmodium vivax and
Plasmodium yoelii adhesive proteins involved in host cell selection and invasion.
Proceedings of the National Academy of Sciences. 2000;97: 9648–9653.
doi:10.1073/pnas.160469097

50. Triglia T, Thompson J, Caruana SR, Delorenzi M, Speed T, Cowman AF.
Identification of Proteins from Plasmodium falciparum That Are Homologous to
Reticulocyte Binding Proteins in Plasmodium vivax. Petri WA, editor. Infect Immun.
2001;69: 1084–1092. doi:10.1128/IAI.69.2.1084-1092.2001

51. Rathore S, Dass S, Kandari D, Kaur I, Gupta M, Sharma YD. Basigin Interacts with
*Plasmodium vivax*Tryptophan-rich Antigen PvTRAg38 as a Second Erythrocyte
Receptor to Promote Parasite Growth. J Biol Chem. 2017;292: 462–476.
doi:10.1074/jbc.M116.744367

52. Emerson V, Holtkotte D, Pfeiffer T, Wang I-H, Schnölzer M, Kempf T, et al.
Identification of the Cellular Prohibitin 1/Prohibitin 2 Heterodimer as an Interaction
Partner of the C-Terminal Cytoplasmic Domain of the HIV-1 Glycoprotein. J Virol.
2010;84: 1355–1365. doi:10.1128/JVI.01641-09

53. Cornillez-Ty CT, Liao L, Yates JR, Kuhn P, Buchmeier MJ. Severe acute
respiratory syndrome coronavirus nonstructural protein 2 interacts with a host
protein complex involved in mitochondrial biogenesis and intracellular signaling. J
Virol. 2009;83: 10314–10318. doi:10.1128/JVI.00842-09

54. Kuadkitkan A, Wikan N, Fongsaran C, Smith DR. Identification and characterization
of prohibitin as a receptor protein mediating DENV-2 entry into insect cells.
Virology. 2010;406: 149–161. doi:10.1016/j.virol.2010.07.015

55. Alam MS, Rathore S, Tyagi RK, Sharma YD. Host-parasite interaction: multiple
sites in the *Plasmodium vivax* tryptophan-rich antigen PvTRAg38 interact with the
erythrocyte receptor band 3. Griesinger C, editor. FEBS Lett. 2016;590: 232–241.
doi:10.1002/1873-3468.12053

56. Yipp BG, Anand S, Schollaardt T, Patel KD, Looareesuwan S, Ho M. Synergism of
multiple adhesion molecules in mediating cytoadherence of *Plasmodium
falciparum*–infected erythrocytes to microvascular endothelial cells under flow.
Blood. 2000;96: 2292–2298. doi:10.1182/blood.V96.6.2292

57. Vogt AM, Barragan A, Chen Q, Kironde F, Spillmann D, Wahlgren M. Heparan
sulfate on endothelial cells mediates the binding of *Plasmodium falciparum*–
infected erythrocytes via the DBL1α domain of PfEMP1. Blood. 2003;101: 2405–
2411. doi:10.1182/blood-2002-07-2016

58. Vigan-Womas I, Guillotte M, Juillerat A, Hessel A, Raynal B, England P, et al.
Structural Basis for the ABO Blood-Group Dependence of *Plasmodium falciparum*
Rosetting. Rowe JA, editor. PLoS Pathog. 2012;8: e1002781.
doi:10.1371/journal.ppat.1002781

59. Kessler A, Dankwa S, Bernabeu M, Harawa V, Danziger SA, Duffy F, et al. Linking
EPCR-Binding PfEMP1 to Brain Swelling in Pediatric Cerebral Malaria. Cell Host &
Microbe. 2017;22: 601-614.e5. doi:10.1016/j.chom.2017.09.009

60. Esser C, Bachmann A, Kuhn D, Schuldt K, Förster B, Thiel M, et al. Evidence of
promiscuous endothelial binding by *Plasmodium falciparum*-infected erythrocytes.
Cell Microbiol. 2014;16: 701–708. doi:10.1111/cmi.12270

61. Blomqvist, K. Thawing of glycerolyte-frozen parasites with NaCl. Methods in
Malaria Research. MR4/ATCC, Manassas. Methods in Malaria Research. Methods
in Malaria Research. MR4/ATCC, Manassas. 2008. p. 15.

62. Rangel GW, Clark MA, Kanjee U, Lim C, Shaw-Saliba K, Menezes MJ, et al.
Enhanced ex vivo *Plasmodium vivax* Intraerythrocytic Enrichment and Maturation

for Rapid and Sensitive Parasite Growth Assays. Antimicrob Agents Chemother.
2018;62: e02519-17. doi:10.1128/AAC.02519-17

63. Snounou G, Viriyakosol S, Xin Ping Zhu, Jarra W, Pinheiro L, Do Rosario VE, et al.
High sensitivity of detection of human malaria parasites by the use of nested
polymerase chain reaction. Molecular and Biochemical Parasitology. 1993;61:
315–320. doi:10.1016/0166-6851(93)90077-B

64. HaileMariam M, Eguez RV, Singh H, Bekele S, Ameni G, Pieper R, et al. S-Trap,
an Ultrafast Sample-Preparation Approach for Shotgun Proteomics. J Proteome
Res. 2018;17: 2917–2924. doi:10.1021/acs.jproteome.8b00505

65. Matulis D. Selective Precipitation of Proteins. CP Protein Science. 2016;83.
doi:10.1002/0471140864.ps0405s83

66. Omasits U, Ahrens CH, Müller S, Wollscheid B. Protter: interactive protein feature
visualization and integration with experimental proteomic data. Bioinformatics.
2014;30: 884–886. doi:10.1093/bioinformatics/btt607

67. Hallgren J, Tsirigos KD, Pedersen MD, Almagro Armenteros JJ, Marcatili P,
Nielsen H, et al. DeepTMHMM predicts alpha and beta transmembrane proteins
using deep neural networks. Bioinformatics; 2022 Apr.
doi:10.1101/2022.04.08.487609

68. Teufel F, Almagro Armenteros JJ, Johansen AR, Gíslason MH, Pihl SI, Tsirigos
KD, et al. SignalP 6.0 predicts all five types of signal peptides using protein
language models. Nat Biotechnol. 2022;40: 1023–1025. doi:10.1038/s41587-021-
01156-3

69. Pierleoni A, Martelli PL, Casadio R. PredGPI: a GPI-anchor predictor. BMC
Bioinformatics. 2008;9: 392. doi:10.1186/1471-2105-9-392

70. Ashland, OR: Becton, Dickinson and Company. FlowJo™ Software (for Windows)
Version 10.8.1. 2023.

71. Deans A-M, Nery S, Conway DJ, Kai O, Marsh K, Rowe JA. Invasion Pathways
and Malaria Severity in Kenyan *Plasmodium falciparum* Clinical Isolates. Infect
Immun. 2007;75: 3014–3020. doi:10.1128/IAI.00249-07

72. Yu F, Haynes SE, Nesvizhskii AI. IonQuant Enables Accurate and Sensitive Label-
Free Quantification With FDR-Controlled Match-Between-Runs. Molecular &
Cellular Proteomics. 2021;20: 100077. doi:10.1016/j.mcpro.2021.100077

Supporting information captions

S1 Text. Supporting methods.

S1 Table. *P. vivax* merozoite candidate receptors

S1 Fig. Purity Analysis of Recombinant Proteins.

Recombinant proteins BSG, TfR1, and PHB2 were separated on a 4–12%

polyacrylamide gel and stained with Coomassie blue. The observed bands correspond

to the expected molecular weights for each protein: BSG (35 kDa), TfR1 (~77.4 kDa),

and PHB2 (~35.7 kDa), confirming their purity. The Kaleidoscope molecular weight

marker (Bio-Rad) is included in the left lane as a reference.

S2 Fig. Reticulocyte enrichment.

(A, B) Reticulocyte percentage in the initial whole blood sample. (C, D) Reticulocyte

percentage after Percoll density (70%-62%) enrichment. (E) Purity of reticulocytes

(CD71$^+$ CD45$^-$) after sorting collection. This high-purity reticulocyte sample was not

evaluated with brilliant cresyl blue due to the small total volume of sample obtained (~

50 µL). (B, D) Blood smears stained with brilliant cresyl blue to identify reticulocytes (red

arrows). Scale bars are 10 µm.

S3 Fig. Differentiation of erythroid cells.

(A-D) BEL-A cells. (E-H) JK-1 cells. (A, E) Proerythroblasts (20-25 µm) were identified

by their large nucleus with fine chromatin, prominent nucleoli, and basophilic cytoplasm.

These cells progressed to (B, F) Basophilic erythroblasts (16-18 µm), characterized by a

slightly smaller nucleus with condensed chromatin and cytoplasm with pronounced

basophilia due to increased ribosomes synthesizing Hb. Subsequently, cells

differentiated into (C, G) polychromatophilic erythroblasts (12-15 µm), which exhibited a

more condensed nucleus with clusters of heterochromatin. The final nucleated stage,

(D, H) orthochromatic erythroblast, indicated by the red arrow in D (10-12 µm), was

noted for its pyknotic nucleus and a cytoplasm rich in acidophilic Hb. At this stage, the

nucleus is expelled to form reticulocytes (8-10 μm), which then mature into erythrocytes

(7-8 μm) through the final step of erythroid differentiation.

S4 Fig. Gene Ontology (GO) term analysis of plasma membrane proteins.

Scatterplots showing the GO terms: cellular component, molecular function, and

biological process of enriched plasma membrane proteins. The bubble color indicates

the count of the GO term, which was obtained from the DAVID functional annotation

charts. The size of the bubbles indicates the frequency of the GO term in the Gene

Ontology Annotation (GOA) database. The axes of the plot have no intrinsic meaning.

REVIGO (version 1.8.1) was used to generate these scatterplots by reducing the

dimensionality of a pairwise semantic similarity matrix of GO terms using

Multidimensional Scaling (MDS). The resulting projection is not nonlinear. The guiding

principle is that semantically similar GO terms should remain close together in the plot.

S5 Fig. Topology of *P. vivax* merozoite receptor candidates.

(A-J) Predicted locations of potential parasite receptors are illustrated for the signal

peptide (yellow), extracellular domain (blue), intracellular domain (mauve), and

membrane-embedded domain (red).

S6 Fig. Reticulocyte enrichment process.

(A) Initial percentage of reticulocytes in the whole blood sample (0.11%). (B)

Percentage of reticulocytes after enrichment using a 70%-62% Percoll density gradient

(10.5%). (C) Final percentage of reticulocytes (CD45$^-$ and CD98$^+$) following FACS

sorting (87.5%).

A

B *P. vivax*

bp M *PvSx* *Pv* NCh NC

1000

C *P. falciparum*

bp M *PvSx* *Pf* NCh NC

1000

D *P. malariae*

bp M *PvSx* *Pm* NCh NC

1000

E

| *PvSx* | 1 | TTCTAGCTTAATCCACATAACTGATACTTCGTATCGACTTTGTGCGCATTTTGCTATTAT | 60 |
| XR_003001206.1 | 669 | TTCTAGCTTAATCCACATAACTGATACTTCGTATCGACTTTGTGCGCATTTTGCTATTAT | 728 |

| *PvSx* | 61 | GTGTTCTTTTAATTAAAATGATTCTTTTTAAGGACTTTCTTTGCTTCGGCTTGGA | 115 |
| XR_003001206.1 | 729 | GTGTTCTTTTAATTAAAATGATTCTTTTTAAGGACTTTCTTTGCTTCGGCTTGGA | 783 |

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6