MODELING TISSUE-RESIDENT MACROPHAGE DEVELOPMENT FROM MOUSE PLURIPOTENT STEM CELLS

Ann K. Baako[1,2,3], Ragavi Vijayakumar[1], Daniel Medina-Cano[1], Zhaoquan Wang[2], Jesús Romero-Pichardo[2], Kelvin Fadojutimi[2], Stephanie C. Do[1], Yuan Lin[1], Mohammed Islam[1], Sanjana Dixit[1], Alissa J. Trzeciak[2], Justin S.A. Perry[2#], Thomas Vierbuchen[1,2,3#]

AFFILIATIONS
[1] Developmental Biology Program, Sloan Kettering Institute, Memorial Sloan Kettering Cancer Center, New York, NY, USA
[2] Immunology Program, Sloan Kettering Institute, Memorial Sloan Kettering Cancer Center, New York, NY, USA
[3] Neuroscience Program, Weill Cornell Graduate School of Medical Sciences, Cornell University, New York, NY, USA

#These authors contributed equally
*For correspondence: vierbuct@mskcc.org (T.V.), perryj@mskcc.org (J.S.A.P.)

SUMMARY
Tissue-resident macrophages (TRMs) are innate immune cells that participate in tissue development, homeostasis, and immune surveillance. Extensive efforts have been made to recapitulate TRM development from pluripotent stem cells (PSCs) *in vitro* to study molecular and cellular mechanisms of TRM development and to create cellular models of disease. However, available PSC models of mouse TRM development exhibit low overall efficiencies of TRM generation, produce heterogeneous off-target populations, and rely upon undefined media components, thus limiting their reproducibility, scalability, and widespread application as an experimental platform for TRM biology. To address these important limitations, we developed an efficient and reproducible protocol to faithfully recapitulate the stepwise differentiation of mouse PSCs (epiblast stem cells) into unspecialized, proliferative TRMs through the pro-definitive hematopoietic program under defined conditions. These immature TRMs can stably integrate into developing mouse neural organoids *in vitro* and acquire features of microglia. In addition, PSC-derived immature TRMs can stably engraft into the lung niche *in vivo* and adopt alveolar macrophage characteristics. This new platform for modeling mouse TRM development represents a powerful experimental model system for studying TRM function and dysfunction in development and disease.

INTRODUCTION
During early embryogenesis, tissue-resident macrophages (TRMs) – professional phagocytes of the innate immune system that are responsible for engulfing and digesting unwanted material – colonize nearly all developing organs. TRMs co-develop with their respective tissues, acquiring specialized functions that support the specific needs of each tissue during development and homeostasis [1,2]. For example, in the brain, microglia regulate neural circuit formation by pruning excess synapses and clearing apoptotic cells [3,4]. In the lung, alveolar macrophages clear inhaled material and recycle surfactant to maintain proper respiratory function throughout life [5,6]. It is now appreciated that TRMs also play important roles in the pathophysiology of many diseases, including Alzheimer's disease, schizophrenia, and atherosclerosis [1,7]. Given the diverse and widespread functions of TRMs in tissues throughout the body and in various pathological states, there is significant interest in identifying strategies to modulate TRM function and to generate specific TRM populations from pluripotent stem cells (PSCs) for applications in regenerative medicine [8-11].

Pioneering studies in mouse and human models have established that TRMs originate from
erythro-myeloid progenitors (EMPs) born in the yolk sac (referred to as yolk sac hematopoiesis
or pro-definitive hematopoiesis) rather than from bona fide hematopoietic stem cell (HSCs),
which are generated slightly later during definitive hematopoiesis [1,12,13]. In mice, early
mesodermal cells are generated from the posterior primitive streak region and migrate into the
developing yolk sac around embryonic day 6.5-7.0 (E6.5-E7.0), where they differentiate into
extra-embryonic mesoderm [14,15]. Extra-embryonic mesoderm further differentiates into yolk sac
mesoderm that gives rise to the yolk sac endothelium, a subset of which will begin to generate
hemogenic endothelial cells beginning at ~E8.25 [2,16,17]. Yolk sac hemogenic endothelial cells
undergo an endothelial-to-hematopoietic transition (EHT) to produce EMPs, which can either
differentiate locally in the yolk sac into pre-macrophages (pMacs) or enter the nascent
circulation (after E9.5) and colonize the fetal liver [18,19]. Within the fetal liver, EMPs differentiate
into pMacs as well as other hematopoietic progeny that support embryonic development,
including erythrocytes, megakaryocytes, granulocytes, and mast cells [1,17-23]. pMacs exit the fetal
liver and migrate throughout the embryo, seeding developing tissues, undergoing local
expansion, and then further differentiating into specialized, mature TRMs in response to tissue-
specific instructive cues. Importantly, some TRM populations, such as microglia and
Langerhans cells (TRMs that inhabit skin and other epithelial tissues), persist throughout life in
mice with minimal replacement from infiltrating HSC-derived macrophages [24-27].
71
The complex and dynamic nature of yolk sac hematopoiesis makes it challenging to
characterize the underlying molecular and cellular mechanisms. *In vitro* differentiation of
pluripotent stem cells (PSCs) into yolk sac hematopoietic progenitors has emerged as a
powerful experimental system for modeling TRM development [28-31]. For example, human PSC
(hPSC) protocols can recapitulate features of yolk sac hematopoiesis, generating pre-
hematopoietic mesoderm, hemogenic endothelial cells, EMPs, immature TRMs, and specialized
TRMs such as microglia-like cells [32-37]. These reduced complexity models have helped to
delineate critical signaling pathways and transcriptional regulatory mechanisms that orchestrate
specification of mesodermal populations into yolk sac hemogenic endothelium and EMPs [34,35,38-
40]. In addition, generation of human TRMs *in vitro* can be used to generate cellular models of
TRM function and disease biology and can serve as a potential source of cells for next-
generation cell therapies [33,35,41].
84
Although much of our understanding of TRM development and function has come from studies
in mouse models, protocols for differentiating mouse PSCs into TRMs remain significantly less
advanced compared to analogous protocols for human PSCs, and thus they have not been
widely adopted. For example, current protocols for modeling mouse yolk sac hematopoiesis
generate heterogeneous populations of mesoderm in 3D embryoid bodies, require purification of
intermediate populations to achieve a significant yield of desired populations, and utilize
undefined media components such as serum [42-44]. In contrast, there are now hPSC protocols for
stepwise modeling of yolk sac hematopoiesis that more closely recapitulate the relevant
developmental transitions, generate desired populations with moderate/high efficiency, and use
defined media components [35].
95
The gap in protocol development between mouse and human PSC systems can be attributed in
part to differences in PSC culture methods commonly used for each species. Mouse PSCs are
typically cultured in the naïve pluripotent state, whereas human PSCs are cultured in the primed
pluripotent state [45-47]. As a result, naïve mouse PSCs must first transition from a naïve to a
primed pluripotent state to effectively respond to differentiation signals and differentiate into
germ layer progenitors [46,48]. This naïve-to-primed transition involves rapid proliferation and is
inherently asynchronous, thus introducing significant heterogeneity at this initial stage that

complicates the subsequent stages of any differentiation protocol [49]. Recent work from our
group and others suggests using mouse primed-state PSCs (epiblast stem cells [EpiSCs]) as a
starting point for directed differentiation can overcome some of these issues, enabling the
development of robust and efficient differentiation models [50-52].
107
Here, we establish a defined protocol for the stepwise specification of mouse EpiSCs into EMPs
and subsequently into a > 90% pure population of immature TRMs in 2D adherent culture
without the need for purification or co-culture with other cell types. By optimizing the timing and
strength of Wnt and TGF-β signaling, we identify conditions for efficient specification of pre-
hematopoietic mesoderm (PHM), leading to increased yields of EMPs and immature TRMs.
This protocol is reproducible across multiple genetically diverse PSC lines and can be readily
scaled to generate large quantities of EMPs and immature TRMs. Immature TRMs exhibit
surface markers, gene expression profiles, and functional properties characteristic of yolk sac-
derived TRMs *in vivo*. These immature TRMs can integrate into mouse cerebral cortex
organoids and the *in vivo* lung niche, where they adopt features of mature TRMs. This new
platform for modeling mouse TRM development will enable paired *in vitro* and *in vivo* studies
that leverage the full suite of mouse genetic tools, and thus should be a powerful experimental
model system for mechanistic studies of TRM function and dysfunction in development and
disease.
122
DESIGN
Current state-of-the-art protocols for modeling yolk sac hematopoiesis and generating TRMs
from mouse PSCs have significant shortcomings compared to analogous protocols that have
been developed for human PSCs. First, current protocols start from PSCs cultured in the naïve
pluripotent state (mouse embryonic stem cells), which must first transition into a primed
pluripotent state before further differentiation can proceed [45,46]. This initial step is inherently
stochastic, making downstream differentiation stages difficult to control. Additionally, cells at this
stage exhibit high proliferation rates and increased frequency of apoptosis, further complicating
differentiation efforts [46]. Second, current protocols perform differentiation in 3D embryoid body
(EB) cultures, which are more heterogeneous compared to 2D adherent cultures. This
heterogeneity impacts reproducibility, leading to variable differentiation outcomes across
technical replicates and different cell lines. The heterogeneity and asynchronous nature of
differentiations also complicates efforts to model intermediate steps of pro-definitive
hematopoiesis. Finally, many existing protocols rely on serum, which contains complex,
undefined components that can introduce batch-to-batch variability, making it more difficult to
study the specific signaling pathways and other cues critical for each stage of differentiation.
Given these shortcomings, we sought to develop an improved protocol for modeling mouse yolk
sac hematopoiesis (pro-definitive hematopoiesis) and TRM differentiation *in vitro*. Such a model
would be of interest to researchers studying TRM biology and development *in vivo* as it would
serve as a complementary reductionist model for in-depth mechanistic studies.
143
To establish a new mouse model of yolk sac hematopoietic differentiation, we began by
adapting a recently developed 2D, serum-free protocol for generating yolk sac-derived TRMs
from hPSCs [35]. hPSCs are generally cultured in the primed pluripotent state, corresponding to
the pluripotent cells in the post-implantation epiblast, which are competent to directly
differentiated into progenitors of each of the three embryonic germ layers [45,53]. Mouse PSCs can
also be cultured in the primed pluripotent state (EpiSCs), but mouse EpiSCs have not been
broadly adopted as a starting point for *in vitro* directed differentiation studies [47,54]. We recently
demonstrated that hPSC protocols can be adapted for the differentiation of mouse EpiSCs into
definitive endoderm and neuroectoderm [50,55]. Based on these results, we chose to adapt an
hPSC TRM differentiation protocol for directed differentiation of mouse EpiSCs into TRMs [35].

This required careful consideration of species-specific differences, including 1) mouse
development progresses more rapidly than human development, 2) extra-embryonic
mesodermal progenitors experience distinctive signaling environments in mouse compared to
human due to differences in yolk sac development, and 3) the surface marker profile used to
define early yolk sac mesoderm subpopulations in human does not translate directly to mouse.
Consequently, adapting the protocol involved refining timing and developmental cues and
validating markers to align with mouse-specific developmental processes.
161
162 **RESULTS**
163
Identification of conditions for generating yolk sac pre-hematopoietic mesoderm (PHM)
from mouse epiblast stem cells
Starting at ~E6.25, pluripotent cells in the epiblast ingress through the posterior primitive streak
and give rise to nascent mesoderm cells, a subset of which migrate proximally towards the
extra-embryonic ectoderm to become extra-embryonic mesoderm [14,15,56]. Extra-embryonic
mesodermal cells further differentiate into several distinct populations by E7.75-E8.0, including
the allantois core domain, allantois outer mesenchyme, and yolk sac mesoderm [57]. The yolk sac
mesoderm can be distinguished by expression of FOXF1 and GATA6, and by the absence of
T/BRA and SOX17 [57]. A subset of yolk sac mesoderm further differentiates into arterial and
venous endothelial cells that make up the yolk sac endothelium, and some of these endothelial
cells further differentiate into hemogenic endothelium [58-60]. Finally, hemogenic endothelial cells
undergo an endothelial-to-hematopoietic transition (EHT) to produce multipotent erythro-myeloid
progenitors (EMPs), which give rise to embryonic erythroid and myeloid lineages, including
tissue-resident macrophages [61] (**Fig. 1A**). We sought to recapitulate these stepwise
developmental transitions, starting from mouse epiblast stem cells, which are equivalent to
pluripotent epiblast cells in the pre- or peri-gastrulation mouse embryo (**Fig. 1B,**
Supplementary Fig. 1A).
181
However, in mice, the exact surface marker profile of yolk sac PHM that gives rise to hemogenic
endothelium remains incompletely defined, complicating the assessment of specification
efficiency during early differentiation stages. Previous studies in mouse embryos and in
embryoid bodies derived from mouse PSCs found that the $KDR^+ PDGFR\alpha^-$ population is
enriched for cells with hematopoietic potential [42,44,62-64]. Furthermore, a recent study reported
that mouse and human PHM cells are $KDR^+ CXCR4^+$ [65]. Based on these data, we chose to use
$KDR^+ PDGFR\alpha^-$ and $KDR^+ CXCR4^+$ marker profiles for evaluating the efficiency of PHM
induction from EpiSCs.
190
We first sought to identify the appropriate signaling conditions to specify PHM from EpiSCs. We
began by differentiating EpiSCs into T/Bra^+ posterior primitive streak cells using an established
serum-free cocktail of growth factors – FGF2, TGF-β (Activin A), BMP4, and Wnt (CHIR99021)
[35,36,66]. This generates an average of ~80% T/Bra^+ posterior primitive streak-like cells in 24 hours
(n = 3 distinct EpiSC lines) (**Fig. 1C and Supplementary Fig. 1A-C**). To further specify
posterior primitive streak progenitors into $KDR^+ PDGFR\alpha^-$ PHM, we maintained high levels of
BMP4, given its expression throughout the extra-embryonic ectoderm during gastrulation [14], and
examined the impact of modulating the levels of Wnt and TGF-β signaling during the second
day of differentiation.
200
We first modulated Wnt signaling during day 2 (D2) and evaluated $KDR^+ PDGFR\alpha^-$ PHM
generation. To do this, we kept Activin A constant at 10 ng/mL and FGF2 and BMP4 at 10
ng/mL and 40 ng/mL respectively. We tested four different Wnt conditions – Block Wnt (4 nM
LGK-974), endogenous Wnt signaling only (no addition of Wnt inhibitor and no addition of

CHIR99021), moderate Wnt activation (0.5 μM CHIR99021), and high Wnt activation (2 μM
CHIR99021). We found that endogenous Wnt levels yielded the highest number of KDR⁺
PDGFRα⁻ cells (25.8%, **Fig.1C, D**). Interestingly, this result is distinct from what has been
observed in hPSC differentiation, where it is critical to inhibit Wnt signaling shortly after primitive
streak differentiation to specify PHM [35,39,40].
210
Having established the optimal level of Wnt signaling, we next investigated the effect of different
levels of TGF-β pathway activity on the KDR⁺ PDGFRα⁻ population. These experiments
indicated that 10 ng/mL of Activin A generates the highest frequency of KDR⁺ PDGFRα⁻ PHM
cells, and that ~87% of the KDR⁺ PDGFRα⁻ PHM cells in this condition also express CXCR4
(**Fig. 1C, E, F**). Further increasing Activin A (40 ng/mL) caused an increase in the KDR⁺
PDGFRα⁺ population, which is thought to be enriched for cardiogenic mesoderm [42].
Interestingly, the frequency of KDR⁺ CXCR4⁺ cells is more sensitive to the levels of TGF-β
signaling, with the highest fraction of KDR⁺ CXCR4⁺ cells observed at the highest levels of
Activin A (40 ng/mL), and much lower levels observed in conditions where TGF-β is low or
actively blocked (**Supplementary Fig. 1D**). Based on these results, we chose to proceed using
the 10 ng/mL Activin A condition to maximize the KDR⁺ PDGFRα⁻ CXCR4⁺ population.
222
These optimized conditions generated ~40% KDR⁺ PDGFRα⁻ PHM cells at 48 hours.
Importantly, specification of PHM from EpiSCs was reproducible between technical replicates
(the same cell line differentiated in independent experiments) and from a second EpiSC line
from an outbred genetic background, DO-202 (B6/129 n = 13 technical replicates, DO-202 n = 4
technical replicates) (**Fig. 1F**). Further characterization of differentiating cultures at 48 hours
revealed cells expressing markers of yolk sac mesoderm/PHM (FOXF1, GATA6) surrounding
clusters of T/Brachyury⁺ cells, which are likely primitive streak-like cells or off-target extra-
embryonic mesoderm subtypes that express T/Brachyury (**Supplementary Fig. 1E**). These
optimized conditions for specification of PHM-enriched extra-embryonic mesoderm from EpiSCs
provide a strong foundation for modeling subsequent stages of yolk sac hematopoiesis.
233
Efficient generation of erythro-myeloid progenitors from pre-hematopoietic yolk-sac
mesoderm
After optimizing conditions for generation of yolk sac PHM, we next sought to determine
whether PHM can be further differentiated into yolk sac hemogenic endothelium and EMPs. To
generate EMPs, we adapted differentiation conditions previously used for hPSC differentiation
into EMPs, reducing the total time allotted for each step of differentiation to account for the
faster pace of murine development compared to human. (**Fig. 2A**). We first induced hemogenic
endothelium generation by adding VEGF to PHM cultures. Once hemogenic endothelium had
been generated, we sequentially supplemented media with SCF, IL-6, TPO, and IL-3, cytokines
known to induce hematopoiesis and promote EMP expansion [35]. In the developing mouse
embryo, EMPs emerge from yolk sac hemogenic endothelial cells via an endothelial-to-
hematopoietic transition and can be isolated based on their expression of the surface markers
Kit, CD41, and CD16/32 [67]. Using this surface marker profile, we monitored the generation of
EMPs in differentiating cultures between D2 and D7 (**Fig. 2B**). Semi-adherent EMPs were first
observed at low frequency at D5 (~6%) and peaked at ~20% by D7-D8, comparable to the
~17% EMP frequency reported in a similar hPSC-based model [35]. Under these conditions,
essentially all EMPs also expressed the pan-myeloid marker CD45 and the transcription factor
RUNX1, which is required for EMP generation from yolk sac hemogenic endothelium (**Fig. 2B,**
C) [68]. Importantly, we found that EMP generation was reproducible across technical replicates
and in a second EpiSC line, DO-202, albeit at slightly lower efficiency (n = 4 independent
differentiation experiments for each cell line, **Fig. 2B**). Taken together, these data demonstrate

that PHM generated from EpiSCs can be efficiently and reproducibly differentiated into EMPs
over a period of 3-5 days.
257
Differentiation of EMPs into immature TRMs and characterization of immature TRMs
To further differentiate EMPs into pMacs and immature TRMs, we dissociated differentiating
cultures at D8 and replated cells onto Matrigel-coated tissue culture plates in serum-free media
containing M-CSF and IL-34, growth factors that both signal through the CSF1 receptor
(CSF1R) to stimulate TRM differentiation and proliferation during embryonic development (**Fig.**
3A). After two days in these conditions (D10 overall), round, vacuolated, immature TRMs with
filopodia were present throughout the culture (**Fig. 3B**). Immature TRMs are highly proliferative
and continue to divide for at least 8 days under these conditions, giving rise to nearly pure
populations of TRMs that express the pan-macrophage markers PU.1 and IBA1 by D16 (**Fig.**
3B). In addition, D16 TRMs uniformly express the pan-myeloid marker CD45 and the pan-
macrophage markers CD11b and CX_3CR1. A subset of TRMs (~16%) express P2RY12, a
receptor that is preferentially expressed on microglia and on some Kupffer cells *in vivo* [69-71] but
generally not highly expressed on TRMs *in vitro* [72-74]. Importantly, we observed efficient
generation of CX_3CR1^+ cells at D16 across multiple, independent differentiations from three
distinct EpiSC lines (B6/129 [n = 6 technical replicates from independent differentiations], DO-
202 [n = 3 technical replicates from independent differentiations], and DO-223 [n = 3 technical
replicates from independent differentiations], **Fig. 3C**). Beginning with 20,000 EpiSCs in a 96-
well plate, each differentiation yielded approximately 360,000 CX_3CR1^+ immature TRMs per cell
line.
277
To further evaluate the properties of immature TRMs, we performed RNA-seq on CX_3CR1^+-
sorted immature TRMs from three distinct EpiSC lines (B6/129, DO-202, DO-223; n = 2
technical replicates per cell line). Consistent with the flow cytometry data, immature TRMs
exhibited robust expression of macrophage-enriched genes (*Csf1r*, *Mpeg1*, *Slc6a6*, *Trem2*) as
well as modest expression of TRM-enriched genes including *Tmem119* and *P2ry12* (**Fig. 3D**). In
contrast, genes that are expressed specifically in mature, functionally specialized subtypes of
TRMs, such as *Sall1* (microglia), *Nfatc4* (kidney TRMs), and *Spi-c* (spleen TRMs) were
expressed at negligible levels, consistent with prior work indicating that TRMs *in vitro* fail to
express markers characteristic of mature TRMs *in vivo* [72-74] (**Fig. 3D**). Immature TRMs also do
not express Hox genes (*Hoxa5, Hoxa9, Hoxb4, Hoxb5, Hoxb8, Hoxc6*) that are normally
expressed by HSC-derived macrophages [75-77], consistent with their generation via EMPs in the
pro-definitive hematopoietic program (**Fig. 3D)**. In addition, to evaluate whether immature TRMs
were more similar to pMacs or embryonic TRMs, we examined the expression of gene
signatures characteristic of *in vivo* pMacs from E9.5 and E10.25 yolk sac and head (n = 50
genes) and macrophages from E10.25 and E10.5 yolk sac and head (n = 71 genes, **Fig. 3E**) [19].
In vitro-derived immature TRMs shared transcriptional features with both *in vivo* pMacs and
embryonic macrophages, supporting their identity as an intermediate TRM state fated to mature
into specialized TRMs.
296
297 We then examined the functional capacity of immature TRMs to phagocytose CypHer 5E-
298 labeled substrates. The immature TRMs phagocytosed synaptosomes and apoptotic neurons
299 within 45 minutes of exposure to the substrates (**Fig. 3F, Supplementary Video**).
300
To evaluate the potential of immature TRMs to acquire features of more specialized, tissue-
specific macrophage populations *in vitro,* we chose to culture immature TRMs in conditions that
promote acquisition of some features of microglia identity. To do this, at D11 we switched
immature TRMs into media containing IL-34, TGF-β, and cholesterol based on a serum-free
cocktail (TIC media) that was developed to promote primary microglia survival *in vitro* [72].

Specifically, we made a slight modification to this media composition by maintaining our
macrophage base media but supplementing with the same concentrations of the key cytokines,
lipids and acids found in the original TIC media. We refer to our media as the B27-TIC media
and draw comparisons with two other media compositions – the original TIC media and
immature TRM media (**Supplementary Fig. 2A**).

Both TIC and B27-TIC media conditions led to reduced proliferation and enhanced ramification
of TRMs, whereas cells in immature TRM conditions remained rounded and continued to
proliferate (**Supplementary Fig. 2A**). The highest degree of ramification was observed when
cells in the two TIC conditions were split at D16 and replated onto collagen-coated tissue culture
plates (**Supplementary Fig. 2A**). Surprisingly, we observed that attachment and survival of
TRMs/microglia-like cells was significantly better in the B27-TIC media compared to the TIC
condition, and thus we chose to use B27-TIC for subsequent experiments. We then compared
transcriptomes of immature TRMs and microglia-like cells from three distinct EpiSC lines by
isolating CX_3CR1^+ cells by FACS in each condition and performing bulk RNA-seq. Principal
component analysis (PCA) revealed that TRMs in each condition separated by cell line on PC1,
and then by media condition on PC2, suggesting that the differences in transcriptome between
genetically distinct EpiSC lines were larger than the differences between TRMs in each of the
two media conditions (**Supplementary Fig. 2B**). Compared to immature TRMs, the B27-TIC
microglia-like cells expressed higher levels of *in vivo* microglia signature genes, including
P2ry12, *Tmem119*, *Mertk*, *Tgfbr1*, *Hexb*, and *Sall3* (**Supplementary Fig. 2C**). B27-TIC
microglia-like cells also had strong expression of genes expressed in primary microglia isolated
from the brain at E14.5 and postnatal day 4/5 (P4/5) [78], further suggesting that they can acquire
features of immature microglia (**Supplementary Fig. 2D**).

To compare B27-TIC microglia-like cells to primary microglia more directly, we isolated primary
$CD11b^+$ microglia from P4 mice and cultured them in B27-TIC for 6 days. We then examined
expression of surface markers broadly expressed by TRMs (CD45, CD11b, CX_3CR1) and more
specifically enriched on microglia (P2RY12). Cultured primary microglia and B27-TIC microglia-
like cells expressed similar levels of CX_3CR1, CD11b, and P2RY12 within their respective
$CD45^+$ populations (**Supplementary Fig. 2E**). In addition to their transcriptional and phenotypic
similarities to neonatal microglia, microglia-like cells displayed functional properties similar to
their *in vivo* counterparts. Microglia-like cells successfully phagocytosed synaptosomes,
apoptotic Jurkat cells, and the yeast antigen zymosan. Fixed confocal imaging confirmed that
the microglia-like cells also expressed CD68, a marker of macrophage activation
(**Supplementary Fig. 2F**). Thus, by culturing immature TRMs in conditions that more closely
mimic the CNS environment, immature TRMs generated from mouse EpiSCs can be further
induced to acquire some features of microglia, exhibiting general similarity to *ex vivo* cultured
microglia from neonatal mice.

Immature TRMs integrate into developing cortical organoids and adult mouse lung niche
The terminal differentiation and maintenance of functionally specialized TRMs requires
continuous instructive signaling from their unique tissue microenvironments. Previous studies
suggest that these tissue-specific signals are difficult to replicate in simple tissue culture
models, and thus *ex vivo* and PSC-derived TRMs are more similar to immature, unspecialized
TRMs [72]. More complex models of developing tissues, such as 3D tissue organoids, can more
closely recapitulate the complex signaling environment found in tissues *in vivo*. Therefore, we
next sought to determine if immature TRMs can colonize mouse cerebral cortical organoids and
acquire characteristics of microglia. We used a cerebral cortical organoid differentiation protocol
established in our laboratory to generate cortical organoids [55]. These organoids contain only
neuroectodermal cell types, and thus do not contain microglia (**Supplementary Fig. 3**). We

added immature TRMs (D28) to D10 cortical organoids, a time point that corresponds
approximately to the mid-to-late stages of cortical neurogenesis *in vivo* (~E15.5-E17.5). Cortical
organoids with integrated TRMs were then cultured for an additional 14-21 days in cortical
organoid media supplemented with IL-34 and TGF-B to promote proliferation and microglial
differentiation of transplanted immature TRMs (**Fig. 4A**). Two weeks post-transplantation,
approximately 7% of the cells within the cortical organoids expressed the microglia marker
CD11b, with this population increasing to around 40% after an additional week of culture (**Fig.
4B**). The relative increase in abundance of transplanted TRMs over time likely reflects both
microglial proliferation and death of neuronal and glial cells within the organoids.
Immunofluorescence staining of tissue-cleared cortical organoids demonstrated that
transplanted TRMs successfully integrated into the neuroepithelium of the developing
organoids, proliferated, migrated throughout the organoids to tile the developing neuroepithelial
tissue, and acquired ramified morphologies similar to microglia in developing cerebral cortex *in
vivo* (**Fig. 4C, Supplementary Fig. 3**). These experiments provide a proof-of-concept that
mouse EpiSCs can be used to generate chimeric cerebral cortical organoids containing
microglia-like TRMs.
373
Finally, we sought to assess the capacity of immature TRMs to engraft into specific tissue
niches in adult mice. We selected the lung as our model due to the availability of a well-
established, efficient protocol for adoptive cell transfer. We first depleted the endogenous
alveolar macrophage population in CD45.1 congenic mice using clodronate liposomes [79]. We
then transplanted immature CD45.2$^+$ TRMs intratracheally and allowed them to colonize lung
tissue for 3 days or 7 days (**Fig. 4D**). By day 3, approximately 25% of the exogenous CD45.2
TRMs present in the lungs had reduced CD11b expression, which is typically observed in
alveolar macrophages. Among these CD45.2$^+$ CD11b$^-$ cells, approximately 65% also expressed
the alveolar macrophage surface markers CD11c and SiglecF, suggesting that transplanted
immature TRMs had successfully integrated into the alveolar niche and acquired features of
differentiated alveolar macrophages. Similar results were observed 7 days after transplantation,
suggesting stable engraftment of EpiSC-derived TRMs (**Fig. 4E, F**).
386
387 Taken together, these data demonstrate that immature TRMs can integrate into two distinct
388 specialized tissue environments and acquire characteristics of functionally specialized, mature
389 TRMs, highlighting the potential utility of PSC-derived immature TRMs for studying TRM
390 development and function in various tissue contexts.
391
DISCUSSION
Here, we developed a defined (serum-free), adherent (2D) protocol for recapitulating mouse
yolk sac hematopoiesis (pro-definitive hematopoiesis) starting from primed pluripotent stem
cells (EpiSCs). This protocol can generate > 90% pure population of immature TRMs in 16 days
without the need for purification or co-culture with other cell types. This new protocol addresses
several of the limitations of EB-based approaches, leading to improved differentiation efficiency,
scalability, and reproducibility across distinct cell lines. Given these advantages, this protocol
represents a powerful new model system for studying molecular and cellular mechanisms of
yolk sac hematopoiesis and TRM differentiation. To facilitate adoption of this protocol by other
laboratories, we have included a detailed, step-by-step protocol (see **Supplementary
Experimental Procedures**).
403
In vitro directed differentiation models enable a variety of experimental designs that are not
currently feasible *in vivo*, including large-scale genetic or chemical screens, live imaging, and
biochemical studies that require large numbers of cells. Our improved *in vitro* model of mouse
yolk sac hematopoiesis will thus allow for the application of these powerful experimental

approaches to dissect mechanisms of cell fate specification that have proven difficult to study *in*
vivo. Furthermore, using mouse EpiSCs means it is possible to take advantage of the extensive
mouse genetic tools that have been developed to study yolk sac hematopoiesis, including
knockout mouse lines and reporter alleles. Unlike human PSC models, findings from mouse
PSC models can also be rigorously evaluated *in vivo* under controlled conditions. Given these
advantages, we believe that this new mouse EpiSC model for studying yolk sac hematopoiesis
and TRM differentiation can complement and empower existing model systems and will thus be
of broad interest.
416
There is significant interest in incorporating TRMs into various tissue organoid models to better
recapitulate the cellular complexity of tissues *in vivo* and to examine the crosstalk between
TRMs and their host tissues [33]. We show that mouse TRMs generated from mouse EpiSCs can
integrate into developing mouse cerebral cortical organoids, generating the first chimeric neuro-
immune organoid model derived completely from mouse PSCs. In the future, it will be
interesting to use this system to model mechanisms of microglial development,
neuroinflammation, and neurodegenerative disease. With additional development, this tractable
in vitro model could also serve as a scalable platform for genetic or chemical screening [80-82].
425
Perhaps the most translationally relevant application of our system is the adoptive transfer of *in*
vitro-derived cells into *in vivo* tissue niches. This approach serves as a powerful platform to
investigate disease pathogenesis and test therapeutic efficacy *in vivo*. For instance, transferring
TRMs into mouse brains or lungs could assess their ability to rescue phenotypes in models of
Alzheimer's disease or respiratory disease. Unlike human cell transplantation into mice, which
requires complex immunosuppression techniques to prevent xenogeneic rejection [83,84], mouse-
to-mouse adoptive transfers bypass this requirement when cells are matched to the host strain.
This technical simplicity allows for more straightforward experimental designs and reproducible
outcomes, offering new opportunities for studying therapeutic interventions.
435
436 **Limitations of the Current Study**
437 While this new protocol can generate PHM and EMPs with reasonable efficiency, off-target
438 mesodermal populations are still produced. Further optimization will be necessary to improve
439 PHM and EMP yields. In addition, we have not characterized the differentiation of EMPs into
440 other lineages (megakaryocytes, granulocytes, mast cells) [67].
441
442 Although we observed that culturing immature TRMs in B27-TIC media increased expression of
443 some microglia-enriched genes, this *in vitro* model does not fully recapitulate the gene
444 expression signatures of microglia *in vivo* (consistent with previous studies) [72]. Achieving fully
445 mature TRM identities *in vitro* will require exposure to additional niche-specific cues, including
446 organ-specific extracellular matrix components or biomechanical cues [85].
447
448 Finally, while our adoptive transfer studies confirm that *in vitro*-derived TRMs can integrate into
449 host tissues, the long-term stability and functional equivalence of these cells compared to
450 endogenous TRMs will require further investigation. Future longitudinal studies tracking
451 engrafted TRMs in vivo will be essential for assessing their durability and evaluating their
452 therapeutic potential.

Figure 1: Identification of conditions for generating yolk sac pre-hematopoietic mesoderm (PHM) from mouse EpiSCs

Figure 1: Identification of conditions for generating yolk sac pre-hematopoietic mesoderm (PHM) from mouse EpiSCs

A. Overview of developmental stages from pluripotency through yolk sac hematopoiesis and TRM differentiation, including relevant stages of embryonic development in mouse (bottom row). **B**. Overview of approach for differentiation of EpiSCs into immature TRMs, including stage-specific marker genes. **C**. Summary of the first two stages of the TRM differentiation protocol. The optimal levels of Wnt and TGF-β signaling to promote KDR(+) PDGFRa(-) yolk sac mesoderm/PHM differentiation during Stage 2 were not known. **D**. Representative flow cytometry plots (left) of D2 PHM yield observed at different levels of Wnt pathway activation. CHIR99021 was used to activate the canonical Wnt pathway, and LGK479 was used to inhibit Wnt signaling. Activin A (AA) was kept constant at 10 ng/mL during stage 2 for each condition. Summary of flow cytometry data across 3 independent differentiation experiments (right). Significance was determined using ordinary one-way ANOVA, corrected for multiple comparisons with Tukey test (***P < 0.001). **E**. Representative flow cytometry plots (left) showing D2 PHM yield observed at different levels of TGF-β pathway activation. Activin A was used to stimulate TGF-β signaling and SB431542 was used to block TGF-β signaling. Only endogenous Wnt signaling (same as the No Wnt condition in [D] was used in Stage 2). Summary of flow cytometry data across 3 independent differentiation experiments (right). Significance was determined using ordinary one-way ANOVA, corrected for multiple comparisons with Tukey test (*P < 0.05; **P < 0.01; ***P < 0.001).

F. Representative flow cytometry plots (left) measuring PHM yield at the end of Stage 2 in optimized PHM conditions (No Wnt, 10 ng/mL Activin A). Percentage of CXCR4 within the KDR(+) PDGFRa(-) population is also shown. Summary of flow cytometry data from 2 distinct cell lines (right). B6/129 EpiSCs (n = 13 independent differentiations [technical replicates]). DO-202 EpiSCs (n = 4 independent differentiations [technical replicates])

Figure 2: Efficient generation of erythro-myeloid progenitors from pre-hematopoietic yolk sac mesoderm

Figure 2: Efficient generation of erythro-myeloid progenitors from pre-hematopoietic yolk sac mesoderm
A. Detailed overview of signaling conditions for hemogenic endothelium differentiation and hematopoiesis. These conditions are adapted (with modifications) from Guttikonda et al., 2021. **B**. Evaluation of EMP emergence during in vitro hematopoiesis (left). Generation of Kit(+) CD41(+) CD16/32(+) EMPs and myeloid marker expression (CD45) was evaluated using flow cytometry at the indicated timepoints. Flow cytometry plots are gated for Kit(+) cells (top row). Under these conditions, essentially all EMPs also express CD45 (bottom row). Quantification of EMP generation (D8) across 4 independent differentiation experiments from the indicated cell lines (right). **C**. Immunofluorescence staining of differentiating cultures at D8 for RUNX1, a transcription factor expressed in yolk sac hemogenic endothelia and EMPs. EHT = endothelial-to-hematopoietic transition. Scale bars = 25 μM.

Figure 3: Differentiation of EMPs into immature TRMs and characterization of immature TRMs

Figure 3: Differentiation of EMPs into immature TRMs and characterization of immature TRMs
A. Overview of conditions used for differentiation of EMPs into immature TRMs and TRM expansion. **B**. Brightfield images of immature TRMs at D10 and D16 (top row) and immunofluoresence staining immature TRMs (D16) for pan-macrophage markers IBA1 and PU.1 (bottom row). **C**. Assessment of TRM surface markers CX_3CR1, CD11b, CD45, and P2RY12 by flow cytometry to evaluate immature TRM differentiation efficiency (left). Summary of differentiation efficiency (% CX_3CR1(+) cells) obtained from the indicated EpiSC lines (right). Each dot represents an independent differentiation experiment. **D**. Heatmap of gene expression (RNA-seq) of a panel of cell type-specific marker genes in CX_3CR1(+) immature TRMs purified by FACS at D16. Log2 normalized counts (transcripts per million [TPM]) are plotted for each gene (average expression level from n = 2 independent differentiations [technical replicates] from each of the indicated cell lines). **E**. Violin plots of expression levels (log2 normalized counts, TPM) of individual genes from signatures of yolk sac-derived pMacs and yolk sac-derived macrophages obtained from E10.25 and E10.5 embryos (Mass et. al., 2016). Each dot represents a single gene, and the value for the expression level of the gene is the average expression across TRMs from all of the 3 EpiSC lines. **F**. Confocal microscopy images (brightfield overlaid with fluorescence) of D20 immature TRMs containing pH-sensitive CypHer 5E-labeled phagocytosed synaptosomes (left) and apoptotic neurons (right). Arrows indicate engulfed cellular material. Scale bars = 25 μm.

Figure 4: Immature TRMs integrate into developing cortical organoids and adult mouse lung niche

Figure 4: Immature TRMs integrate into developing cortical organoids and adult mouse lung niche
A. Overview of strategy for co-culture of immature TRMs and mouse cerebral cortex organoids generated from EpiSCs.
B. Quantification of CD11b(+) TRMs in cortical organoids using flow cytometry at D24 and D32. **C**. Confocal imaging of tissue-cleared cortical organoids with integrated TRMs. Immunofluorescence staining for TRM markers PU.1 and IBA1. High magnification image (right) showing ramified morphology of TRMs. **D**. Overview of experimental design for depletion of endogenous alveolar macrophages and subsequent intratracheal delivery of immature TRMs into mouse lungs. CD45.1 mice were used so that transplanted CD45.2-expressing TRMs can be distinguished from CD45.1(+) host macrophages. **E**. Representative flow cytometry plots for lung tissue from mice transplanted with TRMs at the indicated timepoints. Alveolar macrophages are CD11b(-) SiglecF(+) CD11c(+). **F**. Summary of results from lung transplantation experiments at D3 and D7 post-transplant. Individual dots represent results from a single mouse (biological replicates).

Supplementary Figure 1: Identification of conditions for generating yolk sac pre-hematopoietic mesoderm (PHM) from mouse EpiSCs

Supplementary Figure 1: Identification of optimal conditions for generating yolk sac pre-hematopoietic mesoderm (PHM) from EpiSCs

A. Summary of the first stage of the TRM differentiation protocol. **B**. Immunofluorescence staining for brachyury (T) was used to evaluate the percentage of cells that have acquired a primitive streak-like identity at the end of stage 1 (d1). **C**. Quantification of Brachyury (T) positive cells at the end of stage 1 across 3 distinct EpiSC lines (B6/129, DO-202, DO-223). Three independent differentiation experiments were performed for each EpiSC line, with two technical replicates per cell line. **D**. Representative flow cytometry plots showing D2 KDR(+) CXCR4 (+) yield across TGF-β conditions (left). Summary of flow cytometry data across 3 independent differentiation experiments (right). Significance was determined using ordinary one-way ANOVA, corrected for multiple comparisons with Tukey test (not significant [ns] P > 0.05; **P < 0.01; ***P < 0.001). **E**. Immunofluorescence staining for additional markers of PHM (FOXF1, GATA6) in differentiating cultures at the end of Stage 2 (D2). Scale bars = 100 μm.

Supplementary Figure 2: Characterization of TRMs cultured in *ex vivo* microglia maintenance conditions

Supplementary Figure 2: Characterization of TRMs cultured in *ex vivo* microglia maintenance conditions
A. Overview of different media conditions for culturing immature TRMs (left). Bright field images of TRMs in each condition (right) and immunofluorescence staining for cell proliferation (Ki-67) (far right). **B**. Principal component analysis (PCA) plot of RNA-seq data from D16 TRMs cultured in each of the three media conditions (color coded based on [A]). **C**. MA plot comparing gene expression of D16 immature TRMs and TRMs cultured in B27-TIC. A panel of genes relevant for immature TRMs and microglia are annotated. Statistical significance was calculated using a Wald test and corrected using the Benjamini-Hochberg method; gray dots $P > 0.1$; blue dots $P < 0.1$ **D**. Violin plots of gene expression (RNA-seq) in D16 B27-TIC cells. Gene signatures are derived from microglia isolated at E14.5 or P4/P5 (Hammond et al., 2019). Each dot represents the average expression level of gene (n = 3 biological replicates). **E**. Representative geometric mean fluorescence intensity (gMFI) plots comparing microglia surface marker profiles of primary microglia and B27-TIC microglia-like cells; gated for CD45(+) expression (top row). Quantification of gMFI across biological replicates (bottom row). Statistical significance was determined using Welch's t-test (ns $P > 0.05$). **F**. Confocal imaging of B27-TIC TRMs (D20) phagocytosing pH-sensitive CypHer 5E-labeled synaptosomes (left). Immunofluorescence staining for pan-macrophage marker CD68 and IBA1 in fixed B27-TIC TRMs that were exposed to CypHer 5E-labeled apoptotic Jurkat T cells or AlexaFluor 594-conjugated zymosan beads. Arrows indicate engulfed targets (right). Scale bars = 25 μm. DIV = days *in vitro.*

Supplementary Figure 3: Immature TRMs integrate into developing cortical organoids and adult mouse lung niche

Supplementary Figure 3: Immature TRMs integrate into developing cortical organoids and adult mouse niche
Immunofluorescence staining of D16 cortical organoids for markers of deep-layer cortical neurons (CTIP2), upper-layer neurons (SATB2), telencephalic progenitors (FOXG1) and astrocytes (GFAP). Scale bars = 30 μm

ACKNOWLEDGEMENTS
We would like to thank our colleagues in the Vierbuchen and Perry laboratories for helpful discussions and advice throughout the course of the study. We would also like to thank Tomi Lazarov and Laura Santini for providing helpful comments on the manuscript.

This study would not have been possible without the core facilities at MSKCC. We would like to specifically thank members of the Flow Cytometry Core facility and the Integrated Genomics Operation Core for their advice and assistance with experiments and data analysis throughout the course of this project.

Some figure panels were created using the commercial version of Biorender.

AUTHOR CONTRIBUTIONS
Conceptualization: A.K.B., J.S.A.P., T.V.
Methodology: A.K.B., R.V., Z.W., J.E.R.P., K.F., M.I., S.D, S.C.D., Y.L., A.T.
Formal Analysis: A.K.B., R.V., D.M.C., Y.L.
Investigation: A.K.B., R.V.
*Resources***:** J.S.A.P., T.V.
Data Curation: A.K.B., R.V.
Writing – original draft: A.K.B., J.S.A.P., T.V.
Writing – review and editing: A.K.B., J.S.A.P., T.V.
Supervision: J.S.A.P., T.V.
Project Administration: J.S.A.P., T.V.
Funding Acquisition: J.S.A.P., T.V.

DATA AVAILABILITY
Source data and processed files are available in the Gene Expression Omnibus under the accession number GSE296070.

DECLARATION OF INTERESTS
J.S.A.P is a co-founder of Atish Technologies.

FUNDING
This study was made possible by financial support to T.V. from the following sources: Startup funding for the Vierbuchen Lab provided by Sloan Kettering Institute for Cancer Research (MSKCC) and the Josie Robertson Investigator Program (T.V.), an NIH Cancer Center Support Grant (NIH P30 CA008748). The work was also supported by grants to J.S.A.P. from the NIH (NCI 5R00CA237728; NIGMS 1DP2GM146337), a Pew Biomedical Scholars Award, a pilot award from the CSBC Research Center for Cancer Systems Immunology at MSKCC (NCI 5U54CA209975). A.K.B. is supported by an MSKCC MERIT Mandel Fellowship.

MATERIALS AND METHODS

Differentiation of EpiSCs into immature TRMs.
Mouse EpiSCs were generated from ESCs as described in previous work from our lab [50]. The three EpiSC cell lines used were 1) C57BL/6J X 129S1/SvImJ F1 hybrid genetic background (B6/129), 2) DO-202 [86], and 3) DO-223 [86]. EpiSCs were karyotyped to assess chromosomal integrity and tested for mycoplasma to rule out bacterial contamination. EpiSCs were cultured on gelatin-coated dishes with irradiated mouse embryonic fibroblasts in N2B27 media (50% DMEM-F12, 50% Neurobasal medium supplemented with 0.5% N2 supplement, 1% B27 supplement without vitamin A, 2 mM GlutaMAX, 1% penicillin-streptomycin, 0.1% 2-

510   mercaptoethanol) supplemented with 20 ng/mL Activin A, 12.5 ng/mL heat-stable bFGF (FGF2),
511   and 175 nM NVP-TNKS656. Media was changed daily, and cells were passaged every ~48
512   hours at a 1:6 ratio after dissociation with Accutase into small clumps of 3-5 cells. All cells were
513   maintained at 37°C in 5% CO_2 under normoxic conditions.
514
EpiSCs between passages 8-12 were cultured to confluency and detached from the fibroblast
layer using 0.1 U/µL collagenase IV, followed by resuspension in Accutase for 1-2 minutes to
obtain a single-cell suspension. Plates for differentiation were sequentially coated first with 16.7
µg/mL Fibronectin in PBS$^{-/-}$, washed twice with PBS$^{-/-}$, and then coated with 10 µg/mL
Laminin521 in PBS$^{+/+}$ overnight. Cells were plated at a density of 60,000 cells/cm^2 in Lo FABC-
R media, which consisted of chemically defined medium 2 (CDM2) (See supplementary Table 1
for recipe) supplemented with 5 ng/mL FGF2, 30 ng/mL BMP4, 5 ng/mL Activin A, 3 µM CHIR
99021, and 50 nM Chroman-1 and 5 µM Emricasan (See Supplementary Table 1 for detailed
information on how to prepare media) .
524
After 24 hours, the medium was switched to Hi FAB media (CDM supplemented with 10 ng/mL
FGF2, 10 ng/mL Activin A, and 40 ng/mL BMP4). On day 2, media was replaced with VF media
(Essential 6 medium containing 15 ng/mL VEGF and 5 ng/mL FGF2). VF media was refreshed
at the end of day 3. On day 4, cells were cultured in VFSI media (Essential 6 medium
supplemented with 15 ng/mL VEGF, 5 ng/mL FGF2, 200 ng/mL SCF, and 20 ng/mL IL-6). On
day 5, cells were transferred to SITI media (Essential 6 medium containing 100 ng/mL SCF, 10
ng/mL IL-6, 30 ng/mL TPO, and 30 ng/mL IL-3), with daily media changes until day 7 or 8. Cells
were never washed between media changes.
533
On day 7 or 8, cells were dissociated into single cells with Accutase and replated in
macrophage media at a density of 60,000–120,000 cells per cm^2 on Matrigel-coated plates.
Macrophage media consisted of 75% IMDM with GlutaMAX, 25% F12 with GlutaMAX, 1:100
B27 supplement without vitamin A, 100 µg/mL penicillin-streptomycin, 100 ng/mL mouse M-
CSF, and 100 ng/mL mouse IL-34. Macrophage media was changed three times a week, and
cells were passaged every 7-8 days by simply pipetting to detach them from the plates and
replating at 60,000 cells per cm^2.
541
542   Additional details about reagents and differentiation media compositions are provided in
543   Supplementary Table 1 and the Key Resources Table.
544
545   **Fluorescence-Activated Cell Sorting (FACS) analysis**
546   For 2D cultured cells, cell dissociation was performed with Accutase for 1-2 minutes, followed
547   by washing in FACS buffer (1% BSA, 5 mM EDTA). Organoid cells were dissociated using
548   Accutase containing 30 µg/mL DNase I for 30-45 minutes. After dissociation, cells were washed
549   and resuspended in FACS buffer with the appropriate antibodies.
550
551   Day 16+ macrophages were incubated with Fc block (unconjugated mouse anti-CD16/32 Bio X
552   Cell, 1:50) for 10 minutes to prevent nonspecific Fc receptor binding before antibody staining.
553   Cells were then washed, resuspended in FACS buffer, and filtered through 40 µm filters prior to
554   analysis on an Attune Nxt flow cytometer. Data was analyzed using FlowJo v10.10.0.
555
556   Sorting of CX$_3$CR1$^+$ macrophages was done at the MSKCC Flow Cytometry Core Facility using
557   either BD FACSAria or BD FACSymphony S6 cell sorters.
558
FACS antibodies and dilutions used were as follows:

Day 2 cells: KDR-PE (BioLegend, 1:200), PDGFRα-PE/Cy7 (BioLegend, 1:200), CXCR4-Alexa
Fluor 647 (BioLegend, 1:200)
Day 8 cells: Kit-BV421 (BD Biosciences, 1:200), CD41-FITC (BioLegend, 1:200), CD16/32-
BV711 (BioLegend, 1:200), CD45-Alexa Fluor 700 (BioLegend, 1:200)
Day 16+ macrophages: unconjugated anti-mouse CD16/32, CX₃CR1-BV785 (BioLegend,
1:200), CD11b-FITC (BioLegend, 1:200), CD45-Alexa Fluor 700 (BioLegend, 1:200), P2RY12-
PE (BioLegend, 1:200)
567
568　Detailed information on all FACS antibodies is available in Supplementary Table 2.
569
Immunostaining of 2D cells
Cells were washed twice with PBS-/- and fixed with 4% paraformaldehyde (PFA) for 15 minutes
at room temperature. After fixation, cells were rinsed twice with PBS-/- and permeabilized with
PBS-/- containing 1% Triton-X for 10 minutes at room temperature. Cells were blocked for 30
minutes with PBS-/- with 0.3% Triton-X and 5% BSA. Primary antibodies, diluted in PBS
containing 0.1% Triton-X (PBST) with 1% BSA, were then added overnight at 4°C. The following
primary antibodies were used: rabbit anti-Brachyury/T (Abcam, 1:1000), goat anti-Brachyury/T
(R&D Systems, 1:200), goat anti-FOXF1 (R&D Systems, 1:400), rabbit anti-GATA6 (Cell
Signaling Technology, 1:500), rabbit anti-VE-Cadherin (ThermoScientific, 1:200), rabbit anti-
Runx1 (ThermoScientific, 1:400), goat anti-Iba1 (Novus Biologicals, 1:500), rabbit anti-Pu.1
(ThermoScientific, 1:200), rat anti-Ki-67 (Invitrogen, 1:500). Detailed information about
antibodies can be found in Supplementary Table 3.
582
583　After overnight incubation with primary antibodies, cells were washed three times with PBST for
584　10 minutes each. Secondary antibodies, diluted 1:500 in PBST with 1% FBS, were added for 2
585　hours at room temperature. After three additional 10-minute washes with PBST, cells were
586　stained with DAPI (1 µM in PBST) for 15 minutes at room temperature. Cells were then rinsed
587　with PBST and imaged using a Leica Dmi8 microscope.
588
Differentiation of immature TRMs into microglia-like cells.
After 3-4 days of proliferation in macrophage media, around day 11, cells were transitioned to
microglia media without additional passaging. Two formulations of microglia media were tested:
the first, TIC media, is an established serum-free medium for culturing primary mouse and rat
microglia[72]. This media contains DMEM-F12, 2 mM GlutaMAX, 100 µg/mL penicillin-
streptomycin, 1X Insulin-Transferrin-Selenium, and 1 mg/mL N-acetyl cysteine (NAc),
supplemented with 2 µg/mL TGF-β, 100 ng/mL IL-34, 1.5 µg/mL cholesterol, 0.1 mg/mL oleic
acid, 0.001 µg/mL gondoic acid, and 10 µg/mL heparan sulfate.
597
The second formulation, a modified TIC media we refer to as B27-TIC, comprised of a base of
75% IMDM with GlutaMAX, 25% F12 with GlutaMAX, and 1:100 B27 supplement without
vitamin A, supplemented identically to TIC media. After 5-6 days in TIC media, cells were split
and replated at a density of 120,000 cells/cm^2 on Matrigel- or collagen IV-coated plates, with
collagen IV-coated plates yielding more highly ramified cells. Cells in B27-TIC media adhered
better to coated tissue culture plates.
604
605　Additional details about reagents and differentiation media compositions are provided in the
606　KEY RESOURCES TABLE and Supplementary Table 1.
607
608
609

Primary microglia isolation and culture
Neonatal brains were collected from P4 WT C57BL/6J mice and dissociated using the Miltenyi
gentleMACS Octo Dissociator with Heaters, brain dissociation kits (Miltenyi Biotec, 130-096-
427, 130-107-677, 130-092-628), and gentleMACS C Tubes (Miltenyi Biotec, 130-093-237),
following the manufacturer's instructions. Microglia were then purified using CD11b MicroBeads
(Miltenyi Biotec, 130-097-142), LS Columns (Miltenyi Biotec, 130-042-401), and the
QuadroMACS separator (Miltenyi Biotec, 130-091-051) following manufacturer's instructions.
Purified microglia were cultured in B27-TIC media. Cells were maintained under standard
conditions at 37°C in 5% CO_2 within a normoxic tissue culture incubator.

Bulk RNA sequencing
Approximately 500,000 -1,000,000 CX3CR1$^+$ immature macrophages or microglia-like cells
were sorted from three distinct mouse cell lines. Sorted cells were pelleted by centrifugation,
frozen, and sent to the Memorial Sloan Kettering Cancer Center (MSKCC) Integrated Genomics
Operation Core (IGO) for RNA extraction and quality control. The core performed RNA-polyA
sequencing, generating 30–40 million reads per sample.

Resulting FASTQ files were processed using Salmon[87] for transcript mapping and quantification
against the mm10 mouse genome. Downstream analyses were conducted in RStudio, using the
packages DESeq2, pcaExplorer, and pheatmap[88,89]. Statistical significance was assessed using
the Wald test and corrected with the Benjamini–Hochberg method.

Phagocytosis assays
Macrophages were first plated on collagen-IV-coated 8-well glass chamber slides (Nunc,
155409 for live imaging, 154534 for fixed imaging) and allowed to adhere overnight before
addition of phagocytosis targets.

For the synaptosome phagocytosis assay, synaptosomes were isolated from wild-type adult
mice using the Syn-PER Synaptic Protein Extraction Reagent (ThermoScientific, 87793)
following the manufacturer's protocol. Synaptosomes were stained with CypHer 5E (GE Life
Sciences, PA15401) via agitation, washing, and sonication. A BCA assay confirmed the staining
concentration (10 mM CypHer 5E/50 μL). Stained synaptosomes (20 μL) were added to
macrophages, and live imaging was performed for 45 minutes.

For the apoptotic neuron phagocytosis assay, apoptosis was induced in Neuro-2a cells (ATCC,
CCL-131) using 0.25 μM staurosporine (Cayman Chemical, 81590) for 12-15 hours. Apoptotic
neurons were stained with 1 μM CypHer 5E in serum-free HBSS for 45 minutes at 37°C, then
washed in DMEM for 20 minutes at 37°C. These neurons were resuspended in either immature
TRM media or B27-TIC and incubated with macrophages at a 1:1 ratio for 45 minutes during
live imaging.

For apoptotic Jurkat cells, apoptosis was induced with 150 mJ/cm^2 UVC irradiation followed by a
4-hour incubation at 37°C and 5% CO_2. Cells were stained with 1 μM CypHer 5E in serum-free
HBSS for 45 minutes at 37°C, washed, and incubated in serum-complete medium for 25
minutes. Macrophages were fed apoptotic Jurkat cells at a 1:1 ratio for 45 minutes. Post-
incubation, cells were washed, fixed with 4% PFA, and permeabilized with 0.3% Triton-X for 20
minutes. After blocking with 4% goat serum and 1% BSA, cells were incubated overnight at 4°C
with primary antibodies: goat anti-Iba1 (FUJIFILM Wako Chemicals, 1:500) and rabbit anti-
CD68 (ThermoScientific, 1:250). Secondary antibodies (1:500) were applied for 1 hour at room
temperature. Cells were washed, mounted with Prolong Gold, and imaged the next day.

661 For zymosan phagocytosis, AlexaFluor 594-conjugated zymosan particles (ThermoScientific,
662 Z23374) were added to macrophages at a 1:1 ratio, followed by the same fixation and staining
663 protocol used for Jurkat cell phagocytosis.
664
665 Live imaging of phagocytosis was conducted using time-lapse confocal microscopy on a Zeiss
666 Axio Observer.Z1-7 with a Z PIEZO stage, heating/gas-controlled insert, Zeiss 20x PlanApo
667 objective, LSM 980 with Airyscan.2 multiplex detector. Images were captured every 30 seconds.
668 Fixed cell imaging was also performed on the same system. All data was analyzed with Zen
669 Blue software (Zeiss) and FIJI.
670
Generation of cerebral cortical organoids and TRM transplantation
Cortical organoids devoid of macrophages were first generated from EpiSCs using a previously
described protocol[55]. Briefly, EpiSCs were seeded at 1,000 cells per microwell in an AggreWell
400 plate (34415, STEMCELL Technologies) using EB formation media consisting of 50 nM
chroman-1, 5 µM emricasan, 400 nM LDN-193189, 10 µM SB-431542, 1 µM cyclopamine, and
100 nM LGK-974 in N2B27 (B27 without vitamin A) media. After 24 hours, EBs were transferred
to the same media, but without chroman-1 and emricasan, and embedded in Matrigel. To
prepare the Matrigel mix, 66.7 µL of EBs in media were combined with 100 µL of Matrigel using
wide-bore tips, then plated in a 6-well plate without touching well edges and incubated at 37°C
for 30 minutes. Following incubation, 3 mL of warm media was added on top. On day 2, media
was replaced with plain N2B27 containing 1 µM cyclopamine and 1% KSR for 24 hours. On day
3, media was replaced with N2B27 supplemented with 1 µM cyclopamine, 3 µM CHIR99201,
and 1% KSR. On day 4, organoids were recovered from Matrigel using Cell Recovery Solution
(Corning, 354270). Wells were washed with PBS, and 2–3 mL of Cell Recovery Solution was
added, ensuring the Matrigel dome was lifted for efficient organoid retrieval. Plates were
incubated at 4°C for 30 minutes, followed by two washes in organoid media (N2B27 with 20%
KSR). Finally, cortical organoids were transferred to Petri dishes and maintained in N2B27 with
20% KSR while shaking at 65 rpm (Infors HT, Celltron benchtop shaker). Cortical organoid
media (N2B27 with 20% KSR) was changed every other day until day 10, when immature TRMs
were added.
691
692 To integrate immature TRMs, a single organoid was transferred to an ultra-low attachment U-
693 bottom 96-well plate. Day 20-28 immature TRMs (50,000 cells per organoid) were added in
694 cortical organoid media supplemented with 100 ng/mL IL-34 and 2 ng/mL TGF-β. The immature
695 TRMs were allowed to spontaneously migrate into the organoids overnight at 37°C, 5% CO_2.
696
The organoids were then pooled and transferred to 10 cm plates, which were placed on the
orbital shaker. Any remaining immature TRMs from the 96-well plate were added to the 10 cm
plate to maximize integration. These neuroimmune organoids were cultured for an additional 2-3
weeks in organoid media supplemented with IL-34 (100 ng/mL) and TGF-β (2 ng/mL), with
media changes twice a week. Organoids without immature TRMs were cultured in parallel as
controls.
703
Immunostaining of organoids
Organoids were collected in 1% BSA-coated tubes, washed twice with PBS-/- and fixed in 4%
paraformaldehyde for 45 minutes at 4°C. After fixation, samples were washed twice with PBS-/-
and permeabilized using 0.5% Triton X-100 for 15 minutes at 4°C. Blocking was carried out for
15 minutes in Organoid Washing Solution (OWS; adapted from Dekkers et al., 2019), composed
of 0.2% Triton X-100, 0.02% SDS, and 0.2% BSA in PBS-/-, at 4°C.
710

Organoids were incubated overnight at 4°C with primary antibodies diluted in OWS. The
following day, they were washed three times with OWS for 2 hours each under constant
shaking, followed by overnight incubation with secondary antibodies and DAPI (1 µM) in OWB
at 4°C. The organoids were then washed three times with OWB over a total of 6 hours while
shaking. Finally, organoids were mounted in DeepClear solution (CelExplorer Labs, DC-201)
and imaged using a Leica SP8 confocal microscope. Image analysis was performed with Imaris
software.
718
719   **Adoptive transfer of immature TRMs into mouse lung**
720   To deplete endogenous alveolar macrophages, 8-week-old CD45.1 congenic male and female
721   mice were anesthetized with isoflurane and administered 50 µL of clodronate liposomes (Fisher
722   Scientific, NC0337390) intratracheally. After 3 days, mice were anesthetized again and given
723   300,000 immature TRMs in 50 µL of sterile PBS intratracheally.
724
On days 3 and 7 post-transfer, mice were euthanized and perfused with a 21G needle
containing 40 mL of cold PBS-/- (Corning, 21-040-CM) and 5 mM EDTA (Invitrogen, 15575-
038). One lung per mouse was dissected, minced into small fragments, and digested for 30
minutes at 37°C in enzyme solution comprising PBS with 160 Wünsch U/mL collagenase D
(Sigma, 11088866001), 5% heat-inactivated FBS (Sigma, 12306C), and 10 mM HEPES (Gibco,
15630-080) (referred to as PFH buffer). The reaction was stopped with 10 mM EDTA for 5
minutes.
732
Digested lung tissue was further dissociated mechanically using a 70 µm cell strainer and an
18G needle with 3-mL syringe plunger in 6-well plates containing 4 mL of cold PFH buffer. The
resulting single-cell suspensions were transferred to 15-mL tubes and pelleted by centrifugation
at 1500 rpm for 5 minutes at 4°C. Pellets were resuspended in 38% isotonic Percoll (10% 10X
PBS, 90% Percoll; GE Healthcare, 17-0891-01) in PFH buffer to gradient separate leukocytes
by centrifugation at 2000 rpm for 30 minutes at room temperature with no brake.
739
740   To evaluate the differentiation of transferred immature TRMs into mature alveolar macrophages,
741   lung cell suspensions were incubated with Fc block for 10 minutes, followed by staining with the
742   following antibodies: CD45.2-AlexaFluor 700 (BioLegend, 1:400), CD11b-PE (Thermo Scientific,
743   1:800), CD11c-APC (BioLegend, 1:800), and SiglecF-BV421 (BioLegend, 1:400).
744
745   Detailed information on antibodies is provided in Supplementary Table 2.
746
Statistical analysis and reproducibility
Statistical tests were performed using GraphPad Prism 10 or Rstudio. Experiments with
statistical analyses were performed with a minimum of three independent differentiation
experiments (technical replicates) or biological replicates (different cell lines). For experiments
involving live mice, each data point represents results from a single mouse (lung adoptive
transfer) or from cells pooled from 4 different mice (*ex vivo* microglia cultures).

SUPPLEMENTARY INFORMATION

Key Resources Table
Supplementary Experimental Procedures: Step-by-step protocol for the directed differentiation of mouse EpiSCs into immature TRMs and microglia-like cells
Supplementary Table 1: 2D Differentiation media compositions
Supplementary Table 2: List of FACS antibodies
Supplementary Table 3: List of IF antibodies
Supplementary Video: Phagocytosis of CypHer 5E-labelled synaptosomes by immature TRMs

REFERENCES

1. Lazarov, T., Juarez-Carreno, S., Cox, N., and Geissmann, F. (2023). Physiology and diseases of tissue-resident macrophages. Nature *618*, 698-707. 10.1038/s41586-023-06002-x.
2. Mass, E., Nimmerjahn, F., Kierdorf, K., and Schlitzer, A. (2023). Tissue-specific macrophages: how they develop and choreograph tissue biology. Nat Rev Immunol *23*, 563-579. 10.1038/s41577-023-00848-y.
3. Tay, T.L., Savage, J.C., Hui, C.W., Bisht, K., and Tremblay, M.E. (2017). Microglia across the lifespan: from origin to function in brain development, plasticity and cognition. J Physiol *595*, 1929-1945. 10.1113/JP272134.
4. Paolicelli, R.C., and Ferretti, M.T. (2017). Function and Dysfunction of Microglia during Brain Development: Consequences for Synapses and Neural Circuits. Front Synaptic Neurosci *9*, 9. 10.3389/fnsyn.2017.00009.
5. Hussell, T., and Bell, T.J. (2014). Alveolar macrophages: plasticity in a tissue-specific context. Nat Rev Immunol *14*, 81-93. 10.1038/nri3600.
6. Lambrecht, B.N. (2006). Alveolar macrophage in the driver's seat. Immunity *24*, 366-368. 10.1016/j.immuni.2006.03.008.
7. Park, M.D., Silvin, A., Ginhoux, F., and Merad, M. (2022). Macrophages in health and disease. Cell *185*, 4259-4279. 10.1016/j.cell.2022.10.007.
8. Gao, C., Jiang, J., Tan, Y., and Chen, S. (2023). Microglia in neurodegenerative diseases: mechanism and potential therapeutic targets. Signal Transduct Target Ther *8*, 359. 10.1038/s41392-023-01588-0.
9. Shibuya, Y., Kumar, K.K., Mader, M.M., Yoo, Y., Ayala, L.A., Zhou, M., Mohr, M.A., Neumayer, G., Kumar, I., Yamamoto, R., et al. (2022). Treatment of a genetic brain disease by CNS-wide microglia replacement. Sci Transl Med *14*, eabl9945. 10.1126/scitranslmed.abl9945.
10. Lim, P.N., Cervantes, M.M., Pham, L.K., and Rothchild, A.C. (2021). Alveolar macrophages: novel therapeutic targets for respiratory diseases. Expert Rev Mol Med *23*, e18. 10.1017/erm.2021.21.
11. Walsh, A.E., and Lukens, J.R. (2025). Harnessing microglia-based cell therapies for the treatment of neurodegenerative diseases. Curr Opin Immunol *94*, 102552. 10.1016/j.coi.2025.102552.
12. Ginhoux, F., Greter, M., Leboeuf, M., Nandi, S., See, P., Gokhan, S., Mehler, M.F., Conway, S.J., Ng, L.G., Stanley, E.R., et al. (2010). Fate mapping analysis reveals that adult microglia derive from primitive macrophages. Science *330*, 841-845. 10.1126/science.1194637.

809   13.   Hoeffel, G., and Ginhoux, F. (2018). Fetal monocytes and the origins of tissue-
810   resident macrophages. Cell Immunol *330*, 5-15. 10.1016/j.cellimm.2018.01.001.
811   14.   Arnold, S.J., and Robertson, E.J. (2009). Making a commitment: cell lineage
812   allocation and axis patterning in the early mouse embryo. Nat Rev Mol Cell Biol
813   *10*, 91-103. 10.1038/nrm2618.
814   15.   Kinder, S.J., Tsang, T.E., Quinlan, G.A., Hadjantonakis, A.K., Nagy, A., and Tam,
815   P.P. (1999). The orderly allocation of mesodermal cells to the extraembryonic
816   structures and the anteroposterior axis during gastrulation of the mouse embryo.
817   Development *126*, 4691-4701. 10.1242/dev.126.21.4691.
818   16.   Palis, J., Chan, R.J., Koniski, A., Patel, R., Starr, M., and Yoder, M.C. (2001).
819   Spatial and temporal emergence of high proliferative potential hematopoietic
820   precursors during murine embryogenesis. Proc Natl Acad Sci U S A *98*, 4528-
821   4533. 10.1073/pnas.071002398.
822   17.   Palis, J., Robertson, S., Kennedy, M., Wall, C., and Keller, G. (1999).
823   Development of erythroid and myeloid progenitors in the yolk sac and embryo
824   proper of the mouse. Development *126*, 5073-5084. 10.1242/dev.126.22.5073.
825   18.   Hoeffel, G., and Ginhoux, F. (2015). Ontogeny of Tissue-Resident Macrophages.
826   Front Immunol *6*, 486. 10.3389/fimmu.2015.00486.
827   19.   Mass, E., Ballesteros, I., Farlik, M., Halbritter, F., Gunther, P., Crozet, L., Jacome-
828   Galarza, C.E., Handler, K., Klughammer, J., Kobayashi, Y., et al. (2016).
829   Specification of tissue-resident macrophages during organogenesis. Science
830   *353*. 10.1126/science.aaf4238.
831   20.   Tober, J., McGrath, K.E., and Palis, J. (2008). Primitive erythropoiesis and
832   megakaryopoiesis in the yolk sac are independent of c-myb. Blood *111*, 2636-
833   2639. 10.1182/blood-2007-11-124685.
834   21.   Li, Z., Liu, S., Xu, J., Zhang, X., Han, D., Liu, J., Xia, M., Yi, L., Shen, Q., Xu, S.,
835   et al. (2018). Adult Connective Tissue-Resident Mast Cells Originate from Late
836   Erythro-Myeloid Progenitors. Immunity *49*, 640-653 e645.
837   10.1016/j.immuni.2018.09.023.
838   22.   Gentek, R., Ghigo, C., Hoeffel, G., Bulle, M.J., Msallam, R., Gautier, G., Launay,
839   P., Chen, J., Ginhoux, F., and Bajenoff, M. (2018). Hemogenic Endothelial Fate
840   Mapping Reveals Dual Developmental Origin of Mast Cells. Immunity *48*, 1160-
841   1171 e1165. 10.1016/j.immuni.2018.04.025.
842   23.   Gomez Perdiguero, E., Klapproth, K., Schulz, C., Busch, K., Azzoni, E., Crozet,
843   L., Garner, H., Trouillet, C., de Bruijn, M.F., Geissmann, F., and Rodewald, H.R.
844   (2015). Tissue-resident macrophages originate from yolk-sac-derived erythro-
845   myeloid progenitors. Nature *518*, 547-551. 10.1038/nature13989.
846   24.   Hashimoto, D., Chow, A., Noizat, C., Teo, P., Beasley, M.B., Leboeuf, M., Becker,
847   C.D., See, P., Price, J., Lucas, D., et al. (2013). Tissue-resident macrophages
848   self-maintain locally throughout adult life with minimal contribution from
849   circulating monocytes. Immunity *38*, 792-804. 10.1016/j.immuni.2013.04.004.
850   25.   Alliot, F., Godin, I., and Pessac, B. (1999). Microglia derive from progenitors,
851   originating from the yolk sac, and which proliferate in the brain. Brain Res Dev
852   Brain Res *117*, 145-152. 10.1016/s0165-3806(99)00113-3.
853   26.   Jenkins, S.J., and Allen, J.E. (2021). The expanding world of tissue-resident
854   macrophages. Eur J Immunol *51*, 1882-1896. 10.1002/eji.202048881.

27. Hoeffel, G., Chen, J., Lavin, Y., Low, D., Almeida, F.F., See, P., Beaudin, A.E.,
Lum, J., Low, I., Forsberg, E.C., et al. (2015). C-Myb(+) erythro-myeloid
progenitor-derived fetal monocytes give rise to adult tissue-resident
macrophages. Immunity *42*, 665-678. 10.1016/j.immuni.2015.03.011.
28. Wiles, M.V., and Keller, G. (1991). Multiple hematopoietic lineages develop from
embryonic stem (ES) cells in culture. Development *111*, 259-267.
10.1242/dev.111.2.259.
29. Nakano, T., Kodama, H., and Honjo, T. (1994). Generation of
lymphohematopoietic cells from embryonic stem cells in culture. Science *265*,
1098-1101. 10.1126/science.8066449.
30. Kennedy, M., Firpo, M., Choi, K., Wall, C., Robertson, S., Kabrun, N., and Keller,
G. (1997). A common precursor for primitive erythropoiesis and definitive
haematopoiesis. Nature *386*, 488-493. 10.1038/386488a0.
31. Choi, K., Kennedy, M., Kazarov, A., Papadimitriou, J.C., and Keller, G. (1998). A
common precursor for hematopoietic and endothelial cells. Development *125*,
725-732. 10.1242/dev.125.4.725.
32. Claes, C., Van den Daele, J., and Verfaillie, C.M. (2018). Generating tissue-
resident macrophages from pluripotent stem cells: Lessons learned from
microglia. Cell Immunol *330*, 60-67. 10.1016/j.cellimm.2018.01.019.
33. Cakir, B., Kiral, F.R., and Park, I.H. (2022). Advanced in vitro models: Microglia in
action. Neuron *110*, 3444-3457. 10.1016/j.neuron.2022.10.004.
34. Atkins, M.H., Scarfo, R., McGrath, K.E., Yang, D., Palis, J., Ditadi, A., and Keller,
G.M. (2022). Modeling human yolk sac hematopoiesis with pluripotent stem cells.
J Exp Med *219*. 10.1084/jem.20211924.
35. Guttikonda, S.R., Sikkema, L., Tchieu, J., Saurat, N., Walsh, R.M., Harschnitz,
O., Ciceri, G., Sneeboer, M., Mazutis, L., Setty, M., et al. (2021). Fully defined
human pluripotent stem cell-derived microglia and tri-culture system model C3
production in Alzheimer's disease. Nat Neurosci *24*, 343-354. 10.1038/s41593-
020-00796-z.
36. Abud, E.M., Ramirez, R.N., Martinez, E.S., Healy, L.M., Nguyen, C.H.H.,
Newman, S.A., Yeromin, A.V., Scarfone, V.M., Marsh, S.E., Fimbres, C., et al.
(2017). iPSC-Derived Human Microglia-like Cells to Study Neurological
Diseases. Neuron *94*, 278-293 e279. 10.1016/j.neuron.2017.03.042.
37. Douvaras, P., Sun, B., Wang, M., Kruglikov, I., Lallos, G., Zimmer, M., Terrenoire,
C., Zhang, B., Gandy, S., Schadt, E., et al. (2017). Directed Differentiation of
Human Pluripotent Stem Cells to Microglia. Stem Cell Reports *8*, 1516-1524.
10.1016/j.stemcr.2017.04.023.
38. Choi, K.D., Vodyanik, M.A., Togarrati, P.P., Suknuntha, K., Kumar, A., Samarjeet,
F., Probasco, M.D., Tian, S., Stewart, R., Thomson, J.A., and Slukvin, II (2012).
Identification of the hemogenic endothelial progenitor and its direct precursor in
human pluripotent stem cell differentiation cultures. Cell Rep *2*, 553-567.
10.1016/j.celrep.2012.08.002.
39. Sturgeon, C.M., Ditadi, A., Awong, G., Kennedy, M., and Keller, G. (2014). Wnt
signaling controls the specification of definitive and primitive hematopoiesis from
human pluripotent stem cells. Nat Biotechnol *32*, 554-561. 10.1038/nbt.2915.

40.   Bredemeyer, A.L., Amrute, J.M., Koenig, A.L., Idol, R.A., He, L., Luff, S.A., Dege,
C., Leid, J.M., Schilling, J.D., Hinson, J.T., et al. (2022). Derivation of extra-
embryonic and intra-embryonic macrophage lineages from human pluripotent
stem cells. Development *149*. 10.1242/dev.200016.
41.   Shah, Z.H., Tian, L., Li, Z.X., Jin, L.W., Zhang, J.Y., Li, Z.L., Barr, T., Tang, H.J.,
Feng, M.Y., Caligiuri, M.A., and Yu, J.H. (2024). Human anti-PSCA CAR
macrophages possess potent antitumor activity against pancreatic cancer. Cell
Stem Cell *31*. 10.1016/j.stem.2024.03.018.
42.   Kattman, S.J., Witty, A.D., Gagliardi, M., Dubois, N.C., Niapour, M., Hotta, A.,
Ellis, J., and Keller, G. (2011). Stage-specific optimization of activin/nodal and
BMP signaling promotes cardiac differentiation of mouse and human pluripotent
stem cell lines. Cell Stem Cell *8*, 228-240. 10.1016/j.stem.2010.12.008.
43.   Pandya, H., Shen, M.J., Ichikawa, D.M., Sedlock, A.B., Choi, Y., Johnson, K.R.,
Kim, G., Brown, M.A., Elkahloun, A.G., Maric, D., et al. (2017). Differentiation of
human and murine induced pluripotent stem cells to microglia-like cells. Nat
Neurosci *20*, 753-759. 10.1038/nn.4534.
44.   Harland, L.T.G., Simon, C.S., Senft, A.D., Costello, I., Greder, L., Imaz-
Rosshandler, I., Gottgens, B., Marioni, J.C., Bikoff, E.K., Porcher, C., et al.
(2021). The T-box transcription factor Eomesodermin governs haemogenic
competence of yolk sac mesodermal progenitors. Nat Cell Biol *23*, 61-74.
10.1038/s41556-020-00611-8.
45.   Du, P., and Wu, J. (2024). Hallmarks of totipotent and pluripotent stem cell states.
Cell Stem Cell *31*, 312-333. 10.1016/j.stem.2024.01.009.
46.   Morgani, S., Nichols, J., and Hadjantonakis, A.K. (2017). The many faces of
Pluripotency: in vitro adaptations of a continuum of in vivo states. BMC Dev Biol
17, 7. 10.1186/s12861-017-0150-4.
47.   Weinberger, L., Ayyash, M., Novershtern, N., and Hanna, J.H. (2016). Dynamic
stem cell states: naive to primed pluripotency in rodents and humans. Nat Rev
Mol Cell Biol *17*, 155-169. 10.1038/nrm.2015.28.
48.   Smith, A. (2017). Formative pluripotency: the executive phase in a developmental
continuum. Development *144*, 365-373. 10.1242/dev.142679.
49.   Kalkan, T., Olova, N., Roode, M., Mulas, C., Lee, H.J., Nett, I., Marks, H., Walker,
R., Stunnenberg, H.G., Lilley, K.S., et al. (2017). Tracking the embryonic stem
cell transition from ground state pluripotency. Development *144*, 1221-1234.
10.1242/dev.142711.
50.   Medina-Cano, D., Corrigan, E.K., Glenn, R.A., Islam, M.T., Lin, Y., Kim, J., Cho,
H., and Vierbuchen, T. (2022). Rapid and robust directed differentiation of mouse
epiblast stem cells into definitive endoderm and forebrain organoids.
Development *149*. 10.1242/dev.200561.
51.   Edri, S., Hayward, P., Baillie-Johnson, P., Steventon, B.J., and Martinez Arias, A.
(2019). An epiblast stem cell-derived multipotent progenitor population for axial
extension. Development *146*. 10.1242/dev.168187.
52.   Najm, F.J., Zaremba, A., Caprariello, A.V., Nayak, S., Freundt, E.C., Scacheri,
P.C., Miller, R.H., and Tesar, P.J. (2011). Rapid and robust generation of
functional oligodendrocyte progenitor cells from epiblast stem cells. Nat Methods
8, 957-962. 10.1038/nmeth.1712.

53. Rossant, J., and Tam, P.P.L. (2017). New Insights into Early Human
Development: Lessons for Stem Cell Derivation and Differentiation. Cell Stem
Cell *20*, 18-28. 10.1016/j.stem.2016.12.004.
54. Tesar, P.J., Chenoweth, J.G., Brook, F.A., Davies, T.J., Evans, E.P., Mack, D.L.,
Gardner, R.L., and McKay, R.D. (2007). New cell lines from mouse epiblast share
defining features with human embryonic stem cells. Nature *448*, 196-199.
10.1038/nature05972.
55. Medina-Cano, D., Islam, M.T., Petrova, V., Dixit, S., Balic, Z., Yang, M.G.,
Stadtfeld, M., Wong, E.S., and Vierbuchen, T. (2024). A MOUSE ORGANOID
PLATFORM FOR MODELING CEREBRAL CORTEX DEVELOPMENT AND CIS-
REGULATORY EVOLUTION IN VITRO. bioRxiv, 2024.2009.2030.615887.
10.1101/2024.09.30.615887.
56. Parameswaran, M., and Tam, P.P. (1995). Regionalisation of cell fate and
morphogenetic movement of the mesoderm during mouse gastrulation. Dev
Genet *17*, 16-28. 10.1002/dvg.1020170104.
57. Morgani, S.M., Metzger, J.J., Nichols, J., Siggia, E.D., and Hadjantonakis, A.K.
(2018). Micropattern differentiation of mouse pluripotent stem cells recapitulates
embryo regionalized cell fate patterning. Elife *7*. 10.7554/eLife.32839.
58. Frame, J.M., Fegan, K.H., Conway, S.J., McGrath, K.E., and Palis, J. (2016).
Definitive Hematopoiesis in the Yolk Sac Emerges from Wnt-Responsive
Hemogenic Endothelium Independently of Circulation and Arterial Identity. Stem
Cells *34*, 431-444. 10.1002/stem.2213.
59. Yzaguirre, A.D., and Speck, N.A. (2016). Insights into blood cell formation from
hemogenic endothelium in lesser-known anatomic sites. Dev Dyn *245*, 1011-
1028. 10.1002/dvdy.24430.
60. Goldie, L.C., Lucitti, J.L., Dickinson, M.E., and Hirschi, K.K. (2008). Cell signaling
directing the formation and function of hemogenic endothelium during murine
embryogenesis. Blood *112*, 3194-3204. 10.1182/blood-2008-02-139055.
61. Lange, L., Morgan, M., and Schambach, A. (2021). The hemogenic endothelium:
a critical source for the generation of PSC-derived hematopoietic stem and
progenitor cells. Cell Mol Life Sci *78*, 4143-4160. 10.1007/s00018-021-03777-y.
62. Biben, C., Weber, T.S., Potts, K.S., Choi, J., Miles, D.C., Carmagnac, A.,
Sargeant, T., de Graaf, C.A., Fennell, K.A., Farley, A., et al. (2023). In vivo clonal
tracking reveals evidence of haemangioblast and haematomesoblast contribution
to yolk sac haematopoiesis. Nat Commun *14*, 41. 10.1038/s41467-022-35744-x.
63. Lugus, J.J., Park, C., Ma, Y.D., and Choi, K. (2009). Both primitive and definitive
blood cells are derived from Flk-1+ mesoderm. Blood *113*, 563-566.
10.1182/blood-2008-06-162750.
64. Shalaby, F., Ho, J., Stanford, W.L., Fischer, K.D., Schuh, A.C., Schwartz, L.,
Bernstein, A., and Rossant, J. (1997). A requirement for Flk1 in primitive and
definitive hematopoiesis and vasculogenesis. Cell *89*, 981-990. 10.1016/s0092-
8674(00)80283-4.
65. Lazarov, T., Loyher, P.-L., Yang, H., Choo, Z.-N., Deng, Z., Nowotschin, S., Kuo,
Y.-Y., Zhou, T., Alberdi-Gonzalez, A., Stumm, R., et al. (2024). Identification of a
mesodermal progenitor for the pro-definitive angio-hematopoietic lineage.
bioRxiv, 2024.2008.2024.609533. 10.1101/2024.08.24.609533.

66.　Endoh, M., and Niwa, H. (2022). Stepwise pluripotency transitions in mouse stem
cells. EMBO Rep *23*, e55010. 10.15252/embr.202255010.
67.　McGrath, K.E., Frame, J.M., Fegan, K.H., Bowen, J.R., Conway, S.J.,
Catherman, S.C., Kingsley, P.D., Koniski, A.D., and Palis, J. (2015). Distinct
Sources of Hematopoietic Progenitors Emerge before HSCs and Provide
Functional Blood Cells in the Mammalian Embryo. Cell Rep *11*, 1892-1904.
10.1016/j.celrep.2015.05.036.
68.　Chen, M.J., Yokomizo, T., Zeigler, B.M., Dzierzak, E., and Speck, N.A. (2009).
Runx1 is required for the endothelial to haematopoietic cell transition but not
thereafter. Nature *457*, 887-891. 10.1038/nature07619.
69.　Bennett, M.L., Bennett, F.C., Liddelow, S.A., Ajami, B., Zamanian, J.L., Fernhoff,
N.B., Mulinyawe, S.B., Bohlen, C.J., Adil, A., Tucker, A., et al. (2016). New tools
for studying microglia in the mouse and human CNS. Proc Natl Acad Sci U S A
113, E1738-1746. 10.1073/pnas.1525528113.
70.　Butovsky, O., Jedrychowski, M.P., Moore, C.S., Cialic, R., Lanser, A.J., Gabriely,
G., Koeglsperger, T., Dake, B., Wu, P.M., Doykan, C.E., et al. (2014).
Identification of a unique TGF-beta-dependent molecular and functional signature
in microglia. Nat Neurosci *17*, 131-143. 10.1038/nn.3599.
71.　Ishimaru, M., Yusuke, N., Tsukimoto, M., Harada, H., Takenouchi, T., Kitani, H.,
and Kojima, S. (2014). Purinergic signaling via P2Y receptors up-mediates IL-6
production by liver macrophages/Kupffer cells. J Toxicol Sci *39*, 413-423.
10.2131/jts.39.413.
72.　Bohlen, C.J., Bennett, F.C., Tucker, A.F., Collins, H.Y., Mulinyawe, S.B., and
Barres, B.A. (2017). Diverse Requirements for Microglial Survival, Specification,
and Function Revealed by Defined-Medium Cultures. Neuron *94*, 759-773 e758.
10.1016/j.neuron.2017.04.043.
73.　Gosselin, D., Link, V.M., Romanoski, C.E., Fonseca, G.J., Eichenfield, D.Z.,
Spann, N.J., Stender, J.D., Chun, H.B., Garner, H., Geissmann, F., and Glass,
C.K. (2014). Environment drives selection and function of enhancers controlling
tissue-specific macrophage identities. Cell *159*, 1327-1340.
10.1016/j.cell.2014.11.023.
74.　Okabe, Y., and Medzhitov, R. (2014). Tissue-specific signals control reversible
program of localization and functional polarization of macrophages. Cell *157*,
832-844. 10.1016/j.cell.2014.04.016.
75.　Ng, E.S., Azzola, L., Bruveris, F.F., Calvanese, V., Phipson, B., Vlahos, K., Hirst,
C., Jokubaitis, V.J., Yu, Q.C., Maksimovic, J., et al. (2016). Differentiation of
human embryonic stem cells to HOXA(+) hemogenic vasculature that resembles
the aorta-gonad-mesonephros. Nat Biotechnol *34*, 1168-1179. 10.1038/nbt.3702.
76.　Hills, D., Gribi, R., Ure, J., Buza-Vidas, N., Luc, S., Jacobsen, S.E., and
Medvinsky, A. (2011). Hoxb4-YFP reporter mouse model: a novel tool for tracking
HSC development and studying the role of Hoxb4 in hematopoiesis. Blood *117*,
3521-3528. 10.1182/blood-2009-12-253989.
77.　Steens, J., and Klein, D. (2022). HOX genes in stem cells: Maintaining cellular
identity and regulation of differentiation. Front Cell Dev Biol *10*, 1002909.
10.3389/fcell.2022.1002909.

78.   Hammond, T.R., Dufort, C., Dissing-Olesen, L., Giera, S., Young, A., Wysoker, A.,
1038   Walker, A.J., Gergits, F., Segel, M., Nemesh, J., et al. (2019). Single-Cell RNA
1039   Sequencing of Microglia throughout the Mouse Lifespan and in the Injured Brain
1040   Reveals Complex Cell-State Changes. Immunity *50*, 253-271 e256.
1041   10.1016/j.immuni.2018.11.004.
1042   79.   Thepen, T., Van Rooijen, N., and Kraal, G. (1989). Alveolar macrophage
1043   elimination in vivo is associated with an increase in pulmonary immune response
1044   in mice. J Exp Med *170*, 499-509. 10.1084/jem.170.2.499.
1045   80.   Ormel, P.R., Vieira de Sa, R., van Bodegraven, E.J., Karst, H., Harschnitz, O.,
1046   Sneeboer, M.A.M., Johansen, L.E., van Dijk, R.E., Scheefhals, N., Berdenis van
1047   Berlekom, A., et al. (2018). Microglia innately develop within cerebral organoids.
1048   Nat Commun *9*, 4167. 10.1038/s41467-018-06684-2.
1049   81.   Sabate-Soler, S., Nickels, S.L., Saraiva, C., Berger, E., Dubonyte, U., Barmpa,
1050   K., Lan, Y.J., Kouno, T., Jarazo, J., Robertson, G., et al. (2022). Microglia
1051   integration into human midbrain organoids leads to increased neuronal
1052   maturation and functionality. Glia *70*, 1267-1288. 10.1002/glia.24167.
1053   82.   Zhang, W., Jiang, J., Xu, Z., Yan, H., Tang, B., Liu, C., Chen, C., and Meng, Q.
1054   (2023). Microglia-containing human brain organoids for the study of brain
1055   development and pathology. Mol Psychiatry *28*, 96-107. 10.1038/s41380-022-
1056   01892-1.
1057   83.   Kirkiles-Smith, N.C., Harding, M.J., Shepherd, B.R., Fader, S.A., Yi, T., Wang, Y.,
1058   McNiff, J.M., Snyder, E.L., Lorber, M.I., Tellides, G., and Pober, J.S. (2009).
1059   Development of a humanized mouse model to study the role of macrophages in
1060   allograft injury. Transplantation *87*, 189-197. 10.1097/TP.0b013e318192e05d.
1061   84.   Yin, L., Wang, X.J., Chen, D.X., Liu, X.N., and Wang, X.J. (2020). Humanized
1062   mouse model: a review on preclinical applications for cancer immunotherapy. Am
1063   J Cancer Res *10*, 4568-4584.
1064   85.   Dolan, M.J., Therrien, M., Jereb, S., Kamath, T., Gazestani, V., Atkeson, T.,
1065   Marsh, S.E., Goeva, A., Lojek, N.M., Murphy, S., et al. (2023). Exposure of iPSC-
1066   derived human microglia to brain substrates enables the generation and
1067   manipulation of diverse transcriptional states in vitro. Nat Immunol *24*, 1382-
1068   1390. 10.1038/s41590-023-01558-2.
1069   86.   Glenn, R.A., Do, S.C., Guruvayurappan, K., Corrigan, E.K., Santini, L., Medina-
1070   Cano, D., Singer, S., Cho, H., Liu, J., Broman, K., et al. (2024). A Pluripotent
1071   Stem Cell Platform for in Vitro Systems Genetics Studies of Mouse Development.
1072   bioRxiv. 10.1101/2024.06.06.597758.
1073   87.   Patro, R., Duggal, G., Love, M.I., Irizarry, R.A., and Kingsford, C. (2017). Salmon
1074   provides fast and bias-aware quantification of transcript expression. Nat Methods
1075   *14*, 417-419. 10.1038/nmeth.4197.
1076   88.   Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold
1077   change and dispersion for RNA-seq data with DESeq2. Genome Biol *15*, 550.
1078   10.1186/s13059-014-0550-8.
1079   89.   Marini, F., and Binder, H. (2019). pcaExplorer: an R/Bioconductor package for
1080   interacting with RNA-seq principal components. BMC Bioinformatics *20*, 331.
1081   10.1186/s12859-019-2879-1.
1082