Controlling the synchronization and symmetry breaking of coupled bacterial pili on active biofilm carpets

Authors

Baha Altın[,1], Yusuf Ilker Yaman[1,2], Enes Talha Günay[1] Alp Ünlü[1], Yiğithan Gediz[1], Neslihan Gedik[3], Bora Karadaş[,1], Mustafa Başaran[1,2], Coşkun Kocabaş[4], Şahin Özdemir[5,6], and Aşkın Kocabaş[1,7]†

[1]Department of Physics, Koç University, Sarıyer, 34450 Istanbul, Türkiye

[2]Current address: Sciences and Engineering Program, Harvard University, Cambridge 02138, MA

[3]Department of Physics, Undergraduate Program, Boğaziçi University, 34342 Bebek, Istanbul Türkiye

[4]Department of Materials, University of Manchester, Manchester, M13 9PL, UK

[5]Department of Engineering Science and Mechanics, The Pennsylvania State University, University Park, Pennsylvania 16802, USA

[6]Department of Electrical and Computer Engineering, Saint Louis University, Saint Louis, Missouri 63103, USA

[7]Koç University Surface Science and Technology Center, Koç University, Sarıyer, 34450 Istanbul, Türkiye

†Corresponding author: **akocabas@ku.edu.tr**

Abstract

In the low Reynolds number regime, active biological systems utilize nonreciprocal cyclic activities to achieve motility, as seen in the spinning of bacterial flagella and the beating of cilia. Coupling among these active mechanical components leads to synchronization, and emergence of metachronal waves. Here, we report that biofilms of *Pseudomonas nitroreducens* form active carpets surfaces textured with various topological defects generating large-scale propagating waves. On these active structures, nonreciprocally coupled extension and retraction activities of bacterial pili drives the collective oscillations. Surprisingly, this collective behavior exhibits broken left-right symmetry. We discover that this symmetry-breaking process is primarily driven by an aging-related frequency gradient across the biofilm. Leveraging these insights, we further demonstrate the ability to control the collective dynamics of these waves, including symmetry breaking, transitions from spiral waves into target and propagating plane waves by manipulating the biofilm properties. Overall, our findings illuminate the fundamental role of nonreciprocally interacting active components in regulating synchronization, collective dynamics, and symmetry-breaking phenomena in biological systems.

Introduction

At low Reynolds numbers regime, overcoming reversibility constrain to gain motility, is a challenging task which requires innovative solutions. Microorganisms utilize a variety of specialized active mechanical components to solve these difficulties by performing nonreciprocal motion[1-4]. Typical examples include the rotating helical flagella and beating elastic cilia. These active mechanical systems have the ability to execute cyclic motions, thereby breaking time-reversibility constraints[5]. In addition, due to the coupling among these active microstructures, complex synchronization phenomena also known as metachronal waves could emerge[6-8]. From these perspective coupling, synchronization and collective behaviors are intrinsically related and ubiquitous in nature. Such widespread dynamical processes are also biologically essential, as seen in cilia-driven mucus transport in lungs[9], nodal flow in developing embryos[10] , central motor generators[11] and spinal cord development[12]. These intricate synchronization behaviors share the same underlying dynamical principles that surprisingly align with non-Hermitian physics of active systems[13-16]. All these active systems are capable of injecting and transferring energy across the system by breaking certain symmetries.

Recent advances, especially those in the field of active matter physics[17-22], have illuminated critical non-Hermitian features, such as chiral states where PT symmetry is broken. In this broken mode system could sustain the phase difference between fundamental fields or the modes which further trigger the directional energy transfer between the active compartments and generate travelling waves. Notably this non-Hermitian process is particularly driven by nonreciprocity in the coupled

mechanical system[8,17,23]. Although non-Hermiticity has its roots in quantum mechanics[14,24] and the

fundamental principles have been broadly studied in photonics[25], electronics, acoustics,

optomechanics[26,27], superconducting qubits[28], trapped ions[29,30], single-spin systems[31], and in light-

matter interactions[32]. This process is often achieved not only using strong nonlinearities, but also

utilizing nonreciprocal interactions and coupling[33-35]. On the other hand, nonreciprocity is a very

common feature in biological systems. Such as hydrodynamic interactions[36] or prey predator[37]

relations could easily break action reaction symmetry and results in exotic collective behaviors.

Extending the basic concepts from non-Hermitian physics to biological domain holds promise for

shedding light on the fundamental principles of symmetry breaking in active and, more crucially,

in complex biological systems.

Various biological models with specific active micromechanical components have been studied.

The most notable biological platforms are ciliated epithelial cell[38], bacterial carpet[39,40], starfish

embryo[41] social amoeba[42-44] and walking placozoa[45]. Addition to flagellum or cilia, bacteria could

also use nano filaments known as type IV pili to gain motility on surfaces where the rotating

flagella do not work effectively. Recent studies also highlighted the importance of pili controlling

the collective behaviors of bacteria[46-49] and also large-scale oscillations in the form of biofilm.

During our recent experiments, we noted striking dynamic patterns and unidirectionally

propagating waves on the surface of *Pseudomonas Nitroreducens (PN)* bacterial biofilms. These

unexpected collective activities ranging from spiral to planar wave formations, interestingly

aligned with recent predictions of active matter physics[23,50]. These activities are reminiscent of

metachronal waves observed in various biological systems. We also observed that these collective

activities are driven by the coupled pili activity of the bacteria. Recently, large-scale pili-driven

activities and spiral waves were also identified in *Pseudomonas aeruginosa*[49]. The main distinction

of our observed metachronal wave lies in its localized behavior on the surface and broken left-

right symmetry across the biofilm, which we refer to as the active biofilm carpet. In this study, we

aim to provide a comprehensive experimental approach including biological, physical and

theoretical tools to be able to understand and control these collective behaviors and also the

symmetry breaking process of these biological systems.

Bacterial biofilms develop into multicellular communities characterized by interdependent

biological structures. Understanding the dynamics of these dense biological formations is vital,

given that their collective actions can amplify their pathogenic potential. The collective movement

of pili and its variability among strains offers a crucial foundation for understanding the influence

of pili activities on pathogenicity. Moreover, pili is also a primary target protein for next-

generation medications and antibacterial therapies. In essence, comprehending the effect of pili

dynamics on the collective behavior of biofilms can greatly influence human health by preventing

severe infections.

Figure 1. Emergence of unidirectionally propagating waves on bacterial biofilms. (a-b) Optical imaging shows spiral, planar (a), and radially shrinking waves (b) propagating unidirectionally on biofilm surfaces. Left-right or radial symmetry is generally broken on growing biofilm surfaces. Scale bar 250 um. (c) Time response of optical scattering signal indicating the firing state of coupled pili dynamics. (d) Schematic representation of the coupling mechanism of pili on biofilm surfaces, modeled as an active solid. The oscillatory extension and retraction of pili act as active units. (e) Collective behavior of elastically coupled active biofilm surfaces characterized by local displacement (U) and pili polarization(P). Propagating waves remain localized on the surface and travel toward the direction of the sharp rising edge. f) Optical images of the leading edge of the growing biofilm with fingering instabilities. PN and PA14 show strong fingers compared to PAO1.

Results

Emergence of metachronal waves on active biofilm carpets

Bacterial biofilms are commonly used in various experiments particularly to investigate host-pathogen interactions. During our recent screen of biofilm library for nematode *C.elegans[51]*, we observed that *Pseudomonas Nitroreducens* (PN, Materials and Methods) bacterial biofilms exhibit rhythmic activities approximately 10 hours post-inoculation (Figure1a-b). In their firing phase, faint propagating pairs of spiral wave patterns emerged on the biofilm's surface. These waves have a periodicity of around 2-7 minutes (Figure1c). We also determined that the visibility of these waves improves dramatically with oblique contrast and polarized light-based microscopy (Figure1a-b, Supplementary Video1-2). Later on, these waves were very dynamic and converged to various combinations of spiral, target and planar waves. The most striking feature of these waves is their broken left-right symmetries. They, often unidirectionally propagate from right to left (edge to the center) or radially in inward direction across the colony (Figure1a-b). Furthermore, the contrast of the wave's switches between the dark to bright sharp lines depending on the direction of propagation. This suggest that the front kink of these waves scatters the light asymmetrically (Supplementary Figure 1). We observed that these waves predominantly propagate on the biofilm's surface rather than its bulk region (Figure1e). This critical feature is particularly revealed by the nature of our optical imaging systems which is very sensitive to the surface topography. To validate this observation, we employed cytoplasmic GFP for imaging. Yet, no waves were detected in the green fluorescence channel (Supplementary Figure2). Similarly, when we labeled extracellular DNA with a dye, the waves remained unobserved in the fluorescence signal (Supplementary Figure2). Moreover, close-up imaging (100X) highlighted the bacterial displacement together with clear optical dark or bright contrast on the surface (Supplementary Video3). Imaging growing

biofilm starting from a single bacterium clarifies that oscillations first starts from dense regions

and gradually propagate through the entire colony (Supplementary Video4). Further motivated by

recent studies[48,49], we also tested the *Pseudomonas aeruginosa* strain PA14 under the same

conditions. Similarly, PA14 exhibited weak but comparable wave patterns, though within a shorter

time window (Supplementary Figure3).

To delve deeper into these phenomena, our next focus was on the necessary media, given that the

original experiments were conducted on *C. elegans* using Nematode Growth Media (NGM) plates

(see Materials and Methods). Interestingly, we found that these activities were exclusive to NGM.

Substituting NGM with a standard LB plate did not reproduce the same dynamics; instead, the

bacteria grow rapidly. Yet, upon omitting yeast extract and replacing bactotryptone with

bactopeptone, a major nutritional component, from the LB media PN resumed its activity

(Supplementary Figure5). These findings underscore the significance of nutrient restriction as a

primary factor. Remarkably, these observations are unexpected, but become evident due the use

of NGM plate, which is not common in microbiology experiments.

Although PA14 generates similar waves we observed that PAO1, a close relative of PA14, failed

to produce waves on both NGM and LB plates (Supplementary Figure4). Worth noting, PAO1 is

also an important human pathogen causing *Cystic Fibrosis* (CF). Finally, through this

comprehensive screening effort, we successfully identified a group of bacteria, offering the

potential to examine various factors, from genetics to physical domain, influencing the emergence

of these spiral waves. Both PA01 and PA14 possess robust genetic toolkits, while PN strain display

a clear and prolonged active carpet state for detailed physical investigations. The other critical

difference between all these biofilms is the finger formation around the leading edge. Unlike

PAO1, PN and PA14 form very strong instabilities leading unstable extensions (Figure1f).

We then shifted our attention to the genetic distinctions between PN, PA14 and PAO1. Initially,

we first explored the contribution of bacterial pili and flagella as an active mechanical structure.

We found that PA14:ΔPilB and PA14:ΔPilA mutations eliminated the waves but not PA14:ΔFilK

(Supplementary Figure3). This suggests that activity is exclusively originating from pili dynamics

and the original dynamics is similar to the recent work[49], but activities are localized to the surface.

We then explored the potential of hyper piliation, which might amplify the coupling of pili on the

biofilm surfaces. Yet, the PAO1 ΔPilH mutant, known for its polarized and hyper piliation

condition[46,47], did not manifest the waves but provided only small fluctuations on the surface

(Supplementary Figure4). Then, we pinpointed that a significant variation was in the pili subunit

groups[52]. While PA14 possesses group 3 Pilin, PAO1 has group 2 Pilin. It's essential to highlight

that all these bacteria have type IV pili, though they might be categorized into separate groups.

Currently, our hypothesis posits that diverse pili types might exhibit varied physical characteristics

such as elasticity or rigidity. Our understanding of PN at the genetic level remains limited but it

shares strong similarity with PA. Nonetheless, the various genome sequences of PN are accessible.

Our analysis of their pili related genomic region revealed a resemblance to the group 4 and 5 pili

(Supplementary Figure 6). All these firing pili groups 3 and 4, 5 have additional accessory genes

in the genome and they may contribute to the expression or folding of these pilins. Together with

the unstable edge fingering (Figure1f), the comparison of PN and PA strains support the idea that

group of pili and particularly pili expression play a critical role in driving waves on these bacterial

biofilms.

Figure 2. Numerical modeling of coupled pili dynamics as an active carpet. *(a-c) Numerical simulation results based on the nonreciprocal Kuramoto phase-field model. Nonreciprocal coupling term (α) among bacteria drives the emergence of spiral waves, while the excitability term (b) of the mechanical system induces pulsatile behavior. (d) Numerical simulations of local oscillation frequencies under varying conditions (α = 0, α = 0.2π, and b = 0.1π). (e) Experimental measurements of pulses on biofilm surfaces confirming pulsatile responses. (f-g) Numerical simulations of elastically coupled biofilm structures using the active solid model. Active solids exhibit large-scale spiral and planar wave formations. Displacement (U) and pili polarization (P) fields highlight the essential phase difference necessary for wave propagation and limit-cycle oscillations.*

Modelling of propagating spiral waves

To get more intuition about the dynamics of the waves and emergence of the broken symmetry

across the biofilm, we next focused on the mathematical modelling. From an active matter

perspective, spiral waves can emerge in non-equilibrium and excitable media[53-55] as well as in

coupled oscillatory systems with delays[56]. Despite these systems exhibiting similar wave patterns,

the internal mechanisms that trigger the waves vary significantly. Recent studies particularly on

the concept of nonreciprocity provides comprehensive approach[23,50]. It is important to note that

coupling between active mechanical units could introduce nonreciprocity into the system due to

local symmetry-breaking processes. Inspired from the previous studies on the dynamics of

cilia[17,57], and recent robotic system[58] we proposed that coupled active pili system could also show

similar limit cycles oscillations defined by bacterial displacement (**U**) and the polarization of the

bacterial pili (**P**) (Figure1e). To study these phenomena, we first utilized a minimal

phenomenological phase field model that includes a nonreciprocal coupling term. The oscillatory

dynamics of the model are driven by oscillatory extension and retraction processes. The Kuramoto

based model[23,56,59], which effectively captures the phase dynamics of coupled oscillators, is

extended here by incorporating a delay term (α) that disrupts the odd symmetry—representing

action and reaction symmetry between interacting oscillators—and enriches the system's dynamic

repertoire. This modified model is described by the equation:

208
$$\frac{d\emptyset_i}{dt} = w_0 - b\,sin(\emptyset_i) - K\sum_j sin(\emptyset_i - \emptyset_j - \alpha)$$

Here, $\emptyset$ represents the local phase of individual pili oscillations (The phase of parameter space

defined by **U** and **P**, Figure 1e). We assume interactions only among nearest neighbors, where w_0

represents the uniform intrinsic oscillation frequency of an isolated oscillator. The term α

specifically break odd interaction symmetry between pili, facilitating the formation of travelling

waves. Note that traveling waves indicate this broken PT symmetry between these fields. It's

important to emphasize that the detailed biophysical mechanisms behind this nonreciprocity

remain unclear. Several possibilities, such as force relaxation dynamics or hydrodynamics

interactions, could also result in similar nonreciprocal behavior[60]. Our phenomenological model

accurately captures the principal features of these phenomena. While reciprocally coupled

oscillators ($\alpha = 0$) tend to achieve a globally synchronized state (Figure 2a), nonreciprocity favors

the formation of traveling waves (Figure 2b-c, Supplementary Video5). The excitability term (**b**)

shapes the pulsative nature of the waves (illustrated in Figure 2d-e) by amplifying the effects of

specific phases within the coupling mechanism[56,61,62].

222
$$\partial_t U = \pi P + F_e$$

223
$$\partial_t P = (P \times \partial_t U) \times P + \frac{1 - P^2}{2}\partial_t U - D_r P$$

Next, we turned our focus to continuum simulations to capture the essential properties of these

dynamic processes (Supplementary Video6). The recently developed active solid model provides

a broad and robust framework for studying collective behaviors of interacting active components,

spanning from robotic systems to human crowds[58,63]. The core idea of this model is based on two

nonreciprocally coupled fields: displacement (**U**), representing bacterial displacement within the

biofilm lattice, and polarization (**P**), representing the orientation of pili, which exert active forces

whose direction can be modulated by the displacement field (Figure1d). The dynamics of the active

biofilm lattice can be described by coupled equations, where $\mathbf{F_e}$ denotes elastic force and D_r

represents polarization relaxation. The underlying rationale of these equations is that retracted pili

exert active forces on bacteria, which are elastically coupled to the biofilm structure. Local biofilm

displacement can guide the growth direction of pili extension, influenced by local deformation.

Importantly, since pili are anchored to the substrate, their orientation dynamically adjusts based

on extension processes. Our simulations demonstrate that the active solid model successfully

captures critical dynamical phenomena, such as the formation of spiral and planar waves (Figure

2 f-g). As anticipated, fields $\mathbf{U}$ and $\mathbf{P}$ develop a characteristic phase difference, indicative of limit

cycle oscillations. Moreover, analysis of vector configurations around defect cores reveals

directional reversals of displacement $\mathbf{U}$ on opposite sides of defects, highlighting the fundamental

mechanism underlying defect formation, where $\mathbf{U}$ diminishes, and the phase ($\emptyset$) becomes

undefined.

Figure 3. Controlling transitions between spiral, target, and planar waves. *(a-c) Numerical simulations illustrating transitions from spiral waves to target and planar waves. Pairs of spiral waves merge, forming topologically neutral target waves, which eventually give rise to planar waves dominating the biofilm surface. (d-f) Optical imaging of biofilm surfaces demonstrating similar transitions between spiral, target, and planar waves experimentally triggered by adding a water droplet. Red arrows indicate wave propagation direction. (g-i) Controlled recovery of spiral waves achieved by heating biofilm surfaces, removing excess moisture, and facilitating re-emergence of spiral waves.*

Controlling the transition between the waveforms on active biofilm carpet

Extensive parameter testing has revealed that our models not only capture the emergence of spiral

waves but also predicts transitions into target and plane wave solutions which we commonly

observe in grooving biofilms. During the transition to target waves, two oppositely spinning spiral

waves with topological charges of +1 and -1 merge, forming a symmetrically expanding target

wave. Subsequently, fast oscillating plane waves progressively dominate the entire simulation

domain (Figure3 a-c, Supplementary Video5).

Motivated by our simulation results, we hypothesize that our active carpet system could also

facilitate similar transitions between different forms of traveling waves[64,65], offering a means to

externally control the dynamics of the system and may explain the details of broken symmetries.

We observed that adding a water droplet to the biofilm surface led to the transformation of spiral

waves into target and planar waves (Figure 3d-f). To recover the spiral waves, we increased the

surface temperature by gently heating the system. As expected, with the temperature rise, the

leading edge of the planar waves first became noisy, and multiple target waves randomly appeared

then finally converge to spiral waves (Supplementary Video7). These findings are in interesting

similarity with recent theoretical predictions in different nonreciprocal systems[50,64].

Additionally, by incorporating the water-soluble polymer polyethylene glycol (PEG) into the

biofilm, we managed to not only alter wave propagation patterns but also direct the movement of

inward growing target waves. Intriguingly, the addition of small drop of PEG modified the intrinsic

oscillation period of the pili dynamics, creating a radial gradient profile largely due to the slow

evaporation rate and uneven deposition of PEG on the surface. This inward wave guidance is

particularly controlled by frequency gradient (Figure 4a-d). The center has slow oscillations than

the edge. This gradient profile is the critical feature to understand the basics of broken symmetry

observed in the circular biofilms. In naturally growing colony oscillation frequency is also varying

(Figure 4a-b). We hypnotize that this variation is age dependent where the center of the colony is

slowing down due to the nutrition or oxygen depletion process. Before delving deeper into this

interesting symmetry-breaking process, we investigated how the aging of the biofilm affected the

oscillation dynamics. This is because symmetry breaking occurs at late stage where the aging is

critical. To do so, we first measured the oscillation periods as the uniform colony aged. We

observed that the periods of these spiral waves continuously increased (Figure 5a-b). These results

suggest that the intrinsic oscillation dynamics of the Pili decreased, likely due to depleted nutrients

or oxygen, during first 24 hours.

Figure 5: Age-Dependent Dynamics of Oscillations. *a) Sample image showing multiple pairs of spiral waves on a uniform biofilm surface. Scale bar: 100 μm. b) Period of the oscillations increases as the biofilm ages. Error bar shows the S.D. N= 10 measurements.*

Further, we examined the growing biofilm starting from a thin inoculation strip, we reproducibly

achieved a broken left-right symmetry for plane waves across the colony (Supplementary Video8).

The waves propagated from the edges toward the center. This asymmetry became particularly

pronounced a few days after inoculation, during which the colony edges continued to grow while

the center remained stationary (Figure 6a, b). To finally confirm this observation, we numerically

solve the coupled oscillators under varying frequency. Similarly, we found similar broken left-

right symmetry where the planar waves propagating towards the slower region of the biofilm

center (Figure 6c, d, Supplementary Video9).

Figure 6. Left-right symmetry breaking in naturally growing biofilms. (a) Representative image of a bacterial biofilm at a late growth stage (3 days post-inoculation). Metachronal waves propagate towards the biofilm center, with the leading edge displaying chaotic dynamics. Following growth cessation, spiral waves gradually converge into planar waves propagating inward. (b) Oscillation period increases towards the biofilm center. Error bar indicates the S.D. N= 10 measurements (c-d) Numerical simulation demonstrating the formation of planar waves and the left-right symmetry-breaking process, driven by a spatially varying intrinsic oscillation frequency(d), capturing dynamics observed in naturally growing biofilms.

Controlling left-right symmetry breaking

As a final step we focused on how to control the symmetry breaking and dynamics of metachronal waves. We noticed that temperature adjustments also reversibly effect the oscillatory behavior of the waves which increased the oscillation frequency (Figure 7a, b). This suggest that temperature could be also used as a complementary technique to dynamically guide the wave propagation by controlling the frequency gradients. To do so we embedded metallic pipes in the bulk agar plate and tuned the local temperatures (Supplementary Figure7). Setting different temperatures in these pipes creates temperature gradient (∇T~10°C/cm). This gradient profile converted spiral waves quickly to planar waves, propagating from warmer (31°C) to cooler (21°C) areas. Similarly, when we switched the temperature profiles wave propagation also reversed (Figure 7c-d). This result indicates that spatial temperature variations can effectively dictate wave propagation directions. These comprehensive observations confirm that the developed frequency gradients of intrinsic Pili oscillations is the critical feature breaking symmetry and controlling the propagation directions of the waves.

Figure 7: Temperature-Controlled Dynamics of Metachronal Waves and symmetry breaking.
a) Optical pulses from spiral waves on the biofilm surface as temperature increases from 21°C
(T_L) to 31°C (T_H). b) An increase in temperature raises the oscillation frequency c, d) Controlling
the propagation of the waves by creating a temperature gradient. Waves propagate from the
warmer area (fast oscillating, T_H) to the colder region (slow oscillating, T_L). d) Reversing the
temperature gradient changes the direction of propagations of the waves.

Discussion

Synchronization is a ubiquitous phenomenon in biological systems. A classic hallmark signature

of this dynamical process is global synchronization, where all components of the system tend to

merge in phase, as observed in fireflies, clocks and metronomes. Hydrodynamics or intrinsic

activities of the interactions, however, introduce nonreciprocity into these tightly coupled systems,

fundamentally altering the synchronization process. This action reaction asymmetry drives the

emergence of large-scale propagating waves, often emerge as spiral waves in two dimensions.

Remarkably, diverse systems from proteins in cell membrane to social amoebae exhibit similar

dynamics, highlighting the special importance of these waves in biological systems particularly

due to their transport capabilities.

From a symmetry perspective, nonreciprocity is the key feature in understanding these phenomena.

Recent advancements in active matter physics have bridged various seemingly unrelated fields

around this concept. Particularly, non-Hermitian physics, including PT-symmetry, provides a

beneficial macroscopic metric for understanding a system's response through the classical

formalism borrowed from quantum physics and optics. Despite the diverse nature of the dynamic

processes and the varying types of interactions—from predator-prey dynamics, specific chemical

reactions, to elasticity of active metamaterials—the common feature across these systems is to

maintain phase difference between the fundamental fields that support limit cycle oscillations

enabling collective motility and transport capabilities to the interacting active system.

We have observed that bacterial biofilm surfaces also known as active carpet exhibit similar

dynamics due to coupled and cyclic pili extension and retraction activities. Although these

nanoscale activities make the biophysical mechanisms complex and demanding to elucidate, the

fundamental characteristic of the process remains consistent: the cyclic motion of bacterial pili

forms limit cycle oscillations in phase space to disrupt local or action-reaction symmetry among

the interacting components.

Using a phenomenological model, we have concentrated on identifying critical parameters that

control the system dynamics and synchronization. We have also developed methods to manipulate

and control the collective dynamics and propagation of metachronal waves by adjusting these

critical parameters. From the perspective of active matter, the controllability of densely coupled

active mechanical components is crucial, especially at low Reynolds numbers where controlling

turbulence and managing long-range transport is vital. Biological systems can effectively address

these challenges and precisely control active and programable transport[66]. A deeper understanding

of the underlying principles and critical symmetries governing this controllability could provide

new insights into the complexities of biological systems. We speculate that these waves could

promote active diffusion of oxygen and DNA. Further biophysical investigations are needed to

delve deeper into these fascinating collective behaviors and its biological significance.

Furthermore, the dynamical similarities between non-Hermitian physics and PT symmetry and the

metachronal waves due to nonreciprocity are remarkable. Finally, we also hypothesize that the

concepts developed in no-Hermitian physics could bring new perspective to dissect the

complexities of these collective behaviors. Finally, our findings also raise several questions,

particularly regarding the biological significance of these waves in the physiology of the biofilm.

Rhythmic activities[67,68] play a critical role in bacterial communities. We hypothesize that this

synchronization may facilitate rhythmic behavior across the biofilm or enable expression and the

transport of specific biomolecules under stressful conditions. Further studies are needed to

elucidate the mechanisms underlying this emerging phenomenon.

Sharp Rising edge   Sharp Falling edge

OCC illumination

Supplementary Figure 1

Supplementary Figure 1. Asymmetric surface deformation drives directional propagation. OCC

imaging provides clear dark and bright optical contrasts, originating from oblique illumination and

asymmetric scattering by surface waves. An oil droplet was used to investigate the shape-

dependent propagation mechanism. Under oblique illumination, sharp rising edges generate dark

contrasts, while falling edges create bright contrasts. Sharp edges correspond to rapid pili

retraction, followed by gradual recovery.

Supplementary Figure 2

402 **Supplementary Figure 2.** GFP labeling and extracellular DNA staining were used to verify the

403 origin of the optical scattering, specifically to distinguish between surface versus bulk oscillations

404 in the biofilm. Neither fluorescence imaging method revealed oscillatory signals. Simultaneously

405 recorded gray signals, indicating the presence of original waves on the biofilm, were superimposed

406 for comparison.

Supplementary Figure 3

Supplementary Figure 3. In PA14 strains, mutations in pilB and fliK were introduced to identify the processes responsible for driving the oscillations. The pilB mutation abolished wave formation, whereas wave emergence persisted in the fliK mutant, which lacks functional flagella.

Supplementary Figure 4

427 **Supplementary Figure 4**. The PAO1 strain did not generate surface waves. Additionally, the

428 hyperpiliated pilH mutant also failed to exhibit oscillatory wave behavior.

Supplementary Figure 5

433 **Supplementary Figure 5.** Regular LB plates do not provide sufficient conditions for wave

434 generation in PN. However, eliminating yeast extract and replacing tryptone with a low

435 concentration (0.2×) peptone restored wave formation.

Supplementary Figure 6

Supplementary Figure 6. Comparative analysis of the genomic region encoding pili subunit

groups reveals that PA14 contains Group 3 pili, while PN exhibits a Group 4 and 5-like structure.

Both strains also encode accessory proteins associated with unknown pilus function.

Supplementary Figure 7. Optical and thermal images of an NGM plate equipped with metallic

pipes. A closed-loop water pumping-heating system was used to control the local temperature and

create a temperature gradient between the pipes. The spacing between the pipes was varied to

identify the optimal gradient profile.

Data availability

The critical experimental data generated or analyzed during this study are provided as supporting

video files. We did not generate additional data sets.

Software availability

All the codes used in the study will be available online.

Materials and Methods:

Analysis of Pilin Proteins and Accessory Genes: Genetic analyses were conducted using the complete genome sequence of Pseudomonas nitroreducens strain L4 chromosome[69] (NCBI Reference Sequence: NZ_CP120376.1). The reference genomic region between pilB and tRNA-thr was specifically compared.

Imaging System: Time-lapse imaging was performed using multiple microscopy techniques, selected for their capability to visualize surface topography effectively. We observed that oblique and polarized illumination techniques significantly enhanced the visibility of surface waves. During the evaporation process, the roughness of the biofilm significantly varies. During the evaporation process, the roughness of the biofilm varies significantly. We observed that oblique illumination is more effective on rough surfaces, whereas polarized imaging provides clearer results on flat surfaces. Oblique imaging was primarily conducted using Zeiss Smartzoom digital microscopes and Nikon Stereo SMZ18 microscopes equipped with contrast illumination. Additional polarized imaging utilized a Zeiss Axio Imager M2m. Successive images were captured at intervals of 10 seconds. Fluorescence imaging for GFP was conducted using 488 nm blue-light excitation with GFP emission filters, and DNA staining was visualized using RFP emission filters and imaged with an Andor EMCCD camera. The typical optical magnification was 10X.

Image Processing: To enhance wave visibility, successive images taken at 10-second intervals were subtracted, and the resulting false-colored difference images were superimposed onto the original ones. The time-dependent responses of surface pulses were analyzed using ImageJ software.

Bacterial growth and Preparation: The Pseudomonas nitroreducens (PN) bacterial strain used in this study was initially isolated from environmental samples and identified through 16S rRNA sequencing. Verification was performed through comparison with the corresponding strain obtained from DSMZ. For optical imaging, PN bacteria were grown overnight in LB broth at 21 °C on a shaker, and subsequently, 100 μl drops of bacterial suspension were placed onto 10-cm, 1-day-old, 1.4% solid agar, NGM plates. To ensure slow evaporation of excess moisture, humidity was maintained at approximately 80%, and plate lids were partially left open. Nematode growth medium (NGM) plates were prepared following standard protocols.

Numerical Simulation: Numerical simulations based on the Kuramoto model were performed using custom-developed MATLAB code with finite difference algorithms. The original code for the active solid model was obtained from prior literature and modified for improved visualization. Through extensive parameter exploration, we determined that large lattice structures with fixed boundary conditions effectively reproduced spiral and planar wave dynamics.

Author contributions: B.A. and A.Ko. initiated the project and performed the experiments. B.A., Y.I.Y., and M.B. Y.G. developed and executed numerical modeling. B.K. conducted 16S DNA sequencing and GFP labeling of PN. N.G. performed genome sequence analysis. E.T.G. and A.Ü. developed and conducted imaging for the temperature gradient control system. S.K.Ö. and C.K. conceptualized non-Hermitian and PT-symmetry principles. A.Ko. and B.A. drafted the manuscript, with contributions from all authors toward the final manuscript.

Acknowledgments: This work was supported by Airforce Office of Scientific Research (AFOSR) under Grant No: FA9550-22-1-0431 with Program Manager Ali Sayir and Tübitak Project No: 121C159. S.K.O acknowledges the support from Air AFOSR Multidisciplinary University

504 Research Initiative (MURI) Award No. FA9550-21-1-0202 **.** We thank Zeiss, Turkey division and

505 Kenan Doğru for digital microscopy installation and training.

Competing interests

507 The authors declare no competing interests.

Video Captions

Video 1. Time-lapse imaging of active *Pseudomonas nitroreducens* biofilm surfaces generating
metachronal waves. Global left-right symmetry is broken, and merging spiral waves at the edge
drive unidirectional planar waves toward the biofilm center. The wave wavelength is
approximately 100 µm. Associated with Figure 1a.

Video 2. Time-lapse imaging of circularly symmetric active biofilm surfaces generating inward-
propagating metachronal waves. Associated with Figure 1b.

Video 3. High magnification (100X) time-lapse imaging of biofilm surfaces illustrating bacterial
displacement and optical contrast changes.

Video 4. Time-lapse imaging of biofilm growth initiated from a single bacterium. After growth
cessation, dense regions begin firing first, and waves dominate the biofilm surface.

Video 5. Numerical simulation using the nonreciprocal Kuramoto model. Colors indicate
oscillation phases. Spiral waves merge to form target waves, eventually converging into planar
waves. Associated with Figure 2c.

Video 6. Numerical simulation based on the active solid model. Emergence of spiral and planar
waves within the lattice. Associated with Figure 2e.

Video 7 Time-lapse imaging of controlling the transition between waves to target and spiral by
manipulating the moisture of biofilm surface. Associated with Figure 3d-g

Video 8 Time-lapse imaging of a bacterial biofilm with broken left-right symmetry. Associated
with Figure 6.

Video 9. Time-lapse imaging of controlling propagating planar waves under a varying temperature
gradient. Associated with Figure 7.

Video 10. Numerical simulation of propagating planar waves influenced by a frequency gradient.
Associated with Figure 7c.

Competing interests

Authors declare no competing interests.

541 **Tables**

Strain	Parent	Operation	Genotype
BAK132	PN	Isolated from environmental samples, transformed pUCP18-MCSgfpmut3	*ampR*
BAK133	PA14	Received from F. M . Ausubel lab	
BAK134	LD2222	Received from Lars Dietrich Lab[70]	PA14 ΔpilB
BAK135	LD2221	Received from Lars Dietrich Lab[70]	PA14 ΔflgK
BAK136	PAO1 PilH	Received from Joanne Engel	ΔPilH
BAK	PAO1	Received from DSMZ 10145GFP	

543 *Table 1: List of strains used in this study.*

References

1   Wadhwa, N. & Berg, H. C. Bacterial motility: machinery and mechanisms. *Nat Rev Microbiol* **20**,
161-173, doi:10.1038/s41579-021-00626-4 (2022).
2   Lauga, E. Bacterial Hydrodynamics. *Annu Rev Fluid Mech* **48**, 105-130, doi:10.1146/annurev-
fluid-122414-034606 (2016).
3   Lopez, D. & Lauga, E. Dynamics of swimming bacteria at complex interfaces. *Phys Fluids* **26**,
doi:Artn 071902

10.1063/1.4887255 (2014).
4   Kearns, D. B. A field guide to bacterial swarming motility. *Nat Rev Microbiol* **8**, 634-644,
doi:10.1038/nrmicro2405 (2010).
5   Purcell, E. M. Life at low Reynolds number. *American Journal of Physics* **45**, 3-11,
doi:10.1119/1.10903 %J American Journal of Physics (1977).
6   Byron, M. L. *et al.* Metachronal Motion across Scales: Current Challenges and Future Directions.
Integr Comp Biol **61**, 1674-1688, doi:10.1093/icb/icab105 (2021).
7   Leoni, M. & Liverpool, T. B. Hydrodynamic synchronization of nonlinear oscillators at low
Reynolds number. *Phys Rev E* **85**, doi:ARTN 040901

10.1103/PhysRevE.85.040901 (2012).
8   Golestanian, R., Yeomans, J. M. & Uchida, N. Hydrodynamic synchronization at low Reynolds
number. *Soft Matter* **7**, 3074-3082, doi:10.1039/c0sm01121e (2011).
9   Sleigh, M. A., Blake, J. R. & Liron, N. The Propulsion of Mucus by Cilia. *Am Rev Respir Dis* **137**,
726-741, doi:DOI 10.1164/ajrccm/137.3.726 (1988).
10 Hirokawa, N., Tanaka, Y., Okada, Y. & Takeda, S. Nodal flow and the generation of left-right
asymmetry. *Cell* **125**, 33-45, doi:10.1016/j.cell.2006.03.002 (2006).
11 Cohen, A. H., Holmes, P. J. & Rand, R. H. The Nature of the Coupling between Segmental
Oscillators of the Lamprey Spinal Generator for Locomotion - a Mathematical-Model. *J Math Biol*
13, 345-369, doi:Doi 10.1007/Bf00276069 (1982).
12 Yaman, Y. I. & Ramanathan, S. Controlling human organoid symmetry breaking reveals signaling
gradients drive segmentation clock waves. *Cell* **186**, 513-+, doi:10.1016/j.cell.2022.12.042
(2023).
13 Wu, Q. *et al.* Active metamaterials for realizing odd mass density. *P Natl Acad Sci USA* **120**,
doi:ARTN e2209829120

10.1073/pnas.2209829120 (2023).
14 Bender, C. M. & Boettcher, S. Real spectra in non-Hermitian Hamiltonians having PT symmetry.
Physical Review Letters **80**, 5243-5246, doi:DOI 10.1103/PhysRevLett.80.5243 (1998).
15 Chen, Y. Y., Li, X. P., Scheibner, C., Vitelli, V. & Huang, G. L. Realization of active metamaterials
with odd micropolar elasticity. *Nature Communications* **12**, doi:ARTN 5935

10.1038/s41467-021-26034-z (2021).
16 Fruchart, M., Scheibner, C. & Vitelli, V. Odd Viscosity and Odd Elasticity. *Annual Review of*
Condensed Matter Physics **14**, 471-510, doi:10.1146/annurev-conmatphys-040821-125506
(2023).
17 Hickey, D. J., Golestanian, R. & Vilfan, A. Nonreciprocal interactions give rise to fast cilium
synchronization in finite systems. *P Natl Acad Sci USA* **120**, doi:ARTN e2307279120

10.1073/pnas.2307279120 (2023).

18   Marchetti, F. B. a. M. C. Nonreciprocal Pattern Formation of Conserved Fields. *Phys. Rev. X* **14**,
021014.
19   Fruchart, M., Hanai, R., Littlewood, P. B. & Vitelli, V. Non-reciprocal phase transitions. *Nature*
592, doi:10.1038/s41586-021-03375-9 (2021).
20   Dinelli, A. *et al.* Non-reciprocity across scales in active mixtures. *Nat Commun* **14**, 7035,
doi:10.1038/s41467-023-42713-5 (2023).
21   You, Z. H., Baskaran, A. & Marchetti, M. C. Nonreciprocity as a generic route to traveling states.
P Natl Acad Sci USA **117**, 19767-19772, doi:10.1073/pnas.2010318117 (2020).
22   Tang, E., Agudo-Canalejo, J. & Golestanian, R. Topology Protects Chiral Edge Currents in
Stochastic Systems. *Phys Rev X* **11**, doi:ARTN 031015

10.1103/PhysRevX.11.031015 (2021).
23   Uchida, N. & Golestanian, R. Synchronization and Collective Dynamics in a Carpet of Microfluidic
Rotors. *Physical Review Letters* **104**, doi:ARTN 178103

10.1103/PhysRevLett.104.178103 (2010).
24   Hatano, N. & Nelson, D. R. Localization transitions in non-Hermitian quantum mechanics.
Physical Review Letters **77**, 570-573, doi:DOI 10.1103/PhysRevLett.77.570 (1996).
25   Ozdemir, S. K., Rotter, S., Nori, F. & Yang, L. Parity-time symmetry and exceptional points in
photonics. *Nat Mater* **18**, 783-798, doi:10.1038/s41563-019-0304-9 (2019).
26   El-Ganainy, R. *et al.* Non-Hermitian physics and PT symmetry. *Nat Phys* **14**, 11-19,
doi:10.1038/Nphys4323 (2018).
27   Miri, M. A. & Alù, A. Exceptional points in optics and photonics. *Science* **363**, 42-+, doi:ARTN
eaar7709

10.1126/science.aar7709 (2019).
28   Naghiloo, M., Abbasi, M., Joglekar, Y. N. & Murch, K. W. Quantum state tomography across the
exceptional point in a single dissipative qubit. *Nat Phys* **15**, 1232-+, doi:10.1038/s41567-019-
0652-z (2019).
29   Zhang, J. W. *et al.* Dynamical control of quantum heat engines using exceptional points. *Nature*
Communications **13**, doi:ARTN 6225

10.1038/s41467-022-33667-1 (2022).
30   Ding, L. Y. *et al.* Experimental Determination of

-Symmetric Exceptional Points in a Single Trapped Ion. *Physical Review Letters* **126**, doi:ARTN 083604

10.1103/PhysRevLett.126.083604 (2021).
31   Wu, Y. *et al.* Observation of parity-time symmetry breaking in a single-spin system. *Science* **364**,
878-+, doi:10.1126/science.aaw8205 (2019).
32   Ergoktas, M. S. *et al.* Topological engineering of terahertz light using electrically tunable
exceptional point singularities. *Science* **376**, 184-+, doi:10.1126/science.abn6528 (2022).
33   Soleymani, S. *et al.* Chiral and degenerate perfect absorption on exceptional surfaces. *Nature*
Communications **13**, doi:ARTN 599

10.1038/s41467-022-27990-w (2022).
34   Zhong, Q. *et al.* Sensing with Exceptional Surfaces in Order to Combine Sensitivity with
Robustness. *Physical Review Letters* **122**, doi:ARTN 153902

10.1103/PhysRevLett.122.153902 (2019).

35   Lau, H. K. & Clerk, A. A. Fundamental limits and non-reciprocal approaches in non-Hermitian
636   quantum sensing. *Nature Communications* **9**, doi:ARTN 4320

637 10.1038/s41467-018-06477-7 (2018).
638 36   Tlusty, T. Exceptional topology in ordinary soft matter. *Phys Rev E* **104**, doi:ARTN 025002

639 10.1103/PhysRevE.104.025002 (2021).
640 37   Biktashev, V. N., Brindley, J., Holden, A. V. & Tsyganov, M. A. Pursuit-evasion predator-prey
641   waves in two spatial dimensions. *Chaos* **14**, 988-994, doi:10.1063/1.1793751 (2004).
642 38   Loiseau, E. *et al.* Active mucus-cilia hydrodynamic coupling drives self-organization of human
643   bronchial epithelium. *Nat Phys* **16**, 1158-+, doi:10.1038/s41567-020-0980-z (2020).
644 39   Guzmán-Lastra, F., Löwen, H. & Mathijssen, A. J. T. M. Active carpets drive non-equilibrium
645   diffusion and enhanced molecular fluxes. *Nature Communications* **12**, 1906,
646   doi:10.1038/s41467-021-22029-y (2021).
647 40   Kim, M. J. & Breuer, K. S. Use of bacterial carpets to enhance mixing in microfluidic systems. *J
648   Fluid Eng-T Asme* **129**, 319-324, doi:10.1115/1.2427083 (2007).
649 41   Tan, T. H. *et al.* Odd dynamics of living chiral crystals. *Nature* **607**, 287-+, doi:10.1038/s41586-
650   022-04889-6 (2022).
651 42   Gregor, T., Fujimoto, K., Masaki, N. & Sawai, S. The Onset of Collective Behavior in Social
652   Amoebae. *Science* **328**, 1021-1025, doi:10.1126/science.1183415 (2010).
653 43   Dreher, A., Aranson, I. S. & Kruse, K. Spiral actin-polymerization waves can generate amoeboidal
654   cell crawling. *New J Phys* **16**, doi:Artn 055007

655 10.1088/1367-2630/16/5/055007 (2014).
656 44   Ford, H. Z., Manhart, A. & Chubb, J. R. Controlling periodic long-range signalling to drive a
657   morphogenetic transition. *Elife* **12**, doi:ARTN e83796

10.7554/eLife.83796 (2023).
45   Bull, M. S., Kroo, L. A. & Prakash, M. in *arXiv e-prints*   arXiv:2107.02930 (2021).
46   Kühn, M. J. *et al.* Mechanotaxis directs

aeruginosa twitching motility. *P Natl Acad Sci USA* **118**, doi:ARTN e2101759118

10.1073/pnas.2101759118 (2021).
47   Kühn, M. J. *et al.* Two antagonistic response regulators control

polarization during mechanotaxis. *Embo J* **42**, doi:10.15252/embj.2022112165 (2023).
48   Xu, H. R., Huang, Y. L., Zhang, R. & Wu, Y. L. Autonomous waves and global motion modes in
666   living active solids. *Nat Phys* **19**, 46-+, doi:10.1038/s41567-022-01836-0 (2023).
667 49   Liu, S. Q., Li, Y., Wang, Y. H. & Wu, Y. L. Emergence of large-scale mechanical spiral waves in
668   bacterial living matter. *Nat Phys* **20**, doi:10.1038/s41567-024-02457-5 (2024).
669 50   Rana, N. & Golestanian, R. Defect Solutions of the Nonreciprocal Cahn-Hilliard Model: Spirals
670   and Targets. *Physical Review Letters* **133**, 078301, doi:10.1103/PhysRevLett.133.078301 (2024).
671 51   Demir, E., Yaman, Y. I., Basaran, M. & Kocabas, A. Dynamics of pattern formation and
672   emergence of swarming in Caenorhabditis elegans. *Elife* **9**, doi:ARTN e52781

673 10.7554/eLife.52781 (2020).
674 52   Kus, J. V., Tullis, E., Cvitkovitch, D. G. & Burrows, L. L. Significant differences in type IV pilin allele
675   distribution among Pseudomonas aeruginosa isolates from cystic fibrosis (CF) versus non-CF
676   patients. **150**, 1315-1326, doi:https://doi.org/10.1099/mic.0.26822-0 (2004).

677 53   Meron, E. Pattern-Formation in Excitable Media. *Phys Rep* **218**, 1-66, doi:Doi 10.1016/0370-
678   1573(92)90098-K (1992).
679 54   Tan, T. H. *et al.* Topological turbulence in the membrane of a living cell. *Nat Phys* **16**, 657-662
680   %U https://ui.adsabs.harvard.edu/abs/2020NatPh..2016..2657T, doi:10.1038/s41567-020-0841-
681   9 %@ 1745-2473 (2020).
682 55   Aranson, I. S. & Rabinovich, M. I. Dynamics of spiral waves in non-equilibrium media. *Journal of
683   Physics A: Mathematical and General* **23**, 299, doi:10.1088/0305-4470/23/3/014 (1990).
684 56   Paullet, J. E. & Ermentrout, G. B. Stable Rotating Waves in Two-Dimensional Discrete Active
685   Media. *SIAM Journal on Applied Mathematics* **54**, 1720-1744, doi:10.1137/S0036139993250683
686   %U https://epubs.siam.org/doi/abs/10.1137/S0036139993250683 (1994).
687 57   Chakrabarti, B., Furthauer, S. & Shelley, M. J. A multiscale biophysical model gives quantized
688   metachronal waves in a lattice of beating cilia. *P Natl Acad Sci USA* **119**, doi:ARTN e2113539119

689   10.1073/pnas.2113539119 (2022).
690 58   Baconnier, P. *et al.* Selective and collective actuation in active solids. *Nat Phys* **18**, 1234-+,
691   doi:10.1038/s41567-022-01704-x (2022).
692 59   Kim, P. J., Ko, T. W., Jeong, H. & Moon, H. T. Pattern formation in a two-dimensional array of
693   oscillators with phase-shifted coupling. *Phys Rev E* **70**, doi:ARTN 065201

694   10.1103/PhysRevE.70.065201 (2004).
695 60   Gupta, R. K., Kant, R., Soni, H., Sood, A. K. & Ramaswamy, S. Active nonreciprocal attraction
696   between motile particles in an elastic medium. *Phys Rev E* **105**, doi:ARTN 064602

697   10.1103/PhysRevE.105.064602 (2022).
698 61   Kim, S., Park, S. H. & Ryu, C. S. Multistability in coupled oscillator systems with time delay.
699   *Physical Review Letters* **79**, 2911-2914, doi:DOI 10.1103/PhysRevLett.79.2911 (1997).
700 62   Ohta, H. & Sasa, S. Critical phenomena in globally coupled excitable elements. *Phys Rev E* **78**,
701   doi:ARTN 065101

702   10.1103/PhysRevE.78.065101 (2008).
703 63   Gu, F. C., Guiselin, B., Bain, N., Zuriguel, I. & Bartolo, D. Emergence of collective oscillations in
704   massive human crowds. *Nature* **638**, doi:10.1038/s41586-024-08514-6 (2025).
705 64   Rana, N. & Golestanian, R. Defect interactions in the non-reciprocal Cahn-Hilliard model. *New J
706   Phys* **26**, doi:ARTN 123008

707   10.1088/1367-2630/ad9859 (2024).
708 65   Wang, G. Q. *et al.* Target-wave to spiral-wave pattern transition in a discrete Belousov-
709   Zhabotinsky reaction driven by inactive resin beads. *Phys Rev E* **82**, doi:ARTN 045201

710   10.1103/PhysRevE.82.045201 (2010).
711 66   Yang, F., Liu, S. C., Lee, H. J., Phillips, R. & Thomson, M. Dynamic flow control through active
712   matter programming language. *Nat Mater* **24**, doi:10.1038/s41563-024-02090-w (2025).
713 67   Sartor, F. & Kovács, A. T. Rhythmic Spatial Self-Organization of Bacterial Colonies. *Mbio* **13**,
714   doi:10.1128/mbio.01703-22 (2022).
715 68   Eelderink-Chen, Z. *et al.* A circadian clock in a nonphotosynthetic prokaryote. *Sci Adv* **7**,
716   doi:ARTN eabe2086

717   10.1126/sciadv.abe2086 (2021).
718 69   Liu, H. Y., Zhuang, L. B. & Zeng, Q. C. Complete genome sequence data of

719   L4, an endophyte isolated from cotton plants. *Data Brief* **55**, doi:ARTN 110639

720   10.1016/j.dib.2024.110639 (2024).
721   70   Hölscher, T. *et al.* Motility, Chemotaxis and Aerotaxis Contribute to Competitiveness during
722   Bacterial Pellicle Biofilm Development. *J Mol Biol* **427**, 3695-3708,
723   doi:10.1016/j.jmb.2015.06.014 (2015).