Virus glycoprotein nanodisc platform for vaccine design

Kimmo Rantalainen[1,2,3], Alessia Liguori[1,2,3], Gabriel Ozorowski[,3,4], Claudia Flynn[1,2,3], Jon M. Steichen[1,2,3], Olivia Swanson[1,2,3,4], Patrick J. Madden[3,5], Sabyasachi Baboo[4], Swastik Phulera[4], Anant Gharpure[4], Danny Lu[1,2,3], Oleksandr Kalyuzhniy[1,2,3], Patrick Skog[1,2,3], Sierra Terada[1,2,3], Monolina Shil[3,5], Jolene K. Diedrich[4], Erik Georgeson[1,2,3], Ryan Tingle[1,2,3], Saman Eskandarzadeh[1,2,3], Wen-Hsin Lee[4], Nushin Alavi[1,2,3], Diana Goodwin[1,2,3], Michael Kubitz[1,2,3], Sonya Amirzehni[1,2,3], Sunny Himansu[6], Devin Sok[1,2,3], Jeong Hyun Lee[1,2,3], John R. Yates III[4], James C. Paulson[1,3,6], Shane Crotty[3,5,7], Torben Schiffner[1,2,3,8,*], Andrew B. Ward[,3,4,*], William R. Schief[1,2,3,6,*]

Affiliations

[1] Department of Immunology and Microbiology, The Scripps Research Institute, La Jolla, CA, USA.

[2] IAVI Neutralizing Antibody Center, The Scripps Research Institute, La Jolla, CA, USA.

[3] Consortium for HIV/AIDS Vaccine Development (CHAVD), The Scripps Research Institute, La Jolla, CA, USA.

[4] Department of Integrative Structural and Computational Biology, The Scripps Research Institute, La Jolla, CA 92037, USA.

[5] Center for Vaccine Innovation, La Jolla Institute for Immunology, La Jolla, CA 92037, USA.

[6] Moderna Inc., Cambridge, MA, 02139, USA.

[7] Division of Infectious Diseases, Department of Medicine, University of California San Diego, La Jolla, CA 92037, USA.

[8] Institute for Drug Discovery, Leipzig University Medical Faculty, Leipzig, 04103, Germany

*Corresponding authors. Email: torbens@scripps.edu (T.S.), andrew@scripps.edu (A.B.W.), or schief@scripps.edu (W.R.S.)

Abstract

Transmembrane glycoproteins of enveloped viruses are the targets of neutralizing antibodies and essential vaccine antigens. mRNA-LNP technology allows in situ production of transmembrane glycoproteins upon immunization, but biophysical characterization of transmembrane antigens and in vitro analysis of post-immunization antibody responses typically rely on soluble proteins. Here, we present a methodological platform for assembling transmembrane glycoprotein vaccine candidates into lipid nanodiscs. We demonstrate the utility of the nanodiscs in HIV membrane proximal external region (MPER)-targeting vaccine development by binding assays using surface plasmon resonance (SPR), ex vivo B cell sorting with fluorescence-activated cell sorting (FACS), and by determining the structure of a prototypical HIV MPER-targeting immunogen nanodisc in complex with three broadly neutralizing antibodies (bnAbs), including the MPER bnAb 10E8, to 3.5 Å by cryogenic electron microscopy (cryo-EM), providing a template for structure-based immunogen design for MPER. Overall, the platform offers a tool for accelerating the development of next-generation viral vaccines.

Summary

In the past two decades, nanodiscs have emerged as means to encapsulate transmembrane proteins into a stable and more native-like environment. In most cases, proteins are extracted from the membrane by detergent solubilization and then assembled into a nanodisc by removal of the detergent in presence of lipid molecules and a scaffold protein. Since the first apolipoprotein-derived membrane scaffold proteins (MSPs) were introduced, several variations and alternative scaffolds have been described, allowing for a variety of disc sizes and versatile experimental approaches[1-5]. Structural studies of membrane proteins have perhaps benefitted the most from nanodiscs, resulting in important advances in understanding of membrane protein structure-function relationship[6, 7]. They have also been used successfully in a variety of functional and biophysical approaches such as surface plasmon resonance (SPR)[8-10].

Viral glycoproteins of enveloped viruses are membrane proteins and essential targets for vaccine development. Structure-based vaccine design was an integral part of the successful

COVID-19 vaccine development[11-13], and a significant portion of the groundwork stemmed from

decades of HIV vaccine development[14]. Structure-based methods, combined with rapidly

advancing computational protein design approaches and mRNA lipid nanoparticle (mRNA-LNP)

technology, are now revolutionizing iterative vaccine development[15, 16]. Immunogen delivery by

mRNA-LNP expands the available immunogens to more native-like constructs that may contain

the membrane-proximal epitopes such as the recently developed cleavage independent NFL

trimers expressing HIV Env[17] and the MD39.3 gp151 Env trimer used in the HVTN302 clinical

trial (NCT05217641), but recombinant transmembrane glycoprotein production for biophysical

characterization of the immunogens remains challenging. In most studies, membrane proteins are

truncated before the transmembrane domain to obtain soluble ectodomains with higher expression

levels and easier handling and applicability in downstream analytical methods. The regions

excluded typically include the membrane-proximal external region (MPER), transmembrane

domain (TM), and intracellular C-terminal domain (CT). In influenza HA, the anchor region

analogous to HIV MPER joins the ectodomain as flexible linkers to three transmembrane helices

and allows the trimer to tilt in relation to bilayer surface[18]. Both Env MPER and HA anchor have

highly conserved epitopes for neutralizing antibodies (nAbs) making them attractive targets for

vaccine development[19-23]. Significant progress in MPER targeted HIV vaccine development has

been made recently with peptide-liposome formulations showing induction of heterologous

neutralizing antibody B cell lineages in humans, and with mRNA delivered germline-targeting

(GT) epitope scaffold efficiently inducing antibody precursors in non-human primates (NHPs) and

mice[24, 25]. Analysis of antibodies elicited by these vaccines as well as further immunogen

development would benefit from inclusion of the full epitope in recombinantly expressed proteins.

For example, recombinant proteins matching the transmembrane immunogens delivered as mRNA

are required for accurate immunogen assessment in iterative vaccine design methods, such as SPR

and structural studies. Importantly, transmembrane versions of Env may also be required for more

accurate representations of the glycan shield, as many neutralizing antibodies bind specifically to

glycans, and glycoproteins expressed as transmembrane proteins have been shown to have glycan

shield compositions closer to the native glycan shield[26-28].

Some uses of nanodiscs to support vaccine development have been demonstrated, particularly in efforts to characterize the structures of HIV MPER-targeting HIV antibody epitopes and transmembrane domains of virus glycoproteins in their native membrane bilayer environment[29-33]. The nanodisc assembly platform presented here provides the reproducibility, scalability, and accurate replication of the vaccine candidate properties that enable routine use of transmembrane viral glycoproteins in key methods employed in iterative, rational vaccine design. We demonstrate use of nanodiscs to measure antibody affinities by SPR, as FACS probes to sort antigen-specific B cells from mouse and non-human primate (NHP) immunization experiments, and for solving the cryo-EM structure of a prototype HIV boost vaccine candidate in complex with three broadly neutralizing antibodies (bnAbs) that target distinct sites of vulnerability on the HIV Env surface. We highlight the benefits of the approach by reporting the structure of the entire protein epitope of an HIV MPER-targeting bnAb 10E8. Finally, we show the adaptability of the platform to other glycoproteins by assembling engineered Ebola virus glycoprotein in nanodiscs using identical conditions to HIV Env.

Results

Glycoprotein Nanodisc Assembly and Analysis Workflow

We set to establish a workflow that provides material suitable for methods commonly used in rational, iterative vaccine design (Fig. 1). The workflow was set up using the engineered and highly expressing HIV Env clone BG505 MD39.3 gp151[34,35] that is currently being tested in a phase 1 clinical trial HVTN 302 (NCT05217641). A linker, followed by an HRV3C protease cleavage site and a strep tag was added to the intracellular C-terminus, leading to a nanodisc assembly construct base that will hereon be referred to as Env gp151 ND (Fig. 2a-b). This construct was then used to optimize the throughput and reproducibility of nanodisc production, leading to a standardized workflow (Fig. 2c). In short, transmembrane glycoproteins (GPs) were expressed on the surface of human FreeStyle 293-F cells, followed by detergent solubilization with TX-100-containing buffer. As the transmembrane domain of viral glycoproteins is composed of only three helices – one per protomer – detergent solubilization by TX-100 was not expected to disturb the quaternary

structure of the engineered GP trimer as indicated by the binding of structure specific antibodies

to extracted GPs[27, 28]. The detergent-solubilized glycoprotein was then assembled into nanodiscs

using the MSP1D1 scaffold with or without a biotin tag. While a variety of scaffold proteins are

available, we opted to reduce variables and focused on using the standard MSP1D1 scaffold for

all purposes. The developed "batch assembly" approach, resembling previously described "on-

column"[36] and "on-bead"[37] methods, facilitated reproducibility and efficient assembly while GPs

remained bound to the matrix. Empty nanodiscs for control experiments were produced by

omitting GP in the assembly step and using His-tag of the scaffold protein with NiNTA affinity

purification prior to SEC. Utilizing this workflow, up to 12 samples were routinely processed

simultaneously in five days from transfection to purified nanodisc. One liter of cells yielded

approximately 100 to 700 µg of pure GP nanodiscs, which was sufficient for one to three endpoint

assays. For example, a single production batch produced an EBOV GP nanodiscs, an HIV Env

nanodisc-Fab complex for cryo-EM structural studies, and eight FACS probes for B cell sorting.

Final GP nanodisc preparations were stable for at least three months at +4°C based on absorbance

at 280 nm, presence of trimeric GP in nanodisc in negative stain (ns) EM 2D class averages, and

activity in SPR analysis (Fig 3 b).

The workflow was then piloted in an immunogen development project aiming to increase

the affinity of MPER targeting antibody 10E8 and therefore guiding immune responses to MPER

domain. In the native membrane context, the C-terminal MPER helix that is targeted by the heavy

chain complementarity determining region 3 (CDRH3) of 10E8 is sterically occluded (Fig 2b) and

one of the design goals was to improve the MPER access. To this end, N-linked glycosylation sites

at positions N88, N618, and N625 (HxB2 numbering) adjacent to the epitope were removed at the

base of the trimer. An additional R696S mutation that improved 10E8 binding while maintaining

trimeric Env conformation was discovered by mammalian surface display library screening.

Earlier structural analysis indicated that R696 resides at the intersection of the three

transmembrane helices[29, 38]; therefore, this mutation was predicted to disrupt helix interactions and

further open the trimer base. The resulting construct will be referred to as Env gp151 MPER ND.

A negative control for MPER antibody binding and a KO FACS probe for epitope-specific B cell

sorting, designated as Env gp151 MPER-KO ND, was also generated (Fig. 2a). This construct
included MPER mutations W672A, F673R, T676R, N677I, L679M, W680E, Y681W, and
W683D, which were previously reported to collectively prevent the binding of MPER-targeting
antibodies while preserving the overall trimeric structure of Env and other bnAb epitopes[24], or
screened for this study to improve the expression construct. Prototypical Ebola GP immunogen
EBOV GP Mayinga ND was based on a construct with trimer stabilizing mutations T577P and
K588F[39], to which native EBOV GP TM domain and the same intracellular elements as in the HIV
Env constructs were added. Assembled nanodiscs generally exhibited broad peaks in SEC spectra
indicative of extensive glycosylation and variation in hydrodynamic radius that may result from
positional heterogeneity of Env in the disc, and >95% purity and bands of the expected molecular
weight as examined by SDS-PAGE (Supplementary Fig. 1a-b). Ns EM 2D class averages revealed
features typical of nanodiscs with trimeric GP with or without Fabs (Supplementary Fig. 4a). In
addition to the MPER targeting immunogen constructs and EBOV GP Mayinga ND, a germline-
targeting HIV vaccine candidate N332-GT5 gp151[40, 41] was assembled into nanodiscs using the
standardized workflow to test general applicability (Supplementary Fig. 2a). Mass spectrometric
glycan analysis using the DeGlyPHER method[42] was done prior to and after nanodisc assembly to
ascertain site-specific glycan occupancy and processivity of Env constructs. This analysis
confirmed the presence of all expected glycans as well as a high proportion of complex glycans in
transmembrane constructs characteristic of native, transmembrane HIV Env (Supplementary Fig.
1c). Germline-targeting immunogen N332-GT5 showed a higher proportion of complex type
glycans in soluble version of the immunogen, matching closely the glycan composition of the
transmembrane version. All transmembrane Envs showed higher glycan occupancy in gp41
subunit.
164
165 **GP nanodisc binding kinetics to antibodies measured by SPR**
166 SPR is an essential method in iterative vaccine design that measures antibody affinities to
167 engineered immunogens, assessing the effect of the designed features as well as affinities of
168 antibodies induced by vaccination (Fig 1). We set to establish scalable and reproducible conditions

for measuring antibody binding kinetics against transmembrane GPs in nanodiscs in three distinct SPR modalities (Fig. 3a). Modality A utilized an anti-affinity tag capture strategy in which anti-strep-tag antibody was covalently immobilized on the chip surface, glycoprotein nanodiscs were captured using the intracellular strep tag, and Fab analytes were employed to study monovalent interactions. Modality B employed a low surface density IgG capture strategy, in which anti-human IgG capture antibody was first covalently immobilized, and glycoprotein-specific IgG ligands were captured at reduced capture time to limit the ligand density. Glycoprotein nanodiscs were then injected as analytes. In modality C, nanodisc capture is followed by IgG flow as analyte instead of Fabs, allowing a high throughput scouting approach with low nanodisc sample consumption but with reduced kD accuracy due to increased bivalent interaction. Establishing these three modalities for nanodisc GPs allowed flexibility in experiment design based on the need in iterative vaccine development step and sample availability.

Next, we used the GP nanodisc SPR modalities to establish baseline kinetics for neutralizing and non-neutralizing monoclonal HIV Env antibodies spanning diverse epitopes (Fig. 3b-d, Supplementary Fig. 2b-d). Modality A measured 250 – 100 nM affinity for 10E8 to Env gp151 ND design base construct (Fig. 3b) and slightly higher affinity of 36 – 42 nM using modality B, indicating that the low-density capture reduced but did not eliminate the avidity effects of Modality B. Using modality A, we observed 70-fold increase in 10E8 affinity (250 nM vs 3.6 nM) to Env gp151 MPER ND as a result of removal of the MPER epitope-proximal glycans and the mutation R696S in TM (Fig. 3c). Kinetic analysis indicated that the improved affinity in the engineered immunogen was primarily attributed to a decreased off-rate (8.1×10^{-3} s^{-1} for Env gp151 ND vs 3.5×10^{-4} s^{-1} for Env gp151 MPER ND. Fig. 3d). Incorporation of MPER-KO mutations completely abrogated 10E8 binding. Additional MPER targeting bnAbs and bnAbs targeting other epitopes on the surface of Env were also tested for more comprehensive antigenic profiling. Expectedly, this confirmed binding of MPER targeting antibodies to Env gp151 ND and no binding to Env gp140 (Supplementary Fig. 2b), and similar affinities of non-MPER bNabs to soluble Env gp140 and Env gp151 ND (Supplementary Fig. 2c), with reduced or absent binding of non-neutralizing antibodies RM19R and RM20A3 to Env gp151 ND as compared to Env gp140

(Supplementary Fig 2d). These two antibodies target the base of the trimer, which is exposed in
Env gp140, whereas in Env gp151 ND access is presumably restricted by MPER and TM domains
as well as the lipid surface of the nanodisc. Finally, we employed modality C to confirm the
binding of two Ebola GP glycan cap specific antibodies EBOV-296 and 13C6 to EBOV GP
nanodiscs (Fig. 3e). Collectively, these data demonstrate that GP nanodiscs and SPR can be used
to characterize the antigenic landscape of the designed transmembrane immunogen, inform
immunogen selection for in vivo studies, identify suitable complexes for structural studies by cryo-
EM and support GP nanodisc use as B cell sorting probes in FACS (Fig 1).
205

FACS sorting of B Cells from immunized animal models using nanodisc GP probes
Single B cell sorting and sequencing is routinely used to analyze B cell responses induced in vivo.
Typically, biotin-tagged variants of the immunogen are used with streptavidin. This allows
conjugation with multiple fluorochromes to increase selection specificity. By including a matching
sorting probe with KO mutations in the targeted epitope, epitope-specific responses can be
separated from off-target responses against unrelated epitopes. Soluble GP protein constructs
solely consisting of the ectodomain are most often used as antigenic baits. For immunization
studies where membrane-proximal epitopes are of interest, such as the HIV MPER, soluble probes
would not be able to extract all appropriate responses to the targeted epitope due to several
challenges which include 1) the hydrophobicity of the epitope regions which make bait production
challenging, 2) non-native or incomplete structural conformation of the epitope in shorter peptide
representations of the antigen, 3) missing lipid membrane components which constitute a portion
of the native epitope. Here, we used biotinylated MSP1D1 scaffold to generate nanodisc GP FACS
probes. We first tested Env gp151 ND probe binding to FACS compensation beads conjugated to
10E8, DH511 or BG18 bnAbs (Fig. 4a). Background signal was measured as binding to either
MSP1D1 scaffold protein alone, empty nanodisc assembled with DOPC lipids, or to empty
nanodisc with mixture of neutral and charged lipids. Env gp151 ND exhibited substantially higher
signal compared to controls, confirming that adequate separation from background signal can be
detected. We detected no binding with any of the tested probes to unimmunized C57BL/6 (B6)

mouse splenocytes, and a strong signal against engineered Ramos cells expressing the bnAb VRC01 as its B cell receptor (BCR) [43]. This data indicated that Env nanodisc probes were suitable for detection of B cells from mice by flow cytometry that specifically target Env.

We then proceeded with a pilot immunization experiment in mice to explore if vaccine elicited antigen-specific B cells could be identified using Env nanodiscs probes. We utilized transgenic mice expressing the human D3-3 and J_H6 genes that allow mice to express long HCDR3s, including HCDR3 precursors of HIV MPER bnAbs 10E8, DH511 and LN01[24]. Mice were intramuscularly injected with 10 µg of mRNA-LNPs encoding HIV Env gp151 MPER ND construct. Animals were sacrificed, and spleens and lymph nodes were harvested 6 weeks post immunization. B cells were stained with immunogen-matched (WT) nanodisc probes on two distinct fluorophores and Env gp151 MPER-KO ND nanodisc probe with KO mutations in the MPER to identify MPER-specific cells (Fig. 4b and Supplementary Fig. 3). Among class-switched memory B cells (defined as $CD19^+IgD^-IgM^-$), 0.7% of cells were determined to be antigen-specific, compared to 0.15% within the IgD^+IgM^+ naïve B cell population (Fig. 4c). This ~4.7-fold increase was statistically significant (p=0.0177), indicating that the mRNA immunization elicited HIV Env gp151 MPER -specific responses that could be detected by Env nanodisc probes. The proportion of WT Env nanodisc-binding B cells that selectively bound to the 10E8 epitope (%KO$^-$ of WT^{++} B cells) was indistinguishable between the naïve and memory B cell compartments, suggesting that the immunogen did not elicit MPER-specific responses, and that further engineering or inclusion of a prior priming immunogen in the immunization regimen would be required to achieve that capacity.

Antigen-specific (WT^{++}) mouse memory B cells were sorted and their BCRs sequenced to assess the robustness of nanodisc probes in the selection of antigen-specific BCRs. Eight antibodies from each animal were randomly selected for monoclonal antibody production. An additional 12 antibodies with long (>=19 AA) CDRH3s were selected and duplicate antibodies (containing identical heavy- and light chains) were removed, resulting in 54 total antibodies. Genes encoding these antibodies were synthesized and 49 out of 54 antibodies were successfully expressed, purified and used in SPR modality B to measure affinities against Env nanodiscs (Fig.

4d). At 1 μM analyte concentration, no non-specific responses to empty nanodiscs were detected.

Out of the tested antibodies, 47% bound to soluble gp140 matching the immunogen ectodomain

and 65% to Env gp151 MPER ND or Env gp151 MPER-KO ND. Considerable binding to

ectodomain, similar binding to MPER targeting immunogen and its MPER KO version and overall

similar geometric mean affinity to tested Env nanodiscs provided evidence that the elicited

antibodies were not targeting the MPER peptide but bound to the exposed base epitope or other

off-target epitopes. Nevertheless, using the nanodiscs as FACS sorting probes and in SPR, we were

able to sort and measure affinities of several antibodies that were specific to the transmembrane

Env gp151 version (e.g. Ab_38, Ab_48, Ab_79. Supplementary Fig. 3b).

Non-human primates (NHPs) are an important model for late-stage pre-clinical vaccine

development and are often used as final validation before clinical trials. Rhesus macaques, like

humans, have extensive genetic diversity within the immunoglobulin loci and are also capable of

making long CDRH3s [44-46]. To test whether nanodisc-based FACS tetramer probes could be

successfully used in RMs, experiments were carried out using PBMCs from animals immunized

previously with soluble Env MD39[47]. PBMCs were stained with the nanodisc probes to determine

whether there was substantial background binding to the lipid component of the nanodisc probes.

Pre-immunization PBMCs showed low background binding of IgD⁻ memory B cells to nanodisc,

while PBMCs from 2-weeks post an MD39 booster immunization showed high binding to the

nanodisc probes (Fig. 5a). Next, the nanodisc probes were compared directly to soluble Env gp140

protein probes in RMs that had been immunized with MD39.3 gp151 mRNA (matching Env gp151

ND sequence) 4-weeks prior[48]. Across two animals there were no differences in the frequency of

tetramer⁺⁺ cells between the Env gp151 ND nanodisc and Env gp140 soluble protein probes (Fig.

5b). Taken together these data show that nanodiscs bearing stabilized HIV Env constructs can be

deployed as tetramer probes for the evaluation of pre-clinical HIV B cell responses in NHPs.

GP nanodiscs in cryo-EM structural studies

279   To demonstrate how structural studies of nanodisc GPs can inform vaccine design, we set

280   out to solve the structure of prototype MPER vaccine candidate Env gp151 MPER ND in complex

with 10E8 antibody by cryo-EM. We included two other HIV bnAbs: VRC01, which recognize the CD4 binding site, and BG18, which targets the N332 glycan supersite. This antibody cocktail complex serves three purposes: First, the added Fabs improve the orientation distribution of particles on cryo-EM grids. Second, they will stabilize the trimer and prevent protomers from dissociating. Lastly, a functional HIV vaccine will likely require elicitation of bnAbs targeting several sites to achieve sterilizing immunity. A trimer boost would therefore benefit from being able to engage three or more sites (i.e. N332, CD4 and MPER) in a single immunogen. The high-resolution structure of the 10E8 epitope has thus far been resolved only with x-ray crystallography and in complex with MPER peptide alone[49-52]. From our cryo-EM data we produced two reconstructions: the complex with two 10E8 Fabs at 4.3 Å, and the complex with a single 10E8 Fab at 3.5 Å resolution, which enable atomic level interpretation. We noted that while 10E8, VRC01 and BG18 were all added at the same step after the nanodisc batch assembly at ~10x molar excess, all VRC01 and BG18 sites were occupied but 10E8 binding site was partially underoccupied in all reconstructions, suggesting steric hinderance of the MPER epitope with increasing 10E8 occupancy (Fig. 6a-c). In all reconstructions, 10E8 is wedged between the Env ectodomain and nanodisc lipid bilayer. MPER antibody induced Env tilt angle was evaluated by aligning the low pass filtered maps to the nanodisc representing the bilayer plane. A single 10E8 Fab appears to be accommodated with minimal Env tilting while binding of a second 10E8 Fab pushes Env into a more tilted position (Fig. 6b). Unlike in our previous WT Env AMC011 nanodisc cryo-EM reconstruction, density for the TM domain remained undetected, presumably due to the R696S mutation disrupting the stabilizing connection point of the three helices in the middle of the bilayer[29]. The single 10E8 class was subjected to 3D variability analysis (3DVA), which revealed distinct conformations of the MPER helix with ~30° lateral twist of the Fab in conjunction with an apparent dislocation of the adjacent protomer (Supplementary Fig. 4c-d. Supplementary video 1).

Using the model of Env gp151 MPER ND complex, we were able to resolve novel structural details of the 10E8 epitope including a binding pocket for 10E8 antibody framework region 3 (FRH3), formed by Env gp120 and gp41 subunits (Fig. 6d, inset 5). We identified 14 AA

in 10E8 contacting the C -terminus of gp120, helices α6 – α8 of gp41 in protomer 1, and L660 of

gp41 in protomer 2. These contacts are either in the binding pocket or in the helices forming a

collar around the termini of gp120. Strikingly, this binding pocket appeared to be targeted by the

CDRH3 of another HIV bnAb 3BC315, that was shown to accelerate trimer dissociation[53]. To

confirm this similarity in engaging the binding pocket, we solved the cryo-EM structure of Env

gp140 (BG505 MD39.3) in complex with BG18, VRC01 and 3BC315 Fabs to 3.1 Å resolution

(Fig 6f, Supplemental Fig. 4f-g). 3BC315 engages the binding pocket mainly with H100A, G100B

and Y100E at the tip of the CDRH3. In 10E8, the interaction is driven by I75, S74 and N76 of

FRH3 (Fig 6d, inset 5). The observed similarity of the binding pocket engagement led us to propose

that these antibodies may share a common mechanism for trimer destabilization. As part of the

binding pocket, both antibodies contact W623 in gp41 (Fig 6d, inset 5 and 6f), which is >99%

conserved across >5,000 diverse HIV strains from the LANL database (http://www.hiv.lanl.gov),

and has previously been shown to be part of a tryptophan clasp contributing to stabilizing four-

helix collar around the termini of gp120[54]. Engaging the binding pocket leads to widening of the

gp41 protomer-protomer gap (Fig. 6g), which would eventually lead to destabilization of the

contacts in the collar around the termini of gp120 and premature triggering of the fusion peptide.

This would normally occur during virus entry associated Env rearrangements that follow CD4

receptor binding. Another novel structural feature of the 10E8 antibody was the dynamic

remodeling of the N-terminal MPER helix, most likely driven by interactions with CDRH1 (D30,

N31 and W33) and CDRH2 (G52C, G53, G54, W55. Fig. 6d-e. Supplementary video 1). Root

mean square deviation (RMSD) of this CDRH1 and CDRH2 10E8 paratope region is 1.52 Å

compared to a published x-ray structure of 10E8 Fab in complex with a gp41 peptide (PDB 4G6F),

indicating that epitope presentation in a more native context recapitulates interactions more

faithfully than epitope peptide alone. In contrast, the RMSD of FRH3 between the nanodisc cryo-

EM complex and peptide-bound crystal structure is 0.46 Å. While the 10E8 crystal structure fit

well into the cryo-EM map (Supplementary Fig. 4e), we calculated an RMSD of 9.59 Å of the

entire MPER domain between the two structures, from the N-terminal portion of the MPER that

is connected to the rest of the ectodomain in the nanodisc Env cryo-EM structure. Taken together,

the comparison to previous crystal structures of 10E8 shows that native membrane environment of Env is required for complete structural characterization of the MPER targeting antibody. Observations of the native, dynamic binding mode of an MPER antibody, and structural information of the entire side chain contact network will inform the development of next-generation MPER targeting vaccines.

Discussion

Here we present a rigorously optimized glycoprotein nanodisc platform for vaccine design. We show that the assembled GP nanodiscs enable characterization of transmembrane glycoprotein vaccine candidates with key methods in rational, iterative vaccine development that have traditionally been available only to soluble mimics. Applicability of the platform is demonstrated by using SPR to measure antibody kinetics, FACS to characterize new vaccine designs and in vivo antibody responses from preclinical experiments, and cryo-EM to provide structural guidance for immunogen design.

In addition to confirming the assembly of EBOV GP in nanodiscs with two Ebola-specific antibodies, we characterized the kinetic landscape of several HIV antibodies binding to HIV Env in nanodiscs. These confirmed that glycan shield modifications and transmembrane domain point mutation identified through directed evolution increased MPER-targeting bnAb 10E8 affinity to prototypical HIV Env MPER-targeting immunogen Env gp151 MPER ND as compared to its design predecessor. While the in vivo responses to the engineered immunogen in humanized mouse model appeared to be outside the targeted C-terminal region of MPER, the identified antibodies will provide guidance for further immunogen engineering. Potential targets of the extracted antibodies include the engineered glycan hole or off-target epitopes at the trimer base. Base of the trimer is partly occluded by the membrane, but some off-target epitopes may still be accessible. This was also indicated by EM-based polyclonal epitope mapping (EMPEM) showing that transmembrane gp151 Env administered as mRNA still elicits base-binding antibodies[15,48]. Importantly, using the biotinylated GP nanodisc as FACS probe, we were able to extract antibodies targeting sites that are not fully present in the soluble form of the immunogen. While NHP

immunization was beyond the study's scope, we also confirmed GP nanodiscs can be used to identify antigen-specific responses from NHPs, facilitating future studies. Two promising approaches for overcoming the challenges in eliciting MPER targeting antibodies have been recently described, namely priming with germline-targeting immunogen 10E8-GT12[24, 55] or immunizing with MPER peptide alone in liposome formulation[25]. Both solutions present a more accessible MPER peptide for antibody CDRH3. These immunogens can be followed by a next generation Env boost immunogen designs presenting the complete epitope to guide the maturing antibody toward breadth and higher neutralization potency. The analysis and nanodisc GP FACS methods presented here provide vital guidance for these designs.

The platform enabled us to determine the structure of Env gp151 MPER ND to 3.5 Å, leading to discovery of structural features that will inform the next generation of immunogens. The structure confirms three HIV bnAb binding sites targeted in ongoing vaccine development projects can be engaged by the same immunogen[56,24, 44]. This further demonstrated that the nanodisc platform has the potential to be used in cryoEMPEM to resolve structures of membrane proximal antibody epitopes from a mixture of specificities from post-immunization serum[57]. As multiple epitopes are now being engineered into Env trimers for germline targeting, it is critical to understand if and how they may compete or synergize. Additional value in using GP nanodiscs is that, in contrast to previous structural studies using soluble mimetics of Env[40, 58], this nanodisc-based structure resembles more closely the native transmembrane protein immunogen[48], which may be important i.e. when assessing glycan engagement in the epitope. Glycan analysis of Env Env gp151 ND and Env gp151 MPER ND agreed with earlier studies showing that when Env is expressed as transmembrane protein, glycan processing resembles more closely that of the virus particle with higher proportion of complex type glycans as in soluble forms[26, 27]. Unexpectedly, germline targeting Env immunogen N332 GT5 deviated from the trend by displaying nearly identical glycan composition in soluble and transmembrane versions of the immunogen. The most significant value of the Env gp151 MPER ND complex cryo-EM structure however is in characterizing the MPER epitope in unprecedented detail. We were able to resolve contacts of 10E8 bnAb with the Env ectodomain that have thus far remained elusive in structural studies. A

binding pocket for 10E8 FRH3 in the gp120-gp41 interface of the ectodomain, and CDRH1, 2 and

3 contacts to the entire MPER domain ending at the lipid bilayer interface, form a continuous

network of side chain interactions together spanning 29 AA in the paratope and 26 AA in the

epitope. These details can now be used as a structural template to design immunogens for improved

engagement of MPER antibodies. Finally, 3D variability analysis of the cryo-EM data illustrates

significant remodeling of the N-terminal MPER helix induced by CDRH1 and CDRH2 of 10E8.

While care should be taken when interpreting flexible regions, this captures detailed snapshots of

the native flexibility of the glycoprotein and MPER domain. The data depicts the range of

flexibility and may offer clues how to accommodate native glycoprotein flexibility in vaccine

design.

In conclusion, GP nanodisc platform provides a scalable and reproducible solution for a

more complete and native-like environment for transmembrane glycoprotein vaccine design. This

platform can therefore accelerate the development of next-generation transmembrane GP based

vaccines.

Methods

408 **Methods**

409 This method description aims to provide a complete platform that enables routine use of nanodisc

410 approach for iterative vaccine design using viral glycoproteins. While retaining the versatility of a

411 modular approach, each step is simplified and validated for maximizing reproducibility and

412 scalability.

Protein expression and nanodisc assembly

Protein expression and nanodisc assembly

All glycoprotein and antibody constructs were codon optimized for expression in human cells and

synthesized and cloned into expression vectors pHLSec (glycoproteins) or its variants pCW-CHIg-

hG1, pFabCW pCW-CLig-hL2 or pCW-CLig-hk (IgG heavy chains, Fab heavy chains, lambda

light chains and kappa light chains, respectively) by GenScript Biotech. Antibodies were expressed

and purified from HEK293F or ExpiCHO cells according to manufacturer's instruction using

rProtein A Sepharose Fast Flow resin (Cytiva) as described earlier[24, 44].

Soluble Env GP were purified as described earlier[40]. Transmembrane glycoprotein purification method was modified from the earlier protocol[29]. For one liter of transfected cells, 1 mg of DNA was mixed with 25 ml of OPTI-MEM medium and sterile filtrated through 0.22 μm filter unit. 3 mg of PEI MAX (Polysciences, #24765) was separately mixed with 25 ml of OPTI-MEM, filtrated and then combined with the DNA. Combined DNA and PEI MAX were then added to HEK293F cells at 1 million/ml density and cells were incubated at 37 °C, 8% CO_2, and 80% humidity under shaking at 180 rpm for 3 days. Cells were harvested by centrifugation at 1500 rcf in 4 °C for 15 min, washed with ~300 ml cold PBS, and centrifuged again at 1500 rcf in 4 °C for 15 min. Pelleted cells were lysed with ~5 ml of lysis buffer per gram of cells (50 mM Tris-HCl (pH 7.4), 300 mM NaCl, 0.5% TX-100). Prior to use, approximately 1 protease inhibitor cocktail tablet per 50 ml of lysis buffer, and 2 ml of BioLock (IBA Lifesciences, # 2-0205-050) per liter of cells was added to lysis buffer. Cells were lysed for 1 h at 4 °C in an overhead rotating mixer. Lysed cells were centrifuged for 1 h at 25,000 rcf, followed by filtration of the supernatant through 0.22 μm bottle top filtration unit. This resulted in cleared lysate that was passed through Strep-tactin XT 4Flow matrix (IBA Lifesciences, #2-5010-025) in a gravity flow column. Approx. 400 μL of drained matrix was used per liter of cells. Matrix was then washed three times with 2 ml of wash buffer 1 (50 mM Tris-HCl (pH 7.4), 300 mM NaCl, 0.1% (w/v) CHAPS, 0.03 mg/mL deoxycholate), three times with 2 ml of wash buffer 2 (50 mM Tris/HCl pH 7.4, 500 mM NaCl, 1 mM EDTA, 0.1% DDM, 0.01% CHS, 0.03 mg/ml deoxycholate) and three times 2 ml of wash buffer 3 (50 mM Tris/HCl, pH 7.4, 150 mM NaCl, 0.02% DDM, 0.002% CHS, 0.03 mg/ml deoxycholate). After this, matrix with detergent solubilized GP was transferred to a 5 ml test tube for nanodisc assembly step.

Lipid stocks were prepared prior to disc assembly in 1 mM final concentrations. Premade 10% DDM/ 0.1% CHS stock (Anatrace, #D310-CH210) was used throughout the study. All lipids in this study were prepared from chloroform stocks (Avanti Polar Lipids). Chloroform was first evaporated under gentle nitrogen stream until transparent film was formed around the glass vial. Lipid film was then rehydrated for 1 h in room temperature in lipid rehydration buffer (50 mM Tris/HCl, pH 7.4, 150 mM NaCl, 0.1% DDM, 0.01% CHS) in a volume bringing the lipid stock

concentration to 5 mM. After thorough mixing by vortexing, this stock was diluted further to final lipid stock solution at 1 mM concentration and sonicated with microtip (20-25% intensity, 50% time cycles in 4 °C) until clear. Most lipid stock solutions cleared within 10 – 20 min but with some lipids DDM/CHS amount was increased up to 1%/0.01%. 1 mM lipid stocks were aliquoted and stored up to 6 months at -20°C. A lipid mix roughly following HIV particle lipid composition was used throughout the study and contained 45% DOPC, 7% DOPS, 7% DOPA, 7% DOPE, 7% DOPG, 5% sphingomyelin, 2% PI(4,5)P2, and 20% CHS. Scaffold protein was either produced in-house using standard *E. coli* expression method or purchased from Sigma-Aldrich (membrane scaffold protein 1D1 #M6574 or biotinylated membrane scaffold protein 1D1 BTN #MSP13).

Nanodisc batch assembly was done as follows. Strep-tactin matrix with GP bound was drained by gravity flow, after which gravity flow column was capped and 400 µl of wash buffer 3 was added to the matrix. 50% matrix slurry was then transferred to a 5 ml test tube. Lipids were added at approx. 350X molar excess and scaffold at 6X molar excess in relation to the glycoprotein. Glycoprotein amount bound to the matrix was estimated with a separate initial experiment where protein was eluted after the last wash step. Minor variations were made to glycoprotein:scaffold and glycoprotein:lipid ratios without significant difference in the final yield. Matrix with glycoprotein, lipids and scaffold was incubated for 1 h at 4°C prior to addition of SM-2 bio-beads (Bio-Rad, #1523920). Bio-beads were activated according to the manufacturer's instructions in methanol followed by extensive wash with deionized water. Bio-beads were added to approx. 50% v/v with the nanodisc assembly mixture. i.e. if the total volume of the assembly reaction was 1.5 ml, final volume after bio-bead addition in the 5 ml test tube was ~3 ml. Detergent was then removed by incubation in a rotating shaker overnight at 4°C. The following day an additional ~20% of the total reaction volume of fresh bio-beads were added, followed by an additional 1 h incubation at room temperature to ensure complete removal of detergent. Next, contents of the test tube were transferred to a new gravity flow column with a pipette tip with the first ~2 mm cut off to allow pipetting biobeads and matrix. Mixture was washed in column twice with 4 ml of TBS and eluted with Strep-tactin elution buffer BXT in 500 µl fractions. A280 of eluted fractions were monitored and fractions containing protein were pooled and concentrated to <500 µl for SEC

performed using a Superose 6 Increase 10/300 column (Cytiva) at 0.75ml/min in TBS. SDS-PAGE

and ns EM analysis were used to confirm presence of glycoprotein and scaffold protein, typical

disc appearance in ns EM 2D class averages and glycoprotein on disc.

For structural biology purposes, Fabs were added to the GP nanodiscs after the assembly

while nanodisc was still bound to strep-tactin matrix. After the second addition of bio-beads, each

Fab was added at ~10X molar excess and the mixture was incubated for an additional 2h at room

temperature. Complex elution and following steps were identical as described above for

unliganded glycoprotein nanodiscs. Fractions were pooled after SEC and sample was concentrated

either to 1 mg/ml for FACS, SPR and other purposes, or up to 11 mg/ml for cryo-EM sample

preparation.

Identification of R696S mutation on HIV Env by mammalian directed evolution.

A 250 amino acid long site-saturation mutagenesis (NNK) scan of gp41 was synthesized at SGI-

DNA. A fragment of the library containing NNKs from positions T606 to F699 (94 positions) was

cloned into BG505 MD39.2 gp151. This library was stably integrated into 293T-rtTA3G cell line

using the lentivirus-based mammalian display protocol described previously[34, 40,35]. After inducing

library expression overnight, the cells were stained with 10E8 IgG and PGT145 Fab and those

with highest binding to 10E8 that were also positive for PGT145 were sorted. After three rounds

of sorting, the library DNA was extracted and sequenced by Sanger sequencing. 8 of 19 sequences

had a mutation at position 696, which was mutated from R to S, A, V, I and L. The most frequent

mutation, R696S, was selected for further characterization.

Mass spectrometry

Proteinase K treatment and deglycosylation. HIV Env glycoprotein (when membrane bound,

denatured in 6 M urea) was exchanged to water using Microcon Ultracel PL-10 centrifugal filter.

Glycoprotein was reduced with 5 mM tris(2-carboxyethyl)phosphine hydrochloride (TCEP-HCl)

and alkylated with 10 mM 2-Chloroacetamide in 100 mM ammonium acetate for 20 min at room

temperature (RT, 24 °C). Initial protein-level deglycosylation was performed using 250 U of Endo

H for 5 µg trimer, for 1 h at 37 °C. Glycoprotein was digested with 1:25 Proteinase K (PK) for 30

min at 37 °C. PK was denatured by incubating at 90 °C for 15 min, then cooled to RT. Peptides

were deglycosylated again with 250 U Endo H for 1 h at 37 °C, then frozen at –80 °C and

lyophilized. 100 U PNGase F was lyophilized, resuspended in 20 µl 100 mM ammonium

bicarbonate prepared in $H_2^{18}O$, and added to the lyophilized peptides. Reactions were then

incubated for 1 h at 37 °C, subsequently analyzed by LC-MS/MS.

LC-MS/MS. Samples were analyzed on an Q Exactive HF-X mass spectrometer. Samples were

injected directly onto a 25 cm, 100 µm ID column packed with BEH 1.7 µm C18 resin. Samples

were separated at a flow rate of 300 nL/min on an EASY-nLC 1200 UHPLC. Buffers A and B

were 0.1% formic acid in 5% and 80% acetonitrile, respectively. The following gradient was used:

1–25% B over 160 min, an increase to 40% B over 40 min, an increase to 90% B over another 10

min and 30 min at 90% B for a total run time of 240 min. Column was re-equilibrated with solution

A prior to the injection of sample. Peptides were eluted from the tip of the column and nanosprayed

directly into the mass spectrometer by application of 2.8 kV at the back of the column. The mass

spectrometer was operated in a data dependent mode. Full MS1 scans were collected in the

Orbitrap at 120,000 resolution. The ten most abundant ions per scan were selected for HCD

MS/MS at 25 NCE. Dynamic exclusion was enabled with exclusion duration of 10 s and singly

charged ions were excluded.

Data Processing. Protein and peptide identification were done with Integrated Proteomics

Pipeline (IP2). Tandem mass spectra were extracted from raw files using RawConverter [59] and

searched with ProLuCID[60] against a database comprising UniProt reviewed (Swiss-Prot) proteome

for Homo sapiens (UP000005640), UniProt amino acid sequences for Endo H (P04067), PNGase

F (Q9XBM8), and Proteinase K (P06873), amino acid sequences for the examined proteins, and a

list of general protein contaminants. The search space included no cleavage-specificity.

Carbamidomethylation (+57.02146 C) was considered a static modification. Deamidation in

presence of $H_2^{18}O$ (+2.988261 N), GlcNAc (+203.079373 N), oxidation (+15.994915 M) and N-

terminal pyroglutamate formation (−17.026549 Q) were considered differential modifications.
Data was searched with 50 ppm precursor ion tolerance and 50 ppm fragment ion tolerance.
Identified proteins were filtered using DTASelect2[61] and utilizing a target-decoy database search
strategy to limit the false discovery rate to 1%, at the spectrum level[62]. A minimum of 1 peptide
per protein and no tryptic end per peptide were required and precursor delta mass cut-off was fixed
at 15 ppm. Statistical models for peptide mass modification (modstat) were applied. Census2[63]
label-free analysis was performed based on the precursor peak area, with a 15 ppm precursor mass
tolerance and 0.1 min retention time tolerance. "Match between runs" was used to find missing
peptides between runs. Data analysis using GlycoMSQuant [42] was implemented to automate the
analysis. GlycoMSQuant summed precursor peak areas across replicates, discarded peptides
without NGS, discarded misidentified peptides when N-glycan remnant-mass modifications were
localized to non-NGS asparagines and corrected/fixed N-glycan mislocalization where
appropriate.
546

SPR
The kinetics and binding affinities of antibody-antigen interactions were analyzed using either
ProteOn XPR36 system (Bio-Rad) or Carterra LSA. TBS pH 7.4 (20 mM Tris, 150 mM NaCl)
supplemented with BSA at 1 mg/ml without detergents was used as running buffer in all
experiments. In ProteOn XPR36, HC30M XanTec sensor chips were utilized. In modalities B and
C, anti-human IgG (Fc) antibody (GE, #BR-1008-39) was used for capturing IgG (ligand) at low
densities with nanodisc GP as the analyte. In modality A, anti-Strep-tag antibody (pAb, rabbit,
GenScript, #A00875) was used for capturing Strep-tagged nanodiscs GP (ligand) followed by GP
specific Fab as the analyte. Approximately 6000-8000 response units (RU) of the capture antibody
were covalently attached to the sensor surface using EDC-NHS chemistry. For IgG-antigen
interaction studies, approximately 50 to 100 RUs of IgGs at a concentration of 0.3 μg/ml were
immobilized on each flow cell. For Fab-antigen interaction studies, approximately 300 to 400 RUs
of antigen at 10 μg/ml were captured on each flow cell. Analytes were introduced to the flow cell
at a rate of 30 μl/min for 3 minutes, followed by a dissociation phase of 5 minutes. Regeneration

was performed with 1.7% or 0.85% phosphoric acid, each with a contact time of 60-180 seconds,

repeated four times per cycle. Data analysis was conducted using ProteOn Manager software (Bio-

Rad), including raw sensogram processing, interspot referencing, and column double referencing.

Equilibrium or kinetic fits were performed using the Langmuir model as needed.

Kinetics and affinity of antibody-antigen interactions on Carterra LSA using CMDP Sensor Chip

(Carterra) for capture IgG – flow nanodisc (modality B) was done as follows. Chip surfaces were

prepared according to Carterra software instructions. In a typical experiment approx. 500-700 RU

of capture antibody (SouthernBiotech, # 2047-01) in 10 mM Sodium Acetate pH 4.5 was amine

coupled on CMDP chip placing special care on concentration range of the amine coupling reagents.

We used N-Hydroxysuccinimide (NHS) and 1-Ethyl-3-(3-dimethylaminopropyl)

carbodiimidehydrochloride (EDC) from Amine Coupling Kit (GE, #BR-1000-50). Highest

coupling levels of capture antibody were achieved by using 10 times diluted NHS and EDC during

surface preparation runs as compared to kit instruction (10 ml of water each to give 11.5 mg/ml

and 75 mg/ml respectively according to kit instructions). Thus, the concentrations of NHS and

EDC were 1.15 mg/ml and 7.5 mg/ml and the activation time was reduced to 1 min. The

concentrated stocks of NHS and EDC were stored frozen in -20 °C for up to 2 months without

noticeable loss of activity. The capture antibody was used at concentration 25 µg/ml with 10

minutes contact time. Phosphoric Acid 1.7% was used as a regeneration solution with 60 seconds

contact time and injected three times per cycle. Concentration of ligands were approximately 1

µg/ml and contact time was 5 min. Raw sensograms were analyzed using Kinetics software

(Carterra) with interspot and blank double referencing, and Langmuir model. Analyte

concentrations were quantified on NanoDrop 2000c spectrophotometer using absorption signal at

280 nm. Analytes were buffer exchanged into the running buffer using dialysis. In a typical run,

we covered a broad range of affinities and used two referencing practices depending on the off-

rate of the ligand. For fast off-rate (faster than 9×10^{-3} 1/s), we use automated batch referencing

that includes overlay y-aline and higher analyte concentrations. For slow off-rates (9×10^{-3} 1/s or

less), we use manual process referencing that includes serial y-align and lower analyte

concentrations. After automated data analysis by Kinetics software, we performed additional
filtering to remove datasets with highest response signals smaller than signals from negative
controls using R-script.

FACS

Validation of Env-Nanodisc proteins as fluorescent baits

Streptavidin (SA) conjugated-antigen baits were prepared by combining biotinylated Env-
nanodisc or empty nanodisc with fluorescent SA at room temperature for at least 1 hr in dark at a
bait:SA molar ratio of 2:1. Control breads for FACS were generated by conjugating various bnAbs
to compensation beads. Mouse anti-human IgG (BD Biosciences, # 555784) was first captured on
to anti-mouse IgK compensation beads (BD Biosciences, # 552843), followed by a wash step and
a secondary capture of bnAbs of interest. VRC01-expressing Ramos cells [43], C57BL/6 splenocytes,
and the prepared bnAb-conjugated beads were incubated with the nanodisc GP:SA baits at a bait
concentration of 10-50 nM in the final staining volume for 30 min at 4 °C in the dark. Data were
acquired on a BD FACS Melody, and analzyed in FlowJo v10 (FlowJo, LLC).

Mouse immunization studies

hD3-3/JH6 mice were immunized as previously described[24]. Briefly, homozygous hD3-3/JH6
mice were injected with 10 µg (50 µl total volume) of Moderna mRNA LNPs encoding Env gp151
MPER I.M. under anesthesia (5% isoflurane induction) in the left quadriceps muscle. After six
weeks, mice were euthanized with compressed CO_2 (100%) in a clear chamber to allow for
visualization of respiration and subsequent death via respiratory cessation. Blood was collected
from the chest cavity prior to the removal of the spleen and lymph nodes (mesenteric, inguinal,
and popliteal). Tissues were placed in 3 mL FACS buffer (1 X PBS Ca/Mg^{++} free, 1 mM EDTA,
25 mM HEPES, pH 7.0, 1% heat-inactivated FBS) on ice and tissues were disassociated using the
rough ends of two sandblasted microscope slides, followed by centrifugation (460 rcf for 5 minutes
at 4 °C). Red blood cell lysis was performed using 1 mL of ACK buffer (Quality Biological) for 2
minutes on ice before lysis was halted by adding 14 mL FACS buffer per sample. Post lysis and

centrifugation (460 xg for 5 minutes), cells were resuspended in 3 mL Bambanker freezing

medium (Bulldog Bio) prior to filtration through a cotton-plugged, borosilicate pasteur pipette into

a borosilicate glass test tube. 1 mL filtered-cell solution was subsequently divided into three

cryovials/mouse, which were precooled in a styrofoam rack on dry ice. Cells were stored at -80 °C

for 2-7 days prior to long-term storage in liquid nitrogen. All work followed IACUC guidelines

associated with animal protocol number 20-0001.

Mouse sample preparation and B cell sorting

Mouse frozen mouse splenoctyes and lymphocytes were thawed and stained as previously

described[24, 56]. Briefly, after thawing and counting, total B cells were isolated by negative selection

using the EasySep Mouse Pan-B Cell Isolation Kit (StemCell, #19844) according to

manufacturer's instructions. SA-conjugated baits were prepared as described above. Wildtype

antigen baits were conjugated to BV421-SA (BioLegend, #405225) and AlexaFluor 647SA

(Invitrogen, #S21374), while epitope knockout (KO) baits were conjugated to hashtagged

TotalSeq-C PE SA (BioLegend, #405261). Purified B cells were stained at 4 °C in dark, with

appropriate baits and antibody master-mix consisting of FITC anti-CD19 (BioLegend, #152404),

BV786 anti-IgM (BD Biosciences, #743328), PerCP-Cy5.5 anti-IgD (BD Biosciences, #564273),

APC-Cy7 anti-F4/80 (BioLegend, #123118), APC-Cy7 anti-CD11c (BD Biosciences, #561241),

APC-Cy7 anti-Ly6C (BD Biosciences, #557661), APC-H7 anti-CD8a (BD Biosciences,

#560182), and APC-H7 anti-CD4 (BD Biosciences, #560181), all used at 1:100 dilution and

prepared in FACS buffer (1% v/v heat inactivated FBS, 1 mM EDTA, 1 mM HEPES in 1x DPBS).

The KO bait was first added to cells with the antibody master mix for 15 minutes, followed by the

addition of WT baits for an additional 30 min. Each bait was used at final bait concentration of

100 nM in the staining volume or 0.5 µg per sample. During the addition of antibody master mix,

a unique TotalSeq-C anti-mouse hashtag antibody (BioLegend) was added to each sample at a

concentration of 2.5 µL / up to 20 million cells. Following antibody staining, 1 mL of 1:300

live/dead stain diluted in FACS buffer (Invitrogen, #L34966) was added to each sample for an

additional 15 minutes at 4 °C, then washed with 10 mL of FACS buffer. Cells were resuspended

in a final volume of 500 µL FACS buffer and sorted on a BD FACS Melody. Cells were bulk

sorted into a 4 °C chilled PCR plate well containing 20 µL of 0.2 µm filtered FBS, and processed

for BCR sequencing via the 10x Genomics Single Cell Immune Profiling according to kit

protocols, with minor modifications outlined previously[64]. Pooled libraries were sequenced on an

Illumina NextSeq 2000 using a 100-cycle P3 reagent kit (Illumina, #20040559).

BCR sequence analysis

Sequence analysis Raw sequencing data were demultiplexed, processed into assembled VDJ

contigs and counts matrix files, and assigned to specific animal IDs based on TotalSeq-C antibody

hashtag counts using Cell Ranger (v6.1) and scab as previously described[64]. Gene assignment,

annotation, and formatting into Adaptive Immune Receptor Repertoire (AIRR) format[65] for paired

heavy and light chain antibody sequences was performed using Sequencing Analysis and Data

library for Immunoinformatics Exploration (SADIE) with a previously described custom germline

reference database that included all known mouse antibody genes plus the human IGHD3-3 and

IGHJ6 genes that were knocked into hD3-3/JH6 mice[24]. Eight antibodies from each animal were

randomly selected for in vitro analysis, except for one animal from which only two total

heavy/light pairs were recovered. In addition, 12 antibodies with long HCDR3s (>19 aa) were

selected for in vitro characterization.

NHP Flow Cytometry

Frozen NHP PBMC samples were thawed in RPMI media with 10% FBS, 1% GlutaMAX,

and 1% Penicillin/Streptomycin (R10). NHP samples used in this study were from animals that

were reported in previous studies [47,48]. The recovered cells were then stained with the appropriate

antibody panel. Fluorescent antigen probes were constructed by combining fluorophore-

conjugated streptavidin and biotinylated nanodisc GP or soluble Env probes with an appropriate

volume of PBS in small increments across 45 minutes at room temperature (RT). Thawed cells

were incubated with probes for thirty minutes at 4°C. A master mix of surface staining antibodies

was then added and incubated for thirty minutes at 4°C. Cells were washed and prepared for

acquisition on Cytek Aurora (Cytek Biosciences). The following reagents were used in the

PBMC staining panel: Alexa Fluor 647 Streptavidin (BioLegend #405237), BV421 Streptavidin

(BioLegend #405225), PE Streptavidin (BioLegend #405245), PE-Cy7 Streptavidin (BioLegend

#405206), BV711 Streptavidin (BioLegend #405241), BUV615 Streptavidin (BioLegend

#613013), Viability eFluor780 (1:2000, Invitrogen #65-0865-14), CD3 BV510 (1:100, BD

Biosciences #569488), CD14 BV510 (1:100, BioLegend #301842), CD8a BV510 (1:100, BD

Biosciences #563256), CD16 BV510 (1:100, BioLegend #302048), CD20 (1:100, BUV395 BD

Biosciences #563782), IgG BV605 (1:100, BD Biosciences #563246), and IgD AF488 (1:50,

Southern Biotech #2030-30).

EM

Negative stain EM was used for quality control of nanodisc assembly and for initial assessment of

antibody complexes for cryo-EM. Glycoprotein nanodisc or nanodisc with three fold molar excess

of Fab was applied to 400 mesh size Cu grid at 0.04-0.06 mg/ml concentration, blotted off with

filter paper and stained with 2% uranyl formate for 60 s. NsEM data was collected on a Tecnai

Spirit microscope operating at 120 keV, using a Tietz 4k × 4k TemCam-F416 CMOS camera and

Leginon automated image collection software[66]. Data was processed using Relion 3.0 image

processing pipeline[67].

For cryo-EM, nanodisc was complexed with 10X molar excess of VRC01, BG18 and 10E8 Fabs

after batch nanodisc assembly, while bound to affinity matrix. Eluted complex was then SEC

purified (Superose 6 Increase column. Cytiva), concentrated to 8 mg/ml and frozen on graphene

oxide grids (GO on Quantifoils R1.2/1.3, Cu, 400 mesh, Electron Microscopy Sciences) with 30

sec wait time, blot force of 0, and blot time of 2.5-3 s. Fluorinated Fos-Choline-8 (Anatrace #

F300F) was added to sample immediately prior to freezing to a final concentration of 3 mM. GO

grids, high sample concentration and fluorinated Fos-Choline-8 were necessary to prevent freeze-

denaturation, ensure high particle count per micrograph and prevent orientation bias. 11,871

micrographs were collected at pixel size 0.718 Å using a Thermo Fisher Scientific Glacios 2

microscope operating at 200 kV and equipped with a Thermo Fisher Scientific Falcon 4i camera

using a total dose of 44.89 e-/Å2. Automated data collection was performed using the EPU software (Thermo Fisher Scientific) and images were written in the EER frame format. Micrographs were preprocessed using cryoSPARC Live[66], including motion and CTF correction. Briefly, particles were picked with Topaz[67] after training the neural network with a subset of particles first picked with blob picker. Picked particles were subjected to 3 rounds of 2D classification to obtain a final particle stack of 317,792 particles. These were then extracted from micrographs using a box size of 540 pix. Maps containing one or two copies of 10E8 Fab were separated using heterogenous reconstruction and 3D classification tasks. A class with 7,964 particles containing two copies of 10E8 was refined to 4.3Å resolution. A class with 111,002 particles containing single 10E8 was examined with 3D variability analysis[68]. Further classification resulted in final, cleaned particle stack containing 63,366 particles which was then subjected to Non-Uniform Refinement[69] and used for atomic model building. Data collection and processing stats are summarized in Supplementary Table 1. Atomic model building was initiated by docking in available structures of 10E8 Fab (PDB 4G6F), BG18 Fab (PDB 6DFG), VRC01 Fab (PDB 3NGB) and an AlphaFold3[70] -generated model of the trimer into the map using ChimeraX[71]. Manual adjustment was done using Coot[72] and further refinement was done using Phenix Real Space Refine[73] and Rosetta Relax[74]. Final model statistics are summarized in Supplementary Table 1.

The soluble gp140 (MD39.3) cryo-EM structure was done as follows: VRC01, BG18 and 3BC315 Fabs were added at 4.3X molar excess to purified Env glycoprotein. The complex was purified using a HiLoad Superdex 200 16/600 pg column (Cytiva) and concentrated to 10 mg/ml. Samples were vitrified using a Thermo Fisher Scientific Vitrobot Mark IV operating at 100% humidity, 4°C and 3-6 s blot times, on UltrAuFoil 1.2/1.3-300 holey gold film grids (Electron Microscopy Sciences). Prior to grid application, lauryl maltose neopentyl glycol (LMNG) was added to a final concentration of 0.005 mM. Imaging was performed using the same microscope and conditions as for Env gp151 MPER ND complex. After initial cryoSPARC particle picking using Blob Picker, templates were selected from 2D classification and used for Template Picker. A total of 886,907 particles were extracted using a box size of 576 downsampled to 144. After

additional rounds of 2D classification, an Ab Initio Reconstruction was performed, followed by
Non-Uniform refinement. A single 3BC315 Fab was visible in the C1 reconstruction. A mask was
created over 3BC315 and focused 3D classification performed to remove trimers with no 3BC315
particles. A final stack of 243,503 were re-extracted and downsampled to box size 400 (resulting
in pixel size of 1.034 Å) and subjected to Non-Uniform Refinement which resulted in a 3.1 Å
reconstruction. Model building was performed as above and final model statistics are summarized
in Supplementary Table 1.

Maps and models have been deposited to the EMDB and PDB, respectively, under the
accession codes listed in Supplementary Table 1.

Acknowledgements
We thank Nicole Doria-Rose for providing the VRC01 expressing Ramos cell line. We thank Brian
Briney for providing access to Illumina NextSeq 2000 sequencer. We thank Daniel Murin for
providing Ebola GP specific antibodies.

Funding
This work was supported by National Institute of Allergy and Infectious Diseases UM1 AI144462
(Scripps Consortium for HIV/AIDS Vaccine Development; to J.C.P., T.S., A.B.W., and W.R.S),
R01 AI147826 (to W.R.S.); Bill and Melinda Gates Foundation Collaboration for AIDS Vaccine
Discovery awards (INV-007522 and INV-008813 for the IAVI NAC Center to A.B.W. and
W.R.S.; and INV-002916 to A.B.W.); the IAVI Neutralizing Antibody Center (to A.B.W. and
W.R.S.); and the Alexander von Humboldt Foundation (T.S.).

Competing interests: J.M.S and W.R.S. are inventors on a patent for the BG505 MD39 and N332-
GT5 immunogens (US11203617B2 and US20230190914A1). S.H. and W.R.S. are employees and
shareholders of Moderna, Inc. All other authors declare no competing interests.

Author contributions: K.R. designed nanodisc assembly workflow, performed cryo-EM imaging, data processing and structural analysis, and wrote the manuscript. A.L. and O.K. established conditions and conducted SPR assays. G.O. Supervised cryo-EM imaging and data processing and built 3D models. C.F. performed and J.H.L. and D.S. supervised establishing conditions for mouse B cell FACS experiments. J.M.S., O.S. and T.S. contributed to immunogen design. S.B. and J.K.D. performed, and J.R.Y. and J.C.P. supervised mass spectrometry glycan profiling. P.J.M. and M.S. performed, and S.C. supervised NHP B cell FACS analysis. S.W. and A.G. contributed, and G.O. and A.B.W. supervised cryo-EM imaging, data processing and model building. D.L. performed nanodisc assembly reactions. P.K. and S.T. performed mouse immunizations. D.L., E.G., R.T., S.E., N.A., D.G. and M.K. produced purified proteins. W-H.L. contributed to cryo-EM sample preparation. S.H. provided mRNA immunogens. T.S., A.B.W. and W.R.S. supervised the study. G.O., T.S., A.B.W. and W.R.S. edited the manuscript. All authors reviewed the manuscript.

Figure legends

Figure 1. Nanodisc use in different steps in iterative rational vaccine design. Analytical methods that can directly use nanodiscs as sample material are highlighted.

Figure 2. HIV Env constructs used for method development and general overview of the nanodisc assembly workflow. a) Naming and gene organization of the glycoprotein constructs used for nanodisc assembly. Introduced mutations and intracellular elements are indicated. ND refers to constructs made for nanodisc platform. b) Schematic overview of HIV Env constructs and MPER targeting antibody 10E8 binding. c) Key steps of the 5-day workflow for glycoprotein nanodisc assembly. Glycoprotein was extracted with detergent from cell surface and bound to affinity matrix. Disc assembly was performed "in-batch" while bound to affinity purification matrix. Assembled discs were then eluted for final SEC polishing purification, followed by quality control steps before subjecting to final analytical methods (SPR, FACS, cryo-EM).

Figure 3. Different SPR modalities and kinetic analyses of GP nanodiscs. a) Schematic illustrations of three SPR modalities used throughout the study and their strengths and weaknesses. b) Affinity of MPER targeting 10E8 antibody to Env gp151 ND as measured with modalities A and B. Affinities with modality A were measured after storing the sample either 1 week (white), 1 month (grey) or 3 months (black) at 4°C. c) Affinity of 10E8 against Env constructs using modality A. d) SPR data presented in (c), showing increase in affinity originating from reduced off-rate in Env gp151 MPER ND. e) Modality C was used to scout Ebola specific EBOV-296 and EBOV-13C6 IgG epitope accessibility in EBOV GP nanodiscs. Data shows binding of two anti -Ebola antibodies targeting the glycan cap (EBOV-296 and 13C6). Env gp151 ND and anti-HIV MPER antibody PGZL1 were used as a negative controls.

Figure 4. HIV Env nanodisc FACS probe validation and pilot use in pre-clinical mouse model. a) Nanodiscs were tested for binding to HIV MPER bnAbs 10E8 and DH511 and gp120-specific

bnAb BG18 coupled to FACS compensation beads, B6 mouse splenocytes were used as negative

control cells, and VRC01 expressing Ramos cells as positive control cells. Empty nanodiscs were

assembled with either DOPC lipids (blue) or a mixture of neutral and charged lipids (green). b)

Memory (CD19$^+$IgM$^-$IgD$^-$) and naïve (CD19$^+$IgM$^+$IgD$^+$) B cells from mice 6 weeks after

immunization with mRNA-LNPs encoding Env gp151 MPER ND nanodisc were analyzed by flow

cytometry. Antigen-specific (Env gp151 MPER ND^{++}) and epitope-specific (KO$^-$) B cells were

detected using immunogen-matched WT and 10E8-epitope KO nanodisc probes and WT

nanodisc^{++} cells were sorted for BCR sequencing. The complete gating strategy is shown in

supplementary Fig. 3. c) Percent antigen-specific memory B cells in the naïve vs. memory

compartments of each immunized mouse (top), and proportion of 10E8 epitope-specific (KO$^-$)

cells within each antigen-specific population. Paired t-test, *p < 0.05, ns: p> 0.05. d) Sorted cells

were sequenced and selected antibodies were purified for affinity measurement by SPR using

modality B.

Figure 5. HIV Env nanodisc probe validation for sorting NHP cells. a) Flow cytometry plots

showing CD20$^+$IgD$^-$ memory B cells stained with Env gp151 ND nanodisc on two different

fluorophores. Pre-immunization and 2-weeks post-immunization PBMCs from the same RM are

shown and the frequency of Nanodisc^{++} cells of IgD$^-$ cells is listed. b) Flow cytometry plots

showing CD20$^+$IgG$^+$ memory B cells stained with either soluble Env gp140 protein or Env gp151

ND nanodisc on two different fluorophores. PBMCs from two different RMs 4-weeks post mRNA

Env gp151 immunization were split and stained with Env gp140 protein tetramers (left column)

or matching nanodisc Env gp151 ND tetramers (right column). The frequency of tetramer^{++} cells

of IgG$^+$ cells is listed for each plot.

Figure 6. Structure of Env gp151 MPER ND in complex with bnAbs BG18, VRC01 and 10E8

resolved by cryo-EM. a) Low pass filtered maps of different Fab occupancy states and the location

of the bilayer. BG18 Fab is highlighted in green, VRC01 in pink and 10E8 in blue. b) 90° clockwise

rotated side views showing the Env tilt angle in relation to lipid bilayer surface. c) Top view and

side views of highest resolution maps of both 10E8 occupancy states. Densities are highlighted as

follows: gp120 in blue, gp41 in orange, BG18 in green, VRC01 in pink, HC of 10E8 in white, LC

of 10E8 in dark grey and MPER peptide in red. d) Highest resolution reconstruction with single

10E8 was used for model building. Insets show interfaces of the three bound bnAbs and key amino

acid and glycan interactions with BG18 HC (dark green) and LC (light green) in insets 1-2, VRC01

HC (magenta) and LC (pink) in insets 3-4, and 10E8 HC (white) and LC (dark grey) in insets 5-7.

Glycans are highlighted in purple. Insets 2 and 4 exemplify the electron density used to built

interacting glycans (transparent blue. e) Epitope-paratope analysis of 10E8 binding interface

shown in d. Contacting residues are separated into three components corresponding insets 5-7. f)

Binding pocket of 3BC315 (salmon) CDRH3 with key interactions highlighted. g) gp41 protomer

distance analysis in the nanodisc complex structure with 10E8 Fab, and soluble Env structure with

3BC315 showing widened interface between protomers A and B due to antibody binding.

Figures

Figure 1

Figure 2

Figure 3

Figure 4

852 **Figure 5**

854 **Figure 6**

References

1.   Bayburt, T.H., Grinkova, Y.V. & Sligar, S.G. Self-assembly of discoidal phospholipid
bilayer nanoparticles with membrane scaffold proteins. *Nano Letters* **2**, 853-856 (2002).
2.   Dorr, J.M. et al. The styrene-maleic acid copolymer: a versatile tool in membrane
research. *Eur Biophys J* **45**, 3-21 (2016).
3.   Carlson, M.L. et al. The Peptidisc, a simple method for stabilizing membrane proteins in
detergent-free solution. *Elife* **7** (2018).
4.   Frauenfeld, J. et al. A saposin-lipoprotein nanoparticle system for membrane proteins.
Nat Methods **13**, 345-351 (2016).
5.   Pettersen, J.M., Yang, Y. & Robinson, A.S. Advances in nanodisc platforms for
membrane protein purification. *Trends Biotechnol* **41**, 1041-1054 (2023).
6.   Sligar, S.G. & Denisov, I.G. Nanodiscs: A toolkit for membrane protein science. *Protein
Sci* **30**, 297-315 (2021).
7.   Denisov, I.G. & Sligar, S.G. Nanodiscs for structural and functional studies of membrane
proteins. *Nat Struct Mol Biol* **23**, 481-486 (2016).
8.   Trahey, M. et al. Applications of Lipid Nanodiscs for the Study of Membrane Proteins by
Surface Plasmon Resonance. *Curr Protoc Protein Sci* **81**, 29 13 21-29 13 16 (2015).
9.   Bocquet, N. et al. Real-time monitoring of binding events on a thermostabilized human
A2A receptor embedded in a lipid bilayer by surface plasmon resonance. *Biochim
Biophys Acta* **1848**, 1224-1233 (2015).
10. Gluck, J.M., Koenig, B.W. & Willbold, D. Nanodiscs allow the use of integral membrane
proteins as analytes in surface plasmon resonance studies. *Anal Biochem* **408**, 46-52
(2011).
11. Wang, M.Y. et al. SARS-CoV-2: Structure, Biology, and Structure-Based Therapeutics
Development. *Front Cell Infect Microbiol* **10**, 587269 (2020).
12. Pallesen, J. et al. Immunogenicity and structures of a rationally designed prefusion
MERS-CoV spike antigen. *Proc Natl Acad Sci U S A* **114**, E7348-E7357 (2017).
13. Kirchdoerfer, R.N. et al. Pre-fusion structure of a human coronavirus spike protein.
Nature **531**, 118-121 (2016).
14. Shepherd, B.O., Chang, D., Vasan, S., Ake, J. & Modjarrad, K. HIV and SARS-CoV-2:
Tracing a Path of Vaccine Research and Development. *Curr HIV/AIDS Rep* **19**, 86-93
(2022).
15. Pardi, N., Hogan, M.J., Porter, F.W. & Weissman, D. mRNA vaccines - a new era in
vaccinology. *Nat Rev Drug Discov* **17**, 261-279 (2018).
16. Chaudhary, N., Weissman, D. & Whitehead, K.A. mRNA vaccines for infectious
diseases: principles, delivery and clinical translation. *Nat Rev Drug Discov* **20**, 817-838
(2021).
17. Guenaga, J. et al. mRNA lipid nanoparticles expressing cell-surface cleavage
independent HIV Env trimers elicit autologous tier-2 neutralizing antibodies. *Front
Immunol* **15**, 1426232 (2024).

18. Benton, D.J. et al. Influenza hemagglutinin membrane anchor. *Proc Natl Acad Sci U S A*
115, 10112-10117 (2018).
19. Lopez, C.A., Alam, S.M., Derdeyn, C.A., Haynes, B.F. & Gnanakaran, S. Influence of
membrane on the antigen presentation of the HIV-1 envelope membrane proximal
external region (MPER). *Curr Opin Struct Biol* **88**, 102897 (2024).
20. Liu, H., Su, X., Si, L., Lu, L. & Jiang, S. The development of HIV vaccines targeting
gp41 membrane-proximal external region (MPER): challenges and prospects. *Protein*
Cell **9**, 596-615 (2018).
21. Guthmiller, J.J. et al. Broadly neutralizing antibodies target a haemagglutinin anchor
epitope. *Nature* **602**, 314-320 (2022).
22. Zhang, L. et al. An MPER antibody neutralizes HIV-1 using germline features shared
among donors. *Nat Commun* **10**, 5389 (2019).
23. Krebs, S.J. et al. Longitudinal Analysis Reveals Early Development of Three MPER-
Directed Neutralizing Antibody Lineages from an HIV-1-Infected Individual. *Immunity*
50, 677-691 e613 (2019).
24. Schiffner, T. et al. Vaccination induces broadly neutralizing antibody precursors to HIV
gp41. *Nat Immunol* **25**, 1073-1082 (2024).
25. Williams, W.B. et al. Vaccine induction of heterologous HIV-1-neutralizing antibody B
cell lineages in humans. *Cell* **187**, 2919-2934 e2920 (2024).
26. Cao, L. et al. Differential processing of HIV envelope glycans on the virus and soluble
recombinant trimer. *Nat Commun* **9**, 3693 (2018).
27. Rantalainen, K. et al. Co-evolution of HIV Envelope and Apex-Targeting Neutralizing
Antibody Lineage Provides Benchmarks for Vaccine Design. *Cell Rep* **23**, 3249-3261
(2018).
28. Torrents de la Pena, A. et al. Similarities and differences between native HIV-1 envelope
glycoprotein trimers and stabilized soluble trimer mimetics. *PLoS Pathog* **15**, e1007920
(2019).
29. Rantalainen, K. et al. HIV-1 Envelope and MPER Antibody Structures in Lipid
Assemblies. *Cell Rep* **31**, 107583 (2020).
30. Nakatani-Webster, E., Hu, S.L., Atkins, W.M. & Catalano, C.E. Assembly and
characterization of gp160-nanodiscs: A new platform for biochemical characterization of
HIV envelope spikes. *J Virol Methods* **226**, 15-24 (2015).
31. Bhattacharya, P. et al. Nanodisc-incorporated hemagglutinin provides protective
immunity against influenza virus infection. *J Virol* **84**, 361-371 (2010).
32. Mabrouk, M.T. et al. Circularized Nanodiscs for Multivalent Mosaic Display of SARS-
CoV-2 Spike Protein Antigens. *Vaccines (Basel)* **11** (2023).
33. Yang, S. et al. Dynamic HIV-1 spike motion creates vulnerability for its membrane-
bound tripod to antibody attack. *Nat Commun* **13**, 6393 (2022).
34. Steichen, J.M. et al. HIV Vaccine Design to Target Germline Precursors of Glycan-
Dependent Broadly Neutralizing Antibodies. *Immunity* **45**, 483-496 (2016).

35. Willis, J.R. et al. Human immunoglobulin repertoire analysis guides design of vaccine
priming immunogens targeting HIV V2-apex broadly neutralizing antibody precursors.
Immunity **55**, 2149-2167 e2149 (2022).
36. Puthenveetil, R., Nguyen, K. & Vinogradova, O. Nanodiscs and Solution NMR:
preparation, application and challenges. *Nanotechnol Rev* **6**, 111-126 (2017).
37. Gu, S., Huang, M. & Handel, T.M. On-bead purification and nanodisc reconstitution of
human chemokine receptor complexes for structural and biophysical studies. *STAR*
Protoc **4**, 102460 (2023).
38. Dev, J. et al. Structural basis for membrane anchoring of HIV-1 envelope spike. *Science*
353, 172-175 (2016).
39. Rutten, L. et al. Structure-Based Design of Prefusion-Stabilized Filovirus Glycoprotein
Trimers. *Cell Rep* **30**, 4540-4550 e4543 (2020).
40. Steichen, J.M. et al. A generalized HIV vaccine design strategy for priming of broadly
neutralizing antibody responses. *Science* **366** (2019).
41. Xie, Z. et al. mRNA-LNP HIV-1 trimer boosters elicit precursors to broad neutralizing
antibodies. *Science* **384**, eadk0582 (2024).
42. Baboo, S. et al. DeGlyPHER: An Ultrasensitive Method for the Analysis of Viral Spike
N-Glycoforms. *Anal Chem* **93**, 13651-13657 (2021).
43. Hossain, M.A. et al. B cells expressing IgM B cell receptors of HIV-1 neutralizing
antibodies discriminate antigen affinities by sensing binding association rates. *Cell Rep*
39, 111021 (2022).
44. Steichen, J.M. et al. Vaccine priming of rare HIV broadly neutralizing antibody
precursors in nonhuman primates. *Science* **384**, eadj8321 (2024).
45. Vazquez Bernat, N. et al. Rhesus and cynomolgus macaque immunoglobulin heavy-chain
genotyping yields comprehensive databases of germline VDJ alleles. *Immunity* **54**, 355-
366 e354 (2021).
46. Chernyshev, M., Kaduk, M., Corcoran, M. & Karlsson Hedestam, G.B. VDJ Gene Usage
in IgM Repertoires of Rhesus and Cynomolgus Macaques. *Front Immunol* **12**, 815680
(2021).
47. Phung, I. et al. A combined adjuvant approach primes robust germinal center responses
and humoral immunity in non-human primates. *Nat Commun* **14**, 7107 (2023).
48. Ramezani-Rad, P. et al. Vaccination with mRNA-encoded membrane-bound HIV
Envelope trimer induces neutralizing antibodies in animal models. *bioRxiv* (2025).
49. Irimia, A. et al. Lipid interactions and angle of approach to the HIV-1 viral membrane of
broadly neutralizing antibody 10E8: Insights for vaccine and therapeutic design. *PLoS*
Pathog **13**, e1006212 (2017).
50. Soto, C. et al. Developmental Pathway of the MPER-Directed HIV-1-Neutralizing
Antibody 10E8. *PLoS One* **11**, e0157409 (2016).
51. Rujas, E. et al. Structural basis for broad neutralization of HIV-1 through the molecular
recognition of 10E8 helical epitope at the membrane interface. *Sci Rep* **6**, 38177 (2016).
52. Huang, J. et al. Broad and potent neutralization of HIV-1 by a gp41-specific human
antibody. *Nature* **491**, 406-412 (2012).

53.   Lee, J.H. et al. Antibodies to a conformational epitope on gp41 neutralize HIV-1 by
979   destabilizing the Env spike. *Nat Commun* **6**, 8167 (2015).
980   54.   Pancera, M. et al. Structure and immune recognition of trimeric pre-fusion HIV-1 Env.
981   *Nature* **514**, 455-461 (2014).
982   55.   Ray, R. et al. Affinity gaps among B cells in germinal centers drive the selection of
983   MPER precursors. *Nature Immunology* **25** (2024).
984   56.   Cottrell, C.A. et al. Heterologous prime-boost vaccination drives early maturation of HIV
985   broadly neutralizing antibody precursors in humanized mice. *Sci Transl Med* **16**,
986   eadn0223 (2024).
987   57.   Antanasijevic, A. et al. Polyclonal antibody responses to HIV Env immunogens resolved
988   using cryoEM. *Nat Commun* **12**, 4817 (2021).
989   58.   Barnes, C.O. et al. Structural characterization of a highly-potent V3-glycan broadly
990   neutralizing antibody bound to natively-glycosylated HIV-1 envelope. *Nat Commun* **9**,
991   1251 (2018).
992   59.   He, L., Diedrich, J., Chu, Y.Y. & Yates, J.R., 3rd Extracting Accurate Precursor
993   Information for Tandem Mass Spectra by RawConverter. *Anal Chem* **87**, 11361-11367
994   (2015).
995   60.   Xu, T. et al. ProLuCID: An improved SEQUEST-like algorithm with enhanced
996   sensitivity and specificity. *J Proteomics* **129**, 16-24 (2015).
997   61.   Tabb, D.L., McDonald, W.H. & Yates, J.R., 3rd DTASelect and Contrast: tools for
998   assembling and comparing protein identifications from shotgun proteomics. *J Proteome
999   Res* **1**, 21-26 (2002).
1000   62.   Peng, J., Elias, J.E., Thoreen, C.C., Licklider, L.J. & Gygi, S.P. Evaluation of
1001   multidimensional chromatography coupled with tandem mass spectrometry (LC/LC-
1002   MS/MS) for large-scale protein analysis: the yeast proteome. *J Proteome Res* **2**, 43-50
1003   (2003).
1004   63.   Park, S.K., Venable, J.D., Xu, T. & Yates, J.R., 3rd A quantitative analysis software tool
1005   for mass spectrometry-based proteomics. *Nat Methods* **5**, 319-322 (2008).
1006   64.   Hurtado, J. et al. Efficient isolation of rare B cells using next-generation antigen
1007   barcoding. *Front Cell Infect Microbiol* **12**, 962945 (2022).
1008   65.   Breden, F. et al. Reproducibility and Reuse of Adaptive Immune Receptor Repertoire
1009   Data. *Front Immunol* **8**, 1418 (2017).
1010   66.   Punjani, A., Rubinstein, J.L., Fleet, D.J. & Brubaker, M.A. cryoSPARC: algorithms for
1011   rapid unsupervised cryo-EM structure determination. *Nat Methods* **14**, 290-296 (2017).
1012   67.   Bepler, T. et al. Positive-unlabeled convolutional neural networks for particle picking in
1013   cryo-electron micrographs. *Nat Methods* **16**, 1153-1160 (2019).
1014   68.   Punjani, A. & Fleet, D.J. 3D variability analysis: Resolving continuous flexibility and
1015   discrete heterogeneity from single particle cryo-EM. *J Struct Biol* **213**, 107702 (2021).
1016   69.   Punjani, A., Zhang, H. & Fleet, D.J. Non-uniform refinement: adaptive regularization
1017   improves single-particle cryo-EM reconstruction. *Nat Methods* **17**, 1214-1221 (2020).
1018   70.   Abramson, J. et al. Accurate structure prediction of biomolecular interactions with
1019   AlphaFold 3. *Nature* **630**, 493-500 (2024).

1020   71.   Pettersen, E.F. et al. UCSF ChimeraX: Structure visualization for researchers, educators,
1021   and developers. *Protein Sci* **30**, 70-82 (2021).
1022   72.   Casanal, A., Lohkamp, B. & Emsley, P. Current developments in Coot for
1023   macromolecular model building of Electron Cryo-microscopy and Crystallographic Data.
1024   *Protein Sci* **29**, 1069-1078 (2020).
1025   73.   Afonine, P.V. et al. Real-space refinement in PHENIX for cryo-EM and crystallography.
1026   *Acta Crystallogr D Struct Biol* **74**, 531-544 (2018).
1027   74.   Conway, P., Tyka, M.D., DiMaio, F., Konerding, D.E. & Baker, D. Relaxation of
1028   backbone bond geometry improves protein energy landscape modeling. *Protein Sci* **23**,
1029   47-55 (2014).

1030

1031