TITLE:

High-Throughput Assessment of Vocal Modulation Following Chemogenetic Inhibition in Songbirds

Authors

Chenyi Song (宋晨伊) [1,3], Mizuki Fujibayashi[1,3] and Kentaro Abe[1, 2, *]

Affiliations

[1] *Lab of Brain Development, Graduate School of Life Sciences, Tohoku University.*

[2] *Division for the Establishment of Frontier Sciences, Tohoku University,*
  Katahira 2-1-1, Aoba-ku, Sendai, Miyagi 980-8577, Japan.

[3] These authors contributed equally to this study.

* Corresponding author

Kentaro Abe, Ph. D.

Professor, Lab of Brain Development, Graduate School of Life Sciences, Tohoku University.

Katahira 2-1-1, Aoba-ku, Sendai, Miyagi 980-8577, Japan.

Email address: k.abe@tohoku.ac.jp

Abstract

Chemogenetic tools that enable temporary and repeatable manipulation of neuronal activity have become essential for studying the precise neural underpinning of animal behavior. However, their application has so far been limited to songbirds, which are widely used to study behaviors related to vocal communication. In this study, we applied and evaluated designer receptors exclusively activated by designer drugs (DREADD)-mediated neural suppression in songbird's brains, focusing on its impacts on vocalizations. We found that neuronal activity in zebra finches (*Taeniopygia guttata*) can be effectively suppressed both *in vitro* and *in vivo* using the inhibitory DREADD by its ligand deschloroclozapine (DCZ). By establishing a high-throughput system for recording and analyzing their vocalizations, we systematically assessed the effects of DREADD-mediated suppression on song behavior. Inhibiting HVC in zebra finches led to a reduction in song production number for approximately 90 min and altered phonological features over a longer time course. Notably, the concentration of DCZ required to induce robust effects on song behaviors was higher than doses typically used in mice. Furthermore, suppression of HVC or Area X in Bengalese finches (*Lonchura striata domestica*) produced a distinct effect on song features, suggesting species-specific responses to chemogenetic intervention. These findings refine the use of chemogenetic tools in songbirds and provide new insights into the neural mechanisms underlying vocal communication.

Introduction

Tools for manipulating neuron activity are highly versatile and play a central role in neuroscience studies aimed at uncovering the mechanisms of brain function by linking specific neuronal manipulations to resultant behaviors. Particularly valuable are technologies that enable reversive and repeatable control of defined neuronal populations, thereby allowing precise assessment of their contribution to the behaviors of interest. Moreover, minimally invasive approaches are especially desirable in experiments targeting naturalistic, spontaneous behaviors in freely moving animals. Chemogenetic tools are developed to fulfill these needs, aiming to manipulate neural activity *in vivo* with the least invasive procedures. Among such tools, the muscarinic acetyl choline receptors that was modified to selectively respond to artificial chemical reagent, namely Designer Receptors Exclusively Activated by Designer Drugs (DREADDs) have been widely employed in the field of behavioral neuroscience (Armbruster et al., 2007). Since their development, DREADDs have been utilized in a variety of experimental animal species, providing valuable insights into brain function across a wide range of neural processes

including behavior, learning and memory, emotion, sensory processing, whole body physiological regulation (Roth, 2016; Wiegert et al., 2017). The feature of DREADD to allow manipulation on the activity of specific population of neurons and glia within the brain, repetitively and least invasively, is ideal for songbird studies aiming at revealing the neural mechanisms behind communication which necessitates unstressed, natural behavior of subjects (Fujibayashi and Abe, 2024). However, compared to the widespread usage in other species, the application of DREADD technology in songbirds has been limited to few study so far (Heston et al., 2018).

To establish an experimental framework for DREADD-mediated neuronal manipulation in songbirds, this study reassessed the procedures for its application. We focused on hM4Di, which suppresses endogenous activity upon activation, as its effects are more physiologically interpretable and better suited for manipulating brain function (Armbruster et al., 2007). In contrast, other DREADD subtypes, such as hM3Dq and rM3Ds activation enhance neuronal activity but may induce non-physiological excitation patterns, complicating the interpretation of behavioral outcomes (Alexander et al., 2009). This is particularly important in the context of this study, which aims to evaluate the efficacy of DREADD manipulation in modulating song-related behaviors. We found that the DREADD agonist deschloroclozapine (DCZ) (Nagai et al., 2020), effectively suppressed spontaneous neuronal firing in cultured zebra finch neurons transfected with the inhibitory DREADD, hM4Di, in a manner similar to that observed in mouse neurons. Furthermore, we demonstrated that hM4Di also suppresses neuronal function *in vivo* in the zebra finch brain, as evidenced by the attenuation of auditory-induced neuronal activation. For their song's features, we found a reduction in song bout number, and modulation of the phonological features of vocalizations soon after DCZ administration. However, we observed that this effect required approximately tenfold higher DCZ concentrations *in vivo* compared to those used in rodents. By applying the same procedure in Bengalese finches, we found a similar but distinct effect on song, suggesting a species-specific effect on songs. These findings establish a foundation for employing DREADD technology in songbirds to elucidate the precise neural mechanism underlying interindividual communication.

Results
DREADD mediated suppression of neurons in songbird brain
To evaluate the efficacy of DREADD manipulation in the songbird neurons, we prepared dissociated primary neuron cultures from the telencephalon of zebra finch 1 day post hatch. In these cultures, we expressed the inhibitory DREADD receptor, hM4Di, together

with the genetically encoded calcium indicator GCaMP6s (Fig. 1A,B). After 10–15 days in culture, we applied bicuculline to enhance synaptic transmission among them, which induced periodic fluctuations in the GCaMP6s fluorescent signal driven by spontaneous firing (Fig. 1C). These fluctuations were suppressed by tetrodotoxin (TTX, 1 µM) treatment, confirming that they represent excitatory synaptic transmissions among the cultured neurons. The application of DCZ (4.0 µg/mL and 0.4 µg/mL) significantly reduced the fluctuation in GCaMP6s signals. Notably, there was no significant difference in the effect of DCZ treatments between zebra finch and mouse neurons, even at the lowest concentration (Fig. 1D; two-way ANOVA, mouse versus zebra finch, $F(1, 83) = 0.21$, $P = 0.65$). These findings demonstrate that hM4Di effectively suppresses neuronal activity in zebra finch neurons, consistent with its functionality in rodent neurons.

Next, we investigated the impact of DREADD activation on neural activity *in vivo*. The nidopallium caudomedial (NCM), an auditory region of the songbird brain, is known to exhibit robust neural responses to a playback of conspecific birdsong (Pinaud and Terleph, 2008). Using an adeno-associated vector (AAV, serotype 2/9 or 2/1), inhibitory DREADD, hM4Di was expressed to neurons within the NCM of zebra finch (Fig. 2A). To assess the effectiveness of hM4Di in suppressing neuronal activity, we evaluated auditory responses to conspecific songs and the bird's own song (BOS) by quantifying Egr-1/ZENK protein expression using immunostaining at 90 min following song presentation (Mello et al., 1992) (Fig. 2B). Throughout the experiment, birds were continuously observed by an experimenter nearby to ensure that they refrained from spontaneous vocalization. As anticipated, auditory stimulation elicited a robust induction of Egr-1 expression within NCM (Fig. 2C). Intramuscular injections of PBS or DCZ (either 0.1 or 1.0 mg/kg) 60 min before song stimuli, did not result in significant difference in the overall density of Egr-1 positive neurons following song presentation (Fig. 2D). However, we observed that DCZ treatment selectively attenuated Egr-1 expression within hM4Di-expressing neurons, indicating effective chemogenetic inhibition (Fig. 2E). This suppression exhibited a dose-dependent effect, with significant reduction in Egr-1 expression detected in birds administered with 1.0 mg/kg of DCZ, but not with 0.1 mg/kg dose (Fig. 2E). These findings demonstrate that hM4Di mediated DREADD activation effectively suppresses neuronal activity within the songbird brain *in vivo*.

Establishment of the pipeline for systematic analysis of song features
To evaluate the effects of DREADD manipulation on vocal features with particular focus on the time course of their effects, we developed a pipeline for comprehensive song

analysis over extended time periods (Fig. 3). The bird subjects, injected with an AAV vector to express DREADDs in the target brain nucleus were housed in soundproof chambers and alternately given daily injections of PBS and DCZ over 3–11 cycles (Fig. 3A, B). Songs were recorded using SAP2011, a standard song detection program in songbird research field (Tchernichovski et al., 2000). The recorded audio was processed with SAIBS, a machine learning-based system for automatic syllable annotation (Kawaji et al., 2024). In this study, we performed further customization of SAIBS to extract vocal onset and offset information alongside syllable labels, enabling high-throughput extraction of phonological features of each syllable. Using SAT, a MATLAB interface for SAP2011, this information was exported for downstream analysis (Fig. 3C). This integration allowed efficient large-scale analysis of vocal features across a large volume of syllables spanning the entire experimental period. Songs were extracted based on changes in sound density and the diversity of syllable types within high-density regions, and syllables selected for phonological analysis were obtained from these song segments. The data was pre-processed using an outlier detection algorithm based on principal component analysis (PCA), after which the phonological features for individual syllables were quantitatively analyzed (Fig. 3C). This pipeline allows a systematic and highly efficient framework for evaluating the temporal dynamics of DCZ-induced effects. It facilitates detailed assessment of vocal changes over time, capturing both the immediate consequences of DREADD manipulation and potential long-term post-manipulation plasticity.

Effect of DREADD mediated suppression on the song features of zebra finches

To investigate the effect of DREADD manipulation on song vocalization, we first focus on the effects on song after suppression of HVC (letter-based name), a brain nucleus playing a central role in song articulation (Long and Fee, 2008). Previous ablation studies have demonstrated robust and significant deformation and reduction in song bouts, with the severity of the effect depending on the extent of the affected neuronal population (Chen et al., 2014; Alalawi et al., 2019). To achieve DREADD-mediated suppression of HVC, male birds were bilaterally injected with AAV vectors to express hM4Di in the HVC (Fig. 3A). Birds expressing either hM4Di-mCherry or a control construct (mCherry) were isolated in soundproof chambers, and their vocalizations were continuously recorded and analyzed for entire day, throughout the experiment periods (Fig. 3B). DCZ or PBS was administered via intramuscular injection at approximately 9:30 AM on recording days. Each subject received alternating injections with PBS and DCZ at approximately 1–2-day intervals, and song features were analyzed within the same

individual to allow for within-subject comparison (Fig. 4A). In birds expressing hM4Di in HVC, DCZ injection (1.0 mg/kg) led to a transient reduction in song production, with a marked decrease observed during the 0–90 min period following administration (Fig. 4B). This reduction was not observed in control birds injected with DCZ or in birds expressing hM4Di in brain regions other than HVC, such as Area X, thereby ruling out the possibility that the effect was due to DCZ injection alone. Effect integral over the 0–120 min period, defined as the area between the song production curve and the normalized baseline value, revealed that HVC suppression significantly reduced song production compared to control birds and birds with Area X suppression. At the individual level, the DCZ-induced reduction in song production was consistently observed across eight cycles of DCZ/PBS injections, indicating that the effect is both reversible and reproducible (Fig. 4C). The phonological features of the songs were further analyzed using our pipeline for audio feature analysis (Fig. 3C). Immediately after the injection of DCZ in HVC suppressed birds, we observed a reduction in duration of motifs, a chunk of stereotypic sequence of syllables (Fig. 4D,E) (Glaze and Troyer, 2006). Because we analyzed motifs with identical syllable sequences, the observed effect was not attributable to changes in the syntactic organization within the motifs, but rather to phonological features of the syllables and the gaps between them.

Area X is part of the anterior forebrain pathway and is located upstream of LMAN, which is known to play a role in song learning by creating phonological variation in songs (Fee and Goldberg, 2011). Although Area X is not necessary in adult birds for the motor control in producing songs (Sohrabji et al., 1990), it is known to provide context dependent modulation of the phonological features of syllables (Kojima et al., 2018). Contrary to HVC suppression, we observed no significant effect on song production number by DREADD-mediated suppression of Area X (Fig. 3A,4B). As expected from previous studies (Kojima et al., 2018), we found suppression of Area X resulted in persistent reduction of the Wiener entropy of harmonic syllables during the 0–300 min period (Fig. 4F), the phenotype not observed in HVC suppressed birds. For both HVC suppression and Area X suppression, significant change in length of the syllable (Fig. 4G), extent of frequency modulation (FM) (Fig. 4F), and goodness of pitch was observed (Fig. 4G).

Taken together, the DCZ-induced effect in song features was reversible and reproducible, demonstrating the specificity and consistency of the DREADD-manipulation in songbird brain.

Dose-dependent effect of DREADD on song vocalization

Next, we evaluated the dose of DCZ required to induce the observed phenotype in zebra finches. For this analysis, we focused on the robust reduction in song number observed in the birds expressing hM4Di in HVC (Fig. 4B). We administrated varying doses of DCZ (0.01, 0.1, and 1.0 mg/kg) each day for about 6 rounds, and their effects on song production were assessed compared to vehicle-treated days (Fig. 5A). We observed a clear dose-dependent response: no discernible effect at 0.01 mg/kg, a mild reduction at 0.1 mg/kg, and robust suppression at 1.0 mg/kg (Fig. 5B). In this experiment, a decrease in song number that reached significant difference was detected during the 30–60 min periods following DCZ administration at the 1.0 mg/kg dose (Fig. 5B).

We noticed that the effective concentration of DCZ required to elicit robust song-related effects was approximately ten-fold higher than the dose typically used in monkeys and rodents (~0.1 mg/kg) (Nagai et al., 2020; Nentwig et al., 2022). To confirm the functionality of our hM4Di virus, we transduced the same virus into mouse striatum and analyzed the behavioral effects of DCZ injection (Fig. 5C,D). Following hemispheric virus injection, we facilitated spontaneous behavior by administering the dopamine reuptake inhibitor GBR12909 (13.3 mg/kg) (Lane et al., 2005). Subsequently, we injected 0.1 mg/kg of DCZ and analyzed behavior around 30–40 min after DCZ administration. We found that DCZ injection (0.1 mg/kg, a suboptimal dose for zebra finches) robustly induced ipsilateral rotation behavior in mice relative to the virus-infected hemisphere as observed in previous studies (Yu et al., 2022), verifying that our hM4Di viruses function as expected with 0.1 mg/kg DCZ dose in mice (Fig. 5E). Taken together, these results suggest that DCZ exhibits a dose-dependent effect in songbirds *in vivo*, requiring a higher concentration to produce robust song-related changes compared to its effect in rodents.

Effect of DREADD mediated suppression on the song features of Bengalese finches
To further generalize our findings, we applied DREADD-mediated suppression of the HVC and Area X in the Bengalese finch, another songbird species widely studied as a model for vocal communication (Okanoya, 2004; Sakata et al., 2008; Veit et al., 2021; Kawaji et al., 2024). Compared to the songs of zebra finches, songs of Bengalese finches are relatively longer and more variable in the syntactical organization of syllables (Okanoya, 2004). In contrast to zebra finches, hM4Di-mediated suppression of HVC in Bengalese finches did not exhibit significant effect on song production number when compared to the control-treated birds (Fig. 6A,B). For the phonological features of songs, we observed an immediate reduction (0–30 min) followed by a later extension (180–270 min) of motif duration in both HVC-suppressed and Area X-suppressed individuals (Fig. 6C,D). Similar to our observations in zebra finches, this effect was not attributable to

changes in the syntactic organization within motifs, as our analysis was conducted on motifs with identical syllable sequences. Focusing on harmonic syllables within the motifs, suppression of Area X reduced the syllable's Wiener entropy immediately following injection (0–30 min) (Fig. 6E), a phenotype not observed by HVC suppression. For both HVC suppression and Area X suppression, change in syllable duration (Fig. 6F), frequency modulation (FM) (Fig. 6G) was observed as with zebra finches although the detailed effect differed between two species. Notably, a steady increase in goodness of pitch was observed around 0–210 min period particularly in Area X suppressed birds (Fig. 6H), contrasting to what was observed in zebra finches (Fig. 4). Together, these findings confirm the effectiveness of DREADD manipulation in another songbird species and show some species-specific differences in the phenotypic effect on songs.

Discussion

DREADD-based manipulation strategies are well-suited for behavioral neuroscience research; however, their application in songbird studies has remained limited. This lack of prevalence suggests potential inefficiencies or failure in practically implementing DREADDs in songbird models. The aim of this study was to identify these limitations and develop improved methodologies for successful application of DREADD techniques in songbirds. In this study, we successfully demonstrated the reversible effects of hM4Di-mediated neuronal suppression *in vitro* and *in vivo*. Notably, no differences were observed in DCZ-induced DREADD activation in cultured neurons between mice and zebra finches. However, to elicit phenotypic changes in zebra finch *in vivo*, a DCZ dose of approximately 1.0 mg/kg was required, a dose significantly higher than that commonly used in rodent or primate studies (Nagai et al., 2020; Nentwig et al., 2022). The lack of observable effects at lower concentrations remains unexplained, but interspecies differences in drug metabolism may contribute to this discrepancy. Recent studies have shown that clozapine-N-oxidate (CNO), the originally developed DREADD ligand (Bender et al., 1994), exhibits poor brain penetrance and must be metabolized into clozapine to exert its effects (Gomez et al., 2017). Given that the metabolic profile of CNO varies even among rodent species, such as between rats and mice (Manvich et al., 2018), it is plausible that the limited success of DREADD application in songbird research may be attributed to species-specific differences in ligand metabolism between avian and mammalian systems. In this study, we used deschloroclozapine (DCZ), a newly developed DREADD ligand shown to exhibit superior brain penetrance compared to CNO in mammalian species (Nagai et al., 2020). Previous studies have demonstrated that CNO requires approximately 100-fold higher concentrations than DCZ to achieve a

comparable effects (Nagai et al., 2020; Nentwig et al., 2022), suggesting that using CNO in songbirds may require even higher doses than those required for DCZ. We successfully established that a DCZ dose of 1.0 mg/kg is sufficient to induce robust and reversible neuronal suppression in songbirds. While this finding confirms the effectiveness of DREADD-mediated manipulation in songbird models, the overall efficacy of it depends on both adequate receptor expression and ligand pharmacokinetics. Therefore, further investigation into alternative DREADD ligands—such as Compound 21 (Thompson et al., 2018) or JHU37152 and JHU37160 (Bonaventura et al., 2019) —as well as receptors based on different molecular backbones, such as KORDs (Vardy et al., 2015), may facilitate the development of more efficient and flexible chemogenetic approaches for studying neuronal function in the songbird brain.

Manipulating neural activity in songbird brains is essential for establishing causal relationship between neuronal activity and behavioral outcomes. One of the key advantages of DREADDs is their reversibility and reproductivity. In contrast, traditional approaches used in songbird research have largely relied on irreversible manipulations, such as neuronal ablation via local injection of ibotenic acid (Sohrabji et al., 1990; Foster and Bottjer, 2001), caspase activation (Sánchez-Valpuesta et al., 2019), or chromophore-targeted ablation (Scharff et al., 2000). While effective, these non-reversible methods often lack experimental flexibility and reproducibility within individual subjects, necessitating larger sample size to obtain robust results. Reversible approaches have also been explored in songbirds, including the use of tetrodotoxin (TTX) or $GABA_A R$ agonists (Cardin and Schmidt, 2004; Andalman and Fee, 2009; Hamaguchi and Mooney, 2012), and optogenetic tools (Xiao et al., 2018; Singh Alvarado et al., 2021; Katic et al., 2022) to transiently silence neuronal activity. However, these methods typically require invasive procedures, such as brain surgeries to implant cannulas or optical fibers for the delivery of pharmacological agents or light. Compared to these methods, another significant advantage of DREADD technology is its minimal invasiveness during experimental manipulation of targeted neurons. Unlike traditional approaches, the DREADD-based manipulation requires only a single viral transduction procedure performed well in advance of the experimental sessions. Subsequent manipulation of neuronal activity can be achieved through rapid and minimally invasive methods, such as intramuscular injection or oral administration of the ligand. These features make DREADDs particularly well-suited for studies of natural, undisturbed behaviors such as free vocal communication in songbirds, where minimizing stress and experimental disruption is essential.

By applying DREADD-mediated suppression in the songbird brain, we observed

both similarities and differences in experimental outcomes, compared to those reported using conventional methodologies (Chen et al., 2014; Alalawi et al., 2019; Isola et al., 2020). The discrepancy may, in part, stem from differences in the specific neural target manipulated by each approach. For instance, whereas methods such as ablation or TTX application affects both excitatory and inhibitory neurons, hM4Di expression in our study might restricted to specific neuron types because of the use of CaMKIIα promoter which express transgenes more specifically in excitatory neurons (Dittgen et al., 2004; Scheyltjens et al., 2015). Another possible contributing factor is the temporal nature of the manipulation. Conventional methods often involve chronic or irreversible modulation, while DREADD-based approaches allow for acute and reversible modification, which is often reported to produce distinct phenotypes (Otchy et al., 2015).

We observed the differential effects on song phenotypes following DREADD mediated neuronal suppression between zebra finches and Bengalese finches. These differences may arise from the distinct role of HVC and Area X in song vocalization between the two species. Notably, the HVC in Bengalese finches demonstrates auditory feedback response during singing (Okanoya and Yamaguchi, 1997), a feature absent in zebra finches (Hamaguchi et al., 2014). The absence of an effect on song production frequency in Bengalese finches following DREADD-mediated suppression of HVC— contrasting with the reduction observed in zebra finches—may reflect species specific differences in the functional organization of HVC. In particular, Bengalese finches may perceive discrepancies between intended and actual vocal output, potentially allowing them to compensate for disrupted motor commands. Alternatively, differences in the intrinsic connectivity within the HVC may also contribute to this variation. Neurons within the HVC exhibit highly stereotyped sequential activity during song production, driven by synfire chain networks formed through interconnections among these neurons (Hahnloser et al., 2002; Long et al., 2010). It has been hypothesized that Bengalese finches possess branching, parallel synfire chains (Jin, 2009), which may compensate for the impact of suppressing a small subset of neurons. In contrast, zebra finches are thought to rely on more liner synfire chains, making their vocal output more vulnerable to neuronal suppression occur. These functional difference in HVC neurons may be established by substantial differences in their gene expression profiles between the two species (Kato and Okanoya, 2010). Further studies are needed to reveal the species-specific mechanism of song control and to explore the evolutional origins of these differences.

As a limitation of study, the potential off-target effects of DCZ in avian species remain insufficiently explored. In our experiments, the administration of DCZ at a dose

of 1.0 mg/kg alone did not result in noticeable changes in song phenotypes. However, further detailed analysis of whole-body physiology is required, considering the molecular and genetic differences between mammals and birds. The observed 10-fold differences in the dose required for DREADD activation *in vivo* between mammals and birds suggest the need for more efficient brain-penetrant ligands for use in avian species. Furthermore, it remains unclear whether the difference in the effects of HVC or Area X suppression observed in this study compared to those using non-reversible methods were due to differences in acute or chronic suppression. Further studies utilizing similar chemogenetic tools designed for non-invasive and rapid manipulation of neuronal function will be required to address this question and refine our understanding of the mechanism of song control.

Conflict of interest statements
The authors declare no competing financial interests.

Acknowledgments
This work was supported by Tohoku University Research Program "Frontier Research in Duo" No.2101, JSPS/MEXT KAKENHI JP25H01037, JP24H01218, JP24H02146, JP23K18252, JP21K19424 awarded to K.A. and JP25KJ0541 awarded to M.F. We thank the members of the Abe Lab at Tohoku University for help and fruitful suggestions.

Author Contributions
C.S., M.F. and K.A. conceived and initiated the project. K.A. performed cell culture experiments and data analysis. C.S. performed *in vivo* experiments and data analysis. M.F. developed the methodology, performed experiments, and data analysis. C.S., M.F. and K.A. wrote the initial manuscript. All authors discussed and commented on the manuscript.

Material and Methods
Animals and Cares. The care and experimental manipulation of mice and birds used in this study were reviewed and approved by the institutional animal care and use committee of Tohoku University. All experiments and maintenance were conducted following relevant guidelines and regulations (Olson et al., 2014). Birds were purchased from Asada Chojyu and kept in our aviary under a 14-hour light / 10-hour dark cycle (daytime: 8:00−22:00); water and food were given *ad libitum*.

Cell culture and Ca²⁺ Imaging. Primary cortical neuron culture from mouse (*Mus musculus*, Slc:ICR) was prepared as described in previously (Abe and Abe, 2022). Primary neuron culture from zebra finches (*Taeniopygia guttata*) were prepared from individuals of post hatch day 1. The forebrain of their brain was collected and processed with enzymatic digestion with papain (Roche, #10108014001; 300 U/mL) for 20 min at 37°C. The tissues are titrated to dissociate into single cells and plated onto 24-well plates pretreated with poly-L-lysin (SIGMA-Aldrich, #P2336). Cells are cultured in DMEM/F12 (Fujifilm-Wako) with 10% fatal bovine serum (FBS, Biowest) for initial plating, and then maintained with an equal mixture of DMEM/F12 with 10% FBS and Neurobasal medium (Thermo-Fisher, #21103049) supplemented with B27 plus supplement (20×; Thermo-Fisher, #A3582801), Glutamax-I (100×; Thermo-Fisher, #35050061) and penicillin-streptomycin (100×; Fujifilm-Wako) at 37°C under 5% CO_2. Cells were transfected with plasmids using HilyMax reagent (Dojindo) at 2 days *in vitro* (div). Live imaging was performed at 10–15 div at 37°C, using inverted fluorescence microscope (Axiovert A1 MAT, Zeiss) equipped with a Colibri 7 system (Zeiss) and a stage warmer (Tokai hit). Approximately 30 min prior to the imaging, bicuculline (bicuculline methiodide 200 μM; Tocris #131) was applied to the well. Fluorescent signals were captured at 10 Hz using 20× objective lens. A droplet of DCZ (4.0 or 0.4 μg/mL in 20 μL PBS; Deschloroclozapine dihydrochloride, MedChemExpress, #HY-42110A) or tetrodotoxin (TTX, 1.0 μM, Nakarai-tesque #32775-51) were applied during the imaging at specified timings. Spike numbers from GCaMP6s signals of each neuron were extracted by PCA/ICA method (Mukamel et al., 2009) and Detection of raising event algorithms of Mosaic analysis software (Inscopix).

Surgery and DCZ application. Adult zebra finches, Bengalese finches and mice (*Mus musculus*, C57BL/6J) were anesthetized with medetomidine-midazolam-butorphanol mixture (medetomidine 30 mg/mL, All Japan pharma; midazolam 30 mg/mL, Astellas Pharma; butorphanol tartrate 500 mg/mL Meiji Seika Pharma; NaCl 118 mM Fujifilm-Wako; 100 µL per bird and 350 µL per mouse) (Yamamoto and Abe, 2022). In zebra finches, AAVs (1.0 µL per bird) are injected at the coordinate from the Y sinus: anterior, 0 mm; lateral, 2.2 mm; depth, 0.75 mm at beak angle 40° for HVC; anterior, 0.5 mm; lateral, 0.6 mm; depth 1.0 mm and 0 mm; lateral, 2.2 mm; depth, 2.0 mm at beak angle 40° for NCM; and anterior, 5.5 mm; lateral, 1.2 mm; depth, 3.1 mm and 2.9 mm at beak angle 60° for Area X. In Bengalese finches, AAVs (1.0 µL per bird) were injected at the coordinate from the Y sinus: anterior, 5.75 mm; lateral, 1.2 mm; depth, 2.9 mm and 2.7

mm at beak angle 60° for Area X; and anterior, 0 mm; lateral, 2.0 mm; depth 0.75 mm at beak angle 40° for HVC. For a mouse, AAVs (1.0 µL per mouse) were injected unilaterally into the right striatum at six locations with the coordinate from bregma: anterior, 0 mm; lateral, 2.0 mm; depth, 3.0 mm, 2.6 mm and 2.3 mm; and anterior, 1.0 mm; lateral, 1.8 mm; depth, 3.0 mm, 2.6 mm and 2.3 mm. DCZ/PBS application was performed more than 30 days after the injection. 10 mg/mL stock solution of deschlorocZapine-hydrocloride (DCZ-2HCl) was diluted at desired concentration with 150 µL PBS per bird and injected into the pectoral muscle around 9:30 AM every 1–2 days.

Viruses and plasmid constructs. Plasmid used for primary neuron transfection are following: pAAV-CAG-GCaMP6s (RRID: Addgene_100844, a gift from Douglas Kim & GENIE Project) (Chen et al., 2013), and pAAV-CaMKIIa-hM4Di-mCherry (RRID: Addgene_50477, a gift from Bryan Roth). AAVs used for *in vivo* transduction are following: AAV2/9-CaMKIIa-hM4D(Gi)-mCherry (titer $\geq 1.0 \times 10^{13}$ vg/mL, Addgene viral prep #50477-AAV9; 1.0 µL per bird) for hemispheric injection into NCM in auditory response experiments; 4:1 mixture of AAV2/9-CaMKIIa-hM4D(Gi)-mCherry, and AAV2/9-Syn-EGFP (3.54×10^{13} DNase resistant particle/mL); or AAV2/1-CaMKIIa-hM4D(Gi)-2a-EGFP (1.1×10^{13} DNase resistant particle/mL) and AAV2/9-Syn1-mCherry (1.8×10^{14} DNase resistant particle/mL); or AAV2/9-Syn1-mCherry (1.31×10^{13} DNase resistant particle/mL) and AAV2/9-Syn1-EGFP (3.54×10^{13} DNase resistant particle/mL) for bilateral injection at Area X and HVC for song phenotype analysis; 4:1 mixture of AAV2/9-CaMKIIa-hM4D(Gi)-mCherry, and AAV2/9-Syn1-EGFP for mouse striatum. AAVs are obtained from Addgene, or produced in house as reported previously (Abe and Abe, 2022).

Auditory stimulation and immunostaining. Auditory stimulation was conducted on birds >30 days after AAV injection. Male birds were isolated in a soundproof chamber and exposed to a randomized playback of a mixture of their own song and conspecific songs (5–9 s for a bout) randomly every minute for 30 min. Subjects were sacrificed 90 min after the auditory stimulation. To prevent vocalization of songs and calls before and after the auditory stimulation, the chamber was illuminated with dim red light, and an experimenter continuously monitored the birds near the chamber to ensure they neither sang nor fell asleep. The birds were rapidly euthanized after 90 min and perfused with PBS followed by 4% paraformaldehyde (PFA, Nakarai-tesque) in PBS. The cryo-sections were immunostained with Egr-1 (1:800; SantaCruz, #sc-189) and RFP (1:1000; MBL,

#M165-3MS) followed by 4′,6-diamidino-2-phenylindole (DAPI, 0.33 ng/mL, TCI) staining. Images were taken by BZ-810 or BZ-9000 (Keyence) using 4× lens or 10× lens and the whole section merged images were used for the inspection of the infected area, or by FV4000 (Olympus) using 4×, 10×, and 20× lens, and the Z-projection images were created using cellSens FV (Olympus). The expression cell was detected by visual inspection of the stained images using OpenCV-Python (UK Intel Corporation).

Song recording and analysis pipeline. Songs were recorded using SAP2011 (Tchernichovski et al., 2000) and processed with SAIBS (Kawaji et al., 2024) to classify and annotate syllables, as well as to extract vocal onset and offset time along with syllable labels. Song segments were selected based on changes in sound density and the diversity of syllable types within high-density regions. Syllables used for phonological analyses were collected from these extracted songs. We utilized SAT, a MATLAB interface of SAP2011 to extract vocal features between the vocal onset and offset of each syllable. To exclude miss-annotated syllables, the data was then pre-processed through an outlier detection algorithm using principal component analysis (PCA). For each syllable type, we conducted PCA using the major vocal features of each syllable: duration, average frequency, average frequency modulation, average amplitude modulation, average goodness, and average entropy. After transforming the data, retaining the principal components that explain more than 90% of the variance, the inverse transformation was performed. Then, we calculated the mean squared error (MSE) between the reconstructed data and the original data. Using these MSE scores, anomaly detection was performed using the Smirnov-Grubbs test at a significant level of 0.05. The resultant phonological features for each syllable were analyzed. This outlier detection was conducted using Python.

Analysis of songs. Songs were counted every 30 min for 60 min before and 360 min after the injection of PBS/DCZ. For each individual, song counts on DCZ-injected days were normalized to the mean count on PBS-injected days. Motifs were defined as stereotyped sequences of syllables that consistently appear in zebra finch songs, and as invariant sequences appearing with a frequency of at least once per song in Bengalese finch songs. Motif duration was calculated as the duration between the onset of the first syllable and the offset time of the last syllable. All phonological features extracted by using the pipeline mentioned in Figure 3 were standardized by subtracting the mean PBS value and dividing by the standard deviation of PBS values for each 30-min time bin. For zebra finch data, outliers were excluded from each time bin using the interquartile range method

following standardization. Effect integral was calculated as the area between the line graph and the baseline value of 1 for song counts, and 0 for motif durations. As controls, we prepared subjects expressing control constructs exclusively in either HVC or Area X, but the subjects were pooled together for analysis.

Effect of DREADD inhibition on mouse behavior. Mouse behavioral analysis of DREADD induced hemispheric suppression of the striatum was conducted in an open field box ($40 \times 40 \times 40$ cm) constructed from gray plastic plates. The subject with hemispheric injection of hM4Di virus 35 days in advance, was habituated to the box for 10 min before the test. GBR12909 (13.3 mg/kg in 200 µL of 20% DMSO/PBS, i.p., TCI #G0541) was administered first, followed by either DCZ (0.1 mg/kg in 100 µL PBS, i.p.) or vehicle (100 µL PBS, i.p.) at 40-min intervals. All drugs were administered into the right inguinal region. DCZ and PBS treatments was administered as a consecutive-day pairs, with a 1–2-day interval before repeating the cycle, resulting in three repetitions of each treatment. Behavior was video recorded at a resolution of 480×640 pixels and 30 fps using a USB-camera positioned above the open-field box. From the recorded video, the number of right and left turns was manually counted from 30–40 min after drug administration. A rotation was defined as a movement in which the subject turned more than 180° within a circle with a diameter approximately equal to the body length, without pausing for 1–2 s.

Statistics. The alpha score of 0.05 was used to reject the null hypothesis. Two-way ANOVA with post hoc Dunnett's test, Wilcoxon's rank sum test, and Bonferroni correction were used for multiple comparisons. An independent sample *t*-test was used for the comparison of two data sets. One sample *t*-test was used for the comparison of one dataset and the baseline value. All statistical analyses were conducted using R.

Figure and Legends

Figure 1. DREADD mediated suppression of neuron activation in cultured songbird neurons *in vitro*. (**A**) Schematic diagram of experimental procedures. Neuron culture from mouse embryo and zebra finch chick was prepared and transfected. (**B**) Fluorescent image of a cultured zebra finch neuron at 15 div. (**C**) Time lapse plot of fluorescent signals from single cells expressing GCaMP6s before and after PBS, TTX, and DCZ application. Drugs are applied to the medium at 60 s (colored highlights). (**D**) Ca^{2+} spike number after drug application (120–180 s) normalized against the value before drug application (0–60 s). * $P < 0.05$ against PBS in each cohort, Dunnett's test. The number of cells in each cohort are indicated in the bars. Scale bare, 50 µm.

Figure 2. DREADD mediated suppression of auditory responses in songbird brain *in vivo*. (**A**) Immunostained section showing AAV injection site in adult zebra finch brain. NCM, nidopallium caudomedial. Scale bar, 500 μm. (**B**) Schematic diagram of experimental procedures and auditory stimulation setup. (**C**) Immunostained sections. Left, PBS injected bird kept in silence. Right, DCZ injected bird with song presentation. Scale bars, 100 μm. (**D**) The number of Egr-1(+) cells in each group. *n* = 8 sections per group. n. s., $P > 0.05$ against PBS. *P* values are from the Wilcoxon rank sum test and Bonferroni correction. (**E**) Ratio of Egr-1(+) cells within hM4Di (+) cells. * $P < 0.05$ against PBS. *P* values are from Chi-squared test and Bonferroni correction.

Figure 3. Experimental pipeline for systematic analysis of song features. (**A**) Immunostained sections showing AAV injection site in adult zebra finch brain. Red, hM4Di; blue, DAPI. Scale bars, 100 μm. (**B**) Schematic diagram of experimental procedures. Birds were treated with 150 μL of DCZ or PBS through bilateral pectoral injection around 9:30 a.m. every 1 or 2 days. Their songs were recorded with a microphone and processed using SAP2011 from 60 min before and 360 min after the drug administration. (**C**) Schematic diagram for high-throughput vocal analysis. Syllables from finch songs were annotated by SAIBS and calculated phonological features by SAT. The features produced by SAT were utilized to the PCA outlier detection to eliminate the missed annotations.

Figure 4. DREADD mediated suppression of HVC and Area X in zebra finches. (**A**) Schematics of experiment. Each day, control birds (black; no hM4Di expression), and birds expressing hM4Di in HVC (orange) or Area X (blue) were injected with PBS or DCZ of 1.0 mg/kg. (**B**) Number of songs produced per 30-min bin, normalized to PBS controls for each corresponding bin. Bar graph on the right displays integrated effects over the 0–120 min period. n = 5 (control), 9 (HVC), 7 (Area X) birds. (**C**) Trace showing the number of song bouts across eight DCZ administration cycles in a single subject expressing hM4Di in HVC. (**D**) Representative sonograms from birds injected with PBS / DCZ. For comparison, motifs with minimum (DCZ) and maximum (PBS) duration are shown. Vertical lines indicate the onset and offset of a motif. (**E**) Standardized motif duration. Bar graph on the right displays the integrated effects over the 0–120 min period. n = 6 (control), 16 (HVC), 10 (Area X) motifs recorded from 5 (control), 9 (HVC), 7 (Area X) birds. (**F**–**I**) Standardized phonological features of a harmonic syllable. Winner

entropy (**F**), Syllable duration (**G**), extent of Frequency modulation (**H**), and mean goodness of pitch (**I**). $n = 8$ (control), 11 (HVC), and 12 (Area X) syllables recorded from 4 (control), 7 (HVC), 5 (Area X) birds. Across the panels, values from DCZ-injected days were normalized (**A**, **C**) or standardized (**E–I**) to the corresponding PBS-injected values within each 30-min bin. Line and bar plots represent mean ± SEM across subjects. * $P < 0.05$. *P*-values displayed in bar graph were calculated using Wilcoxon rank-sum test with Bonferroni correction for multiple comparisons, and those in line graphs using one-sample *t*-test.

Figure 5. Dose analysis of DREADD effect on songs. (**A**) Schematic diagram of experimental procedures. Zebra finches were treated with three types of concentration DCZ. 1.0mg/kg, 0.1mg/kg, and 0.01mg/kg. 150 µL DCZ solvent was intramuscularly administrated. (**B**) Number of songs produced per 30-min bin, normalized to PBS controls for each corresponding bin. Bar graph on the right displays integrated effects over the 0–120 min period. $n = 5$ birds. * $P < 0.05$. P-values were calculated using one sample t-test (**C**) Schematic diagram of experimental procedures and behavioral analysis setup. (**D**) Immunostained brain section of a mouse with unilateral hM4Di injection. Scale bare, 1.0 mm. (**E**) Change in the number of ipsilateral turns toward the hM4Di injected hemisphere following DCZ administration. $n = 3$ (PBS) and 4 (DCZ) treatments were applied to a single mouse. * $P < 0.05$. P-values were calculated using Welch's t-test.

Figure 6. DREADD mediated suppression of HVC and Area X in Bengalese finches. (**A**) Schematics of experiment. Each day, control birds (black; no hM4Di expression), and birds expressing hM4Di in HVC (orange) or Area X (blue) were injected with PBS or DCZ of 1.0 mg/kg (**B**) Number of songs produced per 30-min bin, normalized to PBS controls for each corresponding bin. Bar graph on the right displays integrated effects over the 0–120 min period. $n = 4$ (control), 5 (HVC), and 3 (Area X) birds. (**C**) Representative sonograms from birds injected with PBS / DCZ. For comparison, motifs with minimum (DCZ) and maximum (PBS) duration are shown. Vertical lines indicate the onset and offset of a motif. (**D**) Standardized duration of motifs. Bar graph on the right displays the integrated effects over the 0–120 min period. $n = 8$ (control), 12 (HVC), and 7 (Area X) motifs. (**E**–**H**) Standardized phonological features of harmonic syllables within motif. Winner entropy (**E**), Syllable duration (**F**), Frequency modulation (**G**), and mean goodness of pitch (**H**). $n = 11$ (control), 19 (HVC), 16 (Area X) syllables recorded from 4 (control), 5 (HVC), and 3 (Area X) birds. Across the panels, values from DCZ-injected days were normalized (**B**) or standardized (**D**–**H**) to the corresponding PBS-

injected values within each 30-min bin. Line and bar plots represent mean ± SEM across subjects. * $P < 0.05$. *P*-values displayed in bar graphs were calculated using Wilcoxon rank-sum test with Bonferroni correction for multiple comparisons, and those in line graphs using one-sample *t*-test.

References

Abe H, Abe K (2022) PCR-based profiling of transcription factor activity in vivo by a virus-based reporter battery. iScience 25:103927.

Alalawi A, Dodu JC, Woolley-Roberts M, Brodie J, Di Marzo V, Soderstrom K (2019) Cannabidiol improves vocal learning-dependent recovery from, and reduces magnitude of deficits following, damage to a cortical-like brain region in a songbird pre-clinical animal model. Neuropharmacology 158:107716.

Alexander GM, Rogan SC, Abbas AI, Armbruster BN, Pei Y, Allen JA, Nonneman RJ, Hartmann J, Moy SS, Nicolelis MA, McNamara JO, Roth BL (2009) Remote Control of Neuronal Activity in Transgenic Mice Expressing Evolved G Protein-Coupled Receptors. Neuron 63:27–39.

Andalman AS, Fee MS (2009) A basal ganglia-forebrain circuit in the songbird biases motor output to avoid vocal errors. Proceedings of the National Academy of Sciences 106:12518–12523.

Armbruster BN, Li X, Pausch MH, Herlitze S, Roth BL (2007) Evolving the lock to fit the key to create a family of G protein-coupled receptors potently activated by an inert ligand. Proceedings of the National Academy of Sciences 104:5163–5168.

Bender D, Holschbach M, Stöcklin G (1994) Synthesis of n.c.a. carbon-11 labelled clozapine and its major metabolite clozapine-*N*-oxide and comparison of their biodistribution in mice. Nuclear Medicine and Biology 21:921–925.

Bonaventura J et al. (2019) High-potency ligands for DREADD imaging and activation in rodents and monkeys. Nat Commun 10:4627.

Cardin JA, Schmidt MF (2004) Auditory Responses in Multiple Sensorimotor Song System Nuclei Are Co-Modulated by Behavioral State. Journal of Neurophysiology 91:2148–2163.

Chen JR, Stepanek L, Doupe AJ (2014) Differential contributions of basal ganglia and thalamus to song initiation, tempo, and structure. Journal of Neurophysiology 111:248–257.

Chen T-W, Wardill TJ, Sun Y, Pulver SR, Renninger SL, Baohan A, Schreiter ER, Kerr RA, Orger MB, Jayaraman V, Looger LL, Svoboda K, Kim DS (2013)

Ultrasensitive fluorescent proteins for imaging neuronal activity. Nature 499:295–300.

Dittgen T, Nimmerjahn A, Komai S, Licznerski P, Waters J, Margrie TW, Helmchen F, Denk W, Brecht M, Osten P (2004) Lentivirus-based genetic manipulations of cortical neurons and their optical and electrophysiological monitoring in vivo. Proceedings of the National Academy of Sciences 101:18206–18211.

Fee MS, Goldberg JH (2011) A hypothesis for basal ganglia-dependent reinforcement learning in the songbird. Neuroscience 198:152–170.

Foster EF, Bottjer SW (2001) Lesions of a telencephalic nucleus in male zebra finches: Influences on vocal behavior in juveniles and adults. Journal of Neurobiology 46:142–165.

Fujibayashi M, Abe K (2024) A behavioral analysis system MCFBM enables objective inference of songbirds' attention during social interactions. Cell Reports Methods 4:100844.

Glaze CM, Troyer TW (2006) Temporal Structure in Zebra Finch Song: Implications for Motor Coding. J Neurosci 26:991–1005.

Gomez JL, Bonaventura J, Lesniak W, Mathews WB, Sysa-Shah P, Rodriguez LA, Ellis RJ, Richie CT, Harvey BK, Dannals RF, Pomper MG, Bonci A, Michaelides M (2017) Chemogenetics revealed: DREADD occupancy and activation via converted clozapine. Science 357:503–507.

Hahnloser RHR, Kozhevnikov AA, Fee MS (2002) An ultra-sparse code underliesthe generation of neural sequences in a songbird. Nature 419:65–70.

Hamaguchi K, Mooney R (2012) Recurrent Interactions between the Input and Output of a Songbird Cortico-Basal Ganglia Pathway Are Implicated in Vocal Sequence Variability. J Neurosci 32:11671–11687.

Hamaguchi K, Tschida KA, Yoon I, Donald BR, Mooney R (2014) Auditory synapses to song premotor neurons are gated off during vocalization in zebra finches Calabrese RL, ed. eLife 3:e01833.

Heston JB, Simon IV J, Day NF, Coleman MJ, White SA (2018) Bidirectional scaling of

vocal variability by an avian cortico-basal ganglia circuit. Physiological Reports 6:e13638.

Isola GR, Vochin A, Sakata JT (2020) Manipulations of inhibition in cortical circuitry differentially affect spectral and temporal features of Bengalese finch song. Journal of Neurophysiology 123:815–830.

Jin DZ (2009) Generating variable birdsong syllable sequences with branching chain networks in avian premotor nucleus HVC. Phys Rev E Stat Nonlin Soft Matter Phys 80:051902.

Katic J, Morohashi Y, Yazaki-Sugiyama Y (2022) Neural circuit for social authentication in song learning. Nat Commun 13:4442.

Kato M, Okanoya K (2010) Molecular characterization of the song control nucleus HVC in Bengalese finch brain. Brain Research 1360:56–76.

Kawaji T, Fujibayashi M, Abe K (2024) Goal-directed and flexible modulation of syllable sequence within birdsong. Nat Commun 15:3419.

Kojima S, Kao MH, Doupe AJ, Brainard MS (2018) The Avian Basal Ganglia Are a Source of Rapid Behavioral Variation That Enables Vocal Motor Exploration. J Neurosci 38:9635–9647.

Lane EL, Cheetham S, Jenner P (2005) Dopamine Uptake Inhibitor-Induced Rotation in 6-Hydroxydopamine-Lesioned Rats Involves Both D1 and D2 Receptors but Is Modulated through 5-Hydroxytryptamine and Noradrenaline Receptors. J Pharmacol Exp Ther 312:1124–1131.

Long MA, Fee MS (2008) Using temperature to analyse temporal dynamics in the songbird motor pathway. Nature 456:189–194.

Long MA, Jin DZ, Fee MS (2010) Support for a synaptic chain model of neuronal sequence generation. Nature 468:394–399.

Manvich DF, Webster KA, Foster SL, Farrell MS, Ritchie JC, Porter JH, Weinshenker D (2018) The DREADD agonist clozapine N-oxide (CNO) is reverse-metabolized to clozapine and produces clozapine-like interoceptive stimulus effects in rats and mice. Sci Rep 8:3840.

Mello CV, Vicario DS, Clayton DF (1992) Song presentation induces gene expression in the songbird forebrain. Proceedings of the National Academy of Sciences 89:6818–6822.

Mukamel EA, Nimmerjahn A, Schnitzer MJ (2009) Automated Analysis of Cellular Signals from Large-Scale Calcium Imaging Data. Neuron 63:747–760.

Nagai Y et al. (2020) Deschloroclozapine, a potent and selective chemogenetic actuator enables rapid neuronal and behavioral modulations in mice and monkeys. Nat Neurosci 23:1157–1167.

Nentwig TB, Obray JD, Vaughan DT, Chandler LJ (2022) Behavioral and slice electrophysiological assessment of DREADD ligand, deschloroclozapine (DCZ) in rats. Sci Rep 12:6595.

Okanoya K (2004) The Bengalese Finch: A Window on the Behavioral Neurobiology of Birdsong Syntax. Annals of the New York Academy of Sciences 1016:724–735.

Okanoya K, Yamaguchi A (1997) Adult bengalese finches (Lonchura striata var. domestica) require real-time auditory feedback to produce normal song syntax. Journal of Neurobiology 33:343–356.

Olson CR, Wirthlin M, Lovell PV, Mello CV (2014) Proper Care, Husbandry, and Breeding Guidelines for the Zebra Finch, Taeniopygia guttata. Cold Spring Harb Protoc 2014:pdb.prot084780.

Otchy TM, Wolff SBE, Rhee JY, Pehlevan C, Kawai R, Kempf A, Gobes SMH, Ölveczky BP (2015) Acute off-target effects of neural circuit manipulations. Nature 528:358–363.

Pinaud R, Terleph TA (2008) A songbird forebrain area potentially involved in auditory discrimination and memory formation. J Biosci 33:145–155.

Roth BL (2016) DREADDs for Neuroscientists. Neuron 89:683–694.

Sakata JT, Hampton CM, Brainard MS (2008) Social Modulation of Sequence and Syllable Variability in Adult Birdsong. Journal of Neurophysiology 99:1700–1711.

Sánchez-Valpuesta M, Suzuki Y, Shibata Y, Toji N, Ji Y, Afrin N, Asogwa CN, Kojima I, Mizuguchi D, Kojima S, Okanoya K, Okado H, Kobayashi K, Wada K (2019)

Mello CV, Vicario DS, Clayton DF (1992) Song presentation induces gene expression in the songbird forebrain. Proceedings of the National Academy of Sciences 89:6818–6822.

Mukamel EA, Nimmerjahn A, Schnitzer MJ (2009) Automated Analysis of Cellular Signals from Large-Scale Calcium Imaging Data. Neuron 63:747–760.

Nagai Y et al. (2020) Deschloroclozapine, a potent and selective chemogenetic actuator enables rapid neuronal and behavioral modulations in mice and monkeys. Nat Neurosci 23:1157–1167.

Nentwig TB, Obray JD, Vaughan DT, Chandler LJ (2022) Behavioral and slice electrophysiological assessment of DREADD ligand, deschloroclozapine (DCZ) in rats. Sci Rep 12:6595.

Okanoya K (2004) The Bengalese Finch: A Window on the Behavioral Neurobiology of Birdsong Syntax. Annals of the New York Academy of Sciences 1016:724–735.

Okanoya K, Yamaguchi A (1997) Adult bengalese finches (Lonchura striata var. domestica) require real-time auditory feedback to produce normal song syntax. Journal of Neurobiology 33:343–356.

Olson CR, Wirthlin M, Lovell PV, Mello CV (2014) Proper Care, Husbandry, and Breeding Guidelines for the Zebra Finch, Taeniopygia guttata. Cold Spring Harb Protoc 2014:pdb.prot084780.

Otchy TM, Wolff SBE, Rhee JY, Pehlevan C, Kawai R, Kempf A, Gobes SMH, Ölveczky BP (2015) Acute off-target effects of neural circuit manipulations. Nature 528:358–363.

Pinaud R, Terleph TA (2008) A songbird forebrain area potentially involved in auditory discrimination and memory formation. J Biosci 33:145–155.

Roth BL (2016) DREADDs for Neuroscientists. Neuron 89:683–694.

Sakata JT, Hampton CM, Brainard MS (2008) Social Modulation of Sequence and Syllable Variability in Adult Birdsong. Journal of Neurophysiology 99:1700–1711.

Sánchez-Valpuesta M, Suzuki Y, Shibata Y, Toji N, Ji Y, Afrin N, Asogwa CN, Kojima I, Mizuguchi D, Kojima S, Okanoya K, Okado H, Kobayashi K, Wada K (2019)

Corticobasal ganglia projecting neurons are required for juvenile vocal learning but not for adult vocal plasticity in songbirds. Proceedings of the National Academy of Sciences 116:22833–22843.

Scharff C, Kirn JR, Grossman M, Macklis JD, Nottebohm F (2000) Targeted Neuronal Death Affects Neuronal Replacement and Vocal Behavior in Adult Songbirds. Neuron 25:481–492.

Scheyltjens I, Laramée M-E, Van den Haute C, Gijsbers R, Debyser Z, Baekelandt V, Vreysen S, Arckens L (2015) Evaluation of the expression pattern of rAAV2/1, 2/5, 2/7, 2/8, and 2/9 serotypes with different promoters in the mouse visual cortex. Journal of Comparative Neurology 523:2019–2042.

Singh Alvarado J, Goffinet J, Michael V, Liberti W, Hatfield J, Gardner T, Pearson J, Mooney R (2021) Neural dynamics underlying birdsong practice and performance. Nature 599:635–639.

Sohrabji F, Nordeen EJ, Nordeen KW (1990) Selective impairment of song learning following lesions of a forebrain nucleus in the juvenile zebra finch. Behavioral and Neural Biology 53:51–63.

Tchernichovski O, Nottebohm F, Ho CE, Pesaran B, Mitra PP (2000) A procedure for an automated measurement of song similarity. Animal Behaviour 59:1167–1176.

Thompson KJ et al. (2018) DREADD Agonist 21 Is an Effective Agonist for Muscarinic-Based DREADDs in Vitro and in Vivo. ACS Pharmacol Transl Sci 1:61–72.

Vardy E et al. (2015) A New DREADD Facilitates the Multiplexed Chemogenetic Interrogation of Behavior. Neuron 86:936–946.

Veit L, Tian LY, Monroy Hernandez CJ, Brainard MS (2021) Songbirds can learn flexible contextual control over syllable sequencing. Elife 10:e61610.

Wiegert JS, Mahn M, Prigge M, Printz Y, Yizhar O (2017) Silencing Neurons: Tools, Applications, and Experimental Constraints. Neuron 95:504–529.

Xiao L, Chattree G, Oscos FG, Cao M, Wanat MJ, Roberts TF (2018) A Basal Ganglia Circuit Sufficient to Guide Birdsong Learning. Neuron 98:208-221.e5.

Yamamoto H, Abe K (2022) Protocol for viral vector-mediated measurement of

transcription factor activity of mouse brain. STAR Protocols 3:101633.

Yu C, Jiang TT, Shoemaker CT, Fan D, Rossi MA, Yin HH (2022) Striatal mechanisms of turning behaviour following unilateral dopamine depletion in mice. European Journal of Neuroscience 56:4529–4545.