1   Title:
2   Mitochondrial heterogeneity disrupts osteoclast differentiation and bone resorption by
3   impairing respiratory complex I
4

Houfu Leng[1, #], Jiahao Jiang [2, #], Katja Gassner[3], Swati Midha[1,4,5,6], Raquel Justo-Méndez[3],
Jianqing Zheng[1], Timothy Hall[7], Lin Luo[1,8], Suzanne D West[9], Tonia L. Vincent[1], Angus Wann[1,4],
Kashyap A Patel[7], Joanna Poulton[10], Chris A. O'Callaghan[2], Ana Victoria Lechuga-Vieco[1,3,12],
Katja Simon[1,11,12]
[1-] Kennedy Institute of Rheumatology, University of Oxford, Roosevelt Drive, Oxford, OX3 7FY,
UK.
[2-] Centre for Human Genetics, Nuffield Department of Medicine, Wellcome Centre for Human
Genetics, University of Oxford, United Kingdom.
[3-] Institute for Research in Biomedicine (IRB Barcelona). The Barcelona Institute of Science and
Technology, Baldiri Reixac, 10, 08028 Barcelona, Spain
[4-] School of Biological Sciences, University of Southampton, Southampton, Hampshire, SO17
1BJ UK
[5-] Centre for Predictive In Vitro Models, Queen Mary University of London, UK
[6-] Centre for Bioengineering, School of Engineering and Materials Science, Queen Mary
University of London, UK
[7-]Department of Clinical and Biomedical Sciences, Faculty of Health and Life Sciences,
University of Exeter, Exeter, UK.
[8-] CAMS Oxford Institute, Nuffield Department of Medicine, University of Oxford, Oxford, UK
[9-] NHS Highly Specialised Service for Rare Mitochondrial Disorders, Oxford Centre for Genomic
Medicine, Oxford University Hospitals, Oxford, UK
[10-] Department of Women's and Reproductive Health, The Women's Centre, University of
Oxford, Oxford, United Kingdom
[11-] Max-Delbrück-Center for Molecular Medicine, Berlin, Germany
[12-] Corresponding Authors
[#]Contributed equally

30
Summary:

Mitochondria are required to meet the high energy demands of osteoclast differentiation and
adapt to hypoxia, inflammatory signals and oxidative stress. Heteroplasmy, the coexistence of
different mitochondrial DNAs (mtDNAs) in one cell disrupts energy balance and causes
metabolic stress. Using a genetically engineered mouse model, we demonstrate that
heteroplasmy impairs osteoclast differentiation and subsequently reduces bone resorption.
Mechanistically, we find that heteroplasmic cells accumulate complex I subunits of the electron
transport chain in the cytoplasm, which severely disrupts ATP production and leads to
premature termination of osteoclastogenesis. Importantly, we show that stimulating
autophagy and improving mitochondrial quality using spermidine alleviates the dysfunction in
both the murine heteroplasmic model and in human heteroplasmic osteoclasts carrying
m.3242A>G mutations, partially through enhanced mitochondrial selection. Overall, our work
highlights the role of mtDNA in osteoclast function and introduces potential therapeutic
approaches for mitochondrial and bone disorders.

45
46   **Keywords**: Complex I, osteoclasts, mitochondrial DNA, heteroplasmy, spermidine, autophagy
47   **Graphical Abstract**

Introduction

Skeletal homeostasis relies on a network of cells in the bone, namely the bone matrix synthesizing osteoblasts (OB) and the bone resorbing osteoclasts (OC). OC are terminally differentiated multinucleated cells originating from myeloid progenitors in the bone marrow. Engagement of RANKL (receptor activator of nuclear factor kappa-B ligand) mainly secreted by OBs with its cognate receptor RANK expressed on OC precursor cells is required for mature OC formation[1]. OC are vital for bone remodeling, deficiency of which can cause osteopetrosis whilst overactivation can cause osteoporosis or Paget Disease[2]. The demanding environmental conditions during osteoclastogenesis require robust metabolic activity to support differentiation and function[3]. Disruptions in the energy metabolism during OC differentiation cause an imbalance in skeletal homeostasis, leading to the onset and progression of bone-related diseases[3]. The energy required for OC differentiation mainly derives from glycolysis and mitochondrial oxidative metabolism[4,5]. Additionally protons (H+) generated by ATP hydrolysis are required to support acidification and bone collagen degradation[6,7].

The generation of new mitochondrial DNA and proteins has been linked to OC differentiation and function. Indeed, mature OCs have bigger and a larger number of mitochondria, elevated mitochondrial oxygen consumption rates and higher levels of the electron transport chain enzymes[4,8]. Notably, RANKL, has been shown to stimulate mitochondrial biogenesis[8,9] . The extensive transcriptional and metabolic reprogramming and intact mitochondrial function are essential for osteoclast differentiation and to demineralize the bone. The absence of Ndufs4, a subunit of complex I in the mitochondrial electron transport chain (mtETC) encoded by the nuclear genome, has been associated with impaired osteoclast differentiation and function, leading to increased bone density[10]. In addition, it has been reported that ECSIT, a complex I-associated protein, plays a central role in the mitochondrial response to early signaling of RANKL[9].

Although the role of mitochondria in osteoclasts (OCs) is well established, it remains unclear how OCs adapt to mutated mitochondrial DNA (mtDNA). This adaptation becomes particularly relevant in the context of heteroplasmy, the coexistence of multiple mtDNA variants within a cell. Bone tissue generally exhibits low levels of heteroplasmy, which may contribute to the limited research on the functional impact of mtDNA variation in bone-residing cells[11]. While mitochondrial point mutations are present in healthy individuals they are commonly associated with mitochondrial diseases. The pathological m.3243A>G mitochondrial point mutation in the MT-TL1 gene was first identified three decades ago and causes the MELAS syndrome (mitochondrial encephalopathy with lactic acidosis and stroke-like episodes)[12,13]. Over time, a better understanding of the disease has revealed an expanding spectrum of extra-neurological manifestations[14]. Primary mitochondrial diseases have been associated with poor bone health[15] and hearing loss due to the temporal bone structure damage, which has been observed in MELAS disease[16,17]. However, due to the complex genetics, phenotypic variability of the disease, and pathogenic molecular mechanisms being unclear, these associations are difficult to ascertain. To study mitochondrial diseases, the genetic modification of

mitochondrial DNA in mammalian cells would be ideal but this is still highly challenging[18]. To
overcome this, a heteroplasmic mouse model was developed on the C57BL/6 background,
carrying a variable mixture of two mtDNA variants, C57 and NZB. NZB mtDNA differs from C57
mtDNA by several mitochondrial single nucleotide polymorphisms (SNPs), including 12
missense mutations, 4 tRNA mutations, 8 rRNA mutations, and 10 mutations in non-coding
regions, making the NZB variant more efficient at oxidative phosphorylation[19]. When both
variants coexist, cells selectively retain the preferred mitochondria, leading to varying degrees
of heteroplasmy across different tissues and cell types[20]. This process of mitochondrial
selection in postnatal individuals closely mirrors the mutant load distribution observed in
tissues affected by primary mitochondrial diseases[11].

Here we address how mtDNA variability impacts osteoclastogenesis both *in vivo* and *ex vivo*,
using bone marrow samples from heteroplasmic mice and peripheral blood mononuclear cells
(PBMCs) from m.3243A>G patients. We find that mitochondrial DNA heterogeneity in mice
leads to increased bone density attributed to impaired OC formation and function.
Heteroplasmic osteoclasts exhibit impaired function due to reduced oxidative phosphorylation
and ATP production, resulting from the failure of nuclear-encoded complex I subunits to be
imported into mitochondria during the early stages of OC differentiation. We show that
preserving mitochondrial quality through spermidine treatment enhances osteoclast
differentiation from m.3243A>G patients *ex vivo* and restores osteoclast function in
heteroplasmic mice. Our study highlights the crucial role of preserving mitochondrial function
under conditions of mtDNA heterogeneity to maintain skeletal homeostasis. This offers a
promising strategy for treating age-related and mitochondrial disease-associated bone
pathologies.

Results

BL/6$^{C57-NZB}$ mice exhibit increased bone density due to diminished osteoclast function

To investigate if BL/6$^{C57-NZB}$ mice display changes in their bone mass, tibiae were collected from
aged-matched male BL/6^{C57}, BL/6^{NZB} and BL/6$^{C57-NZB}$ mice for micro-CT analysis. BL/6$^{C57-NZB}$ mice
showed an increase in trabecular bone (Fig. 1A). Accordingly, when we quantified bone
parameters, we found a 1.3-fold increase of bone volume over total volume (BV/TV) and a 1.2-
fold increase of trabecular thickness (Tb. Th), accompanied by a significant decrease in bone
surface density (BS/BV) and in trabecular separation (Tb. Sp). The decrease in the BS/BV ratio
in BL/6$^{C57-NZB}$ group was due to a significant increase in BV, as compared to BS. (Fig. 1B-C).
Furthermore, a significant increase of cortical thickness (Ct. Th) and cortical area (Ct. Ar) was
observed in BL/6$^{C57-NZB}$ group (Fig1. C), confirming an increase of the overall bone parameters.
Next, we investigated whether these increased bone parameters lead to a corresponding
increase in bone strength by measuring resistance to fracture. We conducted biomechanical
compression testing of the femur from BL/6^{C57} mice, BL/6^{NZB} mice and BL/6$^{C57-NZB}$ mice (Fig. 1D).
Compared with bones from BL/6^{C57} mice, those from BL/6$^{C57-NZB}$ mice needed 25% more

maximum load to break (Fig. 1D). Therefore, the femur from BL/6[C57-NZB] had increased bone
mechanical strength, resulting in greater resistance to deformation or fracture. This phenotype
may stem from reduced osteoclast-mediated bone resorption or enhanced osteoblast-driven
bone formation.

We first investigated whether bone-resorbing OCs are affected in BL/6[C57-NZB] mice, longitudinal
cross-sections of tibiae were stained with tartrate-resistant acid phosphatase (TRAP) specific
for OC. According to bone histomorphometry, tibiae from BL/6[C57-NZB] mice had significantly
lower signal for TRAP-positive OC when compared to those from BL/6[C57] control mice (Fig. 1E-
F). Consistently, a significant reduction of serum carboxy-terminal cross-linked telopeptide of
type 1 collagen (CTX-1) level was observed in BL/6[C57-NZB] mice, indicating that a lower level of
systemic OC activity and of bone resorption (Fig. 1G). However, no difference was observed in
the frequency of bone marrow OC precursors as demonstrated by expression of surface
markers (CD11b$^-$CD3$^-$B220$^-$CD115$^+$) between the 3 genotypes, indicating a diminished OC
differentiation from BL/6[C57-NZB] OC precursors (Supplementary Fig. 1A). No differences were
observed in serum PINP (serum procollagen type I N-propeptide), an osteoblast-derived
marker reflecting bone formation, among the three groups, suggesting that the observed
increases in bone volume and strength in BL/6[C57-NZB] mice were due to decrease in OC number
and function rather than an increase in osteoblast formation (Fig. 1G).

To confirm that OC and not OB mediate the bone phenotype, we performed bone marrow
chimera experiments, in which the host provides the OB and the donor the OC, as the latter
are of hematopoietic origin. Lethally irradiated BL/6[C57-NZB]and BL/6[C57] mice were transplanted
with BM donor cells from either BL/6[C57] or BL/6[C57-NZB] mice. Bone marrow engraftments were
monitored by bleeding every 2 weeks, with the donor cells expressing the CD45.2 isoform while
the host was CD45.1$^+$ to track donor and host cells (Supplementary Fig. 1B–C). Notably, we
found increased bone density, as measured by BV/TV micro-CT analysis (Fig. 1H-I and
Supplementary Fig. 1D), regardless of the recipient strain. These findings indicate that the
increased bone volume in BL/6[C57-NZB] mice is driven by impaired differentiation and function of
OC which are of hematopoietic origin.

Osteoclastogenesis is significantly compromised during the first days of the differentiation process in heteroplasmic cells

To determine the extent to which osteoclastogenesis contributes to the phenotype observed
in BL/6[C57-NZB] mice, we conducted *in vitro* OC differentiation assays. Bone marrow cells from
the three strains were cultured *in vitro* for 2 days with murine macrophage colony stimulating
factor (M-CSF) to generate macrophages (day 0), followed by culture with murine M-CSF and
RANKL for 5 days to induce OC formation. Notably, bone marrow cells from BL/6[C57-NZB] mice
showed a significant reduction in TRAP-positive OC formation (Fig. 2A)[1].

To understand the molecular mechanisms underlying the differences in osteoclastogenic
capacity across the different mitochondrial genotypes we conducted single-cell RNA
sequencing (scRNA-seq) at various stages of OC differentiation (Fig. 2B). Bone marrow cells

from the three strains were cultured for 2 days with M-CSF to generate macrophages (day 0), followed by culture with M-CSF and RANKL for either 1 day or 3 days to induce OC formation. Successful *in vitro* OC differentiation was validated by cross-referencing reported marker genes (e.g *Acp5* and *Ctsk* in Fig. 2C) and negative regulators of OCs following RANKL treatment (e.g. *Irf8* and *Mafb*)[1] (Supplementary Fig.2A). After applying quality control filters, we analyzed a total of 10,809 cells, comprising 4,028, 3,742 and 3,039 cells on day 0, day 1 and day 3, respectively (Fig. 2D and Fig. 2E). Unsupervised graph-based clustering of these single-cell transcriptomic profiles defined 9 distinct cell populations, including a monocyte-like precursor population and a terminal osteoclast population (Fig. 2D). Other transitional populations included *Cxcl16*^{hi} macrophages, dendritic cells, early- and late-stage pre-osteoclasts, and three proliferative populations (Fig. 2D).

To further infer the linkage between each cell population, we computed a per-cell latent time (Supplementary Fig. 2B) and performed partition-based graph abstraction (PAGA) analysis based on the RNA-velocity dynamics[21,22] (Supplementary Fig. 2C) . These analyses revealed a cell trajectory during osteoclastogenesis, which progresses from monocytic precursors to osteoclasts, passing through *Cxcl16*^{hi} macrophages, early-stage pre-osteoclasts and late-stage pre-osteoclasts (Supplementary Fig. 2B-C). Along this trajectory, we noted that BL/6$^{C57-NZB}$ cells were relatively concentrated in Cxcl16^{hi} macrophages (17.4%), and early-stage pre-osteoclasts (8.0%) compared to homoplasmic BL/6^{C57} (14.0% and 6.5%) or BL/6^{NZB} (14.3% and 6.5%) cells. Consequently, fewer heteroplasmic cells reached the mature osteoclast population (10.2% compared to 14.7% and 14.9% in BL/6^{C57} and BL/6^{NZB} cells, respectively) (Fig. 2E and Supplementary Fig. 2D). The statistical significance of this observation was further validated by differential abundance testing using MiloR [23] (Supplementary Fig. 2D).

Since osteoclastogenesis requires finely tuned cellular adaptive mechanisms[24], we hypothesized that heteroplasmic results in the generation of reduced numbers of mature cells and these are functionally compromised compared to control BL/6^{C57} osteoclasts. Indeed, pathway enrichment analysis revealed key differences in cellular respiration between the genotypes, including downregulation of the tricarboxylic acid cycle and of mitochondria-related functional pathways in differentiated osteoclasts (Fig. 2F). Additionally, there was a downregulation of cellular responses to hypoxic environments and associated oxidative stress, particularly reactive oxygen species. The lack of cellular adaptation to the demanding conditions occurring during the early days of OC differentiation was further validated by the fact that BL/6$^{C57-NZB}$ cells exhibit increased catabolic metabolism, despite the findings that adaptation to hypoxia in OC maturation typically requires enhanced cellular anabolism[25] (Fig. 2F). Collectively, our findings suggest that heteroplasmic cells face challenges in osteoclastogenesis, leading to premature arrest at the macrophage stage, which may be linked to altered metabolic adaptation and pathway dysregulation.

Alterations in complex I in heteroplasmic cells impair oxidative phosphorylation and metabolic adaptation during osteoclast differentiation.

We began with the premise that BL/6^{C57}, BL/6^{NZB} and BL/6$^{C57-NZB}$ cells share the same nuclear
genome but differ in mitochondrial DNA, and that changes in nucleo-mitochondrial crosstalk
could lead to alterations in the mitochondrial electron transport chain[20]. Since mitochondria
serve as a central metabolic hub, we analyzed gene expression of major metabolic pathways
during OC differentiation. At day 0 of *in vitro* OC differentiation, there were no significant
differences in metabolic pathways of BL/6$^{C57-NZB}$ cells compared to wild-type BL/6^{C57}; only few
pathways, including glycolysis, were upregulated at day 1 upon RANKL treatment in BL/6$^{C57-NZB}$
cells (Fig. 3A and Supplementary Fig. 3A). In contrast, after 3 days of differentiation, most
mitochondrial-associated pathways, including glutamate degradation pathway, adenosine
biosynthesis, galactose metabolism, pentose phosphate metabolism, ketogenesis, Krebs cycle,
and pyrimidine biosynthesis, were downregulated in the OC cultures from BL/6$^{C57-NZB}$ mice (Fig.
3A and Supplementary Fig. 3A). Specifically, we evaluated the activation of the oxidative
phosphorylation (OXPHOS) pathway by aggregating the average gene expression of all involved
genes (n = 107 genes as annotated in the KEGG database). OC differentiation increased the
cellular energy demand, as indicated by the elevated OXPHOS scores in pre-osteoclasts and
mature osteoclasts (Fig. 3B). The significant reduction in the OXPHOS score observed in OCs
(Cluster 2, representing the most mature osteoclastic cells) derived from BL/6$^{C57-NZB}$ cells
suggests a disruption in oxidative phosphorylation and potentially compromised mitochondrial
metabolic performance (Fig. 3B).

To validate the scRNA-seq findings, we used Seahorse technology to assess the oxygen
consumption rate in BL/6^{C57}, BL/6^{NZB} and BL/6$^{C57-NZB}$ cells (Fig. 3C and Supplementary Figure 3B).
Consistent with gene expression data, no differences were observed at day 0. However, by day
1 of OC differentiation, cells from BL/6$^{C57-NZB}$ mice exhibited a reduction in maximal
mitochondrial respiratory capacity (Fig. 3C and Supplementary Figure 3B) alongside an increase
in reactive oxygen species (ROS) and in mitochondrial mass, likely as a compensatory response
(Supplementary Figure 3C-D). These results provide further evidence of a compromised
OXPHOS capacity in heteroplasmic cells early in the OC differentiation process. However, by
day 3 following RANKL treatment, heteroplasmic cells exhibit lower mitochondrial mass while
maintaining the same spare respiratory capacity (SRC) (Supplementary Figure 3B-C), likely by
utilizing alternative substrates and relying on complex II (succinate dehydrogenase)[26]. We
analyzed the expression of selected OXPHOS structural subunits by western blot on days 0, 1,
2, 3, and 5 of OC differentiation. Interestingly, each genotype displayed a distinct temporal
pattern in the relative abundance of the proteins analyzed. Thus, wild-type BL/6^{C57}cells,
showed a sequential increase in NDUFB8 (complex I) and SDHB (complex II) expression levels
during differentiation with a peak at day 3 (Supplementary Figure 3E). UQCR2 (complex III) and
MTCOI (CIV) changes are less prominent, remaining elevated all the period. Finally, ATP5A
(complex V) peaked acutely at day 5 (Supplementary Figure 3E). On the other hand, BL/6^{NZB}
cell differentiation resemble that of the previous cells with a sharpen peak of SDHB at day 3
and less intense ATP5A increase (Supplementary Figure 3E). In contrast, BL/6$^{C57-NZB}$ cells
showed a mitigated response in NDUFB8 expression after day 3 (Supplementary Fig. 3E),
supporting previous findings (Fig. 3A). Heteroplasmic cells also showed a very mild increase in

SDHB expression. Together these differences lead to a relevant drop in the relative activation between complex II and complex I representative proteins. Very interestingly, these changes in expression correlated with the significantly higher complex I/II activity ratio characteristic of day 3 BL/6[C57-NZB] cells (Fig. 3D). We then address how this differentiation impacted on ATP synthesis and found that BL/6[C57-NZB] cells exhibited reduced efficiency in ATP production (Fig. 3E). This analysis examined changes in expression on a per-cell basis. Interestingly, when analyzed per mitochondria, heteroplasmic cells show lower levels of NDUFA9, an nDNA-encoded subunit of respiratory complex I, compared to BL/6[C57] and BL/6[NZB], after day 3 of *in vitro* OC differentiation. (Fig. 3F-G). In contrast, ATPβ, a subunit of Complex V (ATPase) responsible for ATP synthesis, showed no differences between genotypes in expression per mitochondrion on day 3 across the different genotypes (Fig. 3F-G). Altogether, the data suggest that dysregulation of complex I may be the primary factor affected in OC differentiation in heteroplasmic cells.

OC differentiation requires sequential import of complex I subunits

Thus, complex I became a key focus of our analysis as it also harbors 14 of the SNPs between the C57 and NZB haplotypes[19].

The upregulation of genes encoded in the mitochondrial genome is not apparent from day 0 to day 1 but becomes evident between day 1 and day 3 in cells derived from wild-type BL/6[C57] mice (Supplementary Fig. 4A). Nuclear-encoded mitochondrial proteins are synthesized by cytosolic ribosomes as precursors and need to be imported into mitochondria[27]. Among them, the complex I subunit NDUFA9 harbors an N-terminal cleavable pre-sequence that facilitates its mitochondrial targeting[28]. We used confocal imaging to determine whether nuclear-encoded subunits of the respiratory complexes were properly imported and successfully assembled into functional respiratory complexes within mitochondria. Strikingly, we observed that a significant proportion of NDUFA9 subunit remains in the cytoplasm on day 1 of differentiation of OC from wild-type BL/6[C57] BL/6[NZB] and BL/6[C57-NZB] mice (Supplementary Fig. 4B). Intriguingly, from day 3 to day 5, the colocalization of complex I with mitochondria increases in wild-type cells, suggesting that greater complex I respiratory complex formation is required during OC development (Fig. 4A-B). However, this increase does not happen to the same extent in BL/6[C57-NZB] cells (Fig 4B). By day 5, this colocalization decreases, and prominent localization of the nDNA-encoded subunit within mitochondria is observed only in wild-type BL/6[C57] osteoclasts (Fig. 4B).

Previous studies indicate that complex I inhibition disrupts mitochondrial protein import, impairs chaperone function, and activates the mitochondrial unfolded protein response (mtUPR) to prevent aggregation of respiratory complex subunits[29]. The analysis of gene expression changes related to the mitochondrial protein import machinery and mitochondrial degradome at day 3 of OC differentiation revealed their pronounced downregulation in BL/6[C57-NZB] cells compared to BL/6[C57] cells (Fig. 4C-D). Some of our previous data already indicate a downregulation of the assembly proteins for respiratory complex I (Figure 2F),

suggesting that the failure to complete protein import into mitochondria leads to decreased
respiratory complex assembly factors in BL/6[C57-NZB] cells compared to BL/6[C57] (Fig. 4E).

Altogether, nuclear-encoded complex I subunits fail to be either imported into mitochondria
or properly degraded in the cytoplasm during OC differentiation in BL/6[C57-NZB]. Both processes
require ATP[27,30,31], which is reduced with heteroplasmy at the stage when OCs would normally
depend on respiratory complex I formation for mitochondrial function.

Mitochondrial selection in heteroplasmic cells is evident during osteoclastogenesis

We previously reported that mitochondrial heteroplasmy can be resolved through non-
random mtDNA segregation specific to tissue and cell types[20]. Considering the substantial
changes in metabolic profiles during osteoclastogenesis[32], we reasoned that the successfully
differentiated osteoclasts might not represent the same heteroplasmic levels as compared to
monocytic precursors. To genotype mitochondrial variants at single-cell resolution during early
stages of OC differentiation, we adapted the mgatk pipeline[33] with deep sequencing of the
scRNA-seq libraries. Due to the relatively low coverage for individual variants, we aggregated
all NZB mtDNA SNPs and calculated an overall mtDNA NZB percentage for each single cell on
day 1 and day 3 of *in vitro* OC differentiation of cells from BL/6[C57-NZB] mice. Cells with low
coverage were subsequently filtered out, leaving an average of 237 sequencing reads per cell
covering any NZB mtDNA variants (Supplementary Fig. 5A). We observed a higher level of NZB
mtDNA transcripts on day 3 compared to day 1 cells, which was independent of the sequencing
depth (Fig. 5A). The percentage of NZB mtDNA slightly increased along the OC differentiation
trajectory (Supplementary Fig. 5B). We further validated these findings using PCR followed by
RFLP (Restriction Fragment Length Polymorphism) analysis in OC cultures at Days 1 and 3. We
targeted the G4276A nucleotide, which forms part of a BamH1 restriction site in C57 mtDNA
that is absent in NZB mtDNA[34,35,36]. Our data showed a shift toward more NZB mtDNA as OC
differentiation progressed (Fig. 5B & Supplementary Fig. 5C). In line with these results, single-
cell mitochondrial genotyping demonstrates the heterogeneity of mitochondrial heteroplasmy
across different clusters, with mature osteoclasts showing significant enrichment for the NZB
mtDNA haplotype (Fig. 5C).

Heteroplasmy leads to exhausted autophagy and OC maturation can be rescued by restoring spermidine levels

The downregulation of the mitochondrial proteases seen in the OC degradome analysis in
heteroplasmic OC cultures (Fig. 4D) may lead to impaired organelle quality control. To
determine whether the selective removal of mitochondria with distinct mtDNA haplotype
profiles within the cytoplasm of BL/6[C57-NZB] cells affects mitochondrial recycling during
osteoclastogenesis, we analyzed the autophagic flux throughout the OC differentiation process.
On day 1, BafA1 treatment increased LC3-II levels and decreased the LC3-I/LC3-II ratio,
indicating strong autophagic flux across all genotypes (Fig. 5D–E). However, by day 3, BafA1,
that was used to block autophagosome-lysosome fusion and assess autophagy flux, no longer
increased LC3-II in BL/6[C57-NZB] cells (Fig. 5D–E). This, together with elevated p62 levels in

heteroplasmic cells, even in the absence of BafA1 (Supplementary Fig. 5D), suggests a block in autophagic flux at day 3. Consistent with this, the expression of *Tfeb,* a key transcription factor involved in autophagy[37] and in the regulation of respiratory complex I[38] was downregulated from monocyte precursors to OCs in *in vitro* OC differentiation cultures derived from both BL/6^{C57} and BL/6^{NZB} mice, but not from BL/6$^{C57-NZB}$ mice (Supplementary Fig. 5E). Overall, this suggests that autophagy is regulated differently with heteroplasmy over the course of differentiation and may contribute to the metabolic phenotype of heteroplasmic OC.

Next, we reasoned that the metabolite spermidine may enhance differentiation of heteroplasmic OCs. We have previously demonstrated that autophagy is regulated by endogenous spermidine levels which control translation of TFEB [39,40]. Spermidine activates mitochondrial translation in myeloid cells[41], and boosts oxidative metabolism by binding to the mitochondrial trifunctional protein (MTP) in T cells[42]. Spermidine given to mice in the drinking water restores autophagy levels when these are low[37]. First, we performed OC differentiation of cells from heteroplasmic mice with spermidine supplementation in the media. Spermidine restored osteoclast formation *in vitro* in heteroplasmic cells compared to untreated cultures (Fig. 5F-G). We also observed a significant increase in NZB mtDNA selection on day 5 (Fig. 5H) without any change in mtDNA content (Supplementary Fig. 5F). The dose used to observe this phenotype in heteroplasmic cells has previously been shown to inhibit OC formation in wild-type cells[43], a finding we reproduced in our analyses (Fig. 5F).

While no changes in total ATP were observed in the spermidine-treated OC cultures on day 1 of differentiation (Supplementary Fig. 5G), treated heteroplasmic cells showed a sharp increase in ATP content on day 3 (Fig. 5I). We observed an increase in mitochondrial complex I expression through confocal imaging in spermidine-treated versus untreated OC cultures from BL/6$^{C57-NZB}$ mice on day 5 (Fig. 5J & Supplementary Fig. 5H-I).

To investigate if spermidine can modulate the phenotype of heteroplasmic OC *in vivo*, we generated BM chimeric mice by transplanting BM donor cells from wild-type BL/6^{C57} or BL/6$^{C57-NZB}$ mice into lethally irradiated BL/6$^{C57-NZB}$ recipients, ensuring that all recipient mice share the same host tissue background and avoiding differences in the host environment that could influence the phenotype. After 8 weeks of spermidine treatment (5 mM in drinking water), bone phenotyping analyses using micro-CT demonstrate that in BL/6$^{C57-NZB}$ BM donor cells spermidine administration led to a decrease of BV/TV in mice, accompanied by an increase in Tb. Sp, confirming an activation of bone catabolism (Fig. 5K-L). Lastly, spermidine-treated chimeric BL/6$^{C57-NZB}$ mice showed decreased bone mechanical strength, reversing the phenotype to resemble wild-type BM (Fig. 5M). Taken together, OC differentiation in heteroplasmic cells is linked to a shift in mitochondrial heteroplasmy toward NZB mtDNA variants. By restoring spermidine levels that may have been depleted due to the extra demand during differentiation, osteoclastogenesis was restored, both *in vitro* and *in vivo*.

m.3243A>G heteroplasmy disrupts human osteoclast differentiation, which can be restored by spermidine

To evaluate the potential for osteoclast formation of cells from people with heteroplasmic mitochondrial disease, we collected blood from middle-aged people with MELAS carrying both healthy and m.3243A>G mutated mtDNA within their cytoplasm. This mutation has been linked previously to bone abnormalities[44]. Using peripheral blood mononuclear cells (PBMCs) from age- and sex-matched healthy donors and people with m.3243A>G heteroplasmy, we established an *in vitro* OC differentiation experiment. Briefly, human PBMCs were isolated on day -3 and were seeded with human M-CSF. On day 0, the cells were induced to differentiate into OCs using human M-CSF and human RANKL. By day 6, mature osteoclasts had formed and were fixed for TRAP analysis. Consistent with our findings in mouse heteroplasmic cells (Fig. 2A), the formation of mature osteoclasts was significantly reduced in m.3243A>G cells (Fig. 6A). Additionally, serum levels of the bone resorption marker CTX-1 were lower in m.3243A>G cells, further indicating a systemic osteoclast deficiency (Fig. 6B). Next, we evaluated the effect of spermidine in human cells on *in vitro* osteoclast differentiation in healthy and m.3243A>G cells (Fig. 6C-D). Treatment with 5 µM spermidine increased the number of TRAP+ osteoclasts (Fig. 6C) and ATP levels (Fig. 6D) in both groups. To further explore the association between m.3243A>G and bone mineral density (BMD) in humans, we analyzed data on 271,849 individuals from the UK Biobank cohort. Our study included individuals with BMD measurements by ultrasound (n = 271,849) and Dual-energy X-ray Absorptiometry (DXA) imaging (n = 46,777), along with whole genome sequencing data. Using whole genome sequencing data, we identified 76 individuals with m.3243A>G at >5% blood heteroplasmy for whom BMD data were available and 15 with DXA imaging (Supplementary Table 2). Linear regression analysis, adjusted for relevant covariates, revealed no statistically significant association between the presence of m.3243A>G and BMD measurements from heel, femoral neck, spine (L1-L4), or total body (all p > 0.19) (Supplementary Table 3). Similarly, we found no significant association with self-reported fractures (OR 0.528, p = 0.245) but confirmed a known strong reported association with diabetes (OR 8.07, p = 8.22×10^{-15}) (Supplementary Table 4). A more sensitive analysis, limiting cases to those with >10% heteroplasmy, yielded similar results (Supplementary Table 4). These results suggest that the blood levels of the m.3243A>G variant may not significantly impact BMD or fracture risk in this population cohort.

Discussion

This work investigates how mitochondrial heteroplasmy disrupts OC differentiation by impairing cellular metabolism and the proper assembly of the mtETC. Using a genetically engineered mouse model, we showed that heteroplasmy prevents complex I subunit import into mitochondria, leading to premature termination of osteoclastogenesis and reduced bone resorption. We further demonstrate that spermidine can partially rescue OC function in both heteroplasmic mice and heteroplasmic human cells carrying the m.3242A>G mutation. Our findings highlight the crucial role of mtDNA in regulating OC differentiation, particularly under challenging metabolic demands.

The metabolic requirements for OC differentiation and function are well studied. During OC formation, glycolysis and glucose uptake via GLUT1 are increased, whereas the knockdown of hypoxia inducible factor 1 alpha (Hif1a) or glucose deprivation inhibit OC function[45]. Specific knockouts of respiratory complex I subunits result in deficiencies in OC maturation[10]. While *Glut1*-mediated glucose metabolism through OXPHOS is critical for osteoclastogenesis, lactate production from glucose is also necessary [46]. Hence, both glycolysis and oxidative phosphorylation are required for OC formation. In fact, they may have two distinct key roles: while oxidative phosphorylation appears to be the primary source for OC formation, glycolytic pathway-related enzymes are distributed near the actin ring, where energy-demanding activities associated with bone matrix degradation take place[4]. In conclusion, metabolic dependencies differ between OC differentiation and active bone resorption: the formation of OCs requires OXPHOS, while bone resorption is dependent on glycolysis[4,6]. In this study, we examined how impaired osteoclast (OC) differentiation occurs when mitochondrial subunits originate from different mitochondrial DNA variants. Our findings in heteroplasmic OC precursors revealed increased complex I activity without a corresponding increase in ATP production. This may be explained by dysfunction in downstream complexes, where electrons donated by complex I are inefficiently transferred through the electron transport chain, leading to an accumulation of electrons and reduced oxidative phosphorylation efficiency, ultimately limiting ATP output. Elevated reactive oxygen species (ROS) levels or shifts in metabolic pathways could further contribute to impaired ATP production.

The assembly of mitochondrial respiratory complex I in mammals requires the coordinated integration of at least 44 subunits encoded by either nuclear and mitochondrial genomes[50]. The presence of complex I subcomplexes (intermediate state) in complex I-deficient cells (i.e. when the complex is not assembled properly) indicates a sequential assembly process[51, 52 ,28]. This involves the complete import of nuclear-encoded subunits into mitochondria, degradation of excess cytoplasmic material, and ultimately, the proper assembly and superassembly of respiratory complex I in the mitochondrial lumen. Our data suggests that mitochondrial protein subunits, in particular complex I, accumulate as unimported mitochondrial proteins in the cytoplasm during osteoclast differentiation. Osteoclastogenesis is a tightly regulated process that demands ATP not only for biomass generation but also for metabolic adaptation to substrate availability at different stages. This process unfolds in a hypoxic tissue environment[47]. Intriguingly, hypoxia-induced mitochondrial stress granules have been identified in C. elegans[48]. Similarly during aerobic growth, yeast cells have been shown to accumulate mitochondrial precursor proteins as MitoStores in the cytoplasm [49]. Hence, one explanation for complex I being retained in the cytoplasm may be that this temporary storage site ensures their rapid import into mitochondria during the later stages of differentiation, particularly under challenging nutritional conditions or hypoxic induction. However, unassembled complex I accumulate in the cytoplasm and are bound for degradation. Previous studies in mammalian cells suggest that stress granules can be cleared either through an autophagy-independent disassembly process or via autophagy-dependent degradation[53, 54]. The impaired autophagic

flux observed in the latter stages during OC differentiation in heteroplasmic BL/6[C57-NZB] cells
may contribute to the cytoplasmic accumulation of complex I.

Mitochondrial heteroplasmy presents a largely unexplored area that may influence cellular
function and bone health. Spermidine has been proposed to enhance autophagy[40,55],
translational activity[37] and mitochondrial metabolism[42,56,57]. Collectively, these findings
highlight spermidine as a promising candidate for improving mitochondrial quality. We and
others have found that in both mice and humans a moiety of spermidine is needed to activate
the translation factor eif5a by hypusination in order to increase autophagic flux and lysosomal
activity[37,58]. Additionally, this translation factor is crucial for mitochondrial translation[41], and in
T cells boosts oxidative metabolism by binding to the mitochondrial trifunctional protein
(MTP)[42]. This implies that cellular spermidine levels may run low if this pathway is repeatedly
triggered such as in heteroplasmy. Spermidine may need to be replenished or else an
'exhausted type' of autophagic flux occurs where LC3- II is no longer degraded, as we found
here. We also observed that in wild-type cells spermidine inhibits OC differentiation. This is in
line with the study that showed supplementation with spermine or spermidine enhances bone
strength by inhibiting osteoclasts, whereas inhibiting polyamine biosynthesis reduces these
benefits[43]. The differentiation of OC, like that of most cell types, requires autophagy and
spermidine may help under homeostatic conditions. In contrast, in heteroplasmic mice, we
found that spermidine enhanced OC differentiation at the dose that was previously reported
to inhibit the process in wild-type cells[43]. Similar results have been observed in Nrf2 knockout
cells[59], suggesting that spermidine levels may be critically low and need to be replenished
under certain stressful situations such as excessive ROS, repeated clearance of either
mitochondria or unwanted complexes I in the cytoplasm.

The association of bone density in people with the m.3243A>G mutation remains inconclusive.
Several factors could explain this outcome. First, nuclear genetic variability, which influences
osteoclast dysfunction, may act as a confounding factor. Our findings were based on inbred
mouse strains, where genetic uniformity allowed clearer interpretation of mitochondrial
effects, and maladaptive interactions are more easily detected. In mouse models, a
mitochondrial-encoded protein can aberrantly interact with a nuclear-encoded protein from a
different genetic background, leading to alterations on the function. In contrast, the genetically
diverse human population introduces a wide range of nuclear genotypes and associated
phenotypic variability, likely masking mitochondrial contributions and reducing the probability
of detecting an association without very large sample sizes. The number of m.3243A>G carriers
with bone measurements in the UK Biobank was very low, highlighting the need for larger
studies to confirm or refute this null association. Future studies should further investigate
these factors, including how nuclear genetic variation and osteoblast involvement contribute
to bone phenotypes in the context of mitochondrial mutations in humans.

Our findings demonstrate that spermidine treatment can restore the assembly process and
protein import leading to improved mitochondrial function in human *in vitro* OC differentiation,
as well as *in vivo* in mice. The m.3243A>G mutation in people with MELAS and the

heteroplasmic BL/6$^{C57-NZB}$ mice are comparable in terms of their mitochondrial genetic
variability, each harboring two distinct mtDNA profiles and corresponding mitochondrial
function profiles. Comparing these two models allowed us to identify shared mechanisms
underlying mitochondria-related abnormalities of osteoclastogenesis and led to the discovery
of spermidine as a modulator of this process. This finding opens the door for the potential use
of spermidine in human heteroplasmic conditions, a therapeutic possibility not previously
reported.

Acknowledgements
We acknowledge all Autophagy in the Immune System group members for their scientific
discussions contributing to this manuscript and Prof. Jose Antonio Enríquez (Spanish National
Center for Cardiovascular Research, CNIC, Madrid) who provided mouse models for our
research and providing thoughtful feedback. We thank Dr. Hanlin Zhang from University of
California, Berkeley for discussions and constructive comments. We thank Jianwei Cui from the
University of Oxford who helped with experiments. We thank Patricia Cotta Moreira and Mino
Medghalchi from the Kennedy Institute of Rheumatology animal facility, and the Parc Cientific
de Barcelona (PCB) animal facility for their excellent care and assistance of animal well-being.
We thank Thomas Conrad, Caroline Braeuning and Sarah Nathalie Vitcetz at the MDC Genomics
Platform (BIH, Charité Campus Mitte, Berlin) for discussions about transcriptomic experiments
and sequencing performed at their facility. Histology was performed in the Kennedy Institute
of Rheumatology Histology Facility by Dr. Ida Parisi. We obtained confocal microscopy
assistance from Christoffer Lagerholm, Senior Advanced Microscopy Manager at the Kennedy
Institute of Rheumatology, and the Advanced Digital Microscopy (ADM) Core Facility at IRB
Barcelona is greatly appreciated. We thank the NHS Highly Specialised Service for Rare
Mitochondrial Disorders (Oxford University Hospitals, UK) for the patient recruitment and
share clinical samples. We thank Dr. Elisabeth Greßler from Max-Delbrück-Center for
Molecular Medicine and Hantao Lou from the University of Oxford for experimental assistance.
Images were created using BioRender.com under the academic license granted to AKS.

This work was supported by: University of Oxford Medical and Life Sciences Translational Fund
MC_PC_17174 and MC_PC_18059 from Wellcome ISSF fund to HL. China Scholarship Council
fund (202006320024) to JJ. UMDF PF-23-0010 to RJ-M. Biotechnology and Biological Sciences
Research Council (BBSRC, BB/X007049/1) to SM. Kennedy Trust Prize Studentship (AZT00050-
AZ04) to JZ. Chinese Academy of Medical Sciences (CAMS) Innovation Fund for Medical
Sciences (CIFMS) 2018-I2M-2-002 to LL. Centre for OA Pathogenesis (Versus Arthritis 20205)
to TLV. KTRR KENN202111 to AW. Oxford Hospitals charity and Lily Foundation "Treatments
for Mitochondrial Disease – 2019/20" to JP. Salary support from the UK NHS Specialist
Commissioners who fund the "Rare Mitochondrial Disorders of Adults and Children" clinical
Service to SDW. Novo Nordisk Foundation (NNF0064142) to COC. This work is funded by
Diabetes UK (19/0005994 and 21/0006335), MRC (MR/T00200X/1) and Wellcome's
Institutional Strategic Support Fund awarded to University of Exeter; KAP and TH is funded by

Wellcome (219606/Z/19/Z); The work is supported by the National Institute for Health Research (NIHR) Exeter Biomedical Research Centre, Exeter, UK. European Molecular Biology Organization Postdoctoral Fellowship (EMBO) ALTF115-2019, "La Caixa" Foundation (ID 100010434, LCF/BQ/PI24/12040005) and funding from MINECO through the Centres of Excellence Severo Ochoa Award and CERCA Programme of the Generalitat de Catalunya to AVL-V. Wellcome Trust Fund 220784/Z/20/Z to AKS.

Author contributions

AKS, A.V.L.-V. and H.L. conceptualized and designed the study.

A.V.L.-V., H.L., and J.J. performed and analyzed experiments, methodology, and investigation.

AKS, A.V.L.-V., H.L. and J.J. discussed analyses.

AKS and A.V.L.-V. supervised the project and interpreted the experimental data.

AKS, A.V.L.-V., H.L. and J.J. wrote the original draft.

AKS, A.V.L.-V., H.L. and J.J. reviewed and edited the manuscript.

C.OC. planned experiments, analyzed data and edited the final manuscript.

K.G. performed mitochondrial DNA copy numbers, respiratory complexes activities and confocal imaging.

R.J-M. performed respiratory complexes analyses.

T.H. and K.A.P. investigated the association between m.3243A>G and bone parameters in individuals from the UK Biobank cohort.

J.Z. and T.V. contributed to the 3D reconstructions of bone images and image analysis.

L.L. performed the isolation of donor PBMC samples, followed by OC differentiation, and autophagy flux assays.

S.W. and A.W. analyzed the bone stiffness parameters.

S.D.W. and J.P. provided samples and contributed to discussions on human and patient data.

All authors reviewed and approved the final manuscript.

Competing interests

The authors declare no competing interests.

References

1. Leng, H. *et al.* Modulating glycosphingolipid metabolism and autophagy improves outcomes in pre-clinical models of myeloma bone disease. *Nat. Commun.* **13**, 7868 (2022).

2. Tao, X. *et al.* Clinical Characteristics and Pathogenic Gene Identification in Chinese Patients With Paget's Disease of Bone. *Front. Endocrinol.* **13**, 850462 (2022).

3. Ledesma-Colunga, M. G., Passin, V., Lademann, F., Hofbauer, L. C. & Rauner, M. Novel

Insights into Osteoclast Energy Metabolism. *Curr. Osteoporos. Rep.* **21**, 660–669 (2023).

4. Lemma, S. *et al.* Energy metabolism in osteoclast formation and activity. *Int. J. Biochem.*

Cell Biol. **79**, 168–180 (2016).

5. Taubmann, J. *et al.* Metabolic reprogramming of osteoclasts represents a therapeutic

target during the treatment of osteoporosis. *Sci. Rep.* **10**, 21020 (2020).

6. Jiménez, J. A., Lawlor, E. R. & Lyssiotis, C. A. Amino acid metabolism in primary bone

sarcomas. *Front. Oncol.* **12**, 1001318 (2022).

7. Motyl, K. J., Guntur, A. R., Carvalho, A. L. & Rosen, C. J. Energy Metabolism of Bone.

Toxicol. Pathol. **45**, 887–893 (2017).

8. Zeng, R., Faccio, R. & Novack, D. V. Alternative NF-κB Regulates RANKL-Induced

Osteoclast Differentiation and Mitochondrial Biogenesis via Independent Mechanisms. *J.*

Bone Miner. Res. **30**, 2287–2299 (2015).

9. Marques-Carvalho, A., Sardão, V. A., Kim, H.-N. & Almeida, M. ECSIT is essential for

RANKL-induced stimulation of mitochondria in osteoclasts and a target for the anti-

osteoclastogenic effects of estrogens. *Front. Endocrinol.* **14**, 1110369 (2023).

10. Jin, Z., Wei, W., Yang, M., Du, Y. & Wan, Y. Mitochondrial Complex I Activity Suppresses

Inflammation and Enhances Bone Resorption by Shifting Macrophage-Osteoclast

Polarization. *Cell Metab.* **20**, 483–498 (2014).

11. Naue, J. *et al.* Evidence for frequent and tissue-specific sequence heteroplasmy in

human mitochondrial DNA. *Mitochondrion* **20**, 82–94 (2015).

12. Goto, Y., Nonaka, I. & Horai, S. A mutation in the tRNALeu(UUR) gene associated with

the MELAS subgroup of mitochondrial encephalomyopathies. *Nature* **348**, 651–653

(1990).

13. Hirano, M. *et al.* MELAS: An original case and clinical criteria for diagnosis. *Neuromuscul.*

Disord. **2**, 125–135 (1992).

14. Ng, Y. S. & Gorman, G. S. Stroke-like episodes in adult mitochondrial disease. in

Handbook of Clinical Neurology vol. 194 65–78 (Elsevier, 2023).

15. Gandhi, S. S., Muraresku, C., McCormick, E. M., Falk, M. J. & McCormack, S. E. Risk

factors for poor bone health in primary mitochondrial disease. *J. Inherit. Metab. Dis.* **40**,

673–683 (2017).

16. Di Stadio, A. *et al.* Hearing impairment in MELAS: new prospective in clinical use of

microRNA, a systematic review. *Orphanet J. Rare Dis.* **13**, 35 (2018).

17. Handzel, O., Ungar, O. J., Lee, D. J. & Nadol, J. B. Temporal bone histopathology in

MELAS syndrome. *Laryngoscope Investig. Otolaryngol.* **5**, 152–156 (2020).

18. Fu, Y. *et al.* Engineering mtDNA deletions by reconstituting end joining in human

mitochondria. *Cell* S0092867425001941 (2025) doi:10.1016/j.cell.2025.02.009.

19. Latorre-Pellicer, A. *et al.* Mitochondrial and nuclear DNA matching shapes metabolism

and healthy ageing. *Nature* **535**, 561–565 (2016).

20. Lechuga-Vieco, A. V. *et al.* Cell identity and nucleo-mitochondrial genetic context

modulate OXPHOS performance and determine somatic heteroplasmy dynamics. *Sci.*

Adv. **6**, eaba5345 (2020).

21. Bergen, V., Lange, M., Peidli, S., Wolf, F. A. & Theis, F. J. Generalizing RNA velocity to

transient cell states through dynamical modeling. *Nat. Biotechnol.* **38**, 1408–1414

(2020).

22. Lange, M. *et al.* CellRank for directed single-cell fate mapping. *Nat. Methods* **19**, 159–

170 (2022).

23. Dann, E., Henderson, N. C., Teichmann, S. A., Morgan, M. D. & Marioni, J. C. Differential

abundance testing on single-cell data using k-nearest neighbor graphs. *Nat. Biotechnol.*

40, 245–253 (2022).

24. Tsukasaki, M. *et al.* Stepwise cell fate decision pathways during osteoclastogenesis at

single-cell resolution. *Nat. Metab.* **2**, 1382–1390 (2020).

25. Indo, Y. *et al.* Metabolic regulation of osteoclast differentiation and function. *J. Bone*

Miner. Res. Off. J. Am. Soc. Bone Miner. Res. **28**, 2392–2399 (2013).

26. Zieliński, Ł. P., Smith, A. C., Smith, A. G. & Robinson, A. J. Metabolic flexibility of

mitochondrial respiratory chain disorders predicted by computer modelling.

Mitochondrion **31**, 45–55 (2016).

27. Wachter, C., Schatz, G. & Glick, B. S. Protein import into mitochondria: the requirement

for external ATP is precursor-specific whereas intramitochondrial ATP is universally

needed for translocation into the matrix. *Mol. Biol. Cell* **5**, 465–474 (1994).

28. Lazarou, M., McKenzie, M., Ohtake, A., Thorburn, D. R. & Ryan, M. T. Analysis of the

assembly profiles for mitochondrial- and nuclear-DNA-encoded subunits into complex I.

Mol. Cell. Biol. **27**, 4228–4237 (2007).

29. Franco-Iborra, S. *et al.* Defective mitochondrial protein import contributes to complex I-

induced mitochondrial dysfunction and neurodegeneration in Parkinson's disease. *Cell*

Death Dis. **9**, 1122 (2018).

30. Thomas, H. E. *et al.* Mitochondrial Complex I Activity Is Required for Maximal

Autophagy. *Cell Rep.* **24**, 2404-2417.e8 (2018).

31. Kulawiak, B. *et al.* The mitochondrial protein import machinery has multiple connections

to the respiratory chain. *Biochim. Biophys. Acta BBA - Bioenerg.* **1827**, 612–626 (2013).

32. Park-Min, K.-H. Metabolic reprogramming in osteoclasts. *Semin. Immunopathol.* **41**,

565–572 (2019).

33. Lareau, C. A. *et al.* Massively parallel single-cell mitochondrial DNA genotyping and

chromatin profiling. *Nat. Biotechnol.* **39**, 451–461 (2021).

34. Latorre-Pellicer, A. *et al.* Regulation of Mother-to-Offspring Transmission of mtDNA

Heteroplasmy. *Cell Metab.* **30**, 1120-1130.e5 (2019).

35. Burgstaller, J. P. *et al.* MtDNA segregation in heteroplasmic tissues is common in vivo

and modulated by haplotype differences and developmental stage. *Cell Rep.* **7**, 2031–

2041 (2014).

36. Johnston, I. G. & Jones, N. S. Evolution of Cell-to-Cell Variability in Stochastic, Controlled,

Heteroplasmic mtDNA Populations. *Am. J. Hum. Genet.* **99**, 1150–1162 (2016).

37. Zhang, H. *et al.* Polyamines Control eIF5A Hypusination, TFEB Translation, and

Autophagy to Reverse B Cell Senescence. *Mol. Cell* **76**, 110-125.e9 (2019).

38. Calabrese, C. *et al.* Mitochondrial translocation of TFEB regulates complex I and

inflammation. *EMBO Rep.* **25**, 704–724 (2024).

39. Alsaleh, G. *et al.* Autophagy in T cells from aged donors is maintained by spermidine and

correlates with function and vaccine responses. *eLife* **9**, e57950 (2020).

40. Puleston, D. J. *et al.* Autophagy is a critical regulator of memory CD8+ T cell formation.

eLife **3**, e03706 (2014).

41. Puleston, D. J. *et al.* Polyamines and eIF5A Hypusination Modulate Mitochondrial

Respiration and Macrophage Activation. *Cell Metab.* **30**, 352-363.e8 (2019).

42. Al-Habsi, M. *et al.* Spermidine activates mitochondrial trifunctional protein and

improves antitumor immunity in mice. *Science* **378**, eabj3510 (2022).

43. Chevalier, C. *et al.* Warmth Prevents Bone Loss Through the Gut Microbiota. *Cell Metab.*

32, 575-590.e7 (2020).

44. Langdahl, J. H. *et al.* Mitochondrial Point Mutation m.3243A>G Associates With Lower

Bone Mineral Density, Thinner Cortices, and Reduced Bone Strength: A Case-Control

Study. *J. Bone Miner. Res. Off. J. Am. Soc. Bone Miner. Res.* **32**, 2041–2048 (2017).

45. Indo, Y. *et al.* Metabolic regulation of osteoclast differentiation and function. *J. Bone*

Miner. Res. Off. J. Am. Soc. Bone Miner. Res. **28**, 2392–2399 (2013).

46. Li, B. *et al.* Both aerobic glycolysis and mitochondrial respiration are required for

osteoclast differentiation. *FASEB J. Off. Publ. Fed. Am. Soc. Exp. Biol.* **34**, 11058–11067

(2020).

47. Zhou, P., Zheng, T. & Zhao, B. Cytokine-mediated immunomodulation of

osteoclastogenesis. *Bone* **164**, 116540 (2022).

48. Sun, C.-L., Van Gilst, M. & Crowder, C. M. Hypoxia-induced mitochondrial stress

granules. *Cell Death Dis.* **14**, 448 (2023).

49. Krämer, L. *et al.* MitoStores: chaperone-controlled protein granules store mitochondrial

precursors in the cytosol. *EMBO J.* **42**, e112309 (2023).

50. Formosa, L. E., Dibley, M. G., Stroud, D. A. & Ryan, M. T. Building a complex complex:

Assembly of mitochondrial respiratory chain complex I. *Semin. Cell Dev. Biol.* **76**, 154–

162 (2018).

51. Guerrero-Castillo, S. *et al.* The Assembly Pathway of Mitochondrial Respiratory Chain

Complex I. *Cell Metab.* **25**, 128–139 (2017).

52. Hoefs, S. J. G. *et al.* Novel mutations in the NDUFS1 gene cause low residual activities in

human complex I deficiencies. *Mol. Genet. Metab.* **100**, 251–256 (2010).

53. Buchan, J. R., Kolaitis, R.-M., Taylor, J. P. & Parker, R. Eukaryotic stress granules are

cleared by autophagy and Cdc48/VCP function. *Cell* **153**, 1461–1474 (2013).

54. Gwon, Y. *et al.* Ubiquitination of G3BP1 mediates stress granule disassembly in a

context-specific manner. *Science* **372**, eabf6548 (2021).

55. Hofer, S. J. *et al.* Spermidine is essential for fasting-mediated autophagy and longevity.

Nat. Cell Biol. **26**, 1571–1584 (2024).

56. Zhou, J. *et al.* Spermidine-mediated hypusination of translation factor EIF5A improves

mitochondrial fatty acid oxidation and prevents non-alcoholic steatohepatitis

progression. *Nat. Commun.* **13**, 5202 (2022).

57. Messerer, J. *et al.* Spermidine supplementation influences mitochondrial number and

morphology in the heart of aged mice. *J. Anat.* **242**, 91–101 (2023).

58. Rossi, D., Kuroshu, R., Zanelli, C. F. & Valentini, S. R. EIF5A and EF -P: two unique

translation factors are now traveling the same road. *WIREs RNA* **5**, 209–222 (2014).

59. Dong, Y. *et al.* A clinical-stage Nrf2 activator suppresses osteoclast differentiation via the

iron-ornithine axis. *Cell Metab.* **36**, 1679-1695.e6 (2024).

Materials and methods

Lead Contact and Materials Availability

Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contacts, Anna Katharina Simon (katja.simon@kennedy.ox.ac.uk) and Ana Victoria Lechuga-Vieco (ana.lechugavieco@irbbarcelona.org).

Materials availability

Animal lines generated in this study will be made available on request, but we may require a payment and/or a completed Materials Transfer Agreement if there is potential for commercial application.

Data and code availability

All data reported in this paper will be shared by the lead contact upon request. Original sequencing data files have been deposited at NCBI-SRA (PRJNA1144799), with a temporary link to access the metadata (https://dataview.ncbi.nlm.nih.gov/object/PRJNA1144799?reviewer=45rhoehhhq1rkj66o92nrv2pds). Links or accession numbers are also listed in the key resources table. The datasets will be publicly available upon the publication of this manuscript. This paper does report original code. The code for 3D bone shape reconstruction/evaluation and OC/OB quantification for in-vitro staining (Supplementary Table 1) can be found in GitHub repository: https://github.com/jianqingzheng/image_processing. Any additional information required to reanalyze the data reported in this work paper is available from the lead contact upon request.

Experimental model and subject details

Animals

All animal experiments were done in compliance with the guidelines established by the UK Animal (Scientific Procedures) Act 1986 and the Institutional Animal Care and Use Committee of the Barcelona Science Park and University of Barcelona. In vivo experiments with animals were undertaken under the UK Home Office Project License PPL 30/3388. Unless otherwise specified, all experiments were performed in 8- to 12-week-old mice in a C57BL/6JOlaHsd background and and a heteroplasmic mouse strain with the nuclear genome of C57BL/6JOlaHsd but mitochondrial DNA of C57+NZB (referred to as BL/6$^{C57-NZB}$)[1]. Mice were housed in specific pathogen-free facilities at the Kennedy Institue of Rheumatology (NDORMS, University of Oxford) and Park Scientific of Barcelona (PCB) under a 12-hour light/dark cycle and had ad libitum access to water and a standard chow diet (SDS diets 801,151; 75% energy from carbohydrates, 17.5% from protein, and 7.5% from fat).

Patient samples

Patient blood samples were obtained with the approval of the United Kingdom National Research Ethics Service (South Central-Berkshire, approval number 20/04, IRAS 162181). Written informed consent was obtained from all participants, and all procedures were carried out in accordance with relevant national and international guidelines and regulations. This research has been conducted using the UK Biobank Resource. This work was carried out under UK Biobank project number 103356.

Methods details

Micro computed tomography (µCT) scan and analysis of bone structure

The PerkinElmer QuantumFX scanner was utilized to scan the tibiae. The region of interest was specified as 100 slices below the growth plate of the proximal tibia and scanned for 3 minutes with an energy of 90KV, 200uA, and

field of view of 10. The resulting bone data from the scan were subjected to analysis through the use of Analyse-12.0 software and Analyse-14.0 software.

Compression testing of femur

After being harvested, the femurs were stored directly in -20°C. To prepare for the compression test, the femurs were rehydrated with PBS at 4°C overnight. Murine right femora were subjected to compression test using Electroforce 5500 (TA Instruments, Germany) on a 200 N load. Bones were secured on the lower plate and compressed. For testing, the bones were pre-loaded with a contact force of -0.5 N and load-displacement curves were generated in displacement control at a cross-head speed of 0.3 mm/s until failure of the diaphyseal region. Ultimate load was recorded for each group as an indicator of bone strength.

Serum and BM tissue processing

On the day of sacrifice, intracardiac puncture was used to collect mouse blood samples using capillary blood collection tubes (16.440, Microvette® CB 300 Z). The collected blood samples were kept on ice for up to 2 hours and then centrifuged at full speed for 5 minutes to extract the serum (supernatant). Serum was stored at −80°C until further analysis. To collect mouse BM cells, the femurs were crushed with a mortar and pestle in PBS-0.1% BSA-2 mM EDTA. The mixture was then filtered through 70μm cell strainers, and the collected cells were stored on ice, ready for flow cytometry without RBC lysis.

ELISA (Enzyme-Linked Immunosorbent Assay)

Mouse serum CTX1 level, mouse serum P1NP level and human serum CTX1 level were measured by ELISA according to the manufacturer's instructions.

Tartrate-Resistant Acid Phosphatase (TRAP) Staining

TRAP staining and counterstained with 0.2% methyl green solution experiments were previously described[2] and conducted according to the manufacturer's instructions. TRAP positive multi-nuclear (>3 nuclei) OC were counted. The Zeiss AxioScan Z1 Slide Scanner at the Wolfson Imaging Centre, WIMM, University of Oxford was used to acquire the images of OC.

TRAP staining and histomorphometry analysis of tibiae

Tibiae were fixed in 4% PFA at 4°C overnight and then decalcified in 10% Ethylenediaminetetraacetic acid (EDTA) decalcification solution for a duration of 14 days (buffer changed every 3 days). After decalcification, the tibiae were paraffin embedded for sectioning using a Leica RM2235 microtome, resulting in sections of 5μm thickness to create histology slides (VWR 631-1553). The sections were processed and stained with TRAP solution, followed by counterstaining with 0.2% methyl green solution. Histology slides were scanned using the Hamamatsu NanoZoomer S210 (WIMM, the University of Oxford). OC parameters from histology slides were manually quantified by a researcher from JingLianWen Technology (https://www.jinglianwen.com) blinded to the mouse experiment groups.

Bone Marrow Transplant and chimera blood lineages analysis

To establish BM chimeras, 8-12-week-old female BL/6[C57-NZB] mice and CD45.1[+] B6.SJL female mice were lethally irradiated at the dose of 11 Gy (550 cGy twice, 4 hours apart) and rested for 1 hour prior to injection. 1.5 million BM cells derived from 8-12-week-old BL/6[C57-NZB] and BL/6[C57] female mice or their respective littermates were transplanted (i.v. injection) into these irradiated mice. Reconstitution effectiveness was evaluated by analysing the percentage of CD45.1[+], CD45.2[+] and CD45.1/2[+] cells in the blood PBMCs of bone marrow-transplanted mice at 2,4, 6 and 8 weeks post-irradiation. To achieve this, we collected 50 microliters of blood from tail vein bleeding and treated it with RBC Lysis Buffer for 5 minutes. After centrifugation, the supernatant was removed, and the cells were stained with antibodies for flow cytometry analysis. Given that previous research[3] demonstrated that three weeks is sufficient for OC replenishment after transplantation, mice were sacrificed 8-9 weeks following lethal irradiation and transplantation.

This ensured comprehensive reconstitution of most cell lineages with the transplanted cells. Both blood and bone samples were collected for further analysis.

Mouse In vitro OC differentiation

Mouse in vitro OC differentiation method was previously described[2]. In brief, 8-week-old mouse BM cells were treated with 50 ng/ml M-CSF in Alpha-MEM culture media with 10% FBS, 1% pen/strep, and 1% L-glutamine for 48 hours (day -2) in 96-well plates. On day 0, the adherent cells were differentiated into OCs with 10 ng/ml mouse M-CSF and 50 ng/ml mouse RANKL. Spermidine was added at the dosage of 10uM. Culture media was refreshed every 3 days and OCs were identified by TRAP staining on day 5. The Olympus BX51 microscope at the Kennedy Institute of Rheumatology and AID ELISpot Reader System (ELR078IFL, AID GmbH) at the Wellcome Centre for Human Genetics were used to acquire the images of OC.

Flow Cytometry

Bone marrow and blood samples were collected from mice and prepared for flow cytometry. Erythrocytes were lysed using eBioscience[TM] 1X RBC Lysis Buffer. Cells were first incubated with FcR blocker in PBS for 20 minutes, followed by incubation with surface-conjugated antibodies for 30 minutes at 4°C. Conjugated antibodies used for surface staining were: anti-CD45.1, anti-CD45.2, anti- CD45R, anti-CD3e, anti-CD11b, anti-CD117 and anti-CD115. Identification of viable cells was done by fixable Zombie Aqua Live/Dead or near-IR dead cell staining. Stained samples were fixed with BD Cytofix[TM] Fixation Buffer for 15 minutes at RT. All samples were washed and stored in PBS containing 5 mM EDTA (Sigma-Aldrich) and 2% FBS (Sigma-Aldrich) before acquisition. To determine the absolute cell count within tissues, 10,000 or 20,000 CountBright[TM] absolute counting beads were added to each sample before flow cytometry to determine the total cell count per bone or blood volume. To ensure accuracy, over 2,000 beads were acquired per tube. Samples were acquired on a FACS LSR II (R/B/V) or Fortessa X20 (R/B/V/YG) flow cytometer (BD Biosciences) using FACSDiva software and analysed with FlowJo (version 10.10, BD Biosciences).

Mitochondrial content and ROS levels during OC differentiation: Mitochondrial content and ROS levels were assessed using Nonyl acridine orange (NAO) and CellROX™ Deep Red dyes. Cells were collected and incubated with 100 nM NAO and 5 µM CellROX™ for 20 minutes at 37°C. Ex/Em: 450/530 nm for NAO and 644/665 nm for CellROX™. After washing, cells were surface-stained and prepared for flow cytometry analysis as described in the previous section.

Autophagy flux (Flow cytometry). Autophagy levels were assessed as previously described[4]. In brief, cells were treated for 2 hours with either bafilomycin A1 (10 nM BafA1) or vehicle control, followed by staining using the FlowCellect[TM] Autophagy LC3 Antibody-based assay kit. After labeling with surface markers (as described earlier), cells were washed with Assay Buffer in 96-well U-bottom plates. Permeabilization was performed with 0.05% saponin, followed by immediate centrifugation. Cells were then incubated with anti-LC3 (FITC) at 1:20 dilution in Assay Buffer for 30 minutes at 4°C, followed by a final wash with 150 µL Assay Buffer. Autophagic flux was determined by calculating the LC3-II median fluorescence intensity as (BafA1 - Vehicle) / Vehicle.

Pre-processing of single-cell RNA-seq data

To collect mouse BM cells, the femurs were crushed with a mortar and pestle in PBS-0.1% BSA-2 mM EDTA. The mixture was then filtered through 70µm cell strainers, and the collected cells were plated for in vitro OC differentiation with 50 ng/ml M-CSF in Alpha-MEM culture media. On day 0, the adherent cells were differentiated into OCs with 10 ng/ml mouse M-CSF and 50 ng/ml mouse RANKL. Cells were harvested at the specified time points and were labeled with Cell Multiplexing Oligos (10x Genomics) according to the manufacturer's instructions. Barcoded sequencing libraries of multiplexed single-cell samples were generated using 10X Genomics Chromium system with 3' chemistry. Due to a technique issue, two of the three biological replicates were pooled together prior to the multiplexing step and were treated as single replicate (rep2) in the downstream analysis.

Computational analysis of single-cell RNA-seq data

Raw gene counts were normalized by sequencing depth and log1p transformed in Seurat. Batch effects were corrected using harmony[5] with n=10,000 variable features, while each sample was treated as an individual batch. Cells were then integrated using n = 30 harmony-corrected principal components, clustered at the resolution of 0.5 and then projected in 2D space using uniform manifold approximation and projection (UMAP). Any clusters that belonged to a non-myeloid lineage or had abnormally high levels of nuclear transcripts (possibly damaged cells with intact nuclei) were discarded, leaving n=10,809 cells for downstream analyses. Cluster marker genes and differentially expressed genes between mouse groups were identified by the Wilcoxon rank-sum test implemented in the Seurat package, and p-values were adjusted using the Benjamini-Hochberg procedure. Pathway enrichment was performed using XGR[6]. Genes associated with oxidative phosphorylation (OXPHOS) were obtained from the KEGG database and were converted to mouse symbols using Ensembl genome database. OXPHOS scores were computed using the "AddModuleScore" function in Seurat, and normalized against a set of n=100 random control genes.

The trajectory of osteoclast differentiation was recovered using information from RNA velocity. In brief, genes with a minimum of 20 spliced and 20 unspliced transcripts were extracted from the Cell Ranger alignment outputs and were analyzed in scVelo[7]. The splicing kinetics were recovered using the dynamical model and n=500 top velocity genes. For consistency, we used the same nearest-neighbor graph constructed from scRNA-seq data for all velocity analysis, and inferred velocities were projected onto the scRNA-seq UMAP. The initial and terminal clusters of the differentiation trajectory, together with a latent time for each cell, were inferred using CellRank[8]. We used the default combination of a velocity kernel (80%, deterministic mode) and a connectivity kernel (20%, computed based on the nearest-neighbor graph) to compute the probabilities for cell-cell transitions. The cell-level transitional probabilities were then aggregated at a cluster level and plotted as a fate map using directed PAGA[9].

For differential abundance test between BL/6[C57], and BL/6[C57-NZB] samples, batch-corrected principal components were used to construct single-cell neighborhoods using Milo[10], with prop = 0.1, k=20 and d=30. In brief, cells from BL/6[C57] or BL/6[C57-NZB] samples were counted within each neighborhood, and the statistical significance of differential abundance was evaluated using a generalized linear model with biological replicates included as covariates. The test results were visualized on a UMAP using the "plotNhoodGraphDA" function.

Mitochondrial genotyping using single-cell RNA-seq data

Genetic variants in mitochondrial transcripts were recovered at single-cell level using the mgatk pipeline[11]. In brief, sequencing reads were first aligned to a hard-masked mm10 reference genome using CellRanger (v7.1.0; 10x Genomics), followed by mgatk execution to compute the per variant coverage across the mitochondrial genome. The per-cell heteroplasmic ratio was calculated by dividing the summed allelic coverage of all variants specific to NZB mitochondria (n=90, Supplementary Table 1) over the total sequencing coverage at these locations. Only BL/6[C57-NZB] cells with more than 10 reads covering these variants were retained (n=3,093) for downstream analyses.

Seahorse analyses

Metabolic assays were performed using a Seahorse XFe96 Analyzer with Seahorse XF Cell Mito Stress Test Kit according to the manufacturer's instructions. Day 0, Day 1 and Day 3 pre-OCs were seeded in a 96-well plate at the density of 25,000, 25,000and 15,000 cells per well. Cells were treated with 10 ng/ml mouse M-CSF and 50 ng/ml mouse RANKL for indicated time prior to the assay. A final concentration of 1 uM oligomycin, 2 uM FCCP and 1 uM Rotenone and Antimycin A was used based on a pilot titration experiment. Cell density was quantified by CyQuant staining post Seahorse for data normalization.

ATP Synthesis

ATP levels were quantified using the ATP Bioluminescence Assay Kit CLS II following the manufacturer's protocol. Briefly, equal numbers of cells from each condition were pelleted, boiled in Tris-EDTA buffer (pH 7.74) for 2 minutes, centrifuged, and the supernatant was analyzed. Luminescence was measured in a CLARIOstar plate reader (BMG LABTECH).

Analysis of Complex I and Complex II Respiratory Activities

Enzymatic activities were measured spectrophotometrically using the Beer-Lambert law as previously described[12]. Briefly, assays were conducted at 32°C with frozen-thawed mitochondria in activity buffers. Complex I activity was initiated with 130 μM NADH and monitored at 340 nm, subtracting rotenone-insensitive rates. NADH:O_2 and NADH:decylubiquinone activities were assessed in sucrose buffer with specific inhibitors. Complex II activity was measured at 600 nm after adding 10 mM succinate.

Western Blot Analyses

Autophagy flux during OC differentiation was assessed by incubating the cells in culture media supplemented with 10 nM Bafilomycin A1 for 2 h, with DMSO served as the vehicle control. Autophagic flux was calculated as the ratio of LC3-I to LC3-II.

Cell lysis was performed on ice using NP-40 lysis buffer containing proteinase inhibitors and Phosphatase Inhibitor Cocktail 2. Afterward, the protein concentration was determined using the Pierce™ BCA Protein Assay Kits. The lysate was then treated with Reducing Laemmli Sample Buffer and 15–30 μg of proteins were loaded onto SDS-PAGE gels. For the separation of LC3-I and LC3-II, a 15% Tris-HCl gel and SDS running buffer were employed, while NuPAGE Novex 4–12% Bis-Tris gradient gels with NuPAGE™ MES SDS Running Buffer (1X) were used for other proteins. Proteins were subsequently transferred to a PVDF membrane and blocked with 5% skimmed milk-PBST. Primary antibodies were incubated overnight in 1% milk, followed by secondary antibody incubation in 1% milk with 0.01% SDS for 1 hour. The following primary antibodies were used: anti-LC3, anti-p62, anti-βActin and Total OXPHOS Rodent WB Antibody Cocktail. Imaging was performed using the Odyssey CLx Imaging System, capturing the signals from IRDye 800CW Donkey Anti-Rabbit IgG (H+L) (1:10,000 dilution) and IRDye 680RD Donkey Anti-Mouse IgG (H+L) (1:10,000 dilution) simultaneously, with varying intensities based on fluorescent exposure. See the Key resources table for a list of primary and secondary antibodies. Data analysis was carried out using Image Studio Lite.

Confocal imaging

Bone marrow cells were seeded with 10 ng/ml mouse M-CSF and 50 ng/ml mouse RANKL and differentiated for 1, 3, and 5 days in μ-Slide 18 Well Ibidi chamber slides (81816, Ibidi GmbH). Cells were fixed with 4% paraformaldehyde for 10 mins and permeabilized with 0.2% Triton X-100 in PBS for another 10 min in room temperature. Cells were blocked with PBS containing 2% BSA and 0.01% Tween20 for 1 h and incubated with the primary antibodies overnight at 4°C. The following antibodies were used to perform immunofluorescence staining in murine and human cells: mouse anti-NDUFA9 (20C11B11B11), anti-NDUFB7 (Polyclonal), anti-ATPB (3D5), anti-TOMM20 (EPR15581-54). After washing with PBS for three times, cells were incubated in 2% BSA-PBS for 1 hour with the following secondary antibodies: anti-mouse IgG AF488, anti-mouse AF647, anti-mouse AF546, anti-rabbit AF488 and anti-rabbit AF568. After washing with PBS and stained with 2 ug/ml DAPI for 10 mins, coverslips were mounted in SlowFade™ Diamond Antifade Mountant. Forty to eighty Z-stacks (0.13 μm) were acquired using a Zeiss LSM 980 Airyscan2 confocal system and the ZenBlue software. Data were analysed using ImageJ/Fiji[14].

Genotyping and Heteroplasmy Analysis

Heteroplasmy levels were assessed as previously described[15].H Briefly, mice were genotyped at postnatal day 21 using DNA extracted from ear notches with the REDExtract-N-Amp™ Tissue PCR Kit. Additionally, DNA from cells undergoing OC differentiation (Day 0, Day 1, Day 3, and Day 5) was isolated using 50 mM NaOH at 95°C, followed by neutralization with Tris-HCl (pH 7.5). The G4276A polymorphism, which introduces a BamHI restriction site in C57 mtDNA but not in NZB mtDNA, was used for genotyping. DNA was PCR-amplified with REDExtract-N-Amp™ PCR ReadyMix using primers: Forward 5′-AAGCTATCGGGCCCATACCCCG-3′ (3862-3884) and Reverse 5′-GTTGAGTAGAGTGAGGGATGGG-3′ (4503-4525), under the following conditions: 95°C (30s), 58°C (30s), 72°

C (45s) for 30 cycles. The PCR product was digested with FastDigest BamHI at 37° C for 20 minutes and analysed via agarose gel electrophoresis. DNA bands were visualised using the Gel Doc XR+ System (Bio-Rad) and quantified with Quantity One Software. The NZB mtDNA proportion was determined by the ratio of undigested 664-bp fragments to the total of 664-bp (undigested) and 414/250-bp (digested) fragments, with heteroduplex formation corrected using a standard curve from mixed C57 and NZB mtDNA samples.

Mitochondrial DNA copy number

Total DNA from cells undergoing osteoclast differentiation (Day 0, Day 1, Day 3, and Day 5) was isolated using 50 mM NaOH at 95°C, followed by neutralization with Tris-HCl (pH 7.5). qRT-PCR was performed on Applied Biosystems ®ViiA ™7 Real-Time PCR System using Power SYBR™ Green Master Mix in a 96-well plate to quantify mtDNA copy number relative to nDNA. The primers used were mtCo2 (mtDNA): COII RTF (3'-CTACAAGACGCCACAT-5', 7037-7052) and COII RTR (3'-GAGGGGGAGAGCAAT-5', 7253-7238), and SDH (nDNA): SDH RTF (3'-TACTACAGCCCCAAGTCT-5', 1026-1043) and SDH RTR (3'-TGGACCCATCTTCTATGC-5', 1219-1202). PCR conditions included an initial denaturation at 95ºC for 2 min, followed by 40 cycles of 95ºC for 15 s and 60ºC for 60 s, with a dissociation curve to confirm specificity. MtDNA copy number was calculated as $2 \times (2^{\Delta Ct})$, with $\Delta Ct = nCt - mtCt$, which is inversely proportional to mtDNA content[16].

Human osteoclast differentiation.

Human peripheral blood mononuclear cells (PBMCs) were isolated from EDTA-blood of healthy donors and MELAS patients. The PBMCs were centrifuged and resuspended in 10% DMSO in FBS and stored in -80° C. On the day of seeding (day -3), 2.5×10^5 cells/well of human PBMCs were seeded in 96-well plates with 100 µL/well of Alpha-MEM culture media containing 100 ng/ml human M-CSF, 10% FBS, 1% pen/strep, and 1% L-glutamine. This setup was maintained at 37°C in a 5.5% CO2 humidified incubator for 72 hours to facilitate the purification of adherent cells through plastic adhesion. On day 0, the cells were induced to differentiate into OCs in 100 µL/well of complete osteoclastogenic media (Alpha-MEM culture media with 10% FBS, 1% pen/strep, and 1% L-glutamine) supplemented with 50 ng/ml human M-CSF and 100 ng/ml human RANKL, and spermidine at 10uM. Subsequently, the culture medium was refreshed on day 2 and day 4. By day 6, mature OCs had formed and were fixed using 4% PFA for TRAP analysis.

Phenotypes and m.3243A>G in UK Biobank

The UK Biobank (UKB) is a population cohort of approximately 500,000 individuals from the UK[17]. As of 2024, it provides whole genome sequencing data alongside comprehensive phenotypic information, including electronic health records from hospitals and general practitioners, as well as self-reported data, death records, and cancer registry data. Heel bone mineral density (BMD) measurements using the Sahara heel ultrasound device are available for 278,835 participants, while DXA imaging-derived BMD data exist for 47,757 participants. Our study utilized the subset of UK Biobank participants with either ultrasound or DXA-derived BMD measurements. Our primary bone phenotype incorporated clinically relevant features, including self-reported fracture (data field 2463) as a surrogate for fragility fracture, consistent with previous publications[18]. We also analyzed direct BMD measurements from heel, spine (L1-L4), femur neck, and total body scans, using the minimum value between left and right measurements where available. The characteristics of the included cohort are detailed in Supplementary Table 2.

Mitochondrial m.3243A>G analysis. We identified the mitochondrial DNA (mtDNA) m.3243A>G variant using mitoHPC on UKB DRAGEN CRAM files[19]. The detailed process of variant and sample filtering is described in our recent publication. We successfully processed 490,412 samples to identify the m.3243A>G variant, a known pathogenic variant associated with diabetes and deafness[19]. For this study, we restricted our analysis to individuals with either heel or DXA imaging data. The characteristics of the included cohort are provided in Supplementary Table 2-4. We employed linear regression models to test the association between m.3243A>G and bone phenotypes, adjusting for age, age squared, sex, weight, principal components, assessment center, and sequencing center. A sensitivity analysis was performed, limiting the sample to individuals with >10%

heteroplasmy. We used Fisher's exact test to analyze the association between self-reported fracture and diabetes with m.3243A>G carrier status, using diabetes as a positive control for this analysis. We used Bonferroni corrected p value (0.05/4 = 0.0125) for linear regressions for significance threshold.

Statistical analysis

The mean±standard error of the mean (SEM) was used to represent all data. When comparing two experimental groups, a two-tailed t-test was employed. For datasets that were normally distributed with equal variances, an unpaired two-tailed Student's t-test was utilized. In experiments that had ≥3 experimental groups, one-way analysis of variance (ANOVA) and multiple comparisons with Tukey's correction were employed. Paired or unpaired one-way ANOVA was utilized for multiple comparisons of normally distributed datasets with one variable. The p-value was applied to evaluate the statistical significance of the hypothesis being tested. GraphPad Prism (San Diego, CA, Version 9.5.1 (528)) was used for statistical analyses. Wilcoxon rank-sum tests were used in single-cell statistical testing in R (v4.2.2).

3D bone reconstruction

To reconstruct and visualize the bone, Algorithm 1 is used (Supplementary Table 1). First, the CT volumes are rotated and translated to a standard position and orientation. Next, binary volumes of the entire skeleton structure and the cortical bones are extracted from the input volumes through thresholding. Subsequently, the connected components in the binary volumes are counted. This step aids in removing the fibular bone and retaining the largest connected cortical bone, as facilitated by Algorithm 2. Thereafter, the largest cortical bone is dilated and then eroded to fill the hole inside the tibia using Algorithm 3. In the final step, the cortical bones with filled holes intersect with the tibial skeleton structure, excluding the fibula, to extract the 3D shape of the tibial bone in the region of interest.

In Algorithm 2, a binary volume V^i and the index k serve as inputs (Supplementary Table 1). The initial step involves identifying and counting the connected components using MATLAB's built-in function "bwconncomp()". In the subsequent step, the connected components within the binary volume are labeled and tallied using MATLAB's built-in functions "labelmatrix()" and "regionprops()". These connected components are then arranged in descending order using MATLAB's "sort()" function. The value of index k is assessed to determine whether it's positive or negative. If k is positive, the k^{th} largest component is retained while others are discarded; if k is negative, the k^{th} largest component is eliminated. The algorithm concludes by outputting the remaining components, V^o. As a result, this methodology is adept at removing the fibular bone and any extraneous fragments that lie outside the tibial bone.

Algorithm 3 takes a binary volume V^i and a series of iteration numbers for dilation or erosion $[t_1, t_2, \cdots t_k]$ as inputs (Supplementary Table 1). The structuring element is initialized using MATLAB's built-in function "strel()". Subsequently, for each iteration number designated for dilation or erosion, the iteration number t is examined. If t is positive, the copied binary volume V^o undergoes t dilation iterations, and if t is negative, V^o is eroded for t times. This algorithm is thus integrated into Algorithm 1 with the aim of filling holes inside the cortical bones and subsequently compressing the volume to adhere to the original bone size.

Quantification of OC and OB from in vitro staining

To quantify the OC and OB parameters from microscopic images, Algorithm 4 is employed to determine the area size and the ratio of the cell region. In Algorithm 4, each image is processed using its corresponding mask for the region of interest and specific thresholding values (Supplementary Table 1). The algorithm starts by normalizing the intensity of the input images. Subsequently, the red channel of the image is extracted through thresholding to identify and isolate the cell areas. The next step involves identifying and counting the connected components with MATLAB's built-in function "bwconncomp()". Following that, the connected components within the binary image are labeled and enumerated using MATLAB's "labelmatrix()" and "regionprops()" functions. For each connected component, excluding the background region, the area size and its ratio in the entire region of interest

are computed based on the areas of the red channels. As a result, this algorithm can accentuate the cell regions in the provided images and compute the OC and OB parameters from the given microscopic imagery.

1. Lechuga-Vieco, A. V. *et al.* Heteroplasmy of Wild-Type Mitochondrial DNA Variants in Mice Causes Metabolic Heart Disease With Pulmonary Hypertension and Frailty. *Circulation* **145**, 1084–1101 (2022).

2. Leng, H. *et al.* Modulating glycosphingolipid metabolism and autophagy improves outcomes in pre-clinical models of myeloma bone disease. *Nat Commun* **13**, 7868 (2022).

3. Bakewell, S. J. *et al.* Platelet and osteoclast β_3 integrins are critical for bone metastasis. *Proc. Natl. Acad. Sci. U.S.A.* **100**, 14205–14210 (2003).

4. Alsaleh, G. *et al.* Autophagy in T cells from aged donors is maintained by spermidine and correlates with function and vaccine responses. *eLife* **9**, e57950 (2020).

5. Korsunsky, I. *et al.* Fast, sensitive and accurate integration of single-cell data with Harmony. *Nat Methods* **16**, 1289–1296 (2019).

6. Fang, H., Knezevic, B., Burnham, K. L. & Knight, J. C. XGR software for enhanced interpretation of genomic summary data, illustrated by application to immunological traits. *Genome Med* **8**, 129 (2016).

7. Bergen, V., Lange, M., Peidli, S., Wolf, F. A. & Theis, F. J. Generalizing RNA velocity to transient cell states through dynamical modeling. *Nat Biotechnol* **38**, 1408–1414 (2020).

8. Lange, M. *et al.* CellRank for directed single-cell fate mapping. *Nat Methods* **19**, 159–170 (2022).

9. Wolf, F. A. *et al.* PAGA: graph abstraction reconciles clustering with trajectory inference through a topology preserving map of single cells. *Genome Biol* **20**, 59 (2019).

10. Dann, E., Henderson, N. C., Teichmann, S. A., Morgan, M. D. & Marioni, J. C. Differential abundance testing on single-cell data using k-nearest neighbor graphs. *Nat Biotechnol* **40**, 245–253 (2022).

11. Lareau, C. A. *et al.* Massively parallel single-cell mitochondrial DNA genotyping and chromatin profiling. *Nat Biotechnol* **39**, 451–461 (2021).

12. Curtabbi, A. *et al.* Regulation of respiratory complex I assembly by FMN cofactor targeting. *Redox Biology* **69**, 103001 (2024).

13. Acín-Pérez, R., Hernansanz-Agustín, P. & Enríquez, J. A. Analyzing electron transport chain supercomplexes. in *Methods in Cell Biology* vol. 155 181–197 (Elsevier, 2020).

14. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**, 676–682 (2012).

15. Lechuga-Vieco, A. V. *et al.* Cell identity and nucleo-mitochondrial genetic context modulate OXPHOS performance and determine somatic heteroplasmy dynamics. *Sci. Adv.* **6**, eaba5345 (2020).

16. Moreno-Loshuertos, R. *et al.* Differences in reactive oxygen species production explain the phenotypes associated with common mouse mitochondrial DNA variants. *Nat Genet* **38**, 1261–1268 (2006).

17. Bycroft, C. *et al.* The UK Biobank resource with deep phenotyping and genomic data. *Nature* **562**, 203–209 (2018).

18. Kemp, J. P. *et al.* Identification of 153 new loci associated with heel bone mineral density and functional involvement of GPC6 in osteoporosis. *Nat Genet* **49**, 1468–1475 (2017).

19. Cannon, S. J. *et al.* Penetrance and expressivity of mitochondrial variants in a large clinically unselected population. *Human Molecular Genetics* **33**, 465–474 (2024).

Figure Legends

Figure 1. Heteroplasmic mice exhibit increased bone density and mechanical strength due to osteoclast dysfunction.

A Representative micro-CT reconstructed images of tibiae from 12-14-week-old male BL/6^{C57}, BL/6^{NZB} and BL/6$^{C57-NZB}$ mice.

B–C Micro-CT analysis of bone morphology parameters: BV/TV (bone volume fraction), Tb.Th (trabecular thickness), BS/BV (bone surface density), Tb.Sp (trabecular separation), Ct.Th (cortical thickness) and Ct.Ar (BL/6^{C57}, $n = 9$; BL/6^{NZB}, $n = 12$; and BL/6$^{C57-NZB}$, $n = 10$; each dot represents tibial measurements from an individual animal).

D Schematic overview illustrating the positioning of femur for compression testing between the load area and the platform load frame (left). Representative load displacement curve of femur from the BL/6^{C57} control group (middle). Dot plot showing the maximum load values for the femur in 12- to 14-week-old males of the indicated genotypes (right). BL/6^{C57}, $n = 8$; BL/6^{NZB}, $n = 8$; and BL/6$^{C57-NZB}$, $n = 7$. Each dot represents femoral measurements from an individual animal).

E Representative TRAP/0.2% methyl green-stained tibial sections showing osteoclasts (red/purple, black arrowheads) on the endocortical bone surface in BL/6^{C57}, BL/6^{NZB} and BL/6$^{C57-NZB}$ mice (n=5, two independent experiments).

F Bone histomorphometric quantification of osteoclasts (OCs), including the percentage of OC surface relative to the total bone surface (Oc.S/BS; left) and average length (right) (BL/6^{C57}, $n = 10$; BL/6^{NZB}, $n = 9$; and BL/6$^{C57-NZB}$, $n = 10$ mice, each dot represents an individual animal).

G ELISA analysis of serum levels of CTX-1 (left) and PINP (right) (n = 16 mice from the indicated genotypes, with each dot representing one animal.).

In A-G: *$p < 0.05$, **$p < 0.01$; ***$p < 0.001$, ****$p < 0.0001$ determined by ANOVA test. Data are presented as mean ± SEM.

H-I Lethally irradiated 8-week-old CD45.1+ B6.SJL and BL/6$^{C57-NZB}$ CD45.1+ mice were reconstituted with BM cells from female BL/6^{C57} CD45.2+ or BL/6$^{C57-NZB}$ CD45.2+ mice. After 9 weeks, micro-CT analysis was performed on tibiae. Left: Representative micro-CT reconstructed images. Right: BV/TV quantification (BL/6^{C57} to CD45.1+ B6.SJL, n = 10; BL/6$^{C57-NZB}$ to CD45.1^{+} B6.SJL, n = 12; BL/6^{C57} to BL/6$^{C57-NZB}$, n = 10; BL/6$^{C57-NZB}$ to BL/6$^{C57-NZB}$, n = 8; each dot represents an individual animal). *$p < 0.05$, two-way ANOVA.

Supplementary Figure 1: Gating strategy and quantification for OC progenitors and bone marrow chimera reconstitution.

A Flow cytometry gating strategy of the OC progenitors (CD11b- B220- CD3- CD115+) in BM cells (left) and percentage of OC progenitors (right) from BL/6^{C57} mice (n=10), BL/6^{NZB} mice (n=8) and BL/6$^{C57-NZB}$ (n=9) mice.

B Assessment of reconstitution efficacy in chimeric mice by evaluating the BM CD45.2+ population. Each dot represents an individual animal. Filled dots: donor cells; Empty dots: endogenous cells.

C Representative flow cytometry plots showing the gating strategy and distribution of CD45.1+/CD45.2+ BM cells from the specified groups.

D Morphological bone parameters in tibiae measured by micro-CT. Tb.Th, BS/BV, Tb.Sp, Ct.Th and Ct.Ar (BL/6^{C57} to CD45.1+ B6.SJL group (n=10); BL/6$^{C57-NZB}$ to CD45.1$^+$ B6.SJL group (n=12); BL/6^{C57} to BL/6$^{C57-NZB}$ group (n=10); BL/6$^{C57-NZB}$ to BL/6$^{C57-NZB}$ group (n=8).

In A-D: each dot represents and individual animal. $*p < 0.05$, $**p < 0.01$; $***p < 0.001$, $****p < 0.0001$ determined by two-way ANOVA test. Data are presented as mean ± SEM.

Figure 2. Impaired osteoclast differentiation, with mitochondrial dysfunction emerging as a hallmark in the first 3 days.

A Bone marrow cells from 8-week-old mice BL/6^{C57} , BL/6^{NZB} and BL/6$^{C57-NZB}$ mice were differentiated into osteoclasts (OCs) using 10 ng/ml M-CSF and 50 ng/ml RANKL. OCs were identified by TRAP staining on day 5 (left) Scale bar: 200 µm. The percentage of OC area relative to the total area was quantified (right). N= 20 mice per genotype (four independent experiments; each dot represents and individual animal). $****p < 0.0001$ determined by one-way ANOVA test. Data are presented as mean ± SEM.

B Schematic of experimental strategy for single-cell transcriptomics of murine osteoclast formation. Bone marrow cells from BL/6^{C57} (n=3), BL/6^{NZB} (n=3), and BL/6$^{C57-NZB}$ (n=3) strains were isolated and cultured with M-CSF (10 ng/ml) for 2 days, followed by M-CSF (10 ng/ml) and RANKL (50 ng/ml) from day 0 to day 3. Single-cell sequencing and analysis were then performed with day 0, day 1 and day 3 cells.

C Expression levels of osteoclast marker genes (*Acp5* and *Ctsk*) from aggregated single cells at each time point, indicating successful *in vitro* osteoclastogenesis.

D Uniform manifold approximation and projection (UMAP) visualization of single-cell transcriptomic profiles (n = 10,809 cells) during osteoclast differentiation. Cells are clustered by transcriptomic similarity and shown at three time points: day 0, day 1, and day 3.

E UMAP visualization of 10,809 cells during osteoclast differentiation, clustered by transcriptomic similarity and shown at day 0, day 1, and day 3 for different mtDNA groups (BL/6^{C57}, BL/6^{NZB}, and BL/6C^{57-NZB}).

F GO-BP pathway enrichment analysis for genes upregulated (red) or downregulated (blue) in BL/6$^{C57\text{-}NZB}$ osteoclasts compared to BL/6^{C57} osteoclasts.

Supplementary Figure 2. Single-cell transcriptomic profiling and trajectory analysis of *in vitro* osteoclast formation.

A UMAP visualization of the expression levels of osteoclasts marker genes (*Acp5*, *Mmp9, Ctsk, Dcstamp,* and *Ocstamp*) and negative regulators (*Irf8*, *Bcl6*, and *Mafb*). Color indicates log-normalized gene expression levels.

B UMAP projection of velocity-latent time estimated from the RNA splicing dynamics. Cells with a latent time of 0 or 1 represent the starting or terminating population infered by RNA velocity, respectively.

C A fate map for osteoclast differentiation using directed probabilistic approximate graph abstraction (PAGA). Each node represents one cluster in Figure 2D and arrows represent aggregated velocity flow concordant to the velocity-inferred trajectory. Dotted lines represent other velocity flow between each pair of clusters. Pie charts denote the relative abundance of BL/6^{C57}, BL/6^{NZB} and BL/6$^{C57\text{-}NZB}$ cells, normalized by the total number of cells in each group. The relative abundance of BL/6$^{C57\text{-}NZB}$ cells for each cluster along the osteoclast differentiation trajectory is labelled next to the pie chart.

D A graph representation of differential abundance tests. The relative enrichment BL/6$^{C57\text{-}NZB}$ cells in early-stage pre-osteoclasts (shown in red), as well as the relative depletion in mature osteoclasts (shown in blue), indicates premature termination of osteoclastogenesis in heteroplasmic cells. Nodes represent neighbourhoods of cells with similar transcriptomic profiles, colored by the relative abundance (log$_2$FC) of BL/6$^{C57\text{-}NZB}$ cells compared to BL/6^{C57}. Node sizes correspond to the number of cells in a neighbourhood. Non-significant neighbourhoods (p>0.05) are colored in white and plotted in the background.

Figure 3. Osteoclast maturation requires OXPHOS function which is disrupted by heteroplasmy, due to impaired localisation of complex I

A Heatmap showing gene expression changes for genes encoding subunits of respiratory ketogenesis, the Krebs cycle, pyrimidine biosynthesis and Complex I in osteoclasts derived from BL/6$^{C57\text{-}NZB}$ compared to BL/6^{C57} mice at day 0, day 1, and day 3 (n=3 mice per genotype).

B Aggregated expression of OXPHOS genes (OXPHOS scores) of *in vitro* OCs from different clusters (1-8) derived from BL/6^{C57}, BL/6^{NZB}, and BL/6$^{6C57\text{-}NZB}$ mice (n=3 animals per genotype). Module expression scores were normalized by subtracting the aggregated expression levels of a randomly selected, expression matched control gene set.

C Oxygen consumption rate (OCR) of intact cells estimated by SeaHorse during OC differentiation at day 0, day 1, and day 3 (3 to 5 independent assays per biological sample)

D Ratio of Complex I to Complex II activities. Respiratory complexes activities were measured spectrophotometrically using 5 µg of total mitochondrial protein per assay. Measurements were performed in four replicates using pre-OCs on days 0 and 3 of differentiation.

E Total ATP levels assessed by luminometry on day 1 and day 3 of osteoclast differentiation from BL/6^{C57}, BL/6^{NZB}, and BL/6$^{C57-NZB}$ mice. Each dot represents *in vitro* osteoclast differentiation from an individual mouse of the indicated genotypes. ** $p < 0.01$, determined using Tukey's multiple comparisons following two-way ANOVA.

F-G Confocal imaging of mitochondrial TOM20, Complex I subunit NDUFA9 and Complex V subunit ATPB in pre-osteoclasts on day 3 of differentiation derived from BL/6^{C57}, BL/6^{NZB}, and BL/6$^{C57-NZB}$ mice. F: Protein expression quantification ratio of NDUFA9 and ATPB per mitochondrion (NDUFA9/TOM20; ATPβ/TOM20), based on marker signals (Integrated Density, IntDen) on day 3 (n=4 mice per genotype). G: Representative confocal images for the indicated markers. Scale bar: 5 µm, n = 4 mice per genotype. * $p < 0.05$, two-tailed Dunnett's test following one-way ANOVA.

In C-F data are presented as mean values ± SEM.

Supplementary Figure 3. Transcriptomic analysis of metabolic pathways and mitochondrial characterization during osteoclast differentiation

A Heatmap showing gene expression changes for genes involved in glycolysis, glutamate degradation, adenosine biosynthesis, and the pentose phosphate pathway in osteoclasts cultures derived from BL/6$^{C57-NZB}$ and BL/6^{C57} mice at day 0, day 1, and day 3 (n=3 mice per genotype).

B Spare respiratory capacity (SRC) calculated by subtracting basal respiration from maximal respiration during *in vitro* osteoclast differentiation (day 0 to day 3) for the indicated genotypes. Data are presented as mean ± SEM (n=5, two independent experiments). * $p < 0.05$, two-tailed Dunnett's test following one-way ANOVA..

C Mitochondrial content was assessed using N-acridine orange (NAO) staining. Live cells were analyzed by FACS during osteoclast differentiation from day 0 to day 3.

D ROS quantification on day 0, day 1 and day 3 during OC differentiation. Left: Quantification of CellROX Deep Red MFI measurements (left) . Each dot represents an individual mouse. Right: FACS histogram plot (left) are representative measurements from untreated (grey), cells derived from BL/6^{C57} (blue), BL/6^{NZB} (red) BL/6$^{C57-NZB}$ (green) mice, where the grey line show background MFI (unstained).

In C-D, data were normalized to wild-type BL/6^{C57}; n= 3 BL/6^{C57} n=3 BL/6^{NZB} and n=5 BL/6$^{C57-NZB}$. * p<0.05, ** p <0.01 determined using Tukey's multiple comparisons following two-way ANOVA.

E Western blot analysis for OXPHOS complexes protein expression in day 0 , day 1 and day 3 of *in vitro* OC differentiation. CI-NDUFB8, CII-SDHB, CIII-UQCRC2, CIV-MTCO1 and CV-ATP5A levels are shown. TOM20 expression was used as mitochondrial protein loading control. Representative image from four independent experiments, n=4 mice per genotype.

In B-G, data are presented as mean values ± SEM.

Figure 4. Impaired import of nuclear-encoded complex I subunits into mitochondria leads to disrupted supercomplex assembly in heteroplasmic cells.

A-B: Confocal imaging of mitochondrial TOM20, Complex I subunit NDUFA9 and DAPI in pre-osteoclasts on day 3 of differentiation (A) and mature osteoclasts on day 5 of differentiation (B) derived from BL/6^{C57}, BL/6^{NZB}, and BL/6$^{C57-NZB}$ mice. Left: Representative confocal images for the indicated markers. Right: Quantification of colocalization of Complex I in the mitochondria (TOM20/NDUFA9) using as a control colocalization between mitochondria and the nucleous (TOM20/DAPI), shown by Pearson's coefficient. Scale bar: 5 µm, n = 4 mice per genotype. * p <0.05, *** p < 0.001, **** p < 0.00001, two-tailed Dunnett's test following one-way ANOVA. In A-B, data are presented as mean values ± SEM.

C-E Heatmap showing gene expression changes for genes involved in the mitochondrial protein import machinery (C), mitochondrial degradome (D) and mitochondrial assembly factors (E) in osteoclasts cultures derived from BL/6$^{C57-NZB}$ and BL/6^{C57} mice at day 0, day 1, and day 3 (n=3 mice per genotype).

In A-C, data are presented as mean values ± SEM.

Supplementary Figure 4. Mitochondrial-encoded genes are upregulated at day 3 of OC differentiation

A Violin plots displaying the average expression levels of mitochondrial DNA-encoded genes in BL/6^{C57} cells at day 0 (n = 1,550), day 1 (n = 1,148), and day 3 (n = 1,292) during OC differentiation. Complex I: mt-Nd1, mt-Nd2, mt-Nd3, mt-Nd4, mt-ND4L, mt-Nd5 and mt-Nd6. Complex III: mt-Cytb. Complex IV: mt-Co1, mt-Co2 and mt-Co3. Complex V (ATPase): mt-Atp6 and mt-Atp8. The y-axis represents the depth-normalized gene expression level in log scale (log count per 10,000 sequencing reads). **** p < 0.0001, Wilcoxon test, with day 0 and day 1 cells as the reference groups.

B Confocal imaging of mitochondrial TOM20 (red), Complex I subunit NDUFA9 (cyan) and DAPI (grey) in pre-osteoclasts on day 1 of differentiation derived from BL/6^{C57}, BL/6^{NZB}, and BL/6$^{C57-NZB}$ mice. Representative confocal images for the indicated markers. Scale bar: 5 µm, n = 4 mice per genotype. blurry

Figure 5. Mitochondrial selection during vitro OC maturation, and exhausted autophagic flux can be restored in heteroplasmic cells with spermidine

A NZB mtDNA levels (estimated as a percentage of total mtDNA) from BL/6$^{C57-NZB}$ mice, showing a shift toward mtDNA^{NZB} haplotype in day 3 cells compared to day 1. Each dot represents one cell and is colored by the percentile-normalized sequencing coverage of mtDNA^{NZB} variants. The colour scale represents sequencing depth. day 1: n = 1,249 cells; day 3: n = 884 cells. Median values are marked by black lines. **** $p < 0.0001$, Wilcoxon test.

B Assessment of the heteroplasmy shift toward NZB mtDNA at day 3 compared to day 1 in pre-OCs derived from BL/6$^{C57-NZB}$ young mice (n=9; each dot represents an individual mouse). The transformed heteroplasmy shift was evaluated using day 0 as the reference for each heteroplasmic mouse. NZB mtDNA levels were quantified by PCR, followed by RFLP (Restriction Fragments Length Polymorphism) to distinguish between C57 and NZB mtDNA variants. * $p < 0.05$, unpaired t test.

C NZB mtDNA levels in BL/6$^{C57-NZB}$ cells across different clusters indicating increased mitochondrial segregation in osteoclasts compared to monocytic precursors. Median values are marked by grey lines. **** $p < 0.0001$, Wilcoxon test, with monocytic precursors as the reference group.

D-E Western Blot analysis of autophagy in primary murine BM cells treated with M-CSF and RANKL for 1 or 3 days. (D) Representative images of membranes showing p62 and LC-3 expression and (E) autophagy flux quantification of the LC-3 bands (LC3-I/LC3-II) at day 1 and 3 of OC differentiation. 10 nM BafA1 was added 2 hours before protein harvest, with actin used as a loading control. n = 8 independent biological samples for DMSO control and BafA. **$p < 0.01$, ****$p < 0.0001$, two-way ANOVA with Sidak's multiple comparisons test.

Quantification of autophagy flux based on LC3-I/LC3-II band intensity from western blot analysis after day 1 of OC differentiation. BafA1 (10 nM) was added 2 hours before protein harvest. n = 8 independent biological samples for both DMSO control and BafA1 treatment.).

F-G Bone marrow cells from 8-week-old mice BL/6^{C57}, BL/6^{NZB} and BL/6$^{C57-NZB}$ mice were differentiated into osteoclasts (OCs) using 10 ng/ml M-CSF and 50 ng/ml RANKL with or without 10 uM spermidine (spd) treatment. F: The percentage of OC area relative to the total area was quantified. Data are representative of two independent

experiments. * p < 0.05, ****p < 0.0001, nonparametric Kruskal-Wallis test. G: OCs were identified by TRAP staining on day 5. Scale bar: 50 μm.

H Assessment of the heteroplasmy shift toward NZB mtDNA at day 5 in OCs derived from BL/6^{C57}-NZB young mice (n=10 for untreated and n=8 for cultures treated with 10 μM spermidine; each dot represents an individual mouse). The transformed heteroplasmy shift was evaluated using day 1 as the reference for each heteroplasmic mouse. NZB mtDNA levels were quantified by PCR, followed by RFLP (Restriction Fragments Length Polymorphism) to distinguish between C57 and NZB mtDNA variants. ** p < 0.01, nonparametric Mann-Whitney test.

I Total ATP levels were measured by luminometry on day 3 of OC differentiation in BL/6^{C57}, BL/6^{NZB}, and BL/6$^{C57-NZB}$ mice, with and without 10 μM spermidine treatment. Data are normalized to untreated cells and each dot represents a sample from an individual mouse of the indicated genotype. ****p < 0.0001, one-way ANOVA and Sidak's multiple comparison test.

J Confocal imaging of mitochondrial TOM20, Complex I subunit NDUFA9, and ATPase subunit ATPB in pre-osteoclasts derived from BL/6^{C57}and BL/6^{NZB}, and BL/6$^{C57-NZB}$ mice with and without spermidine treatment on day 5 of *in vitro* OC differentiation. Left: Protein quantification ratio of NDUFA9 per mitochondrion (NDUFA9/TOM20), based on marker signals (Integrated Density, IntDen)(n=3 mice per genotype). **** p <0.0001, two-tailed Dunnett's test following one-way ANOVA. Right: Representative confocal images for the indicated markers in pre-osteoclasts derived from BL/6$^{C57-NZB}$ mice with and without spermidine treatment on day 5 of *in vitro* OC differentiation. Scale bar: 5 μm.

K-L Lethally irradiated young BL/6$^{C57-NZB}$ mice were reconstituted with bone marrow from BL/6^{C57} or BL/6$^{C57-NZB}$ mice and treated with 5 mM spermidine in drinking water for 8 weeks. (K) Representative micro-CT reconstructed images of tibiae from the bone marrow reconstituted mice. (L) BV/TV, Tb.Th, BS/BV Tb.Sp, Ct.Th and Ct.Ar of tibiae quantification by micro-CT (n = 8 BL/6^{C57}; n = 6 BL/6$^{C57-NZB}$ and n = 7 BL/6$^{C57-NZB}$+Spd; each dot represents an individual animal). The experiment is representative of two independent experiments. *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001, one-way ANOVA followed by Tukey's multiple comparisons test.

M Bending test analysis in bone marrow transplanted mice treated with 5 mM spermidine. Dot plots of maximum femur load from bone marrow-transplanted mice, assessed using the bending test (n = 5 mice per BM cell genotype; each dot represents an independent animal). *p < 0.05, **p < 0.01, one-way ANOVA followed by Tukey's multiple comparisons test.

In B-M, data are presented as mean values ± SEM.

Supplementary Figure 5. Analysis of Mitochondrial Selection and the Role of Autophagy and Spermidine Supplementation in In Vitro and In Vivo Osteoclastogenesis

A Average sequencing depth (bars) and the average heteroplasmic percentage (dots) for selected NZB mtDNA variants in heteroplasmic cells. Colors represent the number of cells with at least 10x coverage for a given variant.

B NZB mtDNA levels (%NZB mtDNA, y axis) plotted against the velocity latent time (x axis, same latent time as shown in Supplementary Figure 2B), demonstrating a slight shift towards NZB mtDNA during osteoclast differentiation. *** $p < 0.001$, Spearman's rank correlation test.

C Assessment of the percentage of NZB mtDNA (% NZB mtDNA) during *in vitro* OC differentiation of heteroplasmic cells (0, 48, and 192 hours after the addition of 10 ng/ml M-CSF and 50 ng/ml RANKL). NZB mtDNA levels were quantified by PCR, followed by RFLP (Restriction Fragment Length Polymorphism) to distinguish between C57 and NZB mtDNA variants. Samples were derived from n = 4 BL/6$^{C57-NZB}$ mice.

D Quantification of p62 expression from Western blot analyses shown in Figure 5D. 10 nM BafA1 was added 2 hours before protein harvest, with actin used as a loading control. n = 8 independent biological samples for DMSO control and BafA1; each dot represents an individual mouse. *$p < 0.05$, **$p < 0.01$, determined using Tukey's multiple comparisons following two-way ANOVA.

E Violin plots displaying the average expression levels of Tfeb in monocytic precursors (Mono) and mature osteoclasts (OC) in BL/6^{C57} (black), BL/6^{NZB} (grey), and BL/6$^{C57-NZB}$ mice (red).*$p < 0.05$, ***$p < 0.001$, two-sided Wilcoxon test.

F Mitochondrial DNA copy numbers in BL/6$^{C57-NZB}$ cell cultures with or without 5 µM spermidine treatment on day 1 and day 3 during osteoclast differentiation (n=4-5 mice). mtDNA copy numbers were normalized to genomic DNA copy numbers, as determined by qPCR ($2-\Delta CT$).

G Total ATP levels were measured by luminometry on day 1 of OC differentiation in BL/6^{C57}, BL/6^{NZB}, and BL/6$^{C57-NZB}$ mice, with and without 10 µM spermidine treatment. Data are normalized to untreated cells and each dot represents a sample from an individual mouse of the indicated genotype. ****$p < 0.0001$, one-way ANOVA and Sidak's multiple comparison test.

H-I Confocal imaging of mitochondrial TOM20, Complex I subunit NDUFA9, and ATPase subunit ATPB in pre-osteoclasts derived from BL/6^{C57}and BL/6^{NZB}, and BL/6$^{C57-NZB}$ mice with and without spermidine treatment on day 5 of *in vitro* OC differentiation. I: Protein quantification ratio of ATPB per mitochondrion (ATPB/TOM20), based on marker signals (Integrated Density, IntDen)(n=3 mice per genotype). **** $p < 0.0001$, *** $p < 0.001$, ** $p < 0.01$, two-tailed Dunnett's test following one-way ANOVA. In A-B, data are presented as mean values ± SEM. J:

Representative confocal images for the indicated markers in pre-osteoclasts derived from BL/6^{C57}and BL/6^{NZB} mice with and without spermidine treatment on day 5 of *in vitro* OC differentiation. Scale bar: 5 µm.

Figure 6. Osteoclast differentiaion in cells from patients with mtDNA heterogeneity (m.3243A>G) is impaired and can be restored with spermidine

A Human PBMCs were seeded and differentiated into osteoclasts (OCs) using 50 ng/ml human M-CSF and 75 ng/ml human RANKL. Left: Representative images of TRAP staining from OC on day 9. Right: Quantification of the percentage of OC area relative to the total area. n=3 healthy donors (HD) and n=5 m.3243A>G patients; 3 technical replicates per condition. * p < 0.05, two-tailed unpaired t-test.

B Quantification of bone resorption marker CTX-1 levels in human serum samples from healthy donors (HD, n=10) and m.3243A>G patients (n=5). *p < 0.05, two-tailed unpaired t-test.

C Human PBMCs were seeded and differentiated into osteoclasts (OCs) in the presence of 50 ng/ml human M-CSF and 75 ng/ml human RANKL, with or without 10 µM spermidine treatment. Left: Representative images of TRAP staining from OC on day 6. Right: Quantification of the percentage of OC area relative to the total area. n=4 healthy donors (HD) and n=4 m.3243A>G patients; 6 technical replicates per condition. ** p < 0.01, determined using Tukey's multiple comparisons following one-way ANOVA. Scale bar: 50 µm.

D Total ATP levels assessed by luminometry on day 5 of osteoclast differentiation from human PBMCs in the presence of 50 ng/ml human M-CSF and 75 ng/ml human RANKL, with or without 10 µM spermidine treatment. Each dot represents *in vitro* osteoclast differentiation from a donor. n=5 healthy donors (HD) and n=5 m.3243A>G patients. * p < 0.05, determined using Tukey's multiple comparisons following one-way ANOVA.

In A-D, data are presented as mean values ± SEM.

Figure 1

Figure 2

Supplementary Fig. 2

Figure 3

Supplementary Fig. 3

Figure 4

A

mt-Nd1 **mt-Nd2** **mt-Nd3** **mt-Nd4** **mt-Nd4l** **mt-Nd5** **mt-Nd6**

mt-Cytb **mt-Co1** **mt-Co2** **mt-Co3** **mt-Atp6** **mt-Atp8**

B

Day 1

Supplementary Fig. 4

Figure 5

Supplementary Fig. 5

A

HD   m.3243A>G

OC area / total area %

HD m.3243A>G

B

CTX (pg/ml)

HD m.3243A>G

C

HD   m.3243A>G Control   m.3243A>G +Spd

OC area / total area %

HD   Control   + Spd

m.3243A>G

D

µM ATP / total protein

HD   Control   + Spd

m.3243A>G

Figure 6

Supplementary Table 1:

Algorithm 1: Bone Reconstruction

Input: $X \in R^{d \times h \times w}$, x_{bone}, $x_{cortical}$, $n_{frag} \in R$

Output: $V \in \{0,1\}^{d \times h \times w}$

---

1. Rotate and translate input volume to a standard position and orientation;

2. Extract the whole bone and cortical bones by thresholding: $V^0 \leftarrow nonzero(X - x_{bone})$, $V^1 \leftarrow nonzero(X - x_{cortical})$;   /* 'nonzero' is a MATLAB function */

3. Remove the small fragments from the cortical bones' volume: $V^1 \leftarrow bwareaopen(V^1, n_{frag})$;

4. Remove the fibular bone (the second largest volume): $V^0 \leftarrow KeepRankedVol(V^0, -2)$, $V^1 \leftarrow KeepRankedVolume(V^1, 1)$;

5. Dilate and erode the cortical bone to fill the hole inside: $V^1 \leftarrow SerialDilateErode(V^1, [12, -12])$;

6. Intersect the two volumes: $V \leftarrow V^0 \cap V^1$;

7. Return V.

Algorithm 2: Keep Ranked Volume

Input: $V^i \in \{0,1\}^{d \times h \times w}$, $k \in Z$

Output: $V^o \in \{0,1\}^{d \times h \times w}$

---

1. Find and count the connected components: $V^{cc} \leftarrow bwconncomp(V^i, 26)$;   /* 'bwconncomp' is a MATLAB function */

2. Creates a label matrix corresponding to each connected component: $L^{cc} \leftarrow labelmatrix(V^{cc})$; /* 'labelmatrix' is a MATLAB function */

3. Calculate parameters of each component: $AreaProp \leftarrow regionprops(V^{cc}, 'Area')$;   /* 'regionprops' is a MATLAB function */

   Sort the areas: $(\cdot, index_{sort}) \leftarrow sort(AreaProp, 'descend')$;

4. Copy the volume: $V^o \leftarrow V^i$;

5. If $k > 0$ do:

6.   Remove the components other than the k^{th} largest volume: $V^o[L^{cc}! = k] \leftarrow 0$;

7. Else do:

8.   Remove the components of the k^{th} largest volume: $V^o[L^{cc} == k] \leftarrow 0$;

9. End

10. Return V^o.

11.

Algorithm 3: Serial Dilate & Erode

Input: $V^i \in \{0,1\}^{d \times h \times w}$, $[t_1 \cdots t_k] \in Z^k$

Output: $V^o \in \{0,1\}^{d \times h \times w}$

Set the parameter size of structure element: $size$;

Initialize a structure element: $se \leftarrow strel('cube', size)$;   /* 'strel' is a MATLAB */

Initialize the volume: $V^o \leftarrow V^i$;

Foreach $i \in [1, k] \cap Z$ do:

Foreach $j \in [1, abs(t_i)] \cap Z$ do:

If $t_i > 0$ do:

7   Dilate the volume to fill the hole inside: $V^o \leftarrow imdilate(V^o, se)$;

8   Elseif $t_i < 0$ do:

9   Erode the volume to fit a specific size: $V^o \leftarrow imerode(V^o, se)$;

10   End

11   End

12 Return V^o.

Algorithm 4: Count Area & Ratio

Input: $I^i \in R^{h \times w}$, $\tau \in R$, $n_{conn} \in Z$, $M^i \in \{0,1\}^{h \times w}$

Output: $M^r \in \{0,1\}^{h \times w}$, $a^r \in R^m$, $r^r \in R^m$

---

1 Normalize the input image: $I^o \leftarrow (I^i - I^i)/(I^i - I^i + 10^{-5})$;

2 Filter the image using a high passing filter: $I^o \leftarrow (1 - G) * I^i$; /* G is a gaussian filter kernel
   function */

   Get the areas of the red channel: $M^r \leftarrow (I^o[\cdots, 1] > \tau[1]) \& (I^o[\cdots, 2] < \tau[2]) \& (I^o[\cdots, 3] < \tau[3])$;

   Find and count the connected components: $M^{cc} \leftarrow bwconncomp(M^i, 8)$;   /* 'bwconncomp'
   is a MATLAB function */

   Creates a label matrix corresponding to each connected component: $L^{cc} \leftarrow labelmatrix(M^{cc})$;
   /* 'labelmatrix' is a MATLAB function */

   Calculate parameters of each component: $AreaProp \leftarrow regionprops(M^{cc}, 'Area')$;   /*
   'regionprops' is a MATLAB function */

   Get the number of components: $m \leftarrow |AreaProp| - 1$;

   Initialize the areas and ratios for each component: $a^r \leftarrow [0]_{m \times 1}$, $r^r \leftarrow [0]_{m \times 1}$

   Foreach $i \in [1, m] \cap Z$ do:

10	$a^r[i] \leftarrow \sum(L^{cc} == i + 1) \cap M^r$;
11	$r^r[i] \leftarrow a^r[i]/\sum(L^{cc} == i + 1)$;
12	End
13	Return M^r, a^r, r^r.

Supplementary Table 2: Characteristics of individuals included in the study by m.3243A>G status.

Characteristics	m.3243A>G carriers N=76	Non-carriers, N=271,773
Age at recruitment, years	54 (13.3)	58 (13)
Female	30 (39)	147778 (54.4)
Individuals with heel BMD	76	271773
Individuals with DXA	15	46762
Blood heteroplasmy of m.3243A>G	0.088 (0.085)	-
Individuals with >5% heteroplasmy	76 (100)	-
Individuals 5-10% heteroplasmy	46 (60)	-
Individuals >10% heteroplasmy	30 (40)	-
Weight, kg	69.1 (21.6)	76.4 (21.0)

Median (IQR) for continuous data and n (%) for categorical data is displayed.

Supplementary Table 3: Association analysis (continuous phenotypes) of m.3243A>G with bone mineral density

Phenotype	Total n	n with >5% heteroplasmy	Effect size in g/cm² >5% heteroplasmy	p-value >5% heteroplasmy	n with >10% heteroplasmy	Effect size in g/cm² >10% heteroplasmy	p-value >10% heteroplasmy
Heel BMD	270568	76	-1.60×10^{-5}	0.99	30	-2.53×10^{-2}	0.29
Femur neck BMD	43129	13	-4.34×10^{-2}	0.19	6	-4.80×10^{-2}	0.29
L1-L4 BMD	42472	13	-4.56×10^{-2}	0.35	6	-5.08×10^{-2}	0.48
Total BMD	44494	14	-3.72×10^{-2}	0.20	7	-3.36×10^{-2}	0.41

Figure 2a: Results from linear regressions with BMD measures. Bonferroni corrected p-value < 0.0125.

Supplementary Table 4: Association analysis (binary phenotypes) of m.3243A>G with self-reported fracture and diabetes

Phenotype	n total	n >5% heteroplasmy	OR >5% heteroplasmy (95% CI)	p-value >5% heteroplasmy	n >10% heteroplasmy	OR >10% heteroplasmy(95% CI)	p-value >10% heteroplasmy
self-reported fracture	270247	76	0.528 (0.140, 1.41)	0.245	30	0.679 (0.0784, 2.69)	1.00
diabetes (Eur only)	253554	69	8.07 (4.86, 13.3)	8.22×10^{-15}	29	16.2 (7.24, 37.9)	2.74×10^{-12}

Figure 2b: Results from Fisher's exact test with self-reported fracture and diabetes.