Warm and cool temperatures decrease early-life telomere length in wild pied flycatchers

Furic C[1,2§], Marciau C[1,3], Hsu B-Y[1,4], Cossin-Sevrin N[1,5], Fleitz J[1], Reichert S[1,2], Ruuskanen S[1,6]* & Stier A[1,2§]*

[1]Department of Biology, University of Turku, Finland

[2]Université de Strasbourg, CNRS, IPHC UMR 7178, F-67000 Strasbourg, France

[3]Institute for Marine and Antarctic Studies, University of Tasmania, Hobart, Australia

[4]Department of Life Science, Tunghai University, Taichung, Taiwan

[5]Institute of Biotechnology, HiLIFE, University of Helsinki, Helsinki, Finland

[6]Department of Biological and Environmental Science, University of Jyväskylä, Finland

§ corresponding author: cmfuri@utu.fi; antoine.stier@iphc.cnrs.fr

* co-last authors

Abstract

Climate change represents a major challenge for avian species. It is characterized by an increase in average ambient temperatures, but also in occurrence of extreme weather events, such as heat weaves and cold snaps. These abrupt temperature changes can modify the immediate and long-term survival prospects of nestling birds, when their thermoregulatory capacities are still not fully developed. While immediate nestling survival can easily be measured, long-term survival is more challenging to evaluate. Early-life telomere length has been suggested as a potential biomarker of future fitness prospects. To evaluate the potential impact of changes in early-life temperature, we thus experimentally increased (*ca.* +2.8°C) and decreased (*ca.* -1.7°C) average nestbox temperatures in wild pied flycatchers (*Ficedula hypoleuca*) during nestling postnatal growth, and measured nestling telomere length before fledging. Shorter telomeres were observed in individuals exposed both to an experimental heating or cooling during growth. Our results suggest that long-term survival prospects or long-term performances of individuals exposed to abrupt changes in early-life temperature may be decreased.

Keywords: endotherm, passerine, telomere, thermoregulation, cellular ageing

Introduction

Climate change is characterized by an increase in average ambient temperatures, but also in occurrence of extreme weather events such as cold snaps and heat waves (IPCC 2014). These abrupt temperature changes can cause direct mortality (McKechnie & Wolf 2009, McKechnie *et al.* 2021) but also sub-lethal consequences that may impact fitness because of delayed or long-lasting effects (Conradie *et al.* 2019).These extreme weather events can represent a major challenge for avian species, especially during early development when altricial nestlings are transitioning from an ectothermic stage to a fully endothermic stage prior to fledging (Price & Dzialowski 2018). As individuals do not fully develop their capacity to regulate body temperature during these early-life stages (Boyles *et al.* 2011), their homeothermy depends on ambient temperature fluctuations and parental brooding throughout this thermoregulatory transition (Katzenberger *et al.* 2015). This temperature-sensitivity exposes individuals to short-term mortality risks caused by temperature variation (Hepp & Kennamer 2012, Moreno *et al.* 2015, Andreasson *et al.* 2018, Ospina *et al.* 2018) but also potential long-lasting or delayed effects on their health or reproduction (Metcalfe & Monaghan 2001, Preston *et al.* 2018) that may affect fitness (Greño *et al.* 2008, Hepp & Kennamer 2012, Berntsen & Bech 2016, Nord & Nilsson 2016, Andreasson *et al.* 2018). Previous studies have demonstrated the effect of heatwaves (Greño *et al.* 2008, Nord & Nilsson 2016, Andreasson *et al.* 2018) or cold snaps (Moreno *et al.* 2015, Ospina *et al.* 2018) events during pre- or postnatal stages on growth and parental brooding behavior (Rodríguez & Barba 2016). However, the potential long-term survival effects of such abrupt changes in temperature can be more difficult to characterize, because long-term recaptures rates are often very low for passerine birds (Andreasson *et al.* 2018, Stier *et al.* 2025).

To elucidate these potential long-term or delayed effects of early-life environment, telomere length may be used as a molecular biomarker for future fitness prospects (Eastwood et al. 2019). Telomeres are the protective structures located at the end of chromosomes, playing a role to maintain chromosome stability (Lange *et al.* 2006). It has been shown that telomeres progressively shorten with age (Remot *et al.* 2022), and that telomere length predicts survival prospects (Wilbourn *et al.* 2018). Telomeres also shorten faster in response to various environmental stressors, especially during early-life (Chatelain *et al.* 2019). For instance, telomere length may be affected by early-life temperature conditions, with experimental elevation of egg incubation temperatures leading to shorter telomeres (Vedder *et al.* 2018, Stier *et al.* 2020a, Hope *et al.* 2023). Yet, the effect of heatwaves on telomere length during postnatal stages have been less characterized. There is some evidence of adverse effects of natural or experimental hot conditions on telomere length in the wild (from postnatal d7 to d14:

Stier *et al.* 2025; from postnatal d0 to d7: Eastwood *et al.* 2022), but no effect of experimental heatwaves on telomere length in captivity (from hatching to postnatal d12: Ton *et al*. 2023). To the best of our knowledge, the effect of postnatal coldsnap on telomere length has never been investigated. Telomere shortening can be accelerated by fast growth and rapid cellular proliferation (Monaghan & Ozanne 2018), high oxidative stress levels (Reichert & Stier 2017), or energetic imbalance (Casagrande & Hau 2019). All these mechanisms may underlie potential effects of early-life temperature on telomere length.

In this study, we investigated the effects of experimental increase and decrease in nest temperature during early postnatal development on telomere length in wild pied flycatchers (*Ficedula hypoleuca*). As study years also differed in terms of ambient thermal conditions (see results), we also used this opportunity to investigate the potential impact of longer-term natural variation in ambient temperature on telomere length. Age- and stress-related telomere shortening have been previously demonstrated in this species and its close relative, the collared flycatcher (*Ficedula albicollis)* (Kärkkäinen *et al.* 2019, 2020, 2021, 2022, Stier *et al.* 2020b). Investigating experimentally the effects of heatwave and coldsnap on postnatal telomere length in a free-living species should help to determine the potential long-term or delayed effects of extreme weather events on fitness. We predict that being exposed to early-life hot environmental conditions should lead to shorter telomeres, as found in Stier *et al*. (2025) and Eastwood *et al.* (2022). However, experiencing a coldsnap may lead either to shorter telomeres, if the coldsnap triggers an increase in metabolic rate or energetic imbalance (Casagrande & Hau 2019, Casagrande *et al.* 2020), or to longer telomeres, if the coldsnap reduces metabolic rate of nestlings still being partly ectothermic (Friesen *et al.* 2022).

Material and methods

General field procedures:

Two nest temperature manipulations were conducted in a nestbox population of pied flycatchers located in the island of Ruissalo (Turku, Finland, location: 60°25'N, 22°10'E). In 2018, a heating experiment was performed, (data partly published in Hsu *et al.* (2020)), and in 2019 a cooling experiment was performed. Nestbox temperature was recorded every 3 minutes with an Ibutton thermochron (0.0625°C accuracy) installed *ca.* 5 cm above the rim of the nest. On the second day after hatching (d2), each individual was identified by clipping nails and half of the nestlings (minimum 2 individuals) with the same age and weight were swapped with another nest for other experimental purposes (*i.e.* cross-fostering to disentangle environmental and genetic effects).

Heating experiment (2018):

In 2018, nestbox temperature was increased for half of the boxes (hereafter, 'heated nests') between postnatal d2 and d8. Heating pads (UniHeat 72h, USA) were installed under the ceiling and replaced every 2nd day in the morning. The other half was considered as controls (hereafter, 'non-heated nests'), non-functional heating pads were used and nests were visited every 2nd day as well to standardize disturbance. At d8 after hatching, individuals were identified and ringed, and at postnatal d13 one blood sample of *ca.* 40μl was collected from the brachial vein using heparinized capillaries. A total of 97 nestlings from 41 nests were analyzed, specifically 42 individuals from 21 heated nests and 55 individuals from 20 non-heated ones. When possible, the nestlings selected for telomere analysis included (1) both a female and male nestling from each nest (2) one of genetic origin and one foster nestling, and otherwise were randomly selected. Since females can actively brood nestlings until postnatal d6 or d7 (Lundberg & Alatalo 1992), they could potentially change their behavior depending on the heating treatment, but no evidence was found in this experiment (previously published in Hsu *et al.* (2020)).

Cooling experiment (2019):

In 2019, nestbox temperature was decreased for half of the boxes (hereafter, 'cooled nests') between d7 and d13 after hatching of the nestlings. The timing of the experiment was chosen to avoid any potential confounding effects of brooding behavior by the female, and knowing that pied flycatchers are not yet fully capable of thermoregulation at d7 (Lundberg & Alatalo 1992). Two frozen (-20°C) cooling pads (VWR 26.7*14*3.2*cm, 909g) were attached on each outer side of the nestbox (protected on the outside with insulating material) and replaced every morning. The other half

(hereafter, 'non-cooled nests') was equipped with the same pads but unfrozen and visited every day too. Before the treatment, at d7 after hatching, individuals were identified and ringed. Morphological measurements (body mass and wing length) were conducted at postnatal d13, and one blood sample of *ca.* 40µl was collected from the brachial vein using heparinized capillaries. A total of 56 nestlings from 48 nests were analyzed, including 25 individuals from 23 cooled nests and 31 individuals from 25 non-cooled nests. The nestlings were selected for telomere analysis by trying to balance the number of female and male nestlings, and having one of genetic origin and one foster nestling.

Relative telomere length analysis:

We measured relative telomere length using a real-time quantitative PCR (qPCR) assay previously used and validated in pied flycatcher using the TRF method (Kärkkäinen *et al.* 2019, 2021). DNA was extracted from whole blood samples using the standard salt extraction alcohol precipitation method (Aljanabi & Martinez 1997). DNA concentration and purity were assessed spectrophotometrically (ND-1000-Spectrophotometer, Thermo Fisher Scientific, Waltham, MA, USA), and DNA integrity checked using gel electrophoresis (50 ng of DNA, 0.8% agarose gel at 100 mV for 60 min, Midori Green staining (NIPPON Genetics Europe, Düren, Germany). Each sample was then diluted to 1.2 ng.µl^{-1}, and the final reaction contained 6 ng of DNA, 300nM of forward and reverse primers and SensiFAST SYBR Lo-ROX master mix (Bioline, London, UK) for a total volume of 12 µl. Tel1b was used as a forward telomere primer (5'-CGGTTTGTTTGGGTTTGGGTTTGGGTTTGGGTTTGGGTT-3'), Tel2b as the reverse one (5'-GGCTTGCCTTACCCTTACCCTTACCCTTACCCTTACCCT-3') and RAG1 as a single copy gene (forward primer 5'-GCAGATGAACTGGAGGCTATAA-3' and reverse primer 5'-CAGCTGAGAAACGTGTTGATTC-3') following Kärkkäinen *et al.* (2021). The qPCR conditions started with an initial denaturation (3 min at 95°C), then 40 cycles with first step of 10s at 95°C, second step of 15s at 58°C and third step of 10s at 72°C, with melting curve analysis in the end performed by QuantStudio™ 12K Flex Real-Time PCR System (Thermo Fisher Scientific, Waltham, MA, USA) using 384-well plates. Each plate contained negative control and three internal standards. Samples from the different experimental treatments were distributed evenly across all plates. To determine the baseline fluorescence, the quantification cycle values and the qPCR efficiencies, we used LinRegPCR (Ruijter *et al.* 2009). Relative telomere length was calculated based on plate-specific efficiencies (average ± SE for telomere and RAG1 reactions were, respectively 95.1 ± 1.6% and 101.7 ± 1.8%) using the mathematical model presented in Pfaffl (2001). Technical repeatability based on triplicate measurements of telomere length was 0.74 (95% CI [0.69, 0.79], *p* < 0.001), and inter-plate repeatability based on samples measured on more than one plate was 0.80 (95% CI [0.67, 0.93], *p* <

0.001). Molecular sexing was conducted using qPCR (adapted from Ellegren & Fridolfsson 1997, Chang *et al.* 2008; see details in Hsu *et al.* (2020)).

Statistical analysis:

Statistical analysis was carried out using R software version 4.3.2 (R Core Team 2014). Normality and homogeneity of residual variances were checked using the R package DHARMa (Hartig 2022).

Effect of the experimental manipulations and year on nestbox temperature:

For each year, to evaluate nestbox temperature (*i.e.* hourly average) differences among groups, we used linear mixed-effects models (LMMs - implemented in the lme4 package (Bates *et al.* 2015), Satterthwaite's degrees of freedom), with group (2018: heated vs. non-heated; 2019; cooled vs. non-cooled) and time of day (*i.e.* treated as a categorical variable) treated as fixed effects. We included random intercepts for date (to account for temporal autocorrelation) and rearing nestbox (to control for repeated measurements).

To investigate natural variation in ambient and nestbox temperatures between 2018 and 2019 (for each year, between June 4th to July 12th), LMMs were used on hourly air temperature (provided by the Finnish Meteorological Institute and collected by Artukainen weather station in Turku, 2 to 4 km from the study sites) and nestbox temperature (*i.e.* only non-heated and non-cooled), with year and time of day included as fixed effects, and date modeled as a random effect to account for repeated measurements within days. Model parameters were estimated using restricted maximum likelihood (REML).

Effect of the experimental manipulations of temperature on growth:

The effect of heating treatment on growth was analysed previously and no effect was detected at any age for body mass and tarsus length (Hsu *et al.* 2020). We conducted similar analyses for the cooling experiment, and no significant effects of the treatment were detected for body mass ($t = 0.39$, $P = 0.70$) and wing length ($t = 0.80$, $P = 0.43$).

Effect of the experimental manipulations of temperature on telomere length:

A LMM on relative telomere length for each dataset was conducted separately to assess the effects of the treatment (heated vs. non-heated nests; cooled vs. non-cooled nests) and other traits of interest (i.e sex, cross fostering, hatching date and brood size). Nest of origin, nest of rearing and

the qPCR plate identities were integrated in the model as random intercepts. No significant sex-dependent effects of temperature treatments on telomere length were found ($P > 0.08$), and thus interactions between sex and treatment were removed from final models and are not discussed further.

Year effect on telomere length:

As experimental manipulations differed markedly between years (*e.g.* timing of temperature manipulation), we do not compare directly the two experiments. Yet, environmental conditions (including ambient temperature, see results) differed between the two study years, and thus comparing the control groups between both years (i.e. non-heated from 2018 and non-cooled from 2019) may be informative regarding the response to longer-term natural variation in temperature. This was conducted using a linear mixed model relatively similar to the ones described above, with telomere length as a response variable, the different traits of interest (i.e year, sex, cross fostering, hatching date and brood size) as fixed effects and three random intercepts (i.e nest of origin, nest of rearing and qPCR plate identities).

Results

Nestbox temperature: experimental effects and natural variation between years

In 2018, the temperature in heated nest boxes was on average higher than in non-heated ones (estimate: +2.83°C, 95% CI: [1.73 ; 3.53], t = 6.09, p < 0.001, Fig 1A). In 2019, the temperature in cooled nestboxes was on average lower than in non-cooled nestboxes (estimate: -1.72°C, 95% CI: [-1.56 ; -0.45], t = -3.52, p < 0.01, Fig 1 B). Yet, there were in both cases strong interactions between time of day and experimental treatments (2018: $F_{23.1}$=31.2, p<0.001; 2019: $F_{23.1}$=103.7, p<0.001), with treatment effect being more pronounced during mid-day (Fig. 1A and 1B). For instance, nestbox temperature at 16:00 was *ca.* 3.0°C higher in heated nests (Fig. 1A) and *ca.* 2.0°C lower in cooled nests (Fig. 1B).

During the experiments (June-July), ambient temperature was significantly higher in 2019 than 2018 (estimate: +1.54°C, 95% CI: [0.93 ; 2.15], t = 4.98, p < 0.001, Fig 1 C). This temperature difference was also observed in nestbox temperatures between non-heated and non-cooled groups with higher temperature in 2019 than 2018 (estimate: 2.16°C, 95% CI: [1.08 ; 3.24], t = 3.92, p < 0.01, Fig 1 D).

Fig 1: Experimental and natural variation in nestbox and ambient temperatures: effect of experimental heating (A) and cooling (B) on nestbox temperature; natural variation between years in ambient (C) and nestbox (D) temperatures. The black bar represents the time of cooling or heating pads replacement. Hourly mean ± SE are reported for 41 nests in 2018 (21 heated vs; 20 non-heated) and 48 nests in 2019 (23 cooled vs. 25 non-cooled).

Effect of the experimental manipulations of nestbox temperature on telomere length

Nestlings from heated nests had significantly shorter telomeres than non-heated ones (LMM: $t_{27.7}$ = 2.85, *P* = 0.008; Fig.2A; Table 1A) and nestlings from cooled nests had shorter telomeres compared to the ones from non-cooled nests (LMM: $t_{36.3}$ = 2.33, *P* = 0.026; Fig. 2B; Table 1B).

Fig. 2: Effect of experimental heating (A) and cooling (B) on relative telomere length in pied flycatcher nestlings at day 13 after hatching. Individual raw data points are presented along with estimated marginal means ± SE from statistical models described in Table 1.

Table 1: Summary of the linear mixed models testing the effects of experimental temperature manipulation ((A) heating treatment in 2018, (B) cooling treatment in 2019) and relevant covariates on relative telomere length of pied flycatcher nestlings at day 13 after hatching. Non-significant interactions between treatment and sex were removed from final models, but are presented in the table for completeness. Singular fit partly occurred for the nest of rearing and the identity of qPCR plate, but they were kept as random intercepts as they were part of the experimental design.

(A) Experimental heating (2018, N = 97 nestlings)

Random effects:		Variance			
Nest of origin (N=26)	Intercept	0.0519			
Nest of rearing (N=41)	Intercept	0			
Plate qPCR (N=6)	Intercept	0			
Residual		0.2178			

Fixed effects:	Estimate	Std. Error	t	df	P (>|t|)
Intercept	0.802	0.127	6.32	45.22	< 0.001
Heating (Non-heated)	**0.140**	**0.049**	**2.85**	**27.67**	**0.008**
Sex (Male)	-0.054	0.046	-1.18	89.35	0.24
Cross-fostering (Yes)	0.062	0.049	1.26	62.65	0.21
Hatching date	0.004	0.003	1.38	39.05	0.18
Brood size	0.023	0.021	1.08	36.86	0.29

Heating (Non-heated) × Sex (Male)	-0.156	0.092	-1.69	88.52	0.094

- -

Sample size: N = 55 nestlings from non-heated nest, N = 42 nestlings from heated nest

___

(B) Experimental cooling (2019, N = 56 nestlings)

Random effects:		Variance
Nest of origin (N=29)	Intercept	0.002
Nest of rearing (N=48)	Intercept	< 0.001
Plate qPCR (N=10)	Intercept	0.023
Residual		0.053

Fixed effects:	Estimate	Std. Error	t	df	P (>|t|)
Intercept	0.477	0.207	2.30	29.93	0.029
Cooling (Non-cooled)	**0.161**	**0.069**	**2.33**	**36.31**	**0.026**
Sex (Male)	0.031	0.071	0.43	44.98	0.67

Cross-fostering (Yes)	-0.020	0.069	-0.28	47.27	0.78
Hatching date	0.016	0.008	2.07	14.90	0.056
Brood size	0.045	0.027	1.68	30.53	0.10
Cooling (Non-cooled) × Sex (Male)	-0.027	0.147	-0.18	45.31	0.86

Sample size: N = 31 nestlings from non-cooled nest, N = 25 nestlings from cooled nest

Natural variation in telomere length between years

We observed no significant association between year and telomere length (LMM: $t_{37.29}$ = -0.26, P = 0.80; Fig. 3, Table 2).

Fig. 3: Effect of year on relative telomere length in pied flycatcher nestlings at day 13 after hatching from control nests. Individual raw data points are presented along with estimated marginal means ± SE from statistical models described in Table 2.

Table 2: Summary of the linear mixed model testing the effects of the year (2018 vs. 2019) and relevant covariates on relative telomere length of pied flycatcher nestlings at day 13 after hatching from control groups (non-heated from 2018 and non-cooled from 2019). Non-significant interaction between year and sex was removed from the final model, but is presented in the table for completeness. Boundary singular fit occurred for the nest of rearing and nest of origin, but they were kept as a random intercept since it was part of the experimental design.

Control groups treatment (2018 and 2019, N = 86 nestlings)

Random effects:		Variance
Nest of origin (N=35)	Intercept	0
Nest of rearing (N=40)	Intercept	0
Plate qPCR (N=16)	Intercept	0.019
Residual		0.057

Fixed effects:	Estimate	Std. Error	t	df	P (>|t|)
Intercept	0.941	0.156	6.05	37.29	< 0.001
Year (2019)	-0.024	0.093	-0.26	5.41	0.80
Sex (Male)	-0.062	0.058	-1.07	76.94	0.28
Cross-fostering (Yes)	-0.065	0.059	-1.10	78.67	0.28

Brood size	0.046	0.028	1.62	71.69	0.11

Year (2019) × Sex (Male)	0.205	0.124	1.65	79.74	0.103

Sample size: N = 55 nestlings from 2018, N = 31 nestlings from 2019

Discussion

In this study, we demonstrated that exposure to experimental changes in environmental temperature (warmer or cooler) during early-life can significantly shorten telomeres in pied flycatcher nestlings. Since avian altricial nestlings transition from ectothermy to endothermy during development (Price & Dzialowski 2018), these results echo the effects of experimental increase of temperature on telomere lengths in ectotherms, such as fishes (Siberian sturgeon (Simide *et al.* 2016), Atlantic Salmon at the embryo stage (McLennan *et al.* 2018)) or squamates (Desert Agama (Zhang *et al.* 2018)). To the best of our knowledge, only one cooling experiment has been conducted in ectotherms, showing no significant effect of a transition from 30°C to 20°C (Eastern mosquitofish (Rollings *et al.* 2014). Yet, it has been recently shown in amphibians that cooler temperature, for a similar growth rate, is associated with shorter telomeres (Burraco *et al.* 2023). So, our observations of shorter telomeres in response to experimental heating and cooling seem consistent with an ectothermic-like phenotype. Indeed, high temperature has also been shown to accelerate telomere shortening in ectothermic or partly ectothermic avian embryo and nestlings (Stier *et al.* 2020a, Stier *et al.* 2025, Eastwood *et al.* 2022), but also in endotherms such as humans (Ni *et al.* 2022). Future studies investigating temperature effects in fully endothermic avian models (*e.g.* after fledging) are needed to establish the potential role of (partial) ectothermy on the effects of temperature on telomere shortening.

In contrast with the results stemming from our experimental manipulation, we observed that while the mean ambient temperature of 2019 was 1.5°C higher than in 2018 (and nestbox temperature 2.2°C higher), there was no significant year effect on telomere length. This could suggest that long-term variation in temperature has less effect on telomere length than abrupt changes mimicked by our experimental conditions, although this remains to be experimentally demonstrated. Indeed, an alternative explanation could be that the higher temperature in 2019 contributed to accelerate telomere shortening, but that higher food availability, usually resulting from warmer environmental temperatures (Avery & Krebs 1984, Veistola *et al.* 1997), compensated for any temperature-related shortening. Indeed, better nutritional conditions have been shown to limit telomere loss, for instance in great tit nestlings (Casagrande *et al.* 2023).

The temperature-related telomere shortening we observed in this study in response to experimentally increased or decreased temperature may be explained by several potential mechanisms. First, telomere shortening has been shown to be accentuated by fast growth and rapid cellular proliferation (Monaghan & Ozanne 2018). This seems unlikely to explain the effects we observed on telomeres since experimental heating (Hsu *et al.* 2020) or cooling (present study) had no

detectable effects on body mass or body size growth. Second, telomere shortening can be accelerated by increased oxidative stress levels (Reichert & Stier 2017), and acute heat or cold exposure have been shown to potentially increase oxidative stress levels (Costantini *et al.* 2012, Stier *et al.* 2014). Yet, this seems unlikely to explain the effects we observed on telomeres since experimental heating had no detectable effects on oxidative stress levels (Hsu *et al.* 2020). Third, telomere shortening can be accelerated by energetic imbalance, because of potential trade-offs between immediate and long-term survival during challenging environmental conditions (Casagrande & Hau 2019). This has been suggested to be mediated by glucocorticoid hormone (*i.e.* "stress hormone") effects on cellular signalling via the mTOR pathway (Casagrande & Hau 2019). Since abrupt changes in temperature can increase glucocorticoid levels in altricial nestlings (Jessop *et al.* 2016, De Bruijn & Romero 2018), it is possible that the telomere shortening could be explained by this mechanism, as already suggested in an experimental heating study in great tit (Stier *et al.* 2025). Yet, measuring glucocorticoid levels and mTOR signalling would be required to ascertain this hypothesis.

To conclude, we have demonstrated that experimental increase or decrease in early-life temperature can shorten telomeres, which may impact long-term survival and fitness. Yet, directly measuring effects of experimental temperature manipulations on fitness would be needed to ascertain this hypothesis, which is challenging in small passerine species having very low survival rates after fledging. Consequently, investigating those effects in long-lived species with high recruitment rate and known relationship between telomere length and fitness related traits (e.g. Alpine swift, Bize *et al.* 2009) would seem promising to better understand the effects of heatwaves and cold-snaps during early-life on telomeres and fitness.

Author contribution:

SRu, AS, CM and BYH designed the study and conducted fieldwork. CM, AS and NCS conducted laboratory work. CF, JF and AS analyzed the data. CF, JF & AS wrote the manuscript, with input from SRu, SRe, NCS & CM. SRu and SRe acquired funding. SRu, SRe and AS supervised early-career researchers involved in this study.

Funding

This study was financially supported by the Academy of Finland (#286278 to SRu and #356397 to SRe). AS and SRe were both supported by a 'Turku Collegium for Science and Medicine' Fellowship and Marie Sklodowska-Curie Postdoctoral Fellowships (#894963 and #101110339). All authors declare no conflict of interest.

Acknowledgements

We are grateful to Lucas Bousseau, Thomas Rossille, Hsiao-Yin Liu, Päivi Kotitalo, Mélanie Crombecque, Axelle Delaunay, Lotta Holmen and Mikaela Hukkanen for their help in the field. The laboratory facilities were supported by Biocenter Finland.

Data accessibility

All data are accessible at the private link: https://figshare.com/s/eb091d4259185e8646a6, and will be publicly available at the time of publication.

Ethical permits

All procedures were approved by the Animal Experiment Committee of the State Provincial Office of Southern Finland (license numbers ESAVI/2902/2018 & ESAVI/5718/2019) and by the Environmental Center of Southwestern Finland (license numbers VARELY/549/2018 & VARELY/924/2019) granted to SRu.

Conflict of interest

The authors declare no competing or financial interests.

References

Aljanabi, S. M. & Martinez, I. 1997. Universal and rapid salt-extraction of high quality genomic DNA for PCR-based techniques. Nucleic Acids Res. 25: 4692–4693. doi:10.1093/nar/25.22.4692

Andreasson, F., Nord, A., Nilsson, J. A. 2018. Experimentally increased nest temperature affects body temperature, growth and apparent survival in blue tit nestlings. Journal of Avian Biology. 1–14. doi:10.1111/jav

Avery, M. & Krebs, J. R. 1984. Temperature and foraging success of great tits Parus major hunting for spiders. Ibis. 126(1):33–38. https://doi.org/10.1111/j.1474-919X.1984.tb03661.x

Bates, D., Maechler, M., Bolker, B., Walker, S. 2015. Fitting linear mixed-effects models using lme4. J. Stat. Softw. 67, 1–48. https://doi.org/10.18637/jss.v067.i01

Berntsen, H. H., & Bech, C. 2016. Incubation temperature influences survival in a small passerine bird. J Avian Biol. 47: 141-145. https://doi.org/10.1111/jav.00688

Bize, P., Criscuolo, F., Metcalfe, N., Nasir, L., Monaghan, P. 2009. Telomere dynamics rather than age predict life expectancy in the wild. Proceedings of the Royal Society B: Biological Sciences. 276, 1679–1683. https://doi.org/10.1098/rspb.2008.1817

Boyles, J. G., Seebacher, F., Smit, B., McKechnie, A. E. 2011. Adaptive Thermoregulation in Endotherms May Alter Responses to Climate Change. Integrative and Comparative Biology. 51: 676–690. doi:10.1093/icb/icr053

Burraco, P., Hernandez-Gonzalez, M., Metcalfe, N. B., Monaghan, P. 2023. Ageing across the great divide: tissue transformation, organismal growth and temperature shape telomere dynamics through the metamorphic transition. Proc Royal Soc B 290, 20222448. doi: 10.1098/rspb.2022.2448.

Casagrande, S. & Hau, M. 2019. Telomere attrition: metabolic regulation and signalling function? Biology Letters. 15: 20180885–11. doi:10.1098/rsbl.2018.0885

Casagrande, S., Stier, A., Monaghan, P., Loveland, J. L., Boner, W., Lupi, S., Trevisi, R., Hau, M. 2020. Increased glucocorticoid concentrations in early life cause mitochondrial inefficiency and short telomeres. Journal Of Experimental Biology. 223, jeb222513-14. doi: 10.1242/jeb.222513.

Casagrande, S., Loveland, J. L., Oefele, M., Boner, W., Lupi, S., Stier, A., Hau, M. 2023. Dietary nucleotides can prevent glucocorticoid-induced telomere attrition in a fast-growing wild vertebrate. Mol. Ecol. 32, 5429–5447. https://doi.org/10.1111/mec.17114

Chang, H. W., Gu, D. L., Su, S. H., Chang, C. C., Cheng, C. A., Huang, H. W., Yao, C.T., Chou, T. C., Chuang, L. Y., Cheng, C. C. 2008. High-throughput gender identification of Accipitridae eagles with real-time PCR using TaqMan probes. Theriogenology. 70(1):83-90. doi: 10.1016/j.theriogenology.2008.02.011

Chatelain, M., Drobniak, S. M., Szulkin, M. 2019. The association between stressors and telomeres in non-human vertebrates: a meta-analysis. Ecology Letters. 99, 21–18. https://doi.org/10.1111/ele.13426

Conradie, S. R., Woodborne, S. M., Cunningham, S. J., McKechnie, A. E. 2019. Chronic, sublethal effects of high temperatures will cause severe declines in southern African arid-zone birds during the 21st century. Proceedings Of The National Academy Of Sciences Of The United States Of America. 116: 14065–14070. doi:10.1073/pnas.1821312116

Costantini, D., Monaghan, P., Metcalfe, N. B. 2012. Early life experience primes resistance to oxidative stress. Journal Of Experimental Biology. 215, 2820–2826. doi: 10.1242/jeb.072231.

De Bruijn, R. & Romero, L. M. 2018. The role of glucocorticoids in the vertebrate response to weather. Gen Comp Endocrinol. 269:11-32. doi: 10.1016/j.ygcen.2018.07.007.

Eastwood, J. R., Hall, M. L., Teunissen, N., Kingma, S. A., Aranzamendi, N. H., Fan, M., Roast, M., Verhulst, S. and Peters, A. 2019. Early-life telomere length predicts lifespan and lifetime reproductive success in a wild bird. Molecular Ecology 28:1127–1137.

Eastwood, J. R., Connallon, T., Delhey, K., Hall, M. L., Teunissen, N., Kingma, S. A., La Porte, A. M., Verhulst, S., Peters, A. 2022. Hot and dry conditions predict shorter nestling telomeres in an endangered songbird: Implications for population persistence, Proc. Natl. Acad. Sci. U.S.A. 119 (25) e2122944119, https://doi.org/10.1073/pnas.2122944119

Ellegren, H. & Fridolfsson, A. K. 1997. Male–driven evolution of DNA sequences in birds. Nat Genet. 17, 182–184. https://doi.org/10.1038/ng1097-182

Friesen, C. R., Wapstra, E., Olsson, M. 2022. Of telomeres and temperature: Measuring thermal effects on telomeres in ectothermic animals. Mol. Ecol. 31, 6069–6086. doi: 10.1111/mec.16154.

Greño, J. L., Belda, E. J., Barba, E. 2008. Influence of temperatures during the nestling period on post-fledging survival of great tit Parus major in a Mediterranean habitat. J Avian Biol. 39: 41–49. doi:10.1111/j.0908-8857.2008.04120.x

Hartig, F. 2022. DHARMa: Residual Diagnostics for Hierarchical (Multi-Level/Mixed) Regression Models. R package version 0.4.6, https://CRAN.R-project.org/package=DHARMa

Hepp, G. R. & Kennamer, R. A. 2012. Warm is better: incubation temperature influences apparent survival and recruitment of wood ducks (Aix sponsa). PLoS ONE. 7: e47777. doi:10.1371/journal.pone.0047777

Hope, S. F., Angelier, F., Ribout, C., Groffen, J., Kennamer, R. A., Hopkins, W. A. 2023. Warmer incubation temperatures and later lay-orders lead to shorter telomere lengths in wood duck (Aix sponsa) ducklings. Journal of Experimental Zoology Part A: Ecological and Integrative Physiology. 339(1), 101–111. https://doi. org/10.1002/jez.2659

Hsu, B. Y., Sarraude, T., Cossin-Sevrin, N., Crombecque, M., Stier, A., Ruuskanen, S. 2020. Testing for context-dependent effects of prenatal thyroid hormones on offspring survival and physiology: an experimental temperature manipulation. Sci Rep. 10, 14563. https://doi.org/10.1038/s41598-020-71511-y

Jessop, T. S., Lane, M. L., Teasdale, L., Stuart-Fox, D., Wilson, R. S., Careau, V., Moore, I. T. 2016. Multiscale Evaluation of Thermal Dependence in the Glucocorticoid Response of Vertebrates. The American Naturalist. 188(3), 342–356. https://www.jstor.org/stable/26519175

Kärkkäinen, T., Teerikorpi, P., Panda, B., Helle, S., Stier, A., Laaksonen, T. 2019. Impact of continuous predator threat on telomere dynamics in parent and nestling pied flycatchers. Oecologia. 191, 757-766. doi:10.1007/s00442-019- 04529-3

Kärkkäinen, T., Bize, P., Stier, A. 2020. Correlation in telomere lengths between feathers and blood cells in pied flycatchers. Journal of Avian Biology. 51(4), Article e02300. https://doi.org/10.1111/jav.02300

Kärkkäinen, T., Teerikorpi, P., Schuett, W., Stier, A., Laaksonen, T. 2021. Interplays between pre- and post-natal environments affect early-life mortality, body mass and telomere dynamics in the wild. J Exp Biol. 224(Pt 1):jeb231290. doi: 10.1242/jeb.231290.

Kärkkäinen, T., Laaksonen, T., Burgess, M., Cantarero, A., Martínez-Padilla, J., Potti, J., Moreno, J., Thomson, R. L., Tilgar, V., Stier, A. 2022. Population differences in the length and early-life dynamics of telomeres among European pied flycatchers. Mol Ecol. 31(23):5966-5978. doi: 10.1111/mec.16312.

Katzenberger, J., Tate, G., Koeslag, A., Amar, A. 2015. Black Sparrowhawk brooding behaviour in relation to chick age and weather variation in the recently colonised cape peninsula, South Africa. Journal of Ornithology. 156(4), 903–913. https://doi.org/10.1007/s1033 6-015-1199-0

Lange, T. D., Lundblad, V., Blackburn, E. H. 2006. Telomeres. Cold Spring Harbor Laboratory Press, New York 576.

Lundberg, A. & Alatalo, R. 1992. The Pied Flycatcher (Poyser, London)

McLennan, D., Armstrong, J. D., Stewart, D. C., Mckelvey, S., Boner, W., Monaghan, P., Metcalfe, N. B. 2018. Telomere elongation during early development is independent of environmental temperatures in Atlantic salmon. J Exp Biol. 221(Pt 11):jeb178616. doi: 10.1242/jeb.178616.

McKechnie, A. E. & Wolf, B. O. 2009. Climate change increases the likelihood of catastrophic avian mortality events during extreme heat waves. Biology Letters. 6(2), 253–256. https://doi.org/10.1098/rsbl.2009.0702

McKechnie, A. E., Rushworth, I. A., Myburgh, F., Cunningham, S. J. 2021. Mortality among birds and bats during an extreme heat event in eastern South Africa. Austral Ecology. doi:10.1111/aec.13025

Metcalfe, N. & Monaghan, P. 2001. Compensation for a bad start: grow now, pay later? Trends in Ecology & Evolution. 16(5):254-260. doi: 10.1016/s0169-5347(01)02124-3.

Monaghan, P. & Ozanne, S. E. 2018. Somatic growth and telomere dynamics in vertebrates: relationships, mechanisms and consequences. Philos Trans R Soc Lond B Biol Sci. 373(1741):20160446. doi: 10.1098/rstb.2016.0446.

Moreno, J., González-Braojos, S., Ruiz-de-Castañeda, R. 2015. A spring cold snap is followed by an extreme reproductive failure event in a mountain population of Pied Flycatchers Ficedula hypoleuca. Bird Study. 62(4), 466–473. https://doi.org/10.1080/00063657.2015.1073680

Ni, W., Wolf, K., Breitner, S., Zhang, S., Nikolaou, N., Ward-Caviness, C. K., Waldenberger, M., Gieger, C., Peters, A., Schneider, A. 2022. Higher Daily Air Temperature Is Associated with Shorter Leukocyte Telomere Length: KORA F3 and KORA F4. Environ. Sci. Technol. 56, 17815–17824. doi: 10.1021/acs.est.2c04486.

Nord, A. & Nilsson, J. A. 2016. Long-term consequences of high incubation temperature in a wild bird population. Biology Letters. 1–4. doi:10.1098/rsbl.2016.0087&domain=pdf&date_stamp=2016-04-05

Ospina, E. A., Merrill, L., Benson, T. J. 2018. Incubation temperature impacts nestling growth and survival in an open-cup nesting passerine. Ecol Evol. 8(6):3270-3279. doi: 10.1002/ece3.3911.

Pfaffl, M. W. 2001. A new mathematical model for relative quantification in real-time RT–PCR. Nucl. Acids Res. 29(9):e45. doi: 10.1093/nar/29.9.e45.

Preston, J. D., Reynolds, L. J., Pearson, K. J. 2018. Developmental Origins of Health Span and Life Span: A Mini-Review. Gerontology. 64: 237–245. doi:10.1159/000485506

Price, E. R. & Dzialowski, E. M. 2018. Development of endothermy in birds: Patterns and mechanisms. Journal of Comparative Physiology B. 188(3), 373–391. https://doi.org/10.1007/s0036 0-017-1135-0

R Core Team. 2014. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. http://www.R-project.org/.

Reichert, S. & Stier, A. 2017. Does oxidative stress shorten telomeres in vivo? A review. Biol Lett. 13(12):20170463. doi: 10.1098/rsbl.2017.0463.

Remot, F., Ronget, V., Froy, H., Rey, B., Gaillard, J.-M., Nussey, D. H., Lemaitre, J.-F. 2022. Decline in telomere length with increasing age across nonhuman vertebrates: A meta-analysis. Molecular Ecology. 31(23), 5917–5932. https://doi.org/10.1111/mec.16145

Rodríguez, S. & Barba, E. 2016. Nestling Growth is Impaired by Heat Stress: an Experimental Study in a Mediterranean Great Tit Population. Zoological studies. 55: e40. doi:10.6620/zs.2016.55-40

Rollings, N., Miller, E., Olsson, M. 2014. Telomeric attrition with age and temperature in Eastern mosquitofish (Gambusia holbrooki). Naturwissenschaften. 101(3):241-4. doi: 10.1007/s00114-014-1142-x.

Ruijter, J. M., Ramakers, C., Hoogaars, W. M. H., Karlen, Y., Bak-ker, O., van den Hoff, M. J., Moorman, A. F. M. 2009. Amplification efficiency: linking baseline and bias in the analy-sis of quantitative PCR data. Nucl. Acids Res. 37: e45–e45. doi: 10.1093/nar/gkp045.

Simide, R., Angelier, F., Gaillard, S., Stier, A. 2016. Age and Heat Stress as Determinants of Telomere Length in a Long-Lived Fish, the Siberian Sturgeon. Physiol Biochem Zool. 89(5):441-7. doi: 10.1086/687378.

Stier, A., Massemin, S., Criscuolo, F. 2014. Chronic mitochondrial uncoupling treatment prevents acute cold-induced oxidative stress in birds. Journal Of Comparative Physiology B-Biochemical Systemic And Environmental Physiology. 184, 1021–1029. doi: 10.1007/s00360-014-0856-6.

Stier, A., Metcalfe, N. B., Monaghan, P. 2020a. Pace and stability of embryonic development affect telomere dynamics: An experimental study in a precocial bird model. Proceedings of the Royal Society B: Biological Sciences. 287(1933), 20201378. https://doi.org/10.1098/rspb.2020.1378

Stier, A., Hsu, B.-Y., Marciau, C., Doligez, B., Gustafsson, L., Bize, P., Ruuskanen, S. 2020b. Born to be young? Prenatal thyroid hormones increase early-life telomere length in wild collared flycatchers. Biology Letters. 16, 20200364–4. doi: 10.1098/rsbl.2020.0364.

Stier, A., Hsu, B.-Y., Cossin-Sevrin, N., Garcin, N., Ruuskanen, S. 2021. From climate warming to accelerated cellular ageing: An experimental study in wild birds. bioRxiv. https://doi.org/10.1101/2021.12.21.473625

Ton, R., Boner, W., Raveh, S., Monaghan, P., Griffith, S. C. 2023. Effects of heat waves on telomere dynamics and parental brooding effort in nestlings of the zebra finch (Taeniopygia castanotis) transitioning from ectothermy to endothermy. Mol Ecol. 32(17):4911-4920. doi: 10.1111/mec.17064.

Vedder, O., Verhulst, S., Zuidersma, E., Bouwhuis, S. 2018. Embryonic growth rate affects telomere attrition: an experiment in a wild bird. J Exp Biol. 221(Pt 15):jeb181586. doi: 10.1242/jeb.181586.

Veistola, S., Lehikoinen, E., Eeva, T. 1997. Weather and breeding success at high latitudes - The Pied Flycatcher Ficedula hypoleuca and the Siberian Tit Parus cinctus. Ornis Fennica. 74. 89-98.

Wilbourn, R. V., Moatt, J. P., Froy, H., Walling, C. A., Nussey, D. H., Boonekamp, J.J. 2018. The relationship between telomere length and mortality risk in non-model vertebrate systems: a meta-analysis. Philosophical transactions of the Royal Society of London Series B, Biological sciences. 373: 20160447–9. doi:10.1098/rstb.2016.0447

Zhang, Q., Han, X., Hao, X., Ma, L., Li, S., Wang, Y., Du, W. 2018. A simulated heat wave shortens the telomere length and lifespan of a desert lizard. J Therm Biol. 72:94-100. doi: 10.1016/j.jtherbio.2018.01.004.