Complex three-dimensional rearing environments amplify compensatory

plasticity following early blindness

Deepa L. Ramamurthy[1, 2], Mackenzie Englund[1], Tanner J. Kovacs[1], Heather Dodson[1],

Leah A. Krubitzer[1,3].

[1]Center for Neuroscience, University of California, Davis, Davis CA 95618

[2]Present address: Helen Wills Neuroscience Institute, University of California, Berkeley,

189 Weill Hall, Berkeley, CA 94720

[3]Lead Contact

*Correspondence: lakrubitzer@ucdavis.edu

SUMMARY

The neocortex has a remarkable capacity to alter its functional organization and

connectivity in response to sensory loss, particularly if this loss occurs early in life. A key

question is whether this cross-modal reorganization is driven by sensory deprivation itself

or through enhanced use of the spared senses. We investigated how different rearing

environments shape neural responses in primary somatosensory cortex (S1) of short-

tailed opossums (*Monodelphis domestica)*, following elimination of visual inputs through

bilateral enucleation in early development. Early blind and sighted littermates were reared

in enriched environments to promote active tactile exploration in three-dimensional (3D)

space, or in highly restricted standard laboratory cages. In adulthood, both enriched

groups showed adaptive changes in exploration patterns and gap crossing behaviors

relative to standard-reared counterparts. Thus, early blind animals showed behavioral

compensation for vision loss, when challenged by complex environments. Enriched

rearing increased selectivity of S1 neural responses to whisker touch and powerfully

altered receptive field shapes such that they were less horizontally anisotropic. This shift

was strongest in enriched early blind animals, enhancing tuning along the horizontal axis

more than standard-reared early blind animals. Thus, alterations in S1 receptive fields

following early blindness were amplified by environmental complexity. Sighted opossums

reared with enrichment also showed similar whisker receptive field plasticity, though to a

slightly lower degree. Together, these results demonstrate the strong influence of the

rearing environment on reorganization of cortex that processes inputs from the spared

senses, underscoring the role of experience in directing compensatory plasticity following

early sensory loss.

INTRODUCTION

Early loss of vision leads to heightened perceptual abilities in the remaining senses, accompanied by structural and functional changes in the nervous system (Goldreich and Kanics, 2003; Wong et al., 2011; Voss, 2011; Morland et al, 2021; see Mezzera and López-Bendito, 2016; Lazzouni and Lepore, 2014 for review). However, the developmental origins of these changes remain poorly understood—it is not clear to what extent sensory experience, movement options and affordances contribute to compensatory changes in brain and behavior following early sensory loss. In this study, we investigate how enhancing tactile experience through rearing in an enriched environment contributes to the neural and behavioral plasticity of whisker–mediated touch in a marsupial, the short-tailed opossum, when vision is lost very early in development (Ramamurthy and Krubitzer, 2018; Ramamurthy et al., 2021).

It is well established that exposure to enriched environments drives plasticity in sensory cortices of juveniles, and even in adult animals with intact brains and bodies. This includes the functional reorganization of sensory representations through changes in cortical territory, functional properties (latency, strength, and selectivity of neural responses), and the morphology and molecular expression patterns of neurons (Seo, 1992; Leah et al., 1995; Coq and Xerri, 1998; Cancedda et al., 2004; Engineer et al., 2004; Polley et al., 2004; Guic et al., 2008; Mégevand et al., 2009; Devonshire et al., 2010; Alwis and Rajan, 2013, LeMessurier et al., 2018, Bibollet-Bahena 2023). Enrichment also counteracts degradation of representations associated with a transiently deprived sensory modality, and promotes recovery from brain injury (Polley et al., 1999; Bartoletti et al., 2004; Sale et al, 2007; Nakashima and Dyck, 2008; Wang et al., 2013;

Greifzu et al., 2014; Zhu et al., 2014; Zheng et al., 2014; Jiang et al., 2015; Kalogeraki et

al., 2017; Levine et al., 2017). Enrichment through naturalistic experiences in accordance

with a species' lifestyle engages animals in an ethologically relevant manner (Landers et

al., 2011; Wilkinson et al, 2010; Branchi et al., 2011; Weissbrod et al, 2013) and is

especially effective in reshaping cortical sensory representations at structural, functional

and molecular levels (Polley et al., 1999; Polley et al., 2004; Frostig, 2006; Gomez-Pinilla

et al., 2011; Landers et al., 2011). Therefore, in this study, we used a complex 3D rearing

environment designed to mimic the natural habitat and lifestyle of short-tailed opossums.

Many small mammals, including short-tailed opossums, exhibit a continuum in

patterns of vertical use of strata in a forest habitat, from the forest floor to the tree canopy

(Jenkins, 1974; Jones et al., 2003; Delciellos and Vieira, 2006; Abreu and Oliveira, 2014;

Shapiro et al., 2014). Short-tailed opossums are primarily terrestrial hunters and foragers

(Jones et al., 2003) but prefer nesting above the ground in rocky outcrops (Macrini, 2004),

using arboreal runways like branches and logs to negotiate complex terrain (Lammers,

2007; Lammer and Gaunter 2008; Ladine and Kissel, 1994; Montgomery, 1980). Short-

tailed opossums are skilled climbers (Saunders et al., 1995; Saunders et al., 1998;

Wilkinson et al., 2010) despite lacking many of the morphological specializations

observed in more arboreal species (Schmitt et al., 2010; Rupert et al., 2014; Goin et al.,

2016; Thomas et al., 2016). While they exhibit some arboreal traits—including some

degree of forelimb specialization and tail prehensility (Jones et al., 2003; Schmitt et al.,

2010; Rupert et al., 2014; Thomas et al., 2016)—short-tailed opossums primarily rely on

behavioral strategies to navigate and maintain balance on arboreal substrates (Lammers,

2004; Lammers, 2007; Lammers, 2009; Lammers and Biknevicius, 2004; Lammers et al.,

2006; Lammers and Gauntner, 2008). Whisker touch is critical for climbing in spatially

complex 3D environments (Ahl, 1986; Arkley et al., 2017), since it guides gap crossing

and forelimb placement (Arkley et al., 2014; Niederschuh et al., 2015; Arkley et al., 2017;

Grant et al., 2018). Peripheral organization of the whisker system is closely associated

with lifestyle, being more well-developed in arboreal than in terrestrial species (Beddard,

1902; Pocock, 1914; Lyne, 1958; Lyne, 1959; Muchlinski, 2013; Grant et al., 2017; Grant

et al., 2018) – in short-tailed opossums, whisker layout is consistent with semi-arboreal

and scansorial marsupial species (Ramamurthy and Krubitzer, 2016; Grant et al., 2013;

Lyne, 1958; Lyne, 1959). Thus, while short-tailed opossums are highly capable of vertical

utilization of the environment, doing so involves behavioral modifications based on

sensory input, including tactile information (Lammers and Biknevicius, 2004; Lammers,

2009).

In prior work, we showed that early loss of visual inputs in short-tailed opossums

(at P4, prior to the formation of retinogeniculate and thalamocortical pathways) enhances

tactile acuity and alters neural coding of whisker inputs in primary somatosensory cortex

(S1), with receptive fields exhibiting greater selectivity along the natural axis of whisker

motion (Ramamurthy & Krubitzer, 2018). Loss of vision at this early developmental stage

also leads to major reorganization of thalamic and cortical connections of somatosensory

cortex relative to sighted animals (Dooley & Krubitzer, 2019). This raises a critical

question: are alterations in sensory coding in somatosensory cortex contingent on the

deprivation of visual inputs itself, which gives rise to the novel anatomical connections

that may form the substrate for these functional changes? Or does tactile experience drive

functional plasticity, and can enrichment amplify the compensatory neural and behavioral

plasticity observed following early blindness? Here, we implement rearing conditions that

encourage naturalistic patterns of tactile exploration in a spatially challenging

environment designed to increase reliance on whisker use. By comparing how this

enriched environment shapes neural coding of whisker touch stimuli in S1 and whisker-

mediated behavior in sighted and early blind animals, we interrogate the role of tactile

experience in directing compensatory plasticity following early blindness.

RESULTS

Blind and sighted opossums both adapt behaviorally to enriched rearing

environments

Following bilateral enucleation on postnatal day 4, opossum mothers and their

litters, including sighted and blind pups, were transferred into a new standard or enriched

home cage environment (**Figure 1A**; **Figure S1A-D**). Prior to P12, sensory experience in

either environment was determined entirely by patterns of behavior of the mother, due to

the obligate attachment of opossum pups. Juvenile animals continue to rely on the mother

for both sustenance and transportation until weaning at P56, but with increasing

independent exploration (**Figure S1E-F**). Opossum mothers always preferentially nested

high above the ground (3/3 mothers of four enriched experimental litters; example in

Video S1), so pups were forced to rely heavily on touch to navigate the enriched

environment. By adulthood, both sighted and early blind animals were highly adapted to

their environments (**Video S2-S3**). Overall patterns of exploration did not differ between

sighted and blind animals (**Figure S2**; **Figure 1B-C**), indicating successful behavioral

compensation by blind animals even in more spatially complex environments. When

placed in a novel arena with the dimensions and basic layout of the enriched home cages,

enriched opossums showed increased exploration of the vertical space compared to

those reared in standard cages (p<0.001, permutation test; **Figure 1D**). In a gap crossing

task conducted in the dark, opossums across all experimental groups performed fewer

successful crossings when gaps were large (**Figure 1E-F**, 12-14 cm; eSC: 0.17, eEB:

0.14, sSC: 0.08, sEB: 0.14; gap crossings per minute across all trials) vs. small (2-10 cm;

eSC: 2.29, eEB: 2.51, sSC: 2.75, sEB: 3.07). Consistent with previous studies in small

mammals (Hutson & Masterton 1986, Voigts et al., 2015, Arkley et al. 2017, Graham &

Socha 2020, Toso & Diamond 2023), gap crossing was heavily dependent on the

whiskers. All experimental groups failed to cross large gaps when whiskers were trimmed

and performed fewer crossings per minute even for small gaps. Notably, enriched groups

were more affected by whisker trimming compared to standard groups (gap crossings per

minute after whisker trimming – eSC: 1.34, eEB: 1.60, sSC: 2.31, sEB: 2.56), indicating

their increased reliance on whisker-based strategies. In general, for all experimental

groups, time taken to successfully finish crossing was longer for large vs. small gaps

(**Figure 1G**; linear mixed-effect model, p<0.001; p<0.001 for all pairwise comparisons,

permutation test), with small but significant increases in crossing times for small gaps

following whisker trimming (linear mixed-effect model, p<0.05; sSC: p<0.05, sEB:

p<0.001, eSC: p<0.001, eEB: p<0.05, permutation test). Differences between groups

based on blindness or rearing conditions were most apparent for the duration to

successfully cross large gaps. Early blind opossums were faster than sighted controls

even under standard rearing conditions (sSC vs. sEB: p<0.05, permutation test), and their

behavioral proficiency was amplified by enriched rearing (eEB vs. sEB: p<0.001,

permutation test). Both sighted and blind enriched groups were significantly faster than

standard-reared animals (p<0.001 for all pairwise comparisons, permutation test) but not

significantly different from each other (p>0.05).

Tactile experience amplifies compensatory neural plasticity in blind opossums

Single-unit recording experiments were performed in 19 opossums reared in

standard conditions (**Figure 2**; 10 sSC, 9 sEB), and 14 opossums reared in enriched

conditions (**Figure 2**; 7 eSC, 7 eEB), acquiring data for a total of 339 neurons across

experimental groups (see **Table S1**; **Figure S3**). 63-73% of units recorded in S1 (sSC:

60, sEB: 64, eSC: 58, eEB: 51) had a significant evoked response (>2 SD above baseline)

to stimulus onset following single whisker deflections, and these neurons were included

in receptive field analyses. Raster plots and peristimulus time histograms (PSTHs) for

whisker-responsive neurons in each group are shown in **Figure 2A-B**.

As reported previously in standard-reared animals (Ramamurthy and Krubitzer,

2018), spontaneous firing rates of S1 neurons were low for all experimental groups but

were not significantly different among experimental groups (**Figure 2C**; mean ± SEM,

sSC: 0.582 ± 0.083 spk/s, sEB: 0.613 ± 0.121 spk/s, eSC: 0.544 ± 0.079 spk/s, eEB:

0.503 ± 0.082 spk/s; linear mixed-effect model, p>0.05). However, responses evoked by

the stimulation of even the most preferred single whisker stimulus (best whisker; BW) in

enriched groups were suppressed in relation to stimulus-evoked firing rates in standard-

reared sighted opossums (**Figure 2D**; mean ± SEM, sSC: 11.4 ± 0.839 spk/s, sEB: 8.39

± 0.591 spk/s, eSC: 7.82 ± 0.748 spk/s, eEB: 4.57 ± 0.52 spk/s; linear mixed-effects

model, eSC vs. sSC: p<0.01, eEB vs. sSC: p<0.001; see **Table S2** for exact p-values for

the fixed-effect of experimental group, and for all pairwise comparisons between groups),

similar to previous observations in standard-reared early blind animals (for details see

Ramamurthy and Krubitzer, 2018). Since relative suppression of BW-evoked responses

was observed in enriched sighted controls, in addition to enriched early blind animals, this

effect was not contingent on the lack of vision itself – rather it was driven by increased

dependence on touch in the enriched groups of opossums. Mean BW-evoked firing rates

in enriched sighted control animals were comparable to those in standard early blind

animals (linear mixed-effects model, sEB vs. eSC: $p>0.05$). However, in enriched early

blind animals, mean BW-evoked firing rates were lower relative to all other experimental

groups (linear mixed-effects model, eEB vs. sEB: $p<0.001$, eEB vs. eSC: $p<0.001$, eEB

vs. sSC: $p<0.001$). Thus, blindness was not required for suppression of whisker-evoked

responses, but the combination of tactile enrichment and vision loss produced the

greatest effect on whisker-evoked firing rates of S1 neurons. This is consistent with

animals relying on their whiskers to a greater extent when reared in a complex

environment in the absence of vision.

Average whisker receptive fields in both sighted and blind enriched groups were

significantly smaller than those in standard-reared sighted controls (mean ± SEM, eEB:

3.22 ± 0.409, eSC: 3.48 ± 0.364, sSC: 5.92 ± 0.427; linear mixed-effects model, eSC vs.

sSC: $p<0.001$, eEB vs. sSC: $p<0.001$) when measured as the number of whiskers that

evoked a significant response above spontaneous firing (**Figure 2E**). We see similar

results when receptive fields are quantified as the mean rank-ordered tuning curve across

neurons (**Figure 2F**; linear mixed-effects model, eSC vs. sSC: $p<0.001$). The finding that

tactile enrichment in sighted control animals leads to a decrease in mean receptive field

size—similar to both standard-reared and enriched early blind animals—supports the

interpretation that this reduction reflects an experience-dependent effect across all three

groups. However, changes in measures of overall receptive field size in enriched early

blind animals relative to standard-reared early blind animals were either not significant

(**Figure 2E**, left, number of whiskers: eEB: 3.22 ± 0.409, sEB: 3.38 ± 0.270; linear mixed-

effects model, eEB vs. sEB: p>0.05) or modest (**Figure 2E**, right, rank-ordered tuning

curve: linear mixed-effects model, p<0.01). This suggests that, under standard rearing

conditions, receptive field plasticity in early blind animals may already have reached a

limit that cannot be further shifted by enrichment.

Enrichment enhances selectivity for whisker touch along the behaviorally relevant

horizontal axis

To better understand the effects of enriched rearing on receptive fields of S1

neurons, we characterized two-dimensional receptive fields for the row (horizontal) and

arc (vertical) dimensions of the whisker pad (**Figure 3A**). Visual inspection of mean 2D

receptive fields revealed clear differences in receptive field anisotropies across

experimental groups (**Figure 3B**). We quantified anisotropies in receptive fields as the

shape index, ranging in value from -1 to 1, where positive values indicated horizontally

anisotropic shapes, negative values indicate vertically anisotropic shapes and values

close 0 indicate more isotropic shapes. Receptive fields for neurons in the S1 whisker

representation in standard-reared sighted control and early blind animals were dominated

by horizontally anisotropic shapes (**Figure 3C-D**). However, for early blind animals reared

in enriched conditions, the receptive field shape distributions were shifted away from

horizontal anisotropy – either when considering only 1° surround whisker (SW) positions

(**Figure 3C**; neurons with shape index > 0.0, eEB: 37.3%, sSC: 68.3%, sEB: 57.8%; linear

mixed-effects model, eEB vs. sSC: p<0.001, eEB vs. sEB: p<0.05) , or when including all

SW (1° - 3°) positions (**Figure 3D**; neurons with shape index > 0.0, eEB: 37.3%, sSC:

81.7%, sEB: 76.6%; linear mixed-effects model, eEB vs. sSC: p<0.001, eEB vs. sEB:

p<0.001).

Receptive field distributions were also shifted away from horizontal anisotropy in

sighted control animals reared in the enriched environment. This shift was not seen when

only 1° SW positions were considered (**Figure 3C**; neurons with shape index > 0.0, eSC:

55.2%; linear mixed-effects model, eSC vs. sSC: p>0.05, eSC vs. sEB: p>0.05) but

becomes apparent when 1° - 3° SW positions are considered (**Figure 3D**; neurons with

shape index > 0.0, eSC: 46.6%; linear mixed-effects model, eSC vs. sSC: p<0.05, eSC

vs. sEB: p>0.05). Thus, rearing in an enriched environment, which was permissive of

more naturalistic patterns of tactile behavior, shifted the shapes of receptive fields of S1

neurons away from the horizontally anisotropic shapes typically found in animals reared

in standard laboratory cages. The shift occurred in both sighted and early blind enriched

animals, but the effect was more pronounced in early blind animals.

The results of receptive field shape analyses pointed to differential effects of

rearing environment on tuning along horizontal and vertical axes of receptive fields. We

quantified neuronal tuning width separately along whisker rows and arcs for each

experimental group (**Figure 3E**). Both sighted and early blind enriched animals showed

sharper tuning along both the vertical and horizontal axes relative to sighted controls. But

there was a striking difference in the magnitude of tuning width changes along the

horizontal vs. vertical axes. This is an amplification of the shift in receptive field

anisotropies seen in early blind animals under standard rearing conditions (Ramamurthy

and Krubitzer, 2018). Along the arc (vertical) axis, tuning was sharpened to the same

extent in the enriched groups as seen in standard-reared early blind animals relative to

sighted controls (**Figure 3E**; %BW response for in-arc whiskers: eEB: 19.4 ± 2.80%, eSC:

18.7 ± 2.13%, sEB: 20.1 ± 1.58%, sSC: 26.7 ± 2.50%; linear mixed-effects model, eEB

vs. sSC: $p<0.05$, eEB vs. sEB: $p>0.05$, eSC vs. sSC: $p<0.05$, eSC vs. sEB: $p>0.05$). But

along the row (horizontal axis), both sighted and early blind enriched animals showed

sharper tuning relative to both groups of standard-reared animals (**Figure 3E**; %BW

response for in-row whiskers: eEB: 13.8 ± 2.07%, eSC: 20.9 ± 1.96%, sEB: 35.1 ± 2.82%,

sSC: 49.1 ± 2.85%; linear mixed-effects model, eEB vs. sSC: $p<0.001$, eEB vs. sEB:

$p<0.001$, eSC vs. sSC: $p<0.001$, eSC vs. sEB: $p<0.001$). There was a trend for sharper

row tuning in enriched early blind animals compared to enriched sighted control animals,

but this was not significant (linear mixed-effects model; eSC vs. eEB: $p>0.05$). We

computed the whisker selectivity index as the normalized difference ratio of the neuronal

response to the best whisker in comparison to responses to in-row or in-arc whiskers.

Population distributions of the selectivity index for S1 neurons (**Figure 3F**) further validate

the anisotropy in the enhancement of selectivity seen for animals reared in enriched

environments (BW vs. in-arc whiskers: linear mixed-effects model, eEB vs. sSC: $p<0.05$,

eEB vs. sEB: $p>0.05$, eSC vs. sSC: $p<0.05$, eSC vs. sEB: $p>0.05$, eEB vs. eSC: $p>0.05$;

BW vs. in-row whiskers: linear mixed-effects model, eEB vs. sSC: $p<0.001$, eEB vs. sEB:

$p<0.001$, eSC vs. sSC: $p<0.001$, eSC vs. sEB: $p<0.001$, eEB vs. eSC: $p>0.05$).

Thus, a spatially complex enriched environment drives significantly greater

improvement in neuronal selectivity for whisker touch in early blind animals especially

along the rostrocaudal axis, which is the primary axis of natural whisking behavior. A

similar improvement in selectivity was observed in sighted control littermates reared in

the enriched environment, supporting the role of tactile experience rather than vision loss

itself in driving these effects. In summary, the results of our current study indicate that

compensatory effects on neural coding in S1 following early blindness do not arise solely

from visual deprivation and can be amplified and directed by tactile experience.

DISCUSSION

Although the effects of enrichment on plasticity within a sensory modality have

been studied at structural, functional, and molecular levels, few studies have examined

its impact on cross-modal compensatory plasticity, focusing mainly on c-fos expression

or synaptic changes (Piche et al., 2004; Mundiñano and Martínez-Millán, 2010; Zheng et

al., 2014). Here, we tested the effects of rearing sighted and early blind short-tailed

opossums in a large and dynamic, spatially complex 3D environment, and examined

effects on on tactile behavior and receptive field configurations of neurons in S1.

Enrichment reduces abnormal stereotypic behaviors (e.g. overgrooming) and

other behavioral indicators of stress, aggression, and enhances learning, memory, motor

performance and social behavior (Dijkhuizen et al., 2024; Ratuski et al., 2024; Cait et al.,

2024; Nakhwa et al., 2024; Caires & Bossolani-Martins, 2023; Mieske et al., 2022;

Kentner et al., 2021; Khoo et al., 2020; Bailoo et al., 2018; Girbovan & Plamondon, 2013;

Amaral et al., 2008; Mason et al., 2007). It can also improve sensory function and

performance, including whisker touch discrimination (Zheng et al., 2021). In our study, we

implemented enrichment intended to increase the use of whisker touch. Since whiskers

are critical for guiding forelimb placement in crossing gaps when small mammals climb

(Ahl, 1986; Arkley et al., 2014; Niederschuh et al., 2015; Arkley et al., 2017; Grant et al.,

2018), we reared opossums in cages with periodically varying, spatially complex climbing

substrates and intermittently introduced novel toys, since novel objects and other

interactive enrichment devices promote active whisking behavior (Nakhwa et al., 2024;

Sofroniew & Svoboda, 2015). We show that in a novel setting, both sighted and blind

opossums showed differential exploration patterns reflecting the geometry of their rearing

environments. Additionally, enriched opossums are faster and more successful in gap

crossing using a whisker-dependent strategy. Thus, opossums use behavioral strategies

shaped by their rearing environments, and our enrichment approach effectively engages

and enhances naturalistic whisker-mediated behavior in both sighted and blind animals,

enabling us to evaluate the effects of enriched rearing on compensatory plasticity

following early vision loss.

We have previously shown that early blindness alters the coding of whisker touch

inputs in opossums, in a manner that supports enhanced tactile discrimination

(Ramamurthy & Krubitzer 2018, Ramamurthy et al., 2021). This involved suppression of

whisker-evoked responses combined with increased spatial selectivity, due to the greater

relative suppression of neural responses to non-preferred stimuli (SW) vs. the preferred

stimulus (BW). These alterations in the receptive fields of neurons in S1 resemble the

effects of naturalistic environmental enrichment on receptive fields of neurons in S1 of

adult rats, described by Polley and colleagues (1999, 2004). Here we demonstrate that

rearing early blind short-tailed opossums in an enriched environment further amplifies

these changes. Since selectivity to whisker stimuli was enhanced not just in early blind

animals but also in sighted control animals reared in enriched conditions, we can infer

that suppression of whisker-evoked responses and increased selectivity of neural

responses in the S1 whisker representation are effects of tactile experience rather than

visual deprivation itself.

A key finding was the effect of the rearing environment on the shapes of receptive

fields of S1 neurons – shape distributions for both sighted and early blind animals

exhibited a prominent shift away from the horizontally anisotropic shapes that dominate

in standard-reared opossums. This was accompanied by a reduction in tuning width along

row (horizontal) vs. arc (vertical) axes in enriched animals – on average, tuning widths

were reduced to a greater extent along the row axis, parallel to the main axis of whisker

movements. This may reflect more naturalistic patterns of whisker use in short-tailed

opossums, enabled by the increase in vertical space and availability of climbing

substrates in the enriched rearing environment. In rats, simulations that modelled

exploratory behavior predicted changes in the spatial patterns of whisker contact

depending on the environment being explored (Hobbs et al., 2015) – whiskers within the

same horizontal row were more likely to come in simultaneous contact during exploration

of flat surfaces, but in more complex and naturalistic environments, the probability

distributions of simultaneous whisker contact shifted away from this row-wise structure.

Such alterations in the patterns of tactile stimuli impinging on the whisker array over the

course of development may, in turn, impact receptive field structure.

Another possibility is that anisotropic effects on receptive fields are inherent to

differences in row vs. arc organization within the neocortex. In rats and mice, it is well-

documented that there is a bias towards preferential connectivity among barrels

corresponding to whiskers in the same row (Bernardo et al., 1990; Hoeflinger et al., 1995;

Kim and Ebner, 1999; Keller and Carlson, 1999), and neurons in barrel cortex typically

display patterns of activity that tend to be elongated along the row axis (Armstrong-James

and Fox, 1987; Simons, 1978; Kleinfeld and Delaney, 1996; Ego-Stengel et al., 2005; Le

Cam et al., 2011). Further, there is evidence for differential effects on responses of S1

neurons along row and arc axes when sensory inputs are altered within the whisker

system, for example, through chronic stimulation (Quairiaux et al., 2007) or selective

trimming (Dubroff et al., 2005; Wallace and Sakmann, 2008) of specific whiskers. Thus,

biases in the functional organization and connectivity within the S1 whisker representation

may favor plasticity, preferentially sharpening tuning along the row axis.

A long-standing question has been whether reorganization following sensory loss

is primarily driven by deprivation of the lost modality or due to increased experience using

the spared senses. Studies in blind humans with Braille reading experience support an

important role for tactile experience in generating tactile spatial acuity enhancement

following blindness (Wong et al., 2011). The relative contributions of deprivation of the

lost modality versus increased or modified use of the spared senses (Goldreich and

Kanics, 2003; Wong et al., 2011; Voss, 2011) to different aspects of cortical

reorganization following sensory loss are still not well understood. Our study provides

clear evidence for the role of experience in driving compensatory neural and behavioral

plasticity following sensory loss, highlighting the importance of therapeutic strategies

engaging tactile exploration (for example through play-based learning of tactile skills,

Hilditch 2024; or the use of haptic devices, Fleck et al., 2025) to maximize neural and

behavioral compensation through touch.

ACKNOWLEDGMENTS

We thank Dr. Cindy Clayton for animal care, Carly Jones for assistance with preliminary

data analysis, and Carlos Pineda for comments on a draft of this manuscript. This

research was supported by NIH NEI 5R01EY034303 and James S. McDonnell

Foundation grant 220020516 to L.A.K and NSF GRFP DGE-1650042 to D.L.R.

AUTHOR CONTRIBUTIONS

D.L.R. and L.A.K. designed the study. D.L.R., M.E., T.J.K., and H.D. performed the

experiments. D.L.R. analyzed the data. D.L.R. and L.A.K. wrote the manuscript.

DECLARATION OF INTERESTS

The authors declare no competing interests.

MATERIALS AND METHODS

Animals

Thirty-three adult short-tailed opossums (*Monodelphis domestica*; age range: 4-16

months; weight range: 61-165g) were used for electrophysiological recording

experiments. Animals belonged to four experimental groups (**Table S1**) across two

rearing paradigms (standard: 's', enriched: 'e'). Data from animals in the standard-rearing

groups were reported in a previous publication (Ramamurthy and Krubitzer, 2018),

however, new analyses of some aspects of the data are additionally presented here for

comparison with data from enriched-rearing groups.

Ethics Statement

All protocols were approved by the Institutional Animal Care and Use Committee

of the University of California, Davis, and experiments were conducted according to the

criteria outlined in the National Institutes of Health *Guide for the Care and Use of*

Laboratory Animals.

Bilateral enucleation

For all experimental groups, bilateral enucleations were performed at P4 using

procedures previously described (Ramamurthy and Krubitzer, 2018). Mothers of

experimental litters were lightly anesthetized with Alfaxan (initial dose: 20 mg/kg;

maintenance doses: 10-50%; IM) to facilitate enucleation of the pups, which are attached

to the mother at this developmental stage. Pups were anesthetized by hypothermia.

Health of both the mother (respiration rate and body temperature) and the pups

(heartbeat, coloration and mobility) was monitored throughout the procedure. An incision

was made in the skin covering the eyes, the eyes were removed under microscopic

guidance, and the skin was repositioned and resealed using surgical glue. Approximately

50% of each litter was bilaterally enucleated. After complete recovery from anesthesia,

mothers along with their attached litters were placed into either standard or enriched

housing, depending on the experimental group.

Rearing paradigms

Mothers of all experimental groups were reared and housed in standard laboratory

cages until bilateral enucleations were performed on the pups at P4. Following the

enucleation procedure, mothers with litters belonging to the standard-rearing groups were

placed in standard laboratory cages, while those belonging to the enriched-rearing groups

were placed in custom-made enriched cages (**Figure S1**). All animals were maintained

on a 14/10 light/dark cycle. Food and water were available *ad libitum* in the cage under

both rearing conditions. All experimental litters were housed together with the mother until

weaning at ~8 weeks (P56), and at that stage they were separated from the mother, but

the whole litter continued to be housed together until adulthood at ~4 months (P120).

Adults (>P120) were individually housed under standard rearing conditions, but in groups

of 2-4 same-sex littermates under enriched conditions (see below).

Standard-rearing: Under standard-rearing conditions, adult animals were singly housed

in standard laboratory cages, in cages similar to those typically used for laboratory

rodents (cage dimensions: 18.5"l x 10"w x 8"h). The home cage was provided with a

nesting cup and nesting material (shredded paper towels).

Enriched-rearing: The enriched rearing environment in our study incorporated features

common to environmental enrichment (EE) paradigms previously used in rodents

(Nithianantharajah and Hannan, 2006; Sale et al., 2009; Bengoetxea et al., 2012), and

additionally takes into consideration the natural lifestyle of short-tailed opossums (**Figure**

S1). Juvenile animals were cohoused until weaning with the mother as well as littermates,

and then with just their littermates from weaning until adulthood, as described above.

Enriched cages were provided with standard bedding and nesting material, as well as

some additional nesting material with a different texture. In addition to the nesting cup

placed on the floor as in standard cages, enriched cages provided the option of nesting

high above the cage floor, in nesting boxes attached to the cage walls. A single nesting

box was provided in the cage at juvenile stages of the experimental litters. In adulthood,

balanced numbers of eSC and eEB littermates were housed together in each enriched

cage, with a maximum of four opossums per cage. Each adult animal was provided an

individual nesting box. Enriched cages were substantially larger (~8x in volume; 29"l x

18"w x 24"h) compared to standard cages to encourage tactile exploration in all three

dimensions. The greatest increase relative to standard cages was along the vertical

dimension: enriched cages were 3x taller than standard cages. Enrichment toys (Bio-

Serv) of different shapes and textures were provided in the cage. Toys were replaced,

and the positions of manzanita branches were changed every 2–3 days to introduce

novelty and promote continual exploration of the environment. .

Behavioral testing

Arena testing: Arena testing was performed in three adult animals from each of the four

experimental groups. The arena consisted of a cage with the dimensions of enriched

cages, containing a single nest box near the top of the arena, with a branch leading up to

it. Animals were placed in the arena and allowed to explore it for 5 minutes in the

darkness, and videos of their behavior were acquired under infrared (IR) illumination.

idTracker software (**Figure S2A**; Romero-Ferrero et al., 2018) was used to extract

movement trajectories of individual animals, which was further analyzed in Matlab.

Exploratory activity for each animal was quantified as the average number of frames in

which they spent in the regions defining horizontal and vertical space within the arena,

and the vertical exploration index was calculated as:

$$Vertical\ Exploration\ Index = \frac{(Vertical\ exporation - Horizontal\ exploration)}{(Vertical\ exporation + Horizontal\ exploration)}$$

Thus, more positive values indicated a shift towards vertical exploration, while more

negative values indicated a shift towards horizontal exploration.

Gap crossing task: Gap crossing was tested in 3-4 adults animals per experimental group.

Animals were allowed to spontaneously cross from one platform ("start") to another

platform ("goal") with either small (2-10 cm) or large gap distances (12-14 cm) separating

them, which were interleaved in a pseudorandom order. Animals were tested for 7 days

with intact whiskers, followed by 1-3 days of testing after whisker trimming. On each trial

animals were allowed to remain on the start platform and attempt crossing for a maximum

of 1 minute. If no crossings were performed, the trial was ended and animals were

removed from the platform for 1 min before returning to the platform for the next trial.

Videos of this behavior were acquired under IR illumination or dim red light, and

DeepLabCut was used for analysis of gap crossings (**Figure S2B-C**). The snout tip, the

base of the tail and the position of the edges of each platform were labeled in a subset of

frames to generate the dataset (260 frames across 13 opossums) on which the network

was trained (200,000 iterations). On a subset of trials, slow-motion video clips in a

zoomed-in field of view were acquired to visually confirm the use of whiskers during gap

crossing. Infrared motion sensors (emitter/detector pairs) placed on either side of the

edge of each platform detected beam breaks and beam break times were recorded as a

secondary method of confirmation.

Surgical procedures

Animals were anesthetized with urethane (initial dose: 1.25 g/kg, 30% in saline, IP;

supplemental doses: 0.125-0.313 g/kg, 30% in saline, IP). Vital signs (respiration and

body temperature) were monitored throughout the experiment. At the beginning of the

surgery, dexamethasone (0.4-2.0 mg/kg; IM) was administered to minimize intracranial

swelling. Lidocaine (2% solution; SC) was injected down the midline of the scalp and

around the ears. The animal was then placed in a stereotaxic frame. Following an incision

at the midline of the scalp and retraction of the temporal muscle, a craniotomy was

performed to expose the entire parietal cortex and the dura was retracted. The surface of

the neocortex was covered with a layer of silicone fluid to prevent desiccation and then

photographed so that electrode penetration sites could be related to patterns of

vasculature. The head of the animal was stabilized by cementing a head post to a skull

screw inserted rostral and contralateral to the craniotomy.

Extracellular recordings and whisker stimulation

Somatosensory receptive fields were first coarsely mapped by recording multiunit

activity while applying tactile stimuli using a handheld probe. The representation of the

whiskers in S1 was localized by progressively testing electrode penetration sites medial

to the rhinarium representation and caudal to the lower jaw/lip representation. After this,

when receptive fields were found to be located on the mystacial or genal whisker pad (),

computer-controlled whisker deflections were used quantitatively measure single unit

receptive fields.

Whisker stimuli consisted of single whisker deflections (2°) delivered using

piezoelectric actuators (4 ms/100 ms/4 ms ramp-hold-return; 5 mm from the whisker

base; 1.4 mm forward excursion). Data were collected for a grid of sixteen whiskers on

the mystacial pad to enable construction of somatotopic receptive fields. Whiskers were

deflected in a pseudorandom order with 50-100 trials collected per whisker (in blocks of

3-50 trials) at each recording site. Extracellular recordings (amplifier gain: 10,000X; A-M

Systems Model 1800 Microelectrode AC Amplifier; A-M Systems, Carlsborg, WA;

sampled at 28 kHz; Power1401, Cambridge Electronic Design Limited, Cambridge, UK)

were made 400-500 μm below the pial surface using tungsten microelectrodes (FHC,

Inc., Bowdoin, ME; 1–5 MΩ at 1 kHz) lowered using a hydraulic microdrive (David Kopf

Instruments, Tujunga, CA). Recordings were bandpass filtered (300-3000 Hz), and spike-

sorting was carried out offline to confirm isolation of single units.

After electrophysiological data was collected, fluorescent probes were used to

mark the location of recording sites, so that these could be related to histologically

identified cortical field boundaries (**Figure S3**).

Histology

At the end of each experiment, animals were injected with an overdose of sodium

pentobarbital (Beuthanasia; 250 mg/kg IP) and transcardially perfused with 0.9% saline,

followed by 2-4% paraformaldehyde in phosphate buffer, and then 2-4%

paraformaldehyde in 10% phosphate-buffered sucrose. The brain was extracted and

post-fixed (1-2 hours in 4% paraformaldehyde in 10% phosphate-buffered sucrose). In

some cases, the cortical hemispheres were flattened and the tissue was sectioned

tangentially at 30 µm using a freezing microtome and processed for myelin staining

(Gallyas, 1979; Dooley et al, 2013) to enable reconstruction of cortical field boundaries.

Reconstructions were drawn using in Adobe Illustrator CS5 (Adobe, San Jose, CA).

Digital images of processed tissue were obtained using either a Nikon Multiphot system

or an Optronics MicroFire digital microscope camera. Contrast and brightness of whole

images were adjusted using Adobe Photoshop CS5 (Adobe, San Jose, CA).

Neural data analysis

Data files were processed blind to experimental group identity. Single-unit isolation

was performed while running the experiment using template-matching procedures in

Spike2 (Cambridge Electronic Design Limited, Cambridge, UK; RRID:SCR_000903), and

unit isolation was verified offline in Spike2, using principal component analysis. Stable

waveforms and firing rates over the course of the recording session, and <0.5% refractory

period violations (interspike interval <1.5 ms) were required for units to be included in the

analysis. Spikes times were then exported to MATLAB (Mathworks, Inc., Natick, MA;

RRID:SCR_001622) and analyzed using custom scripts. Whisker evoked firing rates were

measured as the spike count following stimulus onset (0-100 ms) and were considered

significant if they exceeded prestimulus spike counts in an equivalent time window by two

standard deviations. The *best whisker* (BW) was defined as the whisker with the highest

evoked response, and depending on their proximity to the BW the remaining whiskers

were categorized as *first-order surround whiskers* (1° SW; immediately adjacent to BW),

second-order surround whiskers (2° SW; one position removed from BW) and *third-order*

surround whiskers (3° SW; two positions removed from BW).

One measure of receptive field size used was the number of whiskers that evoked

a significant response in each recorded neuron. Another measure used was the rank-

ordered whisker tuning curve for each neuron, where responses to all whiskers tested

were sorted in descending order of whisker-evoked response magnitude, and normalized

to peak (BW response). Further, mean 2D somatotopic receptive fields were obtained

after normalizing SW-evoked response magnitudes to the BW-evoked response, and

aligning receptive fields of single units such that the BW was at the center of the receptive

field for each unit. 50, 75, and 90% contour lines (for response levels relative to the BW)

were plotted for the mean receptive fields (smoothed by linear interpolation) to facilitate

visualization. Raw values were used for all calculations and statistical comparisons.

Anisotropies in the structure of somatotopic receptive fields were quantified as the

shape index:

544
$$Shape\ index = \frac{(Row\ tuning\ width\ index - Arc\ tuning\ width\ index)}{(Row\ tuning\ width\ index + Arc\ tuning\ width\ index)}$$

If the sum of the row and arc tuning width indices was zero, shape index was

assigned to be zero. Shape indices ranged from -1 to 1, with values >0 indicating broader

tuning along the row compared to the arc (i.e. broader horizontal tuning), and values <0

indicating broader tuning along the arc compared to the row (i.e. broader vertical tuning).

Neuronal selectivity for the best whisker along the row and arc dimensions was

measured by assessing tuning width separately along the horizontal and vertical axes as

the *row tuning width index* and the *arc tuning width index* respectively, either considering

only 1° SW, or 1°, 2°, 3° SW in the row or arc of the BW, as stated below:

553
$$Row\ tuning\ width\ index = \frac{(\text{Mean evoked response of } SW_{row})}{\text{BW evoked response}}$$

554
$$Arc\ tuning\ width\ index = \frac{(\text{Mean evoked response of } SW_{arc})}{\text{BW evoked response}}$$

555   Additionally, the *whisker selectivity index (WSI)* for BW vs. in-row or in-arc whiskers was

556   computed separately for each neuron, considering BW responses relative to 1°, 2°, 3°

557   SW responses.

558
$$WSI_{row} = \frac{(BW\ evoked\ response - SW_{row}\ evoked\ response)}{(BW\ evoked\ response + SW_{row}\ evoked\ response)}$$

559
$$WSI_{arc} = \frac{(BW\ evoked\ response - SW_{arc}\ evoked\ response)}{(BW\ evoked\ response + SW_{arc}\ evoked\ response)}$$

In-row and in-arc selectivity indices ranged from 0 to 1, with values closer to 1

indicating higher selectivity for the BW in the corresponding dimension.

Statistical Analysis

Permutation tests for the difference in means were used to assess differences in

mean between two groups (referred to in brief as 'permutation test'). All statistical tests

for neuronal data use 'n' of cells. We account for interindividual variability using a linear

mixed effects model with the formula:

$$Response\ Variable \sim Experimental\ Group + (1|Sex) + (1|Animal\ ID)$$

where experimental group is modeled as a fixed effect, and sex and animal ID are

modeled as random effects. For gap crossing behavior, an additional model was fit to test

for fixed effects of test condition (large vs. small gaps, intact vs. trimmed whiskers):

$$Response\ Variable \sim Experimental\ Group + Test\ Condition + (1|Sex) + (1|Animal\ ID)$$

where experimental group and test condition are modeled as fixed effects, and sex and

animal ID are modeled as random effects. Exact p-values for the fixed effect of

experimental group and/or test condition in the linear mixed-effects model, and pairwise

comparisons across experimental groups are provided in **Table S2.**

FIGURE LEGENDS

Figure 1. Behavioral differences in opossums reared in enriched environments. A.

Timeline showing experimental manipulations relative to key milestones in the

development of short-tailed opossums. Behavior and electrophysiology experiments were

conducted in adults following the sequence shown in the inset, but ages were not

precisely matched to the exact day. **B.** Example movement trajectories from one animal

in each experimental group (sEB: light red, eEB: dark red, sSC: light blue, eSC: dark blue)

tracked over five minutes during behavioral testing in an enriched arena. **C.** Overlaid

movement trajectories from all animals in each experimental group in the arena test show

a pattern of increased exploration of vertical vs. horizontal space in opossums reared in

enriched cages compared to those reared in standard cages. Plus symbols indicate

center of mass of movement trajectories for each group (colors as in **B**), with data for

animals combined within in each rearing condition in black. **D.** Mean vertical exploration

index across animals. Animals reared in the enriched environement show significantly

higher levels of vertical exploration. **E.** Experimental setup in the gap crossing task. **F.**

Number of gap crossings per minute (across all trials) for large and small gaps (with intact

or trimmed whiskers). Gap crossing was whisker-dependent in all groups. No opossums

successfully crossed large gaps after whisker trimming, so this not plotted. **G.** Duration

for successfully crossing large or small gaps (with intact or trimmed whiskers). All groups

show significantly longer durations for succesful gap crossing on large gaps (12-14 cm)

vs. small gaps (2-10 cm). Upon whisker trimming, all groups show a significant increase

in gap crossing duration for small gaps, consistent with the use of whisker-dependent

strategies for gap crossing, as seen in **(F)**. For larger gaps, there are prominent

differences between experimental groups based on rearing condition and blindness. Early

blind animals crossed gaps faster than sighted controls in standard conditions, and their

performance was further enhanced by enriched rearing.

Figure 2. Enriched rearing amplifies alterations in S1 neuron responses to whisker

touch in early blind animals, while also producing similar shifts in sighted controls.

A. Spike raster plots for example neurons responsive to whisker touch in each of the four experimental groups (left to right, sSC, sEB, eSC, eEB). Examples are shown for neurons with the same best whisker (BW), namely C2. Trials corresponding to A, B, C and D whisker rows are separated by horizontal lines. Within each row the direction of more rostral 'R' vs. more caudal 'C' whiskers is labeled on the y-axis.**B.** Peristimulus time histograms (PSTHs) corresponding to the same neurons shown in **(A)**. PSTHs corresponding to whiskers with response magnitudes >50% the BW response are in black, others are in gray**.** The examples in **(A)** and **(B)** illustrate more selective responses to whisker touch in standard-reared blind opossums, which is further amplified in enriched groups. Additionally, neural responses to whiskers vertically adjacent to the BW are stronger than those for horizontally adjacent whiskers in enriched groups, especially in enriched early blind opossums. **C.** Spontaneous firing rates of S1 neurons were not significantly different across experimental groups. **D.** Firing rates evoked by the BW showed significant differences across experimental groups. Notably, both sighted and early blind enriched groups showed suppression relative to standard-reared sighted controls, similar to that which we previously described in standard-reared early blind animals. The magnitude of suppression of the BW-evoked response was greatest in early blind animals reared in the enriched environment. **E.** Mean receptive field size was significantly smaller in sEB, eSC and eEB animals relative to standard-reared sighted controls. This is seen both in terms of number of whiskers driving significant responses (left) and the rank-ordered tuning curve (right) for whisker-responsive S1 neurons. See also **Table S2**.

Figure 3. Enriched rearing amplifies anisotropies in receptive fields of S1 neurons in early blind animals, improving selectivity for whisker touch along a behaviorally relevant axis. A. Schematic diagram of the opossum mystacial whisker pad indicating whisker rows (light yellow) and arcs (gold), and the hortical and vertical axes used for shape analyses. The schematic shown as an inset in individual panels depicts the grid of 16 whiskers and the surrounding whisker (SW) positions considered in that particular analysis for one example BW position (black filled circle) on the grid. **B.** Mean 2D somatotopic receptive fields for each experimental group (left to right, sSC, sEB, eSC, eEB) with 50, 75, and 90% contour lines (for response levels relative to the BW, smoothed by linear interpolation for display). **C-D.** Receptive field shape distributions for S1 neurons in enriched animals were shifted away from horizontal anisotropic shapes, which are typical of standard-reared animals. **C** shows receptive field shape distributions when only 1° SW positions were considered (left) and fractions of horizontally anisotropic receptive fields corresponding to each group (right). Light gray bars indicate horizontally elongated receptive fields defined as shape indices > 0, while black bars indicate horizontally elongated receptive fields with a stricter cutoff of shape indices > 0.5. **D** shows data for the same neurons when all SW positions (1°-3°) were included. Horizontally anisotropic receptive fields are decreased in both enriched groups of animals, and to the greatest extent in enriched early blind animals. **E.** Quantification of tuning width along the row (horizontal/rostrocaudal axis) and arc (vertical axis) across experimental groups. Enriched groups show sharper tuning along the horizontal axis, to a significantly greater extent than standard-reared early blind animals. Enriched groups also show sharper tuning along the vertical axis compared with standard-reared sighted controls, but only to

the same extent as previously observed in standard-reared early blind animals. **F.**

Distributions of the whisker selectivity index of S1 neurons calculated separately for in-

arc whiskers (left) and in-row whiskers (right), when including all SW positions (1°- 3°).

sEB, eSC and eEB groups all showed significantly greater selectivity for whiskers within

arcs, but were not significantly different from each other. The greatest increases in

whisker response selectivity were seen for whiskers within rows i.e. the

horizontal/rostrocaudal axis, which is the primary axis of natural whisker motion.

Selectivity was greatest for eEB animals followed by eSC and sEB animals, with all three

of these groups showing significantly higher selectivity along the horizontal axis comapred

to sSC animals.

SUPPLEMENTARY FIGURE LEGENDS

Figure S1. Standard and enriched rearing environment used for short-tailed

opossums. A. Comparison of cages used in standard rearing (L) and enriched rearing

paradigms (R). Enriched cages (29"l x 18"w x 24"h) were much larger (~8x in volume)

than standard cages (8.5"l x 10"w x 8"h), especially in the vertical dimension (3x), and

provided with a nesting box high above the ground, in addition to the standard nesting

cup placed on the cage floor. They contained various enrichment objects: enrichment toys

that were regularly switched out and manzanita branches that were regularly switched in

position. They were also provided with a running wheel, and incoporated social housing

(see *Materials and Methods*). **B.** Frontal view of an enriched cage. **C.** Top view of the

same enriched cage as **(B)**. Letters indicate different components of the cage, which are

visible in both views: nesting box (a), nesting cup (b), food (c), water (d), running wheel

(e), manzanita branches (f-l), and enrichment toys (m). **D.** Mother with an attached

experimental litter after placement in the enriched rearing environment. **E-F.** Early blind

weanling climbing up a branch to reach the nesting box .

Figure S2. Behavioral video analyses. A. One frame showing semi-automated

identification of an opossum during arena testing using idtracker.ai. **B.** One frame

showing behavioral tracking in the gap crossing task using DeepLabCut. 'IR1' and 'IR2'

indicate positions of infrared motion sensors mounted on the edge of each platform,

tracked using DeepLabCut to aid in gap crossing analyses. **C.** A zoomed in view showing

an opossum on the start platform, extending its whiskers towards the goal platform prior

to crossing the gap.

Figure S3. **Histological verification of recording sites. A-D.** Myelin stain of a tangential

section of the cortex in a standard-reared sighted (sSC) animal (**A**), a standard-reared

early blind (sEB) animal (**C**), an enriched sighted (eSC) animal (**E**), and an enriched early

blind (eEB) animal (**G**). S1 is readily identified because it is darkly staining compared to

adjacent cortical fields. Reconstructions of myeloarchitectural borders drawn from the

series of myelin-stained sections in sSC (**B**), sEB (**D**), eSC (**F**) and eEB (**G**) animals.   In

B, **D**, **F** and **G** the location of fluorescent probes is indicated by open circles, and the

location of recording sites is indicated by black filled circles. Only recording sites at which

whisker-evoked responses of neurons were quantified are shown here. In **A** – **H**, medial

(M) and rostral (R) directions are indicated by arrows, and scale bar = 250 µm.

Video S1. A nesting mother opossum navigating the enriched home cage environment.

Video shown at 4x speed.

Video S2. Sighted opossum in the arena test. Video shown at 4x speed.

Video S3. Early blind opossum in the arena test. Video shown at 4x speed.

REFERENCES

Abreu MSL, Oliveira LRD (2014) Patterns of arboreal and terrestrial space use by non-

volant small mammals in an Araucaria forest of southern Brazil. Anais da

Academia Brasileira de Ciências 86:807-819.

Ahl A (1986) The role of vibrissae in behavior: a status review. Veterinary research

communications 10:245-268.

Alwis DS, Rajan R (2013) Environmental enrichment causes a global potentiation of

neuronal responses across stimulus complexity and lamina of sensory cortex.

Frontiers in cellular neuroscience 7:124.

Alwis DS, Rajan R (2014) Environmental enrichment and the sensory brain: the role of

enrichment in remediating brain injury. Frontiers in systems neuroscience 8:156.

Antón-Bolaños N, Sempere-Ferràndez A, Guillamón-Vivancos T, Martini FJ, Pérez-Saiz

L, Gezelius H, Filipchuk A, Valdeolmillos M, López-Bendito G (2019) Prenatal

activity from thalamic neurons governs the emergence of functional cortical maps

in mice. Science 364:987-990.

Amaral OB, Vargas RS, Hansel G, Izquierdo I, Souza DO (2008) Duration of

environmental enrichment influences the magnitude and persistence of its

behavioral effects on mice. Physiology & Behavior 93:388-394.

Arkley K, Grant RA, Mitchinson B, Prescott TJ (2014) Strategy change in vibrissal active

sensing during rat locomotion. Current biology : CB 24:1507-1512.

Arkley K, Tiktak GP, Breakell V, Prescott TJ, Grant RA (2017) Whisker touch guides

canopy exploration in a nocturnal, arboreal rodent, the Hazel dormouse

(Muscardinus avellanarius). Journal of comparative physiology A, Neuroethology,

sensory, neural, and behavioral physiology 203:133-142.

Bailoo JD, Murphy E, Boada-Saña M, Varholick JA, Hintze S, Baussière C, Hahn KC,

Göpfert C, Palme R, Voelkl B, Würbel H (2018) Effects of Cage Enrichment on

Behavior, Welfare and Outcome Variability in Female Mice. Front Behav Neurosci

12:232.

Bartoletti A, Medini P, Berardi N, Maffei L (2004) Environmental enrichment prevents

effects of dark-rearing in the rat visual cortex. Nature neuroscience 7:215-216.

Beddard FE (1902) Observations upon the Carpal Vibrissæ in Mammals. Proceedings of

the Zoological Society of London 72:127-136.

Bengoetxea H, Ortuzar N, Bulnes S, Rico-Barrio I, Lafuente JV, Argandona EG (2012)

Enriched and deprived sensory experience induces structural changes and rewires

connectivity during the postnatal development of the brain. Neural plasticity 2012.

Branchi I, D'Andrea I, Santarelli S, Bonsignore LT, Alleva E (2011) The richness of social

stimuli shapes developmental trajectories: Are laboratory mouse pups

impoverished? Progress in neuro-psychopharmacology & biological psychiatry

35:1452-1460.

Caires CRS, Bossolani-Martins AL (2023) Which form of environmental enrichment is

most effective in rodent models of autism? Behavioural Processes 211:1-8.

Cait J, Winder CB, Mason GJ (2024) How much 'enrichment' is enough for laboratory

rodents? A systematic review and meta-analysis re-assessing the impact of well-

resourced cages on morbidity and mortality. Applied Animal Behaviour Science

278:106361.

Cancedda L, Putignano E, Sale A, Viegi A, Berardi N, Maffei L (2004) Acceleration of visual system development by environmental enrichment. The Journal of neuroscience : the official journal of the Society for Neuroscience 24:4840-4848.

Coq JO, Xerri C (1998) Environmental enrichment alters organizational features of the forepaw representation in the primary somatosensory cortex of adult rats. Experimental brain research 121:191-204.

Delciellos AC, Vieira MV (2006) Arboreal walking performance in seven didelphid marsupials as an aspect of their fundamental niche. Austral Ecology 31:449-457.

Devonshire IM, Dommett EJ, Grandy TH, Halliday AC, Greenfield SA (2010) Environmental enrichment differentially modifies specific components of sensory-evoked activity in rat barrel cortex as revealed by simultaneous electrophysiological recordings and optical imaging in vivo. Neuroscience 170:662-669.

Diamond ME, Toso A (2023) Tactile cognition in rodents. Neuroscience & Biobehavioral Reviews 149:105161.

Dijkhuizen S, Van Ginneken LMC, Ijpelaar AHC, Koekkoek SKE, De Zeeuw CI, Boele HJ (2024) Impact of enriched environment on motor performance and learning in mice. Scientific Reports 14:5962.

Dooley JC, Franca JG, Seelke AM, Cooke DF, Krubitzer LA (2013) A connection to the past: *Monodelphis domestica* provides insight into the organization and connectivity of the brains of early mammals. J Comp Neurol 521:3877-3897.

Dooley JC, Krubitzer LA (2019) Alterations in cortical and thalamic connections of somatosensory cortex following early loss of vision. J Comp Neurol 527(10):1675-

1688.

Engineer ND, Percaccio CR, Pandya PK, Moucha R, Rathbun DL, Kilgard MP (2004)

Environmental Enrichment Improves Response Strength, Threshold, Selectivity,

and Latency of Auditory Cortex Neurons. Journal of Neurophysiology 92:73-82.

Fleck JJ, Zook ZA, Clark JP, Preston DJ, Lipomi DJ, Pacchierotti C, O'Malley MK (2025)

Wearable multi-sensory haptic devices. Nature Reviews Bioengineering.

Frostig RD (2006) Functional organization and plasticity in the adult rat barrel cortex:

moving out-of-the-box. Current opinion in neurobiology 16:445-450.

Gallyas F (1979) Silver staining of myelin by means of physical development.

Neurological research 1:203-209.

Girbovan C, Plamondon H (2013) Environmental enrichment in female rodents:

considerations in the effects on behavior and biochemical markers. Behav Brain

Res 253:178-190.

Goin FJ, Woodburne MO, Zimicz AN, Martin GM, Chornogubsky L (2016) South

American Living Metatherians: Physiological Ecology and Constraints. In: A Brief

History of South American Metatherians: Evolutionary Contexts and

Intercontinental Dispersals, pp 37-75. Dordrecht: Springer Netherlands.

Goldreich D, Kanics IM (2003) Tactile acuity is enhanced in blindness. J Neurosci

23:3439-3445.

Gomez-Pinilla F, Ying Z, Agoncillo T, Frostig R (2011) The influence of naturalistic

experience on plasticity markers in somatosensory cortex and hippocampus:

effects of whisker use. Brain research 1388:39-47.

Graham M, Socha JJ (2020) Going the distance: The biomechanics of gap-crossing
786   behaviors. J Exp Zool A Ecol Integr Physiol 333:60-73.

787 Grant RA, Breakell V, Prescott TJ (2018) Whisker touch sensing guides locomotion in
788   small, quadrupedal mammals. Proceedings Biological sciences 285.

789 Grant RA, Delaunay MG, Haidarliu S (2017) Mystacial whisker layout and musculature in
790   the guinea pig (Cavia porcellus): a social, diurnal mammal. The Anatomical Record
791   300:527-536.

792 Greifzu F, Pielecka-Fortuna J, Kalogeraki E, Krempler K, Favaro PD, Schluter OM, Lowel
793   S (2014) Environmental enrichment extends ocular dominance plasticity into
794   adulthood and protects from stroke-induced impairments of plasticity. Proceedings
795   of the National Academy of Sciences of the United States of America 111:1150-
796   1155.

797 Guic E, Carrasco X, Rodriguez E, Robles I, Merzenich MM (2008) Plasticity in primary
798   somatosensory cortex resulting from environmentally enriched stimulation and
799   sensory discrimination training. Biological research 41:425-437.

800 Hilditch J (2021) Pre-braille Implementation into Early Education: Tactile Activities to
801   Introduce Braille Concepts to Kindergartners. In: Handbook of Disability: Critical
802   Thought and Social Change in a Globalizing World (Rioux MH, Buettgen A, Zubrow
803   E, Viera J, eds), pp 1-31. Singapore: Springer Nature Singapore.

804 Hutson K, Masterton R (1986) The sensory contribution of a single vibrissa's cortical
805   barrel. Journal of neurophysiology 56:1196-1223.

Kentner AC, Speno AV, Doucette J, Roderick RC (2021) The Contribution of
Environmental Enrichment to Phenotypic Variation in Mice and Rats. eneuro
8:ENEURO.0539-0520.2021.

Khoo SY-S, Correia V, Uhrig A (2020) Nesting material enrichment reduces severity of
overgrooming-related self-injury in individually housed rats. Laboratory Animals
54:546-558.

Mason G, Clubb R, Latham N, Vickery S (2007) Why and how should we use
environmental enrichment to tackle stereotypic behaviour? Applied Animal
Behaviour Science 102:163-188.

Mieske P, Hobbiesiefken U, Fischer-Tenhagen C, Heinl C, Hohlbaum K, Kahnau P, Meier
J, Wilzopolski J, Butzke D, Rudeck J, Lewejohann L, Diederich K (2022) Bored at
home?—A systematic review on the effect of environmental enrichment on the
welfare of laboratory rats and mice. Frontiers in Veterinary Science 9.

Nakhwa AA, Breakell V, Chambers L, Elder A, McLoughlin E, Roberts S, Todd K, Grant
RA (2024) Implications of tactile enrichment on the behaviour and whisker
movements of four species of carnivorans. Applied Animal Behaviour Science
274:106261.

Jenkins Jr., FA (1974) Tree shrew locomotion and the origins of primate arborealism. In:
Jenkins, F.A. (Ed.), Primate Locomotion. Academic Press, New York, pp. 85–115.

Jiang C, Xu X, Yu L, Xu J, Zhang J (2015) Environmental enrichment rescues the
degraded auditory temporal resolution of cortical neurons induced by early noise
exposure. The European journal of neuroscience 42:2144-2154.

Jones M, Archer M, Dickman C (2003) Predators with Pouches: The Biology of

Carnivorous Marsupials: CSIRO PUBLISHING.

Kalogeraki E, Pielecka-Fortuna J, Lowel S (2017) Environmental enrichment accelerates

ocular dominance plasticity in mouse visual cortex whereas transfer to standard

cages resulted in a rapid loss of increased plasticity. PloS one 12:e0186999.

Ladine TA, Kissell RE (1994) Escape Behavior of Virginia Opossums. The American

Midland Naturalist 132:234-238.

Landers MS, Knott GW, Lipp HP, Poletaeva I, Welker E (2011) Synapse formation in adult

barrel cortex following naturalistic environmental enrichment. Neuroscience

199:143-152.

Lammers AR (2004) The biodynamics of arboreal locomotion in the gray short-tailed

opossum (*Monodelphis domestica*).

Lammers AR (2007) Locomotor kinetics on sloped arboreal and terrestrial substrates in

a small quadrupedal mammal. Zoology (Jena, Germany) 110:93-103.

Lammers AR (2009) The effects of substrate texture on the mechanics of quadrupedal

arboreal locomotion in the gray short-tailed opossum (*Monodelphis domestica*).

Journal of experimental zoology Part A, Ecological genetics and physiology

311:813-823.

Lammers AR, Biknevicius AR (2004) The biodynamics of arboreal locomotion: the effects

of substrate diameter on locomotor kinetics in the gray short-tailed opossum

(*Monodelphis domestica*). The Journal of experimental biology 207:4325-4336.

Lammers AR, Gauntner T (2008) Mechanics of torque generation during quadrupedal

arboreal locomotion. Journal of Biomechanics 41:2388-2395.

Lammers AR, Earls KD, Biknevicius AR (2006) Locomotor kinetics and kinematics on

inclines and declines in the gray short-tailed opossum *Monodelphis domestica*.

The Journal of experimental biology 209:4154-4166.

Lazzouni L, Lepore F (2014) Compensatory plasticity: time matters. Frontiers in human

neuroscience 8:340.

Leah J, Allardyce H, Cummins R (1985) Evoked cortical potential correlates of rearing

environment in rats. Biological Psychology 20:21-29.

Lee H-K (2023) Metaplasticity framework for cross-modal synaptic plasticity in adults.

Frontiers in Synaptic Neuroscience 14.

LeMessurier AM, Laboy-Juárez KJ, McClain K, Chen S, Nguyen T, Feldman DE (2019)

Enrichment drives emergence of functional columns and improves sensory coding

in the whisker map in L2/3 of mouse S1. eLife 8:e46321.

Levine JN, Chen H, Gu Y, Cang J (2017) Environmental Enrichment Rescues Binocular

Matching of Orientation Preference in the Mouse Visual Cortex. The Journal of

neuroscience : the official journal of the Society for Neuroscience 37:5822-5833.

López-Bendito G, Aníbal-Martínez M, Martini FJ (2022) Cross-Modal Plasticity in Brains

Deprived of Visual Input Before Vision. Annu Rev Neurosci 45:471-489.

Lyne A (1958) A thesis on the vibrissae and pelage hairs on the marsupialia and

supporting papers.

Lyne A (1959) The systematic and adaptive significance of the vibrissae in the

Marsupialia. Journal of Zoology 133:79-133.

Montgomery, WI (1980) The use of arboreal runways by the woodland rodents,

Apodemus sylvaticus (L.), A. flavicollis (Melchior) and Clethrionomys glareolus

(Schreber). Mammal Review 10:189-195.

Macrini TE (2004) *Monodelphis domestica*. Mammalian Species:1-8.

Megevand P, Troncoso E, Quairiaux C, Muller D, Michel CM, Kiss JZ (2009) Long-term

plasticity in mouse sensorimotor circuits after rhythmic whisker stimulation. The

Journal of neuroscience : the official journal of the Society for Neuroscience

29:5326-5335.

Mezzera C, Lopez-Bendito G (2016) Cross-modal plasticity in sensory deprived animal

models: From the thalamocortical development point of view. Journal of chemical

neuroanatomy 75:32-40.

Morland AB, Brown HD & Baseler HA (2021) Cortical reorganization: reallocated

responses without rewiring. Current Biology, 31(2), R76-R78.

Muchlinski MN, Durham EL, Smith TD, Burrows AM (2013) Comparative histomorphology

of intrinsic vibrissa musculature among primates: implications for the evolution of

sensory ecology and "face touch". American journal of physical anthropology

150:301-312.

Mundiñano IC, Martínez-Millán L (2010) Somatosensory cross-modal plasticity in the

superior colliculus of visually deafferented rats. Neuroscience 165:1457-1470.

Nakashima AS, Dyck RH (2008) Enhanced plasticity in zincergic, cortical circuits after

exposure to enriched environments. The Journal of neuroscience : the official

journal of the Society for Neuroscience 28:13995-13999.

Niederschuh SJ, Witte H, Schmidt M (2015) The role of vibrissal sensing in forelimb

position control during travelling locomotion in the rat (Rattus norvegicus,

Rodentia). Zoology (Jena, Germany) 118:51-62.

Nithianantharajah J, Hannan AJ (2006) Enriched environments, experience-dependent
plasticity and disorders of the nervous system. Nature reviews Neuroscience
7:697-709.

Piche M, Robert S, Miceli D, Bronchti G (2004) Environmental enrichment enhances
auditory takeover of the occipital cortex in anophthalmic mice. The European
journal of neuroscience 20:3463-3472.

Pocock R (1914) 48. On the Facial Vibrissæ of Mammalia. Journal of Zoology 84:889-
912.

Polley DB, Chen-Bee CH, Frostig RD (1999) Two directions of plasticity in the sensory-
deprived adult cortex. Neuron 24:623-637.

Polley DB, Kvasnak E, Frostig RD (2004) Naturalistic experience transforms sensory
maps in the adult cortex of caged animals. Nature 429:67-71.

Ramamurthy DL, & Krubitzer LA (2016) The evolution of whisker-mediated
somatosensation in mammals: Sensory processing in barrelless S1 cortex of a
marsupial, *Monodelphis domestica*. Journal of Comparative Neurology 524(17),
3587-3613.

Ramamurthy DL, Krubitzer LA (2018) Neural Coding of Whisker-Mediated Touch in
Primary Somatosensory Cortex Is Altered Following Early Blindness. The Journal
of Neuroscience 38:6172-6189.

Ramamurthy DL, Dodson HK, Krubitzer LA (2021) Developmental plasticity of texture
discrimination following early vision loss in the marsupial *Monodelphis domestica*.
Journal of Experimental Biology, 224(9), jeb236646.

Ratuski AS, Améndola L, Makowska IJ, Weary DM (2024) Effects of temporary access to
environmental enrichment on measures of laboratory mouse welfare. Sci Rep
14:15143.

Romero-Ferrero F, Bergomi MG, Hinz RC, Heras FJH, de Polavieja GG (2019)
idtracker.ai: tracking all individuals in small or large collectives of unmarked
animals. Nature Methods 16:179-182.

Rupert JE, Schmidt EC, Moreira-Soto A, Herrera BR, Vandeberg JL, Butcher MT (2014)
Myosin isoform expression in the prehensile tails of didelphid marsupials:
functional differences between arboreal and terrestrial opossums. Anatomical
record (Hoboken, NJ : 2007) 297:1364-1376.

Sale A, Vetencourt JFM, Medini P, Cenni MC, Baroncelli L, De Pasquale R, Maffei L
(2007) Environmental enrichment in adulthood promotes amblyopia recovery
through a reduction of intracortical inhibition. Nature neuroscience 10:679.

Sale A, Berardi N, Maffei L (2009) Enrich the environment to empower the brain. Trends
in neurosciences 32:233-239.

Schmitt D, Gruss LT, Lemelin P (2010) Brief communication: forelimb compliance in
arboreal and terrestrial opossums. American journal of physical anthropology
141:142-146.

Seo ML (1992) Effect of environmental complexity on the latency of cortical vibrissa
potentials. Developmental psychobiology 25:67-76.

Shapiro LJ, Young JW, VandeBerg JL (2014) Body size and the small branch niche: Using
marsupial ontogeny to model primate locomotor evolution. Journal of Human
Evolution 68:14-31.

Sofroniew NJ, Svoboda K (2015) Whisking. Current Biology 25:R137-R140.

Teichert M, Isstas M, Wenig S, Setz C, Lehmann K, Bolz J (2018) Cross-modal refinement

of visual performance after brief somatosensory deprivation in adult mice.

European Journal of Neuroscience 47:184-191.

Thomas DR, Chadwell BA, Walker GR, Budde JE, VandeBerg JL, Butcher MT (2017)

Ontogeny of myosin isoform expression and prehensile function in the tail of the

gray short-tailed opossum (*Monodelphis domestica*). Journal of applied physiology

(Bethesda, Md : 1985) 123:513-525.

van Praag H, Kempermann G, Gage FH (2000) Neural consequences of environmental

enrichment. Nature reviews Neuroscience 1:191-198.

Voss P (2011) Superior Tactile Abilities in the Blind: Is Blindness Required? The Journal

of Neuroscience 31:11745-11747.

Voss P (2019) Brain (re)organization following visual loss. WIREs Cognitive Science

10:e1468.

Wang BS, Feng L, Liu M, Liu X, Cang J (2013) Environmental enrichment rescues

binocular matching of orientation preference in mice that have a precocious critical

period. Neuron 80:198-209.

Wang M, Yu X (2023) Experience-dependent structural plasticity of pyramidal neurons in

the developing sensory cortices. Current Opinion in Neurobiology 81:102724.

Weissbrod A, Shapiro A, Vasserman G, Edry L, Dayan M, Yitzhaky A, Hertzberg L,

Feinerman O, Kimchi T (2013) Automated long-term tracking and social

behavioural phenotyping of animal colonies within a semi-natural environment.

Nature communications 4:2018.

Wilkinson M, Stirton C, McConnachie A (2010) Behavioural observations of singly-housed
grey short-tailed opossums (*Monodelphis domestica*) in standard and enriched
environments. Laboratory animals 44:364-369.

Wong M, Gnanakumaran V, Goldreich D (2011) Tactile spatial acuity enhancement in
blindness: evidence for experience-dependent mechanisms. J Neurosci 31:7028-
7037.

Zheng HJV, Meagher JP, Xu D, Patel YA, O'Connor DH, Kwon HB (2021) Environmental
Enrichment Sharpens Sensory Acuity by Enhancing Information Coding in Barrel
Cortex and Premotor Cortex. eNeuro 8.

Zheng J-J, Li S-J, Zhang X-D, Miao W-Y, Zhang D, Yao H, Yu X (2014) Oxytocin mediates
early experience–dependent cross-modal plasticity in the sensory cortices. Nature
neuroscience 17:391.

Zheng HJV, Meagher JP, Xu D, Patel YA, O'Connor DH, Kwon H-B (2021) Environmental
Enrichment Sharpens Sensory Acuity by Enhancing Information Coding in Barrel
Cortex and Premotor Cortex. eneuro 8:ENEURO.0309-0320.2021.

Zhu X, Wang F, Hu H, Sun X, Kilgard MP, Merzenich MM, Zhou X (2014) Environmental
acoustic enrichment promotes recovery from developmentally degraded auditory
cortical processing. The Journal of neuroscience : the official journal of the Society
for Neuroscience 34:5406-5415.

Figure 1

A
Arena test | Gap crossing *intact* *trim* | *whisker regrowth* | Extracellular recordings

bilateral enucleation   *weaning*   *behavior & electrophysiology*

RGC axons & TCAs reach targets

eye opening

postnatal day 0 4   56   80   120   180

whiskers emerge   early whisker movements   whisking onset   adulthood

pups attached to mother locomotor onset   active independent locomotion

cage changes 1x per week   *sSC*
rearing environment { *standard*   *sEB*
toy changes 2x per week, cage changes 1x per week   *eSC*
enriched   *eEB*

B Example trajectories

sEB eEB   sSC eSC

C Overlaid average trajectories

vertical space
sEB   sSC   eSC
eEB
horizontal space
standard   enriched

D
Vertical Exploration Index
×10⁻³
***
standard   enriched

E
start   gap   goal

F
gap crossings/minute
sSC
sEB
eSC
eEB
12-14 cm   2-10 cm   2-10 cm *trim*

G
successful gap crossing duration (s)
***
***
***
*
*** *
y-axis truncated
***
*
***
*** ***
***
*** ***
n.s.
*** *
12-14 cm 2-10 cm 2-10 cm *trim*   12-14 cm 2-10 cm 2-10 cm *trim*   12-14 cm 2-10 cm 2-10 cm *trim*   12-14 cm 2-10 cm 2-10 cm *trim*

Figure 2

Figure 3

A

B Mean receptive field

sSC   sEB   eSC   eEB

C

shape index

fraction of cells

1° SW

**

sSC — eSC
sEB — eEB

% horizontal

sSC sEB eSC eEB

n.s.
n.s.
***

D

shape index

fraction of cells

1°-3° SW

***

% horizontal

sSC sEB eSC eEB

n.s.
***
***
***
n.s.
n.s.

E

in-arc whiskers   in-row whiskers

% BW response

sSC sEB eSC eEB   sSC sEB eSC eEB

n.s.
*
n.s.

***
***
n.s.

F BW vs. in-arc whiskers   BW vs. in-row whiskers

fraction of cells

whisker selectivity index   whisker selectivity index

*   ***

Figure S1

Figure S2

A

Blob area thresholds: 2715 to 26567 px
1 blob detected of area 16840 px

B

IR1

goal

snout tip

gap

IR2

start

tail
base

C

goal

snout tip

mystacial
whiskers

genal
whiskers

gap

start

Figure S3

Table S1. Experimental Design.

		Visual Condition					
		Sighted (SC)			Early Blind (EB)		
		Sex	ALL	>2 SD	Sex	ALL	>2 SD
Rearing Condition	**Standard (s)**	5F, 5M	86	60	4F, 5M	88	64
	Enriched (e)	3F, 4M	84	58	3F, 4M	81	51

Breakdown of subject information for each experimental group. Sexes of animals used for electrophysiological recording experiments, total number of neurons recorded in S1 per group (ALL) and number of S1 neurons with significant whisker-evoked responses (>2 SD) are listed.

Table S2. Statistical comparisons for key results

Result/ Figure Panel	Fixed Effect p-value	Pairwise comparisons p-values			
Gap crossing duration **Fig. 1G**	Test Condition	sSC	sEB	eSC	eEB
(1) large vs. small	2.77e-03	1e-04	1e-04	1e-04	1e-04
(2) intact vs. trim	0.014	0.039	1e-04	1e-04	0.033

Result/ Figure Panel	Expt. Group	sSC vs.sEB	sSC vs. eSC	sSC vs. eEB	sEB vs. eEB	sEB vs. eSC	eSC vs. eEB
Gap crossing duration **Fig. 1G** (3) across groups	1.15e-03	0.014	1e-04	2.0e-04	0.012	1e-04	0.066
Spontaneous activity **Fig. 2C**	0.841	-	-	-	-	-	-
Whisker-evoked activity **Fig. 2D**	7.16e-10	1.77e-04	0.005	1.86e-11	1.57e-04	0.550	6.90e-05
RF size: # of whiskers **Fig. 2E**	1.58e-07	9.43e-07	4.29e-06	8.86e-07	0.763	0.832	0.621
Rank-ordered whisker tuning curve **Fig. 2F**	2.68e-17	2.09e-09	5.63e-13	1.74e-15	0.006	0.048	0.532
Shape index: 1^{st} order adjacent whiskers **Fig. 3C**	0.004	0.249	0.445	6.84e-04	0.023	0.628	0.003
% horizontal: 1^{st} order adjacent whiskers **Fig. 3D** Alpha with Bonferroni correction = 0.008	-	0.267	0.185	0.001	0.039	0.855	0.083
Shape index: $1^{st} – 3^{rd}$ order adjacent whiskers **Fig. 3E**	3.45e-05	0.093	0.013	2.19e-06	0.001	0.384	0.016
% horizontal: $1^{st} – 3^{rd}$ order adjacent whiskers **Fig. 3F** Alpha with Bonferroni correction = 0.008	-	0.516	9.88e-05	2.00e-06	2.50e-05	7.73e-04	0.340
% BW response: in-arc whiskers **Fig. 3G**	0.042	0.033	0.012	0.024	0.812	0.652	0.848
% BW response: in-row whiskers **Fig. 3G**	8.51e-20	7.10e-05	7.59e-14	1.12e-18	1.25e-08	6.19e-05	0.06
BW vs. in-arc whiskers **Fig. 3H**	0.043	0.08	0.014	0.015	0.419	0.436	0.957
BW vs. in-row whiskers **Fig. 3H**	2.49e-18	2.86e-04	8.18e-12	7.00e-18	1.25e-08	6.19e-05	0.06

Exact p-values for the fixed effect of experimental group and/or test condition in the linear mixed-effects model

(see Methods), and pairwise comparisons across experimental groups.