1 **Complex three-dimensional rearing environments amplify compensatory**

2 **plasticity following early blindness**

3 **Abbreviated title**: Enrichment shapes plasticity after early blindness

4 Deepa L. Ramamurthy[1, 2,] *, Mackenzie Englund[1], Tanner J. Kovacs[1,3], Heather Dodson[1],

5 Leah A. Krubitzer[1].

6 [1]Center for Neuroscience, University of California, Davis, Davis CA 95618

7 [2]Current Affiliation: Department of Psychology, 900 University Ave, Riverside, CA 92521

8 [3]Current Affiliation: University of Nebraska Medical Center, 42nd and Emile Streets

9 Omaha, NE 68198

10 *Corresponding Author**: Deepa L. Ramamurthy, deepar@ucr.edu

11 **Author contributions**: D.L.R. and L.A.K. designed the study. D.L.R., M.E., T.J.K., and

12 H.D. performed the experiments. D.L.R. analyzed the data. D.L.R. and L.A.K. wrote the

13 manuscript.

14 **Number of Figures**: 5, **Tables**: 3, **Extended Data Figures**: 3, **Extended Data Tables**:

15 3

16 **Number of Words for Abstract**: 249, **Introduction**: 750, and **Discussion**: 1059

17 **Funding Sources**: This research was supported by NIH NEI 5R01EY034303 and James

18 S. McDonnell Foundation grant 220020516 to L.A.K and NSF GRFP DGE-1650042 to

19 D.L.R.

20 **Conflicts of interest**: None

21 **Acknowledgments**: We thank Dr. Cindy Clayton for animal care, Carly Jones for

22 assistance with preliminary data analysis, and Carlos Pineda for comments on a draft of

23 this manuscript.

ABSTRACT

The neocortex has a remarkable capacity to alter its functional organization and connectivity in response to sensory loss, particularly if this loss occurs early in life. A key question is whether this cross-modal reorganization is driven by sensory deprivation or by enhanced use of the spared senses. We investigated how different rearing environments shape neural responses in primary somatosensory cortex (S1) of short-tailed opossums (*Monodelphis domestica)*, following elimination of visual inputs through bilateral enucleation in early development. Early blind and sighted littermates were reared in enriched environments to promote active tactile exploration in three-dimensional (3D) space, or in standard laboratory cages. In adulthood, both enriched groups showed adaptive changes in exploration patterns and gap crossing behaviors relative to standard-reared counterparts. Thus, early blind animals showed behavioral compensation when challenged by complex environments. Enriched rearing increased selectivity of S1 neural responses to whisker touch and altered receptive field shapes such that they were less horizontally anisotropic. This shift was strongest in enriched early blind animals, enhancing tuning along the behaviorally relevant horizontal axis more than in standard-reared early blind animals. Thus, alterations in receptive fields of neurons in S1 following early blindness were amplified by environmental complexity. Sighted opossums reared with enrichment also showed similar whisker receptive field plasticity, though to a slightly lower degree. Together, these results demonstrate the strong influence of the rearing environment on reorganization of cortex that processes inputs from the spared senses, underscoring the role of experience in directing compensatory plasticity following early sensory loss.

SIGNIFICANCE STATEMENT

Enhanced perceptual abilities following early sensory loss are often attributed to cross-modal recruitment of cortex linked to the deprived sense. However, plasticity also occurs within cortical areas representing spared modalities. It remains unresolved whether deprivation alone is sufficient to induce such reorganization, or whether experience using the spared sense is required. We show that enriched rearing amplifies neural coding changes in primary somatosensory cortex after early blindness, shaping receptive field geometry and promoting adaptive behavioral strategies aligned with environmental demands. Comparable changes in sighted animals reared under the same conditions reveal that reliance on touch—rather than visual deprivation alone—drives this neural and behavioral plasticity, supporting the critical role of experience in enhancing functional outcomes after early sensory impairment.

INTRODUCTION

Early loss of vision leads to heightened perceptual abilities in the remaining senses, accompanied by structural and functional brain plasticity (Goldreich and Kanics, 2003; Wong et al., 2011; Voss, 2011; Morland et al., 2021; Mezzera and López-Bendito, 2016; Lazzouni and Lepore, 2014). However, the developmental origins of this plasticity remain poorly understood, particularly the extent to which sensory experience, movement options and affordances contribute to compensatory changes in brain and behavior. In this study, we investigate how enhancing tactile experience through rearing in an enriched environment contributes to neural and behavioral plasticity of whisker-mediated touch in a marsupial, the short-tailed opossum, when vision is lost early in development.

Exposure to enriched environments drives plasticity in sensory cortices of juvenile

and adult animals with intact brains and bodies. This includes functional reorganization

of sensory representations through changes in cortical territory, neuronal response

properties (latency, strength, and selectivity), neuronal morphology and molecular

expression (Seo, 1992; Leah et al., 1985; Coq and Xerri, 1998; Cancedda et al., 2004;

Engineer et al., 2004; Polley et al., 2004; Guic et al., 2008; Mégevand et al., 2009;

Devonshire et al., 2010; Alwis and Rajan, 2013, LeMessurier et al., 2018, Bibollet-Bahena

2023). Enrichment also counteracts degradation of representations after transient

sensory deprivation, and promotes recovery from brain injury (Polley et al., 1999;

Bartoletti et al., 2004; Sale et al, 2007; Nakashima and Dyck, 2008; Wang et al., 2013;

Greifzu et al., 2014; Zhu et al., 2014; Zheng et al., 2014; Jiang et al., 2015; Kalogeraki et

al., 2017; Levine et al., 2017). Rearing animals in naturalistic environments aligned with

a species' lifestyle engages ethologically relevant behavior (Landers et al., 2011;

Wilkinson et al, 2010; Branchi et al., 2011; Weissbrod et al, 2013) and is especially

effective in reshaping cortical sensory representations at structural, functional and

molecular levels (Polley et al., 1999; Polley et al., 2004; Frostig, 2006; Gomez-Pinilla et

al., 2011; Landers et al., 2011). Here, we therefore used a complex three-dimensional

rearing environment designed to model aspects of the natural habitat and lifestyle of

short-tailed opossums.

Many small mammals, including short-tailed opossums, routinely use forest

habitats across vertical strata, from floor to canopy (Jenkins, 1974; Jones et al., 2003;

Delciellos and Vieira, 2006; Abreu and Oliveira, 2014; Shapiro et al., 2014). Short-tailed

opossums are terrestrial hunters and foragers (Jones et al., 2003) but prefer nesting

above ground in rocky outcrops (Macrini, 2004), using branches and logs as arboreal

runways to negotiate complex terrain (Lammers, 2007; Lammers and Gauntner 2008;

Ladine and Kissel, 1994; Montgomery, 1980). They are skilled climbers despite lacking

morphological specializations of arboreal species, relying on behavioral strategies to

maintain balance and traverse arboreal substrates (Saunders et al., 1995; Saunders et

al., 1998; Wilkinson et al., 2010; Schmitt et al., 2010; Rupert et al., 2014; Goin et al.,

2016; Thomas et al., 2017; Lammers, 2004; Lammers, 2007; Lammers, 2009; Lammers

and Biknevicius, 2004; Lammers et al., 2006; Lammers and Gauntner, 2008). Whisker

touch is critical for navigating spatially complex 3D environments (Ahl, 1986; Arkley et al.,

2017), guiding gap crossing and forelimb placement (Arkley et al., 2014; Niederschuh et

al., 2015; Arkley et al., 2017; Grant et al., 2018). Peripheral organization of the whisker

system reflects lifestyle demands and is more developed in arboreal than terrestrial

species; in short-tailed opossums, whisker layout is consistent with semi-arboreal and

scansorial marsupials (Beddard, 1902; Pocock, 1914; Lyne, 1958; Lyne, 1959;

Muchlinski, 2013; Grant et al., 2017; Grant et al., 2018; Ramamurthy and Krubitzer, 2016;

Grant et al., 2013). Thus, vertical environmental use depends heavily on behavioral

modifications guided by tactile input (Lammers and Biknevicius, 2004; Lammers, 2009).

Early loss of visual inputs in short-tailed opossums at P4, prior to the formation of

retinogeniculate and thalamocortical pathways, enhances tactile acuity and alters neural

coding of whisker inputs in primary somatosensory cortex (S1), with receptive fields

exhibiting greater selectivity along the principal axis of whisker motion (Ramamurthy &

Krubitzer, 2018). Further, there is major reorganization of thalamic and cortical

connections of somatosensory cortex relative to sighted animals (Dooley & Krubitzer,

2019). This raises a critical question: are alterations in somatosensory coding driven primarily by deprivation-induced anatomical reorganization, or by tactile experience itself? Here, we implement rearing conditions that encourage naturalistic tactile exploration in a spatially challenging environment designed to increase reliance on whisker use. By comparing how an enriched environment shapes neural coding and behavior related to whisker touch in sighted and early blind animals, we interrogate the role of tactile experience in directing compensatory plasticity following early blindness.

MATERIALS AND METHODS

Animals

Thirty-three adult short-tailed opossums of either sex (*Monodelphis domestica*; age range: 4-16 months; weight range: 61-165g) were used for electrophysiological recording experiments. Animals belonged to four experimental groups (**Table 1**), early blind and sighted reared in two conditions (standard: 's', enriched: 'e'). Previously published data from standard-reared animals were reanalyzed for comparison with enriched rearing groups (Ramamurthy and Krubitzer, 2018).

Ethics Statement

All protocols were approved by the Institutional Animal Care and Use Committee, and experiments were conducted according to the criteria outlined in the National Institutes of Health *Guide for the Care and Use of Laboratory Animals*.

Bilateral enucleation

For early blind opossums, bilateral enucleations were performed at P4 (**Figure 1**). Mothers of experimental litters were lightly anesthetized with Alfaxan (initial dose: 20

mg/kg; maintenance doses: 10-50%; IM) to facilitate enucleation of the pups, which are

attached to the mother at this developmental stage. Pups were anesthetized by

hypothermia. Health of both the mother (respiration rate and body temperature) and the

pups (heartbeat, coloration and mobility) was monitored throughout the procedure. An

incision was made in the skin covering the eyes, the eyes were removed under

microscopic guidance, and the skin was repositioned and resealed using surgical glue.

Approximately 50% of each litter was bilaterally enucleated and the remaining littermates

served as sighted controls. After complete recovery from anesthesia, mothers along with

their attached litters were placed into either standard or enriched housing, depending on

the experimental group.

Rearing paradigms

Mothers of all experimental groups were reared and housed in standard laboratory

cages until bilateral enucleations were performed on the pups at P4. Following the

enucleation procedure, mothers with litters belonging to the standard rearing groups were

placed in standard laboratory cages, while those belonging to the enriched rearing groups

were placed in custom-made enriched cages (**Figure 1-1**). All animals were maintained

on a 14/10 light/dark cycle. Food and water were available *ad libitum* in the cage under

both rearing conditions. All experimental litters were housed together with the mother until

weaning at ~8 weeks (P56), and at that stage they were separated from the mother, but

the whole litter continued to be housed together until adulthood at ~4 months (P120).

Adults (>P120) were individually housed under standard rearing conditions, but in groups

of 2-4 same-sex littermates under enriched conditions (see below).

Standard rearing: Under standard rearing conditions, adult animals were singly housed

in standard laboratory cages, in cages similar to those typically used for laboratory

rodents (cage dimensions: 18.5"l x 10"w x 8"h). The home cage was provided with a

nesting cup and nesting material (shredded paper towels).

Enriched rearing: The enriched rearing environment in our study incorporated features

common to environmental enrichment (EE) paradigms previously used in rodents

(Nithianantharajah and Hannan, 2006; Sale et al., 2009; Bengoetxea et al., 2012), and

additionally takes into consideration the natural lifestyle of short-tailed opossums (**Figure**

1-1). Juvenile animals were cohoused until weaning with the mother as well as littermates,

and then with just their littermates from weaning until adulthood, as described above.

Enriched cages were provided with standard bedding and nesting material, as well as

some additional nesting material with a different texture. In addition to the nesting cup

placed on the floor as in standard cages, enriched cages provided the option of nesting

high above the cage floor, in nesting boxes attached to the cage walls. A single nesting

box was provided in the cage at juvenile stages of the experimental litters. In adulthood,

balanced numbers of eSC and eEB littermates were housed together in each enriched

cage, with a maximum of four opossums per cage. Each adult animal was provided with

an individual nesting box. Unlike common enrichment paradigms, which typically

incorporate toys into standard cages, the enriched cages used here were substantially

larger (~8x in volume; 29"l x 18"w x 24"h) compared to standard cages to encourage

tactile exploration in all three dimensions. The greatest increase relative to standard

cages was along the vertical dimension: enriched cages were 3x taller than standard

cages. Enrichment toys (Bio-Serv) of different shapes and textures were provided in the

cage. Toys were replaced, and the positions of manzanita branches were changed every

2–3 days to introduce novelty and promote continual exploration of the environment.

Behavioral testing

<u>Arena testing</u>: Arena testing was performed in three adult animals from each of the four

experimental groups. The arena consisted of a cage with the dimensions of enriched

cages, containing a single nest box near the top of the arena, with a branch leading up to

it. Animals were placed in the arena and allowed to explore it for 5 minutes in complete

darkness, and movies of their behavior were acquired under infrared (IR) illumination.

idTracker software (**Figure 2-1A**; Romero-Ferrero et al., 2019) was used to extract

movement trajectories of individual animals, which was further analyzed in MATLAB. In

each movie frame, we segmented the arena into a vertical zone (cage walls, branches,

and nest area) and a horizontal zone (cage floor). Exploratory activity was quantified from

session-wise binarized matrices (1280 × 720 pixels; height × width) containing the

animal's position in each frame. To assess differences in how animals sampled vertical

space in the arena, binarized occupancy maps were summed across sessions separately

for standard- and enriched-reared animals and collapsed along the horizontal axis,

yielding a per-row estimate of vertical occupancy. Vertical occupancy profiles were then

binned in 20-row increments (20 px per bin; vertical dimension only) and normalized to

the percentage of total vertical occupancy (summing to 1 across the vertical axis). For

each animal, total vertical occupancy (V) and horizontal occupancy (H) were computed

by summing binarized pixel values within the corresponding zone across all frames. A

normalized vertical preference index was calculated for each animal as:

$$Vertical\ Preference\ Index = \frac{(V - H)}{(V + H)}$$

Additional analysis of exploratory behavior in the arena test was performed through

manual scoring in BORIS (Behavioral Observation Research Interactive) software. All

home cage behavior data were analyzed offline by four independent observers such that

multiple epochs were scored by at least two independent observers with a high interrater

reliability (Cohen's κ coefficient of > 0.85; κ = 0 indicates no agreement in scoring between

observers and κ = 1 indicates perfect agreement; Ramamurthy & Krubitzer, 2018; Landis

and Koch, 1977). Arena exploration was initially scored using a binary scale of active

states (locomotion bouts) and inactive states (resting bouts). A switch between an active

and inactive state was only considered to have occurred if a bout lasted for at least 5 s.

Active states were then further scored for crossing events that occurred during natural

exploration of the arena. Crossing events were defined as instances in which an animal

crossed from one surface to another within the arena. Four types of crossing events were

scored: (1) floor to wall/wall to floor (2) wall to wall (3) floor to branch/branch to floor (4)

nest or branch to wall/wall or branch to nest. Since systematic differences were not

observed across crossing event types, only combined summary plots are shown.

<u>Gap crossing task</u>: Gap crossing was tested in 3-4 adult animals per experimental group.

Animals were allowed to spontaneously cross from one platform ("start") to another

platform ("goal") with either small (2-10 cm) or large gap distances (12-14 cm) separating

them, which were interleaved in a pseudorandom order. Animals were tested for 7 days

with intact whiskers, followed by 1-3 days of testing after whisker trimming. On each trial

animals were allowed to remain on the start platform and attempt crossing for a maximum

of 1 minute. If no crossings were performed, the trial was ended, and animals were

removed from the platform for 1 min before returning to the platform for the next trial.

Movies of this behavior were acquired under IR illumination or dim red light, and

DeepLabCut was used for analysis of gap crossings (**Figure 2-1B-C**). The snout tip, the

base of the tail and the position of the edges of each platform were labeled in a subset of

frames to generate the dataset (260 frames across 13 opossums) on which the network

was trained (200,000 iterations). On a subset of trials, slow-motion movie clips in a

zoomed-in field of view were acquired to visually confirm the use of whiskers during gap

crossing. Infrared motion sensors (emitter/detector pairs) placed on either side of the

edge of each platform detected beam breaks and beam break times were recorded as a

secondary method of confirmation.

Surgical procedures

Animals were anesthetized with urethane (initial dose: 1.25 g/kg, 30% in saline, IP;

supplemental doses: 0.125-0.313 g/kg, 30% in saline, IP). Vital signs (respiration and

body temperature) were monitored throughout the experiment. At the beginning of the

surgery, dexamethasone (0.4-2.0 mg/kg; IM) was administered to minimize intracranial

swelling. Lidocaine (2% solution; SC) was injected at the midline of the scalp and around

the ears. The animal was then placed in a stereotaxic frame. Following an incision at the

midline of the scalp and retraction of the temporal muscle, a craniotomy was performed

to expose the entire parietal cortex and the dura was retracted. The surface of the

neocortex was covered with a layer of silicone fluid to prevent desiccation and then

photographed so that electrode penetration sites could be related to patterns of

vasculature. After the craniotomy, the animal was removed from the stereotaxic frame to

permit full access to the face for whisker stimulation. Three stainless steel skull screws

were inserted in the skull contralateral to the recording electrode. The head of the animal

was stabilized by cementing a head post to the skull screw in a rostral location over the

cortex. The other two skull screws were inserted epidurally into the skull over the

cerebellum and over the olfactory bulbs, serving as the reference and ground electrodes,

respectively, and also providing additional anchoring support for the dental cement.

Extracellular recordings and whisker stimulation

Somatosensory receptive fields were first coarsely mapped by recording multiunit

activity while applying tactile stimuli using a handheld probe. The representation of the

whiskers in S1 was localized by progressively testing electrode penetration sites medial

to the rhinarium representation and caudal to the lower jaw/lip representation. After this,

when receptive fields were found to be located on the mystacial or genal whisker pad

(Ramamurthy & Krubitzer, 2016), computer-controlled whisker deflections were used to

quantitatively measure single unit receptive fields.

Whisker stimuli consisted of single whisker deflections (2°) delivered using

piezoelectric actuators (4 ms/100 ms/4 ms ramp-hold-return; 5 mm from the whisker

base; 1.4 mm forward excursion). Data were collected for a grid of sixteen whiskers on

the mystacial pad to enable construction of somatotopic receptive fields. Whiskers were

deflected in a pseudorandom order with 50-100 trials collected per whisker (in blocks of

3-50 trials) at each recording site. Extracellular recordings (amplifier gain: 10,000X; A-M

Systems Model 1800 Microelectrode AC Amplifier; A-M Systems, Carlsborg, WA;

sampled at 28 kHz; Power1401, Cambridge Electronic Design Limited, Cambridge, UK)

were made 400-500 μm below the pial surface using monopolar tungsten microelectrodes

(FHC, Inc., Bowdoin, ME; 1-5 MΩ at 1 kHz) lowered with a hydraulic microdrive (David

Kopf Instruments, Tujunga, CA). Recordings were bandpass filtered (300-3000 Hz), and

spike-sorting was carried out offline to confirm isolation of single units. After

electrophysiological data was collected, fluorescent probes were used to mark the

location of recording sites, so that these could be related to histologically identified cortical

field boundaries (**Figure 3-1**).

Histology

At the end of each experiment, animals were injected with an overdose of sodium

pentobarbital (Beuthanasia; 250 mg/kg IP) and transcardially perfused with 0.9% saline,

followed by 2-4% paraformaldehyde in phosphate buffer, and then 2-4%

paraformaldehyde in 10% phosphate-buffered sucrose. The brain was extracted and

post-fixed (1-2 hours in 4% paraformaldehyde in 10% phosphate-buffered sucrose). In

some cases, the cortical hemispheres were flattened and the tissue was sectioned

tangentially at 30 µm using a freezing microtome, and then processed for myelin staining

(Gallyas, 1979; Dooley et al, 2013) to enable reconstruction of cortical field boundaries.

Reconstructions were drawn using Adobe Illustrator CS5 (Adobe, San Jose, CA). Digital

images of processed tissue were obtained using either a Nikon Multiphot system or an

Optronics MicroFire digital microscope camera. Contrast and brightness of whole images

were adjusted using Adobe Photoshop CS5 (Adobe, San Jose, CA).

Neural data analysis

Data files were processed blind to experimental group identity. Single-unit isolation

was performed while running the experiment using template-matching procedures in

Spike2 (Cambridge Electronic Design Limited, Cambridge, UK; RRID:SCR_000903), and

unit isolation was verified offline in Spike2, using principal component analysis. Stable

waveforms and firing rates over the course of the recording session, and <0.5% refractory

period violations (interspike interval <1.5 ms) were required for units to be included in the

analysis. Spike times were then exported to MATLAB (Mathworks, Inc., Natick, MA;

RRID:SCR_001622) and analyzed using custom scripts. Whisker evoked firing rates were

measured as the spike count following stimulus onset (0-100 ms) and were considered

significant if they exceeded prestimulus spike counts in an equivalent time window by two

standard deviations. The *best whisker* (BW) was defined as the whisker with the highest

evoked response, and depending on their proximity to the BW the remaining whiskers

were categorized as *first-order surround whiskers* (1° SW; immediately adjacent to BW),

second-order surround whiskers (2° SW; one position removed from BW) and *third-order*

surround whiskers (3° SW; two positions removed from BW).

One measure of receptive field size used was the number of whiskers that evoked

a significant response in each recorded neuron. Another measure used was the rank-

ordered whisker tuning curve for each neuron, where responses to all whiskers tested

were sorted in descending order of whisker-evoked response magnitude, and normalized

to peak (BW response). Further, mean 2D somatotopic receptive fields were obtained

after normalizing SW-evoked response magnitudes to the BW-evoked response, and

aligning receptive fields of single units such that the BW was at the center of the receptive

field for each unit. Because *Monodelphis* lacks discrete anatomical cortical barrels

(Ramamurthy & Krubitzer, 2016), 2D receptive fields were aligned relative to the BW

position rather than an anatomically defined "principal whisker" or "columnar whisker". 50,

75, and 90% contour lines (for response levels relative to the BW) were plotted for the

mean receptive fields (smoothed by linear interpolation) to facilitate visualization. Raw

values were used for all calculations and statistical comparisons.

We define anisotropy in receptive fields as the difference in the tuning width of an

S1 neuron's receptive field when measured along different spatial directions of the

whisker pad—across arcs (horizontal axis) and across rows (vertical axis). Anisotropies

in the structure of somatotopic receptive fields were quantified as the *shape index:*

301
$$Shape\ index = \frac{(Row\ tuning\ width\ index - Arc\ tuning\ width\ index)}{(Row\ tuning\ width\ index + Arc\ tuning\ width\ index)}$$

If the sum of the row and arc tuning width indices was zero, the shape index was

assigned to be zero. Shape indices ranged from -1 to 1, with values >0 indicating broader

tuning along the row compared to the arc (i.e. broader horizontal tuning), and values <0

indicating broader tuning along the arc compared to the row (i.e. broader vertical tuning).

Neuronal selectivity for the best whisker along the row and arc dimensions was measured

by assessing tuning width separately along the horizontal and vertical axes as the *row*

tuning width index and the *arc tuning width index* respectively, either considering only 1°

SW, or 1°*,* 2°, 3° SW in the row or arc of the BW, as stated below:

310
$$Row\ tuning\ width\ index = \frac{(\text{Mean evoked response of } SW_{row})}{\text{BW evoked response}}$$

311
$$Arc\ tuning\ width\ index = \frac{(\text{Mean evoked response of } SW_{arc})}{\text{BW evoked response}}$$

312 Additionally, the *whisker selectivity index (WSI)* for BW vs. in-row or in-arc whiskers was

313 computed separately for each neuron, considering BW responses relative to 1°*,* 2°, 3°

314 SW responses.

315
$$WSI_{row} = \frac{(BW\ evoked\ response - SW_{row}\ evoked\ response)}{(BW\ evoked\ response + SW_{row}\ evoked\ response)}$$

316
$$WSI_{arc} = \frac{(BW\ evoked\ response - SW_{arc}\ evoked\ response)}{(BW\ evoked\ response + SW_{arc}\ evoked\ response)}$$

In-row and in-arc selectivity indices ranged from 0 to 1, with values closer to 1

indicating higher selectivity for the BW in the corresponding dimension.

Receptive field shape was quantified at two spatial scales (1° only vs. 1°-3°)

because somatotopic receptive fields of neurons in the S1 whisker representation consist

of the BW (peak response) and a graded, multi-whisker surround shaped by intracortical

integration, including anisotropies along row vs. arc axes. This receptive field organization

has been demonstrated in classical mouse and rat studies (e.g., Simons, 1978; Brecht &

Sakmann, 2002; Fox, 2008), as well as prior work demonstrating similar receptive field

structure in opossum S1 (Ramamurthy & Krubitzer, 2016; Ramamurthy & Krubitzer,

2018).

Since receptive field shape analysis characterizes the overall geometry of the field

around the BW, it cannot be meaningfully computed for a non-continuous portion of the

surround alone (e.g., only 2°/3° SWs); therefore 2°/3° were not analyzed separately from

1° SWs; instead, shape analyses with only 1° SWs characterized local receptive field

configuration, whereas analyses including 1°-3° SWs captured receptive field

configuration shaped by broader spatial integration.

Statistical Analysis

Statistical analyses were performed in MATLAB and R. Permutation tests for the

difference in medians were used to assess differences between two groups (referred to

in brief as 'permutation test'). Fisher's exact tests (two-sided) were used for analyses of

2x2 contingency tables to compare proportions across conditions. All statistical tests for

neuronal data use 'n' of cells. We accounted for interindividual variability using linear

mixed effects models.

To model crossing behavior in the gap-crossing task, we fit a generalized linear

mixed-effects model using 'fitglme' in MATLAB (Poisson distribution; log link). Crossing

counts were modeled with the log of observation time included as an offset to estimate

rates. Fixed effects included experimental group, gap size, whisker trimming, and the

interactions of experimental group with gap size and with whisker trimming, respectively.

Since whisker trimming abolished crossings on 100% of large-gap trials, those trials were

excluded from this analysis. Animal identity was included as a random intercept (1|ID) to

account for repeated measures. The model specification was as follows:

$Response\ Variable \sim Experimental\ Group * Gap + Experimental\ Group * Trim + (1|Animal\ ID)$

The model was estimated using the Laplace approximation for maximum likelihood.

Overdispersion was evaluated via Pearson dispersion; if dispersion exceeded 1.5, we

added an observation-level random effect (1|Obs) to handle overdispersion.

To analyze gap crossing durations on successful trials we fit a linear mixed-effects

model using 'fitlme' in MATLAB with the specification:

$Response\ Variable \sim Experimental\ Group * Gap + Experimental\ Group * Trim + (1|Animal\ ID)$

where experimental group and test condition were modeled as fixed effects, and animal

ID modeled as a random effect.

Planned post hoc comparisons for both models were performed using coefTest

with contrast matrices for group and condition effects (gap size and whisker trimming),

including a difference-in-differences contrast for the blindness × environment interaction.

'coefTest' returned p-values for each contrast, with Satterthwaite degrees of freedom

applied where applicable.

For neural data, experimental group was modeled as a fixed effect and animal

identity as a random effect, using the model specification:

$Response\ Variable \sim Experimental\ Group + (1|Animal\ ID)$

Planned contrasts were performed using 'coefTest' for pairwise group comparisons and

for a difference-in-differences (blindness × environment) contrast, with p-values returned

and Satterthwaite degrees of freedom applied where applicable. Additional models were

fit for key analyses with sex included as a fixed effect to test for differences between

males and females, but no significant sex differences were identified.

Exact p-values for fixed effects of experimental group and/or test condition (gap

size/whisker trimming) obtained from Poisson and Gaussian generalized linear mixed-

effects models, or from Fisher's exact tests for categorical comparisons, with pairwise

contrasts across experimental groups are provided in **Table 2** and **Table 3.**

RESULTS

Behavioral adaptations of blind and sighted opossums in enriched rearing

environments

In short-tailed opossums, postnatal day 0 (P0) is developmentally equivalent to

embryonic stages in rodents (approximately E11 in mouse and E12 in rat; Molnár and

Blakemore, 1995; Molnár et al., 1998). Bilateral enucleations were conducted at P4,

before retinal afferents and thalamocortical axons reached their respective targets (Taylor

and Guillery, 1994; Molnár et al., 1998), completely eliminating both spontaneous retinal

activity and visually evoked input (**Figure 1)**. Following bilateral enucleation on postnatal

day 4, opossum mothers and their litters, including sighted and blind pups, were

transferred into a new standard or enriched home cage environment (**Figure 1**; **Figure 1-**

1A-D). Prior to P12, sensory experience in either environment was determined entirely

by behavioral patterns of the mother, due to the obligate attachment of opossum pups.

Juvenile animals continue to rely on the mother for both sustenance and transportation

until weaning at P56, but with increasing independent exploration (**Figure 1-1E-F**).

Opossum mothers always preferentially nested high above the ground (n = 3 mothers,

producing a total of 4 enriched experimental litters; example in **Movie 1**), so pups were

forced to rely heavily on touch to navigate the enriched environment. By adulthood, both

sighted and early blind animals were highly adapted to their environments (**Movie 2-3**).

To assess how rearing condition and visual condition (blind versus sighted) shape

exploratory behavior, we tested each group in a novel arena matching the dimensions

and basic layout of the enriched home cages (**Figure 2; Figure 2-1A**). We quantified

spatial sampling preferences (**Figure 2A-C** and **2E**), overall locomotor activity (**Figure 2D**

and **2F**), and spontaneous crossings between surfaces during natural exploration (**Figure**

2D and **2G**), capturing both spatial and temporal aspects of behavior. These measures

contextualize and complement the more controlled gap crossing behavior assay

presented in the next section. Movement trajectories were highly similar for sighted and

blind animals within each rearing condition (**Figure 2A-B**; **Figure 2-1A**), consistent with

successful behavioral compensation by blind animals even in more spatially complex

environments. In contrast, rearing environment strongly impacted how animals sampled

the novel arena: enriched opossums explored vertical space more extensively than those

reared in standard cages (p = 0.018, permutation test; **Figure 2C**; **Figure 2E**), though

overall activity levels (p=0.565, permutation test) and durations of naturally occurring

crossing events were not significantly different (p=0.232, permutation test; **Figure 2D**;

Figure 2F-G). Quantifying gap crossing during free exploration has some limitations

which may mask differences in sensory-mediated behavior. Crossing events that

occurred during free exploration, where animals crossed from surface to surface did not

necessarily involve crossing of discrete gaps, and we could not control for gap size across

different crossing events. Further animals were free to adopt varying postures prior to

crossing, and to use varying behavioral strategies involving sensory inputs from multiple

body parts to aid gap crossing during free exploration.

To assess sensorimotor performance under more precisely defined task

parameters, we used a gap crossing task (conducted in darkness) in which gap size was

systematically varied across trials, animals had only one possible path forward (across

the gap), and the constraints of the apparatus required animals to adopt a standardized

posture prior to gap crossing (**Figure 3A**; **Figure 2-1B-C**). Crossing large gaps required

whisker use, as the goal platform was out of reach from the start platform except through

whisker contact. We quantified (1) the fraction of trials resulting in a successful crossing

(**Figure 3B**), (2) crossing rates over time (**Figure 3C**), and (3) the duration of successful

crossing events (**Figure 3D**).

The fraction of successful trials (i.e. the probability of success) was strongly

impacted by gap size (sSC: 8%, sEB: 14%, eSC: 17%, eEB: 14%; percent successful

trials, in comparison to 100% success on small gap trials) and by whisker trimming

(**Figure 3B**, Fisher's exact test, $p < 0.001$; see **Table 2** for exact p-values for the fixed-

effect of experimental group, and for all pairwise comparisons between groups),

eliminating successful crossings at large gaps for all experimental groups. This is

consistent with the role of whiskers in gap crossing described in previous studies of small

mammals (Hutson & Masterton 1986, Voigts et al., 2015, Arkley et al. 2017, Graham &

Socha 2020, Toso & Diamond 2023). All animals crossed on 100% of small gap trials,

with or without whiskers, indicating that other body parts also contributed to small gap

crossings.

Crossing rates, which reflect success rate normalized by time, were significantly

lower for large gaps than for small gaps within each experimental group (**Figure 3C,**

Table 2, **Table 3-1**; $p < 0.001$ in all cases, all p-values from Poisson generalized linear

mixed-effects model) with no significant differences across groups ($p > 0.05$). Although

there was a trend toward decreased crossing rates in all experimental groups with whisker

trimming, trimming effects were not statistically significant for crossing rates ($p > 0.05$).

441   The third metric in the gap crossing task was the duration of successful crossings

442 (**Figure 3D**, **Table 2**, **Table 3-1**). We observed significant fixed effects of experimental

443 group ($p < 0.001$; all p-values from the linear mixed effects model), gap size ($p < 0.001$),

444 and whisker trimming ($p < 0.001$), as well as interactions between group and gap size (p

445 < 0.001) and between group and trimming ($p = 0.01$). Across groups, successful crossings

446 were generally slower for large versus small gaps, with modest but significant increases

447 in crossing times on small gaps after whisker trimming (with one exception; see below).

448 Consistent with **Figure 3B**, this pattern suggests primary reliance on whiskers for large

449 gaps, whereas small gaps can be crossed using strategies that incorporate both whiskers

450 and tactile inputs from other body parts (Ramamurthy et al., 2021).

451   Group differences based on visual condition and rearing condition were most

452 pronounced for large gaps (**Figure 3D**). Early blind opossums crossed large gaps faster

453 than sighted controls under standard rearing (sSC vs. sEB: $p < 0.001$), and this advantage

454 was amplified by enriched rearing (eEB vs. sEB: $p < 0.001$). Both enriched groups were

455 significantly faster than standard-reared animals ($p < 0.001$ for all pairwise comparisons),

456 with enriched blind animals outperforming enriched sighted animals ($p < 0.01$). Enriched

457 blind animals followed the general trends of other groups but were the only cohort without

458 a significant difference between large and small gap durations ($p > 0.05$), reflecting

459 comparatively faster large gap performance. They were also the only group in which

460 whisker trimming did not significantly prolong small gap crossings ($p > 0.05$). Taken

461 together, these results indicate heightened performance supported by whisker use on

462 large gaps, along with adaptive recruitment of other body parts to maintain small gap

463 performance when whiskers are absent.

Tactile experience amplifies compensatory neural plasticity in blind opossums

Single-unit recording experiments were performed in 19 opossums reared in standard conditions (10 sSC, 9 sEB), and 14 opossums reared in enriched conditions (7 eSC, 7 eEB), acquiring data for a total of 339 neurons across experimental groups (see **Table 1**; **Figure 3-1**). 63-73% of units recorded in S1 (sSC: 60, sEB: 64, eSC: 58, eEB: 51) had a significant evoked response (>2 SD above baseline) to stimulus onset following single whisker deflections, and these neurons were included in receptive field analyses. Raster plots and peristimulus time histograms (PSTHs) for whisker-responsive neurons in each group are shown in **Figure 4A-B**. These examples illustrate more selective whisker touch responses in standard-reared blind opossums, with selectivity further amplified in enriched groups.

In general, spontaneous firing rates of S1 neurons were low, consistent with previously reported findings in standard-reared animals (Ramamurthy and Krubitzer, 2018), and were not significantly different among experimental groups (**Figure 4C**, **Table 3**, **Table 4-1**; mean ± SEM, sSC: 0.582 ± 0.083 spk/s, sEB: 0.613 ± 0.121 spk/s, eSC: 0.544 ± 0.079 spk/s, eEB: 0.503 ± 0.082 spk/s; linear mixed-effects model, p>0.05; see **Table 3** for exact p-values for the fixed-effect of experimental group, and for all pairwise comparisons between groups). However, responses evoked by the stimulation of even the most preferred single whisker stimulus (best whisker; BW) in enriched groups were suppressed in relation to stimulus-evoked firing rates in standard-reared sighted opossums (**Figure 4D**, **Table 3**, **Table 4-1**; mean ± SEM, sSC: 11.4 ± 0.839 spk/s, sEB: 8.39 ± 0.591 spk/s, eSC: 7.82 ± 0.748 spk/s, eEB: 4.57 ± 0.52 spk/s; linear mixed-effects model, eSC vs. sSC: p<0.01, eEB vs. sSC: p<0.001), as previously reported in early blind

animals under standard rearing conditions (Ramamurthy and Krubitzer, 2018). Since

relative suppression of BW-evoked responses was observed in enriched sighted controls,

in addition to enriched early blind animals, this effect was not contingent on the lack of

vision itself – rather it was driven by increased dependence on touch in the enriched

groups of opossums. Mean BW-evoked firing rates in enriched sighted control animals

were comparable to those in standard early blind animals (linear mixed-effects model,

sEB vs. eSC: $p>0.05$). However, in enriched early blind animals, mean BW-evoked firing

rates were lower relative to all other experimental groups (linear mixed-effects model,

eEB vs. sEB: $p<0.001$, eEB vs. eSC: $p<0.001$, eEB vs. sSC: $p<0.001$). Thus, blindness

was not required for suppression of whisker-evoked responses, but the combination of

tactile enrichment and vision loss produced the greatest effect on whisker-evoked firing

rates of S1 neurons. This is consistent with animals relying on their whiskers to a greater

extent when reared in a complex environment in the absence of vision.

Average whisker receptive fields in both sighted and blind enriched groups were

significantly smaller than those in standard-reared sighted controls (mean ± SEM, eEB:

3.22 ± 0.409, eSC: 3.48 ± 0.364, sSC: 5.92 ± 0.427; linear mixed-effects model, eSC vs.

sSC: $p<0.001$, eEB vs. sSC: $p<0.001$) when measured as the number of whiskers that

evoked a significant response above spontaneous firing (**Figure 4E**, left, **Table 3**, **Table**

4-1). We see similar results when receptive fields are quantified as the mean rank-ordered

tuning curve across neurons (**Figure 4E**, right, **Table 3**, **Table 4-1**; linear mixed-effects

model, eSC vs. sSC: $p<0.001$). The finding that tactile enrichment in sighted control

animals leads to a decrease in mean receptive field size—similar to both standard-reared

and enriched early blind animals—supports the interpretation that this reduction reflects

an experience-dependent effect across all three groups. However, changes in measures

of overall receptive field size in enriched early blind animals relative to standard-reared

early blind animals were either not significant (**Figure 4E**, left, number of whiskers: eEB:

3.22 ± 0.409, sEB: 3.38 ± 0.270; linear mixed-effects model, eEB vs. sEB: p>0.05) or

modest (**Figure 4E**, right, rank-ordered tuning curve: linear mixed-effects model, p<0.01).

This suggests that, under standard rearing conditions, receptive field plasticity in early

blind animals may already have reached a limit that cannot be further shifted by

enrichment.

Enrichment enhances selectivity for whisker touch along the behaviorally relevant

horizontal axis

To better understand the effects of enriched rearing on receptive field configuration

of S1 neurons, we characterized two-dimensional receptive fields for the row (horizontal)

and arc (vertical) dimensions of the whisker pad (**Figure 5A**). Visual inspection of mean

2D receptive fields revealed clear differences in receptive field anisotropies across

experimental groups (**Figure 5B**). We quantified anisotropies in receptive fields as the

shape index, ranging in value from -1 to 1, where positive values indicated horizontally

anisotropic shapes, negative values indicated vertically anisotropic shapes and values

close to 0 indicated more isotropic shapes. Receptive fields for neurons in the S1 whisker

representation in standard-reared sighted control and early blind animals were dominated

by horizontally anisotropic shapes (**Figure 5C-D**, **Table 3**, **Table 5-1**). However, for early

blind animals reared in enriched conditions, the receptive field shape distributions were

shifted away from horizontal anisotropy – either when considering only 1° surround

whisker (SW) positions (**Figure 5C**; neurons with shape index > 0.0, sSC: 68.3%, sEB:

57.8%, eEB: 37.3%, Fisher's exact test, eEB vs. sSC: p=0.001, eEB vs. sEB: p<0.05;

linear mixed-effects model, eEB vs. sSC: p<0.001, eEB vs. sEB: p<0.05) , or when

including all SW (1° - 3°) positions (**Figure 5D**; neurons with shape index > 0.0, eEB:

37.3%, sSC: 81.7%, sEB: 76.6%, Fisher's exact test, eEB vs. sSC: p<0.001, eEB vs. sEB:

p<0.001; linear mixed-effects model, eEB vs. sSC: p<0.001, eEB vs. sEB: p<0.001).

538   Receptive field distributions were also shifted away from horizontal anisotropy in

sighted control animals reared in the enriched environment. This shift was not seen when

only 1° SW positions were considered (**Figure 5C**; neurons with shape index > 0.0, eSC:

55.2%, Fisher's exact test, eSC vs. sSC: p>0.05, eSC vs. sEB: p>0.05; linear mixed-

effects model, eSC vs. sSC: p>0.05, eSC vs. sEB: p>0.05) but becomes apparent when

1° - 3° SW positions are considered (**Figure 5D**; neurons with shape index > 0.0, eSC:

46.6%, Fisher's exact test, eSC vs. sSC: p<0.001, eSC vs. sEB: p<0.001; linear mixed-

effects model, eSC vs. sSC: p=0.01, eSC vs. sEB: p>0.05). Thus, rearing in an enriched

environment, which was permissive of more naturalistic patterns of tactile behavior,

shifted the shapes of receptive fields of S1 neurons away from the horizontally anisotropic

shapes typically found in animals reared in standard laboratory cages. The shift occurred

in both sighted and early blind enriched animals, but the effect was more pronounced in

early blind animals.

551   The results of receptive field shape analyses pointed to differential effects of

rearing environment on tuning along horizontal and vertical axes of receptive fields. We

quantified neuronal tuning width separately along whisker rows and arcs for each

experimental group (**Figure 5E**, **Table 3**, **Table 5-1**). Both sighted and early blind

enriched animals showed sharper tuning along both the vertical and horizontal axes

relative to sighted controls, but there was a striking difference in the magnitude of tuning

width changes along the horizontal vs. vertical axes.

This constitutes an amplification of shifts in receptive field anisotropies previously

described in early blind animals under standard rearing conditions (Ramamurthy and

Krubitzer, 2018). Along the arc (vertical) axis, tuning was sharpened to the same extent

in the enriched groups as seen in standard-reared early blind animals relative to sighted

controls (**Figure 5E**; %BW response for in-arc whiskers: eEB: 19.4 ± 2.80%, eSC: 18.7 ±

2.13%, sEB: 20.1 ± 1.58%, sSC: 26.7 ± 2.50%; linear mixed-effects model, eEB vs. sSC:

$p<0.05$, eEB vs. sEB: $p>0.05$, eSC vs. sSC: $p<0.05$, eSC vs. sEB: $p>0.05$). But along the

row (horizontal axis), both sighted and early blind enriched animals showed sharper

tuning relative to both groups of standard-reared animals (**Figure 5E**; %BW response for

in-row whiskers: eEB: 13.8 ± 2.07%, eSC: 20.9 ± 1.96%, sEB: 35.1 ± 2.82%, sSC: 49.1

± 2.85%; linear mixed-effects model, eEB vs. sSC: $p<0.001$, eEB vs. sEB: $p<0.001$, eSC

vs. sSC: $p<0.001$, eSC vs. sEB: $p<0.001$). There was a trend for sharper row tuning in

enriched early blind animals compared to enriched sighted control animals, but this was

not significant (linear mixed-effects model; eSC vs. eEB: $p>0.05$). We computed the

whisker selectivity index as the normalized difference ratio of the neuronal response to

the best whisker in comparison to responses to in-row or in-arc whiskers. Population

distributions of the selectivity index for S1 neurons (**Figure 5F**, **Table 3**, **Table 5-1**) further

validate the anisotropy in the enhancement of selectivity seen for animals reared in

enriched environments (BW vs. in-arc whiskers: linear mixed-effects model, eEB vs. sSC:

$p<0.05$, eEB vs. sEB: $p>0.05$, eSC vs. sSC: $p<0.05$, eSC vs. sEB: $p>0.05$, eEB vs. eSC:

p>0.05; BW vs. in-row whiskers: linear mixed-effects model, eEB vs. sSC: p<0.001, eEB

vs. sEB: p<0.001, eSC vs. sSC: p<0.001, eSC vs. sEB: p<0.001, eEB vs. eSC: p>0.05).

Thus, a spatially complex enriched environment drives significantly greater

improvement in neuronal selectivity for whisker touch in early blind animals especially

along the rostrocaudal axis, which is the primary axis of natural whisking behavior. A

similar improvement in selectivity was observed in sighted control littermates reared in

the enriched environment, supporting the role of tactile experience rather than vision loss

itself in driving these effects. In summary, the results of our current study indicate that

compensatory effects on neural coding in S1 following early blindness do not arise solely

from visual deprivation and can be amplified and directed by tactile experience.

DISCUSSION

Although the effects of enrichment on plasticity within a sensory modality have

been studied at structural, functional, and molecular levels, few studies have examined

its impact on cross-modal compensatory plasticity, focusing mainly on c-fos expression

or synaptic changes (Piche et al., 2004; Mundiñano and Martínez-Millán, 2010; Zheng et

al., 2014). The short-tailed opossum is born with a highly immature nervous system,

enabling precise extrauterine manipulations at very early stages in development

(Ramamurthy and Krubitzer, 2018; Ramamurthy et al., 2021). Here, we tested the effects

of rearing sighted and early blind short-tailed opossums in a large and dynamic, spatially

complex 3D environment, and examined effects on tactile behavior and receptive field

configurations of neurons in S1.

Enrichment reduces abnormal stereotypic behaviors (e.g. overgrooming) and

other behavioral indicators of stress and aggression, and enhances learning, memory,

motor performance and social behavior (Dijkhuizen et al., 2024; Ratuski et al., 2024; Cait

et al., 2024; Nakhwa et al., 2024; Caires & Bossolani-Martins, 2023; Mieske et al., 2022;

Kentner et al., 2021; Khoo et al., 2020; Bailoo et al., 2018; Girbovan & Plamondon, 2013;

Amaral et al., 2008; Mason et al., 2007). It can also improve sensory function and

performance, including whisker touch discrimination (Zheng et al., 2021). In our study, we

implemented enrichment intended to increase the use of whisker touch. Since whiskers

are critical for guiding forelimb placement in crossing gaps when small mammals climb

(Ahl, 1986; Arkley et al., 2014; Niederschuh et al., 2015; Arkley et al., 2017; Grant et al.,

2018), we reared opossums in cages with periodically varying, spatially complex climbing

substrates and intermittently introduced novel toys, since novel objects and other

interactive enrichment devices promote active whisking behavior (Nakhwa et al., 2024;

Sofroniew & Svoboda, 2015). We show that in a novel setting, both sighted and blind

opossums showed differential exploration patterns reflecting the geometry of their rearing

environments. Additionally, enriched opossums are faster and more successful in gap

crossing using a whisker-dependent strategy. Thus, opossums use behavioral strategies

shaped by their rearing environments, and our enrichment approach effectively engages

and enhances naturalistic whisker-mediated behavior in both sighted and blind animals,

enabling us to evaluate the effects of enriched rearing on compensatory plasticity

following early vision loss.

It has been previously reported that early blindness alters the coding of whisker

touch inputs in opossums, in a manner that supports enhanced tactile discrimination

(Ramamurthy and Krubitzer, 2018; Ramamurthy et al., 2021). This involved suppression

of whisker-evoked responses combined with increased spatial selectivity, due to the

greater relative suppression of neural responses to non-preferred stimuli (SW) vs. the

preferred stimulus (BW). These alterations in the receptive fields of neurons in S1

resemble the effects of naturalistic environmental enrichment on receptive fields of

neurons in S1 of adult rats (Polley et al., 1999, 2004). Here we demonstrate that rearing

early blind short-tailed opossums in an enriched environment further amplifies these

changes. Since selectivity to whisker stimuli was enhanced not just in early blind animals

but also in sighted control animals reared in enriched conditions, we can infer that

suppression of whisker-evoked responses and increased selectivity of neural responses

in the S1 whisker representation are effects of tactile experience rather than visual

deprivation itself.

A key finding was the effect of the rearing environment on the shapes of receptive

fields of S1 neurons – shape distributions for both sighted and early blind animals

exhibited a prominent shift away from the horizontally anisotropic shapes that dominate

in standard-reared opossums. This was accompanied by a reduction in tuning width along

row (horizontal) vs. arc (vertical) axes in enriched animals – on average, tuning widths

were reduced to a greater extent along the row axis, parallel to the main axis of whisker

movements. This may reflect more naturalistic patterns of whisker use in short-tailed

opossums, enabled by the increase in vertical space and availability of climbing

substrates in the enriched rearing environment. In rats, simulations that modelled

exploratory behavior predicted changes in the spatial patterns of whisker contact

depending on the environment being explored (Hobbs et al., 2015) – whiskers within the

same horizontal row were more likely to come in simultaneous contact during exploration

of flat surfaces, but in more complex and naturalistic environments, the probability

distributions of simultaneous whisker contact shifted away from this row-wise structure.

Such alterations in the patterns of tactile stimuli impinging on the whisker array over the

course of development may, in turn, impact receptive field structure.

Another possibility is that anisotropic effects on receptive fields are inherent to

differences in row vs. arc organization within the neocortex. In rats and mice, it is well-

documented that there is a bias towards preferential connectivity among barrels

corresponding to whiskers in the same row (Bernardo et al., 1990; Hoeflinger et al., 1995;

Kim and Ebner, 1999; Keller and Carlson, 1999), and neurons in barrel cortex typically

display patterns of activity that tend to be elongated along the row axis (Armstrong-James

and Fox, 1987; Simons, 1978; Kleinfeld and Delaney, 1996; Ego-Stengel et al., 2005; Le

Cam et al., 2011). Further, there is evidence for differential effects on responses of S1

neurons along row and arc axes when sensory inputs are altered within the whisker

system, for example, through chronic stimulation (Quairiaux et al., 2007) or selective

trimming (Dubroff et al., 2005; Wallace and Sakmann, 2008) of specific whiskers. Thus,

biases in the functional organization and connectivity within the S1 whisker representation

may favor plasticity, preferentially sharpening tuning along the row axis.

A long-standing question has been whether reorganization following sensory loss

is primarily driven by deprivation of the lost modality or due to increased experience using

the spared senses. Studies in blind humans with Braille reading experience support an

important role for tactile experience in generating tactile spatial acuity enhancement

following blindness (Wong et al., 2011). The relative contributions of deprivation of the

lost modality versus increased or modified use of the spared senses (Goldreich and

Kanics, 2003; Wong et al., 2011; Voss, 2011) to different aspects of cortical

reorganization following sensory loss are still not well understood. Our study provides

clear evidence for the role of experience in driving compensatory neural and behavioral

plasticity following sensory loss, highlighting the importance of therapeutic strategies

engaging tactile exploration (for example through play-based learning of tactile skills,

Hilditch 2024; or the use of haptic devices, Fleck et al., 2025) to maximize neural and

behavioral compensation through touch.

REFERENCES

Abreu MSL, Oliveira LRD (2014) Patterns of arboreal and terrestrial space use by non-

volant small mammals in an Araucaria forest of southern Brazil. Anais da

Academia Brasileira de Ciências 86:807-819.

Ahl A (1986) The role of vibrissae in behavior: a status review. Veterinary research

communications 10:245-268.

Alwis DS, Rajan R (2013) Environmental enrichment causes a global potentiation of

neuronal responses across stimulus complexity and lamina of sensory cortex.

Frontiers in cellular neuroscience 7:124.

Alwis DS, Rajan R (2014) Environmental enrichment and the sensory brain: the role of

enrichment in remediating brain injury. Frontiers in systems neuroscience 8:156.

Antón-Bolaños N, Sempere-Ferràndez A, Guillamón-Vivancos T, Martini FJ, Pérez-Saiz

L, Gezelius H, Filipchuk A, Valdeolmillos M, López-Bendito G (2019) Prenatal

activity from thalamic neurons governs the emergence of functional cortical maps

in mice. Science 364:987-990.

Amaral OB, Vargas RS, Hansel G, Izquierdo I, Souza DO (2008) Duration of

environmental enrichment influences the magnitude and persistence of its

behavioral effects on mice. Physiology & Behavior 93:388-394.

Arkley K, Grant RA, Mitchinson B, Prescott TJ (2014) Strategy change in vibrissal active

sensing during rat locomotion. Current biology : CB 24:1507-1512.

Arkley K, Tiktak GP, Breakell V, Prescott TJ, Grant RA (2017) Whisker touch guides

canopy exploration in a nocturnal, arboreal rodent, the Hazel dormouse

(Muscardinus avellanarius). Journal of comparative physiology A, Neuroethology,

sensory, neural, and behavioral physiology 203:133-142.

Bailoo JD, Murphy E, Boada-Saña M, Varholick JA, Hintze S, Baussière C, Hahn KC,

Göpfert C, Palme R, Voelkl B, Würbel H (2018) Effects of Cage Enrichment on

Behavior, Welfare and Outcome Variability in Female Mice. Front Behav Neurosci

12:232.

Bartoletti A, Medini P, Berardi N, Maffei L (2004) Environmental enrichment prevents

effects of dark-rearing in the rat visual cortex. Nature neuroscience 7:215-216.

Beddard FE (1902) Observations upon the Carpal Vibrissæ in Mammals. Proceedings of

the Zoological Society of London 72:127-136.

Bengoetxea H, Ortuzar N, Bulnes S, Rico-Barrio I, Lafuente JV, Argandona EG (2012)

Enriched and deprived sensory experience induces structural changes and rewires

connectivity during the postnatal development of the brain. Neural plasticity 2012.

Branchi I, D'Andrea I, Santarelli S, Bonsignore LT, Alleva E (2011) The richness of social

stimuli shapes developmental trajectories: Are laboratory mouse pups

713   impoverished? Progress in neuro-psychopharmacology & biological psychiatry

714   35:1452-1460.

715   Brecht M, Roth A, Sakmann B (2003) Dynamic receptive fields of reconstructed pyramidal

716   cells in layers 3 and 2 of rat somatosensory barrel cortex. J Physiol 553:243–265.

717   Caires CRS, Bossolani-Martins AL (2023) Which form of environmental enrichment is

718   most effective in rodent models of autism? Behavioural Processes 211:1-8.

719   Cait J, Winder CB, Mason GJ (2024) How much 'enrichment' is enough for laboratory

720   rodents? A systematic review and meta-analysis re-assessing the impact of well-

721   resourced cages on morbidity and mortality. Applied Animal Behaviour Science

722   278:106361.

723   Cancedda L, Putignano E, Sale A, Viegi A, Berardi N, Maffei L (2004) Acceleration of

724   visual system development by environmental enrichment. The Journal of

725   neuroscience : the official journal of the Society for Neuroscience 24:4840-4848.

726   Coq JO, Xerri C (1998) Environmental enrichment alters organizational features of the

727   forepaw representation in the primary somatosensory cortex of adult rats.

728   Experimental brain research 121:191-204.

729   Delciellos AC, Vieira MV (2006) Arboreal walking performance in seven didelphid

730   marsupials as an aspect of their fundamental niche. Austral Ecology 31:449-457.

731   Devonshire IM, Dommett EJ, Grandy TH, Halliday AC, Greenfield SA (2010)

732   Environmental enrichment differentially modifies specific components of sensory-

733   evoked activity in rat barrel cortex as revealed by simultaneous

734   electrophysiological recordings and optical imaging in vivo. Neuroscience

735   170:662-669.

Diamond ME, Toso A (2023) Tactile cognition in rodents. Neuroscience & Biobehavioral

Reviews 149:105161.

Dijkhuizen S, Van Ginneken LMC, Ijpelaar AHC, Koekkoek SKE, De Zeeuw CI, Boele HJ

(2024) Impact of enriched environment on motor performance and learning in

mice. Scientific Reports 14:5962.

Dooley JC, Franca JG, Seelke AM, Cooke DF, Krubitzer LA (2013) A connection to the

past: *Monodelphis domestica* provides insight into the organization and

connectivity of the brains of early mammals. J Comp Neurol 521:3877-3897.

Dooley JC, Krubitzer LA (2019) Alterations in cortical and thalamic connections of

somatosensory cortex following early loss of vision. J Comp Neurol 527(10):1675-

1688.

Ego-Stengel V, e Souza TM, Jacob V, Shulz DE (2005) Spatiotemporal characteristics of

neuronal sensory integration in the barrel cortex of the rat. J Neurophysiol

93:1450–1467.

Engineer ND, Percaccio CR, Pandya PK, Moucha R, Rathbun DL, Kilgard MP (2004)

Environmental Enrichment Improves Response Strength, Threshold, Selectivity,

and Latency of Auditory Cortex Neurons. Journal of Neurophysiology 92:73-82.

Fleck JJ, Zook ZA, Clark JP, Preston DJ, Lipomi DJ, Pacchierotti C, O'Malley MK (2025)

Wearable multi-sensory haptic devices. Nature Reviews Bioengineering.

Fox K (2003) The origin of cortical surround receptive fields studied in the barrel cortex.

Nat Neurosci 6:631–640. Frostig RD (2006) Functional organization and plasticity

in the adult rat barrel cortex: moving out-of-the-box. Current opinion in

neurobiology 16:445-450.

Gallyas F (1979) Silver staining of myelin by means of physical development.
Neurological research 1:203-209.

Girbovan C, Plamondon H (2013) Environmental enrichment in female rodents:
considerations in the effects on behavior and biochemical markers. Behav Brain
Res 253:178-190.

Goin FJ, Woodburne MO, Zimicz AN, Martin GM, Chornogubsky L (2016) South
American Living Metatherians: Physiological Ecology and Constraints. In: A Brief
History of South American Metatherians: Evolutionary Contexts and
Intercontinental Dispersals, pp 37-75. Dordrecht: Springer Netherlands.

Goldreich D, Kanics IM (2003) Tactile acuity is enhanced in blindness. J Neurosci
23:3439-3445.

Gomez-Pinilla F, Ying Z, Agoncillo T, Frostig R (2011) The influence of naturalistic
experience on plasticity markers in somatosensory cortex and hippocampus:
effects of whisker use. Brain research 1388:39-47.

Graham M, Socha JJ (2020) Going the distance: The biomechanics of gap-crossing
behaviors. J Exp Zool A Ecol Integr Physiol 333:60-73.

Grant RA, Breakell V, Prescott TJ (2018) Whisker touch sensing guides locomotion in
small, quadrupedal mammals. Proceedings Biological sciences 285.

Grant RA, Delaunay MG, Haidarliu S (2017) Mystacial whisker layout and musculature in
the guinea pig (Cavia porcellus): a social, diurnal mammal. The Anatomical Record
300:527-536.

Greifzu F, Pielecka-Fortuna J, Kalogeraki E, Krempler K, Favaro PD, Schluter OM, Lowel
S (2014) Environmental enrichment extends ocular dominance plasticity into

adulthood and protects from stroke-induced impairments of plasticity. Proceedings

of the National Academy of Sciences of the United States of America 111:1150-

1155.

Guic E, Carrasco X, Rodriguez E, Robles I, Merzenich MM (2008) Plasticity in primary

somatosensory cortex resulting from environmentally enriched stimulation and

sensory discrimination training. Biological research 41:425-437.

Hilditch J (2021) Pre-braille Implementation into Early Education: Tactile Activities to

Introduce Braille Concepts to Kindergartners. In: Handbook of Disability: Critical

Thought and Social Change in a Globalizing World (Rioux MH, Buettgen A, Zubrow

E, Viera J, eds), pp 1-31. Singapore: Springer Nature Singapore.

Hutson K, Masterton R (1986) The sensory contribution of a single vibrissa's cortical

barrel. Journal of neurophysiology 56:1196-1223.

Kentner AC, Speno AV, Doucette J, Roderick RC (2021) The Contribution of

Environmental Enrichment to Phenotypic Variation in Mice and Rats. eneuro

8:ENEURO.0539-0520.2021.

Khoo SY-S, Correia V, Uhrig A (2020) Nesting material enrichment reduces severity of

overgrooming-related self-injury in individually housed rats. Laboratory Animals

54:546-558.

Mason G, Clubb R, Latham N, Vickery S (2007) Why and how should we use

environmental enrichment to tackle stereotypic behaviour? Applied Animal

Behaviour Science 102:163-188.

Mieske P, Hobbiesiefken U, Fischer-Tenhagen C, Heinl C, Hohlbaum K, Kahnau P, Meier

J, Wilzopolski J, Butzke D, Rudeck J, Lewejohann L, Diederich K (2022) Bored at

home?—A systematic review on the effect of environmental enrichment on the welfare of laboratory rats and mice. Frontiers in Veterinary Science 9.

Molnár Z, Blakemore C (1995) How do thalamic axons find their way to the cortex? Trends Neurosci 18:389–397. 10.1016/0166-2236(95)93935-Q [DOI] [PubMed] [Google Scholar]

Molnár Z, Knott GW, Blakemore C, Saunders NR (1998) Development of thalamocortical projections in the south american gray short-tailed opossum (Monodelphis domestica). J Comp Neurol 398:491–514. 10.1002/(SICI)1096-9861(19980907)398:4%3C491::AID-CNE3%3E3.0.CO;2-Y

Nakhwa AA, Breakell V, Chambers L, Elder A, McLoughlin E, Roberts S, Todd K, Grant RA (2024) Implications of tactile enrichment on the behaviour and whisker movements of four species of carnivorans. Applied Animal Behaviour Science 274:106261.

Jenkins Jr., FA (1974) Tree shrew locomotion and the origins of primate arborealism. In: Jenkins, F.A. (Ed.), Primate Locomotion. Academic Press, New York, pp. 85–115.

Jiang C, Xu X, Yu L, Xu J, Zhang J (2015) Environmental enrichment rescues the degraded auditory temporal resolution of cortical neurons induced by early noise exposure. The European journal of neuroscience 42:2144-2154.

Jones M, Archer M, Dickman C (2003) Predators with Pouches: The Biology of Carnivorous Marsupials: CSIRO PUBLISHING.

Kalogeraki E, Pielecka-Fortuna J, Lowel S (2017) Environmental enrichment accelerates ocular dominance plasticity in mouse visual cortex whereas transfer to standard cages resulted in a rapid loss of increased plasticity. PloS one 12:e0186999.

Ladine TA, Kissell RE (1994) Escape Behavior of Virginia Opossums. The American

Midland Naturalist 132:234-238.

Landers MS, Knott GW, Lipp HP, Poletaeva I, Welker E (2011) Synapse formation in adult

barrel cortex following naturalistic environmental enrichment. Neuroscience

199:143-152.

Lammers AR (2004) The biodynamics of arboreal locomotion in the gray short-tailed

opossum (*Monodelphis domestica*).

Lammers AR (2007) Locomotor kinetics on sloped arboreal and terrestrial substrates in

a small quadrupedal mammal. Zoology (Jena, Germany) 110:93-103.

Lammers AR (2009) The effects of substrate texture on the mechanics of quadrupedal

arboreal locomotion in the gray short-tailed opossum (*Monodelphis domestica*).

Journal of experimental zoology Part A, Ecological genetics and physiology

311:813-823.

Lammers AR, Biknevicius AR (2004) The biodynamics of arboreal locomotion: the effects

of substrate diameter on locomotor kinetics in the gray short-tailed opossum

(*Monodelphis domestica*). The Journal of experimental biology 207:4325-4336.

Lammers AR, Gauntner T (2008) Mechanics of torque generation during quadrupedal

arboreal locomotion. Journal of Biomechanics 41:2388-2395.

Lammers AR, Earls KD, Biknevicius AR (2006) Locomotor kinetics and kinematics on

inclines and declines in the gray short-tailed opossum *Monodelphis domestica*.

The Journal of experimental biology 209:4154-4166.

Lazzouni L, Lepore F (2014) Compensatory plasticity: time matters. Frontiers in human

neuroscience 8:340.

Leah J, Allardyce H, Cummins R (1985) Evoked cortical potential correlates of rearing

environment in rats. Biological Psychology 20:21-29.

Lee H-K (2023) Metaplasticity framework for cross-modal synaptic plasticity in adults.

Frontiers in Synaptic Neuroscience 14.

LeMessurier AM, Laboy-Juárez KJ, McClain K, Chen S, Nguyen T, Feldman DE (2019)

Enrichment drives emergence of functional columns and improves sensory coding

in the whisker map in L2/3 of mouse S1. eLife 8:e46321.

Levine JN, Chen H, Gu Y, Cang J (2017) Environmental Enrichment Rescues Binocular

Matching of Orientation Preference in the Mouse Visual Cortex. The Journal of

neuroscience : the official journal of the Society for Neuroscience 37:5822-5833.

López-Bendito G, Aníbal-Martínez M, Martini FJ (2022) Cross-Modal Plasticity in Brains

Deprived of Visual Input Before Vision. Annu Rev Neurosci 45:471-489.

Lyne A (1958) A thesis on the vibrissae and pelage hairs on the marsupialia and

supporting papers.

Lyne A (1959) The systematic and adaptive significance of the vibrissae in the

Marsupialia. Journal of Zoology 133:79-133.

Montgomery, WI (1980) The use of arboreal runways by the woodland rodents,

Apodemus sylvaticus (L.), A. flavicollis (Melchior) and Clethrionomys glareolus

(Schreber). Mammal Review 10:189-195.

Macrini TE (2004) *Monodelphis domestica*. Mammalian Species:1-8.

Megevand P, Troncoso E, Quairiaux C, Muller D, Michel CM, Kiss JZ (2009) Long-term

plasticity in mouse sensorimotor circuits after rhythmic whisker stimulation. The

Journal of neuroscience : the official journal of the Society for Neuroscience

29:5326-5335.

Mezzera C, López-Bendito G (2016) Cross-modal plasticity in sensory deprived animal

models: From the thalamocortical development point of view. Journal of chemical

neuroanatomy 75:32-40.

Morland AB, Brown HD & Baseler HA (2021) Cortical reorganization: reallocated

responses without rewiring. Current Biology, 31(2), R76-R78.

Muchlinski MN, Durham EL, Smith TD, Burrows AM (2013) Comparative histomorphology

of intrinsic vibrissa musculature among primates: implications for the evolution of

sensory ecology and "face touch". American journal of physical anthropology

150:301-312.

Mundiñano IC, Martínez-Millán L (2010) Somatosensory cross-modal plasticity in the

superior colliculus of visually deafferented rats. Neuroscience 165:1457-1470.

Nakashima AS, Dyck RH (2008) Enhanced plasticity in zincergic, cortical circuits after

exposure to enriched environments. The Journal of neuroscience : the official

journal of the Society for Neuroscience 28:13995-13999.

Niederschuh SJ, Witte H, Schmidt M (2015) The role of vibrissal sensing in forelimb

position control during travelling locomotion in the rat (Rattus norvegicus,

Rodentia). Zoology (Jena, Germany) 118:51-62.

Nithianantharajah J, Hannan AJ (2006) Enriched environments, experience-dependent

plasticity and disorders of the nervous system. Nature reviews Neuroscience

7:697-709.

Piche M, Robert S, Miceli D, Bronchti G (2004) Environmental enrichment enhances
auditory takeover of the occipital cortex in anophthalmic mice. The European
journal of neuroscience 20:3463-3472.

Pocock R (1914) 48. On the Facial Vibrissæ of Mammalia. Journal of Zoology 84:889-
912.

Polley DB, Chen-Bee CH, Frostig RD (1999) Two directions of plasticity in the sensory-
deprived adult cortex. Neuron 24:623-637.

Polley DB, Kvasnak E, Frostig RD (2004) Naturalistic experience transforms sensory
maps in the adult cortex of caged animals. Nature 429:67-71.

Ramamurthy DL, & Krubitzer LA (2016) The evolution of whisker-mediated
somatosensation in mammals: Sensory processing in barrelless S1 cortex of a
marsupial, *Monodelphis domestica*. Journal of Comparative Neurology 524(17),
3587-3613.

Ramamurthy DL, Krubitzer LA (2018) Neural Coding of Whisker-Mediated Touch in
Primary Somatosensory Cortex Is Altered Following Early Blindness. The Journal
of Neuroscience 38:6172-6189.

Ramamurthy DL, Dodson HK, Krubitzer LA (2021) Developmental plasticity of texture
discrimination following early vision loss in the marsupial *Monodelphis domestica*.
Journal of Experimental Biology, 224(9), jeb236646.

Ratuski AS, Améndola L, Makowska IJ, Weary DM (2024) Effects of temporary access to
environmental enrichment on measures of laboratory mouse welfare. Sci Rep
14:15143.

Romero-Ferrero F, Bergomi MG, Hinz RC, Heras FJH, de Polavieja GG (2019)
idtracker.ai: tracking all individuals in small or large collectives of unmarked
animals. Nature Methods 16:179-182.

Rupert JE, Schmidt EC, Moreira-Soto A, Herrera BR, Vandeberg JL, Butcher MT (2014)
Myosin isoform expression in the prehensile tails of didelphid marsupials:
functional differences between arboreal and terrestrial opossums. Anatomical
record (Hoboken, NJ : 2007) 297:1364-1376.

Sale A, Vetencourt JFM, Medini P, Cenni MC, Baroncelli L, De Pasquale R, Maffei L
(2007) Environmental enrichment in adulthood promotes amblyopia recovery
through a reduction of intracortical inhibition. Nature neuroscience 10:679.

Sale A, Berardi N, Maffei L (2009) Enrich the environment to empower the brain. Trends
in neurosciences 32:233-239.

Schmitt D, Gruss LT, Lemelin P (2010) Brief communication: forelimb compliance in
arboreal and terrestrial opossums. American journal of physical anthropology
141:142-146.

Seo ML (1992) Effect of environmental complexity on the latency of cortical vibrissa
potentials. Developmental psychobiology 25:67-76.

Shapiro LJ, Young JW, VandeBerg JL (2014) Body size and the small branch niche: Using
marsupial ontogeny to model primate locomotor evolution. Journal of Human
Evolution 68:14-31.

Sofroniew NJ, Svoboda K (2015) Whisking. Current Biology 25:R137-R140.

Taylor JS, Guillery RW (1994) Early development of the optic chiasm in the gray short-
tailed opossum, Monodelphis domestica. J Comp Neurol 350:109–121.
10.1002/cne.903500108

Teichert M, Isstas M, Wenig S, Setz C, Lehmann K, Bolz J (2018) Cross-modal refinement
of visual performance after brief somatosensory deprivation in adult mice.
European Journal of Neuroscience 47:184-191.

Thomas DR, Chadwell BA, Walker GR, Budde JE, VandeBerg JL, Butcher MT (2017)
Ontogeny of myosin isoform expression and prehensile function in the tail of the
gray short-tailed opossum (*Monodelphis domestica*). Journal of applied physiology
(Bethesda, Md : 1985) 123:513-525.

van Praag H, Kempermann G, Gage FH (2000) Neural consequences of environmental
enrichment. Nature reviews Neuroscience 1:191-198.

Voss P (2011) Superior Tactile Abilities in the Blind: Is Blindness Required? The Journal
of Neuroscience 31:11745-11747.

Voss P (2019) Brain (re)organization following visual loss. WIREs Cognitive Science
10:e1468.

Wang BS, Feng L, Liu M, Liu X, Cang J (2013) Environmental enrichment rescues
binocular matching of orientation preference in mice that have a precocious critical
period. Neuron 80:198-209.

Wang M, Yu X (2023) Experience-dependent structural plasticity of pyramidal neurons in
the developing sensory cortices. Current Opinion in Neurobiology 81:102724.

Weissbrod A, Shapiro A, Vasserman G, Edry L, Dayan M, Yitzhaky A, Hertzberg L,
Feinerman O, Kimchi T (2013) Automated long-term tracking and social

behavioural phenotyping of animal colonies within a semi-natural environment.

Nature communications 4:2018.

Wilkinson M, Stirton C, McConnachie A (2010) Behavioural observations of singly-housed

grey short-tailed opossums (*Monodelphis domestica*) in standard and enriched

environments. Laboratory animals 44:364-369.

Wong M, Gnanakumaran V, Goldreich D (2011) Tactile spatial acuity enhancement in

blindness: evidence for experience-dependent mechanisms. J Neurosci 31:7028-

7037.

Zheng HJV, Meagher JP, Xu D, Patel YA, O'Connor DH, Kwon HB (2021) Environmental

Enrichment Sharpens Sensory Acuity by Enhancing Information Coding in Barrel

Cortex and Premotor Cortex. eNeuro 8.

Zheng J-J, Li S-J, Zhang X-D, Miao W-Y, Zhang D, Yao H, Yu X (2014) Oxytocin mediates

early experience–dependent cross-modal plasticity in the sensory cortices. Nature

neuroscience 17:391.

Zhu X, Wang F, Hu H, Sun X, Kilgard MP, Merzenich MM, Zhou X (2014) Environmental

acoustic enrichment promotes recovery from developmentally degraded auditory

cortical processing. The Journal of neuroscience : the official journal of the Society

FIGURE LEGENDS

Figure 1. Timeline of experimental manipulations relative to developmental milestones in

short-tailed opossums. The four experimental groups are indicated: sSC (standard-reared

sighted control), sEB (standard-reared early blind), eSC (enriched sighted), and eEB

(enriched early blind). Behavioral and electrophysiological experiments were conducted

in adults following the sequence shown in the inset, but ages of adult animals were not

strictly matched by postnatal day.

Figure 2. Enriched rearing alters spatial patterns of exploratory behavior. A.

Example movement trajectories from one animal in each experimental group (sEB: light

red, eEB: dark red, sSC: light blue, eSC: dark blue) tracked over five minutes during

behavioral testing in an enriched arena. **B.** Overlaid movement trajectories from all

animals in each experimental group in the arena test show a pattern of increased

exploration of vertical vs. horizontal space in opossums reared in enriched cages

compared to those reared in standard cages. Plus symbols indicate the center of mass

of movement trajectories for each group (colors as in **A**), with data for animals combined

within each rearing condition shown in black. **C.** Vertical occupancy profiles during arena

exploration. Normalized distributions of vertical position (height above the arena floor)

during exploration derived from 1280×720 px video frames (20 px bins used for

visualization), pooled across animals (black: enriched, gray: standard; dark gray denotes

regions of overlap between the two distributions. **D.** Ethograms showing the temporal

structure of locomotor activity and spontaneous crossing events during arena exploration.

Horizontal bars indicate periods of activity for representative trials from standard-reared

(top, gray) and enriched-reared (bottom, black) animals. Vertical tick marks denote

spontaneous crossings between surfaces. Time is plotted relative to trial start (mm:ss).

E-G. Summary behavioral metrics during arena exploration in standard and enriched

animals. Distributions of vertical preference index (**E**), fraction of trial time spent active

(**F**), and median crossing duration (**G**). Violin plots show Gaussian kernel density

estimates of the pooled distribution for each category; horizontal lines denote medians.

Points represent individual animals, color-coded by experimental group. Animals reared

in the enriched environment show significantly a signficant increase in vertical prefernce

index. Overall activity levels and duration of crossing events during free exploration were

not signficiantly different.

Figure 3. Enriched rearing alters sensorimotor performance during gap crossing.

A. Experimental setup in the gap crossing task. **B.** Contingency tables showing the

percentage of successful and failed gap crossing trials (0% = white, 100% = black;

intermediate percentages shown in grayscale) for small (2–10 cm) and large (12–14 cm)

gaps under intact and whisker-trimmed conditions, shown separately for each

experimental group (sSC, sEB, eSC, eEB). **C.** Number of gap crossings per minute

(across all trials) for large and small gaps (with intact or trimmed whiskers). Gap crossing

was whisker-dependent in all groups. Data for large gaps after whisker trimming are not

shown, as no successful crossings occurred in this condition. **D.** Duration of successful

crossings for large and small gaps (with intact or trimmed whiskers). Most groups show

significantly longer durations for succesful gap crossing on large gaps (12-14 cm) vs.

small gaps (2-10 cm). Upon whisker trimming, most groups show a significant increase

in gap crossing duration for small gaps, consistent with the use of whisker-dependent

strategies for gap crossing. Enriched blind animals were unique in showing no significant

difference between large and small gap durations and no trimming-related increase in

small gap crossing times, reflecting enhanced performance on large gaps through

whisker use alongside compensatory tactile strategies using other body parts that

preserve performance on small gaps when whiskers are absent. For larger gaps, there

are prominent differences between experimental groups based on rearing condition and

blindness. Early blind animals crossed gaps faster than sighted controls in standard

conditions, and their performance was further enhanced by enriched rearing. Violin plots

(color-coded by experimental group) show Gaussian kernel density estimates of the data

distribution, with medians indicated by circular markers. See also **Table 2**.

Figure 4. Enriched rearing amplifies alterations in S1 neuron responses to whisker

touch in early blind animals and produces similar shifts in sighted controls. A.

Spike raster plots from example neurons responsive to whisker touch in each of the four

experimental groups (left to right: sSC, sEB, eSC, eEB). Examples are shown for neurons

with the same best whisker (BW), C2. Trials corresponding to A, B, C, and D whisker

rows are separated by horizontal lines. Within each row, the direction of more rostral ('R')

versus more caudal ('C') whiskers is indicated on the y-axis. **B.** Peristimulus time

histograms (PSTHs) for the same neurons shown in **(A)**. PSTHs corresponding to

whiskers with response magnitudes >50% of the BW response are plotted in black, all

others are in gray. The examples in **(A)** and **(B)** illustrate more selective responses to

whisker touch in standard-reared blind opossums, which is further amplified in enriched

groups. Additionally, neural responses to whiskers vertically adjacent to the BW are

stronger than those for horizontally adjacent whiskers in enriched groups, especially in

enriched early blind opossums. **C.** Spontaneous firing rates of S1 neurons were not

significantly different across experimental groups. **D.** Firing rates evoked by the BW

showed significant differences across experimental groups. Notably, both sighted and

early blind enriched groups showed suppression relative to standard-reared sighted

controls, similar to effects previously reported in early blind animals under standard

rearing conditions. The magnitude of suppression of the BW-evoked response was

greatest in early blind animals reared in the enriched environment. **E.** Mean receptive

field size was significantly smaller in sEB, eSC and eEB animals relative to standard-

reared sighted controls. This is seen both in terms of number of whiskers driving

significant responses (left) and the rank-ordered tuning curve (right) for whisker-

responsive S1 neurons. See also **Table 3**.

Figure 5. Enriched rearing amplifies anisotropies in receptive fields of S1 neurons

in early blind animals, improving selectivity for whisker touch along a behaviorally

relevant axis. A. Schematic of the opossum mystacial whisker pad indicating whisker

rows (light yellow) and arcs (gold), and the horizontal and vertical axes used for receptive

field shape analyses. Insets in individual panels depict the grid of 16 whiskers and the

surrounding whisker (SW) positions included in each analysis, shown for one example

best whisker (BW) position (black filled circle). **B.** Mean two-dimensional somatotopic

receptive fields for each experimental group (left to right: sSC, sEB, eSC, eEB), shown

with 50%, 75%, and 90% contour lines (response levels relative to the BW, smoothed by

linear interpolation for display). This visualization highlights the degree of horizontal vs.

vertical spread (anisotropy) in the average receptive field configuration for each

experimental group. **C-D.** Receptive field shape distributions for S1 neurons in enriched

animals were shifted away from horizontal anisotropic shapes, which are typical of

standard-reared animals. **C** shows receptive field shape distributions when only 1° SW

positions were considered (left) and fractions of horizontally anisotropic receptive fields

corresponding to each group (right). Light gray bars indicate horizontally elongated

receptive fields defined as shape indices > 0, while black bars indicate horizontally

elongated receptive fields with a stricter cutoff of shape indices > 0.5. **D** shows data for

the same neurons when all SW positions (1-3°) were included. Horizontally anisotropic receptive fields are decreased in both enriched groups of animals, and to the greatest extent in enriched early blind animals. **E.** Quantification of tuning width along the row (horizontal/rostrocaudal axis) and arc (vertical axis) across experimental groups. Enriched groups show sharper tuning along the horizontal axis, to a significantly greater extent than standard-reared early blind animals. Enriched groups also show sharper tuning along the vertical axis compared with standard-reared sighted controls, but only to the same extent as previously reported in standard-reared early blind animals. **F.** Distributions of the whisker selectivity index of S1 neurons calculated separately for in-arc whiskers (left) and in-row whiskers (right), when including all SW positions (1°-3°). sEB, eSC and eEB groups all showed significantly greater selectivity for whiskers within arcs, but were not significantly different from each other. The greatest increases in whisker response selectivity were seen for whiskers within rows i.e. the horizontal/rostrocaudal axis, which is the primary axis of natural whisker motion. Selectivity was greatest for eEB animals followed by eSC and sEB animals, with all three of these groups showing significantly higher selectivity along the horizontal axis compared to sSC animals. See also **Table 3**.

MULTIMEDIA

Movie 1. A nesting mother opossum navigating the enriched home cage environment. Movie shown at 4x speed.

Movie 2. Sighted opossum in the arena test. Movie shown at 4x speed.

Movie 3. Early blind opossum in the arena test. Movie shown at 4x speed.

for Neuroscience 34:5406-5415.

TABLES

Table 1. Experimental Design. Breakdown of subject information for each experimental

group. Sexes of animals used for electrophysiological recording experiments, total

number of neurons recorded in S1 per group (ALL) and number of S1 neurons with

significant whisker-evoked responses (>2 SD) are listed.

Table 2. Statistical comparisons for key behavioral results. Exact p-values for fixed

effects of experimental group and/or test condition obtained from Poisson and Gaussian

generalized linear mixed-effects models, or from Fisher's exact tests for categorical

comparisons, with pairwise contrasts across experimental groups.

Table 3. Statistical comparisons for key electrophysiological results. Exact p-

values for fixed effects of experimental group from Gaussian generalized linear mixed-

effects models, with pairwise contrasts across experimental groups.

EXTENDED DATA FIGURE LEGENDS

Figure 1-1. Standard and enriched rearing environments used for short-tailed

opossums. A. Comparison of cages used in standard (left) and enriched (right) rearing

paradigms. Enriched cages (29"l x 18"w x 24"h) were substantially larger (~8x in volume)

than standard cages (8.5"l x 10"w x 8"h), especially in the vertical dimension (~3x). Each

enriched cage contained a nesting box positioned high above the ground, in addition to

the standard nesting cup on the cage floor. Cages also included various enrichment

objects: regularly repositioned manzanita branches, rotating sets of enrichment toys, a

running wheel, and social housing (see *Materials and Methods*). **B.** Frontal view of an

enriched cage. **C.** Top view of the same enriched cage shown in **(B).** Letters mark

individual components visible in both views: nesting box (a), nesting cup (b), food (c),

water (d), running wheel (e), manzanita branches (f–l), and enrichment toys (m). **D.**

Mother with an attached experimental litter after placement in the enriched rearing

environment. **E–F.** Early blind weanling climbing up a branch to reach the nesting box.

Figure 2-1. Behavioral movie analyses. A. Representative frame showing semi-

automated identification of an opossum during arena testing using idtracker.ai. **B.**

Representative frame showing behavioral tracking in the gap crossing task using

DeepLabCut. "IR1" and "IR2" indicate the positions of infrared motion sensors mounted

on the edge of each platform, tracked with DeepLabCut to aid in gap crossing analyses.

C. Zoomed-in view showing an opossum on the start platform extending its whiskers

toward the goal platform prior to crossing the gap.

Figure 3-1. Histological verification of recording sites. A, C, E, G. Myelin-stained

tangential sections of cortex from a standard-reared sighted (sSC) animal **(A)**, a standard-

reared early blind (sEB) animal **(C)**, an enriched sighted (eSC) animal **(E)**, and an

enriched early blind (eEB) animal **(G)**. Primary somatosensory cortex (S1) is identifiable

as a darkly staining region relative to adjacent cortical fields. **B, D, F, H.** Reconstructions

of myeloarchitectural borders drawn from the series of myelin-stained sections in sSC

(B), sEB **(D)**, eSC **(F)** and eEB **(H)** animals. The locations of fluorescent probe insertions

are shown as open circles, and recording sites are shown as black filled circles. Only sites

at which whisker-evoked neuronal responses were quantified are included. In all panels,

medial (M) and rostral (R) directions are indicated by arrows. Scale bar = 250 µm.

1141

EXTENDED DATA TABLES

Table 3-1. ANOVA marginal tests for fixed effects in Figure 3. F statistics, numerator and denominator degrees of freedom (DF1, DF2), and p-values are reported for each model term in analyses corresponding to **Figure 3**.

Table 4-1. ANOVA marginal tests for fixed effects in Figure 4. F statistics, numerator and denominator degrees of freedom (DF1, DF2), and p-values are reported for each model term in analyses corresponding to **Figure 4**.

Table 5-1. ANOVA marginal tests for fixed effects in Figure 5. F statistics, numerator and denominator degrees of freedom (DF1, DF2), and p-values are reported for each model term in analyses corresponding to **Figure 5**.

Figure 1

Figure 2

A Example trajectories
sEB eEB   sSC eSC

B Overlaid average trajectories
vertical zone
sEB   sSC
horizontal zone
standard   eSC eEB enriched

C Activity profile
standard
enriched
10%
% occupancy

D
standard
activity
crossings
enriched
activity
crossings
0:00   1:00   2:00   3:00   4:00   5:00
time from trial start (mm:ss)

E *
vertical preference index
standard enriched

F n.s.
fraction of trial active
standard enriched

G n.s.
median crossing duration (s)
standard enriched

Figure 3

A

start   gap   goal

B

	sSC	sEB	eSC	eEB

12-14 cm ***

	sSC		sEB		eSC		eEB	
Trim	0%	100%	0%	100%	0%	100%	0%	100%
Intact	8%	92%	14%	86%	17%	83%	14%	86%

2-10 cm

	sSC		sEB		eSC		eEB	
Trim	100%	0%	100%	0%	100%	0%	100%	0%
Intact	100%	0%	100%	0%	100%	0%	100%	0%

success   failure   success   failure   success   failure   success   failure

C

D

Figure 4

Figure 5

Table 1. Experimental Design.

		Visual Condition					
		Sighted (SC)			Early Blind (EB)		
		Sex	ALL	>2 SD	Sex	ALL	>2 SD
Rearing Condition	**Standard (s)**	5F, 5M	86	60	4F, 5M	88	64
	Enriched (e)	3F, 4M	84	58	3F, 4M	81	51

Breakdown of subject information for each experimental group. Sexes of animals used for electrophysiological recording experiments, total number of neurons recorded in S1 per group (ALL) and number of S1 neurons with significant whisker-evoked responses (>2 SD) are listed.

Table 2. Statistical comparisons for key behavioral results.

Result/ Figure Panel	Fixed Effect p-value	Pairwise comparisons p-values			
	Test Condition	sSC	sEB	eSC	eEB
% successful trials **Fig. 2B** Alpha with Bonferroni correction = 0.0125	-	-	-	-	-
(1) large vs. small (2) intact vs. trim		<2.2e-16	<2.2e-16	<2.2e-16	<2.2e-16
# Gap crossings/minute **Fig. 3C** (1) large vs. small	2.4277e-09	2.4277e-09	1.7407e-09	9.8418e-10	2.0062e-09
(2) intact vs. trim	0.5356	0.5356	0.37091	0.42035	0.15936
Gap crossing duration **Fig. 3D** (1) large vs. small	1.1011e-79	8.0737e-92	1.4663e-08	0.0128	0.4872
(2) intact vs. trim	2.888e-07	0.0081	9.6063e-09	7.2433e-05	0.1481

Result/ Figure Panel	Expt. Group	sSC vs.sEB	sSC vs. eSC	sSC vs. eEB	sEB vs. eEB	sEB vs. eSC	eSC vs. eEB
# Gap crossings/minute **Fig. 3C** (3) across groups (large, intact)	0.5927	-	-	-	-	-	-
# Gap crossings/minute **Fig. 3C** (3) across groups (small, intact)		-	-	-	-	-	-
# Gap crossings/minute **Fig. 3C** (3) across groups (small, trim)		-	-	-	-	-	-
Gap crossing duration **Fig. 3D** (3) across groups (large, intact)	6.8796e-51	7.7220e-35	3.6476e-62	3.3333e-72	3.2002e-09	3.8990e-04	0.0021
Gap crossing duration **Fig. 3D** (3) across groups (small, intact)		0.7771	0.0691	2.9099e-14	1.0844e-13	0.1031	7.5517e-05
Gap crossing duration **Fig. 3D** (3) across groups (small, trim)		1.3303e-05	5.6083e-19	6.8960e-15	6.6591e-04	8.6167e-06	0.3183

Exact p-values were obtained from Poisson and Gaussian generalized linear mixed-effects models, or from Fisher's exact tests for categorical comparisons, with pairwise contrasts across experimental groups as indicated (see Methods).

Table 3. Statistical comparisons for key electrophysiological results.

Result/ Figure Panel	Fixed Effect p-value	Pairwise comparisons p-values					
	Expt. Group	sSC vs.sEB	sSC vs. eSC	sSC vs. eEB	sEB vs. eEB	sEB vs. eSC	eSC vs. eEB
Spontaneous activity **Fig. 3C**	0.856	-	-	-	-	-	-
Whisker-evoked activity **Fig. 3D**	4.68e-10	4.53e-04	4.09e-04	1.11e-11	0.939	1.38e-04	3.04e-04
RF size: # of whiskers **Fig. 3E** (left)	1.58-07	9.43e-07	4.29e-06	8.86e-07	0.832	0.763	0.621
Rank-ordered whisker tuning curve **Fig. 3E** (right)	1.55e-25	3.271e-11	1.88e-22	4.04e-19	8.19e-04	0.006	0.618
Shape index: 1^{st} order adjacent whiskers **Fig. 4C** (left)	0.004	0.249	0.445	6.84e-04	0.628	0.023	0.0033
% horizontal: 1^{st} order adjacent whiskers **Fig. 4C** Alpha with Bonferroni correction = 0.008 (right)	-	0.266	0.185	0.001	0.039	0.855	0.083
Shape index: $1^{st} - 3^{rd}$ order adjacent whiskers **Fig. 4D** (left)	3.45e-05	0.093	0.013	2.19e-06	0.384	0.001	0.016
% horizontal: $1^{st} - 3^{rd}$ order adjacent whiskers **Fig. 4D** Alpha with Bonferroni correction = 0.008 (right)	-	0.516	9.88e-05	2.00e-06	2.50e-05	7.73e-04	0.340
% BW response: in-arc whiskers **Fig. 4E** (left)	0.042	0.0325	0.0116	0.0244	0.652	0.812	0.848
% BW response: in-row whiskers **Fig. 4E** (left)	8.51e-20	7.10e-05	7.56e-14	1.12e-18	6.19e-05	1.25e-08	0.056
BW vs. in-arc whiskers **Fig. 4F** (left)	0.043	0.080	0.014	0.015	0.436	0.419	0.957
BW vs. in-row whiskers **Fig. 4F** (right)	2.49e-18	2.86e-04	8.18e-12	7.00e-18	3.03e-04	8.31e-09	0.018

Exact p-values were obtained from Gaussian generalized linear mixed-effects models, with pairwise contrasts across experimental groups as indicated (see Methods).