Repetitive Neuronal Activation Regulates Cellular Maturation State via Nuclear Reprogramming

Tomoyuki Murano[1], Hideo Hagihara[1], Katsunori Tajinda[2], Keizo Takao[3,4], Yoshihiro Takamiya[1], Kaoru Katoh[5,6,7], Alfred J. Robison[8], Mitsuyuki Matsumoto[9], Masakazu Namihira[5], Tsuyoshi Miyakawa[1]*

[1]Division of Systems Medical Science, Center for Medical Science, Fujita Health University, Toyoake, Japan

[2]Astellas Research Institute of America, San Diego, CA, USA

[3]Department of Behavioral Physiology, Faculty of Medicine, University of Toyama, Toyama, Japan

[4]Research Center for Idling Brain Science, University of Toyama, Toyama, Japan

[5]Biomedical Research Institute, National Institute of Advanced Industrial Science and Technology (AIST), Tsukuba, Japan

[6]Artificial Intelligence Research Center, National Institute of Advanced Industrial Science and Technology (AIST), Tsukuba, Japan

[7]Ph.D. Program in Humanics, School of Integrative and Global Majors, University of Tsukuba, Tsukuba, Japan

[8]Department of Physiology and Neuroscience Program, Michigan State University, Michigan, USA

[9]Arialys Therapeutics Inc, La Jolla, CA, USA

*Correspondence:

Tsuyoshi Miyakawa

Division of Systems Medical Science, Institute for Comprehensive Medical Science, Fujita Health University, 1-98 Dengakugakubo, Kutsukake-cho, Toyoake, Aichi 470-1192, Japan

Telephone: +81-562-93-9376

Fax: +81-562-92-5382

E-mail: miyakawa@fujita-hu.ac.jp

Abstract

Neural stimulation, such as electroconvulsive therapy (ECT) and repetitive

transcranial magnetic stimulation (rTMS), are highly effective clinical interventions for a broad

spectrum of psychiatric disorders, including depression and schizophrenia. However, their

mechanism of action at the cellular level remains poorly understood. Here, we modelled ECT

with repeated optogenetic neuronal stimulation in the mouse dentate gyrus, and observed ECT-

like behavioral effects, including decreased depression-like behavior and increased locomotor

activity. At the cellular level, we found dematuration to a long-term stable state, persisting for

more than one month, defined by changes in nuclear structure, gene expression patterns

resembling the G_2/M phase of the cell cycle, and altered neural coding of navigational

information. Moreover, knockout of the G_2/M master regulator Cyclin B rescued some of

behavioral and cellular effects. These findings demonstrate that ECT-like brain stimulation

triggers plasticity of the cellular state, revealing a form of stimulus-regulated nuclear

reprogramming with potential clinical utility.

Introduction

In the adult mammalian brain, strong synchronous neural activity typically has toxic effects on neuronal function and viability. Cumulative electrical activity of sufficiently high amplitude, frequency, and/or duration interferes with cellular metabolism, flooding cells with excess calcium and resulting in metabolic failure and death[1,2]. High-intensity neural activity also results in maladaptive neuronal changes, such as dendritic spine loss[3,4], abnormal neurite arborization[4,5], and toxic excitability[6]. Previous studies have identified an intermediate state of senescence after overactivation, characterized by the dematuration of adult neurons to an immature-like state, including global de-differentiation of gene expression patterns, membrane excitability, and neural plasticity[7–11]. Similar neuronal phenotypes have been observed in genetically engineered mice that exhibit abnormal behaviors mirroring common neuropsychiatric disorders, such as schizophrenia, intellectual disability, and bipolar disorder[7,12–16]. Our recent study investigated the levels of lactate, a surrogate marker of neural excitation, in the brains mouse models relevant to neuropsychiatric disorders, and found all mouse models with immature brain phenotypes investigated in this study showed significant increases in lactate levels, suggesting a potential link between cellular hyperexcitability and the immature phenotype[17]. Additionally, immature-like gene expression patterns have been identified in the brain tissues of human patients with neuropsychiatric disorders, such as epilepsy, Alzheimer's disease, and schizophrenia[11,18–23], where neural hyperexcitability is a known or suspected etiological contribution.

In marked contrast to the detrimental effects of high-intensity neural activity, there are reported beneficial effects of moderate electrical activity on neural circuit function. Neural activity is essential for brain development, learning and memory[24,25], including enhancements in synaptic weight[26,27], neuronal excitability[28,29], and gene expression[30,31]. An interesting clinical application of moderate activity is the use of electrical brain stimulation technologies such as electroconvulsive therapy (ECT), as a class of methods for ameliorating negative symptoms in patients with major depression[32,33]. ECT promotes neuroplastic changes, such as increased adult neurogenesis in the dentate gyrus and hippocampal volume enlargement[34–36] with 6–12 repeated stimulations commonly used for long-term positive clinical outcomes[32,33]. Electroconvulsive stimulation (ECS), the animal equivalent of ECT, also induces molecular and electrophysiological changes, such as enhanced neurogenesis and altered synaptic transmission[9,14,37,38]. Similar changes occur with chronic administration of fluoxetine, a selective serotonin reuptake inhibitor (SSRI)[9,39,40]. Interestingly, repetitive ECS and chronic fluoxetine administration can induce cellular dematuration, suggesting that unspecified cellular plasticity may underpin their antidepressant effects[9,39–41] Likewise, other noninvasive brain

stimulation methods, such as repetitive transcranial magnetic stimulation (rTMS) and

transcranial direct current stimulation (tDCS), can induce neuroplastic changes in the brain and

have promising therapeutic potential [42,43]. However, despite their common use, the specific

cellular mechanisms engaged by these brain stimulation therapies, including clinical ECT,

rTMS, and tDCS, remain poorly understood.

In this study, we aimed to elucidate the primary cellular mechanisms by which brain

stimulation technologies like ECT cause cellular dematuration. We applied an ECT-like

optogenetic stimulation to granule cells of the hippocampal dentate gyrus, a key brain area for

ECT, to produce clinically relevant behavioral effects. This stimulation led to an immature-like

gene expression profile in the brain of mice, and a similar patten was observed in the post-

mortem brains of patients with mood disorders. Our analysis also revealed that repetitive

stimulations induce G_2/M cell cycle re-entry and nuclear structural plasticity with epigenomic

and morphonuclear signatures. These findings reveal a unique immature cellular state induced

by ECT-like neuronal activation, which has long-term stability, and a mechanism of activity-

dependent nuclear reprogramming, offering mechanistic insight into ECT effects and related

applications.

Results

1. *Neuronal activations trigger an immature cellular state in DG granule cells*

We challenged neurons *in vivo* with a stimulation protocol designed to mimic human

ECT, in which seizure duration is a critical determinant of therapeutic efficacy and is therefore

tightly controlled in clinical settings. Conventional methods, such as seizure-inducing drugs

and animal ECS, do not allow precise control over seizure duration, leading to variability in

cellular and network-level effects. To overcome this limitation, we used mice expressing

channelrhodopsin-2 (ChR2) in granule cells (GCs) of the dentate gyrus (DG) (POMC-

Cre::ChR2-EYFP mice) and applied optogenetic stimulation, termed Repetitive Optogenetic

Stimulation (REPOPS), enabling precise control over the frequency and duration of stimulation

(Fig.1a and Extended Data Fig. 1a). This optogenetic approach allowed us to establish a precise

experimental system to study cellular effects confined to a specific target cell population. We

employed a moderate stimulation protocol that did not cause cell death (Extended Data Fig.

1b–d).

REPOPS was performed either three or ten times consecutively (Fig. 1a). The

expression of calbindin (CB), a maturation marker of GCs, was reduced 24 hours after

REPOPS for three days, but returned to a level that was not significantly different from the pre-

stimulus levels within two weeks (Fig. 1b). In contrast, REPOPS for 10 days significantly

decreased CB expression at 24 hours and two weeks after the last stimulation (Fig. 1b). The

decrease in CB was inhibited by the administration of Ca^{2+} blockers (Extended Data Fig. 2a–

c), suggesting that Ca^{2+} influx is crucial for REPOPS-induced dematuration. RNA-seq analysis

showed that the number of genes with significant changes in expression ($P < 0.05$, fold change

> 1.2; vs. No Stim group) decreased over time after 3-day REPOPS, but remained unchanged

after 10-day REPOPS (Fig. 1c). Evidently, neuronal activation repeated a few times induces

short-term transcriptome changes that spontaneously revert to the original state; however, when

neuronal activation is chronically repeated, the changes are sustained for longer periods. To

examine whether REPOPS induces an immature-like gene expression pattern at the

transcriptome level, we assessed the similarity between gene expression patterns of

optogenetically-stimulated DG and those of normal infant mice (GSE113727[11]) by the Running

Fisher test[44], a rank-based nonparametric algorithm that evaluates the degree of overlap

between two gene groups implemented in BaseSpace (Illumina, Cupertino, CA, USA). There

was a striking degree of similarity between the datasets (Overlap $P = 3.9 \times 10^{-52}$) with the same

directional changes in expression (Fig. 1d). Collectively, these results indicate that REPOPS

for 10 consecutive days induces long-lasting immature-like gene expression patterns in the DG,

whereas REPOPS for three days does not.

As REPOPS was designed to model the effects of human ECT, we next asked
whether ECT in human patients also induces an immature-like gene expression pattern. We
reanalyzed RNA-seq data from the post-mortem DG of bipolar disorder and major depressive
disorder patients with and without ECT (SRP241159[45]). Genes differentially expressed
between ECT-treated and non-ECT-treated patients were compared with those changing during
typical hippocampal development (0–5 months vs. 20–39 years; GSE25219[46]). This analysis
revealed a significant overlap between the two datasets (Overlap $P = 1.2 \times 10^{-8}$), with
consistent directional changes (Fig. 1f), suggesting that ECT induces an immature-like gene
expression pattern in the DG of human patients, in line with the pattern observed in REPOPS.
Notably, both REPOPS and human ECT induced long-term alterations in the expression of
synapse-related genes: REPOPS significantly downregulated *Gria1*, *Gria2*, *Camk2a*, and
Camk2b, and upregulated *Arc* (Fig. 1e), while ECT-treated patients exhibited similar alterations
in GRIA1, GRIA2, and ARC compared with non-ECT-treated patients (Fig. 1g), suggesting a
shift toward an immature synaptic state.

2. REPOPS regulates the long-term plasticity of genomic structure

Changes in gene expression are often accompanied by alterations in the chromatin
structure of the associated genomic regions[47]. To investigate whether and how neural
overactivation alters the chromatin accessibility of neurons, we performed ATAC-seq on
optogenetically stimulated DG tissue (Fig. 1h). Volcano plots showing changes in chromatin
accessibility are shown in Fig. 1i. We found significant changes in chromatin accessibility in
806 chromatin regions (748 opened and 58 closed) and 21,853 chromatin regions (15,098
opened and 6,755 closed) in the Stim×3+2wks and Stim×10+2wks groups, respectively (Fig.
1i). These results indicate that repeated neuronal activation for 10 consecutive days, but not for
3 days, induced widespread and persistent reorganization of three-dimensional genome
structure, along with concordant global transcriptome changes.

3. REPOPS initiates long-term anti-depressive behavioral changes in mice

To assess whether ECT-like neuronal activation could alter anti-depressive behavior,
we first performed an open field (OF) test (Fig. 2a, 2b). Locomotor activity was recorded for
30 min each day, and the dorsal DG was optogenetically stimulated for 5 min (20–25 min after
the start) on each stimulation day. As previously reported[48], mice showed an increase in
locomotor activity immediately after stimulation (a1–a3 in Fig. 2a). Interestingly, on the day
after the first stimulation, mice in both Stim×3 and Stim×10 groups exhibited a significant
increase in distance traveled during the first 15 min compared to the No Stim group ($P = 0.0134$,

Day 2 in Stim×10 group; $P = 8.05 \times 10^{-4}$, Day 9 in Stim×3 group; Fig. 2b). In the Stim×3

group, this increase peaked on Day 9 and returned to control levels within two weeks (blue line

in Fig. 2b), whereas in the Stim×10 group, it declined gradually from the last stimulation day

but remained significantly elevated two weeks after the last stimulation (Day 24, $P = 0.00362$;

red line in Fig. 2b). Thus, neuronal activation of the DG repeated for 10 days induced

hyperlocomotor activity persisting for at least two weeks, whereas an increase in locomotor

activity after neuronal activation for three days was transient. We also conducted 24-hour

activity monitoring in the home cage (Fig. 2c). The distance traveled by mice in the Stim×10

group gradually increased after the initial stimulation and was significantly higher than that in

the No Stim and Stim×3 groups ($P < 0.05$, Days 7–36; Fig. 2c). Mice in the Stim×3 group did

not show any significant increase in locomotor activity in the home cage. These results indicate

that repeated neuronal activation causes a significant increase in baseline locomotor activity,

which persists for more than one month. Depression-related behavior was assessed with the

tail suspension test (Fig. 2d). Mice in the Stim×10 group showed significantly less immobility

compared to those in the No Stim ($P = 0.0094$) and Stim×3 groups ($P = 0.0198$). A similar

trend was observed in the forced swim test (Fig. 2e), where Stim×10 mice showed reduced

immobility compared to the No Stim group ($P = 0.0304$) and Stim×3 group ($P = 0.0770$). These

results suggest that repeated neuronal activation mimicking ECT reduces depression-like

behavior in mice (for other behavioral tests, see also Extended Data Fig. 3).

4. *Alterations of neural coding of navigational information*

190   To investigate whether and how neural overactivation affects circuit function, we

performed *in vivo* Ca^{2+} imaging under freely moving conditions combined with optogenetic

stimulation (Fig. 3a, 3b). On each stimulation day, we recorded DG neuron activity and

locomotor activity in the OF during the first 30 min, followed by 5 min of optogenetic

stimulation (590–650 nm, 20 msec pulse, 10 Hz; Fig. 3c). The distance traveled during the first

30 min in the Stim×10 group gradually increased and was significantly greater than in the No

Stim group two weeks after the last stimulation ($P = 0.0358$; Fig. 3d). However, the mean Ca^{2+}

transient rate did not differ significantly between these groups throughout the experiment (Fig.

3e).

199   Navigational information, such as position and speed, is encoded in the population

activity of DG neurons[16,49]. To examine how neural overactivation affects this coding, we

computed spatial and speed information for individual DG neurons. In the No Stim group,

spatial information increased on Days 2–11 and Days 24–26 compared to Day 1 (Cohen's $d =$

0.43 and 0.30), suggesting that OF exposure enhanced spatial coding (Fig. 3f). In contrast, the

Stim×10 group showed reduced spatial information on Days 2–11 and Days 24–26 compared

to the No Stim group (d = –0.50 and –0.45; Fig. 3f). Speed information remained unchanged

in the No Stim group throughout the experiment ($|d|$ < 0.10), but increased in the Stim×10

group on Days 2–11 and Days 24–26, exceeding that in the No Stim group (d = 0.46 and P =

0.63; Fig. 3i). Thus, repeated neuronal activation reduced spatial coding and enhanced speed-

related activity in individual DG neurons for over two weeks.

To examine information encoding at the population level, we predicted mouse

position and speed from the population activity of DG neurons using machine learning methods.

Representative decoding results are shown in Fig. 3g and 3j. In the No Stim group, position

decoding errors decreased on Days 2–11 and Days 24–26 and were significantly smaller than

chance (Fig. 3h). In contrast, the Stim×10 group showed no significant change in position

decoding accuracy, and decoding errors were significantly greater than in the No Stim group

on Days 2–11 and Days 24–26 (P = 0.0133 and 0.0335; Fig. 3h). These findings suggest that

position information coding was enhanced by OF exposure but impaired by neural

overactivation. Speed decoding accuracy did not change significantly during the experiment in

the No Stim group, but was increased by REPOPS in the Stim×10 group and was significantly

higher than in the No Stim group on Days 2–11 and Days 24–26 (P = 0.0363 and 0.0223; Fig.

3k). Thus, repeated neuronal activation reshapes information processing in the DG by

suppressing spatial map formation and enhancing speed-related activity, with unchanged

overall activity levels. This shift may underlie the behavioral changes observed in

optogenetically stimulated mice, such as increased locomotor activity and reduced depression-

like behavior.

5. *REPOPS-driven long-term increases in the expression of cell cycle-related genes*

We next explored potential molecular markers associated with REPOPS-induced

neuronal alterations by conducting gene ontology analysis of RNA-seq data from

optogenetically-stimulated DG. Notably, gene categories significantly altered in the

Stim×10+2wks group—but not in the Stim×3+2wks group—included cell cycle–related terms,

such as *"cell division"* and *"mitotic cell cycle process"* (Extended Data Fig. 4). Expression

levels of representative cell cycle-related genes are shown in Fig. 4a. Genes related to the G_2/M

phase, such as *Ccna2*, *Ccnb1*, *Ccnb2*, and *Cdc25c*, were significantly increased in

Stim×10+2wks compared to the No Stim group, whereas genes related to other phases, such as

Ccne1, *Ccne2*, *Cdk2* (G_1-S phase), and *Ccnd1*, *Cdk4* (G_1 phase), showed no significant changes

(Fig. 4a). We found that cyclin B, an essential regulator for the transition from G_2 to M phase,

was expressed in most DG neurons in Stim×10+2wks, while it was barely detectable in the No

Stim group (Fig. 4b). It was broadly expressed throughout the granule cell layer and was not
restricted to the subgranular zone, where newborn neurons are abundant, indicating that the
majority of cyclin B expression reflects induction in mature neurons, rather than enhanced
neurogenesis.
To assess whether gene expression patterns induced by REPOPS resembled specific
cell cycle phases, we used a publicly available microarray dataset of HeLa cells that captures
gene expression pattern dynamics throughout the cell cycle (Fig. 4c; GSE3497[50]). In this
dataset, genes characteristic of each phase were identified through microarray analysis of HeLa
cells sampled at multiple time points across the cell cycle. We compared these cell cycle phase-
specific gene lists (1–30 hour from S phase) with RNA-seq data from optogenetically
stimulated DG (Fig. 4c) and computed a "similarity score" for each time point, defined as the
$-\log_{10}$ of the overlap P value (Fig. 4d). In Stim×3+24hrs and Stim×10+24hrs, similarity score
peaks were observed at 7, 8, and 20 hours from S phase (scores > 3.0; Figs. 4d), corresponding
approximately to the G_2/M phase. In Stim×3+2wks, these peaks declined (scores < 2.0),
whereas in the Stim×10+2wks group, they remained elevated (scores > 3.0; Figs. 4d). Although
neurons are considered post-mitotic and do not undergo cell cycle progression, these findings
suggest that repeated neuronal activation reinstates G_2/M phase-like transcriptional programs
in mature neurons that persist for over two weeks.
257
6. *Long-term G_2/M phase-like epigenomic plasticity in nuclear structure*
We investigated whether REPOPS can induce changes in nuclear structure, focusing
on three major events that occur during the G_2/M phase: (i) nuclear lamina disruption, (ii) an
increase in the phosphorylation of histone H3 (pH3), and (iii) chromatin condensation.
(i) The nuclear lamina is a dense fibrillar network lining the inner nuclear membrane
and providing mechanical support for chromatin organization, which disrupts during mitosis
due to nuclear division[51,52]. In the No Stim group, most neurons exhibited normal, round-
shaped lamin B1, the main component of the nuclear lamina (Fig. 4e and Extended Data Fig.
5a, 5b). In contrast, neurons in the Stim×10+2wks group showed varying degrees of reduced
expression and/or disrupted lamin B1 structure, indicating that 10 days of REPOPS induced
nuclear lamina disruption (Fig. 4e).
(ii) Increased phosphorylated histone H3 (pH3) is another characteristic change
during mitosis, which is a known "mitosis marker." In the No Stim group, most neurons showed
minimal immunoreactivity of pH3, whereas the Stim×10+2wks group exhibited a significant
increase in pH3 immunoreactivity (Fig. 4f), suggesting that repeated neuronal activation
induces mitosis-like epigenetic modifications of chromatin in mature neurons.

(iii) The chromocenter is an aggregation of heterochromatin domains, which can be

visualized as concentrated dot-like signals in the nucleus by DNA counterstaining (indicated

by red arrowheads in Fig. 4g). It is considered a central site for chromatin condensation during

mitosis[53,54]. The chromocenter ratio–defined as the ratio of total chromocenter area to the

nucleus size–was significantly increased by REPOPS, indicating an enlargement of

heterochromatin domains (Fig. 4g). Thus, REPOPS induces a G_2/M phase-like nuclear states,

characterized by nuclear lamina disruption, increased pH3, and chromatin condensation (see

also Extended Data Fig. 6a, 6b).

7. *Cyclin B mediates REPOPS-induced changes in nuclear structure, cellular immaturity,*

and behavior

To investigate the molecular mechanisms underlying G_2/M phase-like nuclear

alterations, we knocked out the expression of cyclin B–an essential gene for the G_2/M phase

transition–by injecting a mixture of AAV9-SpCas9 and AAV9-sgRNA encoding sequences

targeting *Ccnb1* and *Ccnb2* (CyclinB-KO) into the DG (Fig. 5a, 5b). AAV9-sgRNA encoding

scrambled sequences (Scram-KO) was used as a control. The REPOPS-induced expression of

cyclin B, observed in neurons in the granule cell layer, was effectively eliminated in the

CyclinB-KO group (Extended Data Fig. 7b).

Lamin B1 expression was significantly reduced by REPOPS in both Scram-KO and

CyclinB-KO groups, but the degree of reduction was significantly smaller in the CyclinB-KO

group (Fig. 5c). Similarly, REPOPS significantly increased pH3 signals in both groups, but the

degree of increase was significantly reduced in the CyclinB-KO group (Fig. 5d). The

chromocenter ratio was significantly increased by REPOPS in the Scram-KO group, but not in

the CyclinB-KO group (Fig. 5e). Calbindin expression was significantly decreased in both

groups, but the reduction was significantly smaller in the CyclinB-KO group compared to the

Scram-KO group (Fig. 5f). In the open field test, REPOPS significantly increased locomotor

activity in the Scram-KO group from the day after the first stimulation, which remained

significantly greater than that of the No Stim group two weeks after the last stimulation ($P =$

0.0231, Day 24; Fig. 5g). In contrast, in the CyclinB-KO group, locomotor activity showed a

transient increase but gradually returned to baseline levels by Day 24 ($P > 0.99$; Fig. 5g).

Collectively, these findings demonstrate that cyclin B is a critical molecular mediator linking

repeated neuronal activation to G_2/M phase-like nuclear remodeling, neuronal immaturity, and

prolonged hyper-locomotor activity in mice.

8. *ΔFosB is upregulated by Cyclin B/Cdk1 activity and mediates persistent genomic*

remodeling

310   We next explored mechanisms underlying long-term genomic changes by

performing motif enrichment analysis[55] of ATAC-seq data (Fig. 1h, i). We found that open

chromatin peaks showed the highest similarity to ChIP-seq profiles of AP-1 transcription

factors, dimers composed of Fos and Jun family proteins, such as Fra1, Atf3, Batf, and JunB

(Extended Data Fig. 8a), suggesting that AP-1 mediates long-term genomic remodeling. We

focused on ΔFosB, a truncated FosB isoform, as a candidate driver of these long-term changes

due to its exceptionally long half-life (~208 hours[56]). We analyzed its expression time course

following REPOPS. After 3-day REPOPS, ΔFosB was transiently increased at 24 hours but

returned to baseline at 14 and 40 days (Fig. 6a). In contrast, after 10-day REPOPS, ΔFosB was

elevated at 24 hours and remained high at 14 and 40 days (Fig. 6a), suggesting that chronic

repetition of activation induces long-lasting elevation in ΔFosB. This sustained elevation likely

reflects increased baseline expression of Fosb rather than residual REPOPS-induced ΔFosB

(Fig. 6a Extended Data Fig. 8b). Notably, ΔFosB levels negatively correlated with calbindin

expression (Fig. 6b), which is consistent with previous reports[21], suggesting that prolonged

ΔFosB upregulation contributes to pseudo-immaturity of DG neurons (see also Extended Data

Fig. 9).

326   To examine whether cell cycle reentry is involved in this sustained ΔFosB

upregulation, we analyzed its expression in CyclinB-KO mice after REPOPS. At 24 hours after

10-day REPOPS, ΔFosB was similarly increased in both CyclinB-KO and Scram-KO groups,

indicating that cyclin B is dispensable for acute induction (Fig. 6c). In contrast, at two weeks,

ΔFosB was significantly lower in CyclinB-KO than in Scram-KO, suggesting that cyclin B

expression is necessary for the sustained elevation of ΔFosB. To further assess the role of

Cyclin B/Cdk1, we performed REPOPS while continuously perfusing its modulators—PD-

407824 (a Wee1 inhibitor) or BN82002 (a Cdc25c inhibitor)—via an optofluidic neural probe

(Extended Data Fig. 2a) and measured ΔFosB expression (Fig. 6d). Two weeks after 3-day

REPOPS, ΔFosB was not elevated, but PD-407824 infusion significantly increased its

expression (Fig. 6e). Conversely, after 10-day REPOPS, ΔFosB remained elevated for two

weeks, but this increase was suppressed by BN82002 (Fig. 6e). Calbindin showed a

complementary pattern: it remained unchanged two weeks after 3-day REPOPS but decreased

with PD-407824; whereas it was reduced after 10-day REPOPS but reversed by BN82002 (Fig.

6e). These findings indicate that Cyclin B/Cdk1 activity is essential for sustained elevation of

ΔFosB and the associated reduction in calbindin. Thus, repeated neuronal activation and cell

cycle reentry induce long-lasting and stable changes in activity-dependent transcription factors,

global genomic architecture, and cellular maturity.

Discussion

In this study, we tested the hypothesis that brain stimulation methods, such as electroconvulsive therapy (ECT) and repetitive transcranial magnetic stimulation (rTMS), may achieve beneficial clinical effects by reprogramming cellular state into an immature but stable configuration. We developed an ECT-like optogenetic stimulation, REPetitive OPtogenetic Stimulation (REPOPS), which reproducibly induces "dematuration" as indexed by global changes in the cellular transcriptome and 3D genome structure. Most of these changes were transient when REPOPS was repeated for three consecutive days, but could persist for more than one month after 10 consecutive days of activation, consistent with the 6–12 stimulus repeat regimen for ECT. RNA-seq data of human DG also showed that ECT induces immature-like gene expression patterns in patients, resembling the effects by REPOPS. Underlining the clinically relevant behavioral effects of REPOPS, we discovered a novel mechanism where our stimulation protocol induces cell cycle re-entry resembling the G_2/M-phase-like nuclear structure. These changes were reversed by deletion of Cyclin B, a master regulator of the G_2/M phase transition. We further found that activation of Cyclin B/Cdk1 stabilizes ΔFosB expression, thereby promoting sustained reorganization of genome structure. Our results establish a candidate cellular mechanism underpinning the beneficial clinical effects of ECT and offer a novel cell-based perspective for the development of brain stimulation therapies.

Our findings reveal that optogenetic stimulation of hippocampal granule cells can trigger coordinated changes in gene expression, nuclear structural plasticity, and G_2/M cell cycle re-entry, which collectively define an immature-like cell state. The evidence supporting G_2/M cell cycle re-entry includes the increased expression of M phase-related genes such as Cyclin A2, Cyclin B1, Cyclin B2, and Cdc25c, and changes in nuclear structure, including nuclear lamina disruption, an increase in pH3, and chromocenter enlargement. In the literature, cell cycle re-entry has been implicated in several neuropsychiatric and neurological disorders[57,58]. For example, aberrant expression of cell cycle-related genes, such as Cyclin A, Cyclin B, and Cdk1, has been reported in Alzheimer's disease[57,59–63] and temporal lobe epilepsy[64], as well as in animal models of risk events for these diseases, such as epileptic seizures[65,66], ischemia[67], and encephalitis[68]. Nuclear alterations related to the cell cycle, such as nuclear lamina disruption and increased pH3, are observed in the brains of patients and animal models of Alzheimer's disease[58,69,70] and epileptic seizures[71]. Recent large-scale GWAS studies identified several M phase-related genes (e.g., *CCNB2, CDC25C, CENPM, CENPT, MAD1L1,* and *STAG1*) near the risk loci for schizophrenia[72,73]. While these diseases have diverse etiologies, neural hyperexcitability and cell cycle-like processes may be common

contributors to their pathophysiology[74–77]. This study is the first, to our knowledge, to show a

causal link between chronic neural activation and G_2/M cell cycle re-entry. Interestingly, the

cell cycle re-entry observed in various diseases is typically associated with apoptosis[78,79];

however, our REPOPS method did not induce significant cell death (Extended Data Fig. 1c).

REPOPS triggered nuclear structural dynamics, but they were less pronounced than those in

dividing cells during the late M phase[79,80] and there were no fully compacted chromosomes

(Fig. 4e, 4g), suggesting that neurons were arrested in a G_2 to early M phase-like state for at

least two weeks. Importantly, we do not assert that this immature-like cellular state is a perfect

regression to developmental immaturity but rather an analog state of immaturity, a state that is

phenotypically immature relative to the mature state. Another key finding of this study is the

involvement of ΔFosB in neuronal dematuration. ΔFosB has previously been reported to be

highly expressed in the striatum of addiction models[81,82], as well as in the DG of patients with

epileptic seizures and Alzheimer's disease[21,83]. Owing to its long half-life (~208 hours)[56],

ΔFosB is widely believed to mediate long-lasting cellular changes. However, given this half-

life, stimulus-induced ΔFosB would be expected to decay to ~4% of its peak level within 40

days. Therefore, the marked increase observed in the Stimx10+40day group (Fig. 6a) is

unlikely to reflect residual ΔFosB from prior stimulation, but rather a sustained elevation of its

basal expression (Extended Data Fig. 8b). This persistent up-regulation of ΔFosB likely

contributes to long-lasting 3D genome reorganization and neuronal dematuration. We also

suggest that REPOPS triggers repeated calcium influx that initiates dematuration and cell cycle

re-entry (Extended Data. Fig. 2a—c). Activation of calcium signalling is essential for cell cycle

progression[84,85]; therefore, REPOPS-stimulated neurons may transduce a specific pattern of

calcium signaling that causes cell cycle re-entry. Further studies will clarify this signaling

pattern, and we predict that variations of REPOPS methodology may enable fine-scale external

control of cell dematuration *in vivo*.

ECT is well documented as an effective treatment for psychiatric disorders such as

severe depression and anxiety. The current results demonstrate that cellular dematuration

triggered by an ECT-like stimulus can produce positive behavioral phenotypes, including

decreased depression-like symptoms and increased motor activity, similar to the clinical ECT

effects. An important question is how these behaviors are enabled by neuronal stimulation at

the neural circuit level. REPOPS increased the number of speed cells and the speed coding

accuracy in the DG (Fig. 3i–k). Prior studies indicate that activation of DG neurons can elicit

robust exploration of the environment[48], and our observation of increased speed-related activity

in the DG may account for the increased locomotor activity observed in the open field (Fig. 2a,

2b) and home cage (Fig. 2c). Activation of DG neurons is also known to suppress depression-
like behaviors[86,87], therefore, increased speed-related activities of DG neurons may explain the
link between ECT-like stimulation and the reduction in depression-like behavior observed in
the tail suspension and forced swim tests (Fig. 2d, 2e). In contrast to these beneficial effects, a
potential negative side effect of ECT is memory loss, including retrograde and anterograde
amnesia[88,89]. Indeed, we observed a deficit in remote memory, or the ability to recall
information from neocortex into the hippocampal circuit (Extended Data Fig. 3a and
Supplementary results 2). REPOPS caused long-term (> 2 weeks) alterations in synapse-related
genes, including *Gria1*, *Gria2*, *Camk2a*, *Camk2b*, and *Arc* (Fig. 1e). Reanalysis of RNA-seq
data from human patients with and without ECT also revealed significant changes in the
expression of *GRIA1, GRIA2,* and *ARC* (Fig. 1g). These genes play pivotal roles in synaptic
and homeostatic plasticity[90–92] and their decline may be related to memory impairment due to
neural hyperexcitation. Long-term down-regulation in the expression of synapse-related genes
may interfere with the development of place cells and spatial information coding in the DG
(Fig. 3f–h) and CA1[16,93–95]. Such REPOPS-dependent deficit in neural plasticity may share a
common mechanism with ECT-induced amnesia. Thus, ECT-like REPOPS alters neural coding,
leading to both decreased depressive-like behaviors and memory deficits. Further research will
address the ECT stimulus conditions that do not impair memory and learning, but preserve
beneficial clinical therapeutic effects.

Our findings also provide a novel perspective on the mechanism of action of
antidepressant therapy. In this study, we found that REPOPS-triggered cellular and behavioral
effects share intriguing similarities to previously reported behavioral effects and neuronal
dematuration induced by ECS or selective serotonin reuptake inhibitors (SSRIs)[9,10,39,40].
Chronic SSRI treatment may reactivate the plastic state in the adult brain, comparable to that
observed during development[14,41]. For example, in the adult visual cortex, chronic fluoxetine
reopens a critical period of plasticity for the recovery of visual function[96,97], and, in the
hippocampus, increased neural circuit plasticity can promote fear memory extinction[14,98].
SSRIs may regulate neural plasticity by increasing the environmental susceptibility of neural
circuits, thereby reducing depression via exposure to favorable environments[99,100]. This form
of "meta-plasticity"[101,102], or the induction of changes in neural plasticity itself, can also
manifest in opposite forms by neural hyperactivity. A recent study showed that recurrent
seizures cause a long-term reduction in neuronal excitability, indicating decreased neural
plasticity[103]. In the current study, REPOPS for 10 days inhibited spatial map formation by
exposure to the environment for more than two weeks (Fig. 3f–h), suggesting an altered meta-

plasticity for spatial coding. Thus, neural overactivation and dematuration can alter the meta-
plasticity of neural circuits, which may underlie the effects of antidepressant treatments such
as ECT and rTMS, or the development of neuropsychiatric disorders involving neural
hyperexcitability in their etiology. A key insight from the present work is that the immature
cellular state appears to be bistable; that is, it can serve as a reversible therapeutic intermediate,
as observed in ECT, or consolidate into an irreversible pathological lesion, as seen in disease
models.

In summary, we propose a working model for the mechanism of brain stimulation
therapies in the treatment of psychiatric disorders. This class of brain diseases may represent
neuronal populations trapped in various energy minima of metabolic and/or synaptic
inflexibility, such as the loss of plasticity by the pathological closure of neural circuits to
environmental stimulation, as discussed above. In this model, repetitive brain stimulation
methods like ECT or rTMS trigger patterned calcium influx and signaling to drive cellular
dematuration that temporarily resets cellular metabolic and synaptic homeostasis to restore
normal neural circuit flexibility and beneficial clinical outcomes like anti-depressive behavior.
This immature-like state can persist for more than one month, and excessive neuronal
activation may eventually reinforce rather that reset metabolic or synaptic dysfunction in a
vicious cycle associated with chronic psychiatric disorders associated with hyperactivity.
Therefore, our model suggests that the immature-like cell state may serve as a bistable switch
for cellular health at moderate levels of electrical brain stimulation, or disease progression at
higher levels. We believe that these findings open a window to novel methods and protocols
for artificial neural stimulation that may enable cell reprogramming for brain health.

Figure 1. REPOPS induces long-term changes in gene expression, chromatin accessibility, and locomotor activity in mice

a, Experimental design. POMC-Cre::ChR2-EYFP mice express ChR2 in GCs of the DG. Blue light was delivered via a wireless LED device. Optogenetic stimulation (10-msec pulse at 10 Hz for 5 min/day) was applied for 3 or 10 consecutive days. Brains were collected 24 hours or two weeks after the last stimulation.

b, Immunohistostaining of calbindin, a maturation marker of GCs, in dorsal DG. Scale bar, 500 μm. Bar graph shows calbindin immunoreactivity (10–15 sections from 4 mice in each group).

Error bars represent the Mean ± s.e.m. One-way ANOVA followed by Bonferroni correction

for multiple comparisons, $F_{(4,106)} = 24.9$, $P = 1.5 \times 10^{-14}$, ***$P < 0.001$.

c, RNA-Seq. Volcano plots show fold change and P values distribution after differential

transcript abundance analysis using DESeq2. All stimulation groups were compared with the

No Stim group (n = 6 mice per group). Genes with significantly differential expression ($P <$

0.05; fold change > 1.2) are highlighted in red. Bar graphs indicate number of genes with

significant changes (i.e. red dots in Volcano plots).

d, Overlap between genes altered by REPOPS (Stim×10+2wks vs. No Stim) and during normal

development of the mouse DG (P8 infants vs. 33-week-old adults; GSE113727). Bar graphs

show–$\log_{10}$ of overlap P-values for genes upregulated or downregulated by each condition.

The Bonferroni correction was used to adjust the significance level according to the number of

dataset pairs.

e, mRNA expression levels (FPKM) of representative synapse-related genes. Mean ± s.e.m. (n

= 6 mice/group). One-way ANOVA followed by Bonferroni correction for multiple

comparisons; *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$.

f, Overlap between genes altered in DG of human patients with bipolar disorder and major

depressive disorder (with a history of ECT vs. without ECT; SRP241159[45]) and those altered

during human hippocampal development (0-5 month infants vs. 20-39-year-old adults;

GSE25219[46]).

g, mRNA expression levels (FPKM) of representative synapse-related genes in post-mortem

DG from healthy control subject (n=93), patients without ECT history (n=61), and patients

with ECT history (n=7).

i ATAC-seq. Integrative Genomics Viewer (IGV) visualization of representative gained-open

and gained-closed chromatin profiling coverage.

j, Volcano plot showing the fold change and P value distribution after differential analysis of

chromatin accessibility using DESeq2. Stim×3+2wks and Stim×10+2wks groups were

compared with the No Stim group (n = 4 mice/group). Bar graph shows number of peaks with

a significant change ($P < 0.05$).

Figure 2. REPOPS induces anti-depressive behaviors

a, Distance traveled in the open field during the first 30 min on Day 1 (a1), Day 8 (a2), Day 10 (a3), and Day 24 (a4), corresponding to time points in **b** (red arrowheads). Solid lines show group means (n = 6 mice/group); semitransparent area indicates ± s.e.m. (black, No Stim; blue, Stim×3+2wks; red, Stim×10+2wks). Optogenetic stimulation was introduced for 5 min (20–25 min after the start) on Day 1, Day 8, and Day 10. Asterisk indicates Stim×10+2wks vs. No Stim; One-way ANOVA with Bonferroni correction; *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$.

b, Total distance traveled in the first 15 min of the open field test from Day 1 to Day 24. Two-

way repeated measures ANOVA; Stim type, $F_{(2, 14)} = 9.96$, $P = 2.0 \times 10^{-3}$; Day, $F_{(23, 322)} = 5.08$,

$P = 4.63 \times 10^{-12}$. Stim type × Day, $F_{(46, 322)} = 2.58$, $P = 7.81 \times 10^{-7}$.

c, 24-hour locomotor activity in the home cage. Daily average distance traveled is shown (n =

10, 8, and 10 mice for No Stim, Stim×3, and Stim×10 groups, respectively). Data missing on

Day 28, 29, 37 and 47 due to computer system error. $P < 0.05$ on Day 7–8, 10–16, 20, 22–23,

25–30, 33–36 (Stim×10 vs. No Stim).

d, Immobility (%) in tail suspension test. Values are the Mean ± s.e.m. Two-way repeated-

measures ANOVA results shown in panel. n = 10, 9, and 10 mice. Box plot summarizes average

immobility over 10 min. One-way ANOVA with Bonferroni correction; *$P < 0.05$, **$P < 0.01$,

***$P < 0.001$.

e, Same as **d**, for immobility (%) in the forced swim test.

Figure 3. REPOPS mediates long-term changes in neural coding of navigational information

a, Simultaneous *in vivo* calcium imaging and optogenetic stimulation under free-moving conditions. AAV-syn-GCaMP6f and AAV-syn-ChrimsonR were co-injected into the DG, followed by GRIN-lens implantation. An integrated miniature microscope (nVoke; Inscopix) enabled large-scale cellular-resolution Ca^{2+} imaging and optogenetic stimulation.

b, Left: Expression of GCaMP6f and ChrimsonR in the DG. Right: Max projection of relative fluorescence change ($\Delta F/F$) of Ca^{2+} transients from all imaging frames during the first 30 min,

showing representative active neurons.

c, Top: Experimental design of Ca^{2+} imaging combined with optogenetic stimulation during

the open field test. Each 50-min session consisted of 30-min Ca^{2+} imaging (used for the

following analysis), 5-min optogenetic stimulation, and 15-min post-stimulation recording.

This was repeated for 10 days. No light was introduced to mice in the No Stim group. Bottom:

$\Delta F/F$ traces from 15 representative neurons (scale bar, 10% $\Delta F/F$). Data from 25 to 38 min of

a 50-min session are shown.

d, Distance traveled during the first 30 min

e, Average Ca^{2+} transient rate during the first 30 min.

f, Distribution of spatial information for the No Stim (grey curve) and Stim×10 groups (red

curve). The vertical axis represents the frequency of distribution, with the dotted line indicating

the criterion for place cells (top 95% percentile of the shuffled distribution; see Methods

section). Bar graph indicates proportion of place cells.

g, Representative results of position decoding using Ca^{2+} imaging data. The first 30 min was

split into two 15-min halves for training and test data for decoding. Black/red lines: observed

position; grey/pink lines: decoded position.

h, Decoding accuracy (mae; mean absolute error, cm). Dotted lines: shuffled control. Two-way

repeated measures ANOVA: Stim type, $F_{(1, 8)} = 6.49$, $P = 0.034$; Day, $F_{(2, 16)} = 2.89$, $P = 0.085$;

Stim type × Day, $F_{(2, 16)} = 1.11$, $P = 0.353$. Bonferroni correction for multiple comparisons was

performed, *$P < 0.05$.

i, Same as **f**, but for speed information and speed cells. The dotted line indicates the criterion

for speed cells (top 99% percentile of the shuffled distribution).

j, Same as **g**, but for speed decoding.

k, Same as **h**, but for speed decoding accuracy. Two-way repeated measures ANOVA: Stim

type, $F_{(1, 8)}=4.02$, $P = 0.080$; Day, $F_{(2, 16)} = 4.87$, $P = 0.022$; Day×Stim type, $F_{(2, 16)} = 3.98$, $P =$

0.039. Bonferroni correction for multiple comparisons was performed, *$P < 0.05$.

Figure 4. REPOPS confers G₂/M phase-like gene expression patterns and nuclear structure plasticity

a, Expression of representative cell cycle-related genes. mRNA expression levels (FPKM) are normalized to the control group (No Stim). Data are shown as mean ± s.e.m. (n = 6 mice/group). One-way ANOVA with Bonferroni correction; $*P < 0.05$, $**P < 0.01$.

b, Cyclin B immunostaining (red) in the DG. Beeswarm plot shows cyclin B immunoreactivity

in individual neurons. Blue bars indicate the median values of 147 neurons (No Stim) and 202

(Stim×10+2wks) neurons from three biologically independent samples. ML, molecular layer;

GCL, granule cell layer.

c, Cell cycle phase analysis. To infer the cell cycle-like status of DG neurons after REPOPS,

RNA-seq data from optogenetically stimulated DG (Fig. 1c) were compared with publicly

available reference gene expression profiles representing each phase of the cell cycle. These

reference profiles were derived from microarray data of HeLa cell sampled at 1-hour intervals

(1-30 h) after release from S phase arrest (Whitfield *et al.*, 2002; GSE3497)[50].

d, Similarity in gene expression patterns between optogenetically stimulated DG and gene

expression profiles representing each phase of the cell cycle. Similarity scores were calculated

as –log of the overlap *P* value between RNA-seq data of stimulated DG and gene sets

representing each time point of the cell cycle (1–30 hours from the start of S phase). Positive

scores indicate concordant expression changes (e.g., both upregulated), while negative scores

indicate opposite trends. The positions of the S and M phases on the x-axis reflect cell cycle

timing estimated from the HeLa cell data (GSE3497; Whitfield et al., 2002)[50].

e, Lamin B1 immunostaining (red) in the DG. Enlarged views of boxed regions are shown at

right. Arrows and arrowheads indicate neurons with partial or complete disruption of lamin B1

structure. Scale bars: 20 μm (left), 10 μm (right). Beeswarm plots show immunoreactivity and

circularity of lamin B1 in individual neuron (n = 289 and 320 neurons from three independent

samples). Blue bars, medians. Welch's *t*-tests: immunoreactivity, $t_{(589)} = 4.87$, $P = 1.45 \times 10^{-6}$;

circularity, $t_{(547)} = 5.87$, $P = 7.38 \times 10^{-9}$.

f, Immunostaining of phospho-histone H3 (pH3) (red) in the DG. Scale bars: 200 μm (left), 10

μm (right). Beeswarm plot shows pH3 signal intensity in individual neurons (n = 415 and 459

neurons from three independent samples). Welch's *t*-test: $t_{(858)} = 14.1$, $P < 2.2 \times 10^{-16}$.

g, DNA counterstaining with Hoechst (blue and white pseudo-color) and cyclin B

immunostaining (red) in the DG. Enlarged views of boxed areas are shown on the right. Red

arrowheads indicate chromocenters–an aggregation of the heterochromatin domain visible by

Hoechst staining. White arrowheads/arrows: neurons with large/small chromocenters and

high/low cyclin B expression, respectively. Scale bars: 100 μm (left), 10 μm (right). Beeswarm

plot shows the chromocenter ratio (%) per neurons (n = 147 and 202 neurons from three

independent samples). Welch's *t*-test: $t_{(306)} = 5.46$, $P = 9.38 \times 10^{-8}$.

Figure 5. Cyclin B deletion reverses multiple REPOPS-induced phenotypes

a, Experimental design. AAV9 vectors encoding Cas9 and sgRNAs targeting *Ccnb1* and *Ccnb2* (CyclinB-KO) or a *Scramble* sequence (Scram-KO) were co-injected into the DG, followed by optic fiber implantation. After 7-14 days, optogenetic stimulation was performed for 10 consecutive days. Brains were sampled 2 weeks after the last stimulation. The open field test was conducted from Day 1 (immediately before the first stimulation) to Day 24 (two weeks after the last stimulation).

b, Co-expression of sgRNAs and Cas9 in the DG. sgRNAs were co-expressed with mCherry, and Cas9 was tagged with HA. Most neurons in the granule cell layer showed both mCherry and HA immunoreactivity.

c, Lamin B1 immunostaining in the DG (converted to red for visibility; originally stained with Alexa647). Upper and lower panels show control (Scramble-KO) and *Ccnb1-Ccnb2*-KO (CyclinB-KO) mice, respectively. Scale bars: 20 μm (left), 10 μm (right). Beeswarm plot shows lamin B1 immunoreactivity. Red bars, median. One-way ANOVA, $F_{(3, 679)} = 60.19$, $P < 2.0 \times 10^{-16}$; Bonferroni corrected, ***$P < 0.001$. n = 245, 301, 271, and 246 neurons from three independent samples per group.

d, pH3 immunostaining in the DG. One-way ANOVA, $F_{(3, 679)} = 325.7$, $P < 2.0 \times 10^{-16}$; ***$P < 0.001$; 199, 158, 143, and 183 neurons from five independent samples per group. Scale bar, 20 μm.

e, Chromocenter visualized by Hoechst staining (white, pseudo-color). One-way ANOVA, $F_{(3, 679)} = 39.1$, $P < 2.0 \times 10^{-16}$; ***$P < 0.001$; 199, 158, 143, and 183 neurons from five independent samples per group. Scale bar, 10 μm.

f, Calbindin staining in the DG. One-way ANOVA, $F_{(3, 725)} = 104.3$, $P < 2.0 \times 10^{-16}$; ***$P < 0.001$; 217, 157, 162, and 198 neurons from five biologically independent samples per group. Scale bar, 20 μm.

g, Open field test: total distance traveled during the first 15 min from Day 1 to Day 24. Left: Scramble-KO; Right: CyclinB-KO. Black and red lines represent No Stim and Stim×10+2wks groups, respectively. Solid lines, group means (n = 7 mice/group); semitransparent areas, ±s.e.m. Two-way repeated measures ANOVA: Groups, $F_{(3, 24)} = 4.271$, $P = 0.015$; Day, $F_{(6, 163)} = 4.271$, $P = 0.0204$; Groups×Day, $F_{(20, 163)} = 1.09$, $P = 0.36$.

Figure 6. Cyclin B/Cdk1 activity mediates long-term ΔFosB expression and pseudo-immaturity of neurons

a, Top, ΔFosB immunostaining in the DG. Scale bar, 500 μm. Bottom, time course of ΔFosB expression. Day 0 denotes the first stimulation day. Values were normalized to the mean of the No Stim group. The black dashed line indicates the theoretical decay of ΔFosB (half-life = 208 h); blue and red curves represent exponential fits for the Stim×3 and Stim×10 groups,

respectively. n = 9–11 slices per group. One-sample t-tests were performed between observed

values and the theoretical decay curve; ***$P < 0.001$.

b, Immunostaining of ΔFosB and calbindin in the DG (same data as in panel a). Only No Stim

and Stim×10+40 days are shown. Scale bar, 20 μm. Right, scatter plot showing the correlation

between ΔFosB and calbindin immunoreactivity.

c, ΔFosB immunostaining in the DG of Scram-KO and CyclinB-KO mice (same mice as in

Fig. 5). Bottom panels show enlarged views of the white boxes. Grayscale indicates Hoechst

staining. Scale bar, 20 μm. Beeswarm showing ΔFosB immunoreactivity (n = 129–165 cells

per group). Black bars indicate the median. One-way ANOVA: $F_{(5, 859)} = 270.94$, $P < 2.0 \times$

10^{-16}; post hoc comparisons with Tukey's test.

d, Schematic of Cyclin B/Cdk1 activation (modified from Deibler *et al.* 2010[104]). Cdk1 is

activated as cyclin B accumulates. This activation is inhibited by Wee1 and promoted by

Cdc25c.

e, Top, ΔFosB immunostaining in the DG. Middle, enlarged view of the boxed region.

Grayscale indicates Hoechst staining. Bottom, calbindin immunostaining. Scale bar, 20 μm.

f, Beeswarm plots showing ΔFosB, calbindin, and chromocenter ratio. n = 114–156 cells per

group. Black bars indicate the median. One-way ANOVA: ΔFosB: $F_{(4, 649)} = 405.45$, $P < 2.0 \times$

10^{-16}; Calbindin: $F_{(4, 655)} = 117.65$, $P < 2.0 \times 10^{-16}$; Chromocenter ratio: $F_{(4, 655)} = 14.63$, $P < 2.0$

$\times 10^{-16}$. Multiple comparisons were corrected using Tukey's test.

Materials and Methods

Animals

ROSA26-CAG-stop[flox]-ChR2(H134R)-EYFP (Strain#012569; RRID:IMSR_JAX:012569) and

POMC-Cre (Strain #010714; RRID:IMSR_JAX:010714) were obtained from Jackson

Laboratories (Bar Harbor, Maine). These mouse lines were crossed to generate POMC-

Cre::ChR2-EYFP mice, which express ChR2-EYFP selectively in the granule cells (GCs) of

the dentate gyrus (DG) (Fig. 1a). Mice were used for experiments at 3–4 months of age for

RNA-seq and ATAC-seq experiments, and at 6–12 months for other experiments. Three or four

mice were housed in one cage and switched to individual housing 1–2 weeks prior to the

experiment. They were housed with a 12 hr light/dark cycle (lights on at 7:00 a.m., off at 7:00

p.m.) with access to food and water *ad libitum*. The room temperature was maintained at 23 ±

2°C. All the experiments were conducted during the light period. All animal experiments were

approved by the Institutional Animal Care and Use Committee of Fujita Health University

based on the Law for the Human Treatment and Management of Animals and the Standards

Relating to the Care and Management of Laboratory Animals and Relief of Pain. Every effort

was made to minimize the number of animals used.

Stereotactic Surgery

Optic fiber implantation

For optogenetic stimulation, a blue-light optic fiber was implanted into the dorsal DG of adult

POMC-Cre::ChR2-EYFP mice using conventional methods[105]. Briefly, adult mice were

anesthetized with 1.5–3% isoflurane at an oxygen flow rate of 1 L/min. The head fur was

shaved and the incision site was sterilized with 70% ethanol before the surgical procedure. The

mice were mounted on a stereotaxic device (Narishige, Tokyo, Japan). After the scalp was

incised and pulled aside, a 1.0 mm diameter craniotomy was created with a surgical drill on

the skull above the implantation site. Using the stereotaxic device, a 250 μm diameter optic

fiber (470 nm, fiber diameter 250 μm; TeleC-B-3mm-250μm, Bioresearch Center, Aichi,

Japan) was precisely inserted into the target site (2.0 mm posterior to bregma (AP), 1.5 mm

lateral to midline (ML), and 1.5 mm ventral to bregma (DV)) and fixed to the skull with dental

cement. After implantation, the scalp was sutured and treated with povidone-iodine. To avoid

the effects of surgery on mouse behavior, a recovery period of 5–10 days after surgery was

provided when performing behavioral experiments.

Stereotactic injection of AAV9

For the genome editing experiment (Fig. 5), a mixture of adeno-associated virus (AAV)

carrying Cas9 and sgRNA was injected into the DG immediately before optic fiber

implantation. For dominant-negative inhibition of ΔFosB experiment (Extended Data Fig. 9),

we injected an AAV expressing ΔJunD (a kind gift from A.J. Robison) was injected.

Anesthetization and craniotomy were performed in the same way as described above. A glass-

capillary pipette (25 μm inner-diameter tip) filled with the virus suspended in mineral oil was

inserted into the target site (2.0 mm posterior to bregma (AP), 1.5 mm lateral to midline (ML))

through the surgical drill on the skull. A 200 nL of 1:3 AAV mixture of AAV9-U6-sgRNA

(*Ccnb1-Ccnb2* or Scramble)-hSyn1-mCherry (2.0×10^{13} vector genomes vg/ml) and AAV9-

pMecp2-HA-SpCas9 (2.0×10^{13} vg/ml) was injected at 200 nL/min using a microinjection

pump (Nanoliter 2010, WPI, FL) at three depths (1.5 mm, 2.0 mm, and 2.5 mm). After each

injection, a glass pipette was held in place for 1–2 min before retraction to prevent leakage.

Shortly after virus injection, the optic fiber was embedded in the same site using the procedure

described above.

Implantation of injection cannula

In the experiment that combined optogenetic stimulation and drug injection (Fig. 6d–f and

Extended Data Fig. 2), a guide cannula (24-gauge, 11 mm length, C316G, Plastics One) was

implanted into the dorsal DG using the same procedure as that used for optic fiber implantation.

Optogenetic Stimulation and drug infusion

For optogenetic stimulation under freely moving conditions, we employed a wireless LED

illuminator, Teleopto (BioResearch Center, Nagoya, Japan). A 2.0g infrared receiver (TeleR-2-

P, BioResearch Center) was attached to an optic fiber implanted in the mouse head, whereas in

the control group, a dummy receiver was used as a substitute. This receiver was signalled by

an infrared generator (TeleEmitter-C, BioResearch Center) to light up the optic fiber. The light

intensity varied depending on the custom-made fiber, typically within a range of 3–5 mW. We

set the stimulation protocol to 10 msec pulses at a frequency of 10 Hz, utilizing a pulse

generator (STOmk-2, BioResearch Center). These photostimulation intensities are sufficiently

weak to avoid tissue damage[106].

For optogenetic stimulation with an intracerebral drug infusion (Fig. 6d–f and Extended Data
Fig. 2), we employed injection cannulas (31-gauge, 11 mm length, C316I, Plastics One) and
customized optic fibers designed to fit the implanted guide cannula (Bio Research Center). For
acute injection (Extended Data Fig. 2), the injection cannula was attached to the implanted
guide cannula, and 2uL of the drugs was injected over approximately 5 min. The Ca^{2+} blockers
consisted of 100 mM NBQX (ab120046, Abcam), 100 mM AP5 (ab120003, Abcam), 100 mM
mibefradil (ab120343, Abcam), and 100 mM nimodipine (ab120138, Abcam) diluted in saline
(Extended Data Fig. 2c). Kinase inhibitors, K252a (sc-200517, Santa Cruz), SB218078 (sc-
203692, Santa Cruz), Dasatinib (sc-218081, Santa Cruz), and Ro-32-0432 (sc-3549, Santa
Cruz), were used at the same concentration (1.0 mM). Five minutes after the injection, the
injection cannula was gently detached, and a customized optic fiber was inserted through the
guide cannula. For chronic drug infusion (Fig. 6d–f), an osmotic pump (Alzet model 1004) was
implanted subcutaneously and connected to a guide cannula. The pump was filled with PD-
407824 (sc-203669, Santa Cruz) or BN82002 (#217691, Calbiochem) and delivered the drug
at a constant rate. For the effective infusion of drugs into brain tissues *in vivo*, the concentration
of the solution was set to 100–1000 times higher than that used for bath applications in acute
slice cultures according to previous studies[107]. Stimulation was performed in the same manner
as described above. All procedures were conducted in an awake state.

RNA-Sequencing Library Construction, Sequencing, Mapping, and Data Analysis
The dentate gyrus was rapidly dissected from the adult mouse hippocampus under a dissection
microscope as previously described[108]. Previous studies have shown that such preparations are
highly enriched with mature dentate granule neurons (–90% NeuN$^+$ neurons[30,109]; see also
Extended Data Fig. 7a). Total RNA from dissected tissue samples was extracted using
NucleoSpin RNA (Macherey-Nagel, Duren, Germany), according to the manufacturer's
instructions. The RNA quality was assessed with an Agilent 2200 TapeStation (Agilent
Technologies, Santa Clara, CA, United States). Sequencing libraries were prepared using the
TruSeq Stranded mRNA Sample Prep Kit (Illumina, San Diego, CA, USA) and sequenced as
100-bp paired-end reads on the Illumina HiSeq 2500. Sequenced data were mapped onto the
Mus musculus genome (GRCm38/mm10) using TopHat (v2.0.14). Fragments per kilobase per
million (FPKM) values were computed by Genedata Profiler Genome (v10.1.12). Genes with
low expression (FPKM < 0.1) were excluded from the analysis to avoid an excessively high
FDR due to low expression levels.

ATAC-Sequencing Library Construction, Sequencing, Mapping, and Data Analysis

The dentate gyrus of adult mice was dissected in the same manner as that used for RNA-seq. The dentate gyrus was slowly frozen in a freezing box (Cat. No. 5100-0001; Mr. Frosty Freezing Container, Thermo Fisher Scientific). Flash-frozen tissue was sent to Active Motif to perform the ATAC-seq assay. The tissue was manually dissociated, isolated nuclei were quantified using a hemocytometer, and 100,000 nuclei were tagmented as previously described[110], with some modifications based on Corces et al. 2017[111] using the enzyme and buffer provided in the Nextera Library Prep Kit (Illumina). Tagmented DNA was purified using the MinElute PCR purification kit (Qiagen), amplified with 10 cycles of PCR, and purified using Agencourt AMPure SPRI beads (Beckman Coulter). The resulting material was quantified using the KAPA Library Quantification Kit for Illumina platforms (KAPA Biosystems) and sequenced using PE42 sequencing on a NextSeq 500 sequencer (Illumina). For the analysis of ATAC-seq data, we first aligned reads using the BWA algorithm (mem mode; default settings). Duplicate reads were removed, and only reads mapping as matched pairs and uniquely mapped reads (mapping quality $>= 1$) were used for further analysis. Alignments were extended in silico at their 3'ends to a length of 200 bp and assigned to 32-nt bins along the genome. The resulting histograms (genomic "signal maps") were stored in the bigWig files. Peaks were identified using the MACS 2.1.0 algorithm at a cutoff of P-value 1.0 $\times 10^{-7}$, without the control file, and with the –nomodel option. Peaks on the ENCODE blacklist of known false ChIP-Seq peaks were excluded. Signal maps and peak locations were used as input data for Active Motif's proprietary analysis program, which creates Excel tables containing detailed information on sample comparisons, peak metrics, peak locations, and gene annotations. For differential analysis, reads were counted in all merged peak regions (using Subread) and the replicates for each condition were compared using DESeq2. IGV was used to visualize the raw intensities[112]. Statistical analyses were performed using an in-house R script, unless otherwise specified.

Immunohistochemistry

Immunostaining of brain sample with PFA fixation

The mice were deeply anesthetized with isoflurane and transcardially perfused with PBS,
followed by 4% PFA in PBS. Brains were dissected, immersed overnight in 4% PFA in PBS,
and cryoprotected by incubation in 20% sucrose in PBS for three days at 4℃. After
cryoprotection, the brains were mounted in Tissue-Tek optimal cutting temperature compound
(Miles, Elkhart, IN) and frozen in liquid nitrogen. Brain samples were cut into 10-μm-thick
coronal sections using a microtome (CM1850; Leica Microsystems, Wetzlar, Germany). The
sections were stained with the following antibodies: Calbindin (1:1000, rabbit, polyclonal,
Synaptic Systems Cat# 214 002, RRID:AB_2068199), Cyclin B (1:500, mouse, monoclonal,
Thermo Fisher Scientific Cat# MA1-155, RRID:AB_2536863), phospho-Histone H3 (1:500,
rabbit, polyclonal, Millipore Cat# 06-570, RRID:AB_310177), HA-tag (1:500, rabbit,
polyclonal, Cell Signaling Technology Cat# 3724, RRID:AB_1549585), ΔFosB (1:1000, rabbit,
monoclonal, Cell Signaling Technology Cat# 14695, RRID:AB_2798577), Calbindin (1:1000,
rabbit, polyclonal, Synaptic Systems Cat# 214002, RRID:AB_2068199), NeuN (1:500, mouse,
monoclonal, Millipore, Cat# MAB377X, RRID:AB_2149209), GFAP (1:500, rabbit,
polyclonal, Sigma-Aldrich, Cat# G9269, RRID:AB_477035), Iba1 (1:500, rabbit, polyclonal,
FUJIFILM, Cat# 019-19741, RRID:AB_839504), JunD (1:500, rabbit, polyclonal, Abcam,
Cat# ab28837, RRID:AB_2130167), and tri-methyl-Histone H3 (Lys9me3) (1:500, mouse,
monoclonal, FUJIFILM Cat# MABI0308). Before the first antibody reaction, antigen retrieval
was performed in 120℃ autoclave for 5 minutes in 0.01 M sodium citrate buffer (S1699,
Target Retrieval Solution, Dako). Autoclave at 120℃ is essential, especially for detecting
weak signals of nuclear antigens (e.g., pH3 and Cyclin B). This procedure also eliminates
EYFP/mCherry fluorescence, thus preventing overlap with secondary antibody staining with
Alexa 488/594. After antigen retrieval, sections were cooled at room temperature for 2 hours.
Sections were pre-incubated for 1 hour at room temperature in 5% skim milk in PBST for
blocking, and then incubated overnight at 4℃ in PBS containing the primary antibodies. The
next day, secondary antibody reactions were performed for 1 hour at room temperature using
Alexa Fluor 488-, Alexa Fluor 594-, and Alexa Fluor 647-conjugated secondary antibodies
(Invitrogen). Nuclear staining was performed using Hoechst 33258 (Polysciences, Warrington,
PA).
841
Immunostaining of brain sample with acetone fixation
For Lamin B1 staining (1:200, mouse, monoclonal, Thermo Fisher Scientific Cat# MA1-06103,
RRID:AB_2281281), a method omitting PFA perfusion was employed[113,114]. In this protocol,
brains were perfused with PBS alone, without PFA. Immediately after PBS perfusion, the
dissected brains were mounted in Tissue-Tek compound and frozen in liquid nitrogen. Brain

samples were cut into 10-μm-thick coronal sections using a microtome. Before the first

antibody staining, the brain sections were fixed with ice-cold acetone. Antigen retrieval before

the blocking step was omitted and subsequent procedures were performed in the same manner

as for immunostaining of the fixed brain samples.

TUNEL staining

In Situ Cell Death Detection Kit TMR red (Roche, 12156792910) was used for the detection

of apoptotic cells. Brain sections were rinsed in PBS, permeabilized with 0.1 % Tween for 30

min, and incubated with TUNEL reaction reagent (TdT Enzyme and Labeling Safe Buffer) and

Hoechst for 60 min at room temperature. Embryo brain tissue was used as the positive control.

The localization of TUNEL to the nucleus allows for accurate counting of dead or dying cells.

Image acquisition and analysis

Confocal microscopy (LSM 700; Zeiss) was used to obtain images of stained sections. For

image analysis, the regions of interest (ROIs) were manually delineated using ZEN (Zeiss) and

ImageJ software.

Behavioral tests

Open field test

The open field test was performed in an open field apparatus (40 cm × 40 cm × 30 cm; width,

depth, and height, respectively; O'Hara, Japan) made of opaque white plastic, with the center

area illuminated to 100 lux by an LED light attached above the ceiling. Before each behavioral

experiment, the open field apparatus was cleaned with weakly acidified hypochlorous water

(super-hypochlorous water; Shimizu Laboratory Supplies, Kyoto, Japan) to prevent bias due to

olfactory cues. All behavioral experiments were conducted in a soundproof room. During each

recording session, mouse movements were captured using a CCD camera above the open field

arena. Mouse behavior was monitored using a computer screen located outside the room to

minimize artifacts caused by the experimenter's presence. Mouse behavior was recorded at a

sampling rate of 10 Hz. Mouse images were automatically processed using the ImageJ plugin

(Image OF, freely available on the Mouse Phenotype Database website: http://www.mouse-

phenotype.org/software.html) to obtain the location-time sequence for each mouse. From the

traces of mouse movements, we calculated the total distance traveled by the mice during the

session[115].

Locomotor activity monitoring in home cage

Locomotor activity monitoring was performed on mice that had been operated at least one

month before. The system automatically analysed the locomotor activity of the mice in their

home cage[116]. The system contains a home cage (29 × 18 × 12 cm) and a filtered cage top,

separated by a 13-cm-high metal stand containing an infrared video camera, which is attached

to the top of the stand. Each mouse was individually housed in each home cage, and their

locomotor activity was monitored 24 hours a day. Outputs from the video cameras were fed

into a computer and images from each cage were captured at a rate of one frame per second.

The distance traveled was calculated automatically using the ImageJ plugin[115].

Tail suspension test

The tail suspension test was performed to assess depression-related behavior in mice. Each

mouse was suspended 30 cm above the floor by the tail in a white plastic chamber (44 cm

height × 49 cm length × 32 cm width; O'Hara, Japan) with a video camera mounted on the wall

(O'Hara, Japan). The behavior was recorded for 10 minutes. Images were captured at two

frames per second using a video camera and transferred to a computer. The area (pixels) within

which the mouse moved was measured for each pair of successive frames. When the area was

below a certain threshold, the mouse behavior was judged as "immobile." When the area

equaled or exceeded the threshold, the mouse was considered "moving." The optimal threshold

for this judgment was determined by adjusting it to the amount of immobility measured by a

trained human observer. Immobility lasting for < 2 seconds was not included in the analysis.

Data acquisition and analysis were performed automatically using the ImageJ plugin

software[115].

Porsolt forced swim test

The Porsolt forced swim test was performed to assess depression-related behavior. A Plexiglas

cylinder (20 cm height × 10 cm diameter) was placed in a test chamber (49 cm height × 44 cm

length × 32 cm width; O'Hara, Japan). A video camera was mounted on the ceiling of the test

chamber and positioned directly above the cylinder. The mice were placed into a cylinder,

which was filled with water (approximately 23°C) to a height of 7.5 cm.

The immobility time was recorded for 10 minutes. Images were captured at two frames per

second through the video camera. The immobility status of each mouse was determined in the

same manner as for the tail suspension test. Data acquisition and analysis were performed using

the ImageJ plugin[115].

Contextual and cued fear-conditioning tests

Fear conditioning tests were conducted to assess contextual memory (Extended Data Fig. 3a).

In the conditioning session, each mouse was placed in an acrylic chamber consisting of white

(side) and transparent (front, rear, and top) walls (33 × 25 × 28 cm) with a stainless-steel grid

floor (0.2 cm diameter, spaced 0.5 cm apart; O'Hara, Japan). The mice were allowed to explore

the chamber freely for 120 seconds, and 55 dB white noise then served as the conditioned

stimulus (CS) for 30 seconds. During the last 2 seconds of CS presentation, a mild foot shock

(0.3 mA, 2 seconds) was delivered as the unconditioned stimulus (US). The mice were

subjected to two more CS-US pairings with a 2-minutes interstimulus interval. Approximately

24 hours or one month after conditioning, a context test was conducted for 300 seconds. In the

context test, mice were placed in the same chamber in which they had been conditioned. After

the context test, a cued test with an altered context was performed for 360 seconds. In the cued

test, the mice were placed in a triangular chamber (33 × 29 × 32 cm) made of white plastic

walls and floor, which was located in a different sound-attenuating room and allowed to explore

the triangular chamber for 180 seconds. Then, the CS was presented during the last 180 seconds

of the cued test. In each session, the percentage of freezing and distance traveled was calculated

automatically using the ImageJ plugin[115].

Three chamber social interaction test

The testing apparatus consisted of a rectangular three-chambered box and lid with a video

camera (O'Hara, Japan; Extended Data Fig. 3b). The dividing walls of the chamber were made

of transparent plastic, with small square openings (5 × 3 cm) allowing access to each chamber

(20 × 40 × 47 cm). A small round wire cage (9 cm in diameter, 11 cm in height, with vertical

bars 0.5 cm apart) was located in the corner of chambers. The test mice were first placed in the

middle chamber and allowed to explore the entire test chamber for 10 minutes. Immediately

after the 10-minute period, the test mice were placed in a clean holding cage, and a male

C57BL/6J mouse (stranger mouse) with no prior contact with the test mice was enclosed in

one of the wire cages. The test mice were returned to the middle chamber and allowed to

explore for 10 minutes. The time spent in each chamber and the time spent around each cage

was automatically measured using ImageCSI software[115].

in vivo Ca^{2+} imaging with optogenetic stimulation

Viral constructs

Purified and concentrated adeno-associated virus serotype 5 coding GCaMP6f under the

synapsin promoter (AAV5-syn-GCaMP6f; #100837-AAV5) and adeno-associated virus

serotype 5 coding for ChrimsonR-tdTomato under the synapsin promoter (AAV5-syn-

ChrimsonR-tdT) were obtained from Addgene and UNC Vector Core Facility.

Viral delivery of Ca^{2+} sensor and opsin

For delivery of a fluorescent Ca^{2+} sensor and red-shifted opsin into dDG neurons, AAV5-syn-

GCaMP6f and AAV5-syn-ChrimsonR-tdT were injected into the dorsal DG in adult mice (left

hemisphere; 2.0 mm posterior to bregma [AP], 1.0–1.45 mm lateral to midline [ML], and 2.0

mm ventral to bregma [DV]). Briefly, adult mice (> 8 weeks old) were anesthetized with 1.5

to 3% isoflurane at an oxygen flow rate of 1 L/min. The head fur was shaved and the incision

site was sterilized with 70% ethanol prior to the surgical procedure. Mice were mounted on a

stereotaxic device (catalog no. 51730D, Stoelting), and heat pad (catalog no. BWT-100A,

BRC) was placed underneath each mouse to maintain body temperature at 37 °C. After the

scalp was incised and pulled aside, a 2-mm-diameter craniotomy was created with a surgical

drill on the skull above the injection site. Through a glass-capillary injection pipette (25-µm

inner-diameter tip), a mixture of 300nl of AAV5-Syn-GCaMP6f-WPRE-SV40 (titer 4.3×10^{12}

GC/mL) and AAV5-syn-ChrimsonR-tdT (titer 1.53×10^{12} GC/mL) was injected using a

microinjection pump (Nanoliter 2010, WPI). After the injection, the scalp was sutured and

treated with povidone–iodine. The expression of GCaMP6f is driven by the Syn promoter and

is expected to be induced not only in GCs but also in other cell types. We previously reported

that the majority (approximately 90%) of neurons expressing GCaMP in the DG are GCs[16].

GRIN lens implantation

At least two weeks after viral injection, a gradient refractive index (GRIN) lens (Inscopix

catalog no.1050-004623, 1 mm diameter) was implanted at the same position where the virus

was injected. The mouse was mounted on a stereotaxic device and the scalp was removed.

After exposing the skull and removing the overlying connective tissue, we made a cranial hole

slightly larger than the diameter of the GRIN lens. To secure for an entry path for the GRIN

lens, the cortex and white matter were aspirated using a 26-gauge blunt needle attached to a

vacuum pump; to prevent aspiration of the GCs, the depth of needle tip depth was set at 1.6

mm from the bregma and the tissue below was carefully aspirated. Saline was continuously

applied during aspiration to prevent the tissue from drying. During implantation, the lenses

were held in place and slowly lowered using a ProView Implant kit (nVoke, Inscopix, Palo Alto,

California, USA) attached to the stereotaxic arm. The GRIN lens connected to the manual

stereotaxic arm was slowly lowered (5–10 μm/sec) until the field of view was in focus,

referring to the fluorescence intensity of the GCaMP. After temporarily immobilizing the lens

with ultraviolet-curable resin (Primefil, Tokuyama Dental), it was fixed to the exposed skull

using dental cement（Super-Bond C&B, Sun Medical）. The exposed tip of the GRIN lens was

covered with dental silicone (Dent Silicone-V, Shofu). After lens implantation, analgesic and

anti-inflammatory agents (Flunixin, 2 mg/kg; Fujita) and antibiotics (Tribrissen, 0.12 mL/kg;

Kyoritsu Seiyaku) were injected intraperitoneally.

Baseplate attachment

At least two weeks after GRIN lens implantation, the baseplate of a miniature microendoscope

(nVoke, Inscopix) was attached to the GRIN lens using a previously described method[16].

Briefly, the mice were anesthetized and mounted onto a stereotaxic device, and a baseplate

attached to the miniature microscope was placed on the GRIN lens using a gripper (nVoke

accessory, Inscopix). The optimal location was determined by monitoring the fluorescent

images of GCaMP-expressing neurons (where the largest number of neurons was in focus),

and the baseplate was fixed with ultraviolet-curable resin and dental cement at this position. In

cases where we failed to identify neurons at this stage (usually due to the failure of GRIN-lens

implantation), the baseplate was not attached, and such mice were excluded from the

experiment. After the baseplate attachment, a baseplate cover was placed on the baseplate until

Ca^{2+} imaging was performed.

1010

Ca^{2+} imaging and optogenetic stimulation in freely moving mice

Ca^{2+} imaging of the DG neurons was performed while the mice traveled freely in the OF. Prior

to Ca^{2+} imaging, the mice were lightly anesthetized and a miniature microscope (nVoke,

Inscopix) was mounted onto the baseplate of the mice. The mice were then placed back in their

home cages and transferred to a sound-proof behavioral experiment room. At least 30 min after

recovery from anesthesia, the mice were subjected to OF. The Ca^{2+} signals of their DG neurons

were obtained at a 10-Hz sampling rate with a 1,440×l,080-pixel resolution. Recordings per

day totaled 50 min, consisting of before (0–30 min), during (30–35 min), and after (35–50 min)

optogenetic stimulation; no stimulation was introduced in the control group. These 50-min

recordings were performed daily throughout the experimental period, and the first 30 min

recordings were used for data analysis. The experimental period consisted of before stimulation

(Day 1), during stimulation (Day 2-11), and two weeks after stimulation (Day 24-26). We

applied EX-LED (filtered with a 435–460 nm excitation filter) for the excitation of GCaMP6f

fluorescence (0.4 mW/mm^2 at the bottom of the GRIN lens) and OG-LED (filtered with a 590–

650 nm excitation filter) for the excitation of ChrimsonR (10 mW/mm^2 at the bottom of the

GRIN lens). A previous study reported that EX-LED at 2.0 mW/mm^2 resulted in small current

changes of approximately 10 pA via activation of ChrimsonR under slice preparation

(approximately 130 pA by OG-LED irradiation)[117]. Given that EX-LED used in this study are

much lower than these conditions (–0.4 mW/mm^2) and that under *in vivo* imaging conditions

the EX-LED are scattered in the brain tissue and the current changes are much smaller than in

slice preparation[117], unintended activation of ChrimsonR by EX-LED would be negligible.

1032

Detection of Ca^{2+} transients

To extract the activity patterns of individual DG neurons from the obtained fluorescent images,

we used Inscopix Data Processing Software (IDPS 1.8.0). Briefly, the data files of the raw

sequential fluorescent images obtained by nVoke (Inscopix) were imported to the IDPS, and

movie trimming, data preprocessing, spatial filtering, and motion correction were applied. The

instantaneous fluorescence of the motion-corrected images was then normalized by its average

fluorescence over the entire recording period, producing fluorescence-change ratio ($\Delta F/F$)

images. Individual DG neurons were identified using the automated principal component

analysis (PCA) and independent component analysis (ICA) segmentation of activity traces. For

PCA/ICA segmentation, default parameters, whose values were confirmed to work well across

most of the neuronal activity patterns in the cortex and CA1 of the hippocampus, were used

(number of independent components, 120; number of principal components, 150; ICA max

iterations, 100; ICA random seed, 0; ICA convergence threshold, 0.00001; block size, 1,000;

ICA unmixing dimension, spatial; ICA temporal weights, 0.00; and arbitrary units within the

IDPS). Finally, the temporal pattern of each timing of DG neuron activity was identified by an

event-detection algorithm available in IDPS (event threshold factor, four median absolute

deviations; event shortest decay time, 0.2 s), and the temporal pattern of activity was expressed

as a time series of binarized signals for further analysis.

1051

1052

1053

1054 ***Information coding in individual neurons***

1055

Definitions of the position and speed of mice

(i) Position: The OF arena (which had an area of 40 cm × 40 cm) was represented as 200 ×

200-pixel grid. The position of the mouse was determined from the centroid of its shadow on

the CCD camera, above the OF. We then assigned a label corresponding to the discrete location

of the mouse (e.g., [10, 100]) to each time bin (=1/10sec).

(ii) Speed: From the distance traveled between 1 s before and after a given time point, we

calculated the speed of the mouse at that moment and assigned this speed (cm/s) to each time

bin.

1064

Statistical analysis of spatial and speed information

To quantify the tuning specificities of neurons with respect to the position and speed of the

mouse, we measured their specificity in terms of the information rate of cell activity, and

defined them as (i) spatial and (ii) speed information[16,118]. The Ca^{2+} event rate in Ca^{2+} imaging

was considerably lower than that in electrophysiological recordings (average, approximately

0.05 Hz). If the discretization of position and speed is too fine for the number of events in the

recorded cells, a proper null distribution cannot be obtained when creating shuffled data for

that cell. Therefore, we set the resolution of the discretization of the position and speed to be

lower than those that are commonly used.

(i) We used a 2×2 square grid to measure spatial information and computed the amount of

Shannon information conveyed by a single Ca^{2+} transient about the animal's position. The

spatial information I (bits per Ca^{2+} transient) of a cell was calculated as the mutual information

score between the occurrence of a single Ca^{2+} transient in the cell and the animal's behavioral

state of position using the following formula:

$$I_{(\text{bits/Ca}^{2+} \text{ transient})} = \sum_{i=1}^{N} p_i \frac{r_i}{r} \log_2 \frac{r_i}{r}$$

1079

where i is the bin number corresponding to the physical parameter (in this case, spatial position

in the OF; i = 1–4 from the 2×2 square grid), N is the total number of bins, p is the probability

that the mouse occupied bin i, r_i is the mean transient rate at bin i, and r is the overall mean

transient rate.

(ii) To measure speed information, we applied the same formula to the speed after discretizing

it to a binary state: Run (speed > 1 cm/sec) and Stop (speed < 1 cm/sec).

1086

Definitions of place cells and speed cells

To evaluate the statistical significance of the information encoded in individual neurons, we

employed the random permutation method as previously outlined[16]. The calcium event data

were divided into 1000 segments along the time axis and randomly shuffled to generate
permuted calcium event data. This method disrupts the temporal structures of neural activity
and the temporal correlations between neural activity and behavioral variables. Using this
randomly permuted data, we derived a null distribution for the position and speed information
of neurons. Neurons with original information content exceeding the top 95% and 99% of these
null distributions were classified as place or speed cells, respectively, indicating cells with
significant spatial or speed information[119,120].

1097
1098
1099

Decoding position and speed in the open field
To determine how OF behavioral parameters are encoded in the population activities of DG
neurons, we trained decoders using machine learning methods to predict the position and speed
of mice from the population Ca^{2+} activity. We assigned labels to the discretized behavioral
parameters of the mouse (position (cm) and speed (cm/s)) and the binary values of the Ca^{2+}
signal (0 or 1) of all neurons in each time bin. We then divided the Ca^{2+} imaging data and
behavioral data of the first 30 min (before 5-min of optogenetic stimulation each day) into the
first 15 min and last 15 min halves, which were designated as training and test data, respectively.
For each pair of behavioral parameters (position and speed) and the value of the Ca^{2+} signal in
the training data, we trained the decoders using a machine learning method (the code was
obtained and modified from Glaser *et al*., eNeuro, 2020[121]). The decoding accuracy of the
position was reported as the mean absolute error of the distance between the predicted and
actual positions. In the case of speed, decoding accuracy was reported as the correlation
between the predicted and actual instantaneous speeds. This is because speed, unlike position,
is not limited to a certain range and the mean absolute error of speed depends on the average
locomotion speed of each mouse, which is inappropriate for the evaluation of the decoding
error. If the number of neurons recorded in each session was less than 10, the session was
excluded from the decoding analysis.

1118
1119
1120

DNA Constructs and AAV Vectors for in vivo genome editing
For SpCas9 target selection and single-guide RNA (sgRNA) generation, 20-nt target sequences
were selected to precede a 5'-NGG protospacer-adjacent motif (PAM) sequence. To minimize
off-target effects, CRISPR design tool was used (https://crispr.dbcls.jp/). The backbone

plasmid for sgRNA (PX552; pAAV-U6sgRNA(SapI)_hSyn-GFP-KASH-bGH) was obtained

from Addgene. *Ccnb*-sgRNA (*Ccnb1* and *Ccnb2* were tandemly connected; Fig. 5b) and

Scramble-sgRNA nucleotides were synthesized and cloned into AAV backbone plasmids

(Vigene Biosciences, Inc, USA). To avoid fluorescence overlap with ChR2-EYFP, the GFP

cassette was replaced with mCherry (Fig. 5b). The plasmid for SpCas9 was obtained from

Addgene (PX551; pAAV-pMecp2-SpCas9-spA). Concentrated AAV vectors were produced

from plasmids for *Ccnb*-sgRNA, *Scramble*-gRNA, and SpCas9, with support from the Gunma

University Initiative for Advanced Research Viral Vector Core.

1133

1134

1135

Stimulated Emission Depletion Microscopy (STED) observation

Brain slices were double-immunostained with anti-tri-methyl-Histone H3 (Lys9me3) antibody

(1:500, mouse, monoclonal, MAI0308, FUJIFILM) and anti-phospho-Histone H3 antibody

(1:500, rabbit, polyclonal #06-570, Cell Signaling) as the primary antibodies, followed by

Alexa Fluor 488- and Alexa Fluor 555- conjugated antibodies (Molecular Probes) as the

secondary antibodies. STED images were acquired using a TCS SP8 STED 3X system (Leica

Microsystems, Wetzlar, Germany) with 2ch HyDSMD detectors[122]. A white laser (for

excitation) and a 660 nm laser with a donut beam (for stimulated emission depletion) were used

for STED imaging. For the detection of signals of tri-methyl-Histone H3 (Lys9me3) and

phospho-Histone H3, we used excitation wavelengths of 488 and 561 nm, respectively. Raw

STED images (pixel size:20 nm) were recorded with a 93 × glycerol immersion objective

(Leica HC PL APO 93 × /1.30 GLYC motC STED W) and deconvoluted using the Huygens

software (SVI, Hilversum, Netherlands). Image analysis was performed with a custom-

developed code using the image processing toolbox in MATLAB (MathWorks, R2022a).

1150

1151

1152

Data Collection, Pre-processing, and Meta-analysis

All microarray and RNA-seq data used for meta-analysis in this study were obtained from the

BaseSpace Correlation Engine (https://japan.ussc.informatics.illumina.com/c/nextbio.nb;

Illumina, Cupertino, CA), a publicly available database containing over 100,000 microarray

and RNA-seq datasets. The details of all datasets are described in Supplementary Table1. The

datasets registered in BaseSpace underwent several preprocessing, quality control, and

organization stages. Quality control ensures the integrity of samples and datasets. Genes with

a P value < 0.05 (without correction for multiple testing) were included in the differentially

expressed gene datasets. This sensitivity threshold is typically the lowest used with commercial

microarray platforms and the default criterion in BaseSpace analyses[44]. Correction for multiple

testing was omitted to minimize false negatives at this stage. Expression values were used to

calculate fold changes and P values between the two conditions (infants–adults and treated–

untreated). To determine fold changes, we divided the expression values of the probes/genes

in the test datasets by those in the control datasets. If the fold change was < 1.0, these values

were converted to a negative reciprocal or −1/(fold change). Genes with a t-test P value < 0.05

were imported into the BaseSpace Correlation Engine according to the instructions provided

by the manufacturer. The rank order of these genes was determined based on their absolute

fold-changes. All meta-statistical analyses comparing two datasets were performed in

BaseSpace, and the similarities between any two datasets were evaluated as overlap P values

using the Running Fisher algorithm[44].

1173

1174

1175

Statistical Analyses

Mice of the same age and sex were raised under identical conditions and were randomly

allocated to different experimental groups. Blinding was not applied in this study because the

animals needed to be controlled by the stimulation conditions. The sample sizes were

determined based on those reported in similar publications[30,31]. Welch's t-test was used to

compare the averages. An ANOVA was performed when more than two groups were compared,

followed by the Bonferroni post-hoc method. For correlation analysis, we used Pearson

correlation analysis for normally distributed data. When pooling the data of neurons from

multiple mice per group, a Linear Mixed-Effects Regression model was applied to account for

individual differences per mouse. All results of the statistical analyses are summarized in

Supplementary Table 2.

1187

1188

1189

1190 ***Code and Data Availability***

1191 RNA-seq and ATAC-seq data used in this study were deposited in the GEO database with

1192 accession codes GSE227200 (RNAseq) and GSE227201 (ATAC-seq), respectively. The RNA-

1193 seq data for the post-mortem DG of human patient was obtained from SRP241159. The

1194 personal information regarding ECT history was provided by Astellas Pharma Inc. The raw

1.   Choi, D. W. Excitotoxic cell death. *Journal of Neurobiology* **23**, 1261–
1276 (1992).

2.   Dong, X., Wang, Y. & Qin, Z. Molecular mechanisms of excitotoxicity
and their relevance to pathogenesis of neurodegenerative diseases. *Acta Pharmacol Sin* **30**,
379–387 (2009).

3.   Singh, S. P., He, X., McNamara, J. O. & Danzer, S. C. Morphological
changes among hippocampal dentate granule cells exposed to early kindling-epileptogenesis.
Hippocampus **23**, 1309–1320 (2013).

4.   Lepeta, K. *et al.* Synaptopathies: synaptic dysfunction in neurological
disorders – A review from students to students. *Journal of Neurochemistry* **138**, 785–805
(2016).

5.   Holmes, G. L. & Ben-Ari, Y. The Neurobiology and Consequences of
Epilepsy in the Developing Brain. *Pediatr Res* **49**, 320–325 (2001).

6.   Pan, G. *et al.* Compensatory Mechanisms Modulate the Neuronal
Excitability in a Kainic Acid-Induced Epilepsy Mouse Model. *Frontiers in Neural Circuits* **12**,
(2018).

7.   Hagihara, H., Takao, K., Walton, N. M., Matsumoto, M. & Miyakawa, T.
Immature dentate gyrus: an endophenotype of neuropsychiatric disorders. *Neural Plasticity*
2013, e318596 (2013).

8.   Shin, R. *et al.* The immature dentate gyrus represents a shared phenotype
of mouse models of epilepsy and psychiatric disease. *Bipolar Disorders* **15**, 405–421 (2013).

9.   Imoto, Y., Segi-Nishida, E., Suzuki, H. & Kobayashi, K. Rapid and stable
changes in maturation-related phenotypes of the adult hippocampal neurons by
electroconvulsive treatment. *Mol Brain* **10**, 8 (2017).

10.   Ueno, M. *et al.* The effect of electroconvulsive seizure on survival,
neuronal differentiation, and expression of the maturation marker in the adult mouse
hippocampus. *J. Neurochem.* (2019) doi:10.1111/jnc.14691.

11.   Murano, T., Hagihara, H., Tajinda, K., Matsumoto, M. & Miyakawa, T.
Transcriptomic immaturity inducible by neural hyperexcitation is shared by multiple
neuropsychiatric disorders. *Commun Biol* **2**, 1–11 (2019).

12.   Yamasaki, N. *et al.* Alpha-CaMKII deficiency causes immature dentate
gyrus, a novel candidate endophenotype of psychiatric disorders. *Molecular Brain* **1**, 6 (2008).

13.   Takao, K. *et al.* Deficiency of Schnurri-2, an MHC Enhancer Binding
Protein, Induces Mild Chronic Inflammation in the Brain and Confers Molecular, Neuronal,
and Behavioral Phenotypes Related to Schizophrenia. *Neuropsychopharmacology* **38**, 1409–

1425 (2013).

14.   Umemori, J., Winkel, F., Didio, G., Pou, M. L. & Castrén, E. iPlasticity:
Induced juvenile-like plasticity in the adult brain as a mechanism of antidepressants. *Psychiatry*
Clin Neurosci **72**, 633–653 (2018).

15.   Hagihara, H. *et al.* Forebrain-specific conditional calcineurin deficiency
induces dentate gyrus immaturity and hyper-dopaminergic signaling in mice. *Molecular Brain*
15, 94 (2022).

16.   Murano, T. *et al.* Multiple types of navigational information are
independently encoded in the population activities of the dentate gyrus neurons. *Proceedings*
of the National Academy of Sciences **119**, e2106830119 (2022).

17.   Hagihara, H. *et al.* Large-scale animal model study uncovers altered brain
pH and lactate levels as a transdiagnostic endophenotype of neuropsychiatric disorders
involving cognitive impairment. *eLife* **12**, RP89376 (2024).

18.   Li, B. *et al.* Failure of Neuronal Maturation in Alzheimer Disease Dentate
Gyrus. *J Neuropathol Exp Neurol* **67**, 78–84 (2008).

19.   Walton, N. M. *et al.* Detection of an immature dentate gyrus feature in
human schizophrenia/bipolar patients. *Transl Psychiatry* **2**, e135 (2012).

20.   Hagihara, H., Ohira, K., Takao, K. & Miyakawa, T. Transcriptomic
evidence for immaturity of the prefrontal cortex in patients with schizophrenia. *Molecular*
Brain **7**, 41 (2014).

21.   You, J. C. *et al.* Epigenetic suppression of hippocampal calbindin-D28k
by ΔFosB drives seizure-related cognitive deficits. *Nature Medicine* **23**, 1377 (2017).

22.   Murano, T., Koshimizu, H., Hagihara, H. & Miyakawa, T. Transcriptomic
immaturity of the hippocampus and prefrontal cortex in patients with alcoholism. *Sci Rep* **7**,
44531 (2017).

23.   Tavitian, A., Song, W. & Schipper, H. M. Dentate Gyrus Immaturity in
Schizophrenia. *Neuroscientist* 1073858418824072 (2019) doi:10.1177/1073858418824072.

24.   May, A. Experience-dependent structural plasticity in the adult human
brain. *Trends in Cognitive Sciences* **15**, 475–482 (2011).

25.   Ganguly, K. & Poo, M. Activity-Dependent Neural Plasticity from Bench
to Bedside. *Neuron* **80**, 729–741 (2013).

26.   Malenka, R. C. & Bear, M. F. LTP and LTD: An Embarrassment of Riches.
Neuron **44**, 5–21 (2004).

27.   Citri, A. & Malenka, R. C. Synaptic Plasticity: Multiple Forms, Functions,
and Mechanisms. *Neuropsychopharmacol* **33**, 18–41 (2008).

28.   Zhang, W. & Linden, D. J. The other side of the engram: experience-driven
changes in neuronal intrinsic excitability. *Nat Rev Neurosci* **4**, 885–900 (2003).

29.   Titley, H. K., Brunel, N. & Hansel, C. Toward a Neurocentric View of
Learning. *Neuron* **95**, 19–32 (2017).

30.   Su, Y. *et al.* Neuronal activity modifies the chromatin accessibility
landscape in the adult brain. *Nat Neurosci* **20**, 476–483 (2017).

31.   Fernandez-Albert, J. *et al.* Immediate and deferred epigenomic signatures
of in vivo neuronal activation in mouse hippocampus. *Nature Neuroscience* **22**, 1718–1730
(2019).

32.   Espinoza, R. T. & Kellner, C. H. Electroconvulsive Therapy. *New England*
Journal of Medicine **386**, 667–672 (2022).

33.   Salik, I. & Marwaha, R. Electroconvulsive Therapy. in *StatPearls*
(StatPearls Publishing, Treasure Island (FL), 2024).

34.   Oltedal, L. *et al.* Volume of the Human Hippocampus and Clinical
Response Following Electroconvulsive Therapy. *Biological Psychiatry* **84**, 574–581 (2018).

35.   Nuninga, J. O. *et al.* Volume increase in the dentate gyrus after
electroconvulsive therapy in depressed patients as measured with 7T. *Molecular Psychiatry* **25**,
1559–1568 (2020).

36.   Loef, D. *et al.* Electroconvulsive therapy is associated with increased
immunoreactivity of neuroplasticity markers in the hippocampus of depressed patients. *Transl*
Psychiatry **13**, 1–11 (2023).

37.   Zhao, C., Warner-Schmidt, J., S. Duman, R. & Gage, F. H.
Electroconvulsive seizure promotes spine maturation in newborn dentate granule cells in adult
rat. *Devel Neurobio* **72**, 937–942 (2012).

38.   Kobayashi, K. *et al.* Rapid and lasting enhancement of dopaminergic
modulation at the hippocampal mossy fiber synapse by electroconvulsive treatment. *Journal*
of Neurophysiology **117**, 284–289 (2017).

39.   Kobayashi, K. *et al.* Reversal of hippocampal neuronal maturation by
serotonergic antidepressants. *Proc. Natl. Acad. Sci. U.S.A.* **107**, 8434–8439 (2010).

40.   Ohira, K., Hagihara, H., Miwa, M., Nakamura, K. & Miyakawa, T.
Fluoxetine-induced dematuration of hippocampal neurons and adult cortical neurogenesis in
the common marmoset. *Mol Brain* **12**, 69 (2019).

41.   Nguyen, P. T. *et al.* Antidepressants reactivate developmental plasticity
through   remodeling   of   extracellular   matrix.   2025.01.03.631260   Preprint   at
https://doi.org/10.1101/2025.01.03.631260 (2025).

42.   Lefaucheur, J.-P. *et al.* Evidence-based guidelines on the therapeutic use
of repetitive transcranial magnetic stimulation (rTMS): An update (2014–2018). *Clinical*
Neurophysiology **131**, 474–528 (2020).

43.   Shiozawa, P. *et al.* Transcranial direct current stimulation for major
depression: an updated systematic review and meta-analysis. *International Journal of*
Neuropsychopharmacology **17**, 1443–1452 (2014).

44.   Kupershmidt, I. *et al.* Ontology-Based Meta-Analysis of Global
Collections of High-Throughput Public Data. *PLoS ONE* **5**, e13066 (2010).

45.   Jaffe, A. E. *et al.* Profiling gene expression in the human dentate gyrus
granule cell layer reveals insights into schizophrenia and its genetic risk. *Nat Neurosci* 1–10
(2020) doi:10.1038/s41593-020-0604-z.

46.   Kang, H. J. *et al.* Spatio-temporal transcriptome of the human brain.
Nature **478**, 483–489 (2011).

47.   Klemm, S. L., Shipony, Z. & Greenleaf, W. J. Chromatin accessibility and
the regulatory epigenome. *Nat Rev Genet* **20**, 207–220 (2019).

48.   Kheirbek, M. A. *et al.* Differential Control of Learning and Anxiety along
the Dorsoventral Axis of the Dentate Gyrus. *Neuron* **77**, 955–968 (2013).

49.   Stefanini, F. *et al.* A Distributed Neural Code in the Dentate Gyrus and in
CA1. *Neuron* (2020) doi:10.1016/j.neuron.2020.05.022.

50.   Whitfield, M. L. *et al.* Identification of Genes Periodically Expressed in
the Human Cell Cycle and Their Expression in Tumors. *MBoC* **13**, 1977–2000 (2002).

51.   Dechat, T. *et al.* Nuclear lamins: major factors in the structural
organization and function of the nucleus and chromatin. *Genes Dev.* **22**, 832–853 (2008).

52.   Gruenbaum, Y. & Foisner, R. Lamins: nuclear intermediate filament
proteins with fundamental functions in nuclear mechanics and genome regulation. *Annu Rev*
Biochem **84**, 131–164 (2015).

53.   Maison, C. & Almouzni, G. HP1 and the dynamics of heterochromatin
maintenance. *Nat Rev Mol Cell Biol* **5**, 296–305 (2004).

54.   Probst, A. V. & Almouzni, G. Heterochromatin establishment in the
context of genome-wide epigenetic reprogramming. *Trends in Genetics* **27**, 177–185 (2011).

55.   Heinz, S. *et al.* Simple Combinations of Lineage-Determining
Transcription Factors Prime *cis*-Regulatory Elements Required for Macrophage and B Cell
Identities. *Molecular Cell* **38**, 576–589 (2010).

56.   Chen, J., Kelz, M. B., Hope, B. T., Nakabeppu, Y. & Nestler, E. J. Chronic
Fos-Related Antigens: Stable Variants of ΔFosB Induced in Brain by Chronic Treatments. *J.*

Neurosci. **17**, 4933–4941 (1997).

57.   Herrup, K. & Yang, Y. Cell cycle regulation in the postmitotic neuron: oxymoron or new biology? *Nat Rev Neurosci* **8**, 368–378 (2007).

58.   Frost, B. Alzheimer's disease: An acquired neurodegenerative laminopathy. *Nucleus* **7**, 275–283 (2016).

59.   Mertens, J. *et al.* Age-dependent instability of mature neuronal fate in induced neurons from Alzheimer's patients. *Cell Stem Cell* **28**, 1533-1548.e6 (2021).

60.   Nagy, Z., Esiri, M. M., Cato, A.-M. & Smith, A. D. Cell cycle markers in the hippocampus in Alzheimer's disease. *Acta Neuropathologica* **94**, 6–15 (1997).

61.   Busser, J., Geldmacher, D. S. & Herrup, K. Ectopic Cell Cycle Proteins Predict the Sites of Neuronal Cell Death in Alzheimer's Disease Brain. *J. Neurosci.* **18**, 2801–2807 (1998).

62.   Vincent, I., Jicha, G., Rosado, M. & Dickson, D. W. Aberrant Expression of Mitotic Cdc2/Cyclin B1 Kinase in Degenerating Neurons of Alzheimer's Disease Brain. *J. Neurosci.* **17**, 3588–3598 (1997).

63.   Yang, Y., Mufson, E. J. & Herrup, K. Neuronal Cell Death Is Preceded by Cell Cycle Events at All Stages of Alzheimer's Disease. *J. Neurosci.* **23**, 2557–2563 (2003).

64.   Nagy, Z. & Esiri, M. M. Neuronal cyclin expression in the hippocampus in temporal lobe epilepsy. *Exp Neurol* **150**, 240–247 (1998).

65.   Hernández-Ortega, K. & Arias, C. ERK activation and expression of neuronal cell cycle markers in the hippocampus after entorhinal cortex lesion. *Journal of Neuroscience Research* **90**, 2116–2126 (2012).

66.   Pavlova, T., Stepanichev, M. & Gulyaeva, N. Pentylenetetrazole kindling induces neuronal cyclin B1 expression in rat hippocampus. *Neuroscience Letters* **392**, 154–158 (2006).

67.   Love, S. Neuronal expression of cell cycle-related proteins after brain ischaemia in man. *Neuroscience Letters* **353**, 29–32 (2003).

68.   Jordan-Sciutto, K. L., Wang, G., Murphey-Corb, M. & Wiley, C. A. Cell Cycle Proteins Exhibit Altered Expression Patterns in Lentiviral-Associated Encephalitis. *J. Neurosci.* **22**, 2185–2195 (2002).

69.   Frost, B., Bardai, F. H. & Feany, M. B. Lamin dysfunction mediates neurodegeneration in tauopathies. *Current Biology* **26**, 129–136 (2016).

70.   Ogawa, O. *et al.* Ectopic localization of phosphorylated histone H3 in Alzheimer's disease: a mitotic catastrophe? *Acta Neuropathol* **105**, 524–528 (2003).

71.   Evangelisti, C. *et al.* The wide and growing range of lamin B-related

diseases: from laminopathies to cancer. *Cell. Mol. Life Sci.* **79**, 126 (2022).

72.   Ripke, S. *et al.* Biological insights from 108 schizophrenia-associated
genetic loci. *Nature* **511**, 421–427 (2014).

73.   Trubetskoy, V. *et al.* Mapping genomic loci implicates genes and synaptic
biology in schizophrenia. *Nature* **604**, 502–508 (2022).

74.   Cascella, N. G., Schretlen, D. J. & Sawa, A. Schizophrenia and epilepsy:
Is there a shared susceptibility? *Neuroscience Research* **63**, 227–235 (2009).

75.   Adachi, N. & Ito, M. Epilepsy in patients with schizophrenia:
Pathophysiology and basic treatments. *Epilepsy & Behavior* **127**, 108520 (2022).

76.   Vossel, K. A., Tartaglia, M. C., Nygaard, H. B., Zeman, A. Z. & Miller, B.
L. Epileptic activity in Alzheimer's disease: causes and clinical relevance. *The Lancet*
Neurology **16**, 311–322 (2017).

77.   Targa Dias Anastacio, H., Matosin, N. & Ooi, L. Neuronal
hyperexcitability in Alzheimer's disease: what are the drivers behind this aberrant phenotype?
Transl Psychiatry **12**, 1–14 (2022).

78.   Frade, J. M. & Ovejero-Benito, M. C. Neuronal cell cycle: the neuron itself
and its circumstances. *Cell Cycle* **14**, 712–720 (2015).

79.   Walton, C. C. *et al.* Primary neurons can enter M-phase. *Scientific Reports*
9, 4594 (2019).

80.   Poleshko, A. *et al.* H3K9me2 orchestrates inheritance of spatial
positioning of peripheral heterochromatin through mitosis. *eLife* **8**, e49278 (2019).

81.   Nestler, E. J., Barrot, M. & Self, D. W. ΔFosB: A sustained molecular
switch for addiction. *Proceedings of the National Academy of Sciences* **98**, 11042–11046
(2001).

82.   Nestler, E. J. Molecular basis of long-term plasticity underlying addiction.
Nat Rev Neurosci **2**, 119–128 (2001).

83.   Corbett, B. F. *et al.* ΔFosB Regulates Gene Expression and Cognitive
Dysfunction in a Mouse Model of Alzheimer's Disease. *Cell Reports* **20**, 344–355 (2017).

84.   Pande, G., Kumar, N. A. & Manogaran, P. S. Flow cytometric study of
changes in the intracellular free calcium during the cell cycle. *Cytometry* **24**, 55–63 (1996).

85.   Borowiec, A.-S. *et al.* Calcium channels, external calcium concentration
and cell proliferation. *European Journal of Pharmacology* **739**, 19–25 (2014).

86.   Yun, S. *et al.* Stimulation of entorhinal cortex–dentate gyrus circuitry is
antidepressive. *Nature Medicine* **24**, 658–666 (2018).

87.   Tunc-Ozcan, E. *et al.* Activating newborn neurons suppresses depression

and anxiety-like behaviors. *Nat Commun* **10**, 3768 (2019).

88. Sackeim, H. A. *et al.* The Cognitive Effects of Electroconvulsive Therapy in Community Settings | Neuropsychopharmacology. *Neuropsychopharmacology* **32**, 244–254 (2007).

89. O'Connor, M. *et al.* A Dissociation Between Anterograde and Retrograde Amnesia After Treatment With Electroconvulsive Therapy: A Naturalistic Investigation. *The Journal of ECT* **24**, 146 (2008).

90. Shepherd, J. D. *et al.* Arc/Arg3.1 Mediates Homeostatic Synaptic Scaling of AMPA Receptors. *Neuron* **52**, 475–484 (2006).

91. Turrigiano, G. Homeostatic Synaptic Plasticity: Local and Global Mechanisms for Stabilizing Neuronal Function. *Cold Spring Harb Perspect Biol* **4**, (2012).

92. Okuno, H. *et al.* Inverse Synaptic Tagging of Inactive Synapses via Dynamic Interaction of Arc/Arg3.1 with CaMKIIβ. *Cell* **149**, 886–898 (2012).

93. Rotenberg, A., Mayford, M., Hawkins, R. D., Kandel, E. R. & Muller, R. U. Mice Expressing Activated CaMKII Lack Low Frequency LTP and Do Not Form Stable Place Cells in the CA1 Region of the Hippocampus. *Cell* **87**, 1351–1361 (1996).

94. Cacucci, F., Wills, T. J., Lever, C., Giese, K. P. & O'Keefe, J. Experience-Dependent Increase in CA1 Place Cell Spatial Information, But Not Spatial Reproducibility, Is Dependent on the Autophosphorylation of the α-Isoform of the Calcium/Calmodulin-Dependent Protein Kinase II. *J. Neurosci.* **27**, 7854–7859 (2007).

95. Kitanishi, T. *et al.* Novelty-Induced Phase-Locked Firing to Slow Gamma Oscillations in the Hippocampus: Requirement of Synaptic Plasticity. *Neuron* **86**, 1265–1276 (2015).

96. Vetencourt, J. F. M. *et al.* The antidepressant fluoxetine restores plasticity in the adult visual cortex. *Science* **320**, 385–388 (2008).

97. Hensch, T. K. & Bilimoria, P. M. Re-opening Windows: Manipulating Critical Periods for Brain Development. *Cerebrum* **2012**, (2012).

98. Karpova, N. N. *et al.* Fear Erasure in Mice Requires Synergy Between Antidepressant Drugs and Extinction Training. *Science* **334**, 1731–1734 (2011).

99. Branchi, I. The double edged sword of neural plasticity: Increasing serotonin levels leads to both greater vulnerability to depression and improved capacity to recover. *Psychoneuroendocrinology* **36**, 339–351 (2011).

100. Alboni, S. *et al.* Fluoxetine effects on molecular, cellular and behavioral endophenotypes of depression are driven by the living environment. *Mol Psychiatry* **22**, 552–561 (2017).

101.   Brown, K. A. & Gould, T. D. Targeting metaplasticity mechanisms to
promote sustained antidepressant actions. *Mol Psychiatry* 1–14 (2024) doi:10.1038/s41380-
023-02397-1.

102.   Abraham, W. C. & Bear, M. F. Metaplasticity: the plasticity of synaptic
plasticity. *Trends in Neurosciences* **19**, 126–130 (1996).

103.   Stephens, G. S. *et al.* Persistent ΔFosB expression limits recurrent seizure
activity and provides neuroprotection in the dentate gyrus of APP mice. *Progress in*
Neurobiology 102612 (2024) doi:10.1016/j.pneurobio.2024.102612.

104.   Deibler, R. W. & Kirschner, M. W. Quantitative Reconstitution of Mitotic
CDK1 Activation in Somatic Cell Extracts. *Molecular Cell* **37**, 753–767 (2010).

105.   Sparta, D. R. *et al.* Construction of implantable optical fibers for long-term
optogenetic manipulation of neural circuits. *Nat Protoc* **7**, 12–23 (2012).

106.   Cardin, J. A. *et al.* Targeted optogenetic stimulation and recording of
neurons in vivo using cell-type-specific expression of Channelrhodopsin-2. *Nat Protoc* **5**, 247–
254 (2010).

107.   Yoshimura, Y. *et al.* Involvement of T-type Ca2+ channels in the
potentiation of synaptic and visual responses during the critical period in rat visual cortex.
European Journal of Neuroscience **28**, 730–743 (2008).

108.   Hagihara, H., Toyama, K., Yamasaki, N. & Miyakawa, T. Dissection of
hippocampal dentate gyrus from adult mouse. *J Vis Exp* (2009) doi:10.3791/1543.

109.   Keller, D., Erö, C. & Markram, H. Cell Densities in the Mouse Brain: A
Systematic Review. *Frontiers in Neuroanatomy* **12**, (2018).

110.   Buenrostro, J. D., Giresi, P. G., Zaba, L. C., Chang, H. Y. & Greenleaf, W.
J. Transposition of native chromatin for multimodal regulatory analysis and personal
epigenomics. *Nat Methods* **10**, 1213–1218 (2013).

111.   Corces, M. R. *et al.* An improved ATAC-seq protocol reduces background
and enables interrogation of frozen tissues. *Nat Methods* **14**, 959–962 (2017).

112.   Robinson, J. T. *et al.* Integrative genomics viewer. *Nat Biotechnol* **29**, 24–
26 (2011).

113.   Senda, T., Iizuka-Kogo, A. & Shimomura, A. Visualization of the Nuclear
Lamina in Mouse Anterior Pituitary Cells and Immunocytochemical Detection of Lamin A/C
by Quick-freeze Freeze-substitution Electron Microscopy. *J Histochem Cytochem.* **53**, 497–
507 (2005).

114.   Coffinier, C. *et al.* Deficiencies in lamin B1 and lamin B2 cause
neurodevelopmental defects and distinct nuclear shape abnormalities in neurons. *MBoC* **22**,

4683–4693 (2011).

115.   Shoji, H. & Miyakawa, T. Age-related behavioral changes from young to
old age in male mice of a C57BL/6J strain maintained under a genetic stability program.
Neuropsychopharmacology Reports **39**, 100–118 (2019).

116.   Miyakawa, T. *et al.* Conditional calcineurin knockout mice exhibit
multiple abnormal behaviors related to schizophrenia. *PNAS* **100**, 8987–8992 (2003).

117.   Stamatakis, A. M. *et al.* Simultaneous Optogenetics and Cellular
Resolution Calcium Imaging During Active Behavior Using a Miniaturized Microscope.
Frontiers in Neuroscience **12**, (2018).

118.   Skaggs, W. E., McNaughton, B. L. & Gothard, K. M. An Information-
Theoretic Approach to Deciphering the Hippocampal Code. in *Advances in Neural Information*
Processing Systems 5 (eds. Hanson, S. J., Cowan, J. D. & Giles, C. L.) 1030–1037 (Morgan-
Kaufmann, 1993).

119.   Kropff, E., Carmichael, J. E., Moser, M.-B. & Moser, E. I. Speed cells in
the medial entorhinal cortex. *Nature* **523**, 419–424 (2015).

120.   A novel somatosensory spatial navigation system outside the hippocampal
formation | Cell Research. https://www.nature.com/articles/s41422-020-00448-8.

121.   Glaser, J. I. *et al.* Machine Learning for Neural Decoding. *eNeuro* **7**, (2020).

122.   Tanaka, M. *et al.* Fascin in lamellipodia contributes to cell elasticity by
controlling the orientation of filamentous actin. *Genes to Cells* **24**, 202–213 (2019).

1501

1502

Acknowledgments

We thank the members of the Miyakawa laboratory—Wakako Hasegawa, Chikako Ozeki, Tamaki Murakami, Yoko Kagami and Harumi Mitsuya—for technical assistance, and Hirotaka Shoji for critical input on the design of the Ca^{2+} imaging experiment. We are grateful to Akihiro Yamanaka (Nagoya University and the Chinese Institute for Brain Research) for technical advice on optogenetics. We thank A. Konno, H. Hirai, and the Viral Vector Core at the Gunma University Initiative for Advanced Research for producing AAV9 viral particles. We also acknowledge D. Weinberger and T. Hyde for providing the human DG RNA-seq data. We thank Dr. Charles Yokoyama for his valuable comments, discussions, and editing and for greatly improving our manuscript. This research was supported by the Program for Brain Mapping by Integrated Neurotechnologies for Disease Studies (Brain/MINDS) from Japan Agency for Medical Research and Development, AMED under Grant Number JP21dm0207111, Japan Society for the Promotion of Science (JSPS) Grant-in-Aid for Scientific Research on Innovative Areas (grant nos. JP16H06462), JSPS KAKENHI (grant nos. JP20H00522 and JP22K15204), and the Japan Agency for Medical Research and Development Strategic Research Program for Brain Science Grant JP18dm0107101.

Author Contribution Statement

T. Murano, K. Takao, and T. Miyakawa designed the experiments and analyses. T. Murano performed RNA-seq, ATAC-seq, histological analysis, in vivo genome editing experiments, and data analyses. T. Murano and K. Takao performed the behavioral experiments. T. Murano and Y. Takamiya performed the Ca^{2+} imaging experiments. M. Namihira and K. Katoh performed the STED microscopy observations. K. Tajinda and M. Matsumoto provided RNA-seq data of post-mortem DG of human patients. T. Murano prepared all the figures and wrote the manuscript. T. Miyakawa supervised all aspects of the study. All authors reviewed the manuscript.

Declaration of Interests

The authors declare no competing interests.