Plant age and genotype explain variation in the microbiome of natural *Lotus corniculatus* populations

Frank Reis[1], Katrina Lutap[2], Jun Hee Jung[1], Anna Roeder[3,4], Christiane Roscher[3,4], Walter Durka[4,5], Eric Kemen[2], Oliver Bossdorf[1,*]

[1]Plant Evolutionary Ecology, Institute of Evolution and Ecology, University of Tübingen, Germany

[2]Microbial Interactions in Plant Ecosystems, Center for Plant Molecular Biology, University of Tübingen, Germany

[3]Department Physiological Diversity, Helmholtz Centre for Environmental Research-UFZ, Leipzig, Germany

[4]German Centre for Integrative Biodiversity Research (iDiv), Halle-Jena-Leipzig, Leipzig, Germany

[5]Department of Community Ecology, Helmholtz Centre for Environmental Research-UFZ, Halle, Germany

* Corresponding author: oliver.bossdorf@uni-tuebingen.de

Abstract

In natural populations, plants are associated with a huge diversity of bacteria, fungi and other microbes. There is usually substantial microbiome variation between different plant individuals and populations, and the drivers of this variation are still poorly understood, particularly in wild plants. Here, we were interested in the potential of plant genotype and plant age to explain intraspecific variation in the plant microbiome of *Lotus corniculatus*. In seven natural populations, we genotyped a total of 168 individuals over four years, determined their ages through growth ring analysis, and then sequenced their root, shoot, flower and seed microbiomes. We found that plant genotypes differed both in the diversity and composition of microbes, and that some microbial taxa were associated with particular plant genotypes. The genotype effects tended to be strongest and most consistent for plant-associated bacteria, with the largest plant genotype differences in the microbiome diversity of flowers and seeds. We found less evidence for an effect of plant age on microbiome diversity: the age of plants explained variation in fungi diversity, and it was associated with the abundance of several microbial taxa. Our study indicates that the genotype of a plant and - to a lesser degree its age - influences the diversity and composition of plant-associated microbiota, even in complex natural environments.

Importance

The plant microbiome plays a key role in important plant functions such as pathogen resistance, nutrient uptake, and stress tolerance. To fully understand these processes, it is essential to identify the factors that drive microbiome variation. Most research to date has focused on model or crop species under controlled conditions, leaving open questions about the drivers of microbiome

diversity and composition in natural populations. In this multi-year field study, we examined how plant genotype and age shape the microbiomes of *Lotus corniculatus*. We show that the plant genotype strongly influences microbial diversity and composition, while plant age has subtler but still important effects, particularly on fungal communities. To our knowledge, this is the first study to link plant age with microbiome variation in natural populations. These findings demonstrate that plant traits can shape the microbiome even in complex natural environments.

Introduction

Plants are colonized by a wide range of bacterial, fungal and eukaryotic microorganisms collectively called the plant microbiome. The diversity and composition of this microbiome is strongly shaped by the plant species (1, 2) and the surrounding microbial community, especially the soil microbiome (3, 4), but there are many other biotic and abiotic factors that are known to influence plant microbiome variation among and within species, including climate (5, 6), soil characteristics (7), land use (8, 9), herbivory (10, 11) or plant diseases (12–16). Besides this plethora of external factors, the plant microbiome can also be influenced by the plant itself. Variations in gene expression and plant traits can result in changing physical and chemical properties of the plants, and thus habitat conditions for the microbes, which in turn can alter microbiome diversity and composition. Previous studies have shown that the plant microbiome indeed varies across different plant tissue types such as roots, shoots or flowers (17–19) and also between the inside (endophytic) and outside (epiphytic) microbiome (19, 20). Two other potentially important factors creating microbiome variation among plants of the same species are their genetic differences, and their differences in age or developmental stage.

It is well established that the genetic variation in plants can change the pathogen resistance (21–23), and there is also good evidence that it can influence the overall composition of the plant microbiome, e.g. in soybean (24), potato (10), black cottonwood (25). While these studies were all conducted under controlled lab conditions, there is also evidence from more natural common garden experiments, e.g. for plant genotype effects on the microbiome of *Medicago truncatula* (26) and *Boechera stricta* (27). So far the research on these questions has been largely restricted to crops and model species, and to - usually short-term - experiments, whereas studies on wild plants and natural populations remain rare. Still, extrapolating lab results about plant-microbe interactions to natural environments is often challenging (28) because in natural populations plant microbiomes are influenced by a much larger complexity of biotic and abiotic factors. A true understanding of plant genotype effects, and their relative importance, therefore requires to also study plant genotype-microbiome relationships in natural populations.

Another potentially important but so far very little studied intrinsic factor creating intraspecific variation in plant microbiomes is the age or developmental stage of a plant. As a plant grows and develops, it undergoes physiological and biochemical changes that influence important functional traits, such as defense against herbivory (29, 30) or immune responses to pathogens (31). These changes across different life stages will inevitably also influence plant-microbiome interactions (32).

For example, the root-associated microbiome of rice varies between the vegetative phase and later life stages (33), and very similar results were found for soybean (34) and sugarcane (35). For annual plants like rice or soybean such a comparison between vegetative and reproductive phase covers their entire life cycle. However, many plants, in particular wild ones, are perennial, which creates additional possibilities for age-related microbiome variation e.g. through slower successional or competitive replacement processes . So far, there are hardly any studies of perennial plants examining relationships between plant age and plant microbiome. An exception is the study by Wagner et. al (2016) who found substantial changes in root (and to a lesser extent leaf) bacterial communities when comparing two and four year old *Boechera stricta* plants in a common garden, suggesting that significant microbiome changes can also occur at later life stages of perennials. Considering that most plants are perennial, we clearly need more studies relating plant age to plant microbiome in longer-lived plants.

Understanding the influences of plant age and genotype on microbiome composition may be important not only from a fundamental scientific perspective but also for agriculture and plant conservation, for instance if it helps to better manage beneficial crop microbiomes that increase yield production, stress tolerance or pathogen resistance. The last point is particularly important given that pathogens are a huge threat to global food security (36). More generally, understanding the drivers and consequences of plant microbiome diversity may support the management of plant populations in rapidly changing environments, and the effects of plant age and genotype on plant microbiomes - particularly under realistic conditions - are important elements of this.

Here, we combined amplicon-based metagenomics with growth ring analyses for plant age determination and genotype, and the diversity and composition of plant microbiomes within and among natural plant populations. Our study organism was *Lotus corniculatus*, a widely distributed perennial legume that grows naturally in a broad range of environments in temperate Eurasia and is an important food source for many pollinators, such as bumblebees, honeybees, and many wild bees as well as butterflies, flies and beetles (37–39). As a nitrogen fixer it is part of a complex plant-microbe symbiosis influencing the nutrient dynamics of grassland ecosystems (40), and it therefore plays an important role in agriculture. As *L. corniculatus* is known to have a broad life span range at least up to 14 years (41, 42), and it is known to harbor significant intraspecific genetic diversity (43, 44), the species is a suitable system for asking questions about effects of plant age and genotype. Specifically, we wanted to answer the following questions:

1. Do plant age and genotype affect the diversity of the plant microbiome, and if yes how do these effects vary between different plant organs and microbial groups?
2. Are plant age and genotype related to the composition (beta diversity) of the plant microbiome, and how are these relationships influenced by the plant organs and microbial groups?
3. Which specific microbial taxa are most differentiated between plants of different ages and genotypes?

To answer these questions, we collected *L. corniculatus* plants from seven semi-natural grasslands in the Swabian Jura region in Germany over four consecutive years, and we then determined the ages and genotype of all plants, and analyzed their bacterial, fungal and eukaryotic microbes separately for roots, shoots, flowers and seeds.

Methods

Plant material

We collected *Lotus corniculatus* plants from seven extensively managed meadows in the Swabian Alb region of southern Germany (Fig. 1a) for four consecutive years from 2018 to 2021. Sampling was done in the late summer (August/September), when the plants were still flowering but already carried some ripe fruits, so that both flowers and seeds could be sampled. In each of the seven populations, we carefully excavated six entire plants, resulting in a total of 4 years x 7 populations x 6 plants = 168 plants. In the lab, we separated the plants into roots, shoots, flowers and seeds. To surface sterilize all plant parts we washed them by shaking in Falcon tubes (50 ml for roots and shoots; 25 ml for flowers) and 1.5 ml Eppendorf tubes for seeds for 30 s with autoclaved ultrapure water (UPW), for 60 s with epiphyte wash solution (0.1 % Triton X-100 in 1x TE buffer), for 30 s with 80% EtOH and for 30 s with 2 % NaOCl. After that, we rinsed the samples three times (seeds: five times) for 10 s with UPW, and froze them.

Determination of plant ages

To determine the age of the sampled plants, we cut an approx. 3 cm long piece of the oldest part of the plant, i.e. the root collar as the transition zone between the primary root and the germination stem, from all plants and preserved them in 70% ethanol. In the lab, we placed these samples into a microtome (Reichert, Vienna, Austria) and sliced them into transverse sections. Very small samples with a diameter smaller than 3 mm were clamped using a piece of wood. To harden the soft tissue and keep the samples moist for cutting, the sample and the microtome blade were brushed with 99% ethanol during the cutting process. The microtome sections were then fixed on a microscope slide in a 1:1 mixture of water and glycerol and covered with a cover glass. We used a microscope to count the number of growth rings in the xylem, with ring boundaries defined by marginal parenchyma, semi-ring porosity or both (45) (Fig. 1b). Where necessary we used polarized light to improve the visibility of the growth rings. Since the growth rings were not equally distinct in all specimens, we classified the distinctness of the rings into 1 = very good, 2 = intermediate, 3 = poor and 4 = no annual rings visible (Tab. S1) and used only classes 1 to 3 for our analyses. For the statistical analysis we decided to group the plants into five age groups since the low number of replicates for the very young (2–3 years) and very old plants (6–13 years). Therefore we grouped the plants into age categories: younger than 4, 4, 5, 6, and older than 6 years (Fig. 1d).

Plant genotyping

To genotype the sampled plants, and assess their relatedness, we employed ddRAD (46), a reduced-representation sequencing approach. We extracted plant DNA from all shoot samples using the QIAGEN DNeasy Plant Mini Kit (QIAGEN, 2019). We then double-digested the DNA and followed the library preparation protocol of Peterson et al. (2012) (46) with small modifications as given in Durka et al. (2025) (44). Multiplexed samples were sequenced paired-end (PE 150) on an Illumina Novaseq 6000 platform. After sequencing, we demultiplexed the reads using process_radtags from the Stacks 2.0 pipeline (47) resulting in an average of 6.31 million reads retained per sample. We assembled reads, including three outgroup samples of *L. tenuis* and *L. pedunculatus* from Durka et al. (2025) and called SNPs using dDocent 2.7.8 (48, 49) with default parameters, except of using 0.90 as

ClusteringSimilarity, filtered according to O'Leary et al. 2018 introducing a mean depth threshold of 5, and a minor allele frequency threshold of 0.05 and keeping only 1 SNP per contig. We removed samples not used in this study from the resulting VCF file and re-filtered the remaining samples. The curated VCF file was converted into a fasta file (containing only the SNPs for each sample) with PGDSpider version 2.1.1.5 (50). To construct a Maximum Likelihood tree we used the program IQ-TREE release 2.0.3 (51). We used ModelFinder to select the best-fitting substitution model based on the Bayesian Information Criterion (BIC) (52) which identified the General Time Reversible (GTR) model with empirical base frequencies (F), a proportion of invariable sites (I), and gamma-distributed rate heterogeneity with four categories (G4) as the best-fit model (GTR+F+I+G4). To assess branch support, we also performed ultrafast bootstrap approximation (UFBoot) with 1,000 replicates (53). Finally, we used TreeCluster version 1.0.4 (54) to classify the SNP-based genotypes into a more restricted number of genetic groups (later used in our analyses), based on the total branch length of all leaves in the cluster (Sum Branch) with a threshold of 3.1. We visualized the phylogenetic tree using Interactive Tree of Life (iTOL) (55) and colored the samples by their genetic group assignments. From here on we will refer to these nine genetic groups as genotypes.

Microbiome sequencing

To prepare the plant samples for microbiome sequencing, we first homogenized all frozen samples of soil, roots, shoots, flowers, and seeds with a Precellys 24 Tissue Homogenizer (Bertin Technologies, Montigny-le-Bretonneux, France), and then extracted the DNA using the FastDNA Spin Kit for Soil (MP Biomedicals, Irvine, CA, USA). We then performed a two-step PCR targeting the bacterial 16S rRNA V5-V7 region, the fungal ITS2 region, and the eukaryotic 18S rRNA V9 region, using the primers 799F/1192R, fITS7/ITS4, and F1422/R1797, respectively (Tab. S2) (20). As controls we used blank samples (UPW and blank DNA extraction). To reduce the amplification of mitochondrial and chloroplast rRNA sequences from *L. corniculatus*, we incorporated blocking oligos designed with the R package *AmpStop* (Tab. S2) (56, 57). We pooled the amplified products in equal concentrations, purified them with magnetic bead clean-up, and randomly assigned each to one of eight sequencing batches. We sequenced all pools on a Illumina MiSeq platform with PhiX control, using the MiSeq Reagent Kit v3 (600-cycle), and then processed all microbial 16S rRNA, ITS2, and 18S rRNA amplicon sequences using Mothur (56, 58, 59). To remove the primer sequences from 16S rRNA and 18S rRNA data we used Cutadapt, and to remove it from ITS2 data we used ITSx (60, 61). For the taxonomic classification of the bacterial 16S rRNA reads, we used the Greengenes database (13_8_99 release), for fungal ITS2 reads the UNITE database (02.02.2019 release), and for eukaryotic 18S rRNA the PR2 database (version 4.12.0), all of which included the PhiX genome (62–64).

Data analysis

We carried out all data analyses in R Studio 2024.12.1 (65), and we used *dplyr* for structuring and manipulating data frames (66) *ggplot2* package (67) for creating figures. We used the *vegan* package (68) for calculating the alpha diversity (Shannon index) of bacterial, fungal or eukaryotic microbiomes of each organ in each plant, and then linear models to test for the effects of plant age or genotype. To account for variation in other factors, all models included the plant organ, and its respective interaction with plant genotype or plant age as well as sample site and year.

In addition to variation in microbial alpha diversity, we also analyzed the compositional turnover (beta diversity) between different samples, using the R packages *vegan*, *phyloseq*, and *microbiome* (68–70). We first used the OTU relative abundances for a Principal Coordinate Analysis (PCoA) of the Bray-Curtis dissimilarities between samples, and then a PERMANOVA to test how much of the variation in microbial community composition could be explained by plant age, plant genotype, or their interaction. We also visualized the relative abundance profiles of the ten most abundant classes of bacteria, fungi or eukaryotes for the different *L. corniculatus* genotypes or age groups, using the *microeco* package (69). To further understand the abundance patterns of the three different main groups of microbes, we also created graphs of the frequencies of different levels of commonness of OTUs, occurring in one to nine genotypes, or in one to five age groups.

To identify specific bacterial, fungal or eukaryotic taxa that were associated with particular plant genotypes or plant age groups, we used the R packages *microbiomeMarker* and *phyloseq* (70, 71). We used linear discriminant analysis effect size (LEfSe) (p-value < 0.05, LDA score ≥ 2) (72) to identify microbial genera that were significantly associated with particular *L. corniculatus* genotypes or age groups, either separately for each plant organ, or across organs.

Results

The ddRAD sequencing identified a total of 5739 SNPs across all plant individuals. The Maximum Likelihood tree with best score (LL score = -733814, not shown) and consensus tree (LL score = -733810; Fig. S1) agree in all major branches. Both trees are robust, with most bootstrap values above 50%. Based on the branch lengths of the ML tree, the TreeCluster algorithm grouped the plants into nine genotype groups, each represented by 14 to 26 samples (Fig. 1b, Fig. S1). While six of these groups were significantly overrepresented in some sites (Fig. 1b, χ2 test, $P < 0.001$), all groups occurred across multiple sites (3-6 sites, average 4.4 sites), usually with multiple replicates per site, and the use of the genotype groups therefore allowed us to test for their microbiome associations at least partly independently from site variation. The estimated ages of the same plant individuals ranged from two to 13 years, with most plants being four to six years old (Fig. 1d). We found no significant relationship between plant age and sampling year (χ2 test, $P = 0.076$), but there was a statistical association between plant age and sampling site (χ2 test, $P < 0.001$), with an overrepresentation of the younger than four year old plants in one of the sites. Thus, plant age was only to a small extent confounded with site, allowing us to test for its (largely) independent association with microbiome.

Figure 1. Sampling region, genotypic diversity, and determination of plant age. **a)** Photos of four of the sampling sites in southwest Germany. **b)** Frequencies of *Lotus* genotypes in the seven studied populations. The red borders indicate cases where genotypes are significantly overrepresented in specific populations. **c)** The root crown of a *Lotus* plant, and its cross-section, with the growth rings used to assess plant age. **d)** Age distribution of all sampled plants based on herb chronology. The colors indicate the five age groups we created for the statistical analysis.

Microbiome diversity differs between plant genotypes and ages

Across all sampling sites, years, plant individuals and plant organs (672 samples), the metagenome sequencing identified a total of 4,225 bacterial (16S rRNA), 2,027 fungal (ITS2 rRNA), and 1,773 eukaryotic (18S rRNA) OTUs, clustered at 97% sequence similarity and classified into 113 phyla and 1,542 genera. The most abundant bacterial classes we detected in the *Lotus* plants were Gammaproteobacteria, Alphaproteobacteria, Betaproteobacteria, Actinobacteria, and Bacilli (Fig. 3a). The most abundant fungal classes are, Dothideomycetes, Leotiomycetes, Eurotiomycetes, Sordariomycetes, and the most abundant eukaryotic classes Insecta, Dothideomycetes, Chromadorea_X, and Agaricomycetes (Fig. 3b,c).

Part of the observed microbial diversity differed systematically between plant organs, genotypes and age groups. Specifically, the variation in Shannon diversity (= within-sample) was highly significant between the different plant organs of all genotypes and age groups. In all cases, root diversity was substantially higher than that of other organs, followed by shoot and flower diversity. Notably, seed diversity was higher than that of flowers (Fig. 2a-f, Fig. S2). Both bacterial and fungal communities differed significantly among plant genotypes, with some genotypes harboring consistently higher bacterial or fungal diversity than others, across plant organs (Fig. 2a,b). We also found a significant interaction between genotype and plant organ for bacterial diversity: genotype differences were much more pronounced in *Lotus* flowers and seeds than in shoots and roots (Fig. 2a). A similar,

albeit not statistically significant, pattern was visible for fungal diversity, where also the genotype differences were greatest in flowers (Fig. 2b). In contrast to the flower and seed microbiome, the microbial diversity of roots and shoots was much less variable between plant genotypes, and there were also no genotype differences in the diversity of eukaryotes (Fig. 2c). When comparing the microbiome diversity of different plant age groups, we found a significant age effect for fungal diversity, with younger plants harboring a higher diversity of fungal microbes than older ones (Fig. 2e). However, there were no plant age effects on bacterial or eukaryote diversity (Fig. 2d,f).

Figure 2: The average alpha diversity of bacterial, fungal and eukaryotic microbial communities associated with different genotypes (panels a - c) or different age groups (panels d - f) of *Lotus corniculatus* across seven semi-natural grasslands. Illustrations by S. Nicolai Rühl.

Plant genotype and age explain variation in microbiome composition

The age and genotypes of plants were not only associated with the diversity but also the composition of their microbiome. A large number of taxa associated with *L. corniculatus* were shared among all genotypes and age groups, but we also found taxa that were less widely distributed, or even occurred in only one genotype or age group. Interestingly, the patterns of microbial commonness and rarity strongly differed between bacteria on the one side, and fungi and eukaryotes on the other side. In bacteria the majority of OTUs (2572 out of 4225) occurred in all nine genotypes, while OTUs that occurred in only one or few genotypes were rare. In fungi and eukaryotes, such less widely distributed taxa were much more frequent (Fig. 3g). The results were similar when we examined OTU distribution across age groups: almost 80% of all bacterial OTUs occurred in all age groups, and such that occurred in only one or few age groups were rare, whereas the patterns were much more even in fungi and eukaryotes (Fig. 3h).

Figure 3: Relative abundance profiles of microbial communities associated with different genotypes (G1 - G9; panels a-c) or ages (panels d-f) of *Lotus corniculatus*, separately for bacteria, fungi and eukaryotes. We restricted the plots to the ten most abundant classes in each microbial group . Panels (g) and (h) summarize the frequencies of different levels of commonness across genotypes or ages in the three microbial groups.

The most consistent genotype effects were in roots and shoots, where the PERMANOVA identified significant plant genotype effects in the composition of all three microbial groups, with around 10% of the variation in community composition explained by plant genotype. In addition, there were significant genotype effects for bacterial communities in flowers and seeds, and for fungal communities in flowers (Tab. 1, Fig. S3). Interestingly, these genotype effects in flowers and seeds were even stronger, with over 15% variance explained.

We found no significant main effects of plant age on the composition of bacteria, fungal and eukaryotic communities in any of the four plant organs (Tab. 1). However, there were significant plant age by genotype interactions for fungal communities in roots and shoots and eukaryotic communities in roots, indicating a complex interplay between plant genotype and age in explaining microbial community composition in these cases.

Table 1: Results of PERMANOVA analyses testing the effects of plant genotype, plant age, or their interaction, on the Bray-Curtis dissimilarities of microbial communities associated with the roots, shoots, flowers, and seeds of *Lotus corniculatus* across seven semi-natural grasslands in SW Germany.

Bacteria	Roots			Shoots			Flowers			Seeds		
	R^2	F	Pr(>F)	R^2	F	Pr(>F)	R^2	F	Pr(>F)	R^2	F	Pr(>F)
Plant genotype	0.102	1.53	**0.009**	0.103	1.57	**0.004**	0.174	2.67	**0.001**	0.160	2.59	**0.001**
Plant age	0.033	0.99	0.455	0.043	1.3	0.107	0.026	0.81	0.732	0.049	1.55	0.064
Genotpye x Age	0.234	1.04	0.309	0.231	1.04	0.277	0.180	0.82	0.956	0.204	0.98	0.549
Residual	0.632			0.623			0.620			0.588		

Fungi	Roots			Shoots			Flowers			Seeds		
	R^2	F	Pr(>F)	R^2	F	Pr(>F)	R^2	F	Pr(>F)	R^2	F	Pr(>F)
Plant genotype	0.092	1.41	**0.005**	0.108	1.61	**0.001**	0.173	3.02	**0.001**	0.070	1.04	0.385
Plant age	0.029	0.90	0.71	0.034	1.06	0.351	0.042	1.48	0.103	0.025	0.74	0.844
Genotpye x Age	0.264	1.21	**0.012**	0.248	1.15	**0.036**	0.239	1.23	0.079	0.258	1.12	0.385
Residual	0.615			0.615			0.545			0.646		

Eukaryotes	Roots			Shoots			Flowers			Seeds		
	R^2	F	Pr(>F)	R^2	F	Pr(>F)	R^2	F	Pr(>F)	R^2	F	Pr(>F)
Plant genotype	0.107	1.66	**0.001**	0.090	1.38	**0.008**	0.082	1.21	0.139	0.073	1.13	0.235
Plant age	0.035	1.07	0.264	0.037	1.13	0.257	0.029	0.85	0.641	0.046	1.41	0.076
Genotpye x Age	0.246	1.13	**0.021**	0.245	1.11	0.067	0.244	1.07	0.308	0.264	1.20	**0.043**
Residual	0.612			0.628			0.645			0.617		

Specific microbial taxa associated with plant genotypes and age groups

While a large number of taxa are shared among the plant genotypes and age groups, the LEfSe analyses revealed that some microbe taxa were significantly associated with particular individual plant genotypes or age groups (Fig. 4). Interestingly, these patterns of differential abundance were generally stronger in bacteria and fungi than in eukaryotic microbes, and they also tended to be more frequent with regard to plant genotypes (Fig. 4a) than for plant age groups (Fig. 4b), corroborating the PERMANOVA results with more frequent plant genotype than plant age effects (Tab. 1). Moreover, the LEfSe analyses also found that some plant genotypes and age groups showed much larger numbers of specific microbial taxa associations (e.g. genotypes 2, 3, 7 and 8, and age groups 2/3 and 5) than others. When analyzed separately for each plant organ, the LEfSE results differed significantly between organs, both for plant genotypes and plant age groups (Tab. S3), indicating that the observed microbial taxa associations are to some degree organ-specific.

Figure 4: Microbial taxa significantly associated with specific genotypes (a) or age groups (b) of *Lotus corniculatus*, as identified through LefSe analyses. The three colors represent separate LefSe analyses conducted for bacteria, fungi, and eukaryotes, which were analyzed independently. These groups are presented together to provide a clearer overview of microbial variation across genotypes and age groups; however, the LDA score of taxa from different groups cannot be directly compared with each other. The raw results can be found in the supplementary material (Fig. S4).

Discussion

Plants harbor an astonishing diversity of microbes, but the factors shaping these complex communities throughout a plant's life, particularly in natural populations, remain poorly understood. Here, we show that in natural *Lotus corniculatus* populations both the age and genotype of plants is associated with changes in the diversity and composition of their associated microbiome. This is the case for the entire microbiome but also for individual microbial taxa that are significantly associated with specific genotypes or age groups.

Effect of plant age and genotype on microbiome diversity

We generally found large differences in the average microbiome diversity of different plant organs, a result that has already been described in detail elsewhere (56), and that likely reflects the different properties of plant organs as habitats for microbes, and their different modes of connectedness with the environment (73). Because of this large variation, we included the plant organs as factors in our analysis, or ran analyses separately for different organs.

The alpha diversity of bacterial and fungal microorganisms differed significantly among the studied *Lotus corniculatus* genotypes. Interestingly, this effect was particularly strong in the flowers where we observed the strongest among-genotype variation in bacteria and fungi, whereas the microbiome diversity of roots and shoots was much more stable across genotypes. A possible explanation for this could be that flower traits are generally thought to be under stronger genetic control than vegetative plant traits (74), and that these traits can influence the microbiome, particularly if they affect the abundance and diversity of flower visitors (75), which are known to influence the flower microbiome (76, 77). For example, nectar secretion can vary among *L. corniculatus* genotypes (78), potentially influencing the rate of flower visitation. In the study area, the flower visitors of *L. corniculatus* include various bee species—bumblebees, honeybees, and wild bees—but also butterflies (primarily Lycaenidae) and sometimes flies and beetles (39). This broad range of pollinators may play a role in the observed flower microbiome variation between different plant genotypes. Moreover, the flower microbiome is closely linked to the seed microbiome, particularly for bacteria (79–82), which may explain why we also found considerable among-genotype variation for bacteria in seeds.

Among-genotype variation in microbiome was generally lower in plant roots and shoots than in flowers and seeds. It is well known that the root microbiome is strongly influenced, in part through affecting root traits, by different environmental factors such as soil type, structure and pH, and by the surrounding soil microbiome (1, 3, 4, 83, 84). Together, this may 'override' the influence of host genetics and reduce its relative impact on root microbiomes and, as a consequence, also the neighboring shoot microbiomes.

Compared to the variation among genotypes, there was less variation in microbiome diversity between plants of different ages. Only for the Shannon diversity of fungi there were significant differences between age groups. This is interesting, as many fungi play important roles as plant symbionts or pathogens (85–87), and their interactions with plants are strongly influenced by the plant immune system. The immune response of plants changes throughout their development (31, 88), and this could explain plant age-related variation in fungal microbiome diversity. Our results show that in principle plant age can also influence plant microbiome diversity, although to a lesser degree than plant genotype, and our evidence is limited to fungal microorganisms.

We should point out that comparisons of plant ages and genotypes in natural populations are not without challenges. In a field study, variation in plant age and genotype is inevitably confounded with variation in environment. We tried to account for this by aggregating both genotypes and ages into broader groups, which allowed 'replication' across populations. However, there was still statistical overrepresentation of some genotypes and age groups in some populations. Thus, despite

our efforts both factors have not been fully independent of the sampling population, and we therefore cannot rule out that some of the observed genotype or age effects are overestimates.

Effect of plant genotype and age on microbiome composition

We found that different *L. corniculatus* genotypes were also associated with variation in the composition of the microbiome, i.e. genetic variation in the host not only affected the alpha diversity but also the beta diversity of the microbes. Again, the extent of these effects strongly depended on the plant organ: in roots and shoots the composition of all three major microbial groups (bacteria, fungi and eukaryotes) was affected by plant genotype, whereas in flowers only the composition of bacteria and fungi, and in the seeds only that of bacteria, showed significant plant genotype associations. A plethora of genetically variable plant traits are known to affect the plant microbiome and could thus be underlying these observed composition changes. For example, root exudates are an important determinant of the rhizosphere microbiome (89, 90), and can vary across plant genotypes (91, 92). Intraspecific variation in leaf morphology and leaf chemistry has been shown to influence microbial communities in the phyllosphere (93, 94), and variation in plant defense genes is strongly affecting plant colonization by specific pathogens (95, 96).

In contrast to plant genotype, we found little evidence for effects of plant age on the composition of microbial communities. Only in fungi and eukaryotes, and mainly in roots and shoots, we found a significant plant age by genotype interaction, i.e. age effects were inconsistent and differed among genotypes, or were restricted to particular genotypes. This was counter to our expectation of age-related changes in species composition because of gene expression and functional trait changes during plant development (88). One possible reason for our findings could be that most species composition changes happen at early plant ages, and that the main factors shaping the microbiome such as root exudates (97) stabilize after this initial phase, so that the microbial communities reach an equilibrium fairly early. In our study, we deliberately collected only individuals, which had already formed inflorescences to study flowers and seeds. Thus, our youngest sampled plants were two years old, and we could not analyze the early-stage dynamics in more detail. Clearly, a better understanding of these questions, and of the observed plant age by genotype interactions, requires controlled experiments with longer-term observation of the colonization dynamics of different plant genotypes under standardized environmental conditions.

When looking at the overall frequency patterns of microbes, we found that although many taxa were shared across all plant genotypes and ages, there were still significant numbers of taxa that occurred only in a subset of plant genotypes or age groups, and there was an intriguing difference between bacteria on the one hand, and fungi and eukaryotes on the other hand. While in bacteria the level of generalism was particularly high, i.e. a large fraction of taxa occurred everywhere, frequency distributions of taxa were much more even in fungi and eukaryotes, showing a greater dispersal rates of bacteria than of non-bacteria microbes. Bacteria generally possess dispersal mechanisms unavailable to other microorganisms, allowing them to spread more efficiently. For instance, they can rapidly disperse through fungal hyphae in soil (98) and, in general, exhibit much higher dispersal rates than fungal microbes (99). In contrast, fungal dispersal is more strongly constrained by potential dispersal limitation (100), which may further contribute to the broader distribution of bacterial taxa.

Associations of microbial taxa with specific plant genotypes and age groups

The differential abundance analysis identified a number of significant associations of microbial taxa with specific plant genotypes. Interestingly, these associations were not randomly distributed, but some plant genotypes harbored many more specific microbial taxa than others. Plant genotypes can differ in their production of secondary metabolites, which influence the recruitment or inhibition of specific microbes (101). Additionally, modifications in root morphology can limit microbial attachment and interaction, reducing the presence of specialized microbial associations in the root (102). Changes in these chemical and morphological traits can require greater levels of specialization of associated microbes.

If we look at the genotype associations in more detail, we find that several mutualistic bacteria, such as *Bacillus* (103–105), primarily soil-associated bacteria like *Cloacibacterium*, *Prauserella* and *Rubrobacter*, and some growth-promoting fungi like *Agrocybe* (106, 107) were overrepresented in some plant genotypes. Similar host genotype effects on *Bacillus* have been observed in other studies, e.g. in maize cultivars resistant to corn stalk rot versus non-resistant cultivars (108), or in sweet pepper cultivars (104). While plant interactions with the soil-associated bacteria are not yet fully understood, previous studies have shown that the plant genotype can influence the abundance of specific soil microbes in the rhizosphere (109), and that plants actively recruit genotype-specific beneficial soil microbes (110). We also found several pathogens with differential abundance among plant genotypes, e.g. *Erwinia* bacteria responsible for soft rot (111) and the fungal pathogens *Sclerotinia*, *Alternaria*, *Pseudoidium*, *Golovinomyces* and *Gibberella*. Our results corroborate previous studies with different plant species that also found genetic variation in plant resistance to these pathogens (112–115). In general, plant genetic effects on pathogen resistance are a well known phenomenon, and a key topic in agriculture and crop management (116). They also play an important role in natural populations, and are a cornerstones of the geographic mosaic of coevolution observed in many wild species (117–119).

We also found that some microbial taxa were significantly overrepresented in particular plant age groups, e.g. bacteria from the genera *Burkholderia, Erwinia and Ralstonia* and fungi from the genus *Mycena* in the youngest plant group. *Burkholderia* bacteria are known for their growth-promoting effects (110) and have been shown to benefit maize plants during germination and seedling development (120). Although the studied *Lotus* plants were already beyond these initial stages of development, the observed higher abundances of *Burkholderia* in young plants suggest that these bacteria also play a role in early *Lotus* growth and development. *Mycena* is a saprotrophic fungus opportunistically invading plant roots (121), and *Erwinia* and *Ralstonia* are bacterial pathogens causing wilt in various plant species. The increased abundances of these taxa in young *Lotus* plants suggests these may have a less well developed immune system (122, 123) or other, e.g. root physiological, differences that make them more susceptible to invasion (124). There were also microbial taxa that were particularly abundant in older plants, e.g. the pathogenic fungus *Pyrenochaeta* which is known to cause root lesions (125), and which we found overrepresented in plants older than six years. Our results confirm a previous study with oilseed rape that also observed a higher abundance of *Pyrenochaeta* in later plant developmental stages (126). In general, our data show that while plant age does not have strong effects on the overall composition of microbes, there are some significant influences at the level of individual microbial taxa, even in a long-lived plant such as *Lotus corniculatus*.

Conclusion

Our findings show the host genotype is an important driver of natural microbiome diversity of *L. corniculatus*. This influence is evident not only in overall microbial diversity and composition but also in the abundance of specific microbial genera across different genotypes. We also found evidence of plant age influencing the fungal diversity. Additionally, the abundance of specific microbial genera varies between the different host age groups. With varied genetics or age, the physiology and requirements of the plants change, and due to the close interaction between the microbiome and the host, this can result in an altered optimal microbiome composition. Therefore, it is valuable to combine insights from controlled laboratory experiments with data from natural environments to gain a more comprehensive understanding of the impact of genotype and age on the microbiome. Overall, our results show that genotype, and to a lesser extent age, play important roles in shaping the microbiome diversity and composition of natural *L. corniculatus* populations. This knowledge can also help optimize plant health, improve crop yields, and enhance ecosystem resilience which is crucial for effective ecosystem management and plant breeding practices.

Acknowledgements

We thank the Kemen and Bossdorf labs for help with the field sampling, in particular Elke Klenk for her invaluable assistance in collecting and processing the samples, as well Sabine Silberhorn and Julia Rafalski for their help in the lab. We are also indebted to Ralf Lauterbach and Jörg Hailer from the Schwäbische Alb local management team of the Biodiversity Exploratories for instructions and support during the field work. We have been lucky to be granted access to the Schwäbische Alb plots of the Biodiversity Exploratories (DFG Priority Program 1374) through a cooperation agreement with this project. We thank the managers of the Schwäbische Alb exploratory, Kirsten Reichel-Jung and Julia Bass and all former managers for their work in maintaining these plots and the project infrastructure, Victoria Grießmeier for her support through the central office, Andreas Ostrowski for managing the central database, and Markus Fischer, Eduard Linsenmair, Dominik Hessenmöller, Daniel Prati, Ingo Schöning, François Buscot, Ernst-Detlef Schulze, Wolfgang W. Weisser and the late Elisabeth Kalko for their role in setting up the Biodiversity Exploratories project. We thank the administration of the UNESCO Biosphere Reserve Swabian Alb as well as all landowners for the excellent collaboration. Field work permits were issued by the responsible state environmental office of Baden-Württemberg.

Author contributions

OB, EK, FR and KL designed the study; FR and KL planned and conducted the field sampling, lab work and data analyses, JHJ helpt with the data analysis; WD helped with the library preparation and bioinformatics of the ddRAD sequencing; CR and AR helped with the age determination; FR, KL and OB wrote the manuscript with input from all authors. All authors read and approved the final manuscript.

Competing interests

We declare that there are no conflicts of interest related to this study.

Funding

This project has been funded by the Deutsche Forschungsgemeinschaft (DFG; projects BO 3241/11-1 to OB) as part of the priority programme 2125 Deconstruction and Reconstruction of the Plant Microbiota "DECRyPT".

Data availability

This work is based on data elaborated by a project associated with the Biodiversity Exploratories program (DFG Priority Program 1374). The datasets are publicly available in the Biodiversity Exploratories Information System (http://doi.org/10.17616/R32P9Q).
Sequencing data, metadata, OTU tables, and scripts are available on Biodiversity Exploratories Information System (BExIS) (https://www.bexis.uni-jena.de/) under Dataset ID 31836. Primers and blocking oligos used in this study are listed on Table S3.

References

1. Bonito G, Reynolds H, Robeson II MS, Nelson J, Hodkinson BP, Tuskan G, Schadt CW, Vilgalys R. 2014. Plant host and soil origin influence fungal and bacterial assemblages in the roots of woody plants. Mol Ecol 23:3356–3370.

2. Knief C, Ramette A, Frances L, Alonso-Blanco C, Vorholt JA. 2010. Site and plant species are important determinants of the Methylobacterium community composition in the plant phyllosphere. ISME J 4:719–728.

3. Bulgarelli D, Rott M, Schlaeppi K, Ver Loren van Themaat E, Ahmadinejad N, Assenza F, Rauf P, Huettel B, Reinhardt R, Schmelzer E, Peplies J, Gloeckner FO, Amann R, Eickhorst T, Schulze-Lefert P. 2012. Revealing structure and assembly cues for Arabidopsis root-inhabiting bacterial microbiota. Nature 488:91–95.

4. Schlaeppi K, Dombrowski N, Oter RG, Ver Loren van Themaat E, Schulze-Lefert P. 2014. Quantitative divergence of the bacterial root microbiota in Arabidopsis thaliana relatives. Proc Natl Acad Sci 111:585–592.

5. Chen P, Zhao M, Tang F, Hu Y, Peng X, Shen S. 2019. The effect of environment on the microbiome associated with the roots of a native woody plant under different climate types in China. Appl Microbiol Biotechnol 103:3899–3913.

6. Trivedi P, Batista BD, Bazany KE, Singh BK. 2022. Plant–microbiome interactions under a changing world: responses, consequences and perspectives. New Phytol 234:1951–1959.

7. Tkacz A, Cheema J, Chandra G, Grant A, Poole PS. 2015. Stability and succession of the rhizosphere microbiota depends upon plant type and soil composition. ISME J 9:2349–2359.

8. Estendorfer J, Stempfhuber B, Haury P, Vestergaard G, Rillig MC, Joshi J, Schröder P, Schloter M. 2017. The Influence of Land Use Intensity on the Plant-Associated Microbiome of Dactylis glomerata L. Front Plant Sci 8.

9. Gaube P, Junker RR, Keller A. 2021. Changes amid constancy: Flower and leaf microbiomes along land use gradients and between bioregions. Basic Appl Ecol 50:1–15.

10. Malacrinò A, Karley AJ, Schena L, Bennett AE. 2021. Soil Microbial Diversity Impacts Plant Microbiota More than Herbivory. Phytobiomes J 5:408–417.

11. Malacrinò A, Wang M, Caul S, Karley AJ, Bennett AE. 2021. Herbivory shapes the rhizosphere bacterial microbiota in potato plants. Environ Microbiol Rep 13:805–811.

12. Cui Z, Huntley RB, Zeng Q, Steven B. 2021. Temporal and spatial dynamics in the apple flower microbiome in the presence of the phytopathogen Erwinia amylovora. ISME J 15:318–329.

13. Diskin S, Feygenberg O, Maurer D, Droby S, Prusky D, Alkan N. 2017. Microbiome Alterations Are Correlated with Occurrence of Postharvest Stem-End Rot in Mango Fruit. Phytobiomes J 1:117–127.

14. Kusstatscher P, Cernava T, Harms K, Maier J, Eigner H, Berg G, Zachow C. 2019. Disease Incidence in Sugar Beet Fields Is Correlated with Microbial Diversity and Distinct Biological Markers. Phytobiomes J 3:22–30.

15. Solís-García IA, Ceballos-Luna O, Cortazar-Murillo EM, Desgarennes D, Garay-Serrano E, Patiño-Conde V, Guevara-Avendaño E, Méndez-Bravo A, Reverchon F. 2021. Phytophthora Root Rot Modifies the Composition of the Avocado Rhizosphere Microbiome and Increases the Abundance of Opportunistic Fungal Pathogens. Front Microbiol 11.

16. Wen T, Zhao M, Liu T, Huang Q, Yuan J, Shen Q. 2020. High abundance of Ralstonia solanacearum changed tomato rhizosphere microbiome and metabolome. BMC Plant Biol 20:166.

17. Bernard J, Wall CB, Costantini MS, Rollins RL, Atkins ML, Cabrera FP, Cetraro ND, Feliciano CKJ, Greene AL, Kitamura PK, Olmedo-Velarde A, Sirimalwatta VNS, Sung HW, Thompson LPM, Vu HT, Wilhite CJ, Amend AS. 2021. Plant part and a steep environmental gradient predict plant microbial composition in a tropical watershed. ISME J 15:999–1009.

18. Junker RR, Loewel C, Gross R, Dötterl S, Keller A, Blüthgen N. 2011. Composition of epiphytic bacterial communities differs on petals and leaves. Plant Biol 13:918–924.

19. Trivedi P, Leach JE, Tringe SG, Sa T, Singh BK. 2020. Plant–microbiome interactions: from community assembly to plant health. Nat Rev Microbiol 18:607–621.

20. Agler MT, Ruhe J, Kroll S, Morhenn C, Kim S-T, Weigel D, Kemen EM. 2016. Microbial Hub Taxa Link Host and Abiotic Factors to Plant Microbiome Variation. PLOS Biol 14:e1002352.

21. Deng Y, Ning Y, Yang D-L, Zhai K, Wang G-L, He Z. 2020. Molecular Basis of Disease Resistance and Perspectives on Breeding Strategies for Resistance Improvement in Crops. Mol Plant 13:1402–1419.

22. Karasov TL, Shirsekar G, Schwab R, Weigel D. 2020. What natural variation can teach us about resistance durability. Curr Opin Plant Biol 56:89–98.

23. Thrall PH, Burdon JJ. 2000. Effect of resistance variation in a natural plant host–pathogen metapopulation on disease dynamics. Plant Pathol 49:767–773.

24. Liu F, Hewezi T, Lebeis SL, Pantalone V, Grewal PS, Staton ME. 2019. Soil indigenous microbiome and plant genotypes cooperatively modify soybean rhizosphere microbiome assembly. BMC Microbiol 19:201.

25. Leopold DR, Busby PE. 2020. Host Genotype and Colonist Arrival Order Jointly Govern Plant Microbiome Composition and Function. Curr Biol 30:3260-3266.e5.

26. Brown SP, Grillo MA, Podowski JC, Heath KD. 2020. Soil origin and plant genotype structure distinct microbiome compartments in the model legume Medicago truncatula. Microbiome 8:139.

27. Wagner MR, Lundberg DS, del Rio TG, Tringe SG, Dangl JL, Mitchell-Olds T. 2016. Host genotype and age shape the leaf and root microbiomes of a wild perennial plant. 1. Nat Commun 7:12151.

28. Yu Z, Krause SMB, Beck DAC, Chistoserdova L. 2016. A Synthetic Ecology Perspective: How Well Does Behavior of Model Organisms in the Laboratory Predict Microbial Activities in Natural Habitats? Front Microbiol 7.

29. Boege K, Dirzo R, Siemens D, Brown P. 2007. Ontogenetic switches from plant resistance to tolerance: minimizing costs with age? Ecol Lett 10:177–187.

30. Yang LH, Cenzer ML, Morgan LJ, Hall GW. 2020. Species-specific, age-varying plant traits affect herbivore growth and survival. Ecology 101:e03029.

31. Develey-Rivière M-P, Galiana E. 2007. Resistance to pathogens and host developmental stage: a multifaceted relationship within the plant kingdom. New Phytol 175:405–416.

32. Bressan M, Roncato M-A, Bellvert F, Comte G, Haichar F el Z, Achouak W, Berge O. 2009. Exogenous glucosinolate produced by Arabidopsis thaliana has an impact on microbes in the rhizosphere and plant roots. ISME J 3:1243–1257.

33. Edwards JA, Santos-Medellín CM, Liechty ZS, Nguyen B, Lurie E, Eason S, Phillips G, Sundaresan V. 2018. Compositional shifts in root-associated bacterial and archaeal microbiota track the plant life cycle in field-grown rice. PLOS Biol 16:e2003862.

34. Sugiyama A, Ueda Y, Zushi T, Takase H, Yazaki K. 2014. Changes in the Bacterial Community of Soybean Rhizospheres during Growth in the Field. PLOS ONE 9:e100709.

35. Hamonts K, Trivedi P, Garg A, Janitz C, Grinyer J, Holford P, Botha FC, Anderson IC, Singh BK. 2018. Field study reveals core plant microbiota and relative importance of their drivers. Environ Microbiol 20:124–140.

36. Strange RN, Scott PR. 2005. Plant Disease: A Threat to Global Food Security. Annu Rev Phytopathol 43:83–116.

37. Hennig EI, Ghazoul J. 2011. Plant–pollinator interactions within the urban environment. Perspect

Plant Ecol Evol Syst 13:137–150.

38. van der Kooi CJ, Pen I, Staal M, Stavenga DG, Elzenga JTM. 2016. Competition for pollinators and intra-communal spectral dissimilarity of flowers. Plant Biol 18:56–62.

39. Weiner CN, Werner M, Linsenmair KE, Blüthgen N. 2014. Land-use impacts on plant–pollinator networks: interaction strength and specialization predict pollinator declines. Ecology 95:466–474.

40. van der Heijden MGA, de Bruin S, Luckerhoff L, van Logtestijn RSP, Schlaeppi K. 2016. A widespread plant-fungal-bacterial symbiosis promotes plant biodiversity, plant nutrition and seedling recruitment. ISME J 10:389–399.

41. Roeder A, Schweingruber FH, Fischer M, Roscher C. 2017. Growth ring analysis of multiple dicotyledonous herb species—A novel community-wide approach. Basic Appl Ecol 21:23–33.

42. Poschlod P, Hoffmann J, Bernhardt-Römermann M. Effect of grassland management on the age and reproduction structure of Helianthemum nummularium and Lotus corniculatus populations. Preslia 83:421–435.

43. Abraham EM, Ganopoulos I, Giagourta P, Osathanunkul M, Bosmali I, Tsaftaris A, Papaioannou A, Madesis P. 2015. Genetic diversity of *Lotus corniculatus* in relation to habitat type, species composition and species diversity. Biochem Syst Ecol 63:59–67.

44. Durka W, Michalski SG, Höfner J, Bucharova A, Kolář F, Müller CM, Oberprieler C, Šemberová K, Bauer M, Bernt M, Bleeker W, Brändel S, Bucher SF, Eibes PM, Ewald M, Goldberg R, Grant K, Haider S, Harpke A, Haun F, Kaufmann R, Korell L, Kunzmann D, Lauterbach D, Leib S, Lenzewski N, Loritz H, Madaj A-M, Mainz AK, Meinecke P, Mertens H, Meyer MH, Musche M, Ristow M, Rosche C, Roscher C, Rutte D, Schacherer A, Schmidt W, Schmoldt J, Schneider S, Schwarz J-H, Skowronek S, Socher SA, Stanik N, Twerski A, Weiß K, Weiß M, Wille A, Zehm A, Zidorn C. 2025. Assessment of genetic diversity among seed transfer zones for multiple grassland plant species across Germany. Basic Appl Ecol 84:50–60.

45. Schweingruber FH, Poschlod P. 2005. Growth Rings in Herbs and Shrubs: life span, age determination and stem anatomy. For Snow Landsc Res 79:195–415.

46. Peterson BK, Weber JN, Kay EH, Fisher HS, Hoekstra HE. 2012. Double Digest RADseq: An Inexpensive Method for De Novo SNP Discovery and Genotyping in Model and Non-Model Species. PLOS ONE 7:e37135.

47. Rochette NC, Rivera-Colón AG, Catchen JM. 2019. Stacks 2: Analytical methods for paired-end sequencing improve RADseq-based population genomics. Mol Ecol 28:4737–4754.

48. O'Leary SJ, Puritz JB, Willis SC, Hollenbeck CM, Portnoy DS. 2018. These aren't the loci you'e looking for: Principles of effective SNP filtering for molecular ecologists. Mol Ecol 27:3193–3206.

49. Puritz JB, Hollenbeck CM, Gold JR. 2014. dDocent: a RADseq, variant-calling pipeline designed for population genomics of non-model organisms. PeerJ 2:e431.

50. Lischer HEL, Excoffier L. 2012. PGDSpider: an automated data conversion tool for connecting population genetics and genomics programs. Bioinformatics 28:298–299.

51. Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, Lanfear R. 2020. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era. Mol Biol Evol 37:1530–1534.

52. Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. 2017. ModelFinder: fast model selection for accurate phylogenetic estimates. Nat Methods 14:587–589.

53. Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS. 2018. UFBoot2: Improving the Ultrafast Bootstrap Approximation. Mol Biol Evol 35:518–522.

54. Balaban M, Moshiri N, Mai U, Jia X, Mirarab S. 2019. TreeCluster: Clustering biological sequences using phylogenetic trees. PLOS ONE 14:e0221068.

55. Letunic I, Bork P. 2024. Interactive Tree of Life (iTOL) v6: recent updates to the phylogenetic tree display and annotation tool. Nucleic Acids Res gkae268.

56. Lutap K, Almario J, Mahmoudi M, Reis F, Bossdorf O, Kemen E. 2025. Organ-specific microbiomes in natural Lotus corniculatus populations: Metacommunity dynamics in the plant endosphere. bioRxiv https://doi.org/10.1101/2025.04.15.649008.

57. Mayer T, Mari A, Almario J, Murillo-Roos M, Syed M. Abdullah H, Dombrowski N, Hacquard S, Kemen EM, Agler MT. 2021. Obtaining deeper insights into microbiome diversity using a simple method to block host and nontargets in amplicon sequencing. Mol Ecol Resour 21:1952–1965.

58. Almario J, Mahmoudi M, Kroll S, Agler M, Placzek A, Mari A, Kemen E. 2022. The Leaf Microbiome of Arabidopsis Displays Reproducible Dynamics and Patterns throughout the Growing Season. mBio 13:e02825-21.

59. Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, Lesniewski RA, Oakley BB, Parks DH, Robinson CJ, Sahl JW, Stres B, Thallinger GG, Van Horn DJ, Weber CF. 2009. Introducing mothur: Open-Source, Platform-Independent, Community-Supported Software for Describing and Comparing Microbial Communities. Appl Environ Microbiol 75:7537–7541.

60. Bengtsson-Palme J, Ryberg M, Hartmann M, Branco S, Wang Z, Godhe A, De Wit P, Sánchez-García M, Ebersberger I, de Sousa F, Amend A, Jumpponen A, Unterseher M, Kristiansson E, Abarenkov K, Bertrand YJK, Sanli K, Eriksson KM, Vik U, Veldre V, Nilsson RH. 2013. Improved software detection and extraction of ITS1 and ITS2 from ribosomal ITS sequences of fungi and other eukaryotes for analysis of environmental sequencing data. Methods Ecol Evol 4:914–919.

61. Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads. 1. EMBnet.journal 17:10–12.

62. DeSantis TZ, Hugenholtz P, Larsen N, Rojas M, Brodie EL, Keller K, Huber T, Dalevi D, Hu P, Andersen GL. 2006. Greengenes, a Chimera-Checked 16S rRNA Gene Database and Workbench Compatible with ARB. Appl Environ Microbiol 72:5069–5072.

63. Guillou L, Bachar D, Audic S, Bass D, Berney C, Bittner L, Boutte C, Burgaud G, de Vargas C, Decelle J, del Campo J, Dolan JR, Dunthorn M, Edvardsen B, Holzmann M, Kooistra WHCF, Lara E, Le Bescot N, Logares R, Mahé F, Massana R, Montresor M, Morard R, Not F, Pawlowski J, Probert I, Sauvadet A-L, Siano R, Stoeck T, Vaulot D, Zimmermann P, Christen R. 2013. The Protist Ribosomal Reference database (PR2): a catalog of unicellular eukaryote Small Sub-Unit rRNA sequences with curated taxonomy. Nucleic Acids Res 41:D597–D604.

64. Nilsson RH, Larsson K-H, Taylor AFS, Bengtsson-Palme J, Jeppesen TS, Schigel D, Kennedy P, Picard K, Glöckner FO, Tedersoo L, Saar I, Kõljalg U, Abarenkov K. 2019. The UNITE database for molecular identification of fungi: handling dark taxa and parallel taxonomic classifications. Nucleic Acids Res 47:D259–D264.

65. RStudio Team. 2024. RStudio: Integrated Development for R (R version 4.4.2). PBC, Boston, MA URL.

66. Wickham H, François R, Henry L, Müller K, Vaughan D. 2023. dplyr: A Grammar of Data Manipulation (R package version 1.1.4).

67. Wickham H. 2016. ggplot2. Springer International Publishing, Cham. http://link.springer.com/10.1007/978-3-319-24277-4. Retrieved 16 January 2025.

68. Oksanen J, Simpson GL, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'Hara RB, Solymos P, Stevens MHH, Szoecs E, Wagner H, Barbour M, Bedward M, Bolker B, Borcard D, Borman T, Carvalho G, Chirico M, Durand S, Evangelista HBA, FitzJohn R, Friendly M, Furneaux B, Geoffrey Hannigan G, Mark O. H, Lahti L, McGlinn D, Ouellette M-H, Ribeiro Cunha E, Smith T, Stier A, Ter Braak CJF, Weedon J. 2025. vegan: Community Ecology Package (R package version 2.7-0).

69. Liu C, Cui Y, Li X, Yao M. 2021. microeco: an R package for data mining in microbial community ecology. FEMS Microbiol Ecol 97:fiaa255.

70. McMurdie PJ, Holmes S. 2013. phyloseq: An R Package for Reproducible Interactive Analysis and Graphics of Microbiome Census Data. PLOS ONE 8:e61217.

71. Cao Y, Dong Q, Wang D, Zhang P, Liu Y, Niu C. 2022. microbiomeMarker: an R/Bioconductor package for microbiome marker identification and visualization. Bioinformatics 38:4027–4029.

72. Segata N, Izard J, Waldron L, Gevers D, Miropolsky L, Garrett WS, Huttenhower C. 2011. Metagenomic biomarker discovery and explanation. Genome Biol 12:R60.

73. Huang L, Fu Y, Liu Y, Chen Y, Wang T, Wang M, Lin X, Feng Y. 2024. Global insights into endophytic bacterial communities of terrestrial plants: Exploring the potential applications of endophytic microbiota in sustainable agriculture. Sci Total Environ 927:172231.

74. Campbell DR, Raguso RA, Midzik M, Bischoff M, Broadhead GT. 2022. Genetic and spatial variation in vegetative and floral traits across a hybrid zone. Am J Bot 109:1780–1793.

75. Genung MA, Lessard J-P, Brown CB, Bunn WA, Cregger MA, Reynolds WN, Felker-Quinn E, Stevenson ML, Hartley AS, Crutsinger GM, Schweitzer JA, Bailey JK. 2010. Non-Additive Effects of

Genotypic Diversity Increase Floral Abundance and Abundance of Floral Visitors. PLOS ONE 5:e8711.

76. Keller A, McFrederick QS, Dharampal P, Steffan S, Danforth BN, Leonhardt SD. 2021. (More than) Hitchhikers through the network: the shared microbiome of bees and flowers. Curr Opin Insect Sci 44:8–15.

77. Zemenick AT, Vannette RL, Rosenheim JA. 2021. Linked networks reveal dual roles of insect dispersal and species sorting for bacterial communities in flowers. Oikos 130:697–707.

78. Murrell DC, Shuel RW, Tomes DT. 1982. Nectar production and floral characteristics in birdsfoot trefoil (Lotus corniculatus L.). Can J Plant Sci 62:361–371.

79. Compant S, Mitter B, Colli-Mull JG, Gangl H, Sessitsch A. 2011. Endophytes of Grapevine Flowers, Berries, and Seeds: Identification of Cultivable Bacteria, Comparison with Other Plant Parts, and Visualization of Niches of Colonization. Microb Ecol 62:188–197.

80. Dutta B, Ha Y, Lessl JT, Avci U, Sparks AC, Johnson KL, Walcott RR. 2015. Pathways of bacterial invasion and watermelon seed infection by Acidovorax citrulli. Plant Pathol 64:537–544.

81. Lessl JT, Fessehaie A, Walcott RR. 2007. Colonization of Female Watermelon Blossoms by Acidovorax avenae ssp. citrulli and the Relationship between Blossom Inoculum Dosage and Seed Infestation. J Phytopathol 155:114–121.

82. Mitter B, Pfaffenbichler N, Flavell R, Compant S, Antonielli L, Petric A, Berninger T, Naveed M, Sheibani-Tezerji R, von Maltzahn G, Sessitsch A. 2017. A New Approach to Modify Plant Microbiomes and Traits by Introducing Beneficial Bacteria at Flowering into Progeny Seeds. Front Microbiol 8.

83. Ristova D, Busch W. 2014. Natural Variation of Root Traits: From Development to Nutrient Uptake1. Plant Physiol 166:518–527.

84. Schreiter S, Ding G-C, Heuer H, Neumann G, Sandmann M, Grosch R, Kropf S, Smalla K. 2014. Effect of the soil type on the microbiome in the rhizosphere of field-grown lettuce. Front Microbiol 5.

85. Fisher MC, Gurr SJ, Cuomo CA, Blehert DS, Jin H, Stukenbrock EH, Stajich JE, Kahmann R, Boone C, Denning DW, Gow NAR, Klein BS, Kronstad JW, Sheppard DC, Taylor JW, Wright GD, Heitman J, Casadevall A, Cowen LE. 2020. Threats Posed by the Fungal Kingdom to Humans, Wildlife, and Agriculture. mBio 11:e00449-20.

86. Fisher MC, Henk DA, Briggs CJ, Brownstein JS, Madoff LC, McCraw SL, Gurr SJ. 2012. Emerging fungal threats to animal, plant and ecosystem health. Nature 484:186–194.

87. Martin FM, van der Heijden MGA. 2024. The mycorrhizal symbiosis: research frontiers in genomics, ecology, and agricultural application. New Phytol 242:1486–1506.

88. Hu L, Yang L. 2019. Time to Fight: Molecular Mechanisms of Age-Related Resistance. Phytopathology® 109:1500–1508.

89. Hu L, Robert CAM, Cadot S, Zhang X, Ye M, Li B, Manzo D, Chervet N, Steinger T, van der Heijden MGA, Schlaeppi K, Erb M. 2018. Root exudate metabolites drive plant-soil feedbacks on growth and defense by shaping the rhizosphere microbiota. Nat Commun 9:2738.

90. Zhalnina K, Louie KB, Hao Z, Mansoori N, da Rocha UN, Shi S, Cho H, Karaoz U, Loqué D, Bowen BP, Firestone MK, Northen TR, Brodie EL. 2018. Dynamic root exudate chemistry and microbial substrate preferences drive patterns in rhizosphere microbial community assembly. Nat Microbiol 3:470–480.

91. Micallef SA, Shiaris MP, Colón-Carmona A. 2009. Influence of Arabidopsis thaliana accessions on rhizobacterial communities and natural variation in root exudates. J Exp Bot 60:1729–1742.

92. Pacheco-Moreno A, Bollmann-Giolai A, Chandra G, Brett P, Davies J, Thornton O, Poole P, Ramachandran V, Brown JKM, Nicholson P, Ridout C, DeVos S, Malone JG. 2024. The genotype of barley cultivars influences multiple aspects of their associated microbiota via differential root exudate secretion. PLOS Biol 22:e3002232.

93. Hunter PJ, Hand P, Pink D, Whipps JM, Bending GD. 2010. Both Leaf Properties and Microbe-Microbe Interactions Influence Within-Species Variation in Bacterial Population Diversity and Structure in the Lettuce (Lactuca Species) Phyllosphere. Appl Environ Microbiol 76:8117–8125.

94. Kniskern JM, Traw MB, Bergelson J. 2007. Salicylic Acid and Jasmonic Acid Signaling Defense Pathways Reduce Natural Bacterial Diversity on Arabidopsis thaliana. Mol Plant-Microbe Interactions® 20:1512–1522.

95. Glander S, He F, Schmitz G, Witten A, Telschow A, de Meaux J. 2018. Assortment of Flowering Time and Immunity Alleles in Natural Arabidopsis thaliana Populations Suggests Immunity and Vegetative Lifespan Strategies Coevolve. Genome Biol Evol 10:2278–2291.

96. Ronald PC, Salmeron JM, Carland FM, Staskawicz BJ. 1992. The cloned avirulence gene avrPto induces disease resistance in tomato cultivars containing the Pto resistance gene. J Bacteriol 174:1604–1611.

97. Chaparro JM, Badri DV, Vivanco JM. 2014. Rhizosphere microbiome assemblage is affected by plant development. ISME J 8:790–803.

98. Vieira CK, Marascalchi MN, Rozmoš M, Benada O, Belova V, Jansa J. 2025. Arbuscular mycorrhizal fungal highways – What, how and why? Soil Biol Biochem 202:109702.

99. Larsen S, Albanese D, Stegen J, Franceschi P, Coller E, Zanzotti R, Ioriatti C, Stefani E, Pindo M, Cestaro A, Donati C. 2023. Distinct and Temporally Stable Assembly Mechanisms Shape Bacterial and Fungal Communities in Vineyard Soils. Microb Ecol 86:337–349.

100. Zhang G, Wei G, Wei F, Chen Z, He M, Jiao S, Wang Y, Dong L, Chen S. 2021. Dispersal Limitation Plays Stronger Role in the Community Assembly of Fungi Relative to Bacteria in Rhizosphere Across the Arable Area of Medicinal Plant. Front Microbiol 12.

101. Su Y, Wang J, Gao W, Wang R, Yang W, Zhang H, Huang L, Guo L. 2023. Dynamic metabolites: A

bridge between plants and microbes. Sci Total Environ 899:165612.

102. Robertson-Albertyn S, Alegria Terrazas R, Balbirnie K, Blank M, Janiak A, Szarejko I, Chmielewska B, Karcz J, Morris J, Hedley PE, George TS, Bulgarelli D. 2017. Root Hair Mutations Displace the Barley Rhizosphere Microbiota. Front Plant Sci 8.

103. Idriss EE, Makarewicz O, Farouk A, Rosner K, Greiner R, Bochow H, Richter T, Borriss R. 2002. Extracellular phytase activity of Bacillus amyloliquefaciens FZB45 contributes to its plant-growth-promoting effectaaThe GenBank accession numbers for the sequences determined in this work are AY055219 to AY055226. Microbiology 148:2097–2109.

104. Lozo J, Danojević D, Jovanović Ž, Nenadović Ž, Fira D, Stanković S, Radović S. 2022. Genotype-Dependent Antioxidative Response of Four Sweet Pepper Cultivars to Water Deficiency as Affected by Drought-Tolerant Bacillus safensis SS-2.7 and Bacillus thuringiensis SS-29.2 Strains. 3. Horticulturae 8:236.

105. Tiwari R, Kalra A, Darokar MP, Chandra M, Aggarwal N, Singh AK, Khanuja SPS. 2010. Endophytic Bacteria from Ocimum sanctum and Their Yield Enhancing Capabilities. Curr Microbiol 60:167–171.

106. Albrechtova J, Latr A, Nedorost L, Pokluda R, Posta K, Vosatka M. 2012. Dual Inoculation with Mycorrhizal and Saprotrophic Fungi Applicable in Sustainable Cultivation Improves the Yield and Nutritive Value of Onion. Sci World J 2012:374091.

107. Wu J, Kawagishi H. 2020. Plant growth regulators from mushrooms. J Antibiot (Tokyo) 73:657–665.

108. Xia X, Wei Q, Wu H, Chen X, Xiao C, Ye Y, Liu C, Yu H, Guo Y, Sun W, Liu W. 2024. Bacillus species are core microbiota of resistant maize cultivars that induce host metabolic defense against corn stalk rot. Microbiome 12:156.

109. Yang C, Yue H, Ma Z, Feng Z, Feng H, Zhao L, Zhang Y, Deakin G, Xu X, Zhu H, Wei F. 2022. Influence of plant genotype and soil on the cotton rhizosphere microbiome. Front Microbiol 13.

110. Dilla-Ermita CJ, Lewis RW, Sullivan TS, Hulbert SH. 2021. Wheat Genotype-Specific Recruitment of Rhizosphere Bacterial Microbiota Under Controlled Environments. Front Plant Sci 12.

111. Toth IK, Bell KS, Holeva MC, Birch PRJ. 2003. Soft rot erwiniae: from genes to genomes. Mol Plant Pathol 4:17–30.

112. Abe K, Iwanami H, Kotoda N, Moriya S, Takahashi (Sumiyoshi) S. 2010. Evaluation of apple genotypes and Malus species for resistance to Alternaria blotch caused by Alternaria alternata apple pathotype using detached-leaf method. Plant Breed 129:208–218.

113. Sobiczewski P, Peil A, Mikiciński A, Richter K, Lewandowski M, Żurawicz E, Kellerhals M. 2015. Susceptibility of apple genotypes from European genetic resources to fire blight (Erwinia amylovora). Eur J Plant Pathol 141:51–62.

114. Wegulo SN, Yang XB, Martinson CA. 1998. Soybean Cultivar Responses to Sclerotinia

sclerotiorum in Field and Controlled Environment Studies. Plant Dis 82:1264–1270.

115. Yang Q, Yin G, Guo Y, Zhang D, Chen S, Xu M. 2010. A major QTL for resistance to Gibberella stalk rot in maize. Theor Appl Genet 121:673–687.

116. Crute I, Pink D. 1996. Genetics and Utilization of Pathogen Resistance in Plants. Plant Cell 8:1747–1755.

117. Laine A-L. 2009. Role of coevolution in generating biological diversity: spatially divergent selection trajectories. J Exp Bot 60:2957–2970.

118. Thompson JN. 2005. Coevolution: The Geographic Mosaic of Coevolutionary Arms Races. Curr Biol 15:R992–R994.

119. Thrall PH, Laine A-L, Ravensdale M, Nemri A, Dodds PN, Barrett LG, Burdon JJ. 2012. Rapid genetic change underpins antagonistic coevolution in a natural host-pathogen metapopulation. Ecol Lett 15:425–435.

120. dos Santos IB, Pereira AP de A, de Souza AJ, Cardoso EJBN, da Silva FG, Oliveira JTC, Verdi MCQ, Sobral JK. 2022. Selection and Characterization of Burkholderia spp. for Their Plant-Growth Promoting Effects and Influence on Maize Seed Germination. Front Soil Sci 1.

121. Harder CB, Hesling E, Botnen SS, Lorberau KE, Dima B, von Bonsdorff-Salminen T, Niskanen T, Jarvis SG, Ouimette A, Hester A, Hobbie EA, Taylor AFS, Kauserud H. 2023. Mycena species can be opportunist-generalist plant root invaders. Environ Microbiol 25:1875–1893.

122. Lemessa F, Zeller W. 2007. Pathogenic characterisation of strains of Ralstonia solanacearum from Ethiopia and influence of plant age on susceptibility of hosts against R. solanacearum / Charakterisierung der Pathogenese verschiedener äthiopischer Stämme von Ralstonia solanacearum und Einfluss des Pflanzenalters auf die Anfälligkeit von Wirten gegenüber dem Bakterium. J Plant Dis Prot 114:241–249.

123. Liu Q, Beattie GA, Saalau Rojas E, Gleason ML. 2018. Bacterial wilt symptoms are impacted by host age and involve net downward movement of Erwinia tracheiphila in muskmelon. Eur J Plant Pathol 151:803–810.

124. Ando K, Carr KM, Colle M, Mansfeld BN, Grumet R. 2015. Exocarp Properties and Transcriptomic Analysis of Cucumber (Cucumis sativus) Fruit Expressing Age-Related Resistance to Phytophthora capsici. PLOS ONE 10:e0142133.

125. Hilton S, Bennett AJ, Keane G, Bending GD, Chandler D, Stobart R, Mills P. 2013. Impact of Shortened Crop Rotation of Oilseed Rape on Soil and Rhizosphere Microbial Diversity in Relation to Yield Decline. PLOS ONE 8:e59859.

126. Bennett AJ, Hilton S, Chandler D, Mills P, Bending GD. 2020. Relationships between yield, rotation length, and abundance of *Olpidium brassicae* and *Pyrenochaeta* sp. in the rhizosphere of oilseed rape. Appl Soil Ecol 147:103433.