# Spatial transcriptomics and genetically implicated genes identify putative causal tissue structures for complex traits

Linda Kvastad[1‡*], Aaron Kollotzek[1], Rasool Saghaleyni[2], Chih-Fan Chang[1], Tianze Cao[1] and Tuuli Lappalainen[1,3‡]

[1]Science for Life Laboratory, Department of Gene Technology, KTH Royal Institute of Technology, Stockholm, Sweden.

[2]Department of Life Sciences, National Bioinformatics Infrastructure Sweden, Science for Life Laboratory, Chalmers University of Technology, Gothenburg, Sweden

[3]New York Genome Center, New York, NY, USA.

‡ Corresponding authors: Linda Kvastad (linda.kvastad@ki.se); Tuuli Lappalainen (tuuli.lappalainen@scilifelab.se)

* Current affiliation
Department of Cell and Molecular Biology, Karolinska Institute, Solna, Sweden

## Abstract

Spatially resolved transcriptomics is transforming our understanding of cellular and molecular diversity of tissues. Here, to identify tissue structures that are enriched for putatively causal disease processes, we integrated 31 human and mouse spatial datasets from 8 organs with genes that are genetically implicated in 32 human diseases. Applying our novel approach STEAM, we identified both known and novel roles of tissue structures in human diseases and identified spatially clustered disease gene subsets. For example, in the brain we observed enrichment of neuropsychiatric disease in cortical layers, and immune and barrier dysfunction in Alzheimer's disease, with integration with single-cell data highlighting the complementary insights from the two data types. Spatial coexpression of drug target genes with genetically implicated genes in enriched tissue structures showed potential for spatially informed drug repurposing. Altogether, we show vast potential for integration of genetic discoveries with growing spatial datasets to understand human disease biology.

Spatially resolved transcriptomics (SRT) is a rapidly expanding field that has provided unprecedented insights into cellular and molecular variation within tissues, advancing our understanding of tissue structures and their biological functions[1,2]. While much of the field has focused on identifying structures, annotating, and integrating SRT datasets themselves, future opportunities lie in interrogating the biological roles of the identified tissue structures. One key approach to this is gene set enrichment analysis, which asks whether a set of genes – such as genes implicated in a particular disease – are enriched in SRT datasets.

Genome-wide association studies (GWAS) have been critical for identifying genetic variants associated with complex traits and diseases[3]. As genetic data is effective in pinpointing putatively *causal* disease processes - rather than pathophysiological processes caused by the disease or its symptoms - they are being increasingly leveraged in drug development. Indeed, it has been shown that selecting genetically supported drug targets increases new drugs' clinical development success rate[4–6]. This has inspired major efforts, including the Open Targets platform[7,8] that provides a systematic resource listing genes implicated in diverse traits. However, further functional studies are needed to understand the underlying causal biological mechanisms. GWAS data has been combined with bulk and single-cell RNA sequencing (RNA-seq) and epigenomic marks to identify enrichments of GWAS signal in different tissues and cell types[9–21]. SRT datasets provide emerging opportunities to assess GWAS enrichment across tissue structures in multiple organs, traits and datasets[22]. Previous studies have mainly focused on the direct use of GWAS summary statistics. Here, we utilize the Open Targets platform[7,8], where data from multiple GWAS are distilled into trait-specific ranked gene lists. We then use spatially resolved gene expression from healthy tissues to pinpoint the annotated structures where disease-implicated genes are enriched, implicating these structures as potential sites of casual disease processes.

However, such analysis is complicated by the specific characteristics of SRT datasets. Most notably, it is known to be sparse, and the number of genes and unique molecular identifiers (UMIs) are often unevenly distributed. To overcome these challenges, we developed an approach called **S**patial **T**rait **E**nrichment **A**nalysis with per**M**utation testing (STEAM), a robust and scalable computational approach to measure the enrichment of average gene expression across clusters in a spatial dataset from a given gene list; it calculates a permutation p-value and performs multiple testing corrections based on the number of clusters. For ranked gene lists, STEAM can set an empirically derived threshold beyond which the enrichment signal falls to baseline. Thus, STEAM can perform robust gene set enrichment analysis of spatial structures in SRT datasets.

In this study, we analyzed spatial enrichment of genetically implicated gene sets for 32 complex traits and diseases from Open Targets, and combined these data with 31 previously published SRT datasets from eight organs. Applying STEAM to pinpoint putative causal tissue structures for complex diseases, we discovered numerous tissue structure enrichments. For example, our results for Alzheimer disease (ALZ) imply mechanisms beyond amyloid and tau pathology in barrier and

102 immune dysfunction. Furthermore, we analyzed spatial enrichment of drug target
103 genes for these diseases, providing insights into their overlap with genetic evidence
104 and potentially advancing future drug repurposing efforts.
105

## Results

### Spatial trait enrichment analysis with permutation testing

We analyzed 31 SRT datasets, spanning eight organs[23-35] (Supplementary Table 1).
The spatial structures of each dataset were determined using standard clustering
approaches, which included non-negative matrix factorization to identify spatial gene
expression patterns (Supplementary Fig. 1-4, Supplementary Table 2-3). We
analyzed two biological replicates from humans and/or mice in most organs, and
included multiple developmental time points for the heart, intestine, pancreas, and
spinal cord. For each tissue we selected the SRT dataset of best quality as an index
dataset, treating others as replicates with results shown in supplementary materials,
except for the developmental stages where only one sample per stage was included.
We used the Open Targets platform to compile lists of genetically associated
and drug target genes for 32 complex traits. Many genes are targets for drugs used in
multiple diseases, and some genes are implicated in multiple diseases, but we
observed low sharing between genetically associated and drug target genes
(Supplementary Fig. 5). With our main focus being enrichment in structures within
tissues, we then used previously implicated links between traits and tissues to select
trait-tissue pairs for enrichment analysis[8] (Figure 1a, Supplementary Table 4).
To test whether the median expression level of each gene set is higher than the
null in the spatial structures of each SRT dataset, we applied our new approach,
STEAM. For a robust comparison, STEAM considers the uneven data distribution of
genes and transcripts (Supplementary Fig. 6) by creating null gene sets of the same
sizes as the trait gene lists studied and performing 10,000 permutations to calculate
a p-value for each tissue structure. We applied multiple testing corrections within
each disease-SRT dataset pair, and all the reported p-values are after Bonferroni
correction based on the number of spatial structures in a given SRT dataset (Figure
1b-d). The robustness of the enrichment results given the uneven sparsity of SRT
datasets was verified by lack of correlation between the number of genes or UMIs
and nominal p-values (Supplementary Fig. 7).
In the Open Targets platform[7,8], many complex traits have up to several
hundreds of genetically implicated genes (Figure 1a), with an accompanied ranking
score indicating the estimated degree of support for each gene-trait association.
However, it is not clear whether all or an unknown top tier subset of these genes are
truly trait-associated. To investigate this empirically, we analyzed spatial trait
enrichments with STEAM for the selected trait-tissue pairs using a sliding window
approach, scanning each gene list from top to bottom using a window size of 50
genes. For many traits, we observed a clear enrichment for highly ranked genes that
then decreased to null. We assigned these genes as "top genes" based on a nominally

significant sliding window enrichment for at least one tissue structure above the
0.95-quantile (Figure 1 e).
146

**Figure 1. Illustration of the approach and the STEAM pipeline. a)** Selected trait-tissue pairs, spatial
datasets and gene sets included in the study. **b)** Hematoxylin and Eosin (HE) stained mouse brain coronal tissue
section (left) and clusters from SRT data (right). **c)** Spatial distribution of number of genes (left) and unique
molecular identifiers (UMIs) (right). **d)** Null distribution of the median score for genetically implicated
Alzheimer's genes for cluster 3 in (b). **e)** Median scores for ALZ cluster 3 in (b) in a sliding window of 50 genes,
compared to null distribution ranges marked with dotted lines. ADHD=attention deficit hyperactivity disorder.
AHD= anterior horn disorder. ALS=amyotrophic lateral sclerosis. ALZ=Alzheimer disease. ASD=autism
spectrum disorder. AST=asthma. BIP=bipolar disorder. BRD=bronchial disease. CAD=coronary artery disease.
CD=Crohn's diseases. CHD=congenital heart disease. CHOL=total cholesterol measurement. CIRR=cirrhosis of
liver. CKD=chronic kidney disease. COPD=chronic obstructive pulmonary disease. EPI=epilepsy.
IBD=inflammatory bowel disease. IKD=inherited kidney disorder. INS=insomnia. LD= liver disease. LDL=low
density lipoprotein cholesterol measurement. LID=large intestine disorder. LUD=lung disease. MDD=major
depressive disorder. MI=myocardial infarction. MS=multiple sclerosis. NAFLD=non-alcoholic fatty liver disease.
PAR=parkinson's disease. SCZ=schizophrenia. T1D= type 1 diabetes mellitus. T2D= type 2 diabetes mellitus.
UC=ulcerative colitis.
163

<u>Putative causal tissue structures for complex traits</u>

We investigated complex trait enrichments for selected trait-tissue pairs in SRT
datasets from eight organs (Figure 1 a) and their structures, consisting of 56,516
spatial positions (Figure 2, Supplementary Fig. 8-17). Supplementary Table 5
provides p-values for all trait-structure pairs, including those cited below in the text.
We observed a robust signal of enrichment of all genetically implicated genes in the
selected trait-tissue pairs ($\pi_1$ = 0.55, Figure 2 b) as well as drug targets ($\pi_1$ = 0.69,
Figure 2 c). The much lower signal for other trait-tissue pairs ($\pi_1$ = 0.1 for all
genetically implicated genes, Supplementary Figure 8 c) supported the focus on
selected trait-tissue pairings to maintain statistical power.

Inspecting the proportion of significant structures across tissues for all
genetically associated genes, top genes, and drug target genes of selected trait-tissue
pairs, we found that in most cases, all or none of the structures were enriched.
However, interestingly, we found many trait-tissue pairs (16% to 20% for the three
gene classes) where only some structures are significant, pointing to trait
enrichments that are specific to tissue structures (Figure 2 f). Examples of these
patterns are described below. The signal replicated well between true biological
replicates and across species (when data was available, Supplementary Table 5,
supplementary figures referred to below).
We next assessed how well current drug targets align with genetic evidence.
Structures significantly enriched for genetically implicated genes were often also
enriched for drug targets for the same disease ($\pi_1$ = 0.82, Figure 2 d), indicating that
known drugs target biologically relevant tissue structures. However, the lower
proportion of structures enriched for drug targets that are also enriched for
genetically implicated gene sets ($\pi_1$ = 0.51, Figure 2 e) suggests that drug targets are
often expressed also in other tissue structures than those supported by genetics.
Similar results were observed when comparing top genes and drug targets
(Supplementary Fig. 8 d-g).
192

193
**Figure 2. Summary of gene expression enrichments in tissue structures. a)** Enrichments across traits
194 in all tissues for all genetically implicated genes, top-ranked genetically implicated genes, and drug target genes,
195 visualized for each index sample in each tissue and all included developmental stages. **b-c)** Enrichment nominal
196 p-value distribution of trait-structure pairs, with the proportion of true positives measured as $\pi_1$ statistics for **b)**
197

selected trait-tissue pairs for all genetically implicated genes and **c)** selected trait-tissue pairs for drug targets. **d-**
**e)** Sharing of enrichment signal, with **d)** enrichment p-value for drug target genes for trait-structure pairs that
were significant for genetically implicated genes, **e)** enrichment p-value for genetically implicated genes for trait-
structure pairs that were significant for drug target genes for the same trait. Red dotted line = 0.05. **f)**
Proportions of significant structures per selected trait-tissue pair.

<u>Spatial enrichments of non-brain tissues</u>

For some traits, we observed significant enrichments for genetically implicated genes
across the tissue, indicating a broad role of multiple structures in these traits
(Supplementary Table 5). Examples of this pattern include adult human heart for
congenital heart disease (CHD) (Supplementary Fig. 9 a), liver for low-density
lipoprotein cholesterol measurement (LDL), liver for total cholesterol measurement
(CHOL), and developmental pancreas during post-conceptual weeks (pcw) 18-20 for
type 1 diabetes mellitus (T1D) (Supplementary Fig. 11 e, f).
We also observed cases where a smaller subset of tissue structures showed an
enrichment, which highlights the value of spatial data and pinpoints interesting
features of disease biology. For example, in the adult heart for coronary artery
disease (CAD) we observed enrichment only around the vessels (Supplementary
Figure 9 a), which aligns with the known endothelial dysfunction in CAD[36]. For the
human dataset of the kidney cortex we observed a broad enrichment for inherited
kidney disorder (IKD), in contrast to chronic kidney disease (CKD) where we only
saw enrichment in the tissue structures of glomerulus and proximal tubule (p = 0.04;
Supplementary Fig. 11 a). In mouse kidney samples both IKD and CKD had cortex
enrichments, with CKD having the strongest enrichments in glomerulus and
proximal tubule (Supplementary Fig. 11 c). Interestingly, emerging evidence points
towards complex interactions between glomeruli and tubules underlying progressive
kidney injury, where damage to one tissue structure can cause injury to the other[37].
Our observations for CKD support such a model where two different tissue structures
in the same tissue region, here the kidney cortex, could contribute to dysfunction in
disease. In the lung, we saw enrichment across the parenchyma tissue for broadly
defined lung disease (LUD); this was observed both across the parenchyma datasets
and within the bronchi dataset that included a smaller spatial structure of
parenchyma tissue (Supplementary Fig. 12 b-d). In the intestine, we observed the
strongest enrichment towards the top of the villus for the umbrella category of large
intestine disorder (LID) (Supplementary Fig. 13 a), with similar non-significant
enrichment patterns in all other datasets (Supplementary Fig. 13 b-e). In addition,
using a SRT intestine dataset with a strategy for library preparation that excluded
mitochondrial and ribosomal genes we observed enrichments in lymphoid follicles
for Crohn's diseases (CD), and for ulcerative colitis (UC) in lymphoid follicles and
part of the submucosa, and part of the submucosa for inflammatory bowel disease
(IBD) (Supplementary Fig. 13 b). Similar enrichment patterns in lymphoid follicles
were observed for CD in the other adult human and mouse intestine datasets
(Supplementary Fig. 13 a and e). CD is known to be an immune mediated disorder
with lymphatic dysfunction[38].

 Spatial enrichments across development

SRT data across developmental time points provides an opportunity to understand
how disease-relevant structures manifest during tissue development. For some
datasets and diseases, we observed patterns that were generally consistent between
time points. For the developmental spinal cord datasets, we observed enrichments
for anterior horn disorder (AHD) and amyotrophic lateral sclerosis (ALS) in the
anterior ventral horn gray matter tissue (Supplementary Fig. 14), where others have
previously found loss of synapses in the ventral horn for ALS[39]. In the human heart
from fetal to adulthood, we analyzed two different human developmental time points
and a biopsy from the adult human heart (Supplementary Fig. 15), finding congenital
heart disease (CHD) to have the strongest enrichments across the heart muscle tissue
and CAD around vessel structures.
We analyzed T1D gene set enrichment across developmental ages in SRT
datasets from the human developmental pancreas at 15, 18, and 20 post-conceptual
weeks (pcw) (Figure 3, Supplementary Fig. 11 d-f). While T1D is usually diagnosed in
older children, its developmental etiology is unclear, and it can arise even early in
life[40]. During pcw 15, we only observed significant enrichment for T1D genes in
structures containing a mixture of exocrine cells (expressing *CTRB1* and *CTRB2*) and
endocrine cells (expressing *INS*) (Figure 3 b, c; Supplementary Table 5). In contrast,
during pcw 20, all tissue structures were significantly enriched for T1D, although
exocrine and endocrine cell-rich structures had the smallest p-values (Figure 3 b,
Supplementary Table 5). We further investigated how different subsets of genes
contributed to these trait-specific spatial enrichment patterns. We computed spatial
coexpression as Spearman correlation of the expression level per spot between all
top-ranked genetically associated genes. While most genes did not cluster in a clear
way, we identified a small spatially correlated group of key genes, including *INS*,
*CTRB1* and *CTRB2* (Figure 3 d). The correlation and anti-correlation pattern show
that during pcw 15 and 18, exocrine cells and endocrine cells were spatially colocated
and became spatially separated as development progressed (pcw 20) (Figure 3 d).
Interestingly, while T1D has been seen as a mainly endocrine (β-cell) driven
autoimmune disease, evidence has been mounting for an important role of the
pancreatic exocrine cells[14,41]. Our observations support the importance of both
exocrine and endocrine cells in T1D disease and exemplify reorganization of disease-
relevant cell types during tissue development that can be captured in SRT data.
277

278
**Figure 3. Diabetes type 1 mellitus (T1D) enrichment in the pancreas during human development.**
**a)** HE-stained pancreas sections for post-conceptual week (pcw) 15 (left), pcw 18 (middle) and pcw 20 (right). **b)**
Bonferroni-adjusted p-values for T1D using the top 130 genetically associated genes for the developmental time
points. The red arrows show the location of exocrine and endocrine cells. **c)** Expression of selected genes for the
time points. **d)** Spatial gene-gene correlation heatmap, including all top genes (top) and zoomed-in top left corner
(10 genes) of each heatmap (bottom) for the time points. Post-conceptual weeks (pcw). Scale bar 500µm.
285
286

## Spatial and single-cell enrichment of genes implicated in Alzheimer's and neuropsychiatric disorders

The brain is a highly complex organ with distinct regions performing specific functions. Many neurological and psychiatric disorders have been linked to specific brain regions using neuroimaging approaches[42], and GWAS enrichment analysis implicating causal cell types[9,12–21]. Here, we investigated spatial enrichment patterns for multiple diseases in the brain (Figure 2 a, Supplementary Fig. 16-17) using the mouse index dataset with the highest quality and coverage of a coronal brain section. In epilepsy (EPI) and Parkinson's disease (PAR) we observed a broad enrichment across the tissue (Supplementary Fig. 16 a). We found that top-ranked genetically associated genes for ALZ were enriched for choroid plexus, thalamus, and white matter (Figure 4). These findings align with existing literature, which suggests that alterations in the choroid plexus, such as aging-related changes in cerebrospinal fluid dynamics and cytokine levels[43] play a critical role in ALZ pathology. The thalamus's involvement in early ALZ stages is reported to be characterized by gray matter decline[44], and age and severity of dementia with white matter changes[45], which further supports the relevance of the regions we identified in understanding the disease's progression. We observed some similarities and interesting differences between mice and human datasets: While comprehensive SRT analysis of the human brain is complicated by its size, both species had significant enrichments in their white matter areas, whereas humans also showed enrichment towards the outer layers of the cortex, in particular layer 1. In addition, in one human sample, we observed the strongest enrichment in spatial structures containing B-cell-related genes, e.g., *IGKC* (Figure 4 a, Supplementary Fig. 18 a, b).

For neuropsychiatric disorders that share parts of their etiology as well as genetic hits[46] (Supplementary Fig. 18 c), we found that top-ranked genetically associated genes for attention deficit hyperactivity disorder (ADHD), autism spectrum disorder (ASD), bipolar disorder (BIP), and schizophrenia (SCZ) were enriched in the outer layers of the cortex in the mouse dataset (Figure 4 a). In the human dorsolateral prefrontal cortex[24] we observed a significant enrichment for ASD in layer 2, but not for BIP and SCZ (Supplementary Fig. 17 a). Our results align with previous studies: particularly the ASD enrichment in layer 2/3 has been reported by others[24,47]. While Maynard et al.'s prior pseudo-bulking approach[24] had a lower resolution than ours, it discovered a more pronounced enrichment pattern for BIP and SCZ. Our observations may be due to data sparsity when analyzing sections individually.

In addition to structure enrichments across the entire gene sets, we further investigated how different subsets of genes contributed to these trait-specific spatial enrichment patterns. We computed spatial coexpression as Spearman correlation of the expression level per spot between all top-ranked genetically associated genes for ALZ, ADHD, ASD, BIP, and SCZ in the mouse index dataset (Figure 4 b-c). Some disease-associated genes clustered into spatially colocalized subsets that may contribute to the disease via distinct mechanisms.

**Figure 4. Spatial enrichments of brain related traits. a)** Mouse brain coronal tissue section with HE
staining and annotation of selected regions (top left), and trait-specific Bonferroni adjusted permutation p-values
for each spatial cluster, using top genetically associated genes for ALZ (top 70 genes), ADHD (top 200 genes),
ASD (top 130 genes), BIP (top 80 genes) and SCZ (top 140 genes). **b)** Spatial gene-gene correlation heatmap,

336  including all top genes for each trait. **c)** Spatial gene expression of selected genes from the heatmap in (b). Scale
337  bar 500μm.
338
As each structure in a SRT dataset contains a mixture of multiple cell types, we
hypothesized that SRT and single-cell RNA-seq (scRNA-seq) datasets provide
complementary insights into spatial structures and specific cell types, respectively.
To investigate this, we analyzed the Allen Brain Atlas scRNA-seq dataset of the
mouse cortex and hippocampus[48], unfortunately lacking data for the choroid plexus,
the thalamus and the white matter that had spatial enrichments in ALZ. Thus, we
also analyzed the current most comprehensive human brain single-nuclei RNA-seq
(snRNA-seq) dataset[49]. Cell type deconvolution[50] of the mouse SRT dataset showed
biologically meaningful structure-specific cell type proportions (Figure 5 a, b). Next,
we applied STEAM to the sc/snRNA-seq datasets to investigate the cell type
enrichments of genetically associated genes. To decrease the sparsity of datasets we
applied a meta cell approach (see methods), and to empirically redefine the genetic
top genes we used the sliding window approach (Supplementary Fig. 19 - 23).
Supplementary Table 6 and 7 contain all the p-values for cell type - trait pairs
described below in the text. We also did a complementary enrichment analysis of
genetically associated genes among the genes that were upregulated and
differentially expressed genes (DEGs) in each cell type, using Fisher's exact test
(Figure 5 c, d; Supplementary Fig. 24 and 25).
For ALZ, in mouse we observed the strongest enrichments of top genes in
microglia/perivascular macrophage (Micro-PVM) cells, vascular/leptomeningeal
cells (VLMC), endothelial cells, and astrocytes (Supplementary Fig. 19 a), with
complementary enrichment in both species showing similar results, with the
strongest enrichment in microglia (Figure 5 c, Supplementary Fig. 24 i and 25 a, c).
Of these, VLMC and endothelial cells were quite specific to choroid plexus structures
that were enriched for ALZ genes (Figure 4 a: choroid plexus 1; Figure 5 a: structure
14). In contrast, Micro-PVM cells showed little structure-specific enhanced
contribution and were widely spread across all tissue structures (Figure 5 a and b),
indicating potential contributions in disease processes that can take place across the
brain. The human snRNA-seq dataset recapitulated many of these patterns,
including enrichments in the choroid plexus and vascular cells, and various stages of
developing oligodendrocytes (Supplementary Fig. 21 a and 25 a, c). The cell types
underlying ALZ enrichment in the white matter and in the thalamus were less clear,
but both were enhanced for oligodendrocytes with some ALZ enrichment (structure 2
and 3, respectively in Figure 5 a and b).
For ADHD, ASD, BIP and SCZ, we observed broad significant enrichments of
top genes across the mouse neuronal cell types, astrocytes and oligodendrocytes
(Supplementary Fig. 19 and 20). In addition, the complementary enrichment showed
the strongest signal for the neuronal cell types (Supplementary Fig. 24). The spatial
structures enriched for neuropsychiatric disease genes had a high proportion of these
cell types, such as L4/5 and L2/3 intratelencephalic neurons in the isocortex (IT
CTX) (Figure 5 a, b). In the human snRNA-seq dataset covering cell types from the
whole brain, using the complementary analysis of differential expression, e.g., in SCZ

the strongest signals from top genes were found in hippocampal cornu ammonis

(CA) region CA4, eccentric medium spiny neurons and regions CA1-3 (Figure 5 d),

and for all genetically associated genes in committed oligodendrocyte precursor cells

(OPCs) and medial ganglionic eminence (MGE) interneurons (Supplementary Fig. 25

i). These findings align with previous literature for SCZ enrichment in medium spiny

neurons, hippocampal pyramidal cells, certain interneurons and

oligodendrocytes[15,16], as well as more recent findings of a broad enrichment among

neuronal cell types, and MGE interneurons in SCZ[21]. These examples showcase the

complementary power of utilizing both SRT and scRNA-seq to investigate cell and

tissue enrichments of complex traits. Our observations in mouse and human points

towards possible trait contributions from more than one cell type in the same spatial

structure.

**Figure 5. Single-cell contribution to spatial enrichments in the brain. a)** Cell type composition of the tissue structures of the index mouse coronal tissue section (Figure 1 b, Supplementary Figure 1 a), using the Allen Brain Atlas of the mouse cortex and hippocampus[48] scRNA-seq data. Enriched spatial structures are marked with a color legend, with **b)** selected cell types shown. Scale bar 1mm. **c-d)** Trait enrichment across cell types for top genetically associated genes among differentially expressed genes **c)** for ALZ across subclasses in the mouse cortex and hippocampus[48] and **d)** for SCZ across superclusters in the adult human brain[49]. The full list of mouse cell subclass abbreviations can be found in Supplementary Table 6; those shown in **(b)** are: VLMC = Vascular/leptomeningeal cells. Endo = Endothelial cells. Oligo = Oligodendrocytes. Pvalb = Pvalb positive cells. Micro-PVM = Microglia/perivascular macrophages. L2/3 IT CTX = Layer 2/3 intratelencephalic neurons in the isocortex. Human cell supercluster abbreviations; CA = Cornu Ammonis. MGE = Medial Ganglionic Eminence. CGE = Caudal Ganglionic Eminence. Grey dotted line, adj. p-value = 0.05.

## Potential for spatially informed drug repurposing strategies

Repurposing approved drugs for new diseases provides an important and increasing line of drug development. We investigated the potential for spatial enrichment patterns to provide additional information to identify repurposing opportunities. Using the brain to demonstrate this principle, we leveraged spatial coexpression of all genes genetically implicated in a given disease (Figure 1 a) to pinpoint those that are expressed in structures identified as disease-relevant (Figure 4) and also targets of known drugs, targeting brain or non-brain traits (Figure 6, Supplementary Fig. 26-28, and Supplementary Table 8). For example, ALZ-implicated gene *Kif5b* is a target for selpercatinib and pralsetinib used in cancer treatment. It was spatially colocalized with genes such as *Bin1* expressed in the white matter, which was overall enriched for ALZ genes (Figure 6 a and 4 c). Interestingly, others have recently reported that a decrease in *KIF5B* levels could slow down and/or prevent abnormal tau behavior in animal and cell models[51], making it an interesting drug repurposing candidate. In addition, the spatially colocalized ALZ-implicated gene *Clu* is spatially colocalized in the same structures, is also a target for the cancer drug custirsen, and has been suggested to play a role in amyloid beta peptide clearance in an animal model[52] (Figure 6 a). Voltage-gated potassium and calcium channel genes such as *KCNB1*, *KCNV1*, and *CACNA2D1*, are associated to ADHD, ASD, BIP and SCZ[53−56] and targets for multiple drugs used in diverse diseases such as multiple sclerosis, gastrointestinal disease, cardiovascular disease, and restless legs syndrome. They were spatially coexpressed with other genes of the cortex that showed a clear overall enrichment for these traits (Figure 6 b, Supplementary Fig. 26-28). Similar spatial gene expression patterns were observed for these genes in the human cortex dataset (Supplementary Fig. 29). This supports the recent rekindled interest in application of these drugs in neuropsychiatric disease[57]. Our observations suggest that considering spatial enrichment of genetically implicated genes could be an interesting additional layer of evidence for identifying potential drug targets and drug delivery locations that intervene with causal processes of disease.

437
438 **Figure 6. Spatial patterns of drug targets.** Gene-gene spatial correlation heatmaps for the index mouse
439 brain coronal tissue section, for genes genetically implicated in **a)** ALZ and **b)** BIP, with their drug target status
440 indicated. Spatial gene expression of selected genes from the heat maps are shown on the side, scale bar 500μm.

## Discussion

Localizing causal disease processes in human tissues is a key part of the quest to understand disease biology and developing interventions. Integrating GWAS data with emerging spatial datasets provides a new opportunity for this. Here, we presented STEAM, a generalizable approach for spatial trait enrichment analysis with permutation testing, and applied it to genetically implicated genes and drug targets for diverse diseases and SRT datasets from a large number of tissues and developmental stages, as well as two supporting brain sc/snRNA-seq datasets. While genes for some diseases are enriched tissue-wide, in up to a fifth of our spatial data we found evidence pointing towards tissue structure-specific trait enrichments, highlighting the value of the spatial resolution. Many of these discoveries were consistent with known biology, but we also identified links between diseases and tissue structures that are not well described in prior literature and indicate potentially interesting targets for further follow-up. While here we focused on 38 disease-tissue pairs based on prior indications, we anticipate that larger data sets with better spatial resolution, replicates, and higher accuracy in linking genes to diseases will further boost the power to allow even broader discovery across all tissues.

Given its refined structures with different biological functions, the brain provides a particularly interesting target for spatial enrichment analysis and further exploration of how spatial enrichments compare to single-cell analysis. Our results show clearly the previously known enrichment of neuropsychiatric disorders ADHD, ASD, BIP, and SCZ in the outer layers of the cortex and cell types such as L4/5 and L2/3 IT CTX that are present in these structures. For ALZ we identified spatial enrichments in part of the ventricles for choroid plexus, thalamus, and white matter in the mouse SRT data. In addition, for the human SRT datasets we observed enrichments in cortical structures, where impaired plasticity has been associated with impaired working memory[58]. While we only observed spatial structures containing B-cell-related genes in the human brain samples, we note that others have shown that B-cell depletion in mice reverses disease progression[59]. The sc/snRNA-seq analysis showed ALZ gene enrichment in VLMC and endothelial cells that are both quite specific to the highly vascularized choroid plexus, surrounding and inside blood vessels and with a role in maintaining barrier integrity[60–62]. On the other hand, the cell type well known for high enrichment of ALZ signal, microglia, were present across multiple brain structures. These results suggest ALZ mechanisms beyond amyloid and tau pathology, involving barrier and immune dysfunction. More generally, these results indicate that some disease processes can arise from dysfunction in a particular cell type that may or may not be spatially enriched, while others may derive from dysfunction of multiple colocalized cell types jointly (and perhaps interactively) in a given tissue structure. Our results for CKD in the kidney even suggests a scenario where multiple distinct structures in a tissue region contribute to dysfunction. Spatial correlation analyses further help to identify subsets of disease genes potentially involved in different disease mechanisms.

In addition to our analysis of genetically implicated genes, we also analyzed
spatial enrichments of genes that are drug targets for different diseases. These
enrichments often captured the same structures as those discovered via genetics, but
drug target genes were enriched more broadly in structures not implicated by
genetics. We further used our findings to explore strategies for drug repurposing by
identifying situations where a known drug target is not only genetically associated
with the disease but also expressed in a structure enriched for genes of that disease,
further supporting its biological importance and potential as a repurposing target.
Spatially resolved transcriptomics is still a relatively new field, and many of
the limitations of our study arise from the caveats in the current data. Mouse tissues
may not fully recapitulate human biology, but the size of human organs limits and
complicates their spatial analysis. Differences in enrichment results across intestine
datasets highlight the impact of library preparation methods, particularly the
exclusion of mitochondrial genes with protein-coding probe-based library
preparation. While our approach is planned to be robust to varying gene and UMI
count distributions, low-quality datasets will have reduced power. It would be ideal
to have single-cell resolution in spatial data rather than analysis and integration of
both data types in parallel.
Our approach is based on gene sets compiled by Open Targets – rather than
e.g. heritability enrichments of genome annotations[15,16,21] – as this links more directly
to spatial data, facilitates accounting for its properties, and is a natural fit for multi-
species analysis. However, linking genetic associations to genes and scoring these is
not without problems. We empirically identified the top-ranked gene set that showed
a consistent biological signal of spatial enrichment, suggesting that the disease
contribution of some of the lower-ranked genes may be questionable. We anticipate
that future improvements will allow more accurate gene identification and
prioritization and boost the power and resolution of enrichment analyses.
In summary, our study discovered spatial enrichments for a large number of
traits across many organs, developmental stages and cell types. Integrating the
growing spatial and genetic datasets provided a rich source for biological discovery to
characterize mechanisms of human disease.
516
517

## Methods

### Spatially resolved transcriptomics datasets

A summary of the datasets' names and references to their sources can be found in Supplementary Table 1. For each tissue and developmental stage we selected the best quality SRT dataset as an index dataset, as indicated in Supplementary Table 1. Quality was assessed based on the number of genes and UMIs of the dataset, and the HE image field of view including structures of interest. When biological replicates were available, results were shown in supplementary materials. For one SRT dataset (intestine_human_adult_RRST_349_B1) a markedly different 10X library preparation was used, with a transcriptome-wide protein-coding probe-based capture strategy, e.g., excluding mitochondrial and ribosomal genes.

### Spatial transcriptomics data pre-processing

The brain and kidney dataset from 10X Genomics were pre-processed with SpaceRanger (version 1.1.0). Subsequent analyses were carried out using the STUtility[63] (version 1.1.1) and Seurat[64] (version 4.3.0) R-packages. Filtering was performed using the *InputFromTable* function in STUtility, with slightly different settings depending on library preparation. For Visium fresh frozen, Visium RRST, and SRT 2k array library preparations, all spots containing less than 500 UMI counts were removed, and genes were removed if they were present in less than 5 spots or had a total UMI count below 100. For the SRT 1k array library preparations, all spots containing less than 500 UMI counts were removed, and genes were removed if they were present in less than 5 spots or had a total UMI count below 20.

### Defining spatial structures

Each dataset was normalized using the *SCTransform* function in Seurat, returning the top 3000 variable genes. Spatial gene expression patterns were identified using the *RunNMF* function in STUtility, the number of factors used and factors included for clustering are stated in Supplementary Table 5 for each dataset. Clustering was performed using the *FindNeighbors* function in Seurat, based on NNMF factors, where factors were excluded if they i) did not show a clear spatial gene expression pattern, ii) were driven mainly by only one gene, iii) were driven by mitochondrial and/or ribosomal genes, or iv) were driven by hemoglobin genes. Then using the *FindClusters* function in Seurat, with individual clustering resolution for each SRT dataset (Supplementary Table 5). To detect differential gene expression, each dataset was first normalized and scaled using the *NormalizeData* and *ScaleData* functions in Seurat. Differentially expressed genes for each seurat cluster were calculated using the *FindAllMarkers* function in Seurat, with custom settings, only.pos = TRUE, and min.pct = 0.25. All the differentially expressed genes (DEGs) for each dataset and cluster can be found in Supplementary Table 3.

## Tissue structure annotation

Annotation of enriched tissue structures was done based on the original publications (Supplementary Table 1). For the 10X mouse kidney dataset we annotated the cortex based on the structural HE similarities of the mouse dataset from Ferreira et al[26]. For the 10X mouse coronal brain dataset we annotated the enriched structures based on: i) Allen Brain Atlas of mouse coronal sections[65], ii) marker genes from top DEGs of spatial structures associated with specific brain structures according to the Human Protein Atlas[66] (www.proteinatlas.org), iii) and based on the structural HE similarities and cluster annotations of the coronal mouse dataset from Kleshchevnikov et al[31]. In this article, we interchangeably refer to clusters as structures in spatial datasets and cell types in sc/snRNAseq datasets.

## Open Targets gene lists

For genetically associated genes, the platform uses data source specific association scores and calculates a harmonic sum, which is normalized to create scores between 0 and 1. Evidence is also weighted depending on its source, calibrating its relevance compared to the other data sources[8]. For ChEMBL drug target genes, higher scores are given to targets linked to approved drugs and in a descending order based on their stage in clinical trials, and down-weighted based on failure in clinical trials or severe adverse effects[8]. For the Open Targets versions of each gene list, see Supplementary Table 4.

## Open Targets gene list filtering, pre-processing and sliding windows

Gene lists were filtered for each dataset: if a gene on the list was not present in the dataset (after initial dataset filtering) the gene was removed. All genetically associated gene lists were ranked based on the open target association score, while drug targets gene lists were used as is. For mouse datasets, human gene names were converted using the *useMart* function from the biomaRt[67] (version 2.52.0) R-package, using Ensemble version 109 from February 2023. For cases where multiple human genes had the same mouse homolog, we only kept the open target gene list data with the highest overall association score. Sliding window gene sets had a size of 50 genes, starting from the top of the filtered gene list and moved by 10 genes for each subsequent window. Each window gene set score was compared with a null distribution generated from randomly selected genes (see below). Top gene sets were determined based on a nominally significant sliding window enrichment for at least one tissue structure above the 0.95-quantile of the null in the dataset (if window scores were outside the maximum value of the null distribution, this was selected as a cut-off instead).

## Average gene expression level calculation and permutations

The relative average gene expression level score was calculated individually for each dataset first for the entire tissue slide, using dataset specific filtered gene lists for i)

all genetically associated genes, ii) top genetically associated genes (see above) and iii) drug target associated genes, with the *AddModuleScore* function in Seurat. The pre-filtering of gene lists ensured that all genes used for score calculation were present and contributed to the dataset-wide relative scoring. The average expression level score was then subsetted into the spatially defined tissue structures and the median score from each such structure was extracted. We generated 10,000 null distributions by randomly sampling from all genes in the dataset without replacement using the same number of genes in each gene list. We calculated the mean expression level score for each structure with the *AddModuleScore* function. For each tissue structure (or cluster), a permutation p-value was calculated by comparing the median score, generated from dataset specific filtered gene lists for i) all genetically associated genes, ii) top genetically associated genes or iii) drug target associated genes, to comparable null distributions. We calculated the permutation p-value as:

$$p\ value\ =\frac{(Number\ of\ permuted\ statistics\ \geq\ Observed\ statistics)\ +\ 1}{(Total\ number\ of\ permutations)\ +\ 1}$$

Multiple testing correction, using Bonferroni, was applied to all p-values based on the total number of tissue structures (or clusters) in a dataset.

Downstream analyses

We used the Storey-Tibshirani method[68] implemented in the R-package qvalue (version 2.28.0) with the *pi0est* function and pi0.method set to bootstrap[69] to estimate the proportion of true positives (or non-null hypotheses) among sets of hypothesis tests. This was done by first estimating $\pi_0$, the proportions of true null hypotheses, and calculating $\pi_1 = 1 - \pi_0$.

Spearman correlations were calculated in R using the *cor* function. Heatmaps were generated using the *Heatmap* function from the ComplexHeatmap[70] (version 2.15.4) R-package, using hierarchical clustering with complete linkage to find similar clusters.

Single-cell analyses

Single-cell deconvolution of the SRT dataset (brain_mouse_10X) was performed using the *RunNNLS* function from the Semla[50] R-package, with a randomly downsampled (without replacement, seed 42) subset of cells from the Allen Brain Atlas of the mouse cortex and hippocampus (ref). The subsetted cells were restricted to a maximum of 250 single-cells from each subclass, with the exception of Meis2 (n=20) and VLMC (n=152) from which all available cells were included.

To reduce sparseness and increase the power of the single-cell and single-nuclei datasets, we employed a meta cell strategy. The mouse and human brain datasets

were pre-processed according to the following steps and we used the same cluster annotations as the referred to publications of the respective datasets. The single-cell mouse dataset from Allen Brain Atlas of the mouse cortex and hippocampus (ref) was randomly downsampled as described above. We then applied empirical filtering of cells (200 < genes < 8000, 500 < UMI < 35000, mitochondrial-% <= 10, ribosomal-% <= 15) and removed genes with less than 100 UMI counts. The single-nuclei adult human brain datasets[49] for AllNeurons and AllNonNeurons were randomly downsampled (without replacement, seed 123) to a maximum of 250 single-nucleus from each supercluster where subsetted cells were restricted to a maximum of 250 single-cells from each supercluster. We then merged the subsetted datasets for AllNeurons and AllNonNeurons and applied empirical filtering of cells (200 < genes < 15000, 500 < UMI < 125000, mitochondrial-% <= 5, ribosomal-% <= 4) and removed genes with less than 100 UMI counts.

Before generating meta cells, we removed cell types with less than 50 cells (i.e., Meis2). Datasets were then split based on cell type annotation and individually analyzed in Seurat by normalization (*SCTransform*), identification of variable genes (*FindVariableFeatures*) and principal component analysis (*RunPCA*). Clustering was performed using the *FindNeighbors* function and the top 10 principal components. Then using the *FindClusters* function starting at a resolution of 0.1 and iterating through an increase of 0.1 in resolution until yielding a mean number of cells per cluster of ~10. Each cluster within a cell type was then aggregated into a meta cell, and subsequently all meta cells were merged back into one dataset. We used the same cluster annotations as the publications of the respective datasets.

We applied STEAM to calculate gene set enrichment for ALZ, ADHD, ASD, BIP and SCZ. Here, using the sliding window approach to identify top genes and reduce the signal from non-coexpressing genes. Enrichments across ranked gene lists were tested for multiple correction based on number of windows and cell types, using Benjamini & Hochberg. In addition, we performed a complementary enrichment analysis, testing for enrichment of genetically associated genes among the upregulated differentially expressed genes for each cell type using a one-sided Fisher's exact test with function *fisher.test* and alternative set to greater. The complementary enrichment analysis was performed using all and top genes identified with STEAM.

Spatial drug repurposing candidates

We utilized the drug target gene lists from the Open Targets database[7,8], and computed Spearman correlations to look for disease-associated genes grouping in spatially colocalized subsets, which we previously identified as significantly enriched based on the genetically associated gene lists. We included: all genetically associated genes used to determine the trait-specific enrichment, and all drug target genes from the 32 complex traits within this study that also had a genetic association to the trait investigated. We labeled the genes according to their association with the trait

investigated, as i) genetic, ii) a known drug target, iii) a "brain trait" or iv) a "non-brain trait" as depicted in Figure 1a. We constructed correlation-based heatmaps and searched for clusters containing genes with spatially defined expression profiles that implicated the same tissue structures we previously found to have trait-specific enrichments (Supplementary Table 5). This enabled us to select spatially informed drug repurposing candidates from the heatmap clusters and check which other traits these drug repurposing candidates were associated with (Supplementary Table 8).

## Data availability

Links and metadata to published spatial transcriptomics datasets used in this study can be found in Supplementary Table 1. The 10X Genomics dataset for mouse cortex can be found at [https://www.10xgenomics.com/datasets/mouse-brain-section-coronal-1-standard-1-1-0](https://www.10xgenomics.com/datasets/mouse-brain-section-coronal-1-standard-1-1-0) and for mouse kidney at [https://www.10xgenomics.com/datasets/mouse-kidney-section-coronal-1-standard-1-1-0](https://www.10xgenomics.com/datasets/mouse-kidney-section-coronal-1-standard-1-1-0). The scRNA-seq Allen Brain Atlas of the mouse cortex and hippocampus[48], can be found at [https://portal.brain-map.org/atlases-and-data/rnaseq/mouse-whole-cortex-and-hippocampus-10x](https://portal.brain-map.org/atlases-and-data/rnaseq/mouse-whole-cortex-and-hippocampus-10x). The snRNA-seq of the adult human brain[49], can be found at [https://cellxgene.cziscience.com/collections/283d65eb-dd53-496d-adb7-7570c7caa443](https://cellxgene.cziscience.com/collections/283d65eb-dd53-496d-adb7-7570c7caa443).

## Code availability

The STEAM algorithm's computational workflow (Supplementary Fig. 30) and code can be found on Github ([https://github.com/kvastad/STEAM](https://github.com/kvastad/STEAM)), and the STEAM R-package will be available upon the publication of the manuscript.

## Acknowledgements

We thank Enikő Lázár, Ludvig Larsson, Javier Escudero Morlanes and Lovisa Franzén for valuable discussions. This work was supported by a grant from the Knut and Alice Wallenberg Foundation to SciLifeLab for research in Data-driven Life Science, DDLS (grants: KAW 2020.0239 and KAW 2017.0003), and by the National Bioinformatics Infrastructure Sweden (NBIS) at SciLifeLab. The computations were enabled by resources in project (SNIC 2022-5-560 and NAISS 2023-5-517) provided by the National Academic Infrastructure for Supercomputing in Sweden (NAISS) at UPPMAX, funded by the Swedish Research Council through grant agreement no. 2022-06725.

## Author contribution statement

LK and TL designed the study and drafted the manuscript. LK and AK developed the methodology and code. LK performed the core data analysis. LK, AK, CC, and TC performed initial data analysis. R.S. and L.K. co-designed the R-package. R.S. wrote the R-package. All authors reviewed, read, and approved the final manuscript.

## Ethics declarations

T.L. is an advisor and has equity in Variant Bio, was a paid advisor to GSK and has received speaker honoraria from Abbvie and Merck.

## References

1.  Cheng, M. *et al*. Spatially resolved transcriptomics: a comprehensive review of their technological advances, applications, and challenges. *Journal of Genetics and Genomics* **50**, 625–640 (2023).

2.  Duan, H., Cheng, T. & Cheng, H. Spatially resolved transcriptomics: advances and applications. *Blood Science* **5**, 1–14 (2023).

3.  Visscher, P. M. *et al*. 10 Years of GWAS Discovery: Biology, Function, and Translation. *The American Journal of Human Genetics* **101**, 5–22 (2017).

4.  Nelson, M. R. *et al*. The support of human genetic evidence for approved drug indications. *Nat Genet* **47**, 856–860 (2015).

5.  King, E. A., Davis, J. W. & Degner, J. F. Are drug targets with genetic support twice as likely to be approved? Revised estimates of the impact of genetic support for drug mechanisms on the probability of drug approval. *PLoS Genet* **15**, e1008489 (2019).

6.  Minikel, E. V., Painter, J. L., Dong, C. C. & Nelson, M. R. Refining the impact of genetic evidence on clinical success. *Nature* **629**, 624–629 (2024).

7.  Ochoa, D. *et al*. Open Targets Platform: supporting systematic drug–target identification and prioritisation. *Nucleic Acids Res* **49**, D1302–D1310 (2021).

8.  Ochoa, D. *et al*. The next-generation Open Targets Platform: reimagined, redesigned, rebuilt. *Nucleic Acids Res* **51**, D1353–D1359 (2023).

9.  Slowikowski, K., Hu, X. & Raychaudhuri, S. SNPsea: an algorithm to identify cell types, tissues and pathways affected by risk loci. *Bioinformatics* **30**, 2496–2497 (2014).

10. Skene, N. G. & Grant, S. G. N. Identification of Vulnerable Cell Types in Major Brain Disorders Using Single Cell Transcriptomes and Expression Weighted Cell Type Enrichment. *Front Neurosci* **10**, (2016).

11. Genetic effects on gene expression across human tissues. *Nature* **550**, 204–213 (2017).

12. Ongen, H. *et al*. Estimating the causal tissues for complex traits and diseases. *Nat Genet* **49**, 1676–1683 (2017).

13. Calderon, D. *et al*. Inferring Relevant Cell Types for Complex Traits by Using Single-Cell Gene Expression. *The American Journal of Human Genetics* **101**, 686–699 (2017).

14. Finucane, H. K. *et al*. Heritability enrichment of specifically expressed genes identifies disease-relevant tissues and cell types. *Nat Genet* **50**, 621–629 (2018).

15. Skene, N. G. *et al*. Genetic identification of brain cell types underlying schizophrenia. *Nat Genet* **50**, 825–833 (2018).

16. Bryois, J. *et al.* Genetic identification of cell types underlying brain complex
traits yields insights into the etiology of Parkinson's disease. *Nat Genet* **52**,
482–493 (2020).
17. Jagadeesh, K. A. *et al.* Identifying disease-critical cell types and cellular
processes by integrating single-cell RNA-sequencing and human genetics. *Nat*
*Genet* **54**, 1479–1492 (2022).
18. Zhang, M. J. *et al.* Polygenic enrichment distinguishes disease associations of
individual cells in single-cell RNA-seq data. *Nat Genet* **54**, 1572–1580 (2022).
19. Jia, P., Hu, R., Yan, F., Dai, Y. & Zhao, Z. scGWAS: landscape of trait-cell type
associations by integrating single-cell transcriptomics-wide and genome-wide
association studies. *Genome Biol* **23**, 220 (2022).
20. Xiao, Y. *et al.* An analytical framework for decoding cell type-specific genetic
variation of gene regulation. *Nat Commun* **14**, 3884 (2023).
21. Duncan, L. E. *et al.* Mapping the cellular etiology of schizophrenia and complex
brain phenotypes. *Nat Neurosci* **28**, 248–258 (2025).
22. Song, L., Chen, W., Hou, J., Guo, M. & Yang, J. Spatially resolved mapping of
cells associated with human complex traits. *Nature* (2025)
doi:10.1038/s41586-025-08757-x.
23. Asp, M. *et al.* A Spatiotemporal Organ-Wide Gene Expression and Cell Atlas of
the Developing Human Heart. *Cell* **179**, 1647-1660.e19 (2019).
24. Maynard, K. R. *et al.* Transcriptome-scale spatial gene expression in the
human dorsolateral prefrontal cortex. *Nat Neurosci* **24**, 425–436 (2021).
25. Fawkner-Corbett, D. *et al.* Spatiotemporal analysis of human intestinal
development at single-cell resolution. *Cell* **184**, 810-826.e23 (2021).
26. Ferreira, R. M. *et al.* Integration of spatial and single-cell transcriptomics
localizes epithelial cell–immune cross-talk in kidney injury. *JCI Insight* **6**,
(2021).
27. Hildebrandt, F. *et al.* Spatial Transcriptomics to define transcriptional patterns
of zonation and structural components in the mouse liver. *Nat Commun* **12**,
7046 (2021).
28. Guilliams, M. *et al.* Spatial proteogenomics reveals distinct and evolutionarily
conserved hepatic macrophage niches. *Cell* **185**, 379-396.e38 (2022).
29. Parigi, S. M. *et al.* The spatial transcriptomic landscape of the healing mouse
intestine following damage. *Nat Commun* **13**, 828 (2022).
30. Kuppe, C. *et al.* Spatial multi-omic map of human myocardial infarction.
*Nature* **608**, 766–777 (2022).
31. Kleshchevnikov, V. *et al.* Cell2location maps fine-grained cell types in spatial
transcriptomics. *Nat Biotechnol* **40**, 661–671 (2022).
32. Mirzazadeh, R. *et al.* Spatially resolved transcriptomic profiling of degraded
and challenging fresh frozen samples. *Nat Commun* **14**, 509 (2023).
33. Madissoon, E. *et al.* A spatially resolved atlas of the human lung characterizes a
gland-associated immune niche. *Nat Genet* **55**, 66–77 (2023).
34. Olaniru, O. E. *et al.* Single-cell transcriptomic and spatial landscapes of the
developing human pancreas. *Cell Metab* **35**, 184-199.e5 (2023).

35.  Li, X. *et al.* Profiling spatiotemporal gene expression of the developing human
spinal cord and implications for ependymoma origin. *Nat Neurosci* **26**, 891–
901 (2023).
36.  Medina-Leyte, D. J. *et al.* Endothelial Dysfunction, Inflammation and
Coronary Artery Disease: Potential Biomarkers and Promising Therapeutical
Approaches. *Int J Mol Sci* **22**, 3850 (2021).
37.  Fogo, A. B. & Harris, R. C. Crosstalk between glomeruli and tubules. *Nat Rev*
*Nephrol* **21**, 189–199 (2025).
38.  Zhou, Y.-W. *et al.* Crohn's disease as the intestinal manifestation of pan-
lymphatic dysfunction: An exploratory proposal based on basic and clinical
data. *World J Gastroenterol* **30**, 34–49 (2024).
39.  Aousji, O. *et al.* Patterns of synaptic loss in human amyotrophic lateral
sclerosis spinal cord: a clinicopathological study. *Acta Neuropathol Commun*
**11**, 120 (2023).
40.  Johnson, M. B. *et al.* Type 1 diabetes can present before the age of 6 months
and is characterised by autoimmunity and rapid loss of beta cells. *Diabetologia*
**63**, 2605–2615 (2020).
41.  Bruggeman, B. S. & Schatz, D. A. Type 1 Diabetes: A Disorder of the Exocrine
and Endocrine Pancreas. *J Cell Immunol* **5**, 120–126 (2023).
42.  Crossley, N. A., Scott, J., Ellison-Wright, I. & Mechelli, A. Neuroimaging
distinction between neurological and psychiatric disorders. *British Journal of*
*Psychiatry* **207**, 429–434 (2015).
43.  Čarna, M. *et al.* Pathogenesis of Alzheimer's disease: Involvement of the
choroid plexus. *Alzheimer's & Dementia* **19**, 3537–3554 (2023).
44.  van de Mortel, L. A., Thomas, R. M. & van Wingen, G. A. Grey Matter Loss at
Different Stages of Cognitive Decline: A Role for the Thalamus in Developing
Alzheimer's Disease. *Journal of Alzheimer's Disease* **83**, 705–720 (2021).
45.  Kao, Y.-H., Chou, M.-C., Chen, C.-H. & Yang, Y.-H. White Matter Changes in
Patients with Alzheimer's Disease and Associated Factors. *J Clin Med* **8**, 167
(2019).
46.  Anttila, V. *et al.* Analysis of shared heritability in common disorders of the
brain. *Science (1979)* **360**, (2018).
47.  Velmeshev, D. *et al.* Single-cell genomics identifies cell type–specific molecular
changes in autism. *Science (1979)* **364**, 685–689 (2019).
48.  Yao, Z. *et al.* A taxonomy of transcriptomic cell types across the isocortex and
hippocampal formation. *Cell* **184**, 3222-3241.e26 (2021).
49.  Siletti, K. *et al.* Transcriptomic diversity of cell types across the adult human
brain. *Science (1979)* **382**, (2023).
50.  Larsson, L., Franzén, L., Ståhl, P. L. & Lundeberg, J. Semla: a versatile toolkit
for spatially resolved transcriptomics analysis and visualization.
*Bioinformatics* **39**, (2023).
51.  Selvarasu, K. *et al.* Reduction of kinesin I heavy chain decreases tau
hyperphosphorylation, aggregation, and memory impairment in Alzheimer's
disease and tauopathy models. *Front Mol Biosci* **9**, (2022).

52.    Zhou, J. *et al.* BACE1 regulates expression of Clusterin in astrocytes for
860           enhancing clearance of β-amyloid peptides. *Mol Neurodegener* **18**, 31 (2023).
861    53.    Wulff, H., Castle, N. A. & Pardo, L. A. Voltage-gated potassium channels as
862           therapeutic targets. *Nat Rev Drug Discov* **8**, 982–1001 (2009).
863    54.    Clifton, N. E. *et al.* Developmental Profile of Psychiatric Risk Associated With
864           Voltage-Gated Cation Channel Activity. *Biol Psychiatry* **90**, 399–408 (2021).
865    55.    Um, K. B., Kwak, S., Cheon, S.-H., Kim, J. & Hwang, S.-K. AST-001 Improves
866           Social Deficits and Restores Dopamine Neuron Activity in a Mouse Model of
867           Autism. *Biomedicines* **11**, 3283 (2023).
868    56.    Eltokhi, A. *et al.* Pathogenic gating pore current conducted by autism-related
869           mutations in the Na V 1.2 brain sodium channel. *Proceedings of the National*
870           *Academy of Sciences* **121**, (2024).
871    57.    Harrison, P. J., Tunbridge, E. M., Dolphin, A. C. & Hall, J. Voltage-gated
872           calcium channel blockers for psychiatric disorders: genomic reappraisal. *The*
873           *British Journal of Psychiatry* **216**, 250–253 (2020).
874    58.    Kumar, S. *et al.* Extent of Dorsolateral Prefrontal Cortex Plasticity and Its
875           Association With Working Memory in Patients With Alzheimer Disease. *JAMA*
876           *Psychiatry* **74**, 1266 (2017).
877    59.    Kim, K. *et al.* Therapeutic B-cell depletion reverses progression of Alzheimer's
878           disease. *Nat Commun* **12**, 2185 (2021).
879    60.    Huang, Z. *et al.* Blood-brain barrier integrity in the pathogenesis of
880           Alzheimer's disease. *Front Neuroendocrinol* **59**, 100857 (2020).
881    61.    Choi, J. D. *et al.* Choroid Plexus Volume and Permeability at Brain MRI within
882           the Alzheimer Disease Clinical Spectrum. *Radiology* **304**, 635–645 (2022).
883    62.    Delvenne, A. *et al.* Involvement of the choroid plexus in Alzheimer's disease
884           pathophysiology: findings from mouse and human proteomic studies. *Fluids*
885           *Barriers CNS* **21**, 58 (2024).
886    63.    Bergenståhle, J., Larsson, L. & Lundeberg, J. Seamless integration of image
887           and molecular analysis for spatial transcriptomics workflows. *BMC Genomics*
888           **21**, 482 (2020).
889    64.    Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**,
890           3573-3587.e29 (2021).
891    65.    Lein, E. S. *et al.* Genome-wide atlas of gene expression in the adult mouse
892           brain. *Nature* **445**, 168–176 (2007).
893    66.    Sjöstedt, E. *et al.* An atlas of the protein-coding genes in the human, pig, and
894           mouse brain. *Science (1979)* **367**, (2020).
895    67.    Durinck, S., Spellman, P. T., Birney, E. & Huber, W. Mapping identifiers for the
896           integration of genomic datasets with the R/Bioconductor package biomaRt.
897           *Nat Protoc* **4**, 1184–1191 (2009).
898    68.    Storey, J. D. & Tibshirani, R. Statistical significance for genomewide studies.
899           *Proceedings of the National Academy of Sciences* **100**, 9440–9445 (2003).
900    69.    Storey, J. D., Taylor, J. E. & Siegmund, D. Strong Control, Conservative Point
901           Estimation and Simultaneous Conservative Consistency of False Discovery

902          Rates: A Unified Approach. *J R Stat Soc Series B Stat Methodol* **66**, 187–205
903          (2004).
904   70.   Gu, Z., Eils, R. & Schlesner, M. Complex heatmaps reveal patterns and
905          correlations in multidimensional genomic data. *Bioinformatics* **32**, 2847–
906          2849 (2016).
907