The Evolutionary Landscape of tRNA Modifications in Archaea: Insights from High-Throughput Sequencing

Jesse S. Leavitt[1], Henry Moore[1], Thomas J. Santangelo[2] and Todd M. Lowe[1]

[1]Department of Biomolecular Engineering, Baskin School of Engineering, University of California Santa Cruz, Santa Cruz, CA 95064, USA. [2]Department of Biochemistry and Molecular Biology, Colorado State University, Fort Collins, CO, 80523, USA.

Abstract

Transfer RNA (tRNA) modifications play essential roles in structure, stability and decoding accuracy, yet the evolutionary dynamics and diversity of these modifications remain incompletely understood. Recent advances in high-throughput sequencing, including Ordered Two-Template Relay sequencing (OTTR-seq), now facilitate systematic, transcriptome-wide detection of tRNA modifications at single-base resolution. Here we employed OTTR-seq to comprehensively profile tRNA modifications across nine archaeal species spanning diverse ecological niches, including thermophiles, methanogens, acidophiles, and halophiles. We identified coordinated and mutually exclusive methylation patterns at acceptor stem position 6 and 67 in hyperthermophiles, as well as unexpected clade-specific co-modifications at core positions 10 and 26, demonstrating tolerance of canonical anti-determinants in Thermoprotei. Comparative genomic analyses revealed clear associations between modifications and evolutionary changes in enzyme domain architectures, including divergence in homologs of Trm14, Trm10, Trm11, and Trm1. We also expanded upon previously characterized identity elements in the D-stem, such as the conserved G10oU25 pair critical for Trm11 specificity, providing additional insight into determinants that likely govern enzyme-substrate interactions. Together, these findings offer valuable insights into archaeal tRNA modification biology and evolutionary dynamics, and provide a foundation for guiding targeted biochemical experiments and mechanistic studies.

Keywords

Archaea, reverse-transcription, misincorporation, tRNA modification, enzyme-substrate coevolution

Introduction

Transfer RNAs (tRNAs) are central to translating genetic information into functional proteins in all living systems. Post-transcriptional modifications heavily influence tRNA structure and function, modulating the rate of maturation, molecular stability, and the specificity of interactions with aminoacyl-tRNA synthetases, mRNA codons, and a multitude of other cellular components (Grosjean, 2010; Grosjean et al., 2008; Hernandez-Alias et al., 2022; Schultz & Kothe, 2024; Sokołowski et al., 2018; Yared et al., 2024). These modifications are differentially conserved at distinct positions to impart specific properties (Höfer & Jäschke, 2018; Ontiveros et al., 2019). In Archaea, the complete patterns of tRNA modifications are known for only two archaeal species due to the difficulty and low-throughput nature of traditional biochemical characterization methods. From the comprehensive mapping in *Haloferax volcanii* and *Methanocaldococcus jannaschii* (Gupta, 1984; Yu et al., 2019)*,* along with partial mapping in *Methanococcus maripaludis*, *Pyrococcus furiosus*, and *Sulfolobus acidocaldarius* (Wolff et al., 2023), it is clear that tRNA modifications contribute to adaptation in diverse extreme environments, such as high temperatures or low pH, where these organisms reside (Fluke et al., 2024; Hori et al., 2018; Kowalak et al., 1994; Li et al., 2019; Lorenz et al., 2017; Ohira & Suzuki, 2024; Sas-Chen et al., 2020; Shigi, 2018; Tsai et al., 2025).

The archaeal species in this study span diverse ecological environments: *P. furiosus* thrives at temperatures approaching 100°C in deep-sea hydrothermal vents (Schicho et al., 1993), *S. acidocaldarius* is adapted to acid hot springs with pH values near 2 (Reno et al., 2009), and *H. volcanii* grows in hypersaline environments with salt concentrations exceeding 3M NaCl (Gupta, 1984). Even within methanogens like *M. maripaludis* and *M. jannaschii*, adaptation to anoxic and high-pressure marine environments is evident (Whitman et al., 1987). The chemical stability and structural plasticity conferred by tRNA modifications likely play roles in enabling translation to proceed under these diverse and often extreme physicochemical conditions.

High-throughput tRNA sequencing has become a powerful tool for quantifying tRNA abundance and expression dynamics, complementing data from traditional RNA-seq approaches that often underrepresent structured or modified RNAs. In addition to profiling tRNA expression, specialized tRNA-seq methods now enable the detection of chemical modifications by leveraging the tendency of certain base modifications to disrupt Watson-Crick (W-C) base pairing during reverse transcription (RT), which can be captured using various methods (Behrens et al., 2021; Cozen et al., 2015; Erber et al., 2020; Gustafsson et al., 2022; Kimura et al., 2020; Nakano et al., 2025; Upton et al., 2021; Zheng et al., 2015). While mass spectrometry remains the gold standard for direct chemical identification of modified nucleosides, RT-based sequencing offers a complementary high-throughput, nucleotide-resolution approach.

One of the newest RT-based tRNA seq methods, Ordered Two Template Relay sequencing (OTTR-seq), enhances detection of modification-sensitive misincorporation events while maintaining full-length tRNA coverage (Upton et al., 2021; K. Zhang et al., 2025). OTTR-seq uses a processive reverse transcriptase with a defined template-switching mechanism, improving the uniformity of read coverage across structured RNAs. This makes OTTR-seq

particularly well-suited for transcriptome-wide profiling of tRNA modifications, where detecting

both conserved and context-dependent modification events requires consistent capture of intact,

full-length tRNAs. In archaeal species, where tRNAs are highly structured and often densely

modified, this approach enables a comprehensive view of the modification landscape, revealing

relationships between modification frequency, tRNA sequence, and enzyme specificity.

Much like individual aminoacyl-tRNA synthetases (aaRS) recognize and charge subsets

of tRNAs with a specific amino acid (tRNA isoacceptors) based on distinct sequence and

structural identity elements (Westhof et al., 2022), tRNA modification enzymes recognize

specific sequence motifs and structural elements shared by a subset of tRNAs. Some

modifications, such as N2-methylguanosine (m^2G) or N2,N2-dimethylguansine (m^2_2G) at

position 10, are important early in pre-tRNA maturation, mediating correct folding of the

L-shape tertiary structure (Hori, 2019; Shigi et al., 2006). Transfer RNA modification pathways

are interconnected and show cooperativity with one another, influencing stress response and

disease through modulation of tRNA stability and decoding behavior (Hernandez-Alias et al.,

2022; Sokołowski et al., 2018). To add to this complexity, the tRNA modification enzymes'

precise target specificity evolve across diverse phyla, causing uncertainty or inconsistencies

between studies on the same family of modification enzymes in different model systems

(de Crécy-Lagard et al., 2019).

Of the many tRNA modification types, methylation is the most common, with at least 13

different chemical variations of methylated RNA bases shared among all three domains of life

(Hori, 2014; Jackman & Alfonzo, 2013; Motorin & Helm, 2011). Despite this overlap,

modification enzymes vary widely in their protein domain architectures and substrate

specificities across the tree of life. While some, like TrmI, function as multi-subunit complexes

in eukaryotes but as a homotetramer in Archaea (Roovers et al., 2021), others show variability in domain architectures that influence tRNA recognition (Dixit et al., 2019). These differences impact how modification enzymes selectively recognize and act on diverse tRNA sequences, yet the extent of this variation and its functional consequences remain poorly understood. Investigating these evolutionary differences will provide insight into the determinants of tRNA modification specificity.

Here, we elucidate the landscape of RT-detectable tRNA modifications by applying OTTR-seq across nine archaeal species, including three thermophiles, two methanogens, two acidophiles, and two halophiles, to better understand the dynamic rate of tRNA modification evolution across different environments. In particular, we focus on position-specific modifications that vary between clades, isoacceptor-specific tRNAs, or a combination of the two. Furthermore, we investigate the distribution of corresponding tRNA modification enzyme orthologs and provide detailed co-evolutionary analyses of protein domain architectures with respect to variation in tRNA substrate target sites. Altogether, we find substantial evidence for conservation patterns for m^2G/m^2_2G at position 6, m^1A/m^1G at position 9, m^2_2G/xG at position 10 and 26, which co-vary with the corresponding modification enzymes in these species, making prediction of the presence or absence of these modifications feasible from genome sequence alone.

Results

<u>Detection of base-specific misincorporations reveals tRNA modification dynamics across archaea</u>

We profiled post-transcriptional tRNA modifications across nine archaeal species (Table 1) using OTTR-seq (Upton et al., 2021) to identify reverse transcription (RT)-induced misincorporation signatures of tRNA modifications (Figure 1). Known modification sites in six of these species (*H. volcanii*, *M. jannaschii*, *M. maripaludis*, *P. furiosus, S. acidocaldarius,* and *T. kodakarensis*) (Gupta, 1984; Hirata et al., 2019; Wolff et al., 2023; Yu et al., 2019) provided a validated reference for evaluating modification detection (Additional file: Table S0), while additional comparisons with *Thermococcus sp.* AM4 and *Sulfolobus islandicus* enabled assessment of short-term evolutionary conservation (Figure 1A). We predicted sites of tRNA modification via prominent and statistically significant misincorporation frequencies (MI frequency), defined as the proportion of sequencing reads at a given nucleotide position that differ from the reference base. High misincorporation frequencies define the extent to which a modification disrupts incorporation of the cognate nucleotide during reverse transcription, typically by disrupting Watson Crick (W-C) base pairing. The misincorporation frequency provides a proxy for the presence and stoichiometry of modification events.

To evaluate detection confidence, we implemented base-specific MI frequency thresholds that account for spontaneous misincorporation uniquely induced by each canonical base. Guanine and cytosine exhibited elevated background misincorporation rates, even in the absence of annotated modifications, which required adjusted criteria to reduce false positives. Sites of

potential modification within tRNAs were classified as high-confidence if MI frequency

exceeded 10% for G/C, while lower spontaneous MI frequencies allowed a tighter threshold of

5% for A/U. All high-confidence modification sites must also have statistical significance

(adjusted $p < 0.05$). Positions with significant p-values but misincorporation rates below 10% for

G/C or 5% of A/U were designated as moderate-confidence misincorporation sites. The

distribution of high- and moderate-confidently detected modification is summarized by position

(Figure 1B) and across all tRNAs (Additional file: Fig. S1 and Fig. S2, Table S1).

We observed that select tRNA modifications displayed a range of misincorporation

behaviors. Modifications disrupting the W-C face (e.g., m^1A, m^1I, m^2G, m^2_2G) consistently

showed the highest MI rates (Figure 1C) and deviations from expected patterns revealed

biologically meaningful variation. For example, while m^1A58 typically results in near-complete

misincorporation, Thermoprotei species showed reduced MI rates at A58 in tRNA^{Ala}. Despite

lower MI rates, the read identity (patterns of misincorporation) and read depth supported m^1A

modification at this position, suggesting substoichiometric modifications or tRNA-specific

regulation (Figure 1C, Additional file: Fig. S3 and Fig. S4). Similar observations were made for

A57 in *P. furiosus* Thr^{GGU}, where rates of misincorporation markedly dropped but the patterns of

misincorporation were consistent with m^1I.

Other notable outliers included pseudouridine at position 55, which was consistently

undetectable across species, as expected given its minimal impact on reverse transcription.

However, in *H. salinarum* Ser^{CGA}, we observed an unusually high misincorporation rate and

significant p-value at this position. While pseudouridine alone is unlikely to account for this

signal, it may reflect the presence of a related or modified derivative such as $m^1\Psi$, which has

been reported at nearby positions like 54 but may occur at 55 in this specific context (Additional file: Fig. S4).

An important observation was the characteristic misincorporation behavior of different modification types occurring at the same position and base (Figure 1D, Additional file: Fig. S1). Modifications derived from guanine or adenine that disrupt the W-C face (e.g. m^1A, m^2G, and m^2_2G) tended to exhibit the highest misincorporation rates, making them more readily detectable. The relative distributions of these rates varied by modification type from the same position. In some cases, this allowed us to differentiate between modifications at the same position. For example, at position 10, the misincorporation frequency for m^2_2G was consistently higher than that for m^2G, enabling distinction between these modifications.

Homology-based predictions of tRNA modifications identifies dynamic hotspots

Mapping known modification sites to positions of misincorporations in previously characterized species allowed us to predict their presence in related species lacking experimental annotations. For example, the known modification profile of tRNA[iMet] in *P. furiosus* was mapped to concordant sites of misincorporation in *T. kodakarensis*. In this case, we were able to predict all of the known and detectable modification sites in *T. kodakarensis*, with the exception of 2'-O-methyguanine (Gm) at position 22 (Figure 2A), which has a relatively low misincorporation frequency even in the *P. furiosus* tRNA (Figure 1C). Additional pairwise comparisons across species further supported the predictive power of misincorporation-based inference of site-specific tRNA modifications. Across comparisons, known and predicted misincorporation frequencies of modified homologous tRNAs were highly correlated (Additional

file: Table S2). For species within the same clade, Pearson correlation coefficients ranged from 0.87 to 0.95 (Figure 2B-F, Additional file: Fig S5), and this strong agreement indicates that modified homologous tRNAs frequently share conserved misincorporation patterns, reinforcing the value of using known sites of tRNA modification as predictors for tRNA modification patterns in closely related taxa.

Outliers within these comparisons were particularly informative, revealing partial and variable modification states in homologous tRNAs between species within clades. As an example, (Figure 2B, Additional file: Fig. S6), the known m^2_2G in iMet^{CAU} of *P. furiosus* had a lower and more variable pattern of misincorporation compared to the predicted modification of the tRNA^{iMet} from *T. kodakarensis* at position 26. Interestingly, the predicted modification in *T. kodakarensis* had a larger number of deletions in the patterns of misincorporation. While the triple methylation of guanine (m^2_2Gm) is often associated with deletions and higher rates of misincorporations, we define all likely G-derived modifications at position 26 as "xG" as there are often two or more modification types that occur at this position that can not be consistently distinguished solely through MI frequencies. Similarly, the difference observed at position 57 in Thr^{GGU} shows that the patterns of misincorporation are consistent with m^1I despite variation in rates of misincorporation. In *P. furiosus* vs *T. sp.* AM4 (Figure 2C), we identified additional outliers of known m^2_2G in iMet^{CAU} at position 6, known to have either m^2G or m^2_2G. While *T. sp.* AM4 shows low rates of misincorporation that are often associated with m^2G, we define all predicted G-derived modifications at this position as m^2G/m^2_2G. The difference observed at position 34 in Arg^{UCU} shows that the pattern of misincorporation is consistent with cnm^5U despite being at the lower limits of detection (~1% MI). Comparisons between *H. volcanii* and *H. salinarum* (Figure 2D) revealed several divergent misincorporation levels. The low rate and read

identity of known m^2_2G in Lys^{UUU} in *H. volcanii* supports presence of m^2G, which is consistent

with the known m^2G in Leu^{UAA}. The predicted modifications in *H. salinarum* are consistent with

the high rates of misincorporation often associated with m^2_2G. Due to the differences seen in

these known modifications, we predict all G-derived modifications at position 26 as m^2G/m^2_2G

or xG. Additionally, modifications at position 37, such as t^6A in Lys^{CUU} and Lys^{UUU} showed

variations in the predictions, reflecting either partial modification or differences in

species-specific regulation. In *S. acidocaldarius* vs *S. islandicus* (Figure 2E), position 26

modifications in Glu^{CUC}, Phe^{GAA}, and $iMet^{CAU}$ stood out as underrepresented in *S. islandicus*.

Despite these differences, the read identity patterns of predicted modifications were consistent

with the respective known modifications. These variations may suggest species-specific

differences in the regulation of these modifications. Finally, in *M. jannaschii* vs *M. maripaludis*

(Figure 2F), position 26 in $iMet^{CAU}$ was again identified as an outlier. In this case, the

hyperthermophile (*M. jannaschii*) has identity patterns consistent with m^2_2Gm, whereas the

mesophile (*M. maripaludis*) has patterns consistent with m^2G.

Together, conserved modifications at positions like 6, 26, and 37 appear to be hotspots for

interspecies variation in stoichiometry or type. These comparisons highlight how deviations from

expected misincorporation frequencies can predict species-specific differences in modification

state. Moreover, read identity patterns support the biological plausibility of these differences, as

distinct patterns of misincorporation are often consistent with those of known modifications. Our

observations emphasize the caution required in homology based modification predictions, where

careful contextual interpretation is required. For our downstream analyses, we focus on

modifications that have moderate to high confidence of detection, and use xN notation at

positions where the exact modification type is unable to be inferred.

Clade-specific patterns reveal coordinated modification in the acceptor stem and core regions

Distinct modification patterns in the acceptor stem and tRNA core regions reveal coordinated signatures that vary by clade. In particular, m^2G and m^2_2G modifications at positions 6 and 67 (base pairing in the acceptor stem) are frequently predicted in hyperthermophilic species (Figure 3A). While prior studies have reported individual modifications at these positions in select tRNAs from *M. jannaschii* and *P. furiosus* (Wolff et al., 2023; Yu et al., 2019), the broader pattern of coordination has not been previously recognized. Our expanded analysis reveals that across multiple species, position 6 is more frequently modified than position 67, with misincorporation rates suggesting a predominance of m^2_2G6 over m^2G6. In contrast, m^2G67 is less frequent but consistently observed in distinct tRNA subsets (Additional file: Fig. S2 and Table S3). The two modifications are not predicted to co-occur within the same tRNA (Figure 3B), indicating a likely mutually exclusive pattern of modification. This trend is particularly evident in all Thermococci species and *M. jannaschii* (Methanococci). Most G6 conserved tRNAs are predicted to have a modification across these species (Additional file: Fig. S2 and Table S3). However, certain tRNAs in Thermococci, such as Glu^{UUG} and Ile^{GAU}, possess a C6, precluding the possibility of m^2G or m^2_2G at this position in these cases.

Coordination of modification patterns is also observed between positions 10 and 26, two highly conserved sites of modification across archaeal tRNAs. While the predicted modifications conserved in species from Halobacteria and Methanococci tend to occur in mutually exclusive subsets of tRNAs, co-occurrence of modifications at both sites within the same tRNA is found in Thermococci and Thermoprotei (Additional file: Fig. S7 and Table S4). This pattern is especially pronounced in Thermoprotei, where multiple tRNAs such as $tRNA^{Asp}$, $tRNA^{Glu}$, and $tRNA^{Val}$ are

predicted to carry both m^2G/m^2_2G10 and m^2_2G26 (Additional file: Fig. S2, S7 and Table S4). In contrast, while Thermococci also exhibits co-modification of position 10 and 26, their modification types differ. With lower rates of misincorporation at position 10, their tRNAs may instead carry m^2G10 and m^2_2G26 (Additional file: Fig. S7). These differences are not entirely surprising as these specific clades span two phylums (Crenacrhaota and Euryarchaota). Their preferences in patterns of modifications suggest evolutionary divergence in enzymatic activity and substrate selection.

Within Euryarchaeota, coordinated patterns of G10 and G26 modification appear to be more closely linked to specific environmental variables such as temperature. Predicted G10 substrates typically contain G10oU25 identity elements, consistent with previous observations (Urbonavičius et al., 2006), while G26 modifications tend to occur in tRNAs lacking this pairing (Additional file: Fig. S8, Table S6). Interestingly, in Thermoprotei, several tRNAs predicted to be modified at both positions contain G10oU25 but lack the D-arm motifs (e.g., U13oU22, U13oA22) commonly associated with m^2G10/m^2_2G10 in other clades. These findings suggest that the coordination of modifications at structurally proximal positions may reflect shared structural constraints in mesophilic species within Euryarchaeota, and the clade-specific tuning of enzyme-substrate interactions in thermoacidophiles within Crenarchota.

Conserved and clade-specific tRNA modifications at functional positions support distinct evolutionary adaptations

While several tRNA modifications have been proposed to support adaptation to high-temperature environments, our results reveal additional patterns that suggest clade-specific evolutionary pressures beyond thermophily alone. A notable example is the methylation of

purines at position 9, where we find clear distinctions in modification type and substrate specificity between Thermoprotei and Thermococci, which is consistent with previous work (Krishnamohan & Jackman, 2019). In Thermoprotei, predicted modifications are limited to m^1A9, affecting most A9-containing tRNAs (Additional file: Fig. S2). By contrast, Thermococci species display a broader substrate range, with both m^1A9 and m^1G9 modifications predicted across a more diverse set of tRNAs (Additional file: Fig. S2). These differences coincide with distinct patterns of sequence conservation at position 9, such as the rare occurrence of C9 in specific tRNAs of Thermoprotei, and reflect lineage-specific recognition preferences of Trm10 homologs. Since species from both of these clades thrive in elevated temperatures, the variable targets for these tRNA modifications at conserved positions can be shaped by clade-specific substrate preferences and enzyme evolution, rather than by thermophilic adaptation alone.

Consistent with these patterns, we detected clear misincorporation signatures at position 9 in both clades, supporting the presence of m^1A and m^1G modifications in tRNA- and clade-specific manner (Figure 3C). Although Thermococci and Thermoprotei have a comparable number of clade-conserved m^1A9-modified tRNAs predicted (n=18 and n=19, respectively), the individual tRNAs targeted differ between clades (Additional file: Fig. S2). For example, m^1A9 is uniquely predicted in Pro^{GGG} in Thermoprotei, whereas in Thermococci m^1A9 uniquely occurs in Cys^{GCA} and Ser^{GCU}. G9-containing tRNAs are modified only in Thermococci (n=6), while Thermoprotei tRNAs with G9 remain unmodified. Additionally, Thermoprotei harbor rare C9-containing tRNAs, such as in Ser^{GCU} and tRNA^{Val}, whereas Thermococci harbor A9/G9 in their respective tRNAs, further underscoring clade-specific variation in tRNA sequence contexts. Predicted modifications at position 9 appear restricted to these two thermophilic clades. No evidence for m^1A9 and m^1G9 was found in Halobacteria or Methanococci, which lack detectable

Trm10 homologs (Figure 4A). Together, these patterns highlight a narrow conservation of position 9 modifications.

In addition to the clade-specific patterns observed at A9/G9, several other positions, such as G10 and G26, are modified across archaeal species, although their distribution and frequency vary by lineage and environmental niche (Figure 3C). Modifications at position 10, predicted as m^2G or m^2_2G (xG), are present in all clades but are conserved with greater frequency between species in Thermococci (n=17) and Thermoprotei (n=15). Previous studies have associated m^2_2G10 with high-temperature growth in *T. kodakarensis* (Hirata et al., 2019; Hori, 2023), and our data suggest that the presence of m^2_2G10 does correspond with optimal growth temperatures, although it is not exclusive to thermophiles. For instance, these modifications are also frequent in mesophilic Halobacteria (n=13) and Methanococci species *M. maripaludis* (n=10). Interestingly, tRNA[Pro] in Thermoprotei lacks a predicted modification at position 10 despite containing U25, a context known to promote m^2_2G10 installation for stabilization of the G10oU25 wobble pair (Urbonavičius et al., 2006). This absence suggests that the positioning and frequency of this modification may reflect lineage-specific adaptations that are not necessarily dependent on temperature alone.

Position 26 shows even border conservation, with nearly all species exhibiting high counts of predicted $m^2G/m^2_2G/m^2_2Gm$ (xG) at this site (Figure 3C). Thermoprotei have the largest number of predicted modifications at this position (n=39), followed by Thermococci (n=29), Halobacteria (n=22), and Methanococci (n=15). While m^2_2G and its hyper-modified variant m^2_2Gm have been associated with thermophilic species (Wolff et al., 2023), our results suggest that modification frequency at this position does not correlate strictly with optimal growth temperatures of these species. For example, acidophilic species exhibit high counts of

predicted G26 modification, exceeding those observed in hyperthermophiles. Moreover, although the predicted number of conserved G26 is lowest in Methanococci, the hyperthermophile *M. jannaschii* independently shows broader modification patterns (n=21) (Additional file: Fig. S2). Importantly, across species, not all G26 tRNAs are predicted to be modified. These trends suggest that while G26 is conserved and frequently modified, the extent and specificity of its modification vary by clade, reflecting differences in tRNAs and the substrate recognition properties of Trm11 homologs.

Position 37, located adjacent to the anticodon, is another highly conserved site of modification, with distinct patterns of chemical diversity observed across archaeal clades (de Crecy-Lagard et al., 2010). Predicted modifications at this site fall into two main categories: xA37, which includes t^6A-like derivatives, and xG37, which primarily includes m^1G as well as wyosine-like derivatives. Our analysis shows that xA37 modifications are primarily restricted to Halobacteria, where they are predicted in 10 tRNAs conserved between the two species, with only a single conserved prediction detected in Thermoprotei (Figure 3C). This prediction corresponds to Ser^{GCU}, an expected target across all archaeal species in this work (Additional file: Fig. S2). In addition, it is broadly expected in tRNAs that recognize codons beginning with adenine-ANN codons (W. Zhang & Westhof, 2025). However, we did not detect this highly conserved modification with confidence in Ser^{GCU} from *T. kodakarensis* or *P. furiosus*, likely due to the low misincorporation signal typically associated with t^6A. In contrast, xG37 modifications are more broadly distributed and most prevalent in Thermococci, with up to 22 tRNA conserved between species, compared to lower counts in other clades (Figure 3C). These patterns highlight clade-specific distributions of distinct modifications at conserved sites and reflect both evolutionary divergence and technical limitations in detecting certain modification types.

Conserved modifications continue into the T-arm, particularly at position 57 and 58, sites known to contribute to tRNA structure and function across all domains of life (Anderson & Droogmans, 2005; Oerum et al., 2017; Swinehart & Jackman, 2015). Thermophilic and hyperthermophilic species frequently carry both m^1I57 and m^1A58 (Figure 3C), consistent with their proposed role in structural stabilization. In our analysis, m^1I57 is among the most frequently predicted modifications, occuring in 48-65% of tRNAs per species (16/35 to 31/46). m^1A58 is even more widespread, detected in 77-98% of tRNAs (27/35 to 45/46), with one notable exception: Halobacteria lack any predictions of m^1A58 (n=0). This clade-specific absence contrasts with its near-universal presence in other archaeal groups, suggesting divergence of TrmI homologs which typically methylates both A57 and A58 (Roovers et al., 2021). These trends reinforce the high degree of conservation for modifications at structurally significant positions in the tRNA, while also revealing lineage-specific absence.

Phylogenetic variation in tRNA methyltransferase architectures reveals enzyme specialization across Archaea

Methylation is one of the most common post-transcriptional modifications, found across all domains of life. The methyl donor used by most methyltransferases (MTases) is S-adenosyl-L-methionine (AdoMet), acting as their co-substrate. tRNA modification enzyme homolog distributions provide supporting evidence of methyltransferases having specialized roles that drive clade specific adaptations. The AdoMet-dependent enzymes are divided into five different classes based on their structural folds (I-V) (Bujnicki et al., 2002). The enzymes investigated in this work include three class I MTases (Trm14, Trm11, Trm1) and a class IV SPOUT MTase (Trm10). The phylogenetic relationships and predicted domain architectures of

these enzymes across representative archaeal species suggest these enzymes have evolved independently towards different specificities (Figure 4). Each panel presents a phylogenetic tree depicting the presence, location, and range of predicted functional domains, alongside heatmaps summarizing the confidence of predicted substrates across species.

The THUMP-related Trm14 (m^2G6/m^2_2G6) is distributed exclusively in hyperthermophiles, within the Methanococci and Thermococci clades. Consistent with known archaeal crystal structures (Fislage et al., 2012; Menezes et al., 2011), predicted Trm14 homologs maintain three conserved domains: an N-terminal ferredoxin-like domain (NFLD), a THUMP RNA-binding domain, and a Rossmann fold MTase domain (RFM) (Figure 4A, Additional file: Table S7). However, subtle variation at the N-terminal regions distinguishes homologs across clades, specifically MJ0438 uniquely contains a hit for a predicted RImL ferredoxin-like domain. Although there are variations in domain architecture among these homologs, differences in predicted substrate profiles indicate tRNA sequence divergence as the main factor influencing modification specificity rather than enzyme domain architecture alone. Interestingly, the enzyme responsible for m^2G67 remains unidentified, with previous work suggesting Trm14 involvement (Hirata et al., 2019). Our homolog search revealed the presence of Trm14 paralogs distributed within these clades (Additional file: Table S11). These paralogs may represent alternative candidates responsible for m^2G67.

Previously, the SPOUT domain modification enzyme Trm10 (m^1G9/m^1A9) was thought to be widely conserved across Archaea (Krishnamohan & Jackman, 2019), however with the availability of additional genomes, distribution of predicted Trm10 homologs is in fact quite narrow, unique only to species within Thermococci and Thermoprotei. Within these species, the characterization of TK0422 (TkTrm10) from *T. kodakarensis* has shown that it has remarkable

dual specificity for N1-methylation of purine residues at position 9 (Krishnamohan & Jackman, 2019). Although SPOUT domains are notoriously challenging to model at the sequence level, InterProScan returned domain hits for features characteristic of a bifunctional tRNA (guanine(9)-/adenine(9)-N1)-methyltransferase from the NCBIfam domain database (Figure 4B, Additional file: Table S8). In contrast, Thermoprotei vary in their domain features that diverge from bifunctional characteristics and instead resemble residues of a tRNA (adenine(9)-N1)-methyltransferase. This interesting divergence between domain architectures suggests a functional split across phylogeny, and is further supported by the evidence of predicted target substrates from misincorporation (Figure 4B).

The broadly conserved THUMP-related modification enzyme Trm11 (m^2G10/m^2_2G10) has distribution across all clades. Based on previous characterizations, TK0981 is expected to contain a NFLD, a THUMP domain, and a RFM, much like Trm14, differing only in the linker region connecting the NFLD and RFM domains (Hirata et al., 2016; Urbonavičius et al., 2006). Homologs in Thermoprotei have the conserved C-terminal AdoMet and RMKL-like domains, but the expected N-terminal THUMP domain is completely absent (Figure 4C; Additional file: Table S9). The absence of this important tRNA binding domain, along with predicted modified tRNA substrates, implies that the Trm11 in Thermoprotei has a modular partner that contributes to the binding of its targets. There is a predicted ortholog of the Trm112 activator protein present in both species, which has previously been shown to form a complex with Trm11 in the model *H. volcanii* (van Tran et al., 2018), but its contributions to binding tRNAs has not been characterized. The mode of binding by these divergent Trm11 homologs is not clear, however the predicted targeted substrates in species with divergent N-terminal regions do co-vary, specifically with tRNA[Pro] (Figure 4C).

One major group of highly conserved tRNA methyltransferases is the Trm1 family (m^2G/m^2_2G26). These MTases are unusual in that they do not possess a THUMP domain (Hori, 2023; Ihsanawati et al., 2008). Despite this, Trm1, similar to Trm11, catalyzes a two-step reaction, leading to mono- and dimethylation of guanosine. The protein comprises an N-terminal and C-terminal domain characteristic of class I MTases and uses AdoMet as a ligand for methylation. Predicted homolog domain architectures for this family broadly identifies the expected (guanine(26)-N(2))-dimethyltransferase (Additional file: Table S10) across all species, with AdoMet detected near the N-terminal (Figure 4D). This differs from Trm14 and Trm11, where AdoMet is typically found near the C-terminal. Surprisingly, InterProScan does not detect the AdoMet domain in all species, it diverges from expected residues used by the domain model (Additional file: Table S10). Despite relative similarity between the predicted domains across species, targeted substrates are drastically different (Figure 4D). In Euryarchaeota, there is a biased coordination between m^2G10/m^2_2G10 and m^2_2G26 (Additional file: Fig. S7), which could imply one is the anti-determinant of the other. In Thermoprotei, both G10 and G26 are predicted to be modified with m^2_2G.

tRNA sequence and structure features reveal determinants and anti-determinants of modification enzyme targeting

Positions with predicted modifications correspond to conserved nucleotide features in specific tRNAs across various species. Likewise, shifts in predicted modifications are linked to changes in nucleotides at distinct positions. Our comprehensive analysis across multiple archeal clades expands upon previous observations, reinforcing and broadening the understanding of identity elements critical for modification enzyme recognition. Specifically, the strongest

association of these nucleotides aligns with previously characterized identity elements for Trm11, requiring G10oU25 pair in the D-arm (Urbonavičius et al., 2006). Our broad sampling across archaeal species supports this identity element, reinforcing its conserved functional importance. Among Euryarchaeota species, the frequency of predicted m^2G10/m^2_2G10 modifications correspond with the expected G10oU25 and U13oU22 (Figure 5A). Moreover, our analysis identifies additional correlations, involving other non-Watson-Crick pairs such as U13oG22 and U13oUA22 in Thermococci and Thermoprotei (Additional file: Fig. S8, Table S5).

Non-Watson-Crick paris, particularly GoU paris, are known to induce structural variations essential for protein binding without necessitating direct protein-base contact (Westhof et al., 2020, 2022). Our extensive cross-clade analysis further supports that these structural variations contribute to enzyme recognition. Structurally, residues forming Non-Watson-Crick pairs result in shorter, less stable D-stems, which often correspond with the presence of predicted m^2_2G10 modifications (Figure 5B & E). Conversely, we observed absences of these modifications in tRNAs possessing more stable D-stems, further supporting the functional role that this modification has in the structural integrity of tRNAs. For instance, within Halobacteria, the tRNA^{Trp} deviates from the conserved G10oU25 pair, instead forming a G10=C25 pair, coinciding with the absence of predicted m^2_2G10 modification (Figure 5C-D). This difference highlights the stringent substrate specificity linked to identity elements like U25, essential for facilitating m^2_2G10-U25 interactions, but not m^2_2G10-C25 interactions.

Similarly, nucleotide features specific to predicted targets of G10oU25 containing tRNAs have been identified as anti-determinants for the action of Trm1, as demonstrated previously in *P. furiosus* (Urbonavičius et al., 2006). Our results support this relationship across Euryarchaeota,

showing a strong association between the presence of G10oU25 and the absence of m²$_2$G26 (Figure 5E). Interestingly, Thermoprotei do not exhibit this exclusion pattern and often show co-modification of m²$_2$G10 and m²$_2$G26, in addition to unique targeting of C12=G23 and A12oU23 tRNAs (Additional file: Fig. S8, Table S6). This divergence highlights structural adaptation strategies that may be unique to Thermoprotei. In general, weak D-arms likely depend on m²$_2$G10 to stabilize the tRNA in its functional L-shaped conformation (Steinberg & Cedergren, 1995). In a specific example of the tRNA^{Trp} of *T. kodakarensis*, we see how m²$_2$G10 may prevent an alternative interaction between G10=C26 and likely plays a role in most m²$_2$G10oU25-containing tRNAs (Figure 5F). Collectively, this multi-clade analysis extends our current knowledge on interplay between tRNA sequence, structure, and modification enzymes.

Discussion

Through interrogating high-throughput sequencing datasets alongside genomic analyses we have expanded the scope of our understanding of post-transcriptional tRNA modifications, uncovering novel relationships, evolutionary implications, and insights into the coevolution of tRNA substrates with their modifying enzymes. Our comprehensive analysis across diverse archaeal species reveals distinct modification patterns reflecting adaptive mechanisms shaped by both environmental pressures and phylogenetic constraints.

Our base-specific misincorporation analysis illustrates how modifications differentially impact reverse transcription behavior, influencing their detectability via RT-based methods. Modifications disrupting W-C base pairing, such as m$^2_2$G and m^{1}A, consistently yield higher misincorporation signals, facilitating robust detection. Conversely, modifications like 2'-*O*-methylations (Am, Cm, Gm, Um) remain challenging to detect due to minimal interference

with reverse transcription or shadow signals resulting from proximity to nearby modifications. These are important examples of methodological limitations that warrant caution when inferring presence or absence solely from low misincorporation signals, particularly at known conserved modification sites such as position 56.

In this study, we predicted xG modifications at several positions, finding surprising relationships, such as the coordination of m^2G/m^2_2G at positions 6 and 67 in the acceptor stem of hyperthermophiles. While previous work noted individual modifications at these positions, our comparative approach clearly indicates their mutual exclusivity, suggesting a previously unrecognized functional interdependence. Despite narrow conservation in hyperthermophiles, these modifications may not play a role in thermal-stability, which typically involves T-loop/D-loop interactions at the outside globular corner of the L-shape (Grosjean & Oshima, 2007). Insights from (Wolff et al., 2023) suggest m^2_2G6-C67 disrupts the W-C base pair, resulting in a pronounced propeller twist, likely playing a role in modulating the aminoacylation efficiency of aaRS and tRNA charging. With our new observations of this coordinated relationship, we hypothesize that as the steric disruptions alter between m^2_2G6-C67 and C6-m^2G67, charging efficiencies may be regulated for specific tRNAs. It's tempting to speculate that this unique mechanism of charging regulation could provide these hyperthermophiles with adaptive strategies to alter protein synthesis rates in response to environmental shifts.

Additionally, we identified coordinated modification patterns involving conserved core positions 10 and 26 across multiple archaeal clades. Our results highlight distinct clade-specific modification preferences, particularly in Thermoprotei, where co-occurrence of m^2_2G10 with m^2_2G26 is surprisingly frequent. While Thermococci share the unique co-modification of G10 and G26, their patterns contrast with lower frequencies of misincorporation that suggest pairing

of m^2G10 with m^2_2G26 is predominant. This difference implies evolutionary divergence in enzyme specificity for substrate recognition strategies, possibly linked to adaptations beyond thermal stability. Prior biochemical work supports G10oU25 pairs as a robust identity element of m^2_2G10 by Trm11, while simultaneously acting as an anti-determinant for Trm1-mediated m^2_2G26 modifications (Urbonavičius et al., 2006). Interestingly, our observations in Thermoprotei challenge this established pattern, revealing frequent coexistence of both modifications despite expected anti-determinant, indicating flexibility in recognition mechanisms or alternative stabilization strategies. Structurally, these species may utilize the unusual co-modification of 10 and 26 to fold the D-arm region and stabilize the three-dimensional core structure. As a general rule, methylations promote precise H-bonded pairs (e.g. m^1A favors Hoogsteen pairs, m^2G/m^2_2G favors GoU pairs) and electrostatic charges introduced by the chemical adducts are localized in shielded pockets of the tRNA fold (m^1A, m^7G, G^+). In addition, methylations on the base and/or the ribose affect the hydration shells in complex ways (Auffinger & Westhof, 2001; Wolff et al., 2023). These intricate interactions between position 10 and 26 may reflect nuanced structural and functional roles, suggesting novel adaptations to otherwise unstable D-arm configurations, and ensuring tRNA stability and proper folding in acidic environments with elevated temperatures.

The broad identification of tRNA modification enzyme homolog distributions provides supporting evidence for methyltransferases having specialized roles that drive clade-specific adaptations. Comparative analyses of enzyme domain architectures clearly indicate independent evolutionary trajectories towards distinct chemistries and substrate specificities. A distinct example of this is the divergence of Trm10, where Thermoccoci possess dual-specific m^1A9/m^1G9 MTase functionality, contrasting Thermoprotei's exclusive m^1A9 specificity.

Another example is seen in Trm11, where predicted domain architectures indicate a complete loss of the THUMP domain, specifically within Thermoprotei. The absence of this critical domain, along with differences in predicted modified tRNA substrates, suggests a potential modular partnership with another protein to facilitate substrate binding. In support of this speculation, both Thermoprotei species analyzed here possess predicted orthologs of the Trm112 activator protein, known from studies in *H. volcanii* to form complexes with Trm11(van Tran et al., 2018). Together, the divergence of Trm11 domain architectures, combined with unique targeting of C12=G23 and A12oU23 tRNAs in Thermoprotei, are illustrative examples of enzyme and substrate coevolution. Additional comparative insights from recent biochemical analysis of the Trm1 homolog in humans (TRMT1) further supports nuance specificities of enzyme-substrate recognition. Human TRMT1 independently catalyzes m^2G or m^2_2G in a substrate-dependent manner, governed by specific base pairs in the D-stem (Xiong et al., 2023). These similar structural codes may apply broadly across domains, potentially reflecting deep evolutionary conservation of modification determinants.

Our broad analysis provides evidence reinforcing and expanding upon previously identified tRNA modification determinants and anti-determinants, as well as revealing cases where these established patterns are challenged, particularly in Thermoprotei. However, other modification sites, such as those predicted to carry m^2G6/m^2_2G6, m^1A9/m^1G9, and m^2G67, do not exhibit strong evidence for associations with specific nucleotide residues within the tRNA body. Previous biochemical studies on Trm10 have suggested that substrate preferences may instead be influenced by differences in the overall stability of the folded tRNA structure. It is possible that for enzymes such as Trm10 and Trm14, substrate recognition relies on interactions with the global L-shaped tertiary structure in combination with precise recognition of the

catalytic site nucleotide (Strassler et al., 2022). This distinction demonstrates the diversity of recognition strategies across modification enzymes, and emphasizes that inferring determinants of modifications for many enzymes likely requires integration of tertiary structural context.

Altogether, this work generated a wealth of high-throughput data that lays the groundwork for further research by contributing detailed information on tRNA modification profiles, substrate preferences, enzyme distributions, and evolutionary patterns across archaeal lineages. We present a comprehensive experimental analysis of archaeal tRNA modifications and associated enzymes, enhancing our understanding of their adaptive roles. These results provide a valuable foundation for guiding targeted biochemical validation experiments, structural studies of enzyme-substrate interactions, and development of mechanistic models that explain modification specificity and coordination. Future research should focus on characterizing the mechanisms responsible for these modifications, exploring their regulatory functions in translation, and leveraging archaeal strategies for functional applications.

Conclusions

This study provides a comprehensive experimental analysis of archaeal tRNA modifications and the enzymes predicted to install them. By combining high-throughput sequencing with comparative genomic analysis, we uncovered coordinated and clade-specific modification patterns at key tRNA positions (i.e. 6, 9, 10, and 26) and linked these patterns to evolutionary changes in enzyme specificity and domain architecture. Our findings extend the known repertoire of identity elements and anti-determinants of Trm11 and Trm1 homologs, while also revealing instances where recognition rules are challenged, particularly within Thermoprotei. These results emphasize co-evolution between tRNA structure, modifications, and

their enzymes in the context of extreme environmental adaptation. Beyond mapping, this work establishes a framework for future biochemical validation, predictive modeling, and functional studies of tRNA modifications. Ultimately, the insights gained here advance our understanding of tRNA biology in Archaea and provide new opportunities for leveraging tRNA modifications for functional applications.

Acknowledgments

We are grateful to members of the NSF-funded archaeal epitranscriptomics consortium for providing the RNA samples of many species. For *M. maripaludis*, we thank Taiwo Akinyemi, Elliot Shelton, and William B Whitman, supported by US DOE grant DESC0018028. For *T. kodakarensis* and *T. sp.* AM4, we thank Brett Burkhart. For *S. islandicus*, we thank Rachel Whitaker and Changyi Zhang. For *S. acidocaldarius*, we thank Sonja Albers. For *H. salinarium*, we thank David Crowley. For *M. jannaschii*, we thank Biswarup Mukhopadhyay. We thank Patricia Chan for meaningful comments and edits to the manuscript. This work was supported by funding from the USA National Science Foundation, award #2022065 (to TJS and TML), USA National Institutes of Health, award R35 GM143963 (to TJS), and the USA National Aeronautics and Space Administration, award 80NSSC23K1354 (to TJS).

Methods

Culture and RNA isolation

Cultures of *M. maripaludis*, *M. jannaschii*, *S. islandicus*, *S. acidocaldarius*, *T. kodakarensis, T. sp.* AM4, *P. furiosus*, *H. salinarum* and *H. volcanii* were prepared as described previously (Crowley et al., 2006; Jäger et al., 2014; Liman et al., 2022; Mukhopadhyay et al., 1999; Tang et al., 2009; Wagner et al., 2012; Whitman et al., 1987; C. Zhang et al., 2013). For *T. kodakarensis* and *H. volcanii,* we had three biological replicates available (Additional file: Fig. S9). Total RNA from archaeal cell pellets were extracted as described previously in (Tsai et al., 2025). Briefly, cells were treated with BAN reagent (Molecular Research Center, Inc., Cat #BN191). RNA was purified with equal volume of acid phenol-chloroform with isoamyl alcohol (125:24:1, Thermo Fisher Scientific, Cat #AM9722) was added to the reaction and centrifuged to separate the aqueous phase from the organic phase. RNA was then precipitated with 1.5 volumes of isopropanol and 0.1 volume of sodium acetate (Sigma Aldrich, Cat #S7899) at -20C overnight. Finally, the precipitated RNA pellets were washed with 75% ethanol and dissolved in nuclease-free water. Total RNA was used as input for library construction.

OTTR-seq library construction

Total RNA was used for generating OTTR-seq libraries, as previously described (Upton et al., 2021). Slight adjustments to the protocol were made to improve tRNA capture. Briefly, total PNK-treated RNA was 3' tailed using mutant BoMoC RT in buffer containing only ddATP for 90 minutes at 30℃, with the addition of ddGTP for another 30 minutes at 30℃. This was then

heat-inactivated at 65℃ for 5 minutes, and unincorporated ddATP/ddGTP were hydrolyzed by incubation in 5 mM MgCl2 and 0.5 units of shrimp alkaline phosphatase (rSAP) at 37℃ for 15 minutes. 5 mM EGTA was added and incubated for 100℃ for 5 minutes to stop the reaction and denature GC-rich tRNAs. Reverse transcription was then performed at 37℃ for 30 minutes, followed by heat inactivation at 70℃ for 5 minutes. The remaining RNA and RNA/DNA hybrids were then degraded using 1 unit of RNase A at 50℃ for 10 minutes. cDNA was then cleaned up using a MinElute Reaction CleanUp Kit (Qiagen). To reduce adaptor dimers, cDNA was run on a 9% UREA page gel, and the size range of interest was cut out and eluted into gel extraction buffer (300mM NaCl, 10mM Tris; pH 8.0, 1mM EDTA, 0.25% SDS) and concentrated using EtOH precipitation. Size-selected cDNA was then PCR amplified for 12 cycles using Q5 High-fidelity polymerase (NEB #M0491S) and buffer pack with high GC Enhancer (NEB #B9027S). Amplified libraries were then run on a 6% TBE gel, and the size range of interest was extracted to reduce adaptor dimers further. Gel slices were eluted into a gel extraction buffer (300mM NaCl, 10mM Tris; pH 8.0, 1mM EDTA) followed by concentration using EtOH precipitation. Final libraries were pooled and sequenced using 150 SE and 200 cycle kits on an Illumina NovaSeq.

tRNA Analysis of eXpression (tRAX)

Sequencing adaptors were trimmed from raw reads using cutadapt, v1.18, and read counts were generated for RNA types tRAX (Holmes et al., 2022). Briefly, trimmed reads were mapped to their respective strain genomes: *Methanococcus maripaludis* S2, *Methanocaldococcus jannaschii* DSM 2661, *Sulfolobus islandicus* M.16.4, *Sulfolobus acidocaldarius* DSM 639, *Thermococcus kodakarensis* KOD1, *Thermococcus* sp. AM4,

Pyrococcus furiosus DSM 3638, *Halobacterium salinarum* R1, and *Haloferax volcanii* DS2.

Each of the references were combined with their mature tRNAs obtained from GtRNAdb (Chan

& Lowe, 2016). Trimmed reads were mapped using Bowtie2 in very-sensitive mode with the

following parameters to allow for a maximum of 100 alignments per read: –very-sensitive

–ignore-quals –np 5 -k 100. Mapped reads were then filtered to retain only the "best mapping"

alignments. Raw read counts of tRNAs and other small RNA types were computed using tRNA

annotations from GtRNAdb, and annotations from Refseq. Raw read counts were then

normalized using DESeq2.

tRNA modification analysis

Sites of known tRNA modifications were compiled from Modomics (Cappannini et al.,

2023) and relevant literature (Hirata et al., 2019; Krishnamohan et al., 2019; Wolff et al., 2023;

Yu et al., 2019). Modification types for *M. maripaludis*, *M. jannaschii*, *S. acidocaldarius*, *T.

kodakarensis, P. furiosus*, and *H. volcanii* were curated in a tab delimited table (Supplementary

table) and mapped to their corresponding tRNA sequence alignments. Known modification

positions were then mapped onto orthologous tRNAs in unannotated species within phylogenetic

clades, focusing on sites with conserved reference nucleotides and shared rates of

misincorporation.

To assess whether observed misincorporation frequencies were statistically significant

relative to background, we applied tMAP (https://github.com/Aimann/tMAP)(K. Zhang et al.,

2025) using aligned tRNA reads from tRAX (Holmes et al., 2022) as input. At each position

misincorporation frequency was calculated as the number of non-reference base calls divided by

total read coverage. A beta distribution was modeled using these empirical counts, incorporating

pseudocounts derived from the mean and variance of misincorporation rates at known modified positions. A modification threshold was empirically determined as the 75th percentile of misincorporation rates, stratified by reference base identity. P-values were computed using the cumulative distribution function of the beta distribution, representing the probability of observing a misincorporation frequency equal to or greater than the observed value under the null model.

Modification detection relied on two key metrics: the observed misincorporation rate at each position, and the statistical significance (adjusted p-values) of these observed rates. We implemented a base-specific thresholding strategy that accounts for background misincorporation rates inherent to each base. We observed that cytosine and guanine-derived positions tend to exhibit elevated background rates, even at unannotated sites, exceeding 5% and resulting in statistically significant p-values despite the likely absence of true modification. To reduce risk of false-positive predictions driven by these elevated baselines, we set higher thresholds for levels of confidence in these contexts. Specifically, a position and modification type was classified as high-confidence if it met a base-specific misincorporation (>10% for cytosine and guanine; >5% for adenine and uracil) and an adjusted p-value <0.05. Modifications with significant p-values but misincorporation rates below the respective base-specific threshold were classified as moderate-confidence. This approach preserves sensitivity for modifications with potential to induce RT misincorporation, while improving specificity in contexts where signals vary.

Across species, modification types, and positions we observed that variation in detection was largely attributable to differences in the distribution of modification rates across sequence contexts and tRNA positions. The p-value reflects how a given misincorporation rate is relative

to the expected background for a specific base. Even if two different sites have the same rate, they may differ in statistical significance if one falls within a narrow, low-background distribution and the other does not. While low coverage occasionally limited detection, the primary factor influencing statistical significance was the depth and magnitude of misincorporation at a given site.

In cases where multiple modification types are known to occur at the same position, relative misincorporation frequency was used to infer the likely modification. For example, m_2^2G generally causes higher misincorporation than m^2G at position 10, enabling us to discriminate between these two forms in specific archaeal species. This distinction was also partially observable at positions 6 and 26, though with reduced confidence. In some instances, misincorporation patterns were not sufficiently distinct to unambiguously infer modification type. In such cases, known modification types from closely related species with identical reference bases were used to assign a most-likely modification classification.

Homolog search and domain analysis

To search for archaeal homologs and compare domain architectures, we collected representative tRNA modification enzymes that were previously characterized (Constantinesco et al., 1998; Jackman et al., 2003; Menezes et al., 2011; Urbonavičius et al., 2006). Each representative was queried across 220 archaeal genomes using BLASTp with default parameters, then results were filtered to select hits with the lowest e-value for each species and representative. Domain analysis and annotation of highest scoring hits was performed using the InterProScan software package (v5.65-97.0) with the FunFam (4.3.0), MobiDBLite (v2.0), NCBIfam (v13.0), PANTHER (v18.0), SuperFamily (v1.75), CDD (v3.20), Pfam (v36.0),

SMART (v9.0), PRINTS (v42.0) and PIRSF (v2023_05) databases. As e-values are specific to each InterPro database and each utilizes their own e-value post-processing, Pfam and CDD were used preferably when domains were found. All matches were considered tentative hits (https://interproscan-docs.readthedocs.io/en/latest/FAQ.html).

Phylogenetic analysis

The evolutionary relationships among each of the tRNA modification enzymes were inferred using maximum likelihood phylogenies derived from clustered BlastP hits of representatives and their subsequent multiple sequence alignments. In brief, hits for each representative were clustered using MMseq2 (Steinegger & Söding, 2017) using the parameters --min-seq-id 0.3 --cov-mode 0. Clustered groups were subset and aligned with MAFFT using default parameters, then concatenated and aligned again using the parameters --retree 1 --maxiterate 0.

All maximum likelihood phylogenetic trees were estimated using IQ-TREE 2 (v2.1.0). Selection of the best-fit model of amino acid substitution was inferred for the phylogenies using the ModelFinder function in IQ-TREE (LG+F+R6). Custom python scripts were used to prune trees by keeping only BlastP hits of each representative with the lowest e-value in a genome. Trees of each representative and its best-scoring homologs were plotted using the Interactive Tree Of Life (https://itol.embl.de). Predicted domain architectures were annotated onto the phylogenies in Adobe Illustrator.

tRNA secondary structure

The R2DT software (McCann et al., 2025) was used to predict and illustrate tRNA secondary structures. Adobe illustrator was used to annotate structures with predicted modifications and tertiary interactions.

References

Anderson, J. T., & Droogmans, L. (2005). Biosynthesis and function of 1-methyladenosine in transfer RNA. In H. Grosjean (Ed.), *Fine-Tuning of RNA Functions by Modification and Editing* (Vol. 12, pp. 121–139). Springer. https://doi.org/10.1007/b106364

Auffinger, P., & Westhof, E. (2001). An extended structural signature for the tRNA anticodon loop. *RNA (New York, N.Y.)*, *7*(3), 334–341. https://doi.org/10.1017/s1355838201002382

Behrens, A., Rodschinka, G., & Nedialkova, D. D. (2021). High-resolution quantitative profiling of tRNA abundance and modification status in eukaryotes by mim-tRNAseq. *Molecular Cell*, *81*(8), 1802-1815.e7. https://doi.org/10.1016/j.molcel.2021.01.028

Bujnicki, J. M., Feder, M., Radlinska, M., & Blumenthal, R. M. (2002). Structure Prediction and Phylogenetic Analysis of a Functionally Diverse Family of Proteins Homologous to the MT-A70 Subunit of the Human mRNA:m6A Methyltransferase. *Journal of Molecular Evolution*, *55*(4), 431–444. https://doi.org/10.1007/s00239-002-2339-8

Cappannini, A., Ray, A., Purta, E., Mukherjee, S., Boccaletto, P., Moafinejad, S. N., Lechner, A., Barchet, C., Klaholz, B. P., Stefaniak, F., & Bujnicki, J. M. (2023). MODOMICS: A database of RNA modifications and related information. 2023 update. *Nucleic Acids Research*, gkad1083. https://doi.org/10.1093/nar/gkad1083

Chan, P. P., & Lowe, T. M. (2016). GtRNAdb 2.0: An expanded database of transfer RNA genes identified in complete and draft genomes. *Nucleic Acids Research*, *44*(D1), D184–D189.

https://doi.org/10.1093/nar/gkv1309

Constantinesco, F., Benachenhou, N., Motorin, Y., & Grosjean, H. (1998). The
tRNA(guanine-26,N2-N2) methyltransferase (Trm1) from the hyperthermophilic archaeon
Pyrococcus furiosus: Cloning, sequencing of the gene and its expression in Escherichia
coli. *Nucleic Acids Research*, *26*(16), 3753–3761.

Cozen, A. E., Quartley, E., Holmes, A. D., Hrabeta-Robinson, E., Phizicky, E. M., & Lowe, T. M.
(2015). ARM-seq: AlkB-facilitated RNA methylation sequencing reveals a complex
landscape of modified tRNA fragments. *Nature Methods*, *12*(9), 879–884.
https://doi.org/10.1038/nmeth.3508

Crowley, D. J., Boubriak, I., Berquist, B. R., Clark, M., Richard, E., Sullivan, L., DasSarma, S., &
McCready, S. (2006). The uvrA, uvrB and uvrC genes are required for repair of
ultraviolet light induced DNA photoproducts in Halobacterium sp. NRC-1. *Saline
Systems*, *2*, 11. https://doi.org/10.1186/1746-1448-2-11

de Crecy-Lagard, V., Brochier-Armanet, C., Urbonavicˇius, J., Fernandez, B., Phillips, G., Lyons,
B., Noma, A., Alvarez, S., Droogmans, L., Armengaud, J., & Grosjean, H. (2010).
*Biosynthesis of Wyosine Derivatives in tRNA: An Ancient and Highly Diverse Pathway in
Archaea*. 16.

de Crécy-Lagard, V., Boccaletto, P., Mangleburg, C. G., Sharma, P., Lowe, T. M., Leidel, S. A., &
Bujnicki, J. M. (2019). Matching tRNA modifications in humans to their known and
predicted enzymes. *Nucleic Acids Research*, *47*(5), 2143–2159.
https://doi.org/10.1093/nar/gkz011

Erber, L., Hoffmann, A., Fallmann, J., Betat, H., Stadler, P. F., & Mörl, M. (2020). LOTTE-seq
(Long hairpin oligonucleotide based tRNA high-throughput sequencing): Specific
selection of tRNAs with 3'-CCA end for high-throughput sequencing. *RNA Biology*, *17*(1),
23–32. https://doi.org/10.1080/15476286.2019.1664250

Fislage, M., Roovers, M., Tuszynska, I., Bujnicki, J. M., Droogmans, L., & Versées, W. (2012).

Crystal structures of the tRNA:m 2 G6 methyltransferase Trm14/TrmN from two domains of life. *Nucleic Acids Research*, *40*(11), 5149–5161. https://doi.org/10.1093/nar/gks163

Fluke, K. A., Fuchs, R. T., Tsai, Y.-L., Talbott, V., Elkins, L., Febvre, H. P., Dai, N., Wolf, E. J., Burkhart, B. W., Schiltz, J., Brett Robb, G., Corrêa, I. R., & Santangelo, T. J. (2024). The extensive m5C epitranscriptome of Thermococcus kodakarensis is generated by a suite of RNA methyltransferases that support thermophily. *Nature Communications*, *15*(1), 7272. https://doi.org/10.1038/s41467-024-51410-w

Grosjean, H. (2010). Deciphering synonymous codons in the three domains of life: Co-evolution with specific tRNA modification enzymes. *FEBS Letters*, 13.

Grosjean, H., Gaspin, C., Marck, C., Decatur, W. A., & de Crécy-Lagard, V. (2008). RNomics and Modomics in the halophilic archaea Haloferax volcanii: Identification of RNA modification genes. *BMC Genomics*, 26.

Grosjean, H., & Oshima, T. (2007). How Nucleic Acids Cope with High Temperature. In *Physiology and Biochemistry of Extremophiles* (pp. 39–56). John Wiley & Sons, Ltd. https://doi.org/10.1128/9781555815813.ch4

Gupta, R. (1984). Halobacterium volcanii tRNAs. Identification of 41 tRNAs covering all amino acids, and the sequences of 33 class I tRNAs. *Journal of Biological Chemistry, 259*(15), 9461–9471. https://doi.org/10.1016/S0021-9258(17)42723-2

Gustafsson, H. T., Galan, C., Yu, T., Upton, H. E., Ferguson, L., Kaymak, E., Weng, Z., Collins, K., & Rando, O. J. (2022). *Deep sequencing of yeast and mouse tRNAs and tRNA fragments using OTTR* [Preprint]. Genomics. https://doi.org/10.1101/2022.02.04.479139

Hernandez-Alias, X., Katanski, C. D., Zhang, W., Assari, M., Watkins, C. P., Schaefer, M. H., Serrano, L., & Pan, T. (2022). Single-read tRNA-seq analysis reveals coordination of tRNA modification and aminoacylation and fragmentation. *Nucleic Acids Research*, gkac1185. https://doi.org/10.1093/nar/gkac1185

Hirata, A., Nishiyama, S., Tamura, T., Yamauchi, A., & Hori, H. (2016). Structural and functional

analyses of the archaeal tRNA m ^{2} G/m $^2_2$ G10 methyltransferase aTrm11 provide mechanistic insights into site specificity of a tRNA methyltransferase that contains common RNA-binding modules. *Nucleic Acids Research*, *44*(13), 6377–6390. https://doi.org/10.1093/nar/gkw561

Hirata, A., Suzuki, T., Nagano, T., Fujii, D., Okamoto, M., Sora, M., Lowe, T. M., Kanai, T., Atomi, H., Suzuki, T., & Hori, H. (2019). Distinct Modified Nucleosides in tRNA [Trp] from the Hyperthermophilic Archaeon *Thermococcus kodakarensis* and Requirement of tRNA m ^{2} G10/m $^2_2$ G10 Methyltransferase (Archaeal Trm11) for Survival at High Temperatures. *Journal of Bacteriology*, *201*(21), e00448-19, /jb/201/21/JB.00448-19.atom. https://doi.org/10.1128/JB.00448-19

Höfer, K., & Jäschke, A. (2018). Epitranscriptomics: RNA Modifications in Bacteria and Archaea. *Microbiology Spectrum*, *6*(3), 10.1128/microbiolspec.rwr-0015–2017. https://doi.org/10.1128/microbiolspec.rwr-0015-2017

Holmes, A. D., Howard, J. M., Chan, P. P., & Lowe, T. M. (2022). *tRNA Analysis of eXpression (tRAX): A tool for integrating analysis of tRNAs, tRNA-derived small RNAs, and tRNA modifications* (p. 2022.07.02.498565). bioRxiv. https://doi.org/10.1101/2022.07.02.498565

Hori, H. (2014). Methylated nucleosides in tRNA and tRNA methyltransferases. *Frontiers in Genetics*, 26.

Hori, H. (2019). Regulatory Factors for tRNA Modifications in Extreme- Thermophilic Bacterium Thermus thermophilus. *Frontiers in Genetics*, *10*, 204. https://doi.org/10.3389/fgene.2019.00204

Hori, H. (2023). Transfer RNA Modification Enzymes with a Thiouridine Synthetase, Methyltransferase and Pseudouridine Synthase (THUMP) Domain and the Nucleosides They Produce in tRNA. *Genes*, *14*(2), 382. https://doi.org/10.3390/genes14020382

Hori, H., Kawamura, T., Awai, T., Ochi, A., Yamagami, R., Tomikawa, C., & Hirata, A. (2018).

Transfer RNA Modification Enzymes from Thermophiles and Their Modified Nucleosides in tRNA. *Microorganisms*, *6*(4), 110. https://doi.org/10.3390/microorganisms6040110

Ihsanawati, Nishimoto, M., Higashijima, K., Shirouzu, M., Grosjean, H., Bessho, Y., & Yokoyama, S. (2008). Crystal Structure of tRNA *N2,N2*-Guanosine Dimethyltransferase Trm1 from *Pyrococcus horikoshii*. *Journal of Molecular Biology*, *383*(4), 871–884. https://doi.org/10.1016/j.jmb.2008.08.068

Jackman, J. E., & Alfonzo, J. D. (2013). Transfer RNA modifications: Nature's combinatorial chemistry playground: Transfer RNA modifications. *Wiley Interdisciplinary Reviews: RNA*, *4*(1), 35–48. https://doi.org/10.1002/wrna.1144

Jackman, J. E., Montange, R. K., Malik, H. S., & Phizicky, E. M. (2003). Identification of the yeast gene encoding the tRNA m1G methyltransferase responsible for modification at position 9. *RNA (New York, N.Y.)*, *9*(5), 574–585. https://doi.org/10.1261/rna.5070303

Jäger, D., Förstner, K. U., Sharma, C. M., Santangelo, T. J., & Reeve, J. N. (2014). Primary transcriptome map of the hyperthermophilic archaeon Thermococcus kodakarensis. *BMC Genomics*, *15*(1), 684. https://doi.org/10.1186/1471-2164-15-684

Kimura, S., Dedon, P. C., & Waldor, M. K. (2020). Comparative tRNA sequencing and RNA mass spectrometry for surveying tRNA modifications. *Nature Chemical Biology*, *16*(9), 964–972. https://doi.org/10.1038/s41589-020-0558-1

Kowalak, J. A., Dalluge, J. J., McCloskey, J. A., & Stetter, K. O. (1994). The Role of Posttranscriptional Modification in Stabilization of Transfer RNA from Hyperthermophiles. *Biochemistry*, *33*(25), 7869–7876. https://doi.org/10.1021/bi00191a014

Krishnamohan, A., Dodbele, S., & Jackman, J. (2019). Insights into Catalytic and tRNA Recognition Mechanism of the Dual-Specific tRNA Methyltransferase from Thermococcus kodakarensis. *Genes*, *10*(2), 100. https://doi.org/10.3390/genes10020100

Krishnamohan, A., & Jackman, J. E. (2019). A Family Divided: Distinct Structural and

Mechanistic Features of the SpoU-TrmD (SPOUT) Methyltransferase Superfamily. *Biochemistry*, *58*(5), 336–345. https://doi.org/10.1021/acs.biochem.8b01047

Li, J., Li, H., Long, T., Dong, H., Wang, E.-D., & Liu, R.-J. (2019). Archaeal NSUN6 catalyzes m5C72 modification on a wide-range of specific tRNAs. *Nucleic Acids Research*, *47*(4), 2041–2055. https://doi.org/10.1093/nar/gky1236

Liman, G. L. S., Stettler, M. E., & Santangelo, T. J. (2022). Transformation Techniques for the Anaerobic Hyperthermophile Thermococcus kodakarensis. *Methods in Molecular Biology (Clifton, N.J.)*, *2522*, 87–104. https://doi.org/10.1007/978-1-0716-2445-6_5

Lorenz, C., Lünse, C., & Mörl, M. (2017). tRNA Modifications: Impact on Structure and Thermal Adaptation. *Biomolecules*, *7*(4), 35. https://doi.org/10.3390/biom7020035

McCann, H., Meade, C. D., Williams, L. D., Petrov, A. S., Johnson, P. Z., Simon, A. E., Hoksza, D., Nawrocki, E. P., Chan, P. P., Lowe, T. M., Ribas, C. E., Sweeney, B. A., Madeira, F., Anyango, S., Appasamy, S. D., Deshpande, M., Varadi, M., Velankar, S., Zirbel, C. L., … Petrov, A. I. (2025). R2DT: A comprehensive platform for visualizing RNA secondary structure. *Nucleic Acids Research*, *53*(4), gkaf032. https://doi.org/10.1093/nar/gkaf032

Menezes, S., Gaston, K. W., Krivos, K. L., Apolinario, E. E., Reich, N. O., Sowers, K. R., Limbach, P. A., & Perona, J. J. (2011). Formation of m2G6 in Methanocaldococcus jannaschii tRNA catalyzed by the novel methyltransferase Trm14. *Nucleic Acids Research*, *39*(17), 7641–7655. https://doi.org/10.1093/nar/gkr475

Motorin, Y., & Helm, M. (2011). RNA nucleotide methylation. *Wiley Interdisciplinary Reviews. RNA*, *2*(5), 611–631. https://doi.org/10.1002/wrna.79

Mukhopadhyay, B., Johnson, E. F., & Wolfe, R. S. (1999). Reactor-Scale Cultivation of the Hyperthermophilic Methanarchaeon Methanococcus jannaschii to High Cell Densities. *Applied and Environmental Microbiology*, *65*(11), 5059–5065.

Nakano, Y., Gamper, H., McGuigan, H., Maharjan, S., Li, J., Sun, Z., Yigit, E., Grünberg, S., Krishnan, K., Li, N.-S., Piccirilli, J. A., Kleiner, R., Nichols, N., Gregory, B. D., & Hou,

Y.-M. (2025). Genome-wide profiling of tRNA modifications by Induro-tRNAseq reveals coordinated changes. *Nature Communications*, *16*(1), 1047. https://doi.org/10.1038/s41467-025-56348-1

Oerum, S., Dégut, C., Barraud, P., & Tisné, C. (2017). m1A Post-Transcriptional Modification in tRNAs. *Biomolecules*, *7*(1), 20. https://doi.org/10.3390/biom7010020

Ohira, T., & Suzuki, T. (2024). Transfer RNA modifications and cellular thermotolerance. *Molecular Cell*, *84*(1), 94–106. https://doi.org/10.1016/j.molcel.2023.11.041

Ontiveros, R. J., Stoute, J., & Liu, K. F. (2019). The chemical diversity of RNA modifications. *The Biochemical Journal*, *476*(8), 1227–1245. https://doi.org/10.1042/BCJ20180445

Reno, M. L., Held, N. L., Fields, C. J., Burke, P. V., & Whitaker, R. J. (2009). Biogeography of the Sulfolobus islandicus pan-genome. *Proceedings of the National Academy of Sciences*, *106*(21), 8605–8610. https://doi.org/10.1073/pnas.0808945106

Roovers, M., Droogmans, L., & Grosjean, H. (2021). Post-Transcriptional Modifications of Conserved Nucleotides in the T-Loop of tRNA: A Tale of Functional Convergent Evolution. *Genes*, *12*(2), 140. https://doi.org/10.3390/genes12020140

Sas-Chen, A., Thomas, J. M., Matzov, D., Taoka, M., Nance, K. D., Nir, R., Bryson, K. M., Shachar, R., Liman, G. L. S., Burkhart, B. W., Gamage, S. T., Nobe, Y., Briney, C. A., Levy, M. J., Fuchs, R. T., Robb, G. B., Hartmann, J., Sharma, S., Lin, Q., … Schwartz, S. (2020). Dynamic RNA acetylation revealed by quantitative cross-evolutionary mapping. *Nature*, *583*(7817), 638–643. https://doi.org/10.1038/s41586-020-2418-2

Schicho, R. N., Ma, K., Adams, M. W., & Kelly, R. M. (1993). Bioenergetics of sulfur reduction in the hyperthermophilic archaeon Pyrococcus furiosus. *Journal of Bacteriology*, *175*(6), 1823–1830. https://doi.org/10.1128/jb.175.6.1823-1830.1993

Schultz, S. K., & Kothe, U. (2024). RNA modifying enzymes shape tRNA biogenesis and function. *Journal of Biological Chemistry*, 107488.

https://doi.org/10.1016/j.jbc.2024.107488

Shigi, N. (2018). Recent Advances in Our Understanding of the Biosynthesis of Sulfur

Modifications in tRNAs. *Frontiers in Microbiology*, *9*, 2679.

https://doi.org/10.3389/fmicb.2018.02679

Shigi, N., Suzuki, T., Terada, T., Shirouzu, M., Yokoyama, S., & Watanabe, K. (2006).

Temperature-dependent Biosynthesis of 2-Thioribothymidine of Thermus thermophilus

tRNA. *Journal of Biological Chemistry*, *281*(4), 2104–2113.

https://doi.org/10.1074/jbc.M510771200

Sokołowski, M., Klassen, R., Bruch, A., Schaffrath, R., & Glatt, S. (2018). Cooperativity between

different tRNA modifications and their modification pathways. *Biochimica Et Biophysica*

Acta. Gene Regulatory Mechanisms, *1861*(4), 409–418.

https://doi.org/10.1016/j.bbagrm.2017.12.003

Steinberg, S., & Cedergren, R. (1995). A correlation between N2-dimethylguanosine presence

and alternate tRNA conformers. *RNA (New York, N.Y.)*, *1*(9), 886–891.

Steinegger, M., & Söding, J. (2017). MMseqs2 enables sensitive protein sequence searching for

the analysis of massive data sets. *Nature Biotechnology*, *35*(11), Article 11.

https://doi.org/10.1038/nbt.3988

Strassler, S. E., Bowles, I. E., Dey, D., Jackman, J. E., & Conn, G. L. (2022). Tied up in knots:

Untangling substrate recognition by the SPOUT methyltransferases. *Journal of Biological*

Chemistry, *298*(10), 102393. https://doi.org/10.1016/j.jbc.2022.102393

Swinehart, W. E., & Jackman, J. E. (2015). Diversity in mechanism and function of tRNA

methyltransferases. *RNA Biology*, *12*(4), 398–411.

https://doi.org/10.1080/15476286.2015.1008358

Tang, S.-L., Tarasov, V., Athanasopoulos, V., Bath, C., Wendoloski, D., Ferrer, C., Pfeiffer, M.,

Pohlschroder, M., Santos, F., Allers, T., Eichler, J., Camakaris, H., McAlpine, T., Burns,

D., Porter, K., Cukalac, T., & Russ, B. (2009). *Melissa Holmes Masahiro Kamekura Wan*

Lam Stewart Nuttall Wayne Woods Peter Jablonski Juan Serrano Katrina Ngui Josefa Antón.

Tsai, Y.-L., Wolf, E. J., Fluke, K. A., Fuchs, R. T., Dai, N., Johnson, S. R., Sun, Z., Elkins, L., Burkhart, B. W., Santangelo, T. J., & Corrêa, I. R. (2025). Comprehensive nucleoside analysis of archaeal RNA modification profiles reveals an m7G in the conserved P loop of 23S rRNA. *Cell Reports*, *44*(4), 115471. https://doi.org/10.1016/j.celrep.2025.115471

Upton, H. E., Ferguson, L., Temoche-Diaz, M. M., Liu, X., Pimentel, S. C., Ingolia, N. T., Schekman, R., & Collins, K. (2021). *Low-bias ncRNA Libraries using Ordered Two-Template Relay: Serial Template Jumping by a Modified Retroelement Reverse Transcriptase* [Preprint]. Biochemistry. https://doi.org/10.1101/2021.04.30.442027

Urbonavičius, J., Armengaud, J., & Grosjean, H. (2006). Identity Elements Required for Enzymatic Formation of *N*2,*N*2-dimethylguanosine from *N*2-monomethylated Derivative and its Possible Role in Avoiding Alternative Conformations in Archaeal tRNA. *Journal of Molecular Biology*, *357*(2), 387–399. https://doi.org/10.1016/j.jmb.2005.12.087

van Tran, N., Muller, L., Ross, R. L., Lestini, R., Létoquart, J., Ulryck, N., Limbach, P. A., de Crécy-Lagard, V., Cianférani, S., & Graille, M. (2018). Evolutionary insights into Trm112-methyltransferase holoenzymes involved in translation between archaea and eukaryotes. *Nucleic Acids Research*, *46*(16), 8483–8499. https://doi.org/10.1093/nar/gky638

Wagner, M., van Wolferen, M., Wagner, A., Lassak, K., Meyer, B. H., Reimann, J., & Albers, S.-V. (2012). Versatile Genetic Tool Box for the Crenarchaeote Sulfolobus acidocaldarius. *Frontiers in Microbiology*, *3*, 214. https://doi.org/10.3389/fmicb.2012.00214

Westhof, E., Liang, S., Tong, X., Ding, X., Zheng, L., & Dai, F. (2020). Unusual tertiary pairs in eukaryotic tRNAAla. *RNA*, *26*(11), 1519. https://doi.org/10.1261/rna.076299.120

Westhof, E., Thornlow, B., Chan, P. P., & Lowe, T. M. (2022). Eukaryotic tRNA sequences

present conserved and amino acid-specific structural signatures. *Nucleic Acids Research*, *50*(7), 4100–4112. https://doi.org/10.1093/nar/gkac222

Whitman, W. B., Sohn, S., Kuk, S., & Xing, R. (1987). Role of Amino Acids and Vitamins in Nutrition of Mesophilic Methanococcus spp. *Applied and Environmental Microbiology*, *53*(10), 2373–2378.

Wolff, P., Lechner, A., Droogmans, L., Grosjean, H., & Westhof, E. (2023). Identification of Up47 in three thermophilic archaea, one mesophilic archaeon and one hyperthermophilic bacterium. *RNA*, rna.079546.122. https://doi.org/10.1261/rna.079546.122

Xiong, Q.-P., Li, J., Li, H., Huang, Z.-X., Dong, H., Wang, E.-D., & Liu, R.-J. (2023). Human TRMT1 catalyzes m2G or m22G formation on tRNAs in a substrate-dependent manner. *Science China. Life Sciences*. https://doi.org/10.1007/s11427-022-2295-0

Yared, M.-J., Marcelot, A., & Barraud, P. (2024). Beyond the Anticodon: tRNA Core Modifications and Their Impact on Structure, Translation and Stress Adaptation. *Genes*, *15*(3), Article 3. https://doi.org/10.3390/genes15030374

Yu, N., Jora, M., Solivio, B., Thakur, P., Acevedo-Rocha, C. G., Randau, L., de Crécy-Lagard, V., Addepalli, B., & Limbach, P. A. (2019). tRNA Modification Profiles and Codon-Decoding Strategies in *Methanocaldococcus jannaschii*. *Journal of Bacteriology*, *201*(9), e00690-18, /jb/201/9/JB.00690-18.atom. https://doi.org/10.1128/JB.00690-18

Zhang, C., Cooper, T. E., Krause, D. J., & Whitaker, R. J. (2013). Augmenting the Genetic Toolbox for Sulfolobus islandicus with a Stringent Positive Selectable Marker for Agmatine Prototrophy. *Applied and Environmental Microbiology*, *79*(18), 5539–5549. https://doi.org/10.1128/AEM.01608-13

Zhang, K., Manning, A. C., Lentini, J. M., Howard, J., Dalwigk, F., Maroofian, R., Efthymiou, S., Chan, P., Eliseev, S. I., Yang, Z., Chang, H., Karimiani, E. G., Bakhshoodeh, B., Houlden, H., Kaiser, S. M., Lowe, T. M., & Fu, D. (2025). Human TRMT1 and TRMT1L paralogs ensure the proper modification state, stability, and function of tRNAs. *Cell*

Reports, *44*(1), 115092. https://doi.org/10.1016/j.celrep.2024.115092

Zhang, W., & Westhof, E. (2025). The Biology of tRNA t6A Modification and

Hypermodifications—Biogenesis and Disease Relevance. *Journal of Molecular Biology*,

169091. https://doi.org/10.1016/j.jmb.2025.169091

Zheng, G., Qin, Y., Clark, W. C., Dai, Q., Yi, C., He, C., Lambowitz, A. M., & Pan, T. (2015).

Efficient and quantitative high-throughput tRNA sequencing. *Nature Methods*, *12*(9),

835–837. https://doi.org/10.1038/nmeth.3478

Table 1. Summary of species studied in this paper.

Species	Phylum	Class	Habitat	Temperature Preference	Number of tRNA Genes
Sulfolobus islandicus	Crenarchaeota	Thermoprotei	Volcanic hot springs	Thermoacidophile (75°C)	45
Sulfolobus acidocaldarius	Crenarchaeota	Thermoprotei	Terrestrial hot springs	Thermoacidophile (75°C)	46
Pyrococcus furiosus	Euryarchaeota	Thermococci	Hydrothermal marine sediments	Hyperthermophile (95°C)	46
Thermococcus sp. AM4	Euryarchaeota	Thermococci	Sea hydrothermal vents	Hyperthermophile (80°C)	46
Thermococcus kodakarensis	Euryarchaeota	Thermococci	Sea hydrothermal vents	Hyperthermophile (85°C)	46
Haloferax salinarum	Euryarchaeota	Halobacteria	Hypersaline lakes and salterns	Extreme halophile (40°C)	47
Halobacterium volcanii	Euryarchaeota	Halobacteria	Hypersaline environments	Moderate halophile (42°C)	47
Methanococcus maripaludis	Euryarchaeota	Methanococci	Salt marshes	Mesophile (37°C)	35
Methanocaldococcus jannaschii	Euryarchaeota	Methanococci	Marine hydrothermal vent	Hyperthermophile (85°C)	34

Figure 1. Known archaeal tRNA modifications and their misincorporations (MI) rates. A) Phylogenetic tree of representative archaeal species, annotated in boxplots, where each datapoint represents a given tRNA with annotated modifications. The number of known tRNA modifications per species was curated from Modomics and literature sources. Species wherein no literature sources define tRNA modifications are marked with an asterisk(*) B) Cloverleaf schematic of tRNA secondary structure indicating detection confidence level at positions of known tRNA modifications across all species. Modifications are color-coded by estimated confidence of detection via OTTR-seq, based on average misincorporation frequency (MI) and adjusted p-values. High confidence detection (blue) was classified by base-specific MI rates (mean MI > 10% for cytosine and guanine; mean MI >5% for adenine and uracil) and an adjusted p-value <0.05. Modifications with significant p-values but misincorporation rates below the respective base threshold were classified as moderate-confidence (orange). C) Position-specific MI frequencies stratified by base for unannotated and known modified bases. Each point represents a specific species tRNA. MI rates are plotted on a log scale where red lines indicate mean MI. Background MI for unannotated bases (UNAN) includes both unmodified and unannotated positions; edge lines on data points were omitted for clarity. Modifications are labeled below with associated position and modification type. Adjusted p-values represent the probability of observing a misincorporation frequency equal to or greater than the observed value (grey indicates p-adj > 0.05). D) MI profiles at selected positions where multiple known modification types can occur from the same base (e.g. G6, G10, C17, G26, G37, A37). Each subpanel shows MI distributions stratified by specific chemical modification. Differences in MI frequencies aid in distinguishing closely related modification types such as m^2G vs. m^2_2G at G10. Confidence in detection varies across positions and modification types.

Figure 2. Overview of homology-based tRNA modification predictions. A) Representative example of homology-based predictions for the initiator methionine (tRNA^{iMet}) in *P. furiosus* (top) and *T. kodakarensis* (bottom). The coverage plot depicts the known modification sites indicated (e.g. black) and newly predicted sites highlighted (blue). Mapped-read coverage plot illustrates the read depth and pinpoint sites of modification. B,C,D,E,F) Pearson correlation between known modifications vs homology-predicted modifications. Each point in the scatter plot corresponds to a specific modification type and site from homologous tRNAs. The diagonal line represents perfect correlation (r=1) between known and predicted MI. Outliers (± 2 SD) are highlighted (purple) and annotated with their corresponding tRNA, modification, and position. The read identity (pattern of misincorporation) for select outliers are depicted next to the scatter plot, with known modification position (black) compared to newly predicted (blue) (See all outliers in Supplemental Figures) B) *P. furiosus (Pfu)* vs *T. kodakarensis (Tko)*. Select outliers iMet^{CAU} m$^2_2$G at position 26 and Thr^{GGU} m^{1}I at position 57. C) *P. furiosus (Pfu)* vs *T. species AM4 (AM4)*. Select outliers Met^{CAU} m$^2_2$G at position 6, and Arg^{UCU} cnm^{5}U at position 34. D) *H. volcanii (Hvo)* vs *H. salinarum (Hsa)*. Select outliers Leu^{UAA} m^{2}G, Lys^{UUU} m$^2_2$G at position 26, and Lys^{CUU}, Lys^{UUU} t^{6}A at position 37. E) *S. acidocaldarius (Sac) vs S. islandicus (Sis)*. All outliers Glu^{CUC} and Phe^{GAA} m$^2_2$G, and iMet^{CAU} m$^2_2$Gm at position 26. F) *M. jannaschi (Mja)* vs *M. maripaludis (Mma)*. Outlier iMet^{CAU} m$^2_2$Gm at position 26. For positions with more than one modification type (e.g., m^{2}G/m$^2_2$G$_{26}$) "x" was used. This notation is also used for most predictions, as exact modification types weren't ascertained.

Figure 3. Clade-specific coordination patterns in the acceptor stem and conserved clade-specific tRNA modifications. A) Schematic representation of coordinated modification patterns at the acceptor stem of the tRNA cloverleaf secondary structure, highlighting a mutually exclusive relationship between position 6 (typically m²G or m²₂G) and position 67 (m²G). Observed across multiple hyperthermophiles, the data suggest that these modifications do not

co-occur within the same tRNA, possibly due to functional constraints. B) Paired MI frequencies for position 6 and 67 across four hyperthermophilic archaea: *P. furiosus, T. sp.* AM4*, T. kodakarensis,* and *M. jannaschii.* Each line connects paired MI values for individual tRNAs, with colored dots indicating isotype identity. Only moderate and high-confidence predictions (adjusted p-value < 0.05) are shown. Numbers above each panel indicates the total number of predicted modified tRNAs at positions 6 and 67, respectively. A pattern of coordinated modification is consistently observed. C) Distribution of MI frequencies for select moderate to high-confidence tRNA modifications shared across archaeal clades, grouped by phylogenetic class: Thermoprotei (salmon), Thermococci (tan), Halobacteria (blue), and Methanococci (green). Each violin plot summarizes the frequency and variation of a specific modification (e.g. m^1A9, m^1G9, m^2G10, etc.) across species, with counts (above plots) indicating the number of modified tRNAs per group. This highlights the phylogenetic conservation and variability of tRNA modification patterns across lineages.

Figure 4. Phylogenetic and structural diversity of predicted tRNA modification enzymes and their target substrates across Archaea. Each panel presents a phylogenetic tree of predicted homologs of a specific tRNA-modifying enzyme (left), alongside a heatmap summarizing the confidence level of predicted substrates across species (right). Trees are

constructed from clustered protein sequences colored by archaeal clades: Halobacteria (blue), Thermoprotei (salmon), Methanococci (green), and Thermococci (tan). Domain architecture annotations are shown on the tree branches, indicating key functional domains for three class I MTases (Trm14, Trm11, Trm1) and a class IV SPOUT MTase (Trm10). A) Trm14 homologs are restricted to Methanococci and Thermococci and are associated with predicted m^2G6/m^2_2G6 modifications in specific tRNA substrates. The corresponding heatmap shows predicted modifications for representative tRNAs across genomes. B) Trm10 homologs are exclusive to Thermoprotei and Thermococci and are associated with m^1A9 or m^1G9 modifications. The heatmap displays inferred substrate preferences for these homologs across tRNAs and illustrates the dual-functional properties unique to Trm10 in Thermococci. C) Trm11 homologs are broadly conserved, although THUMP domain truncations are observed in Thermoprotei. These homologs are associated with m^2_2G10 modifications in select substrates. Divergence is most notable in tRNA[Pro]. D) Trm1 homologs show widespread conservation, with Thermoprotei variants displaying divergence in the AdoMet domain. Predicted modification patterns at position 26 include m^2_2G and other variants such as m^2G/m^2_2Gm (xG26). Positions previously known and annotated as m^2G reflect sites with adjusted p-values > 0.05, indicating lower statistical support. Together, these patterns highlight coordinated evolution of enzyme domain architectures and their target specificity across archaeal lineages.

Figure 5. Conserved identity elements correlate with recognition of predicted m²₂G10 and xG26 modified tRNAs. A) Base-pairing identities in D-stems corresponding to high-confidence predictions of m²₂G10 modifications in *T. kodakarensis* (Tko) and *H. volcanii* (Hvo). Colors represent base-pair combinations and include the number of predicted modified tRNAs for a given pair, such as non-Watson-Crick pairings G10oU25 (purple) and U13oU22 (lavender). B) Predicted secondary structure of tRNA[Trp] from *T. kodakarensis*, highlighting key structural identity elements for Trm11-mediated m²₂G10 (G10oU25 and U13oU22). These elements form short, unstable D-arms that facilitate enzyme recognition and modification. C) Base-pair

identities in tRNA D-stems associated with high-confidence predictions of xG26 modifications, where G10=C25 are conserved and consistent with anti-determinants for m^2_2G10. D) Predicted secondary structure of tRNA^{Trp} from *H. volcanii*, highlighting structural elements and the absence of m^2_2G10 with the formation of a stable G10=C25 W-C pair. E) Sequence alignment of tRNA^{Trp} D-tem regions species, illustrating conserved identity elements, particularly G10oU25 versus G10=C25. F) Conceptual illustration of structural instability in the D-arm of tRNA^{Trp} from *T. kodakarensis*. Predicted m^2_2G10 modification stabilizes the functional L-shaped tRNA structure by preventing alternative interactions (e.g. a G10=C26 base pair), promoting proper D-loop and T-loop interactions necessary for canonical folding.