Global genome analysis identifies glycolipids and lipoteichoic acid alanylation as contributors to Group B streptococcal diabetic wound infection

Luke R. Joyce[1], Madeline S. Akbari[1], Dustin T. Nguyen[1], Brady L. Spencer[1,2], Kevin S. McIver[3], Alexander R. Horswill[1,4], Kelly S. Doran[1]* and Rebecca A. Keogh[1]*

[1]Department of Immunology and Microbiology, University of Colorado Anschutz, Aurora, CO, USA.

[2]Department of Microbiology, Immunology, and Cancer Biology, University of Viriginia School of Medicine, Virginia, USA

[3]Cell Biology and Molecular Genetics, Maryland Pathogen Research Institute, University of Maryland, College Park, Maryland, USA

[4]Department of Veterans Affairs, Eastern Colorado Healthcare System, Aurora, CO, USA.

*Correspondence: rebecca.keogh@cuanschutz.edu, kelly.doran@cuanschutz.edu

Short title: Glycolipids and LTA alanylation promote GBS survival in diabetic wounds

Abstract

Individuals with diabetes frequently develop chronic, hyper-inflammatory, non-healing wounds, which are the leading cause of all non-traumatic amputations. Group B *Streptococcus* (GBS) is a prominent bacterium isolated from diabetic wound infections and in a murine model of diabetic wound infection, GBS stimulates an influx of neutrophils into the wound. Utilizing a transposon mutant screen, we identified 291 genes required for fitness during diabetic wound infection including enzymes involved in glycolipid biosynthesis and lipoteichoic acid (LTA) alanylation. GBS mutants lacking either LTA alanylation (Δ*dltA*) or lacking all glycolipids (Δ*iagB*) are attenuated in a murine diabetic wound infection. GBS induces primary and secondary degranulation in primary human neutrophils and the Δ*iagB* mutant is significantly more susceptible to neutrophil killing by cationic antimicrobial peptides and reactive oxygen species than Δ*dltA*. Finally, we found that depletion of neutrophils led to significantly greater bacterial recovery, highlighting the importance of neutrophil defense during diabetic wound infection.

Introduction

Diabetes is a group of metabolic diseases, projected to affect over 700 million people by 2045 (1). A frequent complication of diabetes is the development of wounds on the foot and lower limbs (2, 3). These wounds often become chronic, failing to heal within 4 to 6 weeks, due to an unresolved inflammatory phase (4, 5). This inflammation, primarily driven by neutrophils and macrophages leads to the production of reactive oxygen species (ROS), cationic antimicrobial peptides (CAMPs), matrix metalloproteases, and pro-inflammatory chemokines and cytokines, which perpetuate inflammation and limit the reformation of tissue (6, 7).

Diabetic wounds frequently become colonized by a variety of bacterial species which can result in chronic infection (8–11). While some infections are limited to the wound itself, if the infection is uncontrolled, bacteria can penetrate deeper tissue leading to bone and systemic infections, which drive the need for amputation (2, 12, 13). Some of the most prominent bacterial species identified in diabetic wound infection are *Staphylococcus aureus*, *Enterococcus faecalis*, Group B *Streptococcus* (GBS), *Pseudomonas aeruginosa* and many anaerobic species (8, 9, 14, 15). These infections are often difficult to treat due to high levels of antibiotic resistance and tolerance amongst isolates (8, 16, 17). Interestingly, work by Sook et al. has demonstrated that *S. aureus* evolves antibiotic resistance in a diabetic host, and that hyperglycemia potentiates the expansion of resistant populations *in vivo* (18). Further complicating treatment of diabetic wounds is that bacteria in chronic wounds often live in biofilm communities (19, 20). Biofilms, consisting of bacterial cells, DNA, and proteins often lead to a physical barrier which is difficult to treat by antibiotics or routine debridement of tissues (19). These

biofilms can be mono or polymicrobial in nature, and work by Ch'ng et al. has found that *S. aureus* can assist in *E. faecalis* biofilm formation via heme cross-feeding (21).

GBS is a Gram-positive bacterium which colonizes approximately 25% of all individuals in the gastrointestinal and vaginal tract (22–24). In a healthy host, GBS typically colonizes without any adverse health outcomes; however, GBS can cause urinary tract infections (25) and during pregnancy, GBS can ascend vertically and cause intrauterine infections, leading to inflammation and adverse pregnancy outcomes (26, 27). Further, gestational diabetes mellitus has been identified as a risk factor for increased GBS vaginal colonization (28, 29). In immunocompromised individuals such as neonates, and adults with underlying conditions such as diabetes and cancer, GBS can cause diverse infections including pneumonia, meningitis, cardiovascular disease and sepsis (30–34). GBS is frequently isolated from diabetic wounds, despite it rarely colonizing the wounds of non-diabetic individuals (11, 14, 15, 35–40). In addition, GBS alone is sufficient to cause infection and inflammation in the diabetic wound environment (41). Despite this, little is known about which bacterial factors aid in GBS wound establishment, immune cell evasion, and persistence in this niche.

To address this, we previously developed a murine model of GBS diabetic wound infection and found that GBS infection is exacerbated in a diabetic host when compared to a non-diabetic (41). RNA-seq on wound tissues revealed that diabetic mice upregulate numerous genes encoding neutrophil effectors such as MPO, elastase, the subunits of calprotectin, as well as pro-inflammatory cytokines and chemokines in response to GBS infection when compared to uninfected controls. We further found that the abundance of these neutrophil effectors was significantly higher in diabetic mice in comparison to non-

diabetic upon GBS infection. These data support that GBS encounters high levels of

inflammation in the diabetic wound microenvironment, but currently, little is known

regarding the bacterial factors that promote survival in this niche. To date only the

hemolysin/cytolysin (β-H/C, hemolytic pigment) and the plasminogen binding protein,

PbsP, have been shown to contribute to GBS diabetic wound infection (41).

Here, we observe that GBS infection drives an influx of neutrophils into the diabetic

wound microenvironment, and that GBS and neutrophils co-localize spatially in diabetic

wound tissues. We next utilize a global unbiased approach of transposon sequencing

(Tn-seq) and identify genes that are required for fitness in the diabetic wound

environment. We find 291 genes that are significantly underrepresented in this niche,

including genes involved in metabolism, capsule biosynthesis, the production of

membrane glycolipids and lipoteichoic acid (LTA) alanylation. Membrane lipids are a vital

part of host-pathogen interactions (42) and we have recently characterized the role of

GBS glycolipids during systemic infection and meningitis disease progression (43–45).

Utilizing GBS glycolipid mutants and a strain deficient in LTA alanylation, we confirm our

Tn-seq results by finding that all four mutants are attenuated in diabetic wound infection.

We further show the importance of GBS glycolipids and LTA alanylation to keratinocyte

interactions and survival against neutrophil killing by cationic antimicrobial peptides

(CAMPs) and reactive oxygen species (ROS). Together, these data demonstrate the

global gene requirement for GBS fitness in the diabetic wound and further identify GBS

glycolipids and LTA alanylation as contributors to neutrophil survival in this niche.

Results

Neutrophils are the most abundant cell type GBS interacts with during diabetic wound infection.

The diabetic wound is a hyper inflammatory niche with an abundance of immune cells such as phagocytes present (7, 46), however, it is unknown which immune cell populations are the most abundant in response to GBS infection. To address this, we wounded diabetic mice and inoculated the wounds with either GBS WT or PBS controls ($n = 7$ per group) and at 48 hpi, euthanized and perfused mice before harvesting wound tissues. We used flow cytometry to quantify different populations of immune cells (47) (Figure S1A). The primary populations of immune cells present in both the uninfected (mock) and GBS infected wounds were neutrophils, followed by macrophages (Figure 1A). Upon infection with GBS, we observed a significant decrease in the numbers of live cells and live CD45+ cells as a percent of collected singlets (Figure S1B,C), suggesting greater cell death, inclusive of immune cells, in the infected wounds compared to uninfected controls. Live neutrophils as a percent of collected singlets did not change between uninfected and infected wounds (Figure S1D); however, neutrophils did represent a greater proportion of immune cells as the percent of neutrophils of live CD45+ cells compared to the mock controls (Figure 1B). Additionally, a significant decrease in the percent of macrophages of live CD45+ cells compared to mock controls was observed (Figure 1B). This suggests that despite increased immune cell death, the remaining live immune cell population is skewed further toward neutrophils in infected wounds. This is consistent with our previous RNA sequencing results that demonstrated a greater abundance of neutrophil effectors such as MPO and elastase, calprotectin, and CXCL1 in GBS infected diabetic wound tissues when compared to uninfected mice (41). No

differences in the CD11b-CD11c- (including lymphocytes) and the SSC^{low}/MHCII-
(including monocytes and NK cells) populations as a percent of CD45+ cells were
observed (Figure S1E,F). A significant decrease in the percent of CD24+ cells and
dendritic cells was observed (Figure S1G,H), possibly due to the proportional increase of
neutrophils in the wound. Additionally, we did not observe major alterations in
percentages of immune cell populations in the blood of GBS infected and uninfected
controls (Figure S1I). These data indicate that neutrophils are the most abundant immune
cell type GBS encounters in the wound microenvironment at this time point of infection.

To further confirm the immune profile, we utilized multispectral quantitative imaging
of wound tissues excised from diabetic mice infected with WT GBS fluorescently tagged
with GFP or the mock PBS control. Infected mice developed a robust wound under the
epidermal layer (Figure 1C) when compared to the uninfected control. We then chose a
region of interest spanning the wound and epidermal layer (boxed in white) for further
analysis of immune populations. Pathology of the GBS infected region of interest
confirmed a greater proportion of immune cells when compared to the PBS control. GBS
co-localized with Ly6G positive cells, confirming the influx of neutrophils to the site of
infection (Figure 1D). Conversely, higher levels of F4/80 staining were seen in the PBS
control group. These data are the first to demonstrate the spatial dynamics of GBS and
immune cells during diabetic wound infection and highlight that GBS and neutrophils
interact within this microenvironment.

Genome-wide analysis of GBS factors required for survival in the diabetic wound

We next sought to identify which genetic factors GBS requires to colonize and

survive in the diabetic wound. We utilized a GBS Tn mutant library (48–50) in the CJB111

(Serotype V) background (51, 52) and our previously described diabetic wound infection

model as described in materials and methods (41). Groups of mice (4 per group) were

infected with ~1.5 x 10^7 CFU of the GBS Tn library in biological triplicate. Mice were

monitored over the course of infection, and at 48 hours post infection (hpi) mice were

euthanized and wound tissue collected (Figure 2A). The input and wound recovered

libraries were processed as described in the materials and methods.

To identify transposon insertion sites, sequenced reads were mapped to the GBS

CJB111 genome (CP063198) (Figure S2), which identified 291 genes as significantly

underrepresented (padj < 0.05, log2FC < -0.5) and 213 genes as significantly

overrepresented (padj < 0.05, log2FC > 0.5) in the wound compared to the input library

(Figure 2B) (Table S1). The significant gene hits are distributed across the genome and

were then assigned clusters of orthologous groups of proteins (COGs). The number of

significant gene hits in each COG were normalized to the total number of genes in each

COG, to determine the percentage of genes in each COG that were under- or

overrepresented *in vivo*. COGs including nucleotide transport and metabolism,

carbohydrate transport and metabolism, inorganic ion transport and metabolism, lipid

transport and metabolism, and cell wall/membrane/envelope biogenesis had the greatest

proportion of underrepresented genes in the diabetic wound (Figure 2C). We identified

multiple genes known to contribute to GBS infection as significantly underrepresented

such as genes that are involved in β-H/C biosynthesis, capsule biosynthesis, two-

component regulatory systems, glutamine transport, and purine metabolism (Table 1 and

Table S1).

Interestingly, among the most significantly underrepresented genes was *iagA*

(Figure 2D, Table 1) which is a glycosyltransferase involved in the biosynthesis of the

LTA anchor, diglucosyl-diacylglycerol (Glc$_2$-DAG) in GBS (43, 45) (Figure 2E), and is in

an operon with the first glycolipid biosynthesis gene, *iagB*, that is responsible for glucosyl-

DAG (Glc-DAG) biosynthesis (43). An additional gene in glycolipid biosynthesis pathway,

mprF responsible for synthesizing the amino acylated glycolipid lysyl-glucosyl-DAG (Lys-

Glc-DAG) (44, 72) and multiple genes involved in the alanylation of LTA (*dlt* operon) (65)

were also significantly underrepresented in the diabetic wound compared to input (Figure

2D,E). The two remaining genes in the biosynthetic pathway, namely *iagB* and *ltaS*

genes, were not identified in the Tn-seq dataset. However, upon further investigation, we

observed that there were very few to no transposon insertions present in these genes in

the input libraries, thus making it hard to identify their importance with this approach. GBS

glycolipids have recently been shown to impact GBS bloodstream survival (43) and

disruption and inflammation of the blood-brain barrier (BBB) (44, 45), while alanylation of

LTA has been shown to be important for bloodstream survival (64); yet the contribution

of these pathways to GBS survival in the diabetic wound have yet to be elucidated.

GBS glycolipids and LTA alanylation are required for epithelial cell invasion and

survival in the diabetic wound.

To colonize the diabetic wound, GBS must interact with epidermal keratinocyte

cells, a highly specialized epithelial cell (73). To investigate the direct interaction between

GBS WT CJB111 as well as isogenic glycolipid mutants in *iagB*, (43), *iagA* (43), *mprF*

(44) and *dltA* and human keratinocytes, we used the immortalized human keratinocyte

cell line, N/TERT-2G (74). GBSΔ*iagB* is devoid of all glycolipids, GBSΔ*iagA* lacks Glc$_2$-

DAG the LTA anchor, GBSΔ*mprF* lacks the lysine decorated lipids Lys-Glc-DAG and Lys-

PG, and the GBSΔ*dltA* mutant is unable to add alanine modifications to its LTA polymer

(65) (Figure 2E). *In vitro* assays for adhesion and invasion were performed as described

previously for other cell lines (63, 75). There was no significant difference in the ability of

the mutant strains to adhere to N/TERT cells (Figure 3A); however, there were

significantly fewer mutant bacteria recovered from the intracellular compartment

compared to GBS WT (Figure 3B) indicating GBS glycolipids and LTA alanylation play

roles in invasion of human keratinocytes. This may contribute to GBS persistence in the

wound.

To confirm our Tn-seq results and to elucidate the role of glycolipids and LTA

alanylation during diabetic wound infection, we utilized our established streptozotocin

induced diabetic wound model of infection (76) with GBS WT CJB111, Δ*iagB*, Δ*iagA*,

Δ*mprF* and *dltA* mutant strains. After 48 hpi, all mutants were significantly attenuated in

comparison to WT GBS (Figure 3C). Notably, in addition to having reduced bacterial

burden in the wound, mutant strains were not able to spread systemically compared to

WT GBS as evidenced by significantly reduced recovery of mutant bacteria from the

kidneys (Figure 3D). Additionally, far fewer mice had detectable CFU in the kidneys

compared to WT infected mice (Figure 3E). These data indicate that GBS glycolipids and

LTA alanylation play an important role in GBS wound survival and systemic spread during

diabetic infection.

Next, we assessed how GBS glycolipids and LTA modification would impact the

abundance of neutrophil chemoattractant, CXCL1, and the granule enzyme

myeloperoxidase (MPO) in the diabetic wound at time of death. Wounds from mice

infected with Δ*iagB* and Δ*iagA* had a significant decrease in both CXCL1 compared to

GBS WT whereas wounds infected with Δ*mprF* and Δ*dltA* had no significant difference in

CXCL1 abundance (Figure 3F).   Interestingly, only Δ*iagA* infected wounds had a

significant decrease in the abundance of MPO compared to GBS WT infected wounds,

suggesting high levels of neutrophils were still present in the Δ*iagB, ΔmprF,* and Δ*dltA*

wounds to combat the infection (Figure 3G). These data suggest that as bacterial burdens

decrease in the wound, the neutrophil signaling is reduced and results in lowered

neutrophil recruitment to the wound.

GBS glycolipids and LTA alanylation are vital for surviving neutrophil killing.

To further understand the interaction between GBS and neutrophils, we isolated

primary human neutrophils from four independent donors (two female and two male) to

assess primary human neutrophil killing of GBS in diabetic glucose levels. Primary human

neutrophils were conditioned in media containing 22 mM glucose to mimic hyperglycemia

(46) before being exposed to opsonized bacteria. To assess the role of all GBS

glycolipids, including the LTA anchor, we used a GBSΔ*iagB* mutant and to assess impact

of LTA alanylation we used the GBSΔ*dltA* mutant. A significant decrease in survival was

observed for GBSΔ*iagB* compared to GBS WT, which is consistent to our previous study

(43) of GBSΔ*iagB* susceptibility to neutrophil killing under non-diabetic conditions;

however, no significant difference was observed between GBSΔ*dltA* and GBS WT (Figure

4A). The phenotype of the GBSΔ*iagB* mutant was rescued when the strain was

complemented with a plasmid containing the *iagB* gene *(+*pJIagB). GBSΔ*dltA* is known to

be more susceptible to neutrophil killing and bloodstream clearance (64) so we assessed

opsonophagocytic neutrophil mediated killing over time using the human HL60-neutrophil

like cell line that is routinely maintained and differentiated in high glucose media.

Interestingly, we observed that GBSΔ*iagB* is significantly more susceptible to neutrophil

killing at 1, 2 and 3 h compared to GBS WT, whereas only after 2 and 3 h of infection was

there a significant difference between GBSΔ*dltA* and GBS WT (Figure 4B). Additionally,

after 3 h of exposure there was less than 1% survival for both mutants whereas WT

maintains approximately 15% survival. These data indicate that GBS glycolipids and LTA

alanylation both contribute to thwarting neutrophil clearance.

Glycolipids contribute to GBS survival against CAMPs and ROS.

Neutrophils utilize multiple antimicrobial components to kill bacteria, especially

CAMPs such as cathelicidin and ROS such as hydrogen peroxide and hypochlorite (77–

80). The alanylation of LTA has previously been shown to contribute to GBS evasion of

CAMPs by regulating the density of the cell wall (64, 81). Similarly, we have recently

demonstrated that all three GBS glycolipids aid in the protection against CAMP-mediated

killing, likely through a charge repulsion mechanism (43). Here, we observed GBSΔ*iagB*

is more susceptible to neutrophil killing compared to GBS WT and to GBSΔ*dltA*. To

assess if the same is true with CAMP susceptibility, empty vector strains of GBS WT,

GBSΔ*iagB*, GBSΔ*dltA*, and complement strains GBSΔ*iagB*+pJIagB and

GBSΔ*dltA*+pJDltA were incubated with 20 µM of the mouse cathelicidin, mCRAMP, for

30 min. We observed that the GBSΔ*iagB* mutant exhibited significantly more mCRAMP

sensitivity compared to GBS WT and GBSΔ*dltA* strains, and the complemented strains

restored survival to WT levels (Figure 4C). Possible explanations for this difference in

CAMP susceptibility are alterations in membrane fluidity (82). However, no difference in

membrane fluidity was observed between all strains as measured by anisotropy,

indicating CAMP susceptibility is independent of membrane fluidity in these mutants

(Figure S3A).

ROS is another major mechanism of bacterial killing employed by neutrophils and

diabetic wounds have increased levels of ROS compared to nondiabetic wounds (83, 84);

thus, we hypothesized that glycolipids and LTA alanylation would contribute to ROS

survival. To investigate this, empty vector strains of GBS WT, GBSΔ*iagB*, GBSΔ*dltA*, and

complement strains GBSΔ*iagB*+pJIagB and GBSΔ*dltA*+pJDltA were incubated with

0.08% hypochlorite (HOCl) for 1 h resulting in a significant decrease in survival of the

GBSΔ*iagB* mutant compared to GBS WT, which was rescued to WT survival levels in the

pJIagB complement strain (Figure 4D). No difference in survival was observed between

GBSΔ*dltA,* pJDltA, and GBS WT. To further confirm ROS susceptibility and assess

oxidative DNA damage; GBS WT, GBSΔ*iagB* and GBSΔ*dltA* were exposed to 100 µg of

hydrogen peroxide for 1 h resulting in a significant decrease in survival of the GBSΔ*iagB*

mutant compared to GBS WT. No difference between GBS WT and GBSΔ*dltA* was

observed (Figure 4E). Similarly, we measured oxidative DNA damage via ELISA and

observed a significant increase in fold change of oxidative DNA damage in the GBSΔ*iagB*

strain compared to untreated, GBS WT, and GBSΔ*dltA* strains (Figure 4F). No difference

was observed between GBSΔ*dltA* and GBS WT or untreated cells. These data suggest

that LTA alanylation is important for resisting CAMPs; however, GBS glycolipids promote

resistance to both CAMPs and ROS.

GBS induces primary and secondary degranulation in primary human neutrophils

An important mechanism of neutrophil killing of bacteria is degranulation.

Neutrophils produce four types of granules: azurophilic/primary, specific/secondary,

gelatinase/tertiary and secretory, with high concentrations of CAMPs, ROS, proteinases

and additional antimicrobial factors harbored in the primary and secondary granules (77–

80). To assess induction of degranulation we exposed primary human neutrophils in high

glucose conditions to opsonized GBS WT, Δ*iagB* and Δ*dltA* for 30 min and assessed

degranulation using flow cytometry for established human degranulation markers (85)

(Figure S3B). We observed a significant increase in azurophilic/primary degranulation,

indicated by the increased surface expression of CD63 (Figure 5A) and

specific/secondary degranulation, indicated by the increased surface expression of

CD66b (Figure 5B) and CD15 (Figure S3C), of neutrophils exposed to GBS regardless of

the GBS strain, compared to uninfected control neutrophils. Interestingly, the GBSΔ*iagB*

mutant, even though more susceptible to ROS and CAMPs, induces significantly higher

levels of azurophilic/primary degranulation compared to GBS WT and Δ*dltA* mutant. While

GBS infection resulted in no difference in surface expression of gelatinase/tertiary

degranulation marker CD11b between infected and uninfected neutrophils, we did

observe an increase in surface CD18 expression on infected neutrophils compared to

uninfected neutrophils, suggesting tertiary degranulation is not stimulated or is stimulated

to a lesser extent (Figure S3D). These data indicate that neutrophils recognize GBS,

regardless of the missing glycolipids or LTA alanylations, and trigger primary and

secondary degranulation to combat them.

Neutrophils influence GBS burden in the wound and kidneys of diabetic mice.

Neutrophils are highly abundant during diabetic wound infection and promote high

inflammation in the microenvironment (46, 86), as we have also observed during GBS

infection (Figure 1). We therefore sought to investigate the contribution of neutrophils to

GBS clearance in a diabetic host, and we hypothesized that depletion of neutrophils in

the diabetic wound would lead to enhanced recovery of the GBSΔ*iagB* and GBSΔ*dltA*

mutants in comparison to non-depleted controls. To investigate this, systemic neutrophil

depletion of diabetic mice was performed via intraperitoneal injection of anti-Ly6G

antibody, and non-depleted mice were injected with isotype (anti-IgG2a antibody) control

one day before infection. Neutrophil depletion in wound tissues was confirmed by flow

cytometry (Figure S4). Neutrophil depleted and control diabetic mice were infected with

either GBS WT, GBSΔ*iagB* or GBSΔ*dltA* mutants using our streptozotocin induced

diabetic wound infection model. Depletion of neutrophils lead to a significant increase

bacterial burden of GBS WT, GBSΔ*iagB* and GBSΔ*dltA* in the diabetic wound compared

to the non-depleted controls at 48 hpi (Figure 6A). While GBS burdens were consistently

higher in depleted compared to non-depleted controls, the Δ*iagB* and Δ*dltA* mutants still

exhibited a decreased burden when compared to GBS WT within the neutrophil depleted

group (Figure 6A).

Depletion of neutrophils led to greater GBS dissemination into the kidneys based

on CFU recovered (Figure 6B). Interestingly, significant increases in bacterial burdens in

the kidneys of depleted mice infected with GBSΔ*iagB* and GBSΔ*dltA* compared to non-

depleted mice was observed and a higher percentage of mice had CFU in the kidneys in

the depleted groups vs non-depleted groups (Figure 6C). These data indicate that

neutrophils play an important role in controlling GBS persistence in the diabetic wound

and preventing systemic spread. They further suggest that circulating neutrophils can

control both the GBSΔ*iagB* and GBSΔ*dltA* during invasive infection *in vivo*.

Discussion

The diabetic wound is hyperinflammatory, with high levels of neutrophils and

macrophages potentiating inflammation and limiting reformation of the extracellular matrix

(87). However, prior to this work it was unknown how the presence of GBS in diabetic

wound infection would affect immune cell populations. Using flow cytometry and

multispectral immunohistochemistry, we found that neutrophils are the most abundant

immune cell GBS encounters during initial diabetic wound infection (48 hpi), making up

greater than 65% of all immune cells. We also found a significant increase in the

percentage of neutrophils residing in the diabetic wound infected with GBS compared to

uninfected controls. In addition to flow cytometry, we performed multispectral

immunohistochemistry on GBS and mock infected wounds to visualize the localization of

both GBS and immune cells during diabetic wound infection. Wounds infected with GBS

led to the formation of a large bolus of bacteria within the wound underneath the area of

initial incision. This area contained GBS that seemed to co-localize with neutrophils at the

wound site. This was not the case for all immune populations as macrophages were more

present in the tissue section from mock infected wounds in comparison to GBS infected.

These data are the first to spatially characterize the GBS-host interactions within diabetic

wound infection and demonstrate the proximity of neutrophils and the bacterium within

the wound . Further, it highlights the need for mechanistic studies that identify GBS factors

that contribute to neutrophil survival in this niche.

Despite the growing evidence that GBS is present in diabetic wound infections,

little is known about the genetic factors required for fitness and survival in the wound.

Here, we used a highly saturated Tn-seq library and identified 291 significantly

underrepresented genes in diabetic wound infection, indicating they may be required for

fitness. One of the most significantly underrepresented genes was *iagA*, which encodes

for an enzyme involved in GBS glycolipid biosynthesis and is responsible for synthesizing

the LTA anchor, Glc_2-DAG, (43, 45). Additionally, multiple genes encoding enzymes

involved in the alanylation of GBS LTA were significantly under-represented (65),

suggesting that both LTA anchoring and modification are important for GBS to establish

itself in this niche. Further investigation of the dataset identified *mprF,* another gene in

the GBS glycolipid biosynthesis pathway, and the enzyme responsible for lysinylation of

Glc-DAG and phosphatidylglycerol (44). Two genes in these biosynthetic pathways (*iagB*

and *ltaS*) were not identified in our Tn-seq screen, however, very few transposon

insertions in these genes were present in the input library. *LtaS* is known to be essential

in other Gram-positive bacteria (88, 89); however, *iagB* is not essential thus highlighting

a limitation of large scale datasets and potential for false negative hits. In this work, we

characterize the role of GBS glycolipids and the alanylation of LTA in diabetic wound

infection. Using our isogenic glycolipid mutants (∆*iagB,* ∆*iagA,* ∆*mprF*) and our newly

constructed LTA alanylation mutant (∆*dltA*), we confirmed our Tn-seq results observing

significant attenuation of these mutants in the diabetic wound compared to GBS WT.

Throughout this work, we focused on the mechanisms by which GBS glycolipids and LTA

modification contribute to survival in the diabetic wound using the ∆*iagB* (lacking all

glycolipids) and ∆*dltA* (unable to add alanine modifications to LTA) mutant strains.

Little is known about GBS interaction with keratinocytes, with the only factor known

to aid adherence to keratinocytes being the adhesin, PbsP (41). Interestingly, we

observed the GBS glycolipids and LTA alanylation do not play a role in adherence to

human keratinocytes, but are important for invasion into the intracellular compartment, as

has been observed previously for the BBB endothelial cells (43–45). The glycolipid

mutant, ∆*mprF,* is more susceptible to low pH compared to GBS WT (44) highlighting the

glycolipid mutants may be more susceptible to lysosomal killing compared to GBS WT

once inside the host cell. It is currently unknown what receptors GBS binds to on

keratinocytes, the mechanisms of invasion, and additional GBS factors such as

membrane vesicles that may promote invasion into the intracellular compartment of

keratinocytes. Thus, further investigation into the mechanisms of interactions of GBS and

keratinocytes are warranted, as this may serve as a mechanism to escape from immune

cells in the wound and act as potential reservoirs for persistence colonization of the

diabetic wound after debridement of the wound.

We have previously found that GBS glycolipids are important for survival against

neutrophils, as strains lacking glycolipids have enhanced susceptibility to antimicrobial

peptides. However, these studies were performed with neutrophils cultured in

normoglycemic levels of glucose. Individuals with diabetes are known to have aberrant

neutrophil function, largely driven by hyperglycemia (90). We therefore cultured

neutrophils in media containing levels of glucose that are physiologically relevant in a

diabetic host to assess the role of GBS glycolipids and LTA alanylation to neutrophil

survival in high glucose conditions. Here, we also focused on a second mechanism of

neutrophil killing, degranulation, using our glycolipid devoid strain, Δ*iagB*, and our LTA

alanylation mutant, Δ*dltA*. Using primary human neutrophils exposed to high glucose

conditions, we observed all GBS strains, regardless of mutation, triggered both primary

and secondary degranulation. This suggests that the mechanism by which neutrophils

recognize GBS and elicit degranulation is not dependent on GBS glycolipids or LTA

modification. Neutrophil granules, particularly the primary/azurophilic and

secondary/specific granules, harbor a variety of antimicrobial properties such as CAMPs

and ROS. Due to the hyperglycemia in diabetic individuals, neutrophils are

transcriptionally primed to undergo NETosis (91). Work by Derré-Bobillot et al. identified

a secreted GBS nuclease, NucA involved in the degradation of the DNA matrix, or major

backbone of NETs (92). Further, our group has found that this nuclease is highly

upregulated in bacteria recovered from the diabetic wound in comparison to input (41).

NucA was not identified in the Tn-seq screen, likely because it is a secreted factor that

the rest of the population is still producing, preventing it from being underrepresented;

however, it may be important in the diabetic wound microenvironment, and its contribution

to GBS survival in the diabetic wound is ongoing.

Cationic antimicrobial peptides such as cathelicidin and defensins are small

peptides used by the host to target invading pathogens and are an important antimicrobial

factor in neutrophil granules. We and others have shown that GBS glycolipids and LTA

alanylation are important for resisting CAMP mediated killing although potentially via

different mechanisms (43, 64, 81). Interestingly we observed that GBSΔ*iagB* was

significantly more attenuated than the GBSΔ*dltA* after 30 min of exposure to mouse

cathelicidin, even though the membrane fluidity is the same between these mutants and

GBS WT. A possible explanation is that while the alanylation of LTA is required for proper

cell wall density to mediate CAMP resistance (81), the loss of the glycolipids, including

the LTA anchor, results in a more severe change in cell wall dynamics possibly making

the membrane hyper-attractive to the positively charged CAMPs. Another major

antimicrobial property of neutrophil granules is the production of ROS, especially

hydrogen peroxide which is used by myeloperoxidase to generate hypochlorite. Not only

do neutrophils generate high amounts of ROS, but the diabetic wound contains a large

amount of ROS compared to non-diabetic wounds (83, 84). Like the CAMP phenotype,

we observed GBS devoid of glycolipids is more susceptible to both hypochlorite and

hydrogen peroxide killing and resulting oxidative DNA damage compared to both WT and

Δ*dltA* mutant. This could be due to increased lipid peroxidation of the cellular membrane

lacking glycolipids which may generate even more ROS, resulting in increased damage

to the cell and death (93), but further investigation is needed to identify why glycolipids

may protect against ROS attack.

Among the significantly underrepresented genes identified in this Tn-seq are

multiple factors required for metabolism such as purine metabolism, glutamine

metabolism, and mannose utilization. These genes are important for bacterial

pathogenesis (94) and were recently identified to be important for *Enterococcus*

colonization of non-diabetic wounds (95). We also identified multiple GBS virulence

factors such as the β-H/C *cyl* operon and the type VII secretion system (T7SS) as under-

represented in the diabetic wound. The loss of the GBS β-H/C has been previously shown

to impact bloodstream survival and phagocytic defense, and we have also observed its

importance to diabetic wound persistence (41, 96). Similarly, the role of the T7SS has

recently been implicated in cytotoxicity to host cells and promotes colonization of the

vaginal tract (59, 97). With the high number of neutrophils and macrophages present in

the diabetic wound the T7SS may have important functions for defending against these

immune cells and potentially against other bacterial species that colonize this

polymicrobial environment. Interestingly, multiple two component systems (TCS) were

significantly underrepresented in the Tn-seq. One such TCS, CiaH/R has been implicated

in monitoring cell wall/surface stress (62, 98). GBS TCS 8, which has homology to the

VicR/K (WalR/K in some species) system in other streptococci was also significantly

underrepresented (61, 99). Interestingly three currently uncharacterized TCS systems,

TCS 7, 12 and 19 (61) were also significantly underrepresented. The fact that these TCS

systems were significantly underrepresented suggests the strong need for GBS

adaptation to the diabetic wound microenvironment. It further suggests these TCS

systems may play a pivotal, currently unknown, role in GBS niche adaption, which

requires further investigation.

Overall, using Tn-seq, this work has identified multiple GBS factors that are

required to colonize the diabetic wound and to evade the large number of immune cells

present there (Figure 7). Neutrophils are important first line responder to invading

pathogens and the ability for GBS to evade the multiple killing mechanism they possess

is vital to establishment in this niche. The characterization of the role that GBS glycolipids

and LTA alanylation play during colonization of the wound and evasion of neutrophil killing

highlights important cell wall properties that could be targeted by small molecules to aid

in the clearance of GBS. Future studies aim to elucidate the roles of other genetic factors

identified in the Tn-seq and their contribution to diabetic wound colonization. Ultimately,

this work will provide increased knowledge of GBS pathogenesis and virulence potential,

aiding in future treatments and may have broad implications to other bacterial species

that may colonize the diabetic wound.

Acknowledgements

The authors would like to thank the members of the Doran and Horswill labs for

constructive discussions in the preparation of this manuscript, Dr. Robb Welty at the

Biophysics Shared Research Facility at the University of Colorado Anschutz Medical

Campus for help with anisotropy experiments and the Human Immune Monitoring Shared

Resource, specifically Troy Schedin and Angela Minic (RRID:SCR_021985) within the

University of Colorado Cancer Center (P30CA046934) for their expert assistance in

analysis of diabetic wound tissues. This work was funded in part by the American Heart

Association grant 23POST1013835 to L.R.J, NIH/NIAID F31AI178881 to M.S.A,

483 Author Contributions

Conceptualization: LRJ, KSD, RAK; Methodology: LRJ, MSA, DTN, BLS, RAK;

Validation: LRJ, MSA, DTN, BLS, KSD, RAK ; Formal analysis: LRJ, MSA, DTN, RAK;

Investigation: LRJ, MSA, DTN, RAK; Resources: KSM, ARH, KSD; Writing original draft:

LRJ, RAK; Editing and reviewing: LRJ, MSA, DTN, BLS, KSM, ARH, KSD, RAK;

Visualization: LRJ, DTN, RAK; Project administration: LRJ, KSD, RAK; Funding

acquisition: LRJ, KSM, ARH, KSD, RAK

491 Declaration of interests

492 The authors declare no competing interests

494 Material and Methods

495 Bacterial strains and growth conditions.

496 All strains used in the study are listed in Table S2. GBS strains were grown statically at

497 37°C in Todd-Hewitt broth (THB). *Escherichia coli* strains were grown in lysogeny broth

498 at 37°C or 30°C with rotation at 225 rpm. Erythromycin and spectinomycin (Sigma-Aldrich)

499 were supplemented to media at 300 and 100 µg/mL for *E. coli* or 5 and 100 µg/mL for

500 GBS strains, respectively.

Routine molecular biology techniques

All PCR reactions utilized Phusion polymerase (New England Biolabs). PCR products

and restriction digest products were purified using Qiagen PCR purification kit (Qiagen)

per manufacturer protocols. See Table S3 for primers. Plasmids were extracted using

Qiagen plasmid mini and maxiprep kits (Qiagen) per manufacturer protocols. Restriction

enzyme digests utilized XhoI, PstI, BamHI, and SacII (New England Biolabs) for at least

3 h at 37°C. Ligations utilized T4 DNA ligase (New England Biolabs) at 16°C overnight.

All plasmid constructs were sequence confirmed by Nanopore amplicon sequencing

(Quintara Biosciences).

Construction of *dltA* knockout plasmid

Construction of the *dltA* clean deletion knockout plasmid was performed as previously

described (43, 44). Briefly, approximately 1.5 kb upstream and downstream of the

CJB111 *dltA* (ID870_00720), and the spectinomycin cassette from plasmid JC303 (100)

was amplified using PCR. Plasmid, pMBSacB (101), and the PCR products were

digested using appropriate restriction enzymes and ligated overnight and transformed

into chemically competent *E. coli* MC1061 as described (43). Cultures were pelleted using

an Eppendorf 5810R centrifuge at 3900 x g for 10 min at room temperature. Plasmid was

extracted as described above and sequence verified via Nanopore whole plasmid

sequencing (Quintara Biosciences).

Deletion and complementation of CJB111 *dltA*

GBS electrocompetent cells were generated as described and double crossover homologous recombination knockout strategy was performed as described previously (43, 44). After the second cross-over, GBS was incubated in THB + 0.75 M sucrose and 100 µg/mL spectinomycin at 37°C before plating on THB agar supplemented with 100 µg/mL spectinomycin. Colonies were PCR screened for loss of gene using primers outside of the recombination arms and grown in THB at 37°C and stocked. Sequence confirmation of the mutants was done via Nanopore sequencing (Quintara Biosciences). The pJDltA complementation plasmid was constructed as previously described (43, 44). CJB111 *dltA* gene was amplified, restriction digested and ligated overnight into pDCErm (102) plasmid backbone. Ligations were transformed into *E. coli* for maintenance. Complement plasmids sequence confirmed as above and were electroporated into GBS electrocompetent cells prepared as described above.

Murine model of diabetic wound infection.

8 to 12-week-old male mice from the C57Bl/6J or *Lepr^{db}* background were used throughout the study. Two weeks prior to infection mice were given low dose injections of Streptozotocin (50mg/kg) (CAS-No: 18883-66-4) dissolved immediately before injection in 100 µL of 50 mM sodium citrate buffer (pH 4.5) for 5 consecutive days, followed by a glucose chaser (250 µL) to prevent hypoglycemia (76). Mice then sat for a minimum of one week for blood sugar to normalize before infection. Mice were infected as previously described (76). Briefly, the day before infection mice were anesthetized via isoflurane

inhalation, shaved and treated with Nair on the back. Tail snips were performed and blood

sugar measured via glucometer to ensure all mice had blood glucose levels >200 mg/dl.

The following day, mice were weighed and anesthetized via isoflurane inhalation and

injected with lidocaine intra-dermally. Mice were wounded with a 6 mm biopsy punch and

inoculated with 1 x 10^7 CFU GBS or a PBS control unless otherwise indicated. Wounds

were wrapped with the surgical adhesive Tegaderm for 5 h before adhesive removal and

monitoring of animals. After 48 h of infection, mice were sacrificed and wound as well as

kidneys were harvested for further analysis. Tissues were placed into 500 µl of sterile

PBS in 2.0-ml conical tubes (Thermo Fisher Scientific) along with 1.0-mm-diameter

zirconia/silica beads (BioSpec catalog no. 1107911) and homogenized by bead beating

two times for 60 s in a BioSpec mini bead beater. Tissue homogenates were plated on

GBS CHROMagar [SB282(B)], which allows only for the growth of GBS (in pink) and

Enterococcus spp. (in blue). These experiments were approved by the committee on the

use and care of animals at the University of Colorado–Anschutz Medical Campus protocol

no. 00987. For neutrophil depletion experiments, mice were injected with 200 µg of either

anti-Ly6G (BioXCell, clone 18A, catalog BE0075-1) or anti-IgG2a isotype control antibody

(BioXCell, clone 2A3, catalog BE0089) diluted in buffered reagent diluent (BioXCell,

catalog IP0070) the day before infection.

Diabetic wound transposon library infection

Triplicate cultures of the pooled CJB111 pKrmit transposon library (49) were grown

overnight at 37°C in THB with kanamycin at 300 µg/mL and grown to exponential phase

without kanamycin before libraries were normalized to ~1.5 x 10^7 CFU/10 µL and *Lepr^{db}*

mice were infected as described above. Input library and wound homogenate was plated

in duplicated on CHROMagar Strep B with 300 µg/mL kanamycin and incubated overnight

at 37˚C to collect recovered transposon mutants. Bacterial growth from spread plates

were collected and 4 mice per library pooled together before genomic DNA was extracted

using ZymoBiomics DNA miniprep Kit (Zymo Research).

Transposon library sequencing.

Libraries were prepared and sequenced at the University of Minnesota Genomics Center

(UMGC) according to https://www.protocols.io/view/transposon-insertion-sequencing-tn-

seq-library-pre-rm7vzn6d5vx1/v1. Briefly, genomic DNA was enzymatically fragmented,

and adapters added using the NEB Ultra II FS kit (New England Biolabs), and ~50 ng of

fragmented adapted gDNA was used as a template for enrichment by PCR (16 cycles)

for   the   transposon   insertions   using   mariner-specific

(TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGCCGGGGACTTATCATCCAACC)

and Illumina P7 primers. The enriched PCR products were diluted to 1ng/ul and 10 ul was

used as a template for an indexing PCR (9 cycles) using Nextera_R1 (iP5) and

Nextera_R2 (iP7) primers. Sequencing was performed using 150- base paired-end format

on an Illumina NovaSeq 6000 system to generate ~40 million reads per library.

Tn-seq bioinformatics analyses.

Bioinformatic analyses were performed as previously described (50) Reads were trimmed

using Cutadapt (v 4.2) with the following parameters: sequence length with a minimum of

12 bases, removal of flanking "N" bases, reads were trimmed of 3' "G" bases, and reads

were trimmed with the reverse complemented mariner transposon sequence

(ACTTATCAGCCAACCTGTTA). TRANSIT (v 3.2.7) (103) was used to align trimmed

reads to the CJB111 genome (CP063198) for analysis of transposon insertion sites. The

Transit PreProcessor processed reads using default parameters with the Sassetti

protocol, no primer sequence, and mapped them to the genome using Burrows-Wheeler

Alignment (BWA) (104). Insertion sites were normalized using the total trimmed reads

(TTR) in TRANSIT and analyzed using the site-restricted resampling to compare the

insertion counts recovered in wound vs the input library. Resampling was performed using

default parameters, with the addition of ignoring TA sites within 5% of the 5' and 3' end

of the gene. All sequencing reads have been deposited into NCBI SRA under BioProject

accession PRJNA1232139.

N/TERT adherence, invasion and infection assays.

Human immortalized N/TERT-2G keratinocyte cells (74, 105) were seeded at 1×10^5 cells

in keratinocyte serum-free medium (KSFM, Gibco) supplemented with 0.3 mM calcium

chloride (PromoCell) in a 24-well tissue culture treated plate and incubated at 5% CO_2

and 37°C. Assays to determine the total number of bacteria adhered to host cells or

intracellular bacteria were performed using GBS empty vector and complement strain as

described previously (43, 44). Briefly, exponential phase bacteria were normalized to $1 \times$

10^8 CFU/mL in PBS and 1×10^5 CFU per well at multiplicity of infection (MOI) of 1. The

total cell associated GBS were recovered after 30 min incubation. Cells were washed four

times with 1X PBS (Sigma) and detached by addition of 100 µL of 0.25% trypsin-EDTA

solution (Gibco) and incubation for 5 min at 37˚C before lysing the eukaryotic cells with

the addition of 400 µL of 0.025% Triton X-100 (Sigma) and vigorous pipetting. The lysates

were serially diluted and plated on THB agar and incubated overnight to determine CFU.

Bacterial invasion assays were performed as described above except infection plates

were incubated for 2 h before incubation with 100 µg gentamicin (Sigma) and 5 µg

penicillin (Sigma) supplemented media for an additional 2h to kill all extracellular bacteria,

prior to being trypsinized, lysed, and plated as described. Experiments were performed

in biological triplicate with three technical replicates per experiment.

622   Dissociation of wound tissue and blood for flow cytometry

Single cell suspension preparation was performed using methods previously described

by our group (47). In brief, following euthanasia, mice were perfused with PBS+heparin

and wounds were excised, minced with a razor blade, and suspended in 2.4 mL RPMI

1640 medium (Corning). Digestion enzymes were obtained from Miltenyi Biotec Multi

Tissue Dissociation Kit 1 and resuspended according to manufacturer instructions. 100

µL Enzyme D, 50 µL Enzyme R, 12.5 µL Enzyme A were added to each tissue sample in

RPMI. Tissues were mechanically dissociated using gentleMACS™ C Tubes (Miltenyi

Biotec; protocol Multi H) for 30 sec, incubated on a rocking shaker at 37ᵒC for 30 min,

followed by an additional 30 sec of mechanical dissociation and 30 min incubation at 37˚C

while rocking. The digestion mixture was strained through a 70 µm filter, and the single

cells were washed with RPMI and spun down at 300 x g for 5 min. The cell pellet was

resuspended in Ammonium-Chloride-Potassium red blood cell lysis buffer (150 mM

NH_4Cl, 10 mM $KHCO_3$, 0.1 mM Na_2EDTA) for 1 min at room temperature and diluted in

RPMI. Following centrifugation, the cell pellet was resuspended in MACS staining buffer (1X PBS, 2 mM EDTA, 0.5% BSA) to generate single cell suspensions. Blood was obtained from right ventricle cardiac puncture. Approximately 200 µL were taken from each animal and mixed a small amount of heparin suspended in PBS. Ammonium-Chloride-Potassium red blood cell lysis buffer was added to blood samples to a final volume of 1 mL, and incubated for 3 min. Samples were then spun down at 300 x g for 5 min and the resulting pellet was washed again in red blood cell lysis buffer with a repeat spin. The cell pellet was resuspended in MACS staining buffer to generate single cell suspensions as noted above.

Antibody staining of single cell suspensions for flow cytometry analysis

Antibody staining methods and panels were performed as previously described (47). Fixable Viability Dye eFluor™ 506 (ThermoFisher; 1:800 dilution) in PBS was added to single cell suspensions and allowed to incubate for 15 min at room temperature. Conjugated antibodies listed in Table S4 were then added to wound and blood single cell suspensions along with anti-mouse CD16/32 (ThermoFisher, Fc block, clone 93) for 30 min at room temperature. To analyze human neutrophils, antibodies against surface markers listed in Table S4 were added with human FcR Blocking Reagent (Miltenyi Biotec). Stained neutrophils were then washed and fixed using Foxp3/Transcription Factor Staining Buffer Set (eBioscience) according to the manufacturer's protocol. Cells were analyzed on the LSRFortessa (BD Biosciences) flow cytometer. Data were analyzed with BD FlowJo software version 10.10.0. Single-stained beads (Versacomp Antibody Capture Bead Kit; Beckman Coulter) were used as compensation controls. A detailed

gating strategy is included in Figure S2 for wound and blood and Figure S3 for neutrophil

degranulation.

Multispectral Immunohistochemistry and Image Analysis

Diabetic mice were wounded and infected with either WT CJB111 expressing GFP or

mock infected with PBS as described above. After 48h of infection, animals were

sacrificed and wounds excised and sent to the Human Immune Monitoring Shared

Resource (HIMSR) at the University of Colorado Anschutz Medical Center. Multispectral

immunohistochemistry (mIHC) analyses were performed using Vectra Automated

Quantitative Pathology Systems (Akoya Biosciences) as described previously (106).

Tissues were formalin-fixed, paraffin embedded and sectioned onto slides. Whole tissues

were analyzed with the following antibodies; MsCD11b (Novus, NB110-89474), Msly6G

(CST, 87048S), HuCitH3 (abcam, ab5103), MsGFP/YFP (CST, 2956S), msCD11c (CST,

97585S), MsCD45 (CST, 70257S), MsEpCAM (CST, 93790S), MsF480 (CST, 30325S),

and DAPI. All slides were deidentified and imaged by the Anschutz Medical Campus

Human Immune Monitoring Shared Resource (HIMSR) core (RRID:SCR_021985) on

Akoya Biosciences PhenoImager HT scanner. Regions of interest (ROI) were selected,

and multispectral images were collected with the 20× objective. A training set of nine

representative images was used to train the analyses algorithms for tissue and cell

segmentation and phenotyping using in Form software (Akoya Biosciences).

Representative autofluorescence was measured on an unstained control slide and

subtracted from study slides.

Primary human neutrophil killing.

Primary human neutrophils were isolated from healthy human donors (female $n = 2$, male $n = 2$) per the COMIRB protocol number 17-1926 as previously described (107–109). Neutrophils were resuspended in RPMI (Gibco, no phenol red) with 22 mM glucose to mimic diabetes (46). Mid-exponential phase GBS were normalized to 1 x 10^8 CFU/mL in Hanks' Balanced Salt Solution (HBSS, Gibco), and 1mL GBS was opsonized in 10% pooled human serum (Complement Tech) at 37˚C with rotation for 30 min. Opsonized GBS were washed once with HBSS before being resuspended in 1 mL of 4 x 10^6 neutrophils (MOI of 25) and split into two Eppendorf tubes containing 500 µL each. One tube was incubated at 37˚C, with rotation, for 1 h and the second tube used for time 0 h counts. At each time point, 500 µL of 0.05% Saponin (Sigma) was added to the tube and vortex vigorously to lyse neutrophils. 20 µL was removed in duplicate and serial diluted and 50 µL was spread plate on THA containing plates and incubate at 37˚C overnight for CFU enumeration. Percent survival was calculated as the ratio of CFU at 1 h verses 0 h and normalized to the no neutrophil control (RPMI only tubes) set to 100%.

HL60 Opsonophagocytic killing.

HL60 opsonophagocytic killing assays were performed as previously described (43). Briefly, HL60 cells purchased from ATCC (ATCC CCL-240) were differentiated with 1.25% DMSO (Sigma) in RPMI for 4 days. Differentiated HL60 cells were harvested and checked for > 65% viability before starting the assay in accordance with a well-

characterized protocol. In a 96 well plate (Corning), 50 µL containing ~1 x10^3 CFU of mid-

exponential phase GBS, normalized in HBC buffer (1X HBSS, 0.5% BSA, and 2.2 mM

CaCl$_2$, pH 7.2) was added. 50 µL of either 20% normal human serum (Complement Tech)

or 20% heat-killed human serum (60˚C for >30 min) in HBC buffer were added to each

well and incubated at 37˚C, with shaking, for 15 min. 100 µL of HL60 cells (~5 x10^5) were

added to each well and incubated at 37˚C with shaking, MOI ~0.001, as previously

described. Over a 3h time course, 10 µL's were removed and indicated times, serial

diluted and plated on THB agar for enumeration. Assays were performed in biological

triplicate with technical duplicates per replicate. Percent survival was calculated as the

ratio of CFU in normal serum versus CFU in heat-killed serum, and no HL60 control wells

were set to 100%.

Primary human neutrophil degranulation

716 To measure degranulation in primary human neutrophils, neutrophils were isolated as

717 described above and infected with opsonized GBS at an MOI of 25 in 22 mM glucose as

718 described above for 30 min at 37˚C in a 96 well plate. Antibody staining and flow

719 cytometry analysis is described above using known markers of degranulation (85, 110).

Cationic antimicrobial peptide killing

722 mCRAMP killing assays were performed with concentrations as previously described

723 (43). A total of 2-4 x 10^5 CFU of mid-exponential phase cultures were exposed to 20 µM

724   mCRAMP (AnaSpec) in THB for 30 min at 37˚C. Dilutions were plated for CFU

725   enumeration.

Reactive oxygen species susceptibility and 8-OHdG ELISA

Overnight cultures were normalized to 1 x 10^8 CFU/mL and 2 mL of bacteria were

exposed to 0.08% hypochlorite (Clorox) or 100 µg of hydrogen peroxide (Millipore,

386790) for 1 h at 37˚C before 10 µL was removed for serial dilution and plating for CFU

on THA plates. The rest of the bacteria exposed to hydrogen peroxide were pelleted and

stored at -20 ˚C for 8-OHdG Quantification via OxiSelect Oxidative DNA Damage ELISA

Kit (Cell Biolabs, STA-320) per manufacturer protocol. Biological triplicate was performed.

Anisotropy

The small molecule probe 1,6-diphenyl-1,3,5-hexatriene (Sigma; DPH) was used to

evaluate membrane fluidity as previously described (111, 112). Briefly, pellets from

overnight cultures were washed with phosphate buffered saline (PBS) twice and

resuspended in fresh PBS to ~ 1 x 10^8 CFU/mL in 3 mL of PBS and incubated with 5 µM

of DPH for 15 min at 37°C and 200 rpm and samples were kept at 37˚C using a water

bath until analyzed. Anisotropy measurements were taken on a Jobin Yvon FluoroMax-3

spectrofluorometer with polarizers in the L geometry at the Biophysics Shared Research

Facility at the University of Colorado Anschutz Medical Campus. DPH was excited at 360

nm and the fluorescence intensity was monitored at 430 nm. Samples were maintained

at 37°C using the temperature-controlled sample compartment of the spectrofluorometer.

Measurements were performed on biological quadruplicate. The anisotropy of DPH was

calculated from the differences in the fluorescence intensity detected with the polarizers

in the vertical and horizontal orientations and data is reported in arbitrary units (AU).

Biological quadruplicate was performed.

Statistical analyses.

All statistics were performed using GraphPad Prism software V10. All relevant statistical

analyses are indicated in Figure legends.

Ethics statement. Animal experiments were approved by the Institutional Animal Care

and Use Committee at University of Colorado Anschutz Medical Campus protocol no.

00987 and were performed using accepted veterinary standards. The University of

Colorado Anschutz Medical Campus is AAALAC accredited, and its facilities meet and

adhere to the standards in the Guide for the Care and Use of Laboratory Animals.

Data availability

All sequencing reads have been deposited into NCBI SRA under BioProject Accession

PRJNA1232139

Figure 1. Neutrophils are abundant in the diabetic wound during infection. A) Wound immune cell populations measured by flow cytometry. B) A significant increase of neutrophils (left) and significant decrease of macrophages (right) as a percent of CD45+ immune cells is observed in GBS infected diabetic wounds at 48 h compared to uninfected diabetic wounds. C) Multispectral immunohistochemistry microscopy images of diabetic wound tissues mock or GBS infected. Scale bar 800 μm. D) White boxed region of interested showing CD45 pathology, GFP expressing GBS in the wound, Ly6G+ neutrophil marker, F4/80+ macrophage marker, and merged image. Significance determined by B) two-tailed T-Test; **p<.01; ***p<.001.

Figure 2. Essential GBS Genes During Diabetic Wound Infection. A) Experimental set-up for Tn-seq. B) Circos plot showing from outer ringer to inner ring; dark grey, genome length in Kb; maroon, negative Log2Fold Change (L2FC); red, positive L2FC; statistically significant genes (adj. p-value ≤ 0.05) shown in green and non-significant genes in light grey. C) COG pathway analysis of percent of statistically significant genes in each COG for both over- (red) and under- (maroon) represented genes. (Numbers) indicate the total number of genes in each pathway. D) Linear plot of significantly over (red) and underrepresented (maroon) genes ordered by genomic position. E) Glycolipid biosynthesis and LTA decoration pathway in GBS.

Figure 3. Glycolipid biosynthesis enzymes and LTA alanylation are necessary for epithelial cell invasion and survival in diabetic wound infection. *In vitro* assays for A) total cell associated (adherence) and B) invasion of GBS mutants N/TERTs indicate GBS glycolipids and LTA alanylation are dispensable for adhesion but are important for invasion of skin epithelial cells. Each assay was performed in biological triplicate, with three technical replicates per assay. Mean and SEM indicated. Diabetic mice were inoculated with 1 x 10^7 CFU of GBS WT, Δ*iagB*, Δ*iagA*, Δ*mprF*, and Δ*dltA* isogeneic mutants and bacterial burdens were assessed after 48 h in the C) wound and D) kidneys. Significantly less CFU of all four mutants are recovered from the wound and kidneys of infected mice. E) % of mice in each infection group with detectable CFU in kidneys. F) Chemokine CXCL-1 and G) myeloperoxidase protein production measured by ELISA from wound tissues. Median indicated. GBS WT; *n* = 15, Δ*iagB; n* = 12, Δ*iagA*; *n* = 11, Δ*mprF*; *n* = 16, and Δ*dltA;* n = 11. Significance determined by A,B) One-way ANOVA with Fisher LSD and C,D,E,F) Kruskal-Wallis with Dunn's multiple comparisons test ; *p<.05, **p<.01, ***p<.001, ****p<.0001. ns; not significant.

808
809

Figure 4. GBS susceptibility to neutrophil killing factors. A) Primary human neutrophil killing of GBS WT, GBSΔ*iagB,* pJIagB, GBSΔ*dltA*, and pJDltA in high glucose infected at MOI 25 for 1 h. % Survival calculated as the ratio of CFU with neutrophils versus CFU without neutrophils. B) HL60-neutrophil like cell killing of GBS WT, Δ*iagB* and Δ*dltA* at MOI 0.001 for 3 h. % Survival calculated as the ratio of CFU in normal serum versus CFU in heat-killed serum in the presence of HL60 cells. C) Percent survival of GBS WT, GBSΔ*iagB*, pJIagB, GBSΔ*dltA*, and pJDltA to D) mouse cathelicidin mCRAMP after 30 min and D) hypochlorite (HOCl). E) Percent survival of GBS WT, GBSΔ*iagB*, and GBSΔ*dltA*, to 100 µg H_2O_2 after 1 hr and C) Oxidative DNA damage caused by H_2O_2 measured via ELISA. A) Mean and SEM, B=F) Mean and SD. Significance determined by A) Mann-Whitney B-F) One-way ANOVA with Fishers LSD test; *p<.05, **p<.01, ****p<.0001. Each assay was performed in at least biological triplicate.

824
Figure 5. GBS glycolipids and LTA alanylation is not required for degranulation in
primary human neutrophils. Neutrophil degranulation measured in primary human
neutrophils conditioned in high glucose exposed to opsonized GBS by flow cytometry
after 30 min. A) Azurophilic/primary (CD63$^+$) granules: left; representative histogram of
counts of CD63$^+$ neutrophils and right; median fluorescent intensity (MFI) of CD63.
Specific/secondary granules B) CD66b$^+$. A significant increase in both primary and
secondary degranulation compared to uninfected neutrophils is observed. A-C) Mean and
SEM. A-C) Significance determined by One-way ANOVA with Fisher's LSD test; *p<.05,
**p<.01, ***p<.001, ****p<.0001, ns; not significant. Assay was performed in at biological
quadruplicate.

835
836

837
838

Figure 6. Neutrophils play an important role in combating GBS in the diabetic wound. Control (anti-IgG2a antibody, closed circles) and neutrophil depleted diabetic mice (anti-Ly6G antibody, open circles) were infected with 1 x 10^7 GBS WT, $\Delta iagB$, and $\Delta dltA$ and bacterial burdens were assessed at 48h in the A) wound and B) kidney. Significantly higher bacterial burdens were recovered in the wound and kidney in depleted mice compared to control mice. C) % of mice in each group with CFU in the kidney. Control: GBS WT; $n = 11$, $\Delta iagB$; $n = 13$, and $\Delta dltA$; n = 12. Depleted: GBS WT; $n = 12$, $\Delta iagB$; $n = 13$, and $\Delta dltA$; n = 10. Significance determined by Kruskal-Wallis test with Dunn's multiple comparisons; *p<.05, **p<.01, ****p<.0001.

848
849

850
851
Figure 7. GBS glycolipids and LTA alanylation promote diabetic wound colonization. GBS infects the diabetic wound where it encounters a large population of neutrophils in the wound. Neutrophils are recruited to the wound to control the infection. The diabetic wound is rich in multiple antimicrobial molecules such as cationic antimicrobial peptides (CAMP) and reactive oxygen species (ROS). GBS∆*dltA* is unable to modify lipoteichoic acid (LTA) and is more susceptible to both neutrophil killing and CAMPs compared to GBS WT, whereas *GBS∆iagB* lacking all glycolipids is hypersusceptible to neutrophil killing, CAMPs and ROS attack.

Table 1. Significantly underrepresented genes in the diabetic wound.

Locus Tag	Gene	Description	L2FC	Adj. P-Value[#]	Reference
Virulence Factors					
ID870_059 35	*cylA*	ABC transporter ATP-binding protein	-3.66	0	(41, 53, 54)
ID870_059 30	*cylB*	ABC transporter permease	-2.03	0	
ID870_059 10	*cylJ*	cylJ protein	-0.92	0	
ID870_059 25	*cylE*	cylE protein	-0.9	0	
ID870_059 50	*cylG*	3-oxoacyl-ACP reductase	-0.88	0.0008	
ID870_059 05	*cylK*	CylK protein	-0.78	0	
ID870_059 15	*cylI*	beta-ketoacyl-[acyl-carrier-protein] synthase family protein	-0.61	0	
ID870_053 40	*sodA*	superoxide dismutase	-1.91	0	(55)
ID870_052 45	*prsA*	peptidylprolyl isomerase	-1.52	0	
ID870_033 75	*hylB*	hyaluronate lyase	-0.79	0	(56)
ID870_073 70	*perR*	peroxide-responsive transcriptional repressor PerR	-0.61	0.04178	(57)
ID870_046 25	*srtA*	class A sortase	-1.06	0	(58)
ID870_041 75	*esxA2*	WXG100 family type VII secretion target	-0.85	0.03819	(59)
ID870_041 90	*esaB*	EsaB/YukD family protein	-1.35	0.03125	
ID870_042 20	-	hypothetical protein - transmembrane	-0.73	0.04834	
ID870_042 65	-	hypothetical protein	-0.64	0.00153	
ID870_042 90	-	T7SS lipoprotein protein	-0.73	0	
TCS systems					
ID870_056 70	TCS8 _S/*vic K*	cell wall metabolism sensor histidine kinase VicK	-2	0	(60, 61)

ID870_030 10	TCS1 2_S	HAMP domain-containing histidine kinase	-1.44	0	(60, 61)
ID870_044 55	*ciaR*	response regulator transcription factor	-1.31	0	(62, 63)
ID870_044 60	*ciaH*	HAMP domain-containing histidine kinase	-1.02	0	
ID870_104 25	TCS1 9_R	response regulator transcription factor	-0.86	0.00766	(60, 61)
ID870_057 10	TCS7 _R	response regulator transcription factor	-0.59	0.03302	(60, 61)
Capsule/Cell wall/membrane					
ID870_007 15	*dltX*	teichoic acid D-Ala incorporation-associated protein	-1.68	0.03105	(64, 65)
ID870_007 25	*dltB*	D-alanyl-lipoteichoic acid biosynthesis protein	-2.6	0	
ID870_007 30	*dltC*	D-alanine--poly(phosphoribitol) ligase subunit	-1.73	0.01571	
ID870_034 80	*cpsY*	LysR family transcriptional regulator	-2.83	0	(66– 68)
ID870_035 50	*cpsL*	oligosaccharide flippase family protein	-1.78	0	
ID870_035 60	*neuC*	UDP-N-acetylglucosamine 2-epimerase	-0.58	0.01066	
ID870_035 70	*neuA*	N-acylneuraminate cytidylyltransferase	-0.72	0.00463	
ID870_054 55	*pbp2b*	penicillin-binding protein 2	-1.68	0	
ID870_086 25	*pbp1B*	penicillin-binding protein	-0.85	0	
ID870_087 10	*pbp3*	D-alanyl-D-alanine carboxypeptidase	-0.98	0	
ID870_057 25	*iagA*	glycosyltransferase	-4.87	0	(45)
ID870_075 65	*accD*	acetyl-CoA carboxylase carboxyltransferase subunit beta	-1	0.04354	
ID870_075 75	*fabZ*	3-hydroxyacyl-ACP dehydratase	-1.85	0.03968	
ID870_076 10	*fabH*	ketoacyl-ACP synthase III	-1.51	0	
ID870_087 55	*uppP*	undecaprenyl-diphosphate phosphatase	-0.73	0	
ID870_100 50	*mprF*	bifunctional lysylphosphatidylglycerol flippase/synthetase MprF	-0.73	0	(44)
Metabolism					

ID	Gene	Description	Value	p-value	Ref
ID870_00505	*ahpC*	peroxiredoxin	-2.32	0	
ID870_00585	*purA*	adenylosuccinate synthase	-6	0	(69)
ID870_00850	*glnR*	MerR family transcriptional regulator	-1.67	0.01484	
ID870_01300	*manA*	mannose-6-phosphate isomerase class I	-0.79	0.04253	
ID870_02315	*glnQ*	amino acid ABC transporter ATP-binding protein	-1.47	0	(70)
ID870_02320	*glnP*	ABC transporter substrate-binding protein/permease	-1.17	0	
ID870_04130	*pyrF*	orotidine-5'-phosphate decarboxylase	-3.4	0	
ID870_04135	*pyrE*	orotate phosphoribosyltransferase	-1.54	0	
ID870_04145	*pyrB*	aspartate carbamoyltransferase catalytic subunit	-1.86	0	
ID870_04150	*carA*	glutamine-hydrolyzing carbamoyl-phosphate synthase small subunit	-1.13	0.00215	
ID870_04155	*carB*	carbamoyl-phosphate synthase large subunit	-1.28	0	
ID870_04545	*guaA*	glutamine-hydrolyzing GMP synthase	-6.78	0	(69, 71)
ID870_05480	*ppc*	phosphoenolpyruvate carboxylase	-2.51	0	
ID870_07185	*nrdE*	class 1b ribonucleoside-diphosphate reductase subunit alpha	-1.42	0	
ID870_07190	*nrdI*	class Ib ribonucleoside-diphosphate reductase assembly flavoprotein NrdI	-1.14	0	
ID870_07195	*nrdF*	class 1b ribonucleoside-diphosphate reductase subunit beta	-2.16	0	
ID870_07535	*manN*	PTS mannose/fructose/sorbose transporter family subunit IID	-0.98	0	
ID870_07540	*manO*	DUF956 family protein	-1.52	0.04152	
ID870_09305	*purB*	adenylosuccinate lyase	-7.52	0	(69)
ID870_09315	*purK*	5-(carboxyamino)imidazole ribonucleotide synthase	-1.46	0.00522	
ID870_09320	*purE*	5-(carboxyamino)imidazole ribonucleotide mutase	-2.44	0	

ID870_093 25	*purD*	phosphoribosylamine--glycine ligase	-3.05	0	
ID870_093 90	*purH*	Bifunctional phosphoribosylaminoimidazoleca rboxamide formyltransferase/IMP cyclohydrolase	-3.74	0	
ID870_094 00	*purN*	phosphoribosylglycinamide formyltransferase	-3.43	0	
ID870_094 05	*purM*	phosphoribosylformylglycinamidi ne cyclo-ligase	-2.91	0	
ID870_094 10	*purF*	amidophosphoribosyltransferase	-1.54	0	
ID870_094 20	*purC*	Phosphoribosylaminoimidazolesu ccinocarboxamide synthase	-4.15	0	
ID870_098 00	*guaB*	IMP dehydrogenase	-3.11	0	(69, 71)
ID870_102 70	*nrdD*	anaerobic ribonucleoside-triphosphate reductase	-1.16	0	
ID870_102 90	*nrdG*	anaerobic ribonucleoside-triphosphate reductase activating protein	-1.2	0.00279	

861 # Adj. P-Value <0.0001 is assigned 0 by TRANSIT.

Supplemental Figures

870 **Figure S1: Wound and blood immune panel**

871 **Figure S2. Tn-seq metrics**

872 **Figure S3. Membrane fluidity and primary human neutrophil degranulation.**

873 **Figure S4. Neutrophil depletion of diabetic mice.**

875
876
877 **Supplemental tables:**

878
879 **Table S1. Tn-seq dataset (xlsx)**

880 **Table S2. Strains**

881 **Table S3. Primers**

882 **Table S4. Flow Cytometry antibodies**

883
884
885

886
887
Figure S1. Wound and blood immune panel. A) Density plots of murine wound and
blood single cell suspensions stained with antibody panel. Gates identifying cell
populations of interest are labeled. The antibody panel identifies neutrophil, Cd11b$^-$
CD11c$^-$(containing lymphocytes), SSC^{lo}/MHCII$^-$ (containing monocytes and NK cells),
dendritic cell, and CD24$^+$ (containing eosinophils) populations. % immune cells of CD45$^+$
live cells in diabetic mouse wounds infected with CJB111 or mock control (uninfected)
measured by flow cytometry B) live cells of singlets, C) CD45 cells of singlets, D)
neutrophils of singlets, E) Cd11b$^-$ CD11c$^-$(containing lymphocytes), F) SSC^{lo}/MHCII$^-$
(containing monocytes and NK cells) G) CD24$^+$ cells and H) Dendritic cells. I) Summary
of immune cell populations in the blood of diabetic mice with wounds either infected with
CJB111 or uninfected mock control. Uninfected $n = 7$, CJB111 $n = 7$.   Mean and SEM
indicated. Significance determined by: B-I) Two-tailed students T-Test; *p<.05, ***p<.001,
ns; not significant.
901

903 **Figure S2. Tn-seq metrics.** A) Input CFU (green squares) inoculated into the wound and
904 output CFU (black circles, $n = 4$ mice per library) for each biological replicate at time of
905 euthanasia. The number of B) reads mapped to the genome in each library, genomic TA
906 sites insertions were mapped to and the saturation levels of the transposon libraries.
907 Mean and SD indicated. Each dot represents a biological library.
908

909
Figure S3. Membrane fluidity and primary human neutrophil degranulation. A)
Membrane fluidity measure via anisotropy. AU, Arbitrary Units. Mean and SD. B) Density
plots of isolated human neutrophils stained with antibody panel. Neutrophils were
confirmed by the expression of Live CD45$^+$, CD15$^+$, CD16$^+$, CD14$^-$ singlets. Median
fluorescent intensity (MFI) of C) specific/secondary granules (CD15$^+$) and D)
gelatinase/tertiary granule (CD11b$^+$ and CD18$^+$) degranulation measured in primary
human neutrophils exposed to opsonized GBS by flow cytometry after 30 min. C,D) Mean
and SEM. Significance determined by A) Kruskal-Wallis test with Dunn's multiple
comparisons and C,D) One-way ANOVA with Fishers LSD test,; *p<.05, **p<.01, ns; not
significant. Each assay was performed in at least biological triplicate.

920
921
Figure S4. Neutrophil depletion of diabetic mice. Top: Flow plots of CD45$^+$ live
neutrophils (CD11b$^+$ and Ly6G$^+$) in diabetic mice injected with either α-IgG2a antibody
(control) or α-Ly6G antibody (depleted); Bottom: Percent of neutrophils (Ly6G$^+$)
detected. Mean and SEM. $N = 3$ per group.

926
927

933 **Table S2. Strains and plasmids used in this study.**

Organism	Strain	Description	Ref
S. agalactiae	CJB111	Wilt-type *S. agalactiae* strain, serotype V	(51)
	CJB111Δ*iagB*	*iagB* (ID870_05720) deletion strain	(43)
	CJB111Δ*iagB*(pDCErm)	Empty vector control strain	(43)
	CJB111Δ*iagB*(pJIagB)	Expresses GBS *iagB* in pDCErm	(43)
	CJB111Δ*iagA*	*iagA* (ID870_05725) deletion strain	(43)
	CJB111Δ*mprF*	*mprF* (ID870_10050) deletion strain	(43)
	CJB111Δ*dltA*	*dltA* (ID870_00720) deletion strain	This work
	CJB111Δ*dltA*(pDCErm)	Empty vector control strain	This work
	CJB111Δ*dltA*(pJDltA)	Expresses GBS *dltA* in pDCErm	This work
	CJB111(pDEST-GFP)	Expresses GFP from pDESTErm	This work
E. coli	MC1061	Plasmid cloning host; F⁻, araD139, Δ(araABC-leu)7696, Δ(lac)X74, galU, galK, hsdR2, ($r_K^-m_K^+$), mcrB1, rpsL, (Str^r)	(113)
	MC1061(pJDltA)	Expresses CJB111 *dltA* from $P_{tetM/erm}$ in pDCErm	This work
	MC1061(pDltAKO)	CJB111 *dltA* knockout plasmid	This work

Plasmid	Description	
pMBSacB	Allelic exchange plasmid for *S. agalactiae*. Confers erythromycin resistance and sucrose sensitivity	(101)
pDltAKO	Knockout plasmid containing ~1.5 kb sequence flanking ID870_00720	This work
pDCErm	Constitutive expression vector for streptococcus from $P_{tetM/erm}$	(102)
pJDltA	pDCErm expressing CJB111 *dltA* (ID870_00720)	This work
pJIagB	pDCErm expressing CJB *iagB* (ID870_05720)	(43)
pDEST-GFP	pDESTErm expressing GFP	(114)

936 **Table S3. Primers used in this study.**

Primer	5' – 3' sequence	Use
AUpF_XhoI	ACGTCACTCGAGGACCTTCAGACAATTC	Amplifies upstream of DltA
AUpR_SacII	ACGTCACCGCGGTTGTACCATCTCCTAG	Amplifies upstream of DltA
ADwnF_BamHI	ACGTCAGGATCCGTGATACAGGCATTTTTAG	Amplifies downstream of DltA
AdwnR_PstI	ACGTCACTGCAGGGCTTTCCAACTCAACA	Amplifies downstream of DltA
Aout_F	CACATGAGCTAGAATC	Upstream primer for sequencing
Aout_R	TAGACGTTTAAGCATTGC	Downstream primer for sequencing
Afor	ACGTCACCGCGGATGATACATGATATGATTAAAAC	Amplification of dltA for complementation
Arev	ACGTCAGGATCCTCACTTGTTTACCTCG	Amplification of dltA for complementation

Table S4. Flow cytometry antibodies.

Marker	Fluorophore	Clone	Staining Concentration (µg/mL)	Company	Catalog #
colspan=6 **Murine Diabetic Wound and Blood Panel**
Viability Dye	eF506	--	[1:800 dilution]	Invitrogen	65-0866-18
CD45	BUV395	30-F11	0.50	BD Biosciences	564279
Ly6G	APC	1A8	0.50	Invitrogen	17-9668-82
CD11b	FITC	M1/70	2.50	Invitrogen	11-0112-85
CD11c	PE-Cy	N418	1.00	Biolegend	117317
CD64	BV650	X54-5/7.1	4.00	BD Biosciences	740622
F4/80	BV785	BM8	1.00	Biolegend	123141
Ly6C	BV421	AL-21	0.50	BD Biosciences	562727
MHCII	BUV737	M5/114 15.2	0.10	BD Biosciences	748845
CD24	PerCP-Cy5.5	M1/69	0.20	Biolegend	101824
colspan=6 **Human Neutrophil Degranulation Panel**
Viability	eF506	--	[1:800 dilution]	Invitrogen	65-0866-18
CD45	BUV395	HI30	0.125	Invitrogen	363-0459-41
CD14	BV421	61D3	0.50	Invitrogen	404-0149-41
CD15	APC	MMA	0.125	Invitrogen	17-0158-41
CD16	PE/Cy7	CB16	0.030	Invitrogen	25-0168-41
CD11b	FITC	ICRF44	0.50	BD Biosciences	562793
CD66b	PE	G10F5	0.50	Biolegend	305105
CD63	BV785	H5C6	0.50	Biolegend	353043
CD18	APC/Fire750	TS1/18	1.00	Biolegend	302125
Viability	eF506	--	[1:800 dilution]	Invitrogen	65-0866-18
CD45	BUV395	HI30	0.125	Invitrogen	363-0459-41
CD14	BV421	61D3	0.50	Invitrogen	404-0149-41

940
941

References

1. Sun H, Saeedi P, Karuranga S, Pinkepank M, Ogurtsova K, Duncan BB, Stein C, Basit A, Chan JCN, Mbanya JC, Pavkov ME, Ramachandaran A, Wild SH, James S, Herman WH, Zhang P, Bommer C, Kuo S, Boyko EJ, Magliano DJ. 2022. IDF Diabetes Atlas: Global, regional and country-level diabetes prevalence estimates for 2021 and projections for 2045. Diabetes Res Clin Pract 183:109119.

2. Uccioli L, Izzo V, Meloni M, Vainieri E, Specialist M, Ruotolo V, Specialist M, Giurato L. 2014. Non-healing foot ulcers in diabetic patients: general and local interfering conditions and management options with advanced wound dressings. Proc Int Surg Wound Forum 24:35–42.

3. Hingorani A, Lamuraglia GM, Henke P, Meissner MH, Loretz L, Zinszer KM, Driver VR, Frykberg R, Carman TL, Marston W, Mills JL, Murad MH. 2016. The management of diabetic foot: A clinical practice guideline by the Society for Vascular Surgery in collaboration with the American Podiatric Medical Association and the Society for Vascular Medicine. J Vasc Surg 63:3S-21S.

4. Loots MAM, Lamme EN, Zeegelaar J, Mekkes JR, Bos JD, Middelkoop E. 1998. Differences in cellular infiltrate and extracellular matrix of chronic diabetic and venous ulcers versus acute wounds. J Invest Dermatol 111:850–857.

5. Eming SA, Krieg T, Davidson JM. 2007. Inflammation in wound repair: Molecular and cellular mechanisms. J Invest Dermatol 127:514–525.

6. Singer AJ, Clark RAF. 1999. Cutaneous Wound Healing. N Engl J Med.

7. Grice EA, Segre JA. 2012. Interaction of the microbiome with the innate immune response in chronic wounds. Adv Exp Med Biol 946:55–68.

8. Kalan LR, Meisel JS, Loesche MA, Horwinski J, Soaita I, Chen X, Uberoi A, Gardner SE, Grice EA. 2019. Strain- and Species-Level Variation in the Microbiome of Diabetic Wounds Is Associated with Clinical Outcomes and Therapeutic Efficacy. Cell Host Microbe 25:641-655.e5.

9. Keogh RA, Huyvaert S, Moore GD, Horswill AR, Doran KS. 2024. Virulence characteristics of Gram-positive bacteria isolated from diabetic foot ulcers. FEMS Microbes 5:1–10.

10. Rahim K, Saleha S, Zhu X, Huo L, Basit A, Franco OL. 2017. Bacterial Contribution in Chronicity of Wounds. Microb Ecol 73:710–721.

11. Gardner SE, Hillis SL, Heilmann K, Segre JA, Grice EA. 2013. The neuropathic diabetic foot ulcer microbiome is associated with clinical factors. Diabetes 62:923–930.

12. Jeffcoate WJ, Harding KG. 2003. Diabetic foot ulcers. Lancet 361:1545–1551.

13. Jeffcoate WJ, Price P, Harding KG. 2004. Wound healing and treatments for people with diabetic foot ulcer. Diabetes Metab Res Rev 20:78–89.

14. Perim MC, Borges J da C, Celeste SRC, Orsolin E de F, Mendes RR, Mendes
GO, Ferreira RL, Carreiro SC, Da Silva Pranchevicius MC. 2015. Aerobic
bacterial profile and antibiotic resistance in patients with diabetic foot infections.
Rev Soc Bras Med Trop 48:546–554.

15. Wolcott RD, Hanson JD, Rees EJ, Koenig LD, Phillips CD, Wolcott RA, Cox SB,
White JS. 2016. Analysis of the chronic wound microbiota of 2,963 patients by
16S rDNA pyrosequencing. Wound Repair Regen 24:163–174.

16. ALbeloushi A, Elbehiry A, Marzouk E, Zahran R, Edrees H. 2019. Highlight on
multidrug resistance of enterococcus faecalis recovered from diabetic foot
patients. J Pure Appl Microbiol 13:481–489.

17. Kalan LR, Brennan MB. 2019. The role of the microbiome in nonhealing diabetic
wounds. Ann N Y Acad Sci 1435:79–92.

18. Shook JC, Genito CJ, Darwitz BP, Tyson KJ, Velez AZ, Bridwell SK, Parsons JB,
Rowe SE, Marshall CW, Conlon BP, Thurlow LR. 2025. Diabetes potentiates the
emergence and expansion of antibiotic resistance 1591.

19. James GA, Swogger E, Wolcott R, Pulcini ED, Secor P, Sestrich J, Costerton JW,
Stewart PS. 2008. Biofilms in chronic wounds. Wound Repair Regen 16:37–44.

20. Watters C, Everett JA, Haley C, Clinton A, Rumbaugh KP. 2014. Insulin treatment
modulates the host immune system to enhance Pseudomonas aeruginosa wound
biofilms. Infect Immun 82:92–100.

21. Ch'ng J-H, Muthu M, Chong KKL, Wong JJ, Tan CAZ, Koh ZJS, Lopez D, Matysik
A, Nair ZJ, Barkham T, Wang Y, Kline KA. 2022. Heme cross-feeding can
augment Staphylococcus aureus and Enterococcus faecalis dual species biofilms.
ISME J https://doi.org/10.1038/s41396-022-01248-1.

22. Cowley ES, Chaves IZ, Osman F, Suen G, Anantharaman K, Hryckowian AJ.
2023. Determinants of Gastrointestinal Group B Streptococcus Carriage in Adults
https://doi.org/10.1101/2023.08.17.553755.

23. Wilkinson HW. 1978. Group b streptococcal infection in humans. Annu Rev
Microbiol 41–57.

24. Doran KS, Nizet V. 2004. Molecular pathogenesis of neonatal group B
streptococcal infection: No longer in its infancy. Mol Microbiol 54:23–31.

25. Kline KA, Lewis AL. 2016. Gram-positive uropathogens, polymicrobial urinary
tract infection, and the emerging microbiota of the urinary tract. Urin Tract Infect
Mol Pathog Clin Manag 459–502.

26. Romero R, Dey SK, Fisher SJ. 2014. Preterm labor: one syndrome, many causes.
Science 345:760–5.

27. Vornhagen J, Armistead B, Santana-Ufret V, Gendrin C, Merillat S, Coleman M,
Quach P, Boldenow E, Alishetti V, Leonhard-Melief C, Ngo LY, Whidbey C, Doran
KS, Curtis C, Adams Waldorf KM, Nance E, Rajagopal L. 2018. Group B

streptococcus exploits vaginal epithelial exfoliation for ascending infection. J Clin
Invest 128:1985–1999.

28. Wang H, Li N, Chivese T, Werfalli M, Sun H, Yuen L, Hoegfeldt CA, Elise Powe
C, Immanuel J, Karuranga S, Divakar H, Levitt NaA, Li C, Simmons D, Yang X.
2022. IDF Diabetes Atlas: Estimation of Global and Regional Gestational
Diabetes Mellitus Prevalence for 2021 by International Association of Diabetes in
Pregnancy Study Group's Criteria. Diabetes Res Clin Pract 183:109050.

29. Mercado-Evans V, Zulk JJ, Hameed ZA, Patras KA. 2024. Gestational diabetes
as a risk factor for GBS maternal rectovaginal colonization: a systematic review
and meta-analysis. BMC Pregnancy Childbirth 24:1–16.

30. Zhou Y, Wang L-Q, Yan Q, Lee C-C, Hsu M-H, Liao W-T, Zhang L, Chiu C-H.
2022. Genomic Analysis of Group B Streptococcus from Neonatal Sepsis Reveals
Clonal CC17 Expansion and Virulence- and Resistance-Associated Traits After
Intrapartum Antibiotic Prophylaxis. Clin Infect Dis 75:2153–2160.

31. Uggen E, Olaisen C, Valsø R, Gunnar L, Simonsen S, Magne R, Jensen B, Width
F, Bjørn G, Åsvold O, Ivar T, Nilsen L. 2024. Incidence of invasive infections with
Group B streptococcus in adults in Norway 1996 – 2019 : a nationwide registry -
based case – control study. Infection https://doi.org/10.1007/s15010-024-02210-3.

32. Edwards MS, Baker CJ. 2005. Group B streptococcal infections in elderly adults.
Clin Infect Dis 41:839–847.

33. Melin P. 2011. Neonatal group B streptococcal disease: from pathogenesis to
preventive strategies. Clin Microbiol Infect 17:1294–1303.

34. Nguyen LM, Omage JI, Noble K, McNew KL, Moore DJ, Aronoff DM, Doster RS.
2021. Group B streptococcal infection of the genitourinary tract in pregnant and
non-pregnant patients with diabetes mellitus: An immunocompromised host or
something more? Am J Reprod Immunol https://doi.org/10.1111/aji.13501.

35. NS Raja. 2007. Microbiology of diabetic foot infections in a teaching hospital in
Malaysia: a retrospective study of 194 cases. J Microbiol Immunol Infect 40:39.

36. Heravi FS, Zakrzewski M, Vickery K, Malone M, Hu H. 2020. Metatranscriptomic
Analysis Reveals Active Bacterial Communities in Diabetic Foot Infections. Front
Microbiol 11:1–10.

37. Citron DM, Goldstein EJC, Merriam CV, Lipsky BA, Abramson MA. 2007.
Bacteriology of moderate-to-severe diabetic foot infections and in vitro activity of
antimicrobial agents. J Clin Microbiol 45:2819–2828.

38. Wang Y, Zhang H, Ma G, Tian Z, Wang B. 2022. The contribution of intestinal
Streptococcus to the pathogenesis of diabetic foot ulcers: An analysis based on
16S rRNA sequencing. Int Wound J 19:1658–1668.

39. Blancas D, Santin M, Olmo M, Alcaide F, Carratala J, Gudiol F. 2004. Group B
Streptococcal Disease in Nonpregnant Adults: Incidence, Clinical Characteristics,

1059 and Outcome. Eur J Clin Microbiol Infect Dis 23:168–173.

40. Loesche M, Gardner SE, Kalan L, Horwinski J, Zheng Q, Hodkinson BP, Tyldsley AS, Franciscus CL, Hillis SL, Mehta S, Margolis DJ, Grice EA. 2017. Temporal Stability in Chronic Wound Microbiota Is Associated With Poor Healing. J Invest Dermatol 137:237–244.

41. Keogh RA, Haeberle A, Langouet-Astrie C, Kavanaugh JS, Moore G, Schmidt EP, Horswill AR, Doran KS. 2022. Group B Streptococcus Adaptation Promotes Survival in a Hyper-inflammatory Diabetic Wound Environment. Sci Adv 8.

42. Joyce LR, Doran KS. 2023. Gram-positive bacterial membrane lipids at the host-pathogen interface. PLoS Pathog 19:1–7.

43. Joyce LR, Kim S, Spencer BL, Christensen PM, Palmer KL, Guan Z, Siegenthaler JA, Doran KS. 2024. Streptococcus agalactiae glycolipids promote virulence by thwarting immune cell clearance. Sci Adv 10:1–15.

44. Joyce LR, Manzer HS, Mendonça J da C, Villarreal R, Nagao PE, Doran KS, Palmer KL, Guan Z. 2022. Identification of a novel cationic glycolipid in Streptococcus agalactiae that contributes to brain entry and meningitis. PLoS Biol 20:1–15.

45. Doran KS, Engelson EJ, Khosravi A, Maisey HC, Fedtke I, Equils O, Michelsen KS, Arditi M, Peschel A, Nizet V. 2005. Blood-brain barrier invasion by group B Streptococcus depends upon proper cell-surface anchoring of lipoteichoic acid. J Clin Invest 115:2499–2507.

46. Shrestha S, Lee Y-B, Lee H, Choi Y-K, Park B-Y, Kim M-J, Youn Y-J, Kim S-H, Jung S-J, Song D-K, Jin HK, Bae J-S, Lee I-K, Jeon J-H, Hong C-W. 2024. Diabetes Primes Neutrophils for Neutrophil Extracellular Trap Formation through Trained Immunity. Res (Washington, DC) 7:0365.

47. Spencer BL, Nguyen DT, Marroquin SM, Gapin L, O'Brien RL, Doran KS. 2025. Characterization of the Cellular Immune Response to Group B Streptococcal Vaginal Colonization. bioRxiv Prepr Serv Biol https://doi.org/10.1101/2025.01.29.635275.

48. Burcham LR, Akbari MS, Alhajjar N, Keogh RA, Radin JN, Kehl-Fie TE, Belew AT, El-Sayed NM, McIver KS, Doran KS. 2022. Genomic Analyses Identify Manganese Homeostasis as a Driver of Group B Streptococcal Vaginal Colonization. MBio 13.

49. Burcham LR, Le Breton Y, Radin JN, Spencer BL, Deng L, Hiron A, Ransom MR, Mendonça J da C, Belew AT, El-Sayed NM, McIver KS, Kehl-Fie TE, Doran KS. 2020. Identification of zinc-dependent mechanisms used by Group B Streptococcus to overcome calprotectin-mediated stress. MBio 11:1–18.

50. Akbari MS, Joyce LR, Spencer BL, Brady A, McIver KS, Doran KS. 2025. Identification of glyoxalase A in group B Streptococcus and its contribution to methylglyoxal tolerance and virulence. Infect Immun 0:e0054024.

1099   51.   Flores AR, Galloway-Peña J, Sahasrabhojane P, Saldaña M, Yao H, Su X, Ajami
1100   NJ, Holder ME, Petrosino JF, Thompson E, Ros IMY, Rosini R, Grandi G,
1101   Horstmann N, Teatero S, McGeer A, Fittipaldi N, Rappuoli R, Baker CJ,
1102   Shelburne SA. 2015. Sequence type 1 group B Streptococcus, an emerging
1103   cause of invasive disease in adults, evolves by small genetic changes. Proc Natl
1104   Acad Sci U S A 112:6431–6436.

1105   52.   Spencer BL, Chatterjee A, Duerkop BA, Baker CJ, Doran KS. 2021. Complete
1106   Genome Sequence of Neonatal Clinical Group B Streptococcal Isolate CJB111.
1107   Microbiol Resour Announc 10:14–16.

1108   53.   Pritzlaff CA, Chang JCW, Kuo SP, Tamura GS, Rubens CE, Nizet V. 2001.
1109   Genetic basis for the β-haemolytic/cytolytic activity of group B streptococcus. Mol
1110   Microbiol 39:236–248.

1111   54.   Armistead B, Herrero-Foncubierta P, Coleman M, Quach P, Whidbey C, Justicia
1112   J, Tapia R, Casares R, Millán A, Haidour A, Granger JR, Vornhagen J, Santana-
1113   Ufret V, Merillat S, Adams Waldorf K, Cuerva JM, Rajagopal L. 2020. Lipid
1114   analogs reveal features critical for hemolysis and diminish granadaene mediated
1115   Group B Streptococcus infection. Nat Commun 11:1502.

1116   55.   Rajagopal L. 2009. Understanding the regulation of Group B Streptococcal
1117   virulence factors. Future Microbiol 4:201–221.

1118   56.   Kolar SL, Kyme P, Tseng CW, Soliman A, Kaplan A, Liang J, Nizet V, Jiang D,
1119   Murali R, Arditi M, Underhill DM, Liu GY. 2015. Group B Streptococcus Evades
1120   Host Immunity by Degrading Hyaluronan. Cell Host Microbe 18:694–704.

1121   57.   Dubbs JM, Mongkolsuk S. 2012. Peroxide-sensing transcriptional regulators in
1122   bacteria. J Bacteriol 194:5495–5503.

1123   58.   Lalioui L, Pellegrini E, Dramsi S, Baptista M, Bourgeois N, Doucet-Populaire F,
1124   Rusniok C, Zouine M, Glaser P, Kunst F, Poyart C, Trieu-Cuot P. 2005. The SrtA
1125   sortase of Streptococcus agalactiae is required for cell wall anchoring of proteins
1126   containing the LPXTG motif, for adhesion to epithelial cells, and for colonization of
1127   the mouse intestine. Infect Immun 73:3342–3350.

1128   59.   Spencer BL, Job AM, Robertson CM, Hameed ZA, Serchejian C, Wiafe-Kwakye
1129   CS, Mendonça JC, Apolonio MA, Nagao PE, Neely MN, Korotkova N, Korotkov K
1130   V., Patras KA, Doran KS. 2023. Heterogeneity of the group B streptococcal type
1131   VII secretion system and influence on colonization of the female genital
1132   tractMolecular Microbiology.

1133   60.   Faralla C, Metruccio MM, De Chiara M, Mu R, Patras KA, Muzzi A, Grandi G,
1134   Margarit I, Doran KS, Janulczyk R. 2014. Analysis of two-component systems in
1135   group B streptococcus shows that RgfAC and the novel FspSR modulate
1136   virulence and bacterial fitness. MBio 5:1–12.

1137   61.   Thomas L, Cook L. 2020. Two-component signal transduction systems in the
1138   human pathogen streptococcus agalactiae. Infect Immun 88:1–16.

1139   62.   Quach D, Van Sorge NM, Kristian SA, Bryan JD, Shelver DW, Doran KS. 2009.
1140   The ciar response regulator in group b streptococcus promotes intracellular
1141   survival and resistance to innate immune defenses. J Bacteriol 191:2023–2032.

1142   63.   Spencer BL, Deng L, Patras KA, Burcham ZM, Sanches GF, Nagao PE, Doran
1143   KS. 2019. Cas9 contributes to group b streptococcal colonization and disease.
1144   Front Microbiol 10:1–15.

1145   64.   Poyart C, Pellegrini E, Marceau M, Baptista M, Jaubert F, Lamy MC, Trieu-Cuot
1146   P. 2003. Attenuated virulence of Streptococcus agalactiae deficient in D-alanyl-
1147   lipoteichoic acid is due to an increased susceptibility to defensins and phagocytic
1148   cells. Mol Microbiol 49:1615–1625.

1149   65.   Poyart C, Lamy MC, Boumaila C, Fiedler F, Trieu-Cuot P. 2001. Regulation of D-
1150   alanyl-lipoteichoic acid biosynthesis in Streptococcus agalactiae involves a novel
1151   two-component regulatory system. J Bacteriol 183:6324–6334.

1152   66.   Noble K, Lu J, Guevara MA, Doster RS, Chambers SA, Rogers LM, Moore RE,
1153   Spicer SK, Eastman AJ, Francis JD, Manning SD, Rajagopal L, Aronoff DM,
1154   Townsend SD, Gaddy JA. 2021. Group B Streptococcus cpsE Is Required for
1155   Serotype V Capsule Production and Aids in Biofilm Formation and Ascending
1156   Infection of the Reproductive Tract during Pregnancy. ACS Infect Dis 7:2686–
1157   2696.

1158   67.   Carlin AF, Chang YC, Areschoug T, Lindahl G, Hurtado-Ziola N, King CC, Varki
1159   A, Nizet V. 2009. Group B Streptococcus suppression of phagocyte functions by
1160   protein-mediated engagement of human Siglec-5. J Exp Med 206:1691–1699.

1161   68.   Uchiyama S, Sun J, Fukahori K, Ando N, Wu M, Schwarz F, Siddiqui SS, Varki A,
1162   Marth JD, Nizet V. 2019. Dual actions of group B Streptococcus capsular sialic
1163   acid provide resistance to platelet-mediated antimicrobial killing. Proc Natl Acad
1164   Sci U S A 116:7465–7470.

1165   69.   Rajagopal L, Vo A, Silvestroni A, Rubens CE. 2005. Regulation of purine
1166   biosynthesis by a eukaryotic-type kinase in Streptococcus agalactiae. Mol
1167   Microbiol 56:1329–1346.

1168   70.   Tamura GS, Nittayajarn A, Schoentag DL. 2002. A glutamine transport gene,
1169   glnQ, is required for fibronectin adherence and virulence of group B streptococci.
1170   Infect Immun 70:2877–2885.

1171   71.   Ipe DS, Sullivan MJ, Goh KGK, Hashimi SM, Munn AL, Ulett GC. 2021.
1172   Conserved bacterial de novo guanine biosynthesis pathway enables microbial
1173   survival and colonization in the environmental niche of the urinary tract. ISME J
1174   15:2158–2162.

1175   72.   Joyce LR, Guan Z, Palmer KL. 2021. Streptococcus pneumoniae, s. Pyogenes
1176   and s. agalactiae membrane phospholipid remodelling in response to human
1177   serum. Microbiol (United Kingdom) 167.

1178   73.   Eckert RL, Rorke EA. 1989. Molecular biology of keratinocyte differentiation.

1179   Environ Health Perspect 80:109–116.

1180 74. Smits JPH, Niehues H, Rikken G, Van Vlijmen-Willems IMJJ, Van De Zande
1181   GWHJF, Zeeuwen PLJM, Schalkwijk J, Van Den Bogaard EH. 2017. Immortalized
1182   N/TERT keratinocytes as an alternative cell source in 3D human epidermal
1183   models. Sci Rep 7:1–14.

1184 75. Patras KA, Wang NY, Fletcher EM, Cavaco CK, Jimenez A, Garg M, Fierer J,
1185   Sheen TR, Rajagopal L, Doran KS. 2013. Group B Streptococcus CovR
1186   regulation modulates host immune signalling pathways to promote vaginal
1187   colonization. Cell Microbiol 15:1154–1167.

1188 76. Akbari MS, Keogh RA, Radin JN, Sanchez-Rosario Y, Johnson MDL, Horswill AR,
1189   Kehl-Fie TE, Burcham LR, Doran KS. 2023. The impact of nutritional immunity on
1190   Group B streptococcal pathogenesis during wound infection. MBio 0.

1191 77. Amulic B, Cazalet C, Hayes GL, Metzler KD, Zychlinsky A. 2012. Neutrophil
1192   function: From mechanisms to disease. Annu Rev Immunol 30:459–489.

1193 78. Xu X, Håkansson L. 2002. Degranulation of primary and secondary granules in
1194   adherent human neutrophils. Scand J Immunol 55:178–188.

1195 79. Niyonsaba F, Madera L, Afacan N, Okumura K, Ogawa H, Hancock REW. 2013.
1196   The innate defense regulator peptides IDR-HH2, IDR-1002, and IDR-1018
1197   modulate human neutrophil functions. J Leukoc Biol 94:159–170.

1198 80. Zanetti M, Gennaro R, Romeo D. 1995. Cathelicidins: a novel protein family with a
1199   common proregion and a variable C-terminal antimicrobial domain. FEBS Lett
1200   374:1–5.

1201 81. Saar-Dover R, Bitler A, Nezer R, Shmuel-Galia L, Firon A, Shimoni E, Trieu-Cuot
1202   P, Shai Y. 2012. D-Alanylation of Lipoteichoic Acids Confers Resistance to
1203   Cationic Peptides in Group B Streptococcus by Increasing the Cell Wall Density.
1204   PLoS Pathog 8.

1205 82. Lee TH, Hofferek V, Separovic F, Reid GE, Aguilar MI. 2019. The role of bacterial
1206   lipid diversity and membrane properties in modulating antimicrobial peptide
1207   activity and drug resistance. Curr Opin Chem Biol 52:85–92.

1208 83. Lan CCE, Wu CS, Huang SM, Wu IH, Chen GS. 2013. High-Glucose environment
1209   enhanced oxidative stress and increased interleukin-8 secretion from
1210   keratinocytes. Diabetes 62:2530–2538.

1211 84. Ma XX, Liu QK, Kuai L, Ma X, Luo Y, Luo Y, Song JK, Fei XY, Jiang JS, Wang
1212   MX, Shen F, Ru Y, Li B. 2023. The role of neutrophils in diabetic ulcers and
1213   targeting therapeutic strategies. Int Immunopharmacol 124.

1214 85. Mollinedo F. 2019. Neutrophil Degranulation, Plasticity, and Cancer Metastasis.
1215   Trends Immunol 40:228–242.

1216 86. Wang L, Zhou X, Yin Y, Mai Y, Wang D, Zhang X. 2019. Hyperglycemia induces
1217   neutrophil extracellular traps formation through an NADPH oxidase-dependent

1218   pathway in diabetic retinopathy. Front Immunol 10:1–14.

87.   Ganesh S, Don KR, Priya.a J. 2020. The role of matrix metalloproteinases in wound healing. Int J Pharm Res 12:702–707.

88.   Gründling A, Schneewind O. 2007. Synthesis of glycerol phosphate lipoteichoic acid in Staphylococcus aureus. Proc Natl Acad Sci U S A 104:8478–8483.

89.   Percy MG, Gründling A. 2014. Lipoteichoic acid synthesis and function in gram-positive bacteria. Annu Rev Microbiol 68:81–100.

90.   Dowey R, Iqbal A, Heller SR, Sabroe I, Prince LR. 2021. A Bittersweet Response to Infection in Diabetes; Targeting Neutrophils to Modify Inflammation and Improve Host Immunity. Front Immunol 12:1–21.

91.   Shrestha S, Lee Y-B, Lee H, Choi Y-K, Park B-Y, Kim M-J, Youn Y-J, Kim S-H, Jung S-J, Song D-K, Jin HK, Bae J-S, Lee I-K, Jeon J-H, Hong C-W. 2024. Diabetes Primes Neutrophils for Neutrophil Extracellular Trap Formation through Trained Immunity. Research 7:1–14.

92.   Derré-Bobillot A, Cortes-Perez NG, Yamamoto Y, Kharrat P, Couvé E, Da Cunha V, Decker P, Boissier MC, Escartin F, Cesselin B, Langella P, Bermúdez-Humarán LG, Gaudu P. 2013. Nuclease A (Gbs0661), an extracellular nuclease of Streptococcus agalactiae, attacks the neutrophil extracellular traps and is needed for full virulence. Mol Microbiol 89:518–531.

93.   Li D, Pi W, Sun Z, Liu X, Jiang J. 2022. Ferroptosis and its role in cardiomyopathy. Biomed Pharmacother 153:113279.

94.   Goncheva MI, Chin D, Heinrichs DE. 2022. Nucleotide biosynthesis: the base of bacterial pathogenesis. Trends Microbiol 30:793–804.

95.   Tan CAZ, Chong KKL, Yeong DYX, Ng CHM, Ismail MH, Yap ZH, Khetrapal V, Tay VSY, Drautz-Moses DI, Ali Y, Chen SL, Kline KA. 2024. Purine and carbohydrate availability drive Enterococcus faecalis fitness during wound and urinary tract infections. MBio 15.

96.   Liu GY, Doran KS, Lawrence T, Turkson N, Puliti M, Tissi L, Nizet V. 2004. Sword and shield: Linked group B streptococcal β-hemolysin/cytolysin and carotenoid pigment function to subvert host phagocyte defense. Proc Natl Acad Sci U S A 101:14491–14496.

97.   Spencer BL, Tak U, Mendonça JC, Nagao PE, Niederweis M, Doran KS. 2021. A type VII secretion system in Group B Streptococcus mediates cytotoxicity and virulence. PLOS Pathog 17:e1010121.

98.   Mascher T, Zähner D, Merai M, Balmelle N, De Saizieu AB, Hakenbeck R. 2003. The Streptococcus pneumoniae cia regulon: CiaR target sites and transcription profile analysis. J Bacteriol 185:60–70.

99.   Alves LA, Harth-Chu EN, Palma TH, Stipp RN, Mariano FS, Höfling JF, Abranches J, Mattos-Graner RO. 2017. The two-component system VicRK

regulates functions associated with Streptococcus mutans resistance to complement immunity. Mol Oral Microbiol 32:419–431.

100. Chang JC, Jimenez JC, Federle MJ. 2015. Induction of a quorum sensing pathway by environmental signals enhances group A streptococcal resistance to lysozyme. Mol Microbiol 97:1097–113.

101. Hooven TA, Bonakdar M, Chamby AB, Ratner AJ. 2019. A Counterselectable Sucrose Sensitivity Marker Permits Efficient and Flexible Mutagenesis in Streptococcus agalactiae. Appl Environ Microbiol 85:1–13.

102. Jeng A, Sakota V, Li Z, Datta V, Beall B, Nizet V. 2003. Molecular genetic analysis of a group A Streptococcus operon encoding serum opacity factor and a novel fibronectin-binding protein, SfbX. J Bacteriol 185:1208–17.

103. DeJesus MA, Ambadipudi C, Baker R, Sassetti C, Ioerger TR. 2015. TRANSIT - A Software Tool for Himar1 TnSeq Analysis. PLoS Comput Biol 11:1–17.

104. Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform. Bioinformatics 25:1754–1760.

105. Mills KB, Maciag JJ, Wang C, Crawford JA, Enroth TJ, Keim KC, Dufrêne YF, Robinson DA, Fey PD, Herr AB, Horswill AR. 2024. Staphylococcus aureus skin colonization is mediated by SasG lectin variation. Cell Rep 43.

106. Lang J, Capasso A, Jordan KR, French JD, Kar A, Bagby SM, Barbee J, Yacob BW, Head LS, Tompkins KD, Freed BM, Somerset H, Clark TJ, Pitts TM, Messersmith WA, Gail Eckhardt S, Wierman ME, Leong S, Kiseljak-Vassiliades K. 2020. Development of an adrenocortical cancer humanized mouse model to characterize anti-PD1 effects on tumor microenvironment. J Clin Endocrinol Metab 105:26–42.

107. Miller RJ, Crosby HA, Schilcher K, Wang Y, Ortines R V., Mazhar M, Dikeman DA, Pinsker BL, Brown ID, Joyce DP, Zhang J, Archer NK, Liu H, Alphonse MP, Czupryna J, Anderson WR, Bernthal NM, Fortuno-Miranda L, Bulte JWM, Francis KP, Horswill AR, Miller LS. 2019. Development of a Staphylococcus aureus reporter strain with click beetle red luciferase for enhanced in vivo imaging of experimental bacteremia and mixed infections. Sci Rep 9:1–19.

108. Bhattacharya M, Berends ETM, Chan R, Schwab E, Roy S, Sen CK, Torres VJ, Wozniak DJ. 2018. Staphylococcus aureus biofilms release leukocidins to elicit extracellular trap formation and evade neutrophil-mediated killing. Proc Natl Acad Sci U S A 115:7416–7421.

109. Bhattacharya M, Berends ETM, Zheng X, Hill PJ, Chan R, Torres VJ, Wozniak DJ. 2020. Leukocidins and the Nuclease Nuc Prevent Neutrophil-Mediated Killing of Staphylococcus aureus Biofilms. Infect Immun 88:1–16.

110. Kao PH-N, Ch'ng J-H, Chong KKL, Stocks CJ, Wong SL, Kline KA. 2023. Enterococcus faecalis suppresses Staphylococcus aureus-induced NETosis and promotes bacterial survival in polymicrobial infections. FEMS microbes 4:xtad019.

111. Freeman CD, Hansen T, Urbauer R, Wilkinson BJ, Singh VK, Hines KM. 2024.
Defective pgsA contributes to increased membrane fluidity and cell wall thickening
in Staphylococcus aureus with high-level daptomycin resistance. mSphere
9:e0011524.

112. Tiwari KB, Sen S, Gatto C, Wilkinson BJ. 2021. Fluorescence
PolarizationPolarization (FP) Assay for Measuring Staphylococcus aureus
Membrane Fluidity, p. 55–68. *In* Rice, KC (ed.), Staphylococcus aureus: Methods
and Protocols. Springer US, New York, NY.

113. Casadaban MJ, Cohen SN. 1980. Analysis of gene control signals by DNA fusion
and cloning in Escherichia coli. J Mol Biol 138:179–207.

114. Mu R, Kim BJ, Paco C, Del Rosario YD, Courtney HS, Doran KS. 2014.
Identification of a group B streptococcal fibronectin binding protein, SfbA, that
contributes to invasion of brain endothelium and development of meningitis. Infect
Immun 82:2276–2286.

1311

1312

1313