Chinmo is a novel regulator of differential Hippo signaling response within a single

developing organ

Shreeharsha Tarikere[1,2] *, Tarun Kumar[1,3] *, Guillem Ylla[1,4], Cassandra G. Extavour[1,3,5]

1. Department of Organismic & Evolutionary Biology, Harvard University, Cambridge MA,

USA.

2. Current address: Department of Human Genetics, McGill University, Montréal, Québec,

Canada

3. Howard Hughes Medical Institute, Bethesda MD, USA

4. Current address: Laboratory of Bioinformatics and Genome Biology, Faculty of

Biochemistry, Biophysics and Biotechnology, Jagiellonian University, Kraków, Poland

5. Department of Molecular & Cellular Biology, Harvard University, Cambridge MA, USA

* Contributed equally

Correspondence: guillem.ylla@uj.edu.pl, extavour@oeb.harvard.edu

ABSTRACT

Orchestrated control of proliferation of multiple cell types is essential for building a healthy organ.

Here we use larval ovary development in *Drosophila melanogaster* as a model to understand the

homeostasis of somatic and germ line cells in the formation of the female adult reproductive

organ. We previously showed that the highly conserved Hippo signaling pathway regulates

proliferation of both germ line and somatic cells in the *D. melanogaster* larval ovary. Response to

Hippo signaling appeared to be mediated by different genetic mechanisms in germ line and soma,

but the mechanism allowing distinct responses to the same signaling pathway remained unknown.

Here we perform cell type-specific RNA-Seq of isolated germ line and somatic cells from the

developing ovary at multiple time points spanning the formation of germ line stem cell niches, in

either a Hippo signaling loss- or gain-of-function genetic background. Applying network analysis

to these data revealed a novel regulator of ovarian development, the transcription factor *chinmo*.

Subsequent experimental validation showed that *chinmo* acts as a key germ cell-specific

translator of Hippo signaling in the developing ovary, allowing the Hippo signal to be transduced

in cell type-specific ways in germ line and somatic cells within the same organ.

KEY WORDS: Yorkie, ovariole number, FACS, RNA-Seq, Larva, Ovary, *Drosophila*

INTRODUCTION

Animal organs are not heterogeneous masses of cells. Instead, they are composed of multiple cell types that contribute to the same organ, but have distinct gene expression profiles, morphologies, behaviours, and functions. In most organ systems, all these distinct cell types are somatic cells derived from the same germ layer (Barresi and Gilbert 2023). An important exception to this rule is found in the gonads (ovaries and testes), which contain both somatic cells of mesodermal origin, and cells of the germ line, destined to make eggs and sperm (Extavour 2007). Gonads are typically assembled after the specification of germ layers and of the germ line, but before gametogenesis begins. During gonadogenesis, somatic and germ cells proliferate, exchange neighbors, change their shapes and gene expression profiles, and create the final functional anatomical organization of the gonad. As for other systems of organogenesis, cell-cell signaling is crucial for this process. However, each unique cell type within the developing organ often has a distinct, cell type-appropriate response to the same incoming signals, allowing correct organ formation and function. Here we use the *Drosophila melanogaster* (fruit fly) ovary as a study system to elucidate how different cell types within the same organ can differently transduce the same signaling inputs during development.

A single adult ovary in *D. melanogaster* is made up of an estimated 26 distinct cell types (Rust et al. 2020) that are assembled into 16-23 tubular egg producing subunits called ovarioles (Büning 1994), making it an excellent model to study the regulation of organ size and subunit number (Sarikaya and Extavour 2015; Kumar et al. 2020). Ovariole number varies by up to four orders of magnitude across insect species, and in some insects, including Drosophilids, it is positively correlated with egg-laying and therefore fitness (Church et al. 2021; David 1970). In *Drosophila* species, ovariole number is established in the larval stages and determined by the number of stacks (called terminal filaments: TFs) of a specific type of somatic cell (terminal filament cells: TFCs) (King 1968,Honěk 1993,Markow 2007,Hodin 2009,Green 2012,Sarikaya

2019). The number of these stacks of cells, in turn, is determined by the number and stacking

dynamics of the component TFCs. We previously showed that TFC proliferation, and therefore

ovariole number, is regulated by the Hippo (Hpo) signaling pathway, which affects both TFC and

germ cell numbers (Sarikaya and Extavour 2015).

First discovered as a network of tumor overgrowth factors in *D. melanogaster* (Tapon et

al. 2002; Dong et al. 2007; Xu et al. 1995; Wu et al. 2003), the Hpo signaling pathway regulates

organ size, growth, cell proliferation and cell fate determination in animals. In the developing fruit

fly ovary, this pathway can regulate proliferation both cell-autonomously (based on cell-intrinsic

Hpo signaling activity) and non-cell autonomously (activity in one cell can affect proliferation in a

different cell) (Sarikaya and Extavour 2015). Regulation of cell proliferation in both the soma and

the germ line is essential for the normal development of the ovary, including species-specific

ovariole number. Members of the Hippo signaling pathway are expressed ubiquitously across

multiple distinct ovarian cell types that nevertheless have different proliferation dynamics

(Sarikaya and Extavour 2015). This begs the question of how differential regulation of Hpo

signaling may be achieved despite the potential for ubiquitous activation.

The canonical Hippo signaling pathway functions through regulating shuttling of the

transcriptional activator Yorkie (Yki) in and out of the nucleus. This shuttling is regulated by the

function of the serine-threonine kinase Hpo in conjunction with a cascade of kinases that ends

with the phosphorylation of Yki, preventing its nuclear localisation (Oh and Irvine 2008; Huang et

al. 2005). Under this model, unphosphorylated Yki enters the nucleus to form tissue-specific

transcriptional regulatory complexes (Slattery et al. 2013) with loading proteins like Scalloped (Sd)

and Homothorax (Hth). These transcription cofactor complexes bind to DNA and activate a wide

range of target genes, including many necessary for growth and proliferation (Wu et al. 2008;

Zhang et al. 2008; Zhao et al. 2008). Phosphorylation of Yki by other kinases, including Nemo-

like kinase and AMP-activated protein kinase, can either prevent or enhance Yki's targeting by

the canonical Hpo kinases, allowing a Hpo-independent mode of Yki regulation (Moon et al. 2017;

Wang et al. 2015). Furthermore, changes in the binding dynamics of Sd, the coactivator of Yki,

adds another layer of Yki regulation (Manning et al. 2024). Lastly, mechanical stress and proximity

of cell membranes during cell growth and proliferation can also regulate the Hpo signaling activity

(Zhao et al. 2007). Across animals, the Hpo pathway regulates multiple cellular processes *via* Yki

target genes, including cell size, cell proliferation, adhesion and apoptosis (Zheng and Pan 2019),

while interacting with other signaling pathways such as JAK-STAT, Notch, Hedgehog and EGFR

(Besse 2005,Matsuoka 2013,Sarikaya 2015,Yatsenko 2021,Song 2007,Kumar 2020) .

Previously, we showed that Hpo pathway activity regulates cell proliferation in multiple *D.*

melanogaster ovarian cell types, which in turn can determine the numbers of TFCs, TFs and

ultimately ovarioles (Sarikaya and Extavour 2015). RNAi-mediated knockdown of *hpo* or *wts* in

somatic cells increased TF and TFC number, while RNAi against *yki* decreased the numbers of

both cell types (Sarikaya and Extavour 2015). In this context, we demonstrated that the Hpo

pathway interacts with the EGFR pathway to regulate numbers of another somatic cell type, called

intermingled cells (Sarikaya and Extavour 2015) and with JAK-STAT signaling to regulate TFC

proliferation and the homeostasis between intermingled cells and germ cells (Sarikaya and

Extavour 2015). Furthermore, our RNAi screen in a *hpo* knockdown background in the larval ovary

identified a gene interaction network implicating all known conserved animal signaling pathways

in regulating *D. melanogaster* ovary development and function, including establishment of correct

ovariole number (Kumar et al. 2020).

To our surprise, our previous study showed that in contrast to somatic cells, *hpo*

knockdown did not affect germ cell number (Sarikaya and Extavour 2015). However, *yki*

knockdown reduced germ cell number significantly, as it did in somatic cells. This result suggested

the existence of a Hpo-independent mechanism of Yki activity in the germ cells (Sarikaya and

Extavour 2015). However, a mechanistic explanation for the differential response of germ cells

and somatic cells to these Hpo signaling components remains lacking. Given the variation in

outcomes of Hpo pathway abrogation in the germ cells and the somatic cells, we sought to use

cell type-specific transcriptional profiling to identify distinct gene regulatory responses to *hpo* or

yki activity in soma versus germ line. Here we present the results of this approach, wherein we

analyze stage- and tissue-specific transcriptomic datasets generated in three distinct genetic

backgrounds to assess the role of Hpo signaling and its interacting pathways in the cell types of

the developing *D. melanogaster* larval ovary. We reveal the transcription factor Chinmo as a

putative novel regulator of germ cell-specific Hpo-dependent activity and provide experimental

validation of this regulatory interaction in the developing ovary. Further, we provide evidence that

chinmo regulates the previously observed (Sarikaya and Extavour 2015) non-autonomous effect

that Hpo activity in germ cells exerts on neighboring somatic cells. Thus, our results show how

variation in the expression of cell type-specific targets can produce distinct proliferation response

outcomes to the same signaling inputs within a developing tissue.

RESULTS

***hpo* and *yki* knockdown in somatic and germ line tissues**

D. melanogaster ovaries are made up of subunits called ovarioles (Figure 1A), whose number is

established at the third larval instar stage, 72-120 hours after egg laying (hAEL) at 25ºC (Figure

1B) (King 1970). The number of ovarioles in the adult directly corresponds to the number of TFs

formed in the larval ovary (Figure 1B), because TFs are neither formed nor destroyed during the

pupal phase that separates larva from adult (King 1970). The Hpo signaling pathway regulates

TF number by determining the number of TFCs available to assemble into TFs (Sarikaya and

Extavour 2015). We reasoned that the genetic mechanisms of Hpo regulation of TF number might

be revealed by determining the responses of germ cells and somatic cells to Hpo signaling activity

during TF formation (Sarikaya and Extavour 2015). We therefore selected larvae over three

stages of early ovary morphogenesis as follows: (1) before TF formation (Early = 72 hAEL; Figure

1Bi); (2) during TF formation (Middle (mid) = 96 hAEL; Figure 1Bii); and (3) after TF formation

(Late = 120 hAEL; Figure 1 Biii). We expressed GFP specifically in either the somatic cells (red

cells in Figure 1C) or in the germ cells (blue cells in Figure 1C) using cell type-specific GAL4

drivers (Brand and Perrimon 1993) (see Methods) and isolated somatic and germ cells

respectively using Fluorescence-Activated Cell Sorting (FACS) (Supplemental Figure S1). We

had previously established the gene expression profiles of wild type somatic and germ cells at

these stages (Tarikere et al. 2021). In the present study, we used the UAS/GAL4 system (Brand

and Perrimon 1993) to knock down either *hpo* or *yki* in either somatic or germ cells across the

three developmental stages described above, and then performed tissue-specific RNA-Seq to

identify the transcriptional regulatory effects of these two main components of the Hpo signaling

pathway on larval ovary development and morphogenesis (Figure 1D-E).

The Hpo pathway regulates gene expression differently in the ovarian germ line and soma

Our RNA-Seq study produced 48 independent datasets (Supplemental Table S1) of gene

expression in germ cells and somatic cells in two different knockdown conditions and at three

developmental stages. To identify and understand the variation in gene expression across

samples, we performed a principal component analysis (PCA). Our results showed that 51.5% of

the variation across all datasets was explained by cell type (Figure 2A). The first principal

component clearly separated somatic cells from germ cells (PC1 in Figure 2A). Cell type-specific

gene expression included both genes with known roles in the germ line (e.g. *vasa* (Lasko and

Ashburner 1988) and *bruno* (Kim-Ha et al. 1995) or in the somatic gonad (e.g. *traffic jam* (Li et al.

2003) and *bab2* (Godt and Laski 1995), as well as novel genes without such roles previously

described (Supplemental Table S2). The second greatest impact on transcriptional variation

(PC2) was developmental stage (Figure 2A).

We then performed individual PCA analysis of somatic cells (Figure 2B) and germ cells

(Figure 2C), which allowed us to better observe the effects of the two RNAi knockdown conditions

on the sample clustering. In the somatic cells, most of the variation (32.3%) amongst the samples

was attributable to developmental stage rather than to genotype (stages are indicated as distinct

shapes along PC1 in Figure 2B). Along the PC2 axis, datasets were sorted by genotype, with

both the *hpo* and *yki* loss-of-function treatments segregating in opposite directions relative to the

control across all three developmental stages (colors along PC2 in Figure 2B). By contrast, in the

germ cells, the samples clustered by genotype rather than by stage (colors along PC1 in Figure

2C). The *hpo* and *yki* loss-of-function conditions clustered together independently of

developmental stage, but far from the respective controls (Figure 2C).

These results suggest that *yki* and *hpo* regulate transcriptional activity in ovarian germ

cells and somatic cells differently during larval development. In the soma, the two genes have

distinct and stage-specific impacts on the transcriptomic landscape. In contrast, loss of function

of both genes generates very similar effects on the transcriptomes of the germ line, and these

regulatory impacts do not change significantly across the period of development studied here. To

better understand this variation at the level of individual genes in each cell type sample, we

performed a differential gene expression (DGE) analysis of *hpo* and *yki* knockdown conditions

compared to the corresponding control samples at each stage. In general, we found more

differentially expressed genes in germ cells than in somatic cells, when either *hpo* or *yki* levels

were reduced (Figure 2D, Supplemental Table S3).

In the soma, there was a similar number of upregulated (365) and downregulated (310)

genes (17.7%) under *hpo* knockdown, compared to over 60% more upregulated (231) than

downregulated (143) genes (61.5%) under *yki* knockdown (Figure 2D, Supplemental Table S3).

In contrast, in the germ cells we observed on average over twice as many down-regulated genes

(1,161 and 1,120 under *hpo* and *yki* knockdown respectively) as upregulated genes (421 and 600

under *hpo* and *yki* RNAi respectively) in both knockdown conditions. Interestingly, this much larger

number of differentially expressed *hpo*- and *yki*-responsive genes in the germ line did not

correspond with a notable change of the germ line transcriptional landscape throughout the

developmental period under study (Figure 2C). The soma, on the other hand, exhibited distinct

gene expression responses to Hpo signaling over developmental time (Figure 2B), and these

differential responses were explained by a comparatively small number of genes (Figure 2D). We

had previously observed that in the somatic cells, the cell proliferation phenotypes induced by the

reduction of *hpo* and *yki* activity were diametrically opposed (Sarikaya and Extavour 2015).

Consistently, differential gene expression analysis of *hpo* and *yki* knockdown conditions showed

that in the soma, most of the differentially regulated genes (75%) were exclusive to each genotype

(*hpo RNAi* 55% and *yki* RNAi 20%) (Figure 2E). Of all the differentially expressed genes, the

expression of only four genes (*apterous*, *CG11400*, *wrapper*, and *expansion*) was significantly

affected in opposite ways by *hpo* and *yki* RNAi treatment (Figure 2E). By contrast, in the germ

cells the majority (51%) of the differentially expressed genes were common to both *hpo* and *yki*

loss of function conditions (869 commonly downregulated and 245 commonly upregulated), and

only one gene (*CG34408*) showed an "opposite" expression change (upregulated in *yki* loss of

function and downregulated in *hpo* loss of function) (Figure 2F). These results are consistent with

the above-described observations that *yki* and *hpo* knockdowns induced similar changes in the

germ line transcriptome (Figure 2C), but induced genotype-specific effects on the transcriptome

of the soma (Figure 2B).

Functional prediction analysis of the differentially expressed genes

The differentially expressed genes in the germ cells included more significantly enriched pathways from the Kyoto Encyclopedia of Genes and Genomes (KEGG) and terms from the Gene Ontology (GO) database than the genes differentially expressed in the soma (Figure 3, Supplemental Figure S2). We found that the Hippo, MAPK, Hedgehog, and TGFβ pathways were significantly enriched among the downregulated genes in the germ line under both *hpo* and *yki* knockdown conditions (Figure 3). We additionally found that DNA replication and mismatch repair pathways were enriched within the genes upregulated in response to *hpo* knockdown in germ cells. In the germ line response to *yki* knockdown, we observed enriched GO terms relevant to meiosis and cell cycle regulation (Supplemental Figure S2a). In contrast, somatic ovarian cells displayed very few significantly altered pathways under either knockdown condition (Figure 3). GO analysis revealed that members of some of the same processes that were downregulated in the germ line, were downregulated in somatic *hpo* knockdowns, but with fewer gene counts than those in germ cells (Supplemental Figure S2). These processes included cell adhesion, taxis, motility, tissue growth, cell proliferation, stem cell population maintenance, maintenance of cell number, cell proliferation and cell death.

***yki* is downregulated under *hpo* knockdown in the germ line but not the soma**

Our previous work suggested that Yki functions downstream of Hpo in the soma but independently of Hpo in the germ line (Sarikaya and Extavour 2015). We used our cell-type-specific transcriptomes to determine whether this functional difference was reflected at the level of transcript regulation in one or both of germ line and soma. As expected, we observed a significant reduction (adjusted p-value<0.05) of *hpo* transcript levels across all stages and in both cell types under *hpo* knockdown (Figure 4A). The levels of *yki* transcript were also down-regulated (adjusted p-value<0.05) by *yki* RNAi in all stages and in both cell types, with two apparent exceptions

(Figure 4B). First, we noted that the 1.4-fold downregulation of *yki* under *yki* RNAi in early-stage

somatic cells just missed our significance threshold (adjusted p-value 0.055). Furthermore,

somatic cells at this stage displayed a different transcriptional landscape than controls, consistent

with *yki* knockdown (Figure 2B). Second, we noted that one of the control samples for mid stage

germ cells had anomalously low levels of *yki* transcript compared to other control samples of the

same and other stages (outlined in red in Figure 4B). Removal of this putative outlier revealed a

significant reduction of *yki* under yki RNAi (Supplemental Figure S3). Thus, we conclude that both

RNAi conditions were able to significantly reduce the transcript levels of their respective target

genes, and that our analytical methods were able to detect this reduction.

In terms of the effect of RNAi against these two genes on the transcript levels of the other,

we found that *yki* transcript levels were significantly reduced under *hpo* RNAi in late stage germ

cells, but not at earlier stages, and not in any stage in the soma (Figure 4B).*yki* RNAi (log2FC=-

1) and the *hpo* RNAi (log2FC=-0.8) conditions caused a similar reduction in *yki* transcript in the

germ line at this late stage. In contrast, the knockdown of *yki* did not yield a significant change in

hpo transcript levels in either cell type at any stage (Figure 4A). A possible exception may be the

mid stage germ cells: when the control sample with anomalously low levels of *yki* is excluded,

hpo appears significantly downregulated under *yki* knockdown with a fold change of -1.1

(Supplemental Figure S3). The degree of possible down-regulation of *hpo* in the *yki* RNAi

condition is much weaker (Log2FC=-1.1) than the reduction of *hpo* produced by the *hpo* RNAi

condition at the same stage (Log2FC=-4.1) (Figure 4A). Taken together, our results suggest that

the transcriptional regulatory relationship between *hpo* and *yki* in the larval female germ line is

complex, different from that in the soma, developmental stage-specific, and not directly predicted

from previous observations on the regulatory interactions of their protein products.

Gene regulatory network analysis suggests *chinmo* as a novel mediator of *hpo*-dependent

256

257 *yki* **regulation**

258 To generate hypotheses as to how *hpo* might regulate *yki* expression in the larval germ line, we

259 performed a network analysis. To this end, we built a directed network using the DroID database

260 of protein-protein interactions and transcription factor-encoding genes (Murali et al. 2011; Yu et

261 al. 2008). Additionally, we used publicly available Yki and GAF Chip-seq data from *D.*

262 *melanogaster* embryo, wing disc and S2 cells (Oh et al. 2013) to find possible direct targets of Yki

263 in *D. melanogaster* and added them into the network. From the ChIP-seq Yki data, we identified

264 1,513 peaks of Yki-binding regions, which we mapped to the closest gene, thus identifying 1,118

265 genes putatively regulated by Yki across the three studied cell types. By taking the intersection of

266 sets of putative Yki targets from all three cell types, we hoped to retrieve targets of Yki that might

267 also be relevant in the ovary.

268 In the germ line, 31.4% and 33.4% respectively of all the genes downregulated in either

269 *hpo* or *yki* knockdowns were putative direct targets of Yki (Supplemental Figure S4). Conversely,

270 of the 1,118 putative Yki targets in our network that were differentially expressed, most of them

271 were downregulated (94.59% under *hpo* RNAi and 79.14% under *yki* RNAi) (Supplemental Figure

272 S4). In contrast, in somatic cells only 8.23% and 2.86% of putative Yki target genes were

273 significantly affected by *hpo* and *yki* knockdowns respectively (Supplemental Figure S4).

274 Similarly, among the differentially expressed putative Yki targets the downregulated fraction was

275 much lower in the somatic cells than observed in the germ cells (67.37% were downregulated

276 under *hpo* knockdown, and 50% under *yki* knockdown; Supplemental Figure S4). This is

277 consistent with the overall much higher number of differentially regulated *hpo*- and *yki*-responsive

278 genes in the germ cells relative to somatic cells (Figure 2D)

279 We found that the shortest path linking *hpo* and *yki* in the network was through a predicted

280 interaction of Hpo with a transcription factor called Chronologically inappropriate morphogenesis

(Chinmo), which was predicted to regulate *yki* expression (Figure 4C). Moreover, Chinmo was the

sole candidate in our interaction network that was predicted to directly link both Hpo and Yki

(Figure 4C). Our RNA-Seq data showed that this transcription factor is significantly upregulated

in late stage germ cells under both *hpo* and *yki* loss of function conditions, and conversely, was

significantly downregulated in late stage somatic cells under both RNAi conditions (Supplemental

Figure S5). We therefore considered the hypothesis that differential *chinmo* expression and/or

function between germ line and soma might contribute to the different responses of those cell

types to Hpo pathway activity and sought to test this hypothesis experimentally.

Nuclear Chinmo in the Germ cells is regulated by *yki* expression

Our RNA-Seq results indicated that *chinmo* is expressed at constant levels in wild type germ line

cells throughout the developmental stages studied here, while levels of *chinmo* decreased over

developmental time in the ovarian somatic cells (Supplemental Figure S5). Given that *chinmo*

encodes a transcription factor (Zhu et al. 2006; Chen et al. 2020), we used an anti-Chinmo

antibody to examine the expression of Chinmo protein in the cells of the larval ovary. We detected

high levels of Chinmo protein in the nuclei of the germ cells in mid and late stages of larval

development, but only weak expression in the somatic ovarian cells (Figure 5A).

To determine the functional significance of *chinmo* expression in the larval ovary, we

reduced *chinmo* expression in either the somatic cells or the germ line cells. We used ovariole

number as a measure of effective change in phenotype in comparison with the *RNAi* knockdowns

of our predicted interactors of Chinmo in this context, Hpo and Yki, both of which we previously

identified as regulating ovariole number (Sarikaya and Extavour 2015). We found that, consistent

with the weakly detected Chinmo protein in the soma, reduction of *chinmo* in the somatic cells,

unlike the reduction of either *hpo* or *yki,* caused no significant change in ovariole number (Figure

5B; Supplemental Table S4). In contrast, abrogation of *chinmo* in the germ cells caused a

significant increase in ovariole number, as did reduction of *hpo* in the germ cells (Figure 5B;

Supplemental Table S4). Reducing levels of y*ki* in the germ cells did not cause any change in

ovariole number (Figure 5B; Supplemental Table S4).

Our network model predicted a putative transcriptional regulatory relationship between *yki*

and *chinmo*, and a potential protein-protein interaction between Hpo and Chinmo proteins. We

therefore asked whether levels of *yki* or *hpo* gene activity influenced the levels of Chinmo protein

in the developing ovary. Consistent with our network model, Chinmo protein levels were not

significantly affected by *hpo RNAi* (Figure 5C). Similarly, reduction of *chinmo* or *yki* expression in

somatic cells did not significantly alter detectable Chinmo protein levels in germ cells (Figure 5C;

Supplemental Table S4). However, Chinmo protein levels did significantly increase in the germ

cells upon *hpo* knockdown in somatic cells (Figure 5C; Supplemental Table S4), indicating a non-

cell autonomous role for Hpo in Chinmo regulation in the germ line. Reduction of *chinmo* or *yki* in

the germ line caused a reduction in detectable Chinmo protein in these cells (Figure 5C;

Supplemental Table S4).

To understand the functional outcomes in the ovary of Chinmo levels in the larval germ

cells (Figure 5C; Supplemental Table S4), we quantified germ cells, TFCs, and TFs at the late

larval stage, and ovarioles in the adult, under loss of germ line expression of *chinmo*, *yki* and *hpo*.

Reduction of *chinmo*, but not *yki* or *hpo*, in the germ cells led to a significant increase in germ cell

number when compared to both *GAL4* or *[RNAi]* controls (Figure 5D; Supplemental Table

S4). We found that the increase in ovariole number caused by reduction of *chinmo* or *hpo* in the

germ cells (Figure 5B; Supplemental Table S4) was the result of a non-cell-autonomous increase

in TFC number and in TF number (Figure 5E; Supplemental Table S4). Interestingly, reduction in

chinmo levels in the germ line also caused an increase in the number of TFCs in each TF (Figure

5E; Supplemental Table S4). Consistent with the observation that *yki* reduction in germ cells did

not change ovariole number (Figure 5B; Supplemental Table S4), this genetic background also

had wild type numbers of TFCs and TFs (Figure 5E; Supplemental Table S4).

Taken together, these data are consistent with the hypothesis that Chinmo protein in the

nucleus of germ cells regulates germ cell number cell-autonomously, and that Chinmo protein

levels in the germ line require germ line expression of *yki*. Further, expression of *chinmo* and

hippo in the germ cells also exerts a non-cell autonomous effect on somatic cells, limiting TFC

number, TF number, and hence ovariole number.

DISCUSSION

We have provided evidence for a novel genetic mechanistic hypothesis to explain the differential

response of germ line and somatic cells to the Hpo signaling pathway during the larval

development of the *D. melanogaster* ovary. We propose that the transcription factor Chinmo acts

as a modifier of Hpo-independent cell number regulation of the germ line, but not of the somatic

gonad cells.

The Hippo pathway has distinct effects on transcription in the germ cells and the soma of

the larval ovary

In this study, we knocked down either *hpo* or *yki* using RNAi driven in either the germ cells or the

somatic cells, thus providing us with a lens into the effects of each of these genetic manipulations

on the transcriptomic landscapes in each cell type. The canonical understanding of the Hippo

signaling pathway suggests that the presence of phosphorylated Hpo in the cellular milieu

prevents the nuclear localization of Yki and therefore the downstream expression of its target

genes. Under this model, the absence of Hpo in the cytoplasm allows Yki to remain

unphosphorylated, enter the nucleus, and drive the expression of its downstream targets. This

model therefore predicts a different transcriptional profile with and without Hpo. Our observation

that knockdown of either *hpo* or *yki* in the soma impacted transcript levels of two mutually

exclusive sets of genes supports the canonical understanding of the Hippo signaling pathway

(Figure 2B). In the germ cells, however, this is not the case: the effects of the knockdown of either

hpo or *yki* caused transcriptional changes in a similar set of genes (Figure 2C). This suggests that

hpo and *yki* RNAi have different effects on the transcriptome of the soma, but similar effects on

germ line.

Furthermore, we found that more genes were differentially expressed in the germ line than

in the soma (Figure 2D). The higher number of differentially expressed *yki*-regulated genes in

somatic tissues could mean that *yki* directly or indirectly represses more genes in somatic tissue

than it activates. Again, we observed differences in the effect of *hpo* and *yki* knockdown on germ

cells and somatic cells. The higher number of downregulated genes in the germ cells under these

conditions suggests that *hpo* and *yki* may operate in germ line tissue primarily through promoting

the transcription of genes rather than repressing them. In all genetic backgrounds, we observed

that most of the transcriptional variation in the soma was attributable to the progression of

development (Figure 2B), while in the germ line, the greatest variation between the samples was

due to abrogation of Hippo signaling pathway function (Figure 2C). This suggests that many of

the transcriptome differences we observed in the soma across development reflect the dynamic

morphogenetic changes taking place in the somatic cells of the ovary during the process of

terminal filament determination and organization.

Hpo RNAi in germ cells does not affect cell number via _yki_ repression

The similar transcriptomic impact of *yki* and *hpo* knockdowns in the germ cells is intriguing given

their opposite phenotypic effects on cell proliferation in the soma (Sarikaya and Extavour 2015).

We had previously shown that *hpo* knockdown in somatic cells increased their number, while *yki*

knockdown decreased their number (Sarikaya and Extavour 2015). Similarly, overexpression of

hpo and *yki* in somatic cells and in germ cells also had opposite phenotypic effects, whereby *hpo*

overexpression reduced somatic cell number and *yki* knockdown increased it (Sarikaya and

Extavour 2015). These results were consistent with the canonical model of Hippo pathway activity.

However, although *yki* knockdown in germ cells decreased germ cell number, *hpo* knockdown did

not change it. We suggest that transcriptional regulation of *yki* by *hpo* may contribute to the

previously reported apparent imperviousness of germ cell number to *hpo* knockdown. We noted

that in the germ line, a reduction in *hpo* caused a reduction in *yki* (Figure 4B), but that there was

no detectable regulation of *hpo* by *yki* (Figure 4A). To our knowledge, this is the first reported

example of the transcriptional regulation of *yki* by *hpo.* We hypothesize that when *hpo* is reduced

in germ cells, although nuclear localization of Yki is not inhibited, there is less *yki* transcript. This

could in principle lead to less Yki protein and less efficient triggering of the transcriptional cascade

that would result in a cell number increase. In sum, we hypothesize that a balance of *hpo* and *yki*

transcript levels contributes to regulating the number of germ cells.

Chinmo and its role in the regulation of _hpo_ and _yki_

The transcription factor *chinmo* is the only gene we found in our network analysis predicted to

directly regulate *yki* in response to *hpo* levels, since Hpo and Chinmo proteins have the potential

to interact based on yeast two hybrid data (Yu et al. 2008). Furthermore, our network predicts that

chinmo has the potential to regulate the expression of both *hpo* and *yki*, and that in turn, *yki* could

regulate the expression of Chinmo based on ChIP-seq data (Oh et al. 2013). We hypothesize that

chinmo could be involved in regulating *yki* transcript levels in response to Hpo activity levels,

401 which would provide a mode of control of Hpo by Yki through Chinmo. Previous studies of *chinmo*

402 in the *D. melanogaster* testis and ovaries showed that it plays a key role in inducing male fate in

403 somatic cells of the testis, but that it is not required in ovarian somatic cells (Ma et al. 2016).

404 Consistent with that report, our results show increased expression of *chinmo* in response to *yki*

405 knockdown in the germ line but not in the soma (Supplemental Figure S5).

406 Chinmo is a BTB-Zinc finger transcription factor that is conserved across insects (Spokony

407 and Restifo 2007; Nadimpalli et al. 2015). Chinmo was first identified as a negative regulator of

408 the JAK/STAT signalling pathway (Flaherty et al. 2009; Fisher et al. 2012; Wang et al. 2013) and

409 acts as a transcriptional repressor downregulating the expression of a broad spectrum of genes

410 including *shotgun* (*E-Cadherin*) (Enomoto et al. 2021), β *Spectrin*, *polychaetoid*, *scribble* and

411 *mirror* (Grmai et al. 2021). In some studied contexts, Chinmo prevents the differentiation of cells

412 and promotes their ability to proliferate (Chafino et al. 2023) .

413 Given the low levels of detectable Chinmo in somatic nuclei of the larval ovary (Figure

414 5A), and that adult ovariole number is unchanged by somatic reduction of *chinmo* (Figure 5C), we

415 hypothesize that Chinmo does not act in somatic cells to impact the known regulation of the

416 number of TFCs, TFs or ovarioles by *hpo* or *yki* (Sarikaya and Extavour 2015). In contrast, we

417 detected Chinmo protein in larval germ cell nuclei (Figure 5A), where its expression appears to

418 be dependent on *yki* levels in the germ cells (Figure 5C). Further, we observed that reduction of

419 *chinmo* in germ cells caused an increase in germ cell number (Figure 5D), suggesting that Chinmo

420 acts in larval ovarian germ cells to limit their proliferation. This is reminiscent of the observation

421 that in *chinmo*[ST] loss of function mutants, adult male germ cells overproliferate (Ma 2014) . On

422 the other hand, *chinmo*[-/-] clones in a heterozygous *chinmo*[+/-] background do not (Tseng 2021) .

423 Thus, the cell-autonomous impact of *chinmo* function on germ cell proliferation appears to be

424 context- and sex-specific. Combining the results presented herein with our previous

understanding of Hippo signaling in the ovary, we propose a model for the cell type-specific

regulation of *hpo*-mediated proliferation during larval ovary development (Figure 6).

In this model, Chinmo and Yki function in opposition to each other in the germ line to

regulate germ cell number, limiting or promoting germ cell proliferation respectively. Thus, unlike

in the somatic cells, loss of *hpo* in germ cells does not cause an increase in germ cell number

because Chinmo continues to prevent Yki-dependent transcription. The similarities that we

observed between the transcriptional profiles of the *hpo* and *yki* knockdown in the germ cells

might be explained by the prediction that Chinmo prevents Yki-dependent transcription in these

cells. Thus, we propose that the interplay of Chinmo and Yki levels in the nucleus contributes to

the maintenance of germ cell number, and that disrupting this balance causes a change in germ

cell number. This model is also consistent with the Hpo-independent function of Yki in the germ

cells that was suggested by our previous work (Sarikaya and Extavour 2015).

Cell-non-autonomous effects of Chinmo and Hippo

Reduction of somatic *hpo* causes a non-autonomous increase in the levels of detectable nuclear

Chinmo in the germ cells (Figure 5C). This could be due to *hpo* downregulation in either or both

of TFCs or intermingled cells, given that the GAL4 driver we used here (*tj:Gal4*) is expressed in

both somatic cell types (Sarikaya and Extavour 2015). We note that downregulation of *hpo* in

intermingled cells causes JAK/STAT pathway upregulation in germ cells (Sarikaya and Extavour

2015), and that Chinmo is positively regulated by the JAK/STAT signalling pathway (Flaherty et

al. 2009; Fisher et al. 2012; Wang et al. 2013). It is thus possible that abrogating *hpo* activity in

intermingled cells causes nuclear Chinmo levels to rise in the germ line because of the non-

autonomous upregulation of JAK/STAT activity in germ cells.

Our observation of the non-autonomous impact of germ cell number on multiple aspects

of TFC number and organization (Figure 5B, 5E) is a new and intriguing finding. Reducing *chinmo*

or *hpo* in the germ line led to a significant increase in TFC, TF (Figure 5E) and ovariole number

(Figure 5B), suggesting that germ cell number can positively regulate ovariole number, a

hypothesis that has remained underexplored in the field. In our previous work (Sarikaya and

Extavour 2015) we did not observe a statistically significant increase in germ cells in

nos:GAL4>hpo[RNAi] larval ovaries (Sarikaya and Extavour 2015), which is the same genetic

background we used in the present work. Further, although we did not count ovarioles in our

previous analysis of the *nos:GAL4>hpo*[RNAi] condition, we did not observe a statistically significant

non-autonomous impact of *hpo* knockdown in the germ line on terminal filament cell number. We

attribute these differences to the fact that the control genotypes used for comparison and

calculation of statistical significance in the present work (GAL4 or UAS heterozygotes obtained

by crossing the parent strains to wild type strain Oregon R), were slightly different from the control

genotypes used in our previous study (sibling controls) (Sarikaya 2015) . We and others have

observed that ovariole number, and its contributing developmental processes, are such highly

polygenic traits that even small changes in genetic background can have statistically detectable

effects (Coyne 1991,Thomas-Orillard 1976,Bergland 2008,Orgogozo 2006,Telonis-Scott

2005,WAYNE 2001,Wayne 2002,Kumar 2020) .

Although increasing ovariole number by increasing germ cell number has not, to our

knowledge, previously been reported, there are at least two reported examples of the inverse

situation, where reduced ovariole number is correlated with reduced germ cell number. In one

example, ovaries of adult females carrying the dominant female sterile *ovo*[D1] allele, which lacks

functional adult female germ cells, have fewer ovarioles than wild type (Extavour 2000). This could

in principle be because of a reduction in total germ cell number during larval stages in this

genotype, which is consistent with the interpretation of larval germ line clonal analysis of this allele

(Perrimon 1984). In a second example, agametic adult females generated by irradiating the

posterior pole of embryos (Geigy 1931), have fewer ovarioles than controls (Aboïm 1945).

Finally, we note that reduction in *chinmo* levels in the germ cells also caused an

unexpected increase in the number of TFCs in each TF (Figure 5E). To our knowledge, this is the

first reported evidence of changes in germ line gene influencing TCF number and TF

morphogenesis. Determining the non-autonomous regulatory mechanism that allows larval germ

cells to impact the future fecundity of the adult, by influencing somatic morphogenesis of the

ovary, is an interesting avenue for future research.

MATERIALS AND METHODS

Fly stocks

Flies were reared as previously described (Sarikaya et al. 2012) at 25ºC and 60% humidity on

standard *Drosophila* food with yeast in uncrowded vials. The "wild type" genetic background

referred to herein is Oregon R (Lindsley and Grell 1968). The following fly lines were obtained

from the Bloomington *Drosophila* Stock Center (BDSC): *w[*]; P{w[+mW.hs]=GawB}bab1[Pgal4-*

2]/TM6B, Tb[1] (abbreviated herein as *bab:GAL4*; BDSC stock number 6803); *P{w[+mC]=UAS-*

Dcr-2.D}1, w[1118]; P{w[+mC]=GAL4-nos.NGT}40 (abbreviated herein as *nos:GAL4*; BDSC

stock number 25751); *y w; P{w[+mW.hs]=GawB}NP1624* was obtained from the Kyoto Stock

Centre (abbreviated herein as *tj:*GAL4; stock number K104–055); *y[1] v [1];*

P(y[+t7.7]v[+t1.8]=TRiP. HMS00006) attP2 (abbreviated herein as *hpo[RNAi]*; BDSC stock

number 33614) and *y[1] v[1]; P(y[+t7.7] v[+t1.8]=TRiP.HMS00041)attP2* (abbreviated herein as

yki[RNAi]; BDSC stock number 34067); *y[1] v[1]; P{y[+t7.7] v[+t1.8]=TRiP.JF02341}attP2*

(abbreviated herein as *chinmo[RNAi]*; BDSC stock number 26777). *w[1118], P[UAS Stinger]*

(abbreviated herein as *UAS:Green Stinger I*, (Barolo et al. 2000) used for GFP expression was

a gift from Dr. James Posakony (University of California, San Diego). Crosses (described below

in "Staging larvae") were set with 100-200 virgin females and 50-100 males in a 180 ml bottle

(with 50 ml media per bottle) one day prior to egg laying. For somatic *chinmo* knockdown via

RNAi, we used a *tj:GAL4* driver (Li et al. 2003), rather than the *bab*:*GAL4* driver (Cabrera et al.

2002), as the latter caused lethality when driving *chinmo [RNAi]* (data not shown).

Staging larvae

Uniformly staged larvae were collected from near-synchronously laid eggs. To obtain a desired

genotype, crosses were set as described above in "Fly Stocks" at 25°C at 60% humidity and left

overnight to mate. Hourly egg collections were set up on 60 mm apple juice-agar plates (9 g agar,

10 g sugar, 100 ml apple juice and 300 ml water) with a pea-sized spread of fresh yeast paste

(baker's yeast granules made into a paste with a drop of tap water). Eggs were collected hourly

for eight hours. The first two collection plates were discarded to remove asynchronously laid eggs

that may have been retained inside the females following fertilization. Staged first instar larvae

were collected into vials 24 hours after egg collection. Larvae at 72h hours After Egg Laying (h

AEL) were designated as early stage, at 96h AEL as middle stage and at 120h AEL as late stage

of terminal filament development.

Dissection and dissociation of larval ovaries

Staged larvae were collected for dissection every hour. Fat bodies were dissected out of the

larvae using a pair of forceps. Ovaries, located in the center of the length of each fat body, were

then dissected free of the fat body using an insulin syringe needle (BD 328418). Ovaries thus

dissected clear of fat body were collected in ice-cold DPBS (Thermo Fisher 14190144) and

batches of 20-30 ovaries in DPBS were kept on ice for a maximum of four hours until dissociation

and subsequent FACS processing. For dissociation, dissected ovaries in ice-cold DPBS were

transferred to 0.25% trypsin solution (Thermo Fisher 25200056) for ten minutes at room

temperature in the cavity of a glass spot plate (Fisher Scientific 13-748B). They were then

transferred to another cavity containing 2.5% liberase (5 g liberase reconstituted in 2 ml nuclease

free water; Sigma 5401119001), teased apart with tungsten needles until most of the clumps were

separated, and left in the liberase solution (without agitation) at room temperature for ten minutes.

The tissue in liberase was pipetted gently to uniformly mix and dissociate the cells. The cell

suspension was then transferred to an RNA Lobind tube (Eppendorf 8077-230) and placed on a

vortexer for 1 minute. This sample was immediately subjected to fluorescence activated cell

sorting (FACS) and the sorted cell suspension was collected in an RNA Lobind tube containing

100-200 μl Trizol (Thermo Fisher 15596026).

Flow sorting GFP positive cells

The dissociated cell suspension was sorted in a MoFlo Astrios EQ Cell sorter (Beckman Coulter)

run with Summit v6.3.1 software. The dissociated cell suspension was diluted and a flow rate of

200 events per second was maintained with < 98% sorting efficiency during the sorting process.

A scatter gate (R1) was employed to eliminate debris, and a doublet gate (R2) was used to

exclude non-singlet cells. A 488 nm emission Laser was used to excite the GFP, and the collection

was at 576 nm. The GFP-positive cells were designated in gate R3 and sorted directly into Trizol

(Supplemental Table S1). The resulting cells collected in Trizol were frozen immediately by

plunging the tube in liquid nitrogen and then stored at -80°C until RNA extraction. A single

replicate consisted of at least 1000 cell counts pooled from FACS runs.

RNA extraction

Flow sorted cells stored in Trizol at -80°C were thawed at room temperature and lysed with a motorized pellet pestle (Kimble 749540-0000). A Zymo RNA Micro-Prep kit (Zymo Research R2060) was used to isolate RNA from the homogenized Trizol preparations. An equal volume of molecular grade ethanol (Sigma E7023) was added to Trizol and mixed well with a pellet pestle, then pipetted onto a spin column. All centrifugation steps were done at 10,000g for one minute at room temperature. The column was washed with 400 µl Zymo RNA wash buffer and then treated with Zymo DNase (6 U/µl) for 15 minutes at room temperature. The column was then washed twice with 400 µl Zymo RNA Pre-wash buffer and once with Zymo RNA wash-buffer. The RNA was eluted from the column in 55 µl of nuclease-free water (Thermo Fisher 10977015). The RNA obtained was quantified first using a NanoDrop (Model ND1000) spectrophotometer and then using a high sensitivity kit (Thermo Fisher Q32852) on a Qubit 3.0 Fluorometer (Thermo Fisher Q33216). The RNA was also checked for integrity on a high sensitivity tape (Agilent 5067-5579) with an electronic ladder on an Agilent Tapestation 2200 or 4200.

Library preparation

cDNA libraries were prepared using the Takara Apollo library preparation kit (catalogue # 640096). Extracted RNA samples were checked for quality using Tapestation tapes. 50µl of RNA samples were pipetted into Axygen PCR 8-strip tubes (Fisher Scientific 14-222-252) and processed through PrepX protocols on the Apollo liquid handling system. mRNA was isolated using PrepX PolyA-8 protocol (Takara 640098). The mRNA samples were then processed for cDNA preparation using the PrepX mRNA-8 (Takara 640096) protocol. cDNA products were then amplified for 15 cycles of PCR using longAmp Taq (NEB M0287S). During amplification PrepX RNA-Seq index barcode primers were added to each library to enable multiplexing. The amplified library was then cleaned up using PrepX PCR cleanup-8 protocol with magnetic beads (Aline C-

1003). The final cDNA libraries were quantified using a high sensitivity dsDNA kit (Thermo Fisher

Q32854) on a Qubit 3.0 Fluorometer (Thermo Fisher Q33216). cDNA content and quality were

assessed with D1000 (Agilent 5067-5582) or high sensitivity D1000 tape (Agilent 5067-5584, for

low quantities of cDNA) on an Agilent Tapestation 2200 or 4200.

Sequencing cDNA libraries

Libraries were sequenced on an Illumina HiSeq 2500 sequencer. Single-end 50bp reads were

sequenced on a high-throughput flow cell. Libraries of varying concentrations were normalized to

be equimolar, the concentrations of which ranged between 2-10nM per lane. All the samples in a

flow cell were multiplexed and later separated based on unique prepX indices to yield at least 10

million reads per library. The reads were demultiplexed and trimmed of adapters using the

bcl2fastq2 v2.2 pipeline to yield final fastq data files.

Three RNA-Seq biological replicates were obtained for each condition, each of which had

at least 10 million reads. The only exception was the early-stage germ line under both *hpo* and

yki RNAi knockdowns, for which we only obtained a single biological replicate due to the technical

challenge of obtaining enough germ line cells from these knockdown ovaries. These single

replicates of early germ line stages were not included in the differential expression analysis but

are made publicly available as part of the reported dataset (See Data Availability).

Immunohistochemistry

Larvae were dissected as previously described (Sarikaya et al. 2012; Sarikaya and Extavour

2015) and stained as previously described. For all cell counts, samples were dissected at the

larval-pupal transition stage. Primary antibodies and the concentrations used to stain the larval-

pupal transition stage ovaries are as follows: Mouse anti- Engrailed 4D9 (1:40, Developmental

Studies Hybridoma Bank (DSHB)), Rabbit anti-Vasa (1:3000, gift from A. Nakamura, Kumamoto

University), Rat anti-Chinmo (1:500, gift from E. Bach, New York University; (Wu 2012).

Secondary antibodies used were anti-mouse Alexa Fluor 488, anti-rat Alexa Fluor 488, and anti-

rabbit Alexa Fluor 568, all used at 1:400, and Hoechst 33342 used at 1:10,000 (stock solution 10

mg/mL). All samples were imaged on a Zeiss LSM 880 inverted laser confocal microscope. For

quantification of Chinmo expression levels, all samples were dissected and stained in parallel.

Imaging settings, including laser power, detector gain and step sizes, were kept constant for all

experimental and appropriate control samples.

Quantification of ovariole number, cell number and Chinmo protein level

Ovariole number was quantified as previously described (Kumar et al. 2020) . Germ cells were

counted manually in larval-pupal transition stage ovaries stained for Vasa protein, using the

Manual tracking with TrackMate plugin in Fiji (Schindelin et al. 2012; Tinevez et al. 2017) .

Terminal filaments and terminal filament cells in each terminal filament were counted manually in

larval-pupal transition stage ovaries stained for Engrailed, using the Cell counter plugin in Fiji.

Total terminal filament cell number per ovary was calculated by adding up the number of terminal

filament cells in each terminal filament.

RNA-Seq data processing

The *D. melanogaster* genome assembly and gene annotations were obtained from FlyBase

version dmel_r6.36_FB2020_05 (Larkin et al. 2020) . The reads were aligned with RSEM v1.3.3.

(Li and Dewey 2011) Using STAR v2.7.6a as read aligner, (Dobin et al. 2013) we obtained the

gene counts in each library. Because some of the tissue-specific biological samples were

sequenced in more than one lane or run, the reads were split into multiple fastq files, and the

gene counts belonging to the same biological sample were summed. Gene counts in each dataset

were normalized with the variance stabilizing transformation (VST) method implemented in the

DESeq2 v1.26.0 (Love et al. 2014) R package. Further analyses, such as principal component

analysis, hierarchical clustering, and differential expression analysis, were performed in R using

the VST-normalized counts.

Differential expression analysis

The differential expression analyses were performed with DESeq2 v1.26.0 (Love et al. 2014).

Genes with a Benjamini-Hochberg adjusted p-value lower than 0.05 were selected as differentially

expressed in the corresponding contrast.

Functional predictions

The Gene Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways

enrichment analyses were performed on the differentially expressed genes with the enrichGO

and enrichKEGG functions of the clusterProfiler package (v3.14.3) for R (Yu et al. 2012). The GO

terms were obtained using the R package AnnotationDbi (Carlson 2024) with the database

org.Dm.eg.db v3.10.0. The GO overrepresentation analysis of biological process was performed

against the gene universe of all *D. melanogaster* annotated genes in org.Dm.eg.db, adjusting the

p-values with the Benjamini-Hochberg method, adjusted p-value and q-value cutoff of 0.01, and

a minimum of 30 genes per term. For the KEGG enrichment analysis, p-values were adjusted by

the Benjamini-Hochberg procedure, and an adjusted p-value cutoff of 0.05 was used.

Analysis of publicly available ChIP-Seq data

We obtained the ChIP-Seq from GEO (GSE46305 and GSE38594) (Oh et al. 2013, 2014). We downloaded the coordinates of the 3,749, 6,491, and 3,991 Yorkie-bound peaks from embryo (E8-16h), wing imaginal discs, and S2 cells respectively. The coordinates of the peaks were based on the *D. melanogaster* genome assembly BDGP release 5, and transformed to coordinates for the latest genome version (release 6) with the lifterOver function. The peaks were annotated with ChIPseeker (v.1.22.1) using *D. melanogaster* genome annotations from FlyBase (v6.36). We defined the promoter region as the 1,500 bp flanking the transcription start site, and 69-75% of the peaks were found within the promoter regions of genes thus defined. In total 2,331, 3,931, and 2,247, genes with Yki-bound sites in the defined promoter region were identified, in embryo, wing imaginal discs, and S2 cells respectively. 4,980 genes were unique (appeared only in one dataset), 1,118 were common across the three datasets, and 2,406 were common to at least two datasets. Following the same methodology, we analyzed the GAF (*Trl*) Chip-Seq peaks (GSE38594) from embryo (E8-16h) and wing imaginal discs. Thus, we associated the 6,449 and 6,302 GAF-bound peaks from embryo and wing imaginal discs respectively to the promoters of 4,086 genes, of which 2,095 were common to both datasets.

Network analysis

We obtained the Transcription Factor – Gene (TF-gene) interaction network form the Drosophila Interactions database (www.droidb.org, version DroId_v2018_08), containing 157,462 interactions of 12,323 genes. We added to this network the 1,118 interactions of Yorkie and the 2,095 interactions of GAF predicted from the ChIP-seq analysis. After removing redundancies, we had 158,923 edges and 112,562 nodes in the network. We also obtained 127,353 unique protein-protein interactions (PPI) between 10,632 proteins from DroID v2018_08. The TF-gene and PPI networks were joined, yielding a network with 328,989 edges and 13,337 nodes while

designating as an edge attribute whether the interaction was a TF-gene or PPI. We assigned

directionality to the TF-gene edges because TFs regulate the target and not vice-versa while the

PPI edges were set as bi-directional. We looked for the shortest path in the network in which Hpo

could regulate Yki, and Chinmo emerged as the only possible connector, requiring only two steps.

COMPETING INTERESTS STATEMENT

The authors declare no competing interests.

ACKNOWLEDGEMENTS

We thank the flow cytometry and genomics staff at the Bauer Core Facility of Harvard University

Faculty of Arts and Sciences for advice on FACS sorting and sequencing, and members of the

Extavour lab for helpful discussions. This work was supported by NIH R01 award R01-HD073499-

0 to CGE, by funds from Harvard University, and by the Google Cloud Academic Research Credits

Program. CGE is a Howard Hughes Medical Institute investigator.

AUTHOR CONTRIBUTIONS

ST performed all steps in generating individual transcriptome datasets, including experimental

design, ovary dissection and dissociation, RNA-Seq library preparation and sequencing. GY

performed all statistical and network analyses of transcriptome data. TK performed and analysed

genetic experiments testing the role of *chinmo*. CGE conceived of the project, provided oversight

and funding. All authors drafted and edited the manuscript.

DATA AVAILABILITY

All the raw data are publicly available at NCBI-Gene Expression Omnibus (GEO) database under the accession code GSE250320. Control datasets for early, mid and late-stage germ line and somatic cells are accessible as a part of the NCBI-GEO database under the accession code GSE172015 (GSM5239772-89). The scripts used to process and analyze the data are available at the GitHub repository https://github.com/guillemylla/Ovariole_morphogenesis_RNAseq, commit ID 171064d.

REFERENCES

Barolo S, Carver LA, Posakony JW. 2000. GFP and -Galactosidase Transformation Vectors for Promoter/Enhancer Analysis in Drosophila. *BioTechniques* **29**: 726–732.

Barresi MJF, Gilbert SF. 2023. Developmental Biology.

Besse F, Busson D, Pret A. 2005. Hedgehog signaling controls Soma-Germen interactions during Drosophila ovarian morphogenesis. *Dev Dynam* **234**: 422–431.

Brand AH, Perrimon N. 1993. Targeted gene expression as a means of altering cell fates and generating dominant phenotypes. *Development* **118**: 401–415.

Büning J. 1994. Chapter 3. In *The Insect Ovary, Ultrastructure, previtellogenic growth and evolution*, pp. 31–324.

Carlson M. 2024. AnnotationDbi: Introduction To Bioconductor Annotation Packages. https://bioconductor.org/packages/release/bioc/html/AnnotationDbi.html.

Chen H-M, Marques JG, Sugino K, Wei D, Miyares RL, Lee T. 2020. CAMIO: a transgenic CRISPR pipeline to create diverse targeted genome deletions in Drosophila. *Nucleic Acids Res* **48**: 4344–4356.

Church SH, Medeiros BAS de, Donoughe S, Reyes NLM, Extavour CG. 2021. Repeated loss of variation in insect ovary morphology highlights the role of development in life-history evolution. *Proc Royal Soc B* **288**: 20210150.

David J. 1970. Le nombre d'ovarioles chez Drosophila melanogaster: relation avec la fécondité et valeur adaptative. *Achives de Zoologie experimentale et generale* **111**: 357–370.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR. 2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**: 15–21.

Dong J, Feldmann G, Huang J, Wu S, Zhang N, Comerford SA, Gayyed MF, Anders RA, Maitra A, Pan D. 2007. Elucidation of a Universal Size-Control Mechanism in Drosophila and Mammals. *Cell* **130**: 1120–1133.

Enomoto M, Takemoto D, Igaki T. 2021. Interaction between Ras and Src clones causes interdependent tumor malignancy via Notch signaling in Drosophila. *Dev Cell* **56**: 2223-2236.e5.

Extavour CGM. 2007. Gray anatomy: phylogenetic patterns of somatic gonad structures and reproductive strategies across the Bilateria. *Integr Comp Biol* **47**: 420–426.

Fisher KH, Wright VM, Taylor A, Zeidler MP, Brown S. 2012. Advances in genome-wide RNAi cellular screens: a case study using the Drosophila JAK/STAT pathway. *BMC Genom* **13**: 506.

Flaherty MS, Zavadil J, Ekas LA, Bach EA. 2009. Genome-wide expression profiling in the
Drosophila eye reveals unexpected repression of notch signaling by the JAK/STAT pathway.
Dev Dyn **238**: 2235–2253.

Godt D, Laski FA. 1995. Mechanisms of cell rearrangement and cell recruitment in Drosophila
ovary morphogenesis and the requirement of bric a brac. *Development* **121**: 173–187.

Grmai L, Harsh S, Lu S, Korman A, Deb IB, Bach EA. 2021. Transcriptomic analysis of
feminizing somatic stem cells in the Drosophila testis reveal putative downstream effectors of
the transcription factor Chinmo. *G3 Genes Genomes Genetics* **11**: jkab067-.

Hodin J. 2009. Chapter 11. She Shapes Events As They Come: Plasticity in Female Insect
Reproduction. In *Phenotypic Plasticity of Insects* (ed. D.W. Whitman).

Honěk A. 1993. Intraspecific Variation in Body Size and Fecundity in Insects: A General
Relationship. *Oikos* **66**: 483.

Huang J, Wu S, Barrera J, Matthews K, Pan D. 2005. The Hippo Signaling Pathway
Coordinately Regulates Cell Proliferation and Apoptosis by Inactivating Yorkie, the
Drosophila Homolog of YAP. *Cell* **122**: 421–434.

Kim-Ha J, Kerr K, Macdonald PM. 1995. Translational regulation of oskar mRNA by Bruno, an
ovarian RNA-binding protein, is essential. *Cell* **81**: 403–412.

King R. 1970. *Ovarian Development in Drosophila Melanogaster*. Academic Press.

King RC, Aggarwal SK, Aggarwal U. 1968. The development of the female Drosophila
reproductive system. *J Morphol* **124**: 143–165.

Kumar T, Blondel L, Extavour CG. 2020. Topology-driven protein-protein interaction network
analysis detects genetic sub-networks regulating reproductive capacity. *Elife* **9**: e54082.

Larkin A, Marygold SJ, Antonazzo G, Attrill H, dos Santos G, Garapati PV, Goodman JL,
Gramates LS, Millburn G, Strelets VB, et al. 2020. FlyBase: updates to the Drosophila
melanogaster knowledge base. *Nucleic Acids Res* **49**: D899–D907.

Lasko PF, Ashburner M. 1988. The product of the Drosophila gene vasa is very similar to
eukaryotic initiation factor-4A. *Nature* **335**: 611–617.

Li B, Dewey CN. 2011. RSEM: accurate transcript quantification from RNA-Seq data with or
without a reference genome. *BMC Bioinform* **12**: 323.

Li MA, Alls JD, Avancini RM, Koo K, Godt D. 2003. The large Maf factor Traffic Jam controls
gonad morphogenesis in Drosophila. *Nat Cell Biol* **5**: 994–1000.

Lindsley DL, Grell EH. 1968. *Genetic Variations of Drosophila melanogaster*. Carnegie
Institution of Washington, Washington, DC, USA.

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol* **15**: 550.

Ma Q, Cuevas M de, Matunis EL. 2016. Chinmo is sufficient to induce male fate in somatic cells
of the adult Drosophila ovary. *Development* **143**: 754–763.

Manning SA, Kroeger B, Deng Q, Brooks E, Fonseka Y, Hinde E, Harvey KF. 2024. The
Drosophila Hippo pathway transcription factor Scalloped and its co-factors alter each other's
chromatin binding dynamics and transcription in vivo. *Dev Cell* **59**: 1640-1654.e5.

Markow TA, O'Grady PM. 2007. Drosophila Biology in the Genomic Age. *Genetics* **177**: 1269–
1276.

Matsuoka S, Hiromi Y, Asaoka M. 2013. Egfr signaling controls the size of the stem cell
precursor pool in the Drosophila ovary. *Mech Dev* **130**: 241–253.

Moon S, Kim W, Kim S, Kim Y, Song Y, Bilousov O, Kim J, Lee T, Cha B, Kim M, et al. 2017.
Phosphorylation by NLK inhibits YAP-14-3-3-interactions and induces its nuclear localization.
EMBO Rep **18**: 61–71.

Murali T, Pacifico S, Yu J, Guest S, Roberts GG, Finley RL. 2011. DroID 2011: a
comprehensive, integrated resource for protein, transcription factor, RNA and gene
interactions for Drosophila. *Nucleic Acids Res* **39**: D736–D743.

Nadimpalli S, Persikov AV, Singh M. 2015. Pervasive Variation of Transcription Factor
Orthologs Contributes to Regulatory Network Evolution. *PLoS Genet* **11**: e1005011.

Oh H, Irvine KD. 2008. In vivo regulation of Yorkie phosphorylation and localization.
Development **135**: 1081–1088.

Oh H, Slattery M, Ma L, Crofts A, White KP, Mann RS, Irvine KD. 2013. Genome-wide
Association of Yorkie with Chromatin and Chromatin-Remodeling Complexes. *Cell Rep* **3**:
309–318.

Oh H, Slattery M, Ma L, White KP, Mann RS, Irvine KD. 2014. Yorkie Promotes Transcription by
Recruiting a Histone Methyltransferase Complex. *Cell Rep* **8**: 449–459.

Rust K, Byrnes LE, Yu KS, Park JS, Sneddon JB, Tward AD, Nystul TG. 2020. A single-cell
atlas and lineage analysis of the adult Drosophila ovary. *Nat Commun* **11**: 5628.

Sarikaya DP, Belay AA, Ahuja A, Dorta A, Green DA, Extavour CG. 2012. The roles of cell size
and cell number in determining ovariole number in Drosophila. *Dev Biol* **363**: 279–289.

Sarikaya DP, Extavour CG. 2015. The Hippo Pathway Regulates Homeostatic Growth of Stem
Cell Niche Precursors in the Drosophila Ovary. *Plos Genet* **11**: e1004962.

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S,
Rueden C, Saalfeld S, Schmid B, et al. 2012. Fiji: an open-source platform for biological-
image analysis. *Nat Methods* **9**: 676–82.

Slattery M, Voutev R, Ma L, Nègre N, White KP, Mann RS. 2013. Divergent Transcriptional
Regulatory Logic at the Intersection of Tissue Growth and Developmental Patterning. *PLoS*
Genet **9**: e1003753.

Song X, Call GB, Kirilly D, Xie T. 2007. Notch signaling controls germline stem cell niche
formation in the Drosophila ovary. *Development* **134**: 1071–1080.

Spokony RF, Restifo LL. 2007. Anciently duplicated Broad Complex exons have distinct
temporal functions during tissue morphogenesis. *Dev Genes Evol* **217**: 499–513.

Tapon N, Harvey KF, Bell DW, Wahrer DCR, Schiripo TA, Haber DA, Hariharan IK. 2002.
salvador Promotes Both Cell Cycle Exit and Apoptosis in Drosophila and Is Mutated in
Human Cancer Cell Lines. *Cell* **110**: 467–478.

Tarikere S, Ylla G, Extavour CG. 2021. Distinct gene expression dynamics in germ line and
somatic tissue during ovariole morphogenesis in Drosophila melanogaster. *G3 Genes*
Genomes Genetics.

Tinevez J-Y, Perry N, Schindelin J, Hoopes GM, Reynolds GD, Laplantine E, Bednarek SY,
Shorte SL, Eliceiri KW. 2017. TrackMate: An open and extensible platform for single-particle
tracking. *Methods* **115**: 80–90.

Wang H, Chen X, He T, Zhou Y, Luo H. 2013. Evidence for Tissue-Specific JAK/STAT Target
Genes in Drosophila Optic Lobe Development. *Genetics* **195**: 1291–1306.

Wang W, Xiao Z-D, Li X, Aziz KE, Gan B, Johnson RL, Chen J. 2015. AMPK modulates Hippo
pathway activity to regulate energy homeostasis. *Nat Cell Biol* **17**: 490–499.

Wu S, Huang J, Dong J, Pan D. 2003. hippo Encodes a Ste-20 Family Protein Kinase that
Restricts Cell Proliferation and Promotes Apoptosis in Conjunction with salvador and warts.
Cell **114**: 445–456.

Wu S, Liu Y, Zheng Y, Dong J, Pan D. 2008. The TEAD/TEF Family Protein Scalloped
Mediates Transcriptional Output of the Hippo Growth-Regulatory Pathway. *Dev Cell* **14**: 388–
398.

Xu T, Wang W, Zhang S, Stewart RA, Yu W. 1995. Identifying tumor suppressors in genetic
mosaics: the Drosophila lats gene encodes a putative protein kinase. *Development* **121**:
1053–1063.

Yatsenko AS, Shcherbata HR. 2021. Distant activation of Notch signaling induces stem cell
niche assembly. *Plos Genet* **17**: e1009489.

Yu G, Wang L-G, Han Y, He Q-Y. 2012. clusterProfiler: an R Package for Comparing Biological
Themes Among Gene Clusters. *OMICS: A J Integr Biol* **16**: 284–287.

Yu J, Pacifico S, Liu G, Finley RL. 2008. DroID: the Drosophila Interactions Database, a
comprehensive resource for annotated gene and protein interactions. *BMC Genom* **9**: 461.

Zhang L, Ren F, Zhang Q, Chen Y, Wang B, Jiang J. 2008. The TEAD/TEF Family of
Transcription Factor Scalloped Mediates Hippo Signaling in Organ Size Control. *Dev Cell* **14**:
377–387.

Zhao B, Wei X, Li W, Udan RS, Yang Q, Kim J, Xie J, Ikenoue T, Yu J, Li L, et al. 2007.
Inactivation of YAP oncoprotein by the Hippo pathway is involved in cell contact inhibition
and tissue growth control. *Genes Dev* **21**: 2747–2761.

Zhao B, Ye X, Yu J, Li L, Li W, Li S, Yu J, Lin JD, Wang C-Y, Chinnaiyan AM, et al. 2008. TEAD
mediates YAP-dependent gene induction and growth control. *Gene Dev* **22**: 1962–1971.

Zheng Y, Pan D. 2019. The Hippo Signaling Pathway in Development and Disease. *Dev Cell*
50: 264–282.

Zhu S, Lin S, Kao C-F, Awasaki T, Chiang A-S, Lee T. 2006. Gradients of the Drosophila
Chinmo BTB-Zinc Finger Protein Govern Neuronal Temporal Identity. *Cell* **127**: 409–422.

Figure 1: Experimental design to generate stage-specific transcriptomes of in somatic and germ line larval ovarian cells under wild type, *hpo* and *yki* loss of function conditions. (A) Left to right: Location of larval ovaries (indicated as white circles in the larva); location of adult ovaries in female abdomen (indicated in white within the abdomen); a single adult ovary; a single ovariole; a single germarium with terminal filaments in black. **(B)** The three developmental stages during terminal filament (in black) formation studied herein. **(C)** Somatic (red) and germ line (blue) cells were marked for FACS sorting using appropriate GAL4 drivers as described in Materials & Methods. **(D)** Summary of the tissue-specific RNAi knockdown treatments assessed at each stage in somatic and germ line cells. GFP labelled somatic cells **(E)** and germ cells **(F)** were sorted using FACS at the three developing stages of larval ovaries. Data from early-stage germ cells (grey), were not used for analysis; see Materials & Methods. **(G)** Sorted ovarian cell types were processed for mRNA extraction, cDNA library preparation and high throughput sequencing. h AEL = hours after egg laying.

Figure 2: Distinct transcriptional impact of Hpo signaling on germ line and soma across ovarian development . (A) Principal Component Analysis (PCA) of all samples differentiates germ cells from somatic cells along PC1. (B) PCA of somatic transcriptomes showing clustering by time-point in PC1, and the knockdown effect in PC2. **(C)** PCA of germ line transcriptomes in which control and knockdown libraries cluster separately, but the *hpo* and *yki* knockdowns cluster together. **(D)** Number of significantly upregulated and downregulated genes (adjusted p-value < 0.05) in each cell type with grouped stages when comparing each knockdown to the control library. For somatic cells **(E)** and germ cells **(F)** a Venn diagram shows the differentially up- and down- regulated genes under each knockdown, including number and percentage of genes in each intersection.

866 **Figure 3: Perturbing *hpo* or *yki* levels has a larger transcriptomic impact on germ cells than**
867 **somatic cells.** Shown are significantly enriched (Benjamini-Hochberg adjusted p-value<0.05) KEGG
868 pathways within the groups of significantly upregulated (Up) and downregulated (Down) genes in germ
869 cells and somatic cells under *hpo* and *yki* RNAi conditions (grouping mid and late stages together).

Figure 4: Prediction of chinmo as a novel germ cell-specific candidate regulator of *hpo* and/or *yki*. Expression of **(A)** *hpo* and **(B)** *yki* as vst-normalized counts in germ cells (left panels, blue dots) and somatic cells (right panels, red dots) in each RNA-Seq library (color shades indicate genetic condition). The *hpo* and *yki* transcripts appear significantly downregulated their corresponding RNAi conditions. The *yki* reduction in germ cells mid stages is significant when the control sample with anomalously low levels of *yki* (circled in red) is removed (Supplemental Figure S3). Additionally, *yki* appears significantly downregulated under *hpo* knockdown. Statistical significance levels from the differential expression analysis are codified as *** = adjusted p-value<0.01; ** = adjusted p-value<0.05; [a] = adjusted p-value = 0.055, n.s. = not significant. **(C)** Graph showing *hpo*, *yki*, and their only common direct neighbor in our network of protein-protein interactions, *chinmo*. Protein-protein candidate interaction edges are represented in dark grey, and transcription factor-gene candidate interactions are represented in light grey arrows with pointing to the gene predicted to be regulated by the TF. The TF *chinmo* appears has the potential to regulate both *hpo* and *yki*, can physically interact with *hpo*, and can potentially be regulated by *yki*.

Figure 5: Chinmo regulates germ cell number autonomously and somatic cell number and morphogenesis non-autonomously in the developing ovary. (A) Chinmo is detected in germ cell nuclei at mid and late stages of larval development. Red = Vasa; Chinmo = green, scale bars = 20μm. **(B)** *chinmo RNAi* in the germ cells causes an increase in ovariole number. Percentage difference in adult ovariole number where *UAS:chinmo [RNAi]*, *UAS:hippo [RNAi],* or *UAS:yorkie [RNAi]* are driven in the somatic cells or in the germ cells. n=20 for each genotype. **(C)** Abrogation of Hpo signaling in the germ cells causes a change in Chinmo levels in germ cell nuclei. Percentage difference in detected fluorescence levels of Chinmo protein in larval ovarian germ cell nuclei where *UAS:chinmo [RNAi]*, *UAS:hippo [RNAi]* or *UAS:yorkie [RNAi]* are driven in the somatic cells in the germ cells. n $\geq$ 50 nuclei from five ovaries. **(D)** *chinmo RNAi* in germ cells causes an increase in germ cell number. Percentage difference in germ cell number where *UAS:chinmo [RNAi]*, *UAS:hippo [RNAi]* or *UAS:yorkie [RNAi]* are driven in germ cells. **(E)** *chinmo* RNAi in germ cells causes an increase in Terminal Filament (TF) number, Total Terminal Filament Cell (TFC) number and in the number of TFCs per TF. Percentage differences in TF number and TFC number in each ovary, and the number of TFCs per TF where *UAS:chinmo [RNAi]*, *UAS:hippo [RNAi]* or *UAS:yorkie [RNAi]* were driven in the germ cells. In **B** through **E**, grey bars = comparisons to *Gal4* controls; blacks bars = comparisons to *UAS* controls. Statistical significance was calculated using a student's two-tailed t-test with unequal variance. (** = p<0.01, * = p<0.05, ns = p>0.05).

Figure 6. Hypothesis: Chinmo regulates Hpo signaling activity in the germ cells but not somatic cells. We previously showed that the components of the Hippo signaling pathway are present in both the germ and somatic cells, (Sarikaya 2015), and that the somatic cells (red) function under the influence of the canonical Hippo pathway (Sarikaya 2015). Under this model, the reduction of either *hpo* or *yki* causes a change in germ cell number. In germ cells (blue), Chinmo is present, which adds a secondary Hippo-independent repression of *yki*-dependent cell proliferation. Because our data cannot determine whether Chinmo acts directly on the localization of Yki to the nucleus, or indirectly by repressing cell proliferation downstream of Yki, the Chinmo-Yki relationship is indicated by grey dashed lines.