Adropin protects against cardiac remodeling and metabolic dysfunction in a mouse model of HFpEF

Bellina A.S. Mushala[1,2,3], Michael W. Stoner[1,2,3], Janet R. Manning[1,2,3], Paramesha Bugga[1,2,3], Nisha Bhattarai[1,2,3], Maryam Sharifi-Sanjani[1,2,3], Brenda McMahon[1,$], Amber Vandevender[2,4], Steven J. Mullet[7], Brett A. Kaufman[1,2,3], Sruti S. Shiva[1,2,5], Cheng Zhang[6], Eric Goetzman[8], Stephen Y. Chan[1,3], Stacy L. Gelhaus[6,7], Michael J. Jurczak[2,4], and Iain Scott[1,2,3]*

[1]Vascular Medicine Institute, [2]Center for Metabolism and Mitochondrial Medicine, [3]Division of Cardiology, [4]Division of Endocrinology and Metabolism, and [5]Division of Classical Hematology, Department of Medicine, [6]Department of Pharmacology and Chemical Biology, [7]Health Sciences Mass Spectrometry Core, [8]Department of Pediatrics, University of Pittsburgh, Pittsburgh, PA 15261
[$]Current Address: Mount Aloysius College, Cresson, PA 16630

*To whom correspondence should be addressed:

Iain Scott, PhD, FAHA, FCVS

Department of Medicine

University of Pittsburgh

BST W1048

200 Lothrop Street

Pittsburgh, PA 15261

iain.scott@pitt.edu

Keywords: Adropin, HFpEF, Mitochondria, Fatty Acid Oxidation, Glucose Oxidation, O-GlcNAcylation

ABSTRACT

Cardiometabolic heart failure with preserved ejection fraction (HFpEF) is a heterogenous metabolic disease, which in the heart presents as left ventricle diastolic dysfunction, ventricular stiffness, and myocardial structural remodeling. Deleterious changes in cardiac metabolism are central to HFpEF pathophysiology, and proposed treatments for the disease have focused on repairing these defects. In this study, we used a preclinical mouse model that recapitulates cardiometabolic HFpEF to elucidate the molecular mechanisms driving cardiac dysfunction, and tested whether recombinant Adropin (a liver- and brain-derived endogenous peptide hormone) could reverse observed defects. We show that long-term treatment with Adropin reversed multiple markers of HFpEF-related cardiac dysfunction (including fibrosis, diastolic dysfunction, and cardiomyocyte hypertrophy). Using untargeted metabolomics, we found that Adropin treatment reduced hexosamine biosynthesis pathway activity, leading to a reduction in the inhibitory O-GlcNAcylation of the cardiac fatty acid oxidation enzyme long chain acyl-CoA dehydrogenase (LCAD). Our results suggest that Adropin may restore cardiac metabolic function in HFpEF, and that targeting this pathway may be a novel therapeutic avenue for this disease.

INTRODUCTION

Heart failure is a complex clinical syndrome that results from any structural or functional impairments of ventricular filling and/or ejection of blood, and remains the leading cause of morbidity and mortality worldwide (Heidenreich et al., 2023). Heart failure with preserved ejection fraction (HFpEF), a subtype of heart failure, is primarily characterized by cardiac diastolic dysfunction. This presentation of the disease currently accounts for >50% of all heart failure cases, with a global prevalence of approximately 32 million (Shah et al., 2020; Redfield and Borlaug, 2023). HFpEF is recognized as a heterogenous syndrome, whose underlying pathophysiological mechanisms include several extracardiac abnormalities such as metabolic derangements, arterial hypertension, microvascular endothelial dysfunction, inflammation, and renal insufficiency (Shah et al., 2016, Shah et al., 2020).

Partly due to a lack of specifically-designed drugs for the disease, treatment for HFpEF currently focuses on the management of comorbidities (e.g. obesity, type 2 diabetes, hypertension), with new guidelines recommending the use of glycemia-reducing therapeutics (e.g. SGLT2 inhibitors) that improve cardiovascular outcomes in heart failure (Kittleson et al., 2023). The recent STEP-HFpEF trial in obesity-related HFpEF demonstrated that therapy-mediated reductions in body weight may also be a key tool in disease regression (Kosiborod et al., 2023). These developments suggest that approaches targeting metabolic defects can offer substantial benefits in this disease. On this basis, we examined whether treatment with recombinant Adropin, a brain- and liver-derived endogenous peptide hormone (Kumar et al., 2008), may have beneficial effects in a preclinical model of cardiometabolic HFpEF.

Previous studies have shown that Adropin improves physical activity (Wong et al, 2014, Ganesh-Kumar et al., 2012), attenuates hepatic steatosis and injury (Kumar et al., 2008, Chen et al., 2019), and reduces vascular stiffness in diet- induced obesity (Jurrissen et al., 2022). In the heart, we recently demonstrated that short-term Adropin treatment ($\leq$ 3 days) can improve cardiac fuel

substrate flexibility by restoring glucose oxidation in the diabetic heart (Thapa et al., 2019b). Combined, these beneficial features raised the question of whether this peptide can be used as a therapeutic intervention in HFpEF. Here, we show that long-term treatment (4 weeks) with recombinant Adropin restored cardiac diastolic function, reduced cardiac hypertrophy and fibrosis, and improved whole-body glucose tolerance. These changes were linked to a reversal of metabolic remodeling and improved fatty acid oxidation enzyme activity, suggesting that Adropin treatment mitigates the impact of HFpEF stimuli via improved cardiometabolic function.

MATERIALS AND METHODS

Animal Care and Use

Male C57BL/6J 8-week-old mice were obtained from the Jackson Laboratory and maintained on chow for 4 weeks to acclimate to their new environment. Animals were housed in the University of Pittsburgh animal facility under standard conditions, with *ad libitum* access to water and food, and maintained on a constant 12-hour light/dark cycle. At 12 weeks of age, mice were exposed to normal chow (Chow; 60% carbohydrate, 26% protein, 14% fat; ProLab IsoPro RMH 3000), or a combination of high-fat diet (HFD; 20% carbohydrate, 20% protein, 60% fat; Research Diets D12492) plus N-ω-nitro-L-arginine methyl ester (L-NAME: 0.5 g/L, pH 7.4; ThermoFisher) supplemented in drinking water to induce HFpEF (Schiattarella et al., 2019). After 8 weeks of diet, mice were randomly assigned to a control or treatment group receiving daily intraperitoneal injections of either vehicle or recombinant Adropin (Adr.) at 450 nmol/kg animal weight, prepared in PBS with 0.1% BSA, for the remaining 4 weeks of the study. L-NAME supplemented drinking water was changed weekly, and adropin was made fresh and sterile filtered before each use. Animals were euthanized by isofluorane anesthesia followed by cervical dislocation. Experiments were conducted in compliance with National Institutes of Health guidelines, and followed procedures approved by the University of Pittsburgh Institutional Animal Care and Use Committee.

Echocardiography

Mice were anesthetized using isofluorane (1.5%–2.0% v/v by inhalation) and monitored for cardiac functional parameters in the supine position using a Visual Sonics Vevo 3100. Core temperature was maintained at 37 °C by imaging mice while on a heating pad, and heart rates were kept consistent between experimental groups (~400–500 beats per min). Short axis M-mode was used to assess LV dimensions and motion patterns, pulsed-wave (PW) doppler mode to evaluate blood flow velocities across the mitral valve, and tissue doppler mode to measure mitral annular plane velocity. Doppler profiles were acquired in the parasternal long axis (PLAX) apical 4 chamber view. The left atrial area

was quantified in the apical 4-chamber view by tracing the border of the left atrium. Markers of systolic and diastolic function were calculated using standard echocardiography equations. At the end of the procedure, all mice recovered from anesthesia without difficulties. Image analysis was performed independently by a blinded sonographer, and all parameters were measured at least 3 times with averages presented.

Intraperitoneal Glucose Tolerance Test (IPGTT)

Glucose tolerance tests were performed as previously described with minor modifications (Jurczak et al., 2011). After an overnight fast (~16 h), mice were set-up in restrainers with tails snipped <5mm to prime for blood collection and left to acclimate under a heated bulb for 2 h prior to GTT start time at 9 a.m. Basal plasma samples (t = 0) were collected, and D-glucose was diluted in water (20% final concentration) and sterile filtered for intraperitoneal administration at 1.5 g/kg body weight. Blood glucose was measured by tail bleed at set time points using a Bayer Contour Next EZ handheld glucometer (t = 15, 30, 45, 60, and 120 min).

Histology

Preparation and staining of all histological samples were conducted by the Pitt Biospecimen Core at the University of Pittsburgh. All mice were sacrificed, and tissue harvested for analysis within one hour of completion of the IPGTT study. The apex of the heart was excised from each mouse following euthanasia, fixed in 10% buffered formalin phosphate overnight at shaking incubation, washed 3 times for 5 min with 1X phosphate-buffered saline (PBS), and then transferred to 70% ethanol. Samples were then embedded in paraffin, sectioned into 4 μm slides and stained with hematoxylin and eosin (H&E), wheat germ agglutinin (WGA) or Masson's Trichrome for analysis. Histology sections were visualized with the Evos FL Auto 2 Microscope, observed changes in structure were quantified using ImageJ software, and representative images are shown.

Transcriptomics

Total RNA was isolated from the left ventricle of the heart using the RNeasy Plus Mini Kit (Qiagen REF 74134). Approximately 2 µg RNA was used for bulk RNA sequencing performed by Azenta/GENEWIZ, based on company recommendations to identify differential gene expression patterns between the experimental groups.

Quantitative Metabolomics

Metabolic quenching and polar metabolite pool extraction was performed by adding ice cold 80% methanol (aqueous) at a ratio of 1:15 wt input tissue:vol. (13C1)-creatinine, (D3)-taurine, (D3)-lactate and (D3)-alanine (Sigma-Aldrich) were added to the sample lysates as an internal standard for a final concentration of 10 µM. Samples are homogenized using an MP Bio FastPrep system using Matrix D (ceramic sphere) for 60 s at 60 hz. The supernatant was then cleared of protein by centrifugation at 16,000 g. Cleared supernatant (2 µL) was subjected to online LC-MS analysis. Analyses were performed by untargeted liquid chromatography-high-resolution mass spectrometry (LC-HRMS). Briefly, samples were injected via a Thermo Vanquish UHPLC and separated over a reversed phase. Thermo HyperCarb porous graphite column (2.1×100 mm, 3 µm particle size) maintained at 55 °C. For the 20 min LC gradient, the mobile phase consisted of the following: solvent A (water/0.1% FA) and solvent B (ACN/0.1% FA). The gradient used was: 0-1 min 1% B, increase to 15% B over 5 min, continue increasing to 98% B over 5 min, hold at 98% B for 5 min, re-equillibrate at 1% B for 5 min. The Thermo IDX tribrid mass spectrometer was operated in both positive and negative ion mode, scanning in ddMS2 mode (2 µscans) from 70 to 800 m/z at 120,000 resolution with an AGC target of $2e^5$ for full scan, $2e^4$ for ms2 scans using HCD fragmentation at stepped 15, 35, 50 collision energies. Source ionization setting was 3.0 and 2.4 kV spray voltage, respectively, for positive and negative mode. Source gas parameters were 35 sheath gas, 12 auxiliary gas at 320 °C, and 8 sweep gas. Calibration was performed prior to analysis using the PierceTM FlexMix Ion Calibration Solutions (Thermo Fisher Scientific). Integrated peak areas were then extracted manually using Quan Browser

(Thermo Fisher Xcalibur ver. 2.7). Untargeted differential comparisons were performed using Compound Discoverer 3.0 (Thermo Fisher) to generate a ranked list of significant compounds with tentative identifications from BioCyc, KEGG, and internal compound databases. Purified standards were then purchased and compared in retention time, m/z, along with ms2 fragmentation patterns to validate the identity of significant hits.

Protein Isolation and Immunoblotting

Heart tissues were rapidly harvested following euthanasia, weighed, RV discarded, and LV flash-frozen in liquid nitrogen. For protein isolation, tissues were minced and lysed in CHAPS buffer (1% CHAPS, 150 mM NaCl, 10 mM HEPES, pH 7.4) using a VWR 4-Place Mini Bead Mill, then incubated on ice for ~ 2.5 h. Homogenates were spun at 10,000 g at 4 °C for 10 min, and the supernatants collected for immunoblotting. For immunoblotting, protein lysates were quantitated using a BioDrop µLITE Analyzer, prepared in LDS sample buffer, separated using Bolt SDS-PAGE 4–12% or 12% Bis–Tris Plus gels, and transferred to nitrocellulose membranes (all Invitrogen). Membranes were blocked using SuperBlock (PBS) Blocking Buffer and incubated overnight in the following primary antibodies: mouse O-GlcNAc (9875), rabbit GFAT (D12F4), rabbit OGT (D1D8Q), rabbit OGA (E9C5U), and rabbit GAPDH (2118S) from Cell Signaling Technologies; rabbit LCAD (17526-1-AP) from Protein Tech. Protein loading was confirmed using GAPDH as a loading control. Fluorescent anti-goat or anti-rabbit secondary antibodies (red, 700 nm; green, 800 nm) from LiCor were used to detect expression levels. Images were obtained using Licor Odyssey CLx System and protein densitometry was measured using the LiCor Image Studio Lite Ver. 5.2 Software.

Co-Immunoprecipitation

For co-immunoprecipitation experiments, protein lysates were harvested in CHAPS lysis buffer (1% CHAPS, 150 mM NaCl, 10 mM HEPES, pH 7.4), and 500 µg of protein was incubated overnight at 4 °C with 5 µL mouse O-GlcNAc antibody (Cell Signaling). Immunocaptured proteins were isolated

using Protein-G and Protein-A agarose beads (Cell Signaling Technology, catalog number 9007), washed 5 times with 500 µL 1% CHAPS buffer, and then eluted in LDS sample buffer (Life Technologies) at 95 °C. Samples were separated on 4-12% Bis–Tris Bolt gels, transferred to nitrocellulose membranes, and probed with appropriate antibodies. Images were obtained using LiCor Odyssey CLx System and protein densitometry was measured using the LiCor Image Studio Lite Ver. 5.2 Software.

Biochemical Assays

To assess the activity of long chain acyl-CoA dehydrogenase enzymes (LCAD), homogenized protein samples were incubated with palmitoyl-CoA as described previously (Thapa et al., 2017). Briefly, ~10 µg of protein was incubated with 0.1 M potassium phosphate, 50 µM 2,6- dichlorophenolindophenol, 2 mM phenazine ethosulfate, 0.2 mM N-ethylmaleimide, 0.4 mM potassium cyanide, and 0.1% Triton X-100 at 37 °C for 4 min. The reaction was initiated with 60 µM palmitoyl-CoA, and the rate of absorbance change was measured/observed at 600 nm over 45 min. Activities were converted to moles of substrate oxidized/min/mass of protein. *In vitro* LCAD activity was measured after incubating a reaction mixture of 4 µg of recombinant LCAD, 0.8 mM UDP-GlcNAc (Sigma, Catalog # U4375), 40 µg/mL of recombinant human O-GlcNAc Transferase (rhOGT) (Catalog # 8446-GT) in assay buffer (25 mM Tris, 10 mM $CaCl_2$, pH 7.5 and 10 mM $MgCl_2$) at 37 °C for 30 min.

Statistics

Means ± SEM were calculated for all data sets. Data sets containing three or more groups were analyzed using one-way ANOVA with Tukey's post-hoc testing to determine differences between genotypes and feeding/treatment groups. Data sets containing two groups were analyzed using Student's T-test. Time course data was analyzed using one-way ANOVA with Sidak's post- hoc testing. $P \leq 0.05$ was considered statistically significant. Statistical analyses were performed using GraphPad Prism 9.5 Software.

RESULTS

Circulating Adropin levels are diminished in HFpEF patients

Circulating Adropin levels are known to decrease in patients with type 2 diabetes (Soltani et al., 2023), aging (Yang et al., 2018), and hypertension (Gulen et al, 2016). While these are all comorbidities for HFpEF, there have been no published studies examining circulating Adropin levels in patients with this disease. We therefore measured plasma levels of Adropin in deidentified age- and gender-matched control and HFpEF patient samples (n = 19-20) obtained from a University of Pittsburgh biobank **(Sup. Table 1)**. Consistent with the studies in comorbid disease states, there was a significant decrease in plasma Adropin levels in HFpEF patients relative to non-affected controls **(Fig. 1A)**. These data suggest that a reduction in circulating Adropin levels may be linked to HFpEF disease development or progression.

Recombinant Adropin treatment improves glucose tolerance in HFpEF without changes in body weight

The potential link between reduced circulating Adropin and HFpEF in patients supported the premise that therapeutic restoration of Adropin levels may be protective against the disease. We therefore used a recently developed two-hit mouse model of HFpEF (Schiattarella et al, 2019) to test this hypothesis. Male C57BL/6J mice were placed on either a chow (control) or HFpEF (60% high fat diet plus L-NAME-supplemented drinking water) diet for 8 weeks. After 8 weeks, HFpEF mice were randomly given either vehicle (PBS) or Adropin (450 nmol/kg/day; Thapa et al., 2019b) by intraperitoneal (I.P.) injection once daily for 4 weeks **(Fig. 1B)**. Daily Adropin treatment significantly increased circulating Adropin levels in HFpEF mice (15.9 vs. 2.2 ng/mL, P = 0.0067), demonstrating that administration of recombinant Adropin could restore plasma concentrations to those observed in healthy humans.

The recent STEP-HFpEF Phase 3 clinical trial showed that the GLP-1 receptor agonist semaglutide improved patient-reported outcomes in HFpEF, which was at least partially related to weight loss (Kosiborod et al, 2023). We therefore first examined whether Adropin treatment promoted weight loss in mice, and found that there was no significant difference in body weight between vehicle- and Adropin-treated HFpEF animals **(Fig. 1C,D)**. Our group and others have previously shown that short-term Adropin treatment ($\leq$ 3 days) leads to improved whole-body glucose tolerance in diet-induced obese mice (Gao et al., 2015; Thapa et al., 2019a). We next tested whether long-term Adropin treatment (4 weeks) in HFpEF mice had the same effect, and found that there was a significant decrease in insulin resistance in HFpEF mice treated with Adropin relative to non-treated HFpEF controls **(Fig. 1E,F)**. Combined, these data suggest that Adropin treatment may have a positive effect on whole-body physiology by reducing glucose intolerance. However, unlike semaglutide in humans, this improvement appears to be unrelated to the beneficial effects of weight loss.

Recombinant Adropin treatment limits cardiac diastolic dysfunction in a mouse model of HFpEF

We next examined the effect of Adropin treatment on cardiac functional parameters and tissue structure **(Fig. 2A)**. Mice with HFpEF displayed a significant increase in cardiac weight and a reduction in cardiac output, both of which were absent in Adropin-treated HFpEF mice **(Fig. 2C,D)**. As expected from a HFpEF model, all mouse groups displayed normal systolic function, as measured by left ventricular ejection fraction **(Fig. 2E)**. Both control and Adropin-treated HFpEF groups displayed a non-significant ~25% increase in E/e' ratio, and a significant increase in mitral annulus tissue velocity during early diastole (e'), suggestive of mild-to-moderate diastolic dysfunction **(Fig. 2F,H)**. Adropin-treated mice displayed a significant decrease in isovolumetric relaxation time (IVRT), a significant decrease in myocardial performance index (MPI; also known as Tei index), and a trend towards a decreased E/A ratio relative to untreated HFpEF mice **(Fig. 2G,I,J)**. Combined, these data indicate that Adropin treatment may reduce diastolic dysfunction in HFpEF by improving ventricular relaxation

capacity (IVRT and MPI), whilst having a more limited impact on echocardiography measures of left ventricle filling pressures (such as E/e' ratio and e').

Long-term Adropin treatment prevents cardiomyocyte hypertrophy and reduces cardiac fibrosis

Impairment of cardiac relaxation (such as IVRT) often correlates with cardiac remodeling and increased myocardial fibrosis (e.g. Brilla et al., 2000) in disease states. Histological analysis of cardiac tissue sections demonstrated that untreated HFpEF mice displayed a significant increase in myocardial interstitial fibrosis and cardiomyocyte cross-sectional area, which was completely reversed following long-term Adropin treatment **(Fig. 3A-C)**. The positive impact of Adropin on reducing myocardial fibrosis was evident in bulk RNA-seq analysis of untreated vs. Adropin-treated HFpEF hearts **(Sup. Table 2)**, with extracellular matrix processes being identified as the top hit in pathway analyses **(Fig. 3D)**. Closer examination demonstrated that multiple genes related to extracellular matrix formation and fibrosis (e.g. *Postn, Fn1, Col1a1*) were significantly downregulated in HFpEF hearts after Adropin treatment **(Fig. 3E, Sup. Fig. 1)**. Combined with the earlier studies, these data demonstrate that long-term Adopin treatment in HFpEF promoted whole-body metabolic homeostasis, improved myocardial relaxation capacity, prevented maladaptive cardiac tissue remodeling, and limited cardiomyocyte hypertrophy.

Adropin treatment limits cardiac protein O-GlcNAcylation in HFpEF mice

The decrease in cardiac fibrosis in Adropin-treated HFpEF mice is likely to account for at least some of the functional improvements observed, particularly in terms of myocardial relaxation. However, it is unlikely to explain all of the improvements that were found to result from Adropin treatment, such as improved metabolic homeostasis. To better understand the underlying processes involved, we next performed untargeted metabolomic analyses of hearts from untreated and Adropin-treated HFpEF mice. Our analysis demonstrated that metabolites related to the hexosamine biosynthesis pathway

(HBP) were specifically enriched in Adropin-treated vs. untreated HFpEF hearts **(Fig. 4A)**. As the HBP provides substrates for the generation of UDP-GlcNAc, the co-factor for protein O-GlcNAcylation **(Fig. 4B)**, we next examined the metabolites and enzymes that regulate this pathway.

Detailed analysis of our metabolomic data revealed an accumulation of GlcNAc-6-P, CoA, Uridine, and Cytidine in Adropin-treated HFpEF hearts relative to control and untreated HFpEF mice **(Fig. 4C-F)**. As these metabolites feed into the HBP at different points, their general accumulation in Adropin-treated mice suggests a possible overall reduction in metabolites entering this pathway to generate UDP-GlcNAc, the co-factor for protein O-GlcNAcylation. Corroborating our metabolomic data, immunoblotting using an O-GlcNAc-specific antibody demonstrated that there was a significant decrease in cardiac protein O-GlcNAcylation in Adropin-treated HFpEF hearts relative to untreated HFpEF controls **(Fig. 4G)**. O-GlcNAcylation is controlled at the protein level by the rate-limiting enzyme GFAT, and by the opposing activities of the O-GlcNAc transferase (OGT) and O-GlcNAcase (OGA) which, respectively, add and remove this modification to/from other proteins (e.g., Umapathy et al., 2021, Tran et al., 2020, Prakoso et al., 2022). Unexpectedly, despite the difference in O-GlcNAcylation levels observed in untreated and Adropin-treated HFpEF hearts, there were no significant changes in the abundance of any of these regulatory enzymes **(Fig. 4H-K)**. Combined, these data suggest that part of the function of Adropin in protecting from HFpEF is mediated by its modulatory effect on cardiac protein O-GlcNAcylation. By comparing our metabolomic and biochemical data, it would appear that this modulation occurs at the level of HBP metabolite abundance, rather than as an effect on HBP regulatory enzyme activity.

Adropin-mediated reductions in LCAD O-GlcNAcylation improve its enzymatic activity

Studies in humans and mice have demonstrated that pathways related to cardiac energy metabolism, particularly those in fatty acid utilization, are negatively impacted by HFpEF progression (Tong et al., 2021; Hahn et al., 2023; O'Sullivan et al., 2024). We therefore examined whether specific fatty acid

oxidation enzymes were impacted by the increased protein O-GlcNAcylation observed above. Using O-GlcNAc-specific immunoprecipitation, we observed that the beta oxidation enzyme long-chain acyl-CoA dehydrogenase (LCAD) displayed elevated O-GlcNAcylation in HFpEF hearts, which was reversed by Adropin treatment **(Fig. 5A,B)**. We examined the effect of O-GlcNAcylation on LCAD enzymatic activity *ex vivo*, and found that the ability of LCAD to oxidize C16 palmitoyl-CoA was reduced in vehicle-treated HFpEF hearts, but not those treated with Adropin **(Fig. 5C)**. Combined, there was a significant negative correlation between LCAD O-GlcNAcylation abundance and its enzymatic activity across all cardiac tissues **(Fig. 5D)**. To confirm the regulatory effect of increased O-GlcNAcylation on LCAD activity, we performed a series of *in vitro* assays. Recombinant LCAD activity was significantly decreased in the presence of the O-GlcNAcylation substrate (UDP-GlcNAc) and transferase enzyme (OGT), but not in the presence of the substrate alone **(Fig. 5E)**, demonstrating the regulatory requirement of the modification. Combined, our data suggest that increased O-GlcNAcylation of fatty acid enzymes like LCAD in HFpEF may decrease their enzymatic activity, which can be reversed by Adropin treatment *in vivo*.

DISCUSSION

Previous studies on Adropin signaling in cardiometabolic disease have focused on its effects on fuel energy metabolism, with limited exploration of its role on myocardial structure and function (reviewed in Mushala et al., 2021). Short-term Adropin treatment (≤ 3 days) improved cardiac work and efficiency *ex vivo*, which was accompanied by enhanced insulin signaling in lean mice (Altamimi et al., 2019). In contrast, exposure to a high-fat diet inhibited acute Adropin-driven improvements in contractile function, and insulin stimulation failed to compensate for this impaired cardiac contractility in diet-induced obese mice *ex vivo* (Thapa et al., 2022). In this study, we demonstrate that chronic Adropin treatment *in vivo* restores cardiac function, specifically left ventricular diastolic function, in the failing diabetic myocardium of mice.

Diastolic dysfunction is a hallmark of not only HFpEF, but also diabetic cardiomyopathy (DCM), where relaxation and filling pressure of the left ventricle is impaired (reviewed in Jia et al., 2018). Abnormalities of LV passive elasticity are often caused by structural impediments, such as increased cardiomyocyte diameter and extracellular matrix deposition, which contribute to diastolic dysfunction (Aurigemma et al., 2006). Our data shows that Adropin treatment attenuates left ventricular hypertrophy and fibrosis **(Fig. 3)**, indicative of improved structural remodeling. In addition, transcriptomic analysis demonstrated similar increases in pro-fibrotic pathways as seen in HFpEF patients (Ye et al., 2023) which was attenuated upon Adropin treatment **(Fig. 3; Sup. Fig. 1)**, providing further evidence of restored myocardial structure and function. These data match similar findings in a model of systemic sclerosis, where Adropin treatment significantly reduced fibroblast activation and extracellular matrix deposition (Liang et al., 2024). Therefore, our findings collectively highlight the therapeutic potential of prolonged Adropin treatment in preventing cardiac remodeling and functional decline in HFpEF and/or DCM.

Data presented in this study add to our growing understanding of the transcriptomic and metabolomic signature of the HFD + L-NAME HFpEF model, providing preclinical evidence of known mechanisms underlying the pathogenesis of cardiometabolic HFpEF in patients, such as fuel substrate inflexibility (Hahn et al., 2023). Unlike previous acute studies from our group and others, the ability of Adropin treatment to regulate cardiac substrate utilization was independent of the inhibitory effects of PDK4 or GCN5L1 on pyruvate dehydrogenase activity (Thapa et al., 2019b, Altamimi et al., 2019). Instead, we found that Adropin treatment may reduce metabolite entry into and/or flux through the HBP, which inhibits cardiac protein O-GlcNAcylation to promote fatty acid oxidation **(Figs. 4,5)**. The HBP is a non-oxidative branch of glycolysis where the rate-limiting enzyme, GFAT, uses fructose-6-phosphate and glutamine to catalyze its conversion into glucosamine-6-phosphate, which is subsequently metabolized through a series of reactions utilizing substrates from major metabolic pathways to generate the end-product UDP-GlcNAc (Umapathi et al., 2021). UDP-GlcNAc serves as the co-factor for the attachment of an O-linked β-N-acetyl glucosamine moiety (O-GlcNAc) to serine and threonine residues of nuclear, cytoplasmic, and mitochondrial proteins, using 2-3% of the total cellular glucose found in the heart (Tran et al., 2020).

The dynamic on- and off-cycling of the O-GlcNAc modification is regulated by the activity of two enzymes, O-GlcNAc transferase (OGT) and O-GlcNAcase (OGA), respectively (Prakoso et al., 2022). Several studies have implicated aberrant regulation of the HBP, and consequent maladaptive protein O-GlcNAcylation, in heart failure (e.g., Umapathy et al., 2021, Tran et al., 2020, Prakoso et al., 2022). These studies have identified changes in OGT and OGA enzyme expression as the predominant mechanism that triggers excessive protein O-GlcNAcylation in diabetic hearts. However, here we found that changes in HBP metabolite abundance following Adropin treatment appeared to decrease UDP-GlcNAc availability for cardiac protein O-GlcNAcylation levels **(Fig. 4)**.

In this study, we identified the fatty acid oxidation enzyme LCAD as a target of O-GlcNAcylation in HFpEF mice, and demonstrated that Adropin treatment could limit this deleterious modification **(Fig. 5)**. LCAD O-GlcNAcylation was associated with decreased cardiac enzyme activity in HFpEF mouse hearts, which was restored upon Adropin treatment *ex vivo* **(Fig. 5)**. The inhibitory effect of O-GlcNAcylation on LCAD enzyme activity was confirmed *in vitro* using recombinant protein assays, suggesting that Adropin treatment may augment cardiac fatty acid oxidation activity via this mechanism. Further studies need to be conducted to determine whether Adropin-mediated regulation of cardiac LCAD O-GlcNAcylation regulates fatty acid utilization in the heart *in vivo.*

Our study has three notable limitations. Firstly, the exclusive use of male animals prohibits the investigation into possible sex differences in Adropin-mediated effects observed in the heart. Previous studies using the current HFpEF model have demonstrated that female sex is protective against cardiac dysfunction (Tong et al., 2019). On this basis, our study exclusively examined male mice, and it is unknown whether the findings are relevant for female mice. Epidemiological data show that women display a higher prevalence of diastolic dysfunction (HFpEF) compared to men, particularly as age increases (Gori et al., 2014, Beale et al., 2018, Duca et al., 2018). Sexual dimorphism in metabolic homeostasis influences the development of cardiac dysfunction (Lindman et al., 2014, Lagou et al., 2021), suggesting a potential role of reproductive hormones in cardiometabolic HFpEF. Interestingly, hepatic Adropin production has been shown to be regulated by estrogen (Stokar et al., 2022), suggesting a potential difference in basal circulating Adropin levels between male and female mice. We did not detect any sex differences in circulating Adropin levels between male and female patients, regardless of HFpEF status (data not shown), indicating that the effects of estrogen may not lead to the same response in humans. However, the members of our female patient cohort were most likely to be in a post-menopausal state (mean age 67, range 56-75; **Sup. Table 1**), which may limit our ability to draw this conclusion, and more work in this area is required.

Secondly, our metabolomics data relied on a cross-sectional analysis of HFpEF hearts, which precluded flux measurements of metabolites related to the HBP. While cross-sectional metabolomics data has allowed us to make major advances of our understanding of HFpEF (e.g., Hahn et al, 2023), it does limit our understanding of how metabolite abundance changes in response to disease progression. Future work will likely include the use of stably-labeled isotope tracers for specific metabolites, or the measurement of metabolites entering and exiting the heart (e.g., O'Sullivan et al, 2024), to achieve a better understanding of HBP metabolite flux.

Finally, we acknowledge that despite our use of a well-established "two-hit" HFpEF mouse model (HFD+L-NAME; Schiattarella et al., 2019), we did not observe a significant change in E/e' or E/A ratios during our echocardiography analysis **(Fig. 2)**. While these ratios have become the *de facto* norm for detecting diastolic dysfunction in mice (as discussed in Sun et al., 2024), changes in age, heart rate, and anesthesia levels can affect the observed outcomes (Rottman et al., 2007; Sun et al., 2024). While changes in E/e' and E/A ratio were not observed, we did detect a significant increase in both isovolumetric relaxation time (IVRT) and myocardial performance index (MPI; also known as Tei index) in untreated HFpEF animals, which was absent following Adropin treatment **(Fig. 2)**. Both IVRT and MPI have been well validated as measures of diastolic dysfunction (e.g. Tei et al., 1996; Schnelle et al., 2018), and as such we believe that the improvements observed indicate that Adropin can mitigate aspects of diastolic dysfunction in HFpEF.

In summary, we found that HFpEF-driven disruptions to normal cardiac metabolism led to a potential increase in metabolites entering the HBP, resulting in increased in cardiac protein O-GlcNAcylation, and decreased fatty acid oxidation enzyme activity. This occurred in concert with an increase in diastolic dysfunction, fibrosis, and hypertrophy in HFpEF animals. Long-term Adropin treatment reversed the metabolic and structural remodeling observed in the heart, leading to improved myocardial relaxation capacity. These findings significantly increase our understanding of the

protective effects of Adropin in HFpEF, and highlight this pathway as a potential target for future

therapeutic interventions.

REFERENCES

Altamimi TR, Gao S, Karwi QG, Fukushima A, Rawat S, Wagg CS, Zhang L, Lopaschuk GD. Adropin regulates cardiac energy metabolism and improves cardiac function and efficiency. Metabolism. 2019 Sep;98:37-48. doi: 10.1016/j.metabol.2019.06.005. Epub 2019 Jun 14. PMID: 31202835.

Aurigemma GP, Zile MR, Gaasch WH. Contractile behavior of the left ventricle in diastolic heart failure: with emphasis on regional systolic function. Circulation. 2006 Jan 17;113(2):296-304. doi: 10.1161/CIRCULATIONAHA.104.481465. PMID: 16418449.

Beale AL, Meyer P, Marwick TH, Lam CSP, Kaye DM. Sex Differences in Cardiovascular Pathophysiology: Why Women Are Overrepresented in Heart Failure With Preserved Ejection Fraction. Circulation. 2018 Jul 10;138(2):198-205. doi: 10.1161/CIRCULATIONAHA.118.034271. PMID: 29986961.

Brilla CG, Funck RC, Rupp H. Lisinopril-mediated regression of myocardial fibrosis in patients with hypertensive heart disease. Circulation. 2000 Sep 19;102(12):1388-93. doi: 10.1161/01.cir.102.12.1388. PMID: 10993857.

Chen X, Xue H, Fang W, Chen K, Chen S, Yang W, Shen T, Chen X, Zhang P, Ling W. Adropin protects against liver injury in nonalcoholic steatohepatitis via the Nrf2 mediated antioxidant capacity. Redox Biol. 2019 Feb;21:101068. doi: 10.1016/j.redox.2018.101068. Epub 2018 Dec 6. PMID: 30684890; PMCID: PMC6351233.

Duca F, Zotter-Tufaro C, Kammerlander AA, Aschauer S, Binder C, Mascherbauer J, Bonderman D. Gender-related differences in heart failure with preserved ejection fraction. Sci Rep. 2018 Jan 18;8(1):1080. doi: 10.1038/s41598-018-19507-7. PMID: 29348420; PMCID: PMC5773700.

Ganesh Kumar K, Zhang J, Gao S, Rossi J, McGuinness OP, Halem HH, Culler MD, Mynatt RL, Butler AA. Adropin deficiency is associated with increased adiposity and insulin resistance. Obesity (Silver Spring). 2012 Jul;20(7):1394-402. doi: 10.1038/oby.2012.31. Epub 2012 Feb 9. PMID: 22318315; PMCID: PMC3905465.

Gao S, McMillan RP, Zhu Q, Lopaschuk GD, Hulver MW, Butler AA. Therapeutic effects of adropin on glucose tolerance and substrate utilization in diet-induced obese mice with insulin resistance. Mol Metab. 2015 Jan 17;4(4):310-24. doi: 10.1016/j.molmet.2015.01.005. PMID: 25830094; PMCID: PMC4354928.

Gori M, Lam CS, Gupta DK, Santos AB, Cheng S, Shah AM, Claggett B, Zile MR, Kraigher-Krainer E, Pieske B, Voors AA, Packer M, Bransford T, Lefkowitz M, McMurray JJ, Solomon SD; PARAMOUNT Investigators. Sex-specific cardiovascular structure and function in heart failure with preserved ejection fraction. Eur J Heart Fail. 2014 May;16(5):535-42. doi: 10.1002/ejhf.67. Epub 2014 Feb 23. PMID: 24574260.

Gulen B, Eken C, Kucukdagli OT, Serinken M, Kocyigit A, Kılıc E, Uyarel H. Adropin levels and target organ damage secondary to high blood pressure in the ED. Am J Emerg Med. 2016 Nov;34(11):2061-2064. doi: 10.1016/j.ajem.2016.04.014. Epub 2016 Apr 13. PMID: 27592461.

Hahn VS, Petucci C, Kim MS, Bedi KC Jr, Wang H, Mishra S, Koleini N, Yoo EJ, Margulies KB, Arany Z, Kelly DP, Kass DA, Sharma K. Myocardial Metabolomics of Human Heart Failure With Preserved Ejection Fraction. Circulation. 2023 Apr 11;147(15):1147-1161. doi: 10.1161/CIRCULATIONAHA.122.061846. Epub 2023 Mar 1. PMID: 36856044; PMCID: PMC11059242.

Heidenreich PA, Bozkurt B, Aguilar D, Allen LA, Byun JJ, Colvin MM, Deswal A, Drazner MH, Dunlay SM, Evers LR, Fang JC, Fedson SE, Fonarow GC, Hayek SS, Hernandez AF, Khazanie P, Kittleson MM, Lee CS, Link MS, Milano CA, Nnacheta LC, Sandhu AT, Stevenson LW, Vardeny O, Vest AR, Yancy CW. 2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure: Executive Summary: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. Circulation. 2022 May 3;145(18):e876-e894. doi: 10.1161/CIR.0000000000001062. Epub 2022 Apr 1. PMID: 35363500.

Jia G, Hill MA, Sowers JR. Diabetic Cardiomyopathy: An Update of Mechanisms Contributing to This Clinical Entity. Circ Res. 2018 Feb 16;122(4):624-638. doi: 10.1161/CIRCRESAHA.117.311586. PMID: 29449364; PMCID: PMC5819359.

Jurczak MJ, Lee HY, Birkenfeld AL, Jornayvaz FR, Frederick DW, Pongratz RL, Zhao X, Moeckel GW, Samuel VT, Whaley JM, Shulman GI, Kibbey RG. SGLT2 deletion improves glucose homeostasis and preserves pancreatic beta-cell function. Diabetes. 2011 Mar;60(3):890-8. doi: 10.2337/db10-1328. PMID: 21357472; PMCID: PMC3046850.

Jurrissen TJ, Ramirez-Perez FI, Cabral-Amador FJ, Soares RN, Pettit-Mee RJ, Betancourt-Cortes EE, McMillan NJ, Sharma N, Rocha HNM, Fujie S, Morales-Quinones M, Lazo-Fernandez Y, Butler AA, Banerjee S, Sacks HS, Ibdah JA, Parks EJ, Rector RS, Manrique-Acevedo C, Martinez-Lemus LA, Padilla J. Role of adropin in arterial stiffening associated with obesity and type 2 diabetes. Am J Physiol Heart Circ Physiol. 2022 Nov 1;323(5):H879-H891. doi: 10.1152/ajpheart.00385.2022. Epub 2022 Sep 9. PMID: 36083795; PMCID: PMC9602697.

Kittleson MM, Panjrath GS, Amancherla K, Davis LL, Deswal A, Dixon DL, Januzzi JL Jr, Yancy CW. 2023 ACC Expert Consensus Decision Pathway on Management of Heart Failure With Preserved Ejection Fraction: A Report of the American College of Cardiology Solution Set Oversight Committee. J Am Coll Cardiol. 2023 May 9;81(18):1835-1878. doi: 10.1016/j.jacc.2023.03.393. Epub 2023 Apr 19. PMID: 37137593.

Kosiborod MN, Abildstrøm SZ, Borlaug BA, Butler J, Rasmussen S, Davies M, Hovingh GK, Kitzman DW, Lindegaard ML, Møller DV, Shah SJ, Treppendahl MB, Verma S, Abhayaratna W, Ahmed FZ, Chopra V, Ezekowitz J, Fu M, Ito H, Lelonek M, Melenovsky V, Merkely B, Núñez J, Perna E, Schou M, Senni M, Sharma K, Van der Meer P, von Lewinski D, Wolf D, Petrie MC; STEP-HFpEF Trial Committees and Investigators. Semaglutide in Patients with Heart Failure with Preserved Ejection Fraction and Obesity. N Engl J Med. 2023 Sep 21;389(12):1069-1084. doi: 10.1056/NEJMoa2306963. Epub 2023 Aug 25. PMID: 37622681.

Kumar KG, Trevaskis JL, Lam DD, Sutton GM, Koza RA, Chouljenko VN, Kousoulas KG, Rogers PM, Kesterson RA, Thearle M, Ferrante AW Jr, Mynatt RL, Burris TP, Dong JZ, Halem HA, Culler MD, Heisler LK, Stephens JM, Butler AA. Identification of adropin as a secreted factor linking dietary macronutrient intake with energy homeostasis and lipid metabolism. Cell Metab. 2008 Dec;8(6):468-81. doi: 10.1016/j.cmet.2008.10.011. PMID: 19041763; PMCID: PMC2746325.

Lagou V, Mägi R, Hottenga JJ, Grallert H, Perry JRB, Bouatia-Naji N, Marullo L, Rybin D, Jansen R, Min JL, Dimas AS, Ulrich A, Zudina L, Gådin JR, Jiang L, Faggian A, Bonnefond A, Fadista J, Stathopoulou MG, Isaacs A, Willems SM, Navarro P, Tanaka T, Jackson AU, Montasser ME, O'Connell JR, Bielak LF, Webster RJ, Saxena R, Stafford JM, Pourcain BS, Timpson NJ, Salo P, Shin SY, Amin N, Smith AV, Li G, Verweij N, Goel A, Ford I, Johnson PCD, Johnson T, Kapur K, Thorleifsson G, Strawbridge RJ, Rasmussen-Torvik LJ, Esko T, Mihailov E, Fall T, Fraser RM,

Mahajan A, Kanoni S, Giedraitis V, Kleber ME, Silbernagel G, Meyer J, Müller-Nurasyid M, Ganna A, Sarin AP, Yengo L, Shungin D, Luan J, Horikoshi M, An P, Sanna S, Boettcher Y, Rayner NW, Nolte IM, Zemunik T, Iperen EV, Kovacs P, Hastie ND, Wild SH, McLachlan S, Campbell S, Polasek O, Carlson O, Egan J, Kiess W, Willemsen G, Kuusisto J, Laakso M, Dimitriou M, Hicks AA, Rauramaa R, Bandinelli S, Thorand B, Liu Y, Miljkovic I, Lind L, Doney A, Perola M, Hingorani A, Kivimaki M, Kumari M, Bennett AJ, Groves CJ, Herder C, Koistinen HA, Kinnunen L, Faire U, Bakker SJL, Uusitupa M, Palmer CNA, Jukema JW, Sattar N, Pouta A, Snieder H, Boerwinkle E, Pankow JS, Magnusson PK, Krus U, Scapoli C, de Geus EJCN, Blüher M, Wolffenbuttel BHR, Province MA, Abecasis GR, Meigs JB, Hovingh GK, Lindström J, Wilson JF, Wright AF, Dedoussis GV, Bornstein SR, Schwarz PEH, Tönjes A, Winkelmann BR, Boehm BO, März W, Metspalu A, Price JF, Deloukas P, Körner A, Lakka TA, Keinanen-Kiukaanniemi SM, Saaristo TE, Bergman RN, Tuomilehto J, Wareham NJ, Langenberg C, Männistö S, Franks PW, Hayward C, Vitart V, Kaprio J, Visvikis-Siest S, Balkau B, Altshuler D, Rudan I, Stumvoll M, Campbell H, van Duijn CM, Gieger C, Illig T, Ferrucci L, Pedersen NL, Pramstaller PP, Boehnke M, Frayling TM, Shuldiner AR, Peyser PA, Kardia SLR, Palmer LJ, Penninx BW, Meneton P, Harris TB, Navis G, Harst PV, Smith GD, Forouhi NG, Loos RJF, Salomaa V, Soranzo N, Boomsma DI, Groop L, Tuomi T, Hofman A, Munroe PB, Gudnason V, Siscovick DS, Watkins H, Lecoeur C, Vollenweider P, Franco-Cereceda A, Eriksson P, Jarvelin MR, Stefansson K, Hamsten A, Nicholson G, Karpe F, Dermitzakis ET, Lindgren CM, McCarthy MI, Froguel P, Kaakinen MA, Lyssenko V, Watanabe RM, Ingelsson E, Florez JC, Dupuis J, Barroso I, Morris AP, Prokopenko I; Meta-Analyses of Glucose and Insulin-related traits Consortium (MAGIC). Sex-dimorphic genetic effects and novel loci for fasting glucose and insulin variability. Nat Commun. 2021 Jan 5;12(1):24. doi: 10.1038/s41467-020-19366-9. Erratum in: Nat Commun. 2021 Feb 8;12(1):995. doi: 10.1038/s41467-021-21276-3. PMID: 33402679; PMCID: PMC7785747.

Liang M, Dickel N, Györfi AH, SafakTümerdem B, Li YN, Rigau AR, Liang C, Hong X, Shen L, Matei AE, Trinh-Minh T, Tran-Manh C, Zhou X, Zehender A, Kreuter A, Zou H, Schett G, Kunz M, Distler

JHW. Attenuation of fibroblast activation and fibrosis by adropin in systemic sclerosis. Sci Transl Med. 2024 Mar 27;16(740):eadd6570. doi: 10.1126/scitranslmed.add6570. Epub 2024 Mar 27. PMID: 38536934.

Lindman BR, Dávila-Román VG, Mann DL, McNulty S, Semigran MJ, Lewis GD, de las Fuentes L, Joseph SM, Vader J, Hernandez AF, Redfield MM. Cardiovascular phenotype in HFpEF patients with or without diabetes: a RELAX trial ancillary study. J Am Coll Cardiol. 2014 Aug 12;64(6):541-9. doi: 10.1016/j.jacc.2014.05.030. PMID: 25104521; PMCID: PMC4133145.

Mushala BAS, Scott I. Adropin: a hepatokine modulator of vascular function and cardiac fuel metabolism. Am J Physiol Heart Circ Physiol. 2021 Jan 1;320(1):H238-H244. doi: 10.1152/ajpheart.00449.2020. Epub 2020 Nov 20. PMID: 33216612; PMCID: PMC7847067.

Prakoso D, Lim SY, Erickson JR, Wallace RS, Lees JG, Tate M, Kiriazis H, Donner DG, Henstridge DC, Davey JR, Qian H, Deo M, Parry LJ, Davidoff AJ, Gregorevic P, Chatham JC, De Blasio MJ, Ritchie RH. Fine-tuning the cardiac O-GlcNAcylation regulatory enzymes governs the functional and structural phenotype of the diabetic heart. Cardiovasc Res. 2022 Jan 7;118(1):212-225. doi: 10.1093/cvr/cvab043. PMID: 33576380.

O'Sullivan JF, Li M, Koay YC, Wang XS, Guglielmi G, Marques FZ, Nanayakkara S, Mariani J, Slaughter E, Kaye DM. Cardiac Substrate Utilization and Relationship to Invasive Exercise Hemodynamic Parameters in HFpEF. JACC Basic Transl Sci. 2024 Jan 31;9(3):281-299. doi: 10.1016/j.jacbts.2023.11.006. PMID: 38559626; PMCID: PMC10978404.

Redfield MM, Borlaug BA. Heart Failure With Preserved Ejection Fraction: A Review. JAMA. 2023 Mar 14;329(10):827-838. doi: 10.1001/jama.2023.2020. PMID: 36917048.

Rottman JN, Ni G, Brown M. Echocardiographic evaluation of ventricular function in mice. Echocardiography. 2007 Jan;24(1):83-9. doi: 10.1111/j.1540-8175.2006.00356.x. PMID: 17214630.

Schiattarella GG, Altamirano F, Tong D, French KM, Villalobos E, Kim SY, Luo X, Jiang N, May HI, Wang ZV, Hill TM, Mammen PPA, Huang J, Lee DI, Hahn VS, Sharma K, Kass DA, Lavandero S, Gillette TG, Hill JA. Nitrosative stress drives heart failure with preserved ejection fraction. Nature. 2019 Apr;568(7752):351-356. doi: 10.1038/s41586-019-1100-z. Epub 2019 Apr 10. PMID: 30971818; PMCID: PMC6635957.

Schnelle M, Catibog N, Zhang M, Nabeebaccus AA, Anderson G, Richards DA, Sawyer G, Zhang X, Toischer K, Hasenfuss G, Monaghan MJ, Shah AM. Echocardiographic evaluation of diastolic function in mouse models of heart disease. J Mol Cell Cardiol. 2018 Jan;114:20-28. doi: 10.1016/j.yjmcc.2017.10.006. Epub 2017 Oct 19. PMID: 29055654; PMCID: PMC5807035.

Shah SJ, Borlaug BA, Kitzman DW, McCulloch AD, Blaxall BC, Agarwal R, Chirinos JA, Collins S, Deo RC, Gladwin MT, Granzier H, Hummel SL, Kass DA, Redfield MM, Sam F, Wang TJ, Desvigne-Nickens P, Adhikari BB. Research Priorities for Heart Failure With Preserved Ejection Fraction: National Heart, Lung, and Blood Institute Working Group Summary. Circulation. 2020 Mar 24;141(12):1001-1026. doi: 10.1161/CIRCULATIONAHA.119.041886. Epub 2020 Mar 23. PMID: 32202936; PMCID: PMC7101072.

Soltani S, Beigrezaei S, Malekahmadi M, Clark CCT, Abdollahi S. Circulating levels of adropin and diabetes: a systematic review and meta-analysis of observational studies. BMC Endocr Disord. 2023 Apr 7;23(1):73. doi: 10.1186/s12902-023-01327-0. PMID: 37029398; PMCID: PMC10080945.

Stokar J, Gurt I, Cohen-Kfir E, Yakubovsky O, Hallak N, Benyamini H, Lishinsky N, Offir N, Tam J, Dresner-Pollak R. Hepatic adropin is regulated by estrogen and contributes to adverse metabolic phenotypes in ovariectomized mice. Mol Metab. 2022 Jun;60:101482. doi: 10.1016/j.molmet.2022.101482. Epub 2022 Mar 29. PMID: 35364299; PMCID: PMC9044006.

Sun Q, Güven B, Wagg CS, Almeida de Oliveira A, Silver H, Zhang L, Chen B, Wei K, Ketema EB, Karwi QG, Persad KL, Vu J, Wang F, Dyck JRB, Oudit GY, Lopaschuk GD. Mitochondrial fatty acid oxidation is the major source of cardiac adenosine triphosphate production in heart failure with preserved ejection fraction. Cardiovasc Res. 2024 Mar 30;120(4):360-371. doi: 10.1093/cvr/cvae006. PMID: 38193548.

Thapa D, Xie B, Manning JR, Zhang M, Stoner MW, Huckestein BR, Edmunds LR, Zhang X, Dedousis NL, O'Doherty RM, Jurczak MJ, Scott I. Adropin reduces blood glucose levels in mice by limiting hepatic glucose production. Physiol Rep. 2019 Apr;7(8):e14043. doi: 10.14814/phy2.14043. PMID: 31004398; PMCID: PMC6474842.

Thapa D, Xie B, Zhang M, Stoner MW, Manning JR, Huckestein BR, Edmunds LR, Mullett SJ, McTiernan CF, Wendell SG, Jurczak MJ, Scott I. Adropin treatment restores cardiac glucose oxidation in pre-diabetic obese mice. J Mol Cell Cardiol. 2019 Apr;129:174-178. doi: 10.1016/j.yjmcc.2019.02.012. Epub 2019 Feb 26. PMID: 30822408; PMCID: PMC6486841.

Thapa D, Zhang M, Manning JR, Guimarães DA, Stoner MW, O'Doherty RM, Shiva S, Scott I. Acetylation of mitochondrial proteins by GCN5L1 promotes enhanced fatty acid oxidation in the heart. Am J Physiol Heart Circ Physiol. 2017 Aug 1;313(2):H265-H274. doi: 10.1152/ajpheart.00752.2016. Epub 2017 May 19. PMID: 28526709; PMCID: PMC5582919.

Tei C, Dujardin KS, Hodge DO, Kyle RA, Tajik AJ, Seward JB. Doppler index combining systolic and diastolic myocardial performance: clinical value in cardiac amyloidosis. J Am Coll Cardiol. 1996 Sep;28(3):658-64. doi: 10.1016/0735-1097(96)00202-1. PMID: 8772753.

Tong D, Schiattarella GG, Jiang N, May HI, Lavandero S, Gillette TG, Hill JA. Female Sex Is Protective in a Preclinical Model of Heart Failure With Preserved Ejection Fraction. Circulation. 2019 Nov 19;140(21):1769-1771. doi: 10.1161/CIRCULATIONAHA.119.042267. Epub 2019 Nov 18. PMID: 31738599; PMCID: PMC6993895.

Tong D, Schiattarella GG, Jiang N, Altamirano F, Szweda PA, Elnwasany A, Lee DI, Yoo H, Kass DA, Szweda LI, Lavandero S, Verdin E, Gillette TG, Hill JA. NAD$^+$ Repletion Reverses Heart Failure With Preserved Ejection Fraction. Circ Res. 2021 May 28;128(11):1629-1641. doi: 10.1161/CIRCRESAHA.120.317046. Epub 2021 Apr 22. PMID: 33882692; PMCID: PMC8159891.

Tran DH, May HI, Li Q, Luo X, Huang J, Zhang G, Niewold E, Wang X, Gillette TG, Deng Y, Wang ZV. Chronic activation of hexosamine biosynthesis in the heart triggers pathological cardiac remodeling. Nat Commun. 2020 Apr 14;11(1):1771. doi: 10.1038/s41467-020-15640-y. PMID: 32286306; PMCID: PMC7156663.

Umapathi P, Mesubi OO, Banerjee PS, Abrol N, Wang Q, Luczak ED, Wu Y, Granger JM, Wei AC, Reyes Gaido OE, Florea L, Talbot CC Jr, Hart GW, Zachara NE, Anderson ME. Excessive *O-GlcNAcylation* Causes Heart Failure and Sudden Death. Circulation. 2021 Apr 27;143(17):1687-1703. doi: 10.1161/CIRCULATIONAHA.120.051911. Epub 2021 Feb 17. Erratum in: Circulation. 2021 Apr 27;143(17):e892. doi: 10.1161/CIR.0000000000000976. PMID: 33593071; PMCID: PMC8085112.

Wong CM, Wang Y, Lee JT, Huang Z, Wu D, Xu A, Lam KS. Adropin is a brain membrane-bound protein regulating physical activity via the NB-3/Notch signaling pathway in mice. J Biol Chem. 2014 Sep 12;289(37):25976-86. doi: 10.1074/jbc.M114.576058. Epub 2014 Jul 29. PMID: 25074942; PMCID: PMC4162195.

Yang C, DeMars KM, Candelario-Jalil E. Age-Dependent Decrease in Adropin is Associated with Reduced Levels of Endothelial Nitric Oxide Synthase and Increased Oxidative Stress in the Rat Brain. Aging Dis. 2018 Apr 1;9(2):322-330. doi: 10.14336/AD.2017.0523. PMID: 29896421; PMCID: PMC5963353.

Ye B, Bradshaw AD, Abrahante JE, Dragon JA, Häußler TN, Bell SP, Hirashima F, LeWinter M, Zile MR, Meyer M. Left Ventricular Gene Expression in Heart Failure With Preserved Ejection Fraction-Profibrotic and Proinflammatory Pathways and Genes. Circ Heart Fail. 2023 Aug;16(8):e010395. doi: 10.1161/CIRCHEARTFAILURE.123.010395. Epub 2023 Jul 27. PMID: 37582166; PMCID: PMC10430768.

ACKNOWLEDGEMENTS

This work was supported by: National Institute of Health Fellowships (F31DK134089 and T32HL110849) to B.A.S.M; National Institute of Health Shared Instrumentation Grants (S10OD023402, S10OD032141) to S.L.G.; National Institutes of Health Research Grants (R01HL124021, R01HL122596, R01HL151228) to S.Y.C.; National Institute of Health Research Grants (R01HL147861, R0HL156874) and American Heart Association Established Investigator Award (23EIA1037834) to I.S. The University of Pittsburgh Center for Metabolism and Mitochondrial Medicine is supported by the Pittsburgh Foundation (MR2020 109502) grant to M.J.J. This project used the University of Pittsburgh Medical Center (UPMC) Hillman Cancer Center and Tissue and Research Pathology/Pitt Biospecimen Core shared resource, which is supported in part by award P30CA047904. Echocardiography was carried out by the University of Pittsburgh Rodent Ultrasonography Core, which received funding from the NIH Shared Instrumentation Grant Program (S10OD023684).

DATA AVAILABILITY

All data reported herein can be found in the results or supplementary information sections of this article.

AUTHOR CONTRIBUTIONS

B.A.S.M., M.J.J., and I.S. conceived the study and designed experiments. E.S.G., Y.A.A. and S.Y.C. provided essential biological samples. B.A.S.M, M.W.S., B.M., A.M.V., and S.J.M. performed experiments. B.A.S.M., B.M., S.J.M. and I.S. analyzed data. B.A.S.M. and I.S. prepared figures. J.R.M., P.B., N.B., M.S.S., S.S.S., B.A.K., C.Z., E.S.G, S.Y.C., and S.L.G provided critical input and expertise. B.A.S.M and I.S. drafted the manuscript. B.A.S.M. and I.S. edited and revised the manuscript. All authors approved the final submission.

DISCLOSURES

Figure 1: Recombinant Adropin treatment improves glucose tolerance in HFpEF without changes in body weight. A. Total circulating Adropin levels in age- and gender-matched control and

HFpEF patients. N = 19-20. **B.** Timeline and treatment protocol used in mouse HFpEF studies. **C,D.** Daily and final body weights of mice. N = 8-12. **E,F.** IPGTT time-course and area under-the-curve (AUC). N = 8-12.

Figure 2: Recombinant Adropin treatment limits cardiac diastolic dysfunction in a mouse model of HFpEF. A. Representative echocardiography (M-mode and Doppler) images. **B-J.** Echocardiography measurements of heart rate, cardiac output, calculated left ventricle (LV) mass, ejection fraction, E/e' ratio, E/A ratio, mitral annulus tissue velocity during early diastole (e'), isovolumetric relaxation time (IVRT), and myocardial performance index/Tei index) (MPI/Tei). N = 8-12.

Figure 3: Long-term Adropin treatment preventes cardiomyocyte hypertrophy and reduces cardiac fibrosis. A. Representative wheat germ agglutinin (WGA) and Masson's trichrome images. **B.** Cardiomyocyte cross-sectional area. **C.** Cardiac interstitial fibrosis. **D.** Bulk RNA-seq pathway analysis of mouse hearts using ShinyGO in HFpEF vs. HFpEF+Adr. mice. N = 4. **E.** Fibrosis/ECM gene expression (fold change) in HFpEF vs. HFpEF+Adr. mice. N = 4.

Figure 4: Adropin treatment limits cardiac protein O-GlcNAcylation in HFpEF mice. A. BioCyc

pathway analysis of untargeted metabolites in HFpEF vs. HFpEF+Adr. hearts. HBP = hexosamine

biosynthesis pathway. N = 8. **B.** Schematic of hexosamine biosynthesis pathway (HBP) for the generation of the O-GlcNAcylation substrate, UDP-GlcNAc. **C-F.** Quantification of HBP metabolites GlcNAc-6-P, CoA, Uridine, and Cytidine in untreated and Adropin-treated HFpEF mice. N = 8. **G.** Representative immunoblots of total O-GlcNAcylation and HBP/O-GlcNAcylation regulatory proteins. **H-K.** Quantification of immunoblots. N = 8-12.

Figure 5: Adropin-mediated reductions in LCAD O-GlcNAcylation improve its enzymatic activity. A. Enzymatic function of LCAD. **B.** Immunoprecipitation analysis of LCAD O-GlcNAcylation status, N = 4. **C.** LCAD enzymatic activity from bulk cardiac tissue, N = 4. **D.** Linear regression analysis of the relationship between LCAD O-GlcNAcylation and LCAD activity. **E.** *In vitro* enzymatic activity of recombinant LCAD after exposure to O-GlcNAcylation substrate (UDP-GlcNAc) and/or recombinant O-GlcNAc transferase (OGT), N = 3.