1 **An integrative framework linking molecular signatures and locomotory**
2 **phenotypes in space-induced sarcopenia**
3

Authors and Affiliations: Brendan K. Ball[1+*], Hammad F. Khan[1+*]

1. Weldon School of Biomedical Engineering, Purdue University, West Lafayette, IN

47907, USA

[+]These authors contributed equally to this work

*Correspondence: bbkazu@purdue.edu, khan332@purdue.edu

Keywords: Spaceflight, sarcopenia, transcriptomics, locomotion, translational

modeling, systems biology

ABSTRACT

Age-related skeletal muscle deterioration, referred to as sarcopenia, poses significant risks to astronaut health and mission success during spaceflight, yet its multisystem drivers remain poorly understood. While terrestrial sarcopenia manifests gradually through aging, spaceflight induces analogous musculoskeletal decline within weeks, providing an accelerated model to study conserved atrophy mechanisms. Here, we introduced an integrative framework combining cross-species genetic analysis with physiological modeling to understand mechanistic pathways in space-induced sarcopenia. By analyzing rodent and human datasets, we identified conserved molecular pathways underlying microgravity-induced muscle atrophy, revealing shared regulators of neuromuscular signaling including pathways related to neurotransmitter release and regulation, mitochondrial function, and synaptic integration. Building upon these molecular insights, we developed a physiologically grounded central pattern generator model that reproduced spaceflight-induced locomotion deficits in mice. This multi-scale approach established mechanistic connections between transcriptional changes and impaired movement kinetics while identifying potential therapeutic targets applicable to both spaceflight and terrestrial aging-related muscle loss.

INTRODUCTION

Space exploration is rapidly developing with the increasing number of missions to the International Space Station and beyond. As humans push towards longer spaceflights and journeys to new extraterrestrial planets, understanding the physiological changes due to space travel has become critical. While such concerns are actively studied by researchers, our understanding of the biological burden of spaceflight is largely limited. Moreover, this limitation is further exacerbated by the effects of prolonged exposure to the microgravity environment[1] and cosmic radiation,[2,3] which has been proven to be detrimental to human health.[4]

A well-documented consequence of space exploration is the deterioration of musculoskeletal health. Astronauts who return from space experience muscle atrophy, impaired muscle performance, and loss of strength.[5,6] This process of muscle degradation resembles sarcopenia (SA), a progressive age-related condition that often emerges over several decades and leads to muscle loss and function.[7] While this condition progresses over an individual's lifetime, the development of SA-like pathology is accelerated under microgravity conditions.[8] Despite noted observations of muscle and strength breakdown,[9] the exact mechanisms and pathways by which space-induced SA progresses are poorly understood.

There are major challenges to understanding how conditions in space affect human health. First, there are limited experimental samples and astronauts that have been studied on the International Space Station, often due to space constraints, costs, and

accessibility.[10] This is important because studying physiological changes in space-like

conditions on Earth does not effectively translate to the harsh conditions in microgravity.

Additionally, astronauts on average spend on average 6 months per space mission tour,

and thus there is limited information on the longer-term effects of space travel and

sample size to study such effects.[11] To address this challenge, previous studies aimed

to improve our understanding of spaceflight and SA through the use of *in vitro*[12] and *in*

vivo rodent models.[13] Despite these efforts, these approaches did not account for the

heterogeneity of the human population and the biological differences across mammalian

species. As a result, our understanding of a mechanistic pathway to SA conditions is

largely unknown.

To circumvent this limitation, we leveraged a framework known as Translatable

Components Regression (TransComp-R), a computational model that synthesizes

multi-omics data from preclinical rodent models and human data to predict disease

outcomes in humans.[14,15] The TransComp-R pipeline has been successfully applied to

overcome other biomedical challenges including Alzheimer's disease,[16–20]

osteoarthritis,[1] and inflammatory bowel disease.[14,21] We couple this approach with a

biophysically grounded model of central pattern generator (CPG) networks,[22,23]

therefore enabling multiscale interrogation of SA.

In this study, we show that transcriptomics mouse models under spaceflight conditions

could predict human outcomes of SA. We performed cross-species computational

modeling to synthesize spaceflight mouse and SA human data to identify potential

biological pathways that are involved in muscle degeneration. To link the observed

space-related SA, we built a network of leaky integrate-and-fire neurons to build a

computational CPG model that recapitulated locomotion kinetics in healthy conditions.

Importantly, incorporation of the predictive transcriptomic data into the model generated

similar phenotypic gait responses observed under SA conditions. Thus, this approach

enables an improved understanding of mechanisms associated with space-induced SA

and may potentially inform therapeutic avenues for SA and other muscle-deteriorating

conditions related to spaceflight and on Earth.

RESULTS

Spaceflight mouse principal components separate SA and control conditions in

humans

We accessed RNA-sequencing spaceflight mouse data from the NASA Open Science

Data Repository (OSDR). The publicly available spaceflight dataset (OSD-103)

contained 16-week-old C57BL/6J male mice with a sample size of 6 ground controls

and 6 spaceflight conditions.[24,25] The spaceflight group was housed at the International

Space Station for 37 days before euthanasia and collection of quadricep muscle

samples. The ground control group was housed on Earth for reference. We then

acquired RNA-sequencing data of patients with SA from Gene Expression Omnibus

(GEO). We selected three separate cohort studies that studied SA among different

racial demographics, including Caucasian (GSE111006), Afro-Caribbean (GSE111010),

and Chinese (GSE110016) descent.[26,27] Our integration of multiple cohorts allowed us

to reference our findings across different human backgrounds.

We applied the TransComp-R model to identify PCs that encoded transcriptomics

variation of spaceflight mice that predicted SA outcomes in humans. The TransComp-R

pipeline works by computing a mouse principal component analysis (PCA) space and

projecting human data into space. Mouse principal components that encode

transcriptomics capable of stratifying human disease from control groups by a linear

regression can be biologically interpreted through pathway enrichment analyses and

follow-up studies.

With the TransComp-R model, we first matched homologous gene pairs across the

mouse and all three human datasets (**Figure 1a**). We next applied principal component

analysis (PCA) on the mouse dataset to identify principal components (PCs) that

explained 80% of the cumulative variation and projected the human dataset with the

Chinese cohort with SA and control conditions into the mouse PC space.

Figure 1. Mouse to human modeling workflow. (a) Homologous pairs between mouse and human are first matched. Human SA and control groups (Chinese cohort) are projected into the space flight mouse PCs. The human Caucasian and Afro-Caribbean cohorts were used for downstream validation analysis. **(b)** The PCs selected by LASSO are regressed against human SA and control outcomes to determine significance. **(c)** Significant PCs from the generalized linear model are used for downstream biological analysis. (**d**) Biophysical modeling of motor dynamics. (**e**) Recapitulate behavior deficits in space-induced sarcopenia based on genetic determinants.

After synthesizing the mouse and human transcriptomics data together, we selected PCs that were relevant in predicting human SA condition with LASSO (least absolute shrinkage and selection operator) (**Figure 1b**). The selected PCs from LASSO were used in a generalized linear regression against human SA and control groups. The significant PCs were then interpreted using gene set enrichment analysis (GSEA).

Lastly, we validated our findings with the Caucasian and Afro-Caribbean cohorts that

also contained SA and control data.

After matching for homologous pairs, we identified 12,484 genes shared across the

mouse and human datasets. We constructed a mouse PCA space and projected the

Chinese cohort and identified 7 PCs that explained 80% of the variance (83.4% var.

exp). We next calculated how much variance the mouse PCs explained in the human

data to understand the potential of translation from mouse to human (**Figure 2a**).

Among the seven PCs, we found PC1, PC3, and PC6 explained a higher ratio of human

variance compared to mouse variance. This finding may suggest that PC1, PC3, and

PC6 may have a more pronounced ability to capture variance across species than other

PCs.

Figure 2. Significant mouse PCs distinguish SA and control conditions in humans. (a) The percent variance explained of mouse PCs and how much they explain in human data. **(b)** PC selection using the LASSO. PCs selected by LASSO that were significant in a GLM (p value < 0.05) are denoted with a star. **(c)** Plots of significant mouse PCs separating SA and control in humans.

Of the seven PCs, we identified PC2, PC3, PC4, and PC7 with our LASSO model to be relevant eigenvectors discerning human SA and control patients (**Figure 2b**). We created a generalized linear regression model incorporating these four PCs and found PC3, PC4, and PC6 to be significantly separating the SA and control groups (p = 0.0286, 0.0371, and 0.0253, respectively). We next visualized the three significant PCs and found that the transcriptomics variation encoded on each PC separated the human SA and control groups (**Figure 2c**).

Genes contributing to metabolic and immune-associated pathways are distinguished across patients with sarcopenia

Having identified significant PCs that could distinguish between human SA conditions, we sought to interpret the biological implications using GSEA. Our enrichment analysis showed genes contributing to pathways related to neuroactive ligand receptor interactions, immune pathways, and cellular structures were enriched for SA on mouse PC3 (**Figure 3a**). Genes that were involved with neurodegenerative diseases and metabolic pathways were associated with healthy groups. This indicates that these genes are regulated under healthy conditions and may become disrupted under SA conditions.

Figure 3. KEGG pathway enrichment analysis. Biological pathways enriched on **(a)** PC3, **(b)** PC4, and **(c)** PC6. All pathways are significant with a Benjamini-Hochberg adjusted p<0.01. The gene size represents the number of genes enriched in the biological pathway. Pathways are shown with associations to human sarcopenia (red) and control (blue) based on the directionality of separation for each PC. The bar plots are represented by their respective normalized enrichment score (NES).

On the mouse spaceflight PC4, we found pathways involved with immune signaling and cellular structure to be enriched in the SA group (**Figure 3b**). Furthermore, we found neurodegenerative diseases to be involved with control groups, similar to those found in PC3. Among the pathways enriched on PC6, we found neuroactive ligand receptor interaction and PPAR signaling associated with SA, whereas regulation of actin cytoskeleton, adherens junction, and leukocyte transendothelial migration was enriched in control (**Figure 3c**).

We also performed GSEA using the hallmark curation and found signaling mechanisms

involved with the immune system to be involved with SA (**Supplementary Figure 1**).

Additionally, we found oxidative phosphorylation to be consistently enriched for the

control group in PC3, PC4, and PC6. Such results further support our findings with the

reported KEGG pathway analysis. These findings also show that genes contributing to

the immune responses, as well as mitochondrial function associated with oxidative

phosphorylation may play important roles in the muscle degenerative process

associated with SA in humans.

Significant PC gene loadings from the Chinese cohort are translatable to

Caucasian and Afro-Caribbean demographics

After identifying biological pathways associated with SA and control, we questioned if

our findings were relevant in other human demographics since our TransComp-R model

synthesized a human Chinese cohort with spaceflight gene expression data. Here, we

combined the Caucasian and Afro-Caribbean SA datasets to determine if findings from

these populations were also reflected by our findings from the Chinese cohort.

Different from the Chinese cohort, the Caucasian and Afro-Caribbean datasets

contained further annotations on muscle weakness in the subjects. We first excluded

any patients from the control groups who had any reported history of muscle weakness

or muscle loss. We then combined the dataset and verified that batch effects were not

present with PCA (**Supplementary Figure 2**). We next used the combined

Caucasian/Afro-Caribbean data for a partial least squares discriminant analysis (PLS-

DA) model (n=32 Caucasian, n=23 Afro-Caribbean). Breaking down the combined and

processed demographics, we analyzed 28 healthy Caucasians (Age: 72.6±2.3 standard

deviations), 4 SA Caucasians (Age: 74.3±3.6), 14 Afro-Caribbean (76.4±7.4), and 9 SA

Afro-Caribbean (76.8±7.9) individuals.

We constructed three PLS-DA models for each significant mouse PC that we identified

to be significant from our TranComp-R model. We selected a panel of genes driving the

separation among the PCs by selecting the top 10 and bottom 10 genes ranked by their

scores. We then performed PLS-DA and determined what genes were strongly

contributing to the separation. Genes strongly contributing to the model contained a

calculated variable importance of projection (VIP) score greater than 1.

We demonstrated a separation of control and SA with the combined human dataset and

identified genes that contributed to the separation above average (**Figure 4a**). Within

PC3, the human samples were separated among the first latent variable (LV). On this

LV, we found genes that are strongly associated with SA, which included *DLG3*,

CALB2, *LHFPL6*, and *ADAM33* (**Figure 4b**). The *DLG3* and *CALB2* genes are

associated with neuron signaling, whereas *LHFPL6* and *ADAM33* are involved with cell-

to-cell signaling and communication. Oppositely, genes such as *PPP1R12B* and *CNTF*

are associated with the control groups. These genes are involved in helping muscle

contraction through myosin phosphatase activity and enhancing survival factors for

neuronal cells, respectively.

Figure 4. Gene expression driving separation among Chinese SA cohorts is also reflected in Caucasian and Afro-Caribbean populations. (a) PLS-DA on the top 10 and bottom 10 genes identified in the spaceflight PC3. **(b)** The loadings of the first two latent variables for PC3. **(c)** PLS-DA for PC4. **(d)** Loadings for PC3. **(e)** PLS-DA for PC6. **(f)** Loadings for PC6. Loadings with a VIP>1 are labeled with a star. The color on the loadings represents the highest contribution to SA or control by the specific gene.

We also performed PLS-DA on mouse PC4 and demonstrated that LV1 discriminated between the control and SA human groups (**Figure 4c**). We identified *PKP2*, *SARDH*, *FMO4*, *CEP85L*, and *TIMP4* genes to associate with SA, whereas *PLIN3* and *TVP23A* were associated with control (**Figure 4d**). These genes have known associations with

cellular structure, metabolism, and vesicle transport. Similar to the first two mouse PCs,

we also found the control group to cluster on the negative score of LV1, and the SA

group to cluster among positive scores (**Figure 4e**). This separation was defined by

CNR1, *GRIN2B*, *LYRM4*, *MOCOS*, *WDFY2*, and *NME5* for SA. For the control groups,

we found *LRFN5* as the only gene contributing to the PLS-DA separation with a VIP>1.

Interestingly, the *CNR1* and *GRIN2B* genes are linked to neurotransmission and brain

function whereas *LYRM4* and *MOCOS* are connected to mitochondrial metabolism.[28–30]

The *LRFN5* gene we found associated with the control group is associated with cell-to-

cell communication among neuronal cells.

Multivariate analysis reveals rodent gait deficits related to orbital spaceflight

Space-induced muscle atrophy represents a compelling model for investigating

fundamental mechanisms of sarcopenia. We demonstrated with pathway enrichment

analysis that neuroactive ligand receptor interaction and PPAR signaling were

associated with SA in both human and rodent models, suggesting conserved genetic

determinants for SA.[31] To further augment the relationship between SA changes at the

molecular level, we further presented a multivariate analysis of locomotory deficits in

rodents following prolonged microgravity exposure aboard the International Space

Station.

We accessed the spaceflight behavioral assay available on the NASA database (OSD-

478) which contains a prospective collection of gait data between mice undergoing

spaceflight in orbit aboard the ISS and ground control over 35 days.[32] Pre-flight and

post-flight gait measurements were conducted across a total of 40 mice (20 mice per

group), measuring multiple aspects of locomotion and gait kinematics. We conducted an

unsupervised analysis of effects on gait dynamics across spaceflight and ground control

by projecting the data into a mouse PC subspace to extract behavior dynamics that

were most affected during spaceflight.

We used PCA of gait dynamics to reveal distinct behavioral signatures between pre-

and post-flight conditions (**Figure 5a, b**). We clustered the treatment groups based on

chi-squared confidence ellipse and noted a distinct separation in PC space that was

more pronounced in spaceflight groups compared to ground controls, suggesting

specific microgravity-induced adaptations across mice as opposed to temporal artifacts.

The first PC captured 25.5% of total variance followed by 14.8% and 11.3% for PC 2

and 3, respectively. Approximately 80% of locomotory variation is explained by just

eight components (**Figure 5c**), suggesting strongly shared effects across all mice

undergoing spaceflight.

To investigate behavioral contributions, we conducted a loading analysis of tracked gait

properties during locomotion and identified key gait parameters contribution to PC

variance in our data (**Figure 5d**). For example, among the first two components, we

identified propulsion parameters and stride as primary contributors to observed variance

(**Figure 5d**). We found similar results when conducting systematic testing across all gait

parameters across mice (**Supplementary Figure 3**). To further quantify this, we

selected the most contributing gait parameters across the first 4 PC components and

performed inferential statistics across mice. We found significant alterations in propel

time (p=0.04), stance time (p=0.007), stride length (p<0.001), and swing time (p<0.001)

following orbital habitation (**Figure 5e**). These deficits suggest impairments in both

stride generation and balance feedback mechanisms essential for coordinated

movement.[33] Similar observations were noted in humans undergoing functional gait

tests.[34] Moreover, this analysis demonstrated clear locomotory deficits in mice following

spaceflight exposure. Notably, the propelling force and stride kinematics were strongly

altered compared to ground control. These coordinated changes could suggest

disruption in multiple neural pathways governing both balance maintenance and gait

kinetics, consistent with functional adaptations to microgravity conditions.[34,35]

Figure 5. Behavioral alterations during spaceflight. (a) PC projection of the first and second component axis with behavioral variables under pre- and post-flight conditions with ground controls. **(b)** PC projection of the first and third PC components. **(c)** Scree plot of variance explained for the first 10 PC components. **(d)** A bi-polar plot of the top 4 variable behaviors among the flight and ground control groups. **(e)** Highest weighted behavioral alteration properties determined by PC analysis (n = 40 mice total, One-way ANOVA test).

Physiological modeling of gait characteristics

To understand the physiological mechanisms underlying space-induced locomotory deficits as well as build a correlation between key molecular determinants of SA, we first established a biologically plausible physiological model that captured the essential dynamics of spinal locomotor circuits under normal terrestrial conditions. This computational framework serves as a substrate to later assess microgravity-induced alterations, demonstrating how changes in network activity directly translate to muscle function and ultimately to observable gait parameters.

Our network model consisted of leaky integrate-and-fire neurons following well-established membrane potential and gating dynamics.[22,23,36–38] Importantly, to capture oscillatory dynamics often found during locomotion, we assigned distinct flexor and extensor neural populations that were inhibitory connected to one another. This enabled a subset of neurons to generate patterned activity in the absence of input to both regions thus reflecting CPG dynamics (**Figure 6a**).[36] Simulation of the population model drove characteristic oscillatory patterns in membrane potential dynamics between flexor and extensor circuits under normal gravity conditions (**Figure 6b**), with flexors exhibiting more rapid depolarization consistent with their role in initiating limb movement. The computational network successfully reproduced rhythmic activity reflecting central pattern generators, with flexor neurons (red) showing synchronized burst patterns distinct from extensor neurons (blue) during simulated normal locomotion (**Figure 6c**). We used the cumulative output of each group to infer muscle activation patterns which were used to extract gait dynamics (**Figure 6d**). Gait kinematics exhibited a strong

phase-coherent response (67%) between extensor and flexor periods (**Figure 6e**)

reflecting coordination between muscle groups.

Thus, this computational approach creates a powerful platform connecting three levels

of analysis: neuronal firing patterns, muscle activation dynamics, and resultant gait

kinetics (**Figure 6f-h**). The normal phase relationships, stride distributions, and swing

patterns established here provide quantifiable baselines against which space-induced

neural adaptations can be measured in subsequent experiments. By bridging cellular

neurophysiology to whole-organism locomotion, this model enables mechanistic

investigations into how microgravity disrupts motor control at multiple biological levels.

Figure 6. Motor neuron population model for locomotory behavior. (a) Schematic of the mechanistic physiological model. **(b)** Averaged population membrane dynamics from a leaky integrate-and-fire model. **(c)** Rastergram of 50 neurons spiking during constant input. Neurons are colored based on assigned flexor or extensor dynamics. **(d)** Inferred muscle activity patterns as a function of population output. **(e)** Polar plot of phase relationship between flexor and extensor population dynamics. **(f)** Distribution of stride length with coefficient of variation. **(g)** Distribution of swing times with coefficient of variation. **(h)** Regression of correlation swing time and stride length.

Computational framework linking genetic determinants of sarcopenia to altered gait characteristics

To bridge the gap between identified genetic determinants found by the gene enrichment analysis and the physiology of gait characteristics, we modified our network model to exhibit SA phenotypic behavior. This was primarily informed by the identified ligand-gated genetic disruption (G*RIN2B* and *CNR1*), during spaceflight in both mice and human cohorts (**Figure 7a**). We modified the synaptic transmission probability and conductances while biasing the total input variability into the network of individual neurons in both flexor and extensor populations. We noted that, upon this disruption, an observed reduction in phase-coherence between flexor and extensor dynamics compared to the healthy terrestrial network. (Coherence$_{Healthy}$ = 55%, Coherence$_{SA}$ = 33%, **Figure 7b**). The change in the network activity also drove a significant increase in both inferred gait swing time and stride length ($p < 0.0001$, **Figure 7c, d**), most likely due to increase variability in the CPG network. Thus, these results recapitulate the same gait alterations observed in cohorts of mice that spend a significant time in space flight (**Figure 5e**).

Figure 7. The physiological model illustrates genetic determinants for space-related sarcopenia. (**a**) Schematic showing a physiological model of the motor neuron to muscle activation alongside SA-related gene regulation from normalized enrichment score (NES). (**b**) Healthy and SA population-related phase activity across neurons during inferred muscle activity. Note reduced phase strength during movement. Reported phase coherence for healthy and SA activity is $R_{healthy} = 0.55$ and $R_{sarcopenia} = 0.33$, respectively. (**c**) Modeled swing times of population response under healthy and SA ($p = 0.00$, $t(1423) = -19.13$, Student's T-test). (**d**) Modeled swing times of population response under healthy and SA for calculated stride length ($p = 0.00$, $t(2281) = -7.16$, Student's T-test).

This computational framework provided mechanistic insights connecting space-induced alterations in neural circuitry to the observed locomotor deficits. Our findings suggest that space-induced sarcopenia involves not only peripheral muscle atrophy but also fundamental adaptations in neural control systems, including altered central pattern generator function and disrupted coordination between antagonistic muscle groups. These neural adaptations likely contribute significantly to the gait abnormalities

documented in our behavioral analysis and highlight potential targets for
countermeasure development.

DISCUSSION

Space-induced SA is a major physiological change that affects astronauts. We used

transcriptomics muscle biopsies of mice exposed to microgravity and human

populations with human SA data to understand the effects of space travel on humans.

In this work, we presented a comprehensive framework to understand the biological

pathway and its effect on locomotion phenotypes in space-induced SA. By leveraging

TransComp-R, we used a mouse-to-human modeling approach to identify key biological

pathways related to neurotransmitter release and regulation, synaptic integration, and

mitochondrial dynamics associated with SA conditions. Likewise, a comprehensive

analysis of locomotion behavior in mice exposed to spaceflight showed strong alteration

in gait parameters which we mechanistically linked to a CPG network model

incorporating altered identified molecular pathways associated with SA conditions. Our

results established a framework to link genetic determinants of SA to phenotypically

resulting locomotion behavior.

We identified several immune system pathways altered under conditions of SA, which

has also been observed under spaceflight conditions.[39–41] A study reported immune

dysregulation and the redevelopment of herpes virus among astronauts on a 6-month

orbital space mission.[1] A separate 6-month spaceflight that studied two crew members

identified elevated proinflammatory biomarkers in muscle biopsy post-landing.[42] During

spaceflight, blood-derived cytokines including interleukins and tumor necrosis factor-

alpha were also significantly elevated during flight.[43] Further investigation of such

immune pathways may improve our understanding of spaceflight-induced SA.

Metabolic pathways are actively involved in muscle health as demonstrated by protein

alterations during spaceflight.[44] Stein and colleagues reported that whole-body protein

synthesis was reduced by 45% about preflight measurements.[45] In the same study,

post-flight measurements returned to 91% of the pre-flight levels, indicating a lack of

complete recovery from space travel.[45] Additionally, we identified genes that were

strongly associated with SA through neuromuscular signaling. Neuromuscular junction

aging has been proposed to be involved in sarcopenia pathogenesis.[46] The age-related

denervation at the neuromuscular junction occurs before muscle atrophy[47] and may be

involved in SA pathogenesis.[48]

The biomechanical alterations we noted in our study from the behavioral assay align

with previously documented changes in movement atrophy.[49,50] The propulsion and

stride deficits likely reflect preferential atrophy of fast-twitch extensor muscles that

generate locomotive force, mirroring the fiber-type selectivity observed in terrestrial

sarcopenia but compressed into a remarkably accelerated timeframe (~35 days).[35,51]

The concurrent changes in stride parameters suggest simultaneous adaptation of

central pattern generators and sensorimotor integration circuits. It is of interest that

several key protein pathways were also affected in both humans and mice which could

explain some of the downstream locomotion phenotypes. For example, the presence of

DLG3 and *GRIN2B* suggests alterations of synaptic integration across neurons within a

CPG network. Interestingly, *GRIN2B* was upregulated in the skeletal muscle of aging

mice.[52] Such consequences would be reflected in how neural populations rhythmically

generated activity between flexor and extensor periods. Indeed, when altering synaptic

integration properties along the network in our CPG model, primarily synaptic release

probability and input gain, we noted a clear reduction in firing coherence between flexor

and extensor population, which recapitulates the behavior changes we observed in the

gait dynamics.

Although our study identified several pathways to improve our understanding of space-

induced SA, there are limitations to this work. First, due to limited publicly available data

on mice that have been exposed to spaceflight, the sample sizes are smaller.

Additionally, our mouse and human datasets only contained information from males,

which neglects potential biological outcomes in females. Therefore, sex-based findings

from this work should be cautioned. Additionally, we only analyzed physiological

changes in the quadricep muscle and did not investigate the contribution of other limb

muscles. Future studies that incorporate additional demographic and sex-based

information of both males and females, with the inclusion of muscle biopsies and

relevant tissue will largely build upon the findings we reported.

Likewise, there are limitations to our computational models. One such limitation is in the

methodology of our TransComp-R model. The TransComp-R computational model

selects homologous gene pairs across mouse and human datasets; thus, genes that do

not fit this criterion are omitted. As a result, there is a possibility that these omitted

genes may have also served an active role in the physiological changes to spaceflight-

induced SA. Our CPG network model incorporated leaky integrate-and-fire neurons with

reciprocal connections, synaptic probabilities, and membrane dynamics. However,

additional models can be incorporated to further explore the mechanistic contribution of

protein altered during SA. For example, modeling a biophysically realistic model of

neurons with full ion channel distributions and gating dynamics could be more

informative on how changes at the protein level affect neural dynamics and CPG

activity. For example, one can investigate how excessive synaptic release alters

integration properties of neurons and their firing rate. Further experiments are needed to

causally link the synaptic and protein interactions to behavioral deficits observed during

spaceflight or using alternative models of age-related SA.

Our study investigated the effects of SA as a consequence of spaceflight travel. We

used the TransComp-R approach and identified spaceflight mouse PCs that could

predict changes in human age-associated SA in muscle biopsies. Our findings were

then validated by matching a panel of genetic markers from these PCs in other human

cohorts across different populations. Lastly, our CPG neuron model incorporated with

changes in genetic markers of SA was able to recapitulate healthy and space-induced

SA phenotypes in locomotion. Thus, our findings in this work pave a path to link

molecular pathways to behavioral phenotypes of spaced-induced SA.

MATERIALS AND METHODS

Data selection

Data selection

We selected publicly available transcriptomics data from the NASA OSDR and GEO

database with the prerequisite that the data contained a balanced number of samples

with controls and processed tissue samples from similar quadriceps muscle regions in

both mouse and human.

Our mouse dataset was selected from the NASA OSDR. We focused our selection

criteria on transcriptomics data that contained muscle samples near the quadriceps. We

identified a mouse study (OSD-103) that contained a space flight and ground control

group. We next searched the GEO database and identified three human SA datasets

with controls and biopsy samples, each containing transcriptomics data of biopsy

samples from the vastus lateralis muscle. The datasets represent a demographic

population around the world, including Caucasian (GSE111006), Chinese

(GSE111016), and Afro-Caribbean (GSE111010).

Pre-processing and data normalization

We downloaded human transcriptomics SA data from the GEO database using the

GEO2R platform with Bioconductor tools in R (*GEOquery* ver. 2.70.0, *Biobase* ver.

2.62.0, and *limma* ver. 3.58.1). We log_2 transformed the data and z-score normalized

the data per gene. We matched the human and mouse datasets for homologous one-to-

one gene pairing using the *orthogene* package (ver. 1.8.0). Genes that did not contain

homologous pairing were excluded from the model and further downstream analysis.

Mouse to human computational modeling

We performed TransComp-R by performing PCA on the mouse space flight with both flight and ground control groups. We selected mouse PCs that explained up to a total of 80% cumulative variance to prevent overfitting of the data and inclusion of noise. We projected the human dataset (Chinese cohort GSE111016) into the mouse PC space. The projection of the human data into the mouse PCA space is a matrix multiplication of $X^{s \times n}$ and $Q^{n \times p}$ such that s is the row of human patients, n is the homologous gene pairs shared across both mouse and human, and p is the number of PCs that explain 80% cumulative variance. The equation is also shown:

$$\square^{\square\square\square} = \square^{\square\square\square}\square^{\square\square\square} \tag{1}$$

Feature selection of mouse principal components

To select important features and regularize the mouse PCs that best predicted human SA condition, we performed LASSO across 20 rounds of 5-fold cross-validations by regressing the human SA and control group positions along the mouse PCA space against the human disease condition. Mouse PCs with a non-zero coefficient frequency greater than 10 of 20 rounds were selected for the generalized linear model. We performed the generalized linear regression using the equation:

$$Y \sim \beta_0 + \sum_{i=1}^{k} \beta_i (Xq_i) \tag{2}$$

In the generalized linear model, the Xq_i represents the projected human samples into mouse PCs that were selected by LASSO and Y is the human SA and control status. We selected mouse PCs with a p value less than 0.05 for downstream biological interpretation.

Variance explained in human by mouse principal components

Using principal components encoding mouse variation and human data, we can

calculate how much variation the mouse PCs explain the variation in humans. We

calculated the variance explained by mouse PCs in human data with the equation:

$$\text{Variance Explained}(q_i) = \frac{q_i^T [X^T X] q_i}{tr(X^T X)}$$

(**3**)

Where $X^{s \times n}$ is a human data matrix represented by s number of subjects and n number

of genes, and $Q^{n \times p}$ is the mouse data represented by the loading coefficients of n

number genes and p columns of principal components q_i. The T represents the

transpose of the data matrix.

Gene set enrichment analysis

We selected PCs that significantly distinguished between human SA and control groups

for downstream pathway analysis. To identify enriched pathways, we pre-ranked each

significant PC by their respective loadings score and performed GSEA in R (*msigdbr*

ver. 7.5.1, *fgsea* ver. 1.28.0, *clusterProfiler* ver. 4.10.1).[53–55] We generated a holistic

understanding of the enriched pathways by using the hallmark and KEGG curations

from the Molecular Signatures Database. The parameters we used to identify enriched

biological pathways included a minimum gene size of 5, a maximum gene size of 500,

and an epsilon value of 0. We used the default parameter of 1000 permutations to

determine the statistical significance. We defined biological pathways in both KEGG and

hallmark to be significant if the Benjamini-Hochberg adjusted p value was less than

0.01.

Gene selection for cross-population validation

To ensure our findings are reproducible in other datasets, we used the Caucasian and

Afro-Caribbean datasets (GSE111006 and GSE111010) to determine if the findings in

the Chinese cohort are reflective in other demographics. In both the Caucasian and

Afro-Caribbean datasets, we first excluded control patients with a reported history of low

muscle mass or low muscle strength and/or performance to ensure the control groups

had no potential complications. We then concomitantly merged the two cohorts into a

singular dataset and verified for batch effects visualized by PCA.

Using the Chinese cohort, we filtered for the top 10 and bottom 10 gene loadings from

significant mouse PCs identified in our TransComp-R model to determine if

transcriptomic variation encoded on these PCs could stratify SA and healthy groups in

the Caucasian and Afro-Caribbean populations.

Partial least squares discriminant analysis

We used the *mixOmics* package (ver. 6.26.0) to construct the PLS-DA models for each

significant mouse PC that could significantly stratify human SA and control groups in the

Chinese SA cohort.[56]

We identified the most important genes contributing to the model's predictive ability by

calculating the VIP scores. We used the *mixOmics* package to calculate the VIP scores

for each given LV. The following equation:

$$VIP_k = \sqrt{\frac{K \sum_{a=1}^{A} w_{ak}^2 SSA_a}{A \cdot SSA_{total}}}$$ **(4)**

A given VIP score for a specific gene is calculated by *K* number of genes, *A* number of

PLS-DA components, and w_{ak} weight of the gene *k* in a^{th} LV. SSA_a and SSA_{total} are the

sum of squares in the a^{th} LV and the total sum of squares in all LV components,

respectively. We defined a VIP score greater than 1 to represent genes contributing

above average to the model's performance.

Multivariate analysis of locomotor behavior

We used the OSD-478 from the NASA OSDR for locomotor behavior analysis. Principal

component analysis was applied to 47 locomotor behavioral parameters measured in 35

C57BL/6J mice across four experimental groups: pre-flight spaceflight (pre-flight), post-

flight spaceflight (post-flight), pre-flight ground control (GC pre), and post-flight ground

control (GC post). Parameters included stride kinematics (length, duration, variability),

limb coordination metrics, and ground reaction forces collected during treadmill

locomotion.

Behavior and spaceflight cluster validation

All behavioral and locomotion analysis was performed using MATLAB. Group

separation was visualized in reduced-dimensional space by projecting z-scored data

onto the first three PCs. We calculated 95% confidence ellipses for each experimental

group using:

$$\text{Ellipse semi-axis} = \chi^2_{(0.95,22)} \cdot \lambda_i$$

(5)

where χ=5.991 and λ_i represents the eigenvalue of the i^{th} PC. Multivariate analysis of

variance (ANOVA) with Hotelling's T² post-hoc tests assessed group differences in PC

space (α=0.05, Bonferroni-corrected).

Feature selection and hypothesis testing

Behavioral variables most associated with spaceflight effects were identified by

examining PC loadings. For each of the first three PCs, we selected variables with

absolute loadings exceeding the 95th percentile of loading magnitudes (equivalent to

the top 5 variables per PC).

Motor neuron population model for movement

To model motor neuron population for biophysical extraction of movement, we

simulated a population of 50 leaky integrate-and-fire neurons with equal proportions of

flexor and extensor connectivity to muscle output. Membrane potential dynamics

followed biophysically realistic neurons (**Table 1**). Neuronal populations between

extensor and flexors were reciprocally connected with inhibitory conductance,

generating coupled generation of spiking activity resembling pattern generation circuits

(CPG) for locomotion.

Table 1. Parameters for the CPG model

Variable	Value

Resting potential (mV)	-65
Reset potential (mV)	-70
Membrane resistance (MΩ)	10
Synaptic conductance (nS)	0.4
Inhibitory reversal potential (mV)	-80
Synaptic decay (ms)	5
Mean membrane time constant (ms)	10
Refractory period (ms)	3

We also incorporated "kick and decay" kinetics during reciprocal inputs of neuronal

populations.

$$\tau m \, dV/dt = -(V - EL) + RmI(t)$$

(**6**)

Inference of locomotory activity

Population firing rate activity was temporally smoothed with a 5 ms Gaussian kernel.

Peak population activity bursts between flexor and extensor neurons were used to

calculate the phase difference and cycle lengths which were used to infer stride length

and variability, respectively. The correlation of variation was calculated by taking the

mean and standard deviation of cycle lengths between population bursts. To infer swing

times, the duration of peak responses in the flexor population was calculated and the

differences were taken for each cycle. Similarly, the stance time was taken as the time

from the end of the previous swing cycle and the next swing cycle. The ratio of both

swing and stance times yielded the swing stance ratio.

REFERENCES

1. Yatagai, F., Honma, M., Dohmae, N. & Ishioka, N. Biological effects of space
environmental factors: A possible interaction between space radiation and
microgravity. *Life Sciences in Space Research* **20**, 113–123 (2019).

2. Chancellor, J. C., Scott, G. B. I. & Sutton, J. P. Space Radiation: The Number One
Risk to Astronaut Health beyond Low Earth Orbit. *Life (Basel)* **4**, 491–510 (2014).

3. Cucinotta, F. A. Space Radiation Risks for Astronauts on Multiple International Space
Station Missions. *PLOS ONE* **9**, e96099 (2014).

4. Willey, J. S. *et al.* The individual and combined effects of spaceflight radiation and
microgravity on biologic systems and functional outcomes. *Journal of Environmental*
Science and Health, Part C **39**, 129–179 (2021).

5. Le, H., Rai, V. & Agrawal, D. K. Cholesterol: An Important Determinant of Muscle
Atrophy in Astronauts. *Journal of Biotechnology and Biomedicine* **6**, 67–79 (2023).

6. Trappe, T. A., Tesch, P., Alkner, B. & Trappe, S. Microgravity-induced skeletal muscle
atrophy in women and men: implications for long-duration spaceflights to the Moon
and Mars. *Journal of Applied Physiology* **135**, 1115–1119 (2023).

7. Dhillon, R. J. & Hasni, S. Pathogenesis and Management of Sarcopenia. *Clin Geriatr*
Med **33**, 17–26 (2017).

8. Takahashi, H., Nakamura, A. & Shimizu, T. Simulated microgravity accelerates aging
of human skeletal muscle myoblasts at the single cell level. *Biochemical and*
Biophysical Research Communications **578**, 115–121 (2021).

9. Larsson, L. *et al.* Sarcopenia: Aging-Related Loss of Muscle Mass and Function.
Physiological Reviews **99**, 427–511 (2019).

10. Corlett, T., Stavnichuk, M. & Komarova, S. V. Population analysis of space travelers.

Life Sciences in Space Research **27**, 1–5 (2020).

11. Kunitskaya, A., Piret, J. M., Buckley, N. & Low-Décarie, E. Meta-analysis of health

research data from greater than three months International Space Station missions.

Acta Astronautica **201**, 420–430 (2022).

12. Kim, S., Ayan, B., Shayan, M., Rando, T. A. & Huang, N. F. Skeletal muscle-on-a-

chip in microgravity as a platform for regeneration modeling and drug screening.

Stem Cell Reports **19**, 1061–1073 (2024).

13. Baek, K.-W. *et al.* Rodent Model of Muscular Atrophy for Sarcopenia Study. *J Bone*

Metab **27**, 97–110 (2020).

14. Brubaker, Douglas. K. *et al.* An interspecies translation model implicates integrin

signaling in infliximab-resistant inflammatory bowel disease. *Sci Signal* **13**,

eaay3258 (2020).

15. Brubaker, D. K. & Lauffenburger, D. A. Translating preclinical models to humans.

Science **367**, 742–743 (2020).

16. Bergendorf, A., Park, J. H., Ball, B. K. & Brubaker, D. K. Mouse-to-human modeling

of microglia single-nuclei transcriptomics identifies immune signaling pathways and

potential therapeutic candidates associated with Alzheimer's disease.

2025.02.07.637100 Preprint at https://doi.org/10.1101/2025.02.07.637100 (2025).

17. Ball, B. K., Kuhn, M. K., Fleeman Bechtel, R. M., Proctor, E. A. & Brubaker, D. K.

Differential responses of primary neuron-secreted MCP-1 and IL-9 to type 2

diabetes and Alzheimer's disease-associated metabolites. *Sci Rep* **14**, 12743

(2024).

18. Ball, B. K., Proctor, E. A. & Brubaker, D. K. Cross-Species Modeling Identifies Gene
Signatures in Type 2 Diabetes Mouse Models Predictive of Inflammatory and
Estrogen Signaling Pathways Associated with Alzheimer's Disease Outcomes in
Humans. in *Biocomputing 2025* 426–440 (WORLD SCIENTIFIC, 2024).
doi:10.1142/9789819807024_0031.

19. Ball, B. K., Park, J. H., Proctor, E. A. & Brubaker, D. K. Cross-disease modeling of
peripheral blood identifies biomarkers of type 2 diabetes predictive of Alzheimer's
disease. 2024.12.11.627991 Preprint at https://doi.org/10.1101/2024.12.11.627991
(2024).

20. Lee, M. J. *et al.* Computational Interspecies Translation Between Alzheimer's
Disease Mouse Models and Human Subjects Identifies Innate Immune
Complement, TYROBP, and TAM Receptor Agonist Signatures, Distinct From
Influences of Aging. *Frontiers in Neuroscience* **15**, (2021).

21. Suarez-Lopez, L. *et al.* Cross-species transcriptomic signatures predict response to
MK2 inhibition in mouse models of chronic inflammation. *iScience* **24**, 103406
(2021).

22. Harris-Warrick, R. M. Neuromodulation and flexibility in Central Pattern Generator
networks. *Curr Opin Neurobiol* **21**, 685–692 (2011).

23. Grillner, S. & El Manira, A. Current Principles of Motor Control, with Special
Reference to Vertebrate Locomotion. *Physiol Rev* **100**, 271–320 (2020).

24. Beheshti, A., Ray, S., Fogle, H., Berrios, D. & Costes, S. V. A microRNA signature
and TGF-β1 response were identified as the key master regulators for spaceflight
response. *PLoS One* **13**, e0199621 (2018).

25. McDonald, J. T. *et al.* NASA GeneLab Platform Utilized for Biological Response to

Space Radiation in Animal Models. *Cancers (Basel)* **12**, 381 (2020).

26. Migliavacca, E. *et al.* Mitochondrial oxidative capacity and NAD+ biosynthesis are

reduced in human sarcopenia across ethnicities. *Nat Commun* **10**, 5808 (2019).

27. Membrez, M. *et al.* Trigonelline is an NAD+ precursor that improves muscle function

during ageing and is reduced in human sarcopenia. *Nat Metab* **6**, 433–447 (2024).

28. Lim, S. C. *et al.* Mutations in LYRM4, encoding iron–sulfur cluster biogenesis factor

ISD11, cause deficiency of multiple respiratory chain complexes. *Hum Mol Genet*

22, 4460–4473 (2013).

29. Mori, F. *et al.* Genetic variants of the NMDA receptor influence cortical excitability

and plasticity in humans. *Journal of Neurophysiology* **106**, 1637–1643 (2011).

30. Akashi, K. *et al.* NMDA Receptor GluN2B (GluRε2/NR2B) Subunit Is Crucial for

Channel Function, Postsynaptic Macromolecular Organization, and Actin

Cytoskeleton at Hippocampal CA3 Synapses. *J Neurosci* **29**, 10869–10882 (2009).

31. Ahmadian, M. *et al.* PPARγ signaling and metabolism: the good, the bad and the

future. *Nat Med* **19**, 557–566 (2013).

32. Kwok, A. *et al.* Altered rodent gait characteristics after ~35 days in orbit aboard the

International Space Station. *Life Sciences in Space Research* **24**, 9–17 (2020).

33. Sundaramurthy, A. *et al.* Effect of stride length on the running biomechanics of

healthy women of different statures. *BMC Musculoskelet Disord* **24**, 604 (2023).

34. MULAVARA, A. P. *et al.* Physiological and Functional Alterations after Spaceflight

and Bed Rest. *Med Sci Sports Exerc* **50**, 1961–1980 (2018).

35. Fitts, R. H., Riley, D. R. & Widrick, J. J. Functional and structural adaptations of

skeletal muscle to microgravity. *J Exp Biol* **204**, 3201–3208 (2001).

36. Marder, E. & Bucher, D. Central pattern generators and the control of rhythmic

movements. *Curr Biol* **11**, R986-996 (2001).

37. Dayan, P. & Abbott, L. F. *Theoretical Neuroscience: Computational and*

Mathematical Modeling of Neural Systems. (MIT Press, Cambridge, Mass., 2001).

38. Trappenberg, T. *Fundamentals of Computational Neuroscience*. (OUP Oxford,

2010).

39. Smith, J. K. IL-6 and the dysregulation of immune, bone, muscle, and metabolic

homeostasis during spaceflight. *npj Microgravity* **4**, 1–8 (2018).

40. Okamura, Y. *et al.* Impact of microgravity and lunar gravity on murine skeletal and

immune systems during space travel. *Sci Rep* **14**, 28774 (2024).

41. An, R. *et al.* Influence of the spaceflight environment on macrophage lineages. *npj*

Microgravity **10**, 1–8 (2024).

42. Capri, M. *et al.* Recovery from 6-month spaceflight at the International Space

Station: muscle-related stress into a proinflammatory setting. *FASEB J* **33**, 5168–

5180 (2019).

43. Crucian, B. *et al.* Immune System Dysregulation Occurs During Short Duration

Spaceflight On Board the Space Shuttle. *J Clin Immunol* **33**, 456–465 (2013).

44. Ferrando, A. A., Paddon-Jones, D. & Wolfe, R. R. Alterations in protein metabolism

during space flight and inactivity. *Nutrition* **18**, 837–841 (2002).

45. Stein, T. P., Leskiw, M. J., Schluter, M. D., Donaldson, M. R. & Larina, I. Protein

kinetics during and after long-duration spaceflight on MIR. *American Journal of*

Physiology-Endocrinology and Metabolism **276**, E1014–E1021 (1999).

46. Moreira-Pais, A., Ferreira, R., Oliveira, P. A. & Duarte, J. A. A neuromuscular

perspective of sarcopenia pathogenesis: deciphering the signaling pathways

involved. *GeroScience* **44**, 1199–1213 (2022).

47. Deschenes, M. R., Roby, M. A., Eason, M. K. & Harris, M. B. Remodeling of the

neuromuscular junction precedes sarcopenia related alterations in myofibers.

Experimental Gerontology **45**, 389–393 (2010).

48. Arnold, W. D. & Clark, B. C. Neuromuscular junction transmission failure in aging

and sarcopenia: The nexus of the neurological and muscular systems. *Ageing*

Research Reviews **89**, 101966 (2023).

49. Furukawa, S. *et al.* Findings from recent studies by the Japan Aerospace

Exploration Agency examining musculoskeletal atrophy in space and on Earth. *npj*

Microgravity **7**, 1–10 (2021).

50. Lee, P. H. U., Chung, M., Ren, Z., Mair, D. B. & Kim, D.-H. Factors mediating

spaceflight-induced skeletal muscle atrophy. *American Journal of Physiology-Cell*

Physiology **322**, C567–C580 (2022).

51. Rowan, S. L., Purves-Smith, F. M., Solbak, N. M. & Hepple, R. T. Accumulation of

severely atrophic myofibers marks the acceleration of sarcopenia in slow and fast

twitch muscles. *Experimental Gerontology* **46**, 660–669 (2011).

52. Endo, Y. *et al.* Exercise-induced gene expression changes in skeletal muscle of old

mice. *Genomics* **113**, 2965–2976 (2021).

53. Dolgalev, I. msigdbr: MSigDB Gene Sets for Multiple Organisms in a Tidy Data
Format. (2022).

54. Korotkevich, G. *et al.* Fast gene set enrichment analysis. 060012 Preprint at
https://doi.org/10.1101/060012 (2021).

55. Yu, G., Wang, L.-G., Han, Y. & He, Q.-Y. clusterProfiler: an R Package for
Comparing Biological Themes Among Gene Clusters. *OMICS: A Journal of*
Integrative Biology **16**, 284–287 (2012).

56. Rohart, F., Gautier, B., Singh, A. & Cao, K.-A. L. mixOmics: An R package for 'omics
feature selection and multiple data integration. *PLOS Computational Biology* **13**,
e1005752 (2017).

ACKNOWLEDGEMENTS

BKB is supported by the NSF GRFP (DGE-1842166) and the NIH T32DK101001 from
the National Institute of Diabetes and Digestive and Kidney Diseases. HFK is supported
by the NSF GRFP (DGE-1842166) and the NIH T32-5T32DC016853.

AUTHOR CONTRIBUTIONS

BKB and HFK: Conceptualization, data curation, formal analysis, investigation,
methodology, visualization, project administration, resources, funding acquisition,
writing-original draft, writing-review & editing.

DATA AVAILABILITY

All human transcriptomics data used from the study is available through Gene

Expression Omnibus with accession numbers GSE111006, GSE111010, and

GSE111016. Mouse transcriptomics data is available through the NASA Open Science

Data Repository with accession OSD-103. Mouse behavioral data is available with

accession OSD-478 on the NASA OSDR.

SUPPLEMENTARY INFORMATION AND CODE AVAILABILITY

All code for analysis is available at

https://github.com/bbkazu5/MouseFLTandHumanSA. All supplementary information

including figures and data is also made available on the GitHub repository.

COMPETING INTERESTS

The authors declare no competing interests.

SUPPLEMENTARY INFORMATION

Supplementary Figure 1. Hallmark pathway enrichment analysis. (a) Significantly enriched pathways on mouse PC3, **(b)** PC4, and **(c)** PC6.

Supplementary Figure 2. Principal component analysis of the merged Afro-Caribbean and Caucasian dataset. Human patients are labeled by their disease condition and their association with a demographic cohort.

838
839
840
841

Supplementary Figure 3. Effect size of behaviors pre- and post-flight. All gait parameter effect sizes were calculated between pre- and post-flight time points.

858
859
860