<u>Application of spatial transcriptomics across organoids: a high-resolution spatial</u>
<u>whole-transcriptome benchmarking dataset</u>

Nucera MR[1,2,3*] , Charitakis N[1,2,3*], Leung RF[1,2,3*], Leichter A[1,2], Tuano N[1,2], Walkiewicz M[1,2], Sawant
V[1,2,3], Rowley L[1,2], Scurr M[1,2], Er PX[1,2], Tan KS[1,2], Sutton R[1,2], Ahmad F[1,2], Saxena R[1,2,3], Maytum
A[1,2,3], Turner DL[1,2,3], Voges HK[1,2,3], Nim HT[1,2,3], Sun X[4], Yang B[4], Li KL[4], Ball G[3,5], Elefanty AG[1,2,3],
Lamandé SR[1,2,3], Lawlor KT[1,2,3], Vanslambrouck JM[1,2,3], Mills RJ[1,2,3,6], Ng ES[1,2,3], Stanley EG[1,2,3],
Werder RB[1,2,3], Little MH[1,2,3,7], Elliott DA[1,2,3,8,9#], Porrello ER[1,2,3,9,10#], Faux MC[1,3,11#], Eisenstat
DD[1,3,12#], Velasco S[1,2,3,#], Rossello FJ[1,2,8,13,#,+], Ramialison M[1,2,3,14,#]

9
1 Stem Cell Medicine Theme, Murdoch Children's Research Institute, Parkville, VIC, Australia
2 The Novo Nordisk Foundation Center for Stem Cell Medicine, reNEW Melbourne, Murdoch Children's
Research Institute, Parkville, VIC, Australia
3 Department of Paediatrics, University of Melbourne, Parkville, VIC, Australia
4 BGI Australia, Brisbane, QLD, Australia
5 Developmental Imaging, Murdoch Children's Research Institute, Parkville, VIC, Australia
6 School of Biomedical Sciences, Queensland University of Technology, Brisbane, QLD, Australia
7 The Novo Nordisk Foundation Center for Stem Cell Medicine, reNEW Copenhagen, University of
Copenhagen, Denmark
8 Australian Regenerative Medicine Institute, Monash University, Melbourne, VIC, Australia
9 Melbourne Centre for Cardiovascular Genomics and Regenerative Medicine, Melbourne, VIC, Australia
10 Department of Anatomy and Physiology, School of Biomedical Sciences, The University of Melbourne,
Melbourne, VIC, Australia
11 Department of Surgery, The University of Melbourne, Parkville, VIC, Australia
12 The Royal Children's Hospital, Parkville, VIC, Australia
13 Department of Clinical Pathology, University of Melbourne, Melbourne, VIC, Australia
14 Systems Biology Institute, Clayton, VIC, Australia
[*]co-first authors
[#]co-corresponding authors
[+]lead contact fernando.rossello@mcri.edu.au

Summary

Stem cell-derived organoids hold promise to model tissue-specific disease. To enable this, it

is crucial to assess how transcriptional signatures, cellular organisation and composition of

organoids compare to *in vivo* counterparts. However, technologies which elucidate regional

molecular identity, like spatial transcriptomics, have been challenging to apply to organoids.

This study presents the first systematic profiling of multiple stem cell derived organoid

models (brain, heart muscle, heart valve, kidney, lung, cartilage, and haematopoietic) with

Stereo-seq, a full transcriptome, spatial transcriptomics assay using on-chip in situ RNA

capture at subcellular resolution. It describes optimisation of this assay to characterise

organoids, use of multiple organoid samples on a single chip, assess differences in RNA

capture efficiency compared to reference tissues and its limitations. This study introduces a

bespoke analysis method that partitions samples into regions and further characterises them.

These findings inform future works to characterise organoids using spatial transcriptomics,

providing insights in optimising RNA capture of multiple organoids across a chip and novel

methods for regional analysis.

45
46

Introduction

Pluripotent stem cell (PSC)-derived organoid models are used to investigate human

development and disease pathogenesis and provide a platform to measure drug response [1].

This requires organoids to accurately recapitulate the corresponding in vivo tissue, both

physiologically and in cellular composition, by replicating in vitro the process of native stem

cell development and organization [2], [3].

Spatial transcriptomics (ST) technologies capture gene expression across tissues while

maintaining anatomical and morphological structure. ST encompasses a wide array of

technologies that are divided into imaging- and sequencing-based assays [4]. This study

focuses on the application of Stereo-seq, a sequencing-based technology capable of capturing

transcriptome-wide gene expression at single cell resolution [5]. Stereo-seq profiles an

embedded tissue section using a specialised chip, that simultaneously allows mRNA capture

and imaging. Given the high cost of ST experiments per sample, finding a cost-effective

approach to profile multiple organoids using a single chip is critical. While ST has previously

been applied to a single type of organoid being placed on Stereo-seq chip [6], [7], [8],

technical challenges arise when using small size organoids that cover a limited portion of the

ST chip. This may result in artificially lower mRNA levels or capture of fewer gene

transcripts, defined throughout this manuscript as the total data capture. To date, there are no

studies that directly compare mRNA expression and gene profiling of organoids against in

vivo tissue. Furthermore, the placement of several individual organoids on a single chip has

proven to be viable for certain organoid types [8], yet studies describing the feasibility and

benefits of simultaneous placement of multiple organoid conditions on the same chip is

limited.

This study performs a systematic spatial transcriptome profiling of multiple organoid models

(brain, heart muscle, heart valve, kidney, lung, cartilage, and haematopoietic) using the

Stereo-seq platform compared to two in vivo reference E13.5 mouse head and 21 post-

conception week (PCW) mouse heart tissues. The study describes the successful placement of

multiple organoids of the same tissue type - cultured under different conditions for certain

tissues - on the same chip. Subsequently, the platform's capability to clearly distinguish

transcriptional gene signatures from these different individual organoids on the same Stereo-

seq chip was tested. The effect of size, shape, and cell-type composition of different

organoids on data capture, when compared to one another and to larger, in vivo reference

E13.5 mouse head and 21 PCW mouse heart tissues was investigated. Finally, despite

challenges in cell type annotation linked to challenges in achieving single-cell resolution, this

study illustrates an effective analysis for uncovering region-specific differences across

organoids as an alternative to cell clustering and classification. This approach enables the

identification of spatially resolved biologically relevant gene expression patterns across

multiple organoid models for the first time.

Results

This study systematically profiled the transcriptome of eight human organoid models (brain,

lung, haematopoietic, transwell kidney, suspension kidney, heart valve, cartilage, and heart

muscle) using the Stereo-seq platform. A modified Stereo-seq workflow was designed to

capture multiple conditions where multiple organoids of the same tissue type were co-

embedded and placed on a single Stereo-seq chip (see Methods) (Fig. 1A). To prevent tissue

detachment from the chip surface, observed in cartilage, heart muscle, and kidney organoids,

Stereo-seq chips were coated with poly-L-lysine, as part of our modified Stereo-seq

workflow. To limit transcript diffusion and allow for accurate spatial gene expression

profiling, tissue-specific cell permeabilization also required optimisation. This optimization

was critical to allow processing of multiple organoid samples, derived from the same tissue,

and different culture conditions on the same chip. Despite that suitable permeabilization

conditions could be identified for most models, haematopoietic and suspension kidney

organoid samples displayed higher levels of transcript diffusion, revealed by transcripts

captured outside of tissue locations, despite optimised permeabilization conditions (Fig. 1A).

Two reference tissues (E13.5 mouse head and 21 PCW mouse heart) were also profiled using

the standard Stereo-seq workflow, with the aim to use these datasets as baselines against the

characterised organoid models (Fig. 1A). To account for the custom Stereo-seq workflow and

compensate for differences in RNA capture across organoids, a custom bioinformatics

workflow was developed, including a region-specific analysis, described in detail below (Fig.

1B). This workflow filtered out low quality organoid samples to ensure comparisons to

baseline were only drawn for models for which the protocol optimisation had been

successful. This then allowed investigation of the increased chip coverage on data capture, to

determine compatibility of the Stereo-seq platform with multiple organoid models. RNA

capture quality across the entire chip was then assessed through a variety of metrics:

comparison of the number of captured counts and genes on a per CellBin basis (Fig. S1C-

S1D), captured DNA nanoballs and unique reads (Fig. S1A-B) and sequencing saturation

(Fig. S1C). Firstly, data capture was evaluated on CellBins (equivalent to single cells

visualised across the corresponding immunofluorescence ssDNA image), and the smallest

analytical unit generated during pre-processing via default cell-segmentation methods on the

accompanying immunofluorescence image (ssDNA, Fig. 1A). Although large variation

persisted in total number of transcripts captured between organoids and in vivo tissue

reference (Fig. S1A-S1B), capture range for both number of genes and counts per CellBin

were found to be comparable across organoid models and in vivo reference tissues (Fig. 1C-

1D), irrespective of sequencing saturation. It is worth noting that only kidney suspension

organoids did not reach a saturation of 80% (Fig. S1C). This suggests that the observed

variation could have been introduced by differences in chip coverage area, rather than

organoid type, emphasised by a positive correlation between total UMIs passing QC and

tissue size of organoid samples compared to in vivo reference tissue (Fig. S1D).

The heart valve and kidney suspension protocol models showed a reduced RNA capture
relative to other organoid models, evidenced by a low number of DNA nanoballs captured
under tissue along with low numbers of mapped unique reads (Fig. S1A-S1B). Additionally,
the haematopoietic organoid chips, displayed an unexpectedly broad data capture range (Fig.
S1A-S1B). This was likely due to the placement of a large number of organoids per chip
leading to increased transcript diffusion, mainly evidenced by the presence of UMIs outside
tissue regions of ssDNA (Fig. 1A). Given these issues, downstream analyses focused on
organoid models with uniform size and shape, resulting in an increased RNA capture across
chips. Individual brain, transwell kidney, and cartilage organoid samples placed on the same
chip showed comparable detected number of genes and transcripts captured compared to
E13.5 mouse head and 21 PCW mouse heart in vivo references (Fig. S1E-S1I). For quality
assurance, the percentage of CellBins expressing housekeeping genes GAPDH/Gapdh and
ACTB/Actb across in vivo references and individual brain and cartilage organoid samples
were investigated (Fig. S2A). While cell-to-cell expression levels of housekeeping genes
varied (Fig. S2C), similar to previously reported single-cell RNA-Seq experiments [9], no
clear spatial expression patterns of GAPDH/Gapdh and ACTB/Actb across the organoids
passing QC filtering (Fig. S2A-S2C) were observed, reflecting the expected homogenous
expression nature of these genes. These observations also revealed no systematic differences
introduced by coating chips with poly-L-lysine (Fig. S2C).

In addition to CellBin segmentation, Stereo-seq experiments can also be analysed using a
region-based approach ("square bin") by grouping adjacent transcript capture areas together.
Given the smaller size of the organoids compared to mouse reference tissues and RNA
capture variability across all profiled tissues (Fig. 1), the effect of increasing bin size on
capture of RNA levels, genes, and biological signal was explored next (Fig. 2). The effects of
altering square bin size, which determines the resolution of defined regions, revealed loss of
tissue coverage at larger bin sizes, which could further reveal differences in platform
performance when profiling in vivo reference tissues and organoid models (Fig. 2A-D). For
both in vivo references and different organoid models, when the bin size is increased from 50
to 75, the expected accompanying proportional decrease in the number of bins passing QC
was not observed (Fig. 2Ei-ii).

Our analysis demonstrated that increasing bin size from 50 to 75 to 100 (25 μm^2, 37.5 μm^2
and 50 μm^2 respectively) accompanied by analysis using bespoke filtering parameters to

eliminate inclusion of background signal changes the overall tissue coverage area in both the

in vivo reference tissues and brain and cartilage organoids at larger bin sizes (Fig. 2, Table

S1). However, due to the reduced size of organoids samples in general, this loss of largely

peripheral signal may contribute to downstream challenges to identify cell type compositions

in the organoids.

It is expected that a bin size increase of 50% would produce a 50% decrease in the number of

bins. In both in vivo reference tissues, increasing the bin size by 50% led to the number of

bins decreasing by more than half, while a doubling of bin size leads to only about one-third

of the total number of bins being retained (Fig. 2Ei, Table S1). In mouse head tissue this

created a drop from 18634 bins at bin size 50 to 9024 bins at bin size 75 and 5313 at bin size

100 (Table S1). This disproportionate reduction in the number of bins with increasing bin

size was more noticeable across organoids compared to in vivo reference tissues (Fig. 2Eii,

Table S1). For the brain organoids, bin size 50 results in a tissue area of 67450 μm^2 being

analysed, and this fell to 49350 μm^2 with analysis performed at bin size 100 (Table S1). We

also found that larger bin sizes necessitated stricter filtering parameters to eliminate the

inclusion of background signal in the tissue region, leading to removal of certain tissue areas

for downstream analysis and resulting in fewer bins (Table S1). These asymmetrical changes

were mirrored in the total calculated tissue area at different bin sizes for both in vivo

reference tissues and organoid models (Table S1). In addition, increasing the bin size across

organoid and reference tissues led to changes in the range of number of genes captured per

bin (Fig. 2F), with the range of distribution number of captured genes increasing as bin size

increased.

Overall, RNA capture proved more effective at smaller bin sizes while minimising

background signal across the aforementioned tissues. Additionally, smaller bin sizes allowed

for maximum tissue area to be included in downstream analysis without the addition of

background noise to the dataset in both in vivo reference and organoid tissues. However, the

relatively low number of captured UMIs and genes across individual samples, either with

CellBins or a square bin size of 50 above (Fig. 1C-1D), hindered analysis at a single-cell

resolution using this platform (Fig. 2C-2D).

To further investigate the feasibility of multi-sample placement on single chips, the effect of

normalising samples across the entire chip was compared to a sample-specific normalisation,

performed independently on samples after sub-setting (Fig. S3). Overall, the spatial
expression of housekeeping (GAPDH, ACTB) or organoid marker genes remained unaffected
across both normalisation methods (Fig. S3). Evidence that normalisation did not skew
biological signals is reflected in the expected homogeneous expression of COL2A1, COL9A3
and SPARC across cartilage organoids, given that they are composed almost exclusively of
chondrocyte cells (Fig. S3).

Despite increasing bin sizes and normalisation, data generated from organoid experiments did
not allow for cell clustering and further cell type classification primarily due to insufficient
capture of cell type defining transcripts and genes. This was contrasted with the clear and
accurate spatial profiles of the in vivo samples (Fig. S2B-S2D). Therefore, an alternative,
exploratory analysis method that considers organoid regions was developed. This approach
allows comparison of gene expression profiles between border and core regions of organoids,
a valuable analysis given previous findings observing region-specific transcriptional
differences within organoids [1], [10], [11] (Fig. 3A)

After removing outliers (bins falling outside of the organoid boundaries), our approach
classifies remaining bins into inner and outer regions, based on their distance from the centre
of the organoid (see methods). Gene expression from bins within each region is then
aggregated using a pseudo-bulk strategy, resulting in a stronger signal that overcomes the low
expression of individual bins. We selected brain, transwell kidney, and cartilage organoids
for this analysis, based on comparable shape and size of organoid samples of the same tissue
type, which allowed benchmarking of the developed region-specific analysis method (Fig. 3
and Fig. S4A-S4I).

The initial quality-filtering step measured the distance from each bin located at the extremes
of the organoid boundaries to its k nearest neighbouring bins and removed those that were
isolated from the rest. This method successfully excluded any low-quality regions (based on
UMI counts) that potentially arose from inconsistent tissue adherence to the chip (Fig. 3A-C,
Fig. S4A-S4F). From this analysis, brain and cartilage organoid samples exhibited fewer
outliers than kidney, likely due to their uniform shape and lesser number of detached cells at
the edge of the organoids (Fig S4D-S4F). This analysis subsequently enabled the
identification of edge, border, and core regions within the assessed organoid models (Fig. 3D,
Fig. S4G-S4I). A differential expression analysis between core and border regions of
organoid models, using a pseudo-bulk analysis and organoid samples as replicates, followed.

It should be noted that this approach does not account for spatially variable gene expression.
In cartilage organoids, no differentially expressed genes (DEGs) were identified between
these regions, confirming the homogenous cell population expected in this model [12] (Fig.
S4J). In contrast, brain organoids showed transcriptional differences between border or core
regions, with 13 genes differentially upregulated in core and 13 genes in border regions (Fig.
3E). Gene set enrichment analysis (GSEA) of biological processes showed that border
regions of these organoids were enriched for gene sets related to ATP synthesis and electron
transport (Fig. 3F), whereas core regions were enriched for gene sets related to glycolysis
(Fig. 3G). These observations align with previous studies showing enrichment of glycolytic
gene sets in cortical neural progenitors which are located at the core of brain organoid [13],
while more energy-demanding neuronal populations are found at the borders of organoids at
this developmental stage [13]. These findings demonstrate that the proposed region-specific
approach allowed the identification of spatially and metabolically distinct cell populations in
brain organoid models, which was also observed when using a Slide-seq assay, an alternative
sequencing ST technique [13].

We further expanded the use of our method to on irregularly shaped heart muscle organoids.
These organoids have a "figure eight" shape, with two pole regions on either side of the
organoid and a centre region, amenable to measurement contractile functions (Fig. 4A-4B)
[14]. Heart muscle organoids were cultured under two different conditions, using a previously
published standard serum-free method (control) [15] and a directed maturation method (DM)
for enhanced maturation and contractile force [16] (Fig. 4A-4B). By applying the developed
region-specific analysis, pole regions were filtered out from downstream analysis due to low
cell density and mRNA capture (Fig. 4A-4B). This shows that this method could act as a
secondary quality control based on tissue distribution. Centre regions of the control and DM
organoids were compared using organoid samples as replicates (Fig. 4B, Fig. S4K-S4L),
showing transcriptional differences between the compared conditions (Fig. 4C). This analysis
identified an upregulation of MRPS12, a mitochondrial riboprotein, in the DM-cultured
organoids (Fig. 4D), suggesting a difference in mitochondrial activity in organoids culture in
DM-media. In control heart muscle organoids, a GSEA analysis identified enriched gene sets
implicated in ion channel activities (Fig. 4E), while gene sets linked to mitochondrial
activities, ATP synthesis, and metabolism were upregulated in DM heart muscle organoids
(Fig. 4F). Enrichment of these gene sets reflect recent findings that reported alternative usage
of ion channels and increase in sarcomere maturity of organoids cultured in DM media

indicated through the downregulation of ion channel genes and upregulation of metabolic
genes, respectively [16]. The transcriptional differences captured through ST between serum-
free control and DM heart muscle organoids resemble previous characterisation of these
models, demonstrating the validity of ST assays to capture transcriptional differences of
organoids grown under different conditions.

Discussion

This study assessed the performance of Stereo-seq, a ST by sequencing assay, in eight iPSC-
derived human organoid models and two mouse in vivo tissue references. The main objective
was to validate this technology across multiple organoid tissue types in comparison to
reference tissues, in which spatial gene expression profiles showed anatomical specificity at
single cell resolution.

Across the eight organoid models assessed, cartilage, brain, and transwell kidney organoids
resulted in sample quality comparable to tissue references, which could be attributed to their
size and relatively uniform, round shapes. Despite this, RNA capture of these samples failed
to achieve single cell gene expression resolution. Organoids with smaller surface areas, such
as suspension kidney and haematopoietic organoids (notwithstanding comparable transcript
capture and total gene counts) displayed high amounts of diffuse gene expression captured
outside of the organoid surfaces. We suspect this was likely due to additional challenges
presented by these organoids, cultured in more of a 'liquid-phase' compared to other tissue
types, resulting in transcript diffusion. Another challenge in applying Stereo-seq to these
organoids was their lack of adhesiveness, likely contributing to non-uniform transcript
capture across chips. As a remedy, poly-L-lysine chip coating was used to maximise tissue
adhesion (see Methods). The data generated from heart muscle organoids demonstrated that
this method was effective in improving organoid adhesion and should be considered for all
organoids to improve data quality.

This study tested the feasibility of placing multi-sample placement on a single chip.
Previously, ST technology required single tissue sections to be placed on separate chips to
yield optimal outcomes [6], [7], [8]. Our analysis revealed that placing up to 12 sections on
the same chip, as in the case of heart muscle organoids, still resulted in accurate transcript
capture, both cultured in the same media, and organoids grown in different culture media,
such as the case of control vs DM heart muscle organoids. The ability to maximise coverage

with multiple sections per chip greatly extends the potential use of ST technology for
organoids, ensuring high data value per chip and reduce inter-chip variability.

The expression of housekeeping genes such as GAPDH and ACTB was uniform across
cartilage, brain, and transwell kidney organoids, although only in less than 50% of the
identified CellBins, resembling expression patterns of these genes in in vivo reference tissues.
However, the identification of region-specific transcriptional signatures in organoids was
limited when using CellBins (i.e. single-cell resolution), likely due to the relatively low
number of transcripts and genes captured per CellBin. While the number of transcripts and
genes detected per bin were comparable to other studies utilising Stereo-seq [5], it was found
to be lower than the accepted range for single-cell RNA-seq experiments. Interestingly,
increasing square bin sizes, at the expense of losing single cell resolution, did not resolve the
challenges in detecting region-specific expression patterns in organoids models. This may
also be a product of the unbiased transcript capture technology coupled with low number of
transcripts captured, leading to an overrepresentation of abundant transcripts. This shows that
while whole-transcriptome ST, including Stereo-seq and Visium HD, could capture
transcripts from organoid models, its utility for identifying region-specific expression
patterns at single cell level in these model systems was limited. However, this higher-
throughput approach for characterisation of organoids could be useful for discerning spatially
distinct molecular signatures, as previously shown to be viable using in vivo tissue samples
signatures [5], [17].

Given the smaller size of organoids compared to in vivo tissue, the structural compartments
in these models could be relatively subtle and marked by fewer cells. Coupled with the
challenge of achieving single-cell resolution, this hindered the ability of ST technique in
capturing spatially resolved gene expression in organoids and hence identification of defined
patterning. The use of cryo-sections may also limit the assessment of these organoids, as
structural compartments could be rare and difficult to capture on a single cell-thick layer.
These challenges demonstrate the need to further optimise protocols to apply ST to organoid
models. As mentioned above, improvements could include applying alternative chip coating
methods to further increase tissue adhesion or organoid fixation to better preserve spatial
gene expression profiles.

As standard clustering and cell type classification approaches could not be applied to dataset
from organoids of this study, we developed an alternative, exploratory regional analysis

method that allowed spatial gene expression profiling in organoids between core and border regions. We demonstrated its power in identifying region-specific biological processes as well as its utility as a secondary quality control filter when applied to brain and heart muscle organoids. Given the use of Euclidean distance to calculate outliers, the application of this analysis as a secondary quality control filter to round organoids enables more accurate calculation of cells that have detached from main organoid bodies. As mentioned, the ability to understand region-specific gene expression patterns in these models could be key to elucidating general structural organisation that may be present in these systems. This knowledge could then be used to further advance organoid models to better reflect their in vivo counterparts, especially with regards to progressing organoid patterning for the study of developmental tissue patterning. This could have significant implications in developmental biology, such as identifying processes of cell differentiation. As ST platforms for RNA capture continue to evolve and expand imaging options, an updated Stereo-seq protocol incorporating H&E imaging, a morphology-based analysis could be incorporated into future analytical pipelines.

Overall, this study explores the use of profiling multiple organoid types using ST platform. It demonstrates the feasibility of placing multiple organoids generated under different culture conditions on a single Stereo-seq chip, allowing capture of distinct biological signatures from each model. Furthermore, we illustrate the abilities and limitations of profiling a range of organoid types with the current Stereo-seq platform, identifying potential steps for future optimisation such as use of poly-L-lysine for adhesion. Given the limitation around current RNA capture in allowing organoid-specific cell-type annotation, which is not mitigated by increasing bin size for analysis, we developed a regional-specific analysis workflow for capturing biologically relevant gene expression. This provides a basis for future optimisation of organoid profiling using the Stereo-seq platform, including maximising data capture from a single experiment.

Limitations of Study

Although the application of Stereo-seq ST technology across a wide range of organoids was examined, other ST technologies may present different outcomes when applied to organoid systems. For instance, other sequencing-based ST technology such as 10x Visium HD, also offers spatial gene expression profiling at single cell resolution, and, given the different chemistry and transcript capture technology, it could produce alternative results [18].

Alternatively, probe-based imaging ST technologies could also perform differently, with sub-
cellular resolution and higher specificity. To validate findings reported in this study,
traditionally spatial gene expression profiling methods such in situ hybridisation or
immunofluorescence could be performed to assess the accuracy of the gene expression
patterns identified. Additionally, expanding the range of in vivo tissues to correspond to the
organoid systems included, such as including mouse kidney and lung tissues, would provide
additional points of reference to validate the use of Stereo-seq ST technology on various
organ and organoid systems.

Methods

Animal maintenance and ethics approval

All animals were housed in the Murdoch Children's Research Institute (MCRI) Disease
Model Unit and all animal works performed were conducted under the approval and in
compliance of the MCRI Animal Ethics Committee agreements (SPPL20327). Timed
pregnant mice were sacrificed by cervical dislocation following MCRI AEC approved
protocols at embryonic day 13.5 where the presence of a vaginal plug after mating indicated
(E0.5). E13.5 mouse heads and 21 postconceptional week (PCW) mouse hearts were fresh
frozen in OCT (SAKURA Cat# 4583) and stored at -800C until further processing.

Maintenance of induced pluripotent stem cells (iPSC)

PB001.1 iPSCs (male) [19] were used for haematopoietic organoid differentiation,
CRL1502.2 iPSCs (female) [20] were used for transwell kidney organoid differentiation,
MAFBmTagBFP:GATA3mCherry reporter iPSCs [20] were used for suspension kidney
organoid differentiation, PB10.5 iPSCs (male) [19] were used for heart organoid
differentiation. Brain organoids were generated from PB001.1 [19] PB001.1 harbouring a
heterozygous 1bp insertion at chr5:177,135,105 (c.2insG) and PB001.1 harbouring a
homozygous 4bp deletion at chr5:177,135,106 (c.3_6delGGAT). BU3 NGST CRISPRi
iPSCs (male) [21] were used for lung organoid differentiation. MCRIi019-A iPSCs (female)
[22] were used for cartilage organoid differentiation. Mycoplasma contamination in all
iPSCs were monitored and excluded by regular testing. Karyotyping of all iPSCs were
performed regularly and no significant genomic imbalances were identified.

All iPSCs, with the exception of those used for brain organoids were grown on human
embryonic stem cell (hESC)-qualified Matrigel Matrix (Corning Cat# 354277), with varying

medium conditions. iPSCs for brain organoids were grown on Geltrex™ LDEV-Free
Reduced Growth Factor Basement Membrane Matrix (Thermo Fisher Scientific Cat #
A1413301). iPSCs used for differentiation of cartilage, transwell kidney, suspension kidney,
and haematopoietic organoids were maintained in Essential 8 Medium (Thermo fisher
Scientific Cat# A1517001). iPSCs used for lung organoid differentiations were maintained in
StemFlex Medium (Thermo Fisher Scientific Cat# A3349401), whereas iPSCs for brain and
heart organoid differentiation were maintained in mTESR1 Medium (Stem Cell Technologies
Cat# 85857).

Organoid preparation for Stereo-seq

Eight organoid models were generated in accordance to previously described differentiation
protocols, including lung airway and alveolar type 2 organoids   [23], [24], [25]
haematopoietic organoids [26], heart organoids serum-free controls [14], [16], [27], heart
organoids with directed maturation (DM) [28], cartilage organoids [12], heart valve
organoids [27], transwell kidney organoids [28], suspension kidney organoids [29], and brain
organoids [13], [30]. Specifically, following alterations were included in respective organoids
to generate different experimental conditions. Both iPSC-derived lung organoids at air-liquid
interface were infected on the apical surface with respiratory syncytial virus (RSV-A2, MOI
1) as previously described for SARS-CoV-2 [31]. After 4 hours, the inoculum was removed
and cells washed. Uninfected and infected samples were collected 72 hours post infection.
From day 48, cartilage organoids were either grown without additional treatment for 14 days
or treated as follows. Condition 1: cartilage organoids were cultured without additional
treatment after day 48 for 7 days followed by addition of 10nM Triiodothyronine (Sigma
Aldrich Cat# T6397) for 7 days, condition 2: cartilage organoids were cultured without
additional treatment after day 48 for 7 days, followed by 7 days in osteogenic medium [12]
and condition 3: cartilage organoids were grown with addition of 10nM Triiodothyronine for
7 days followed by 7 days in osteogenic medium. All cartilage organoid conditions were
harvested at day 62. Brain organoids grown from all three cell lines were collected on day 31.

At the time of harvest, organoids were separated from culturing media and washed in PBS,
then placed in pre-chilled OCT. Organoids were then transferred to plastic mould containing
OCT, and frozen in an ethanol or isopropanol slurry. For brain organoids, three organoids
were embedded together and placed on the chip, whereas four transwell kidney, and cartilage
organoids were embedded together and placed on the same chip. Eighteen heart valve

organoids were embedded together and placed on the same chip. Six heart organoids of each condition (serum-free and DM) were embedded together in an array of 12 organoids. Over 20 haematopoietic and suspension kidney organoids were embedded in a single mould. For lung organoids, a pair of airway and alveolar type 2 sections, either infected or non-infected, were embedded in OCT together. However, the uninfected airway section was lost during sample processing, resulting in only infected airway section present and alveolar type 2 sections remained.

Stereo-seq T-chip coating

For the heart muscle, heart valve, and cartilage organoids, the sections were mounted on poly-L-lysine coated Stereo-seq T-chip to enhance adhesion. Stereo-seq T-chips were washed using nuclease-free H2O twice and dried, then coated with 100µL of 0.01% (w/v) poly-L-lysine (Sigma Aldrich Cat# P8920) for 10 minutes at room temperature. Post incubation, the chips were washed with nuclease-free H2O twice, and dried on a 37oC slide dryer.

Stereo-seq sample processing and library preparation

Samples were prepared and processed using the STOmics Transcriptomics kit (STOmics Cat# 111KT114) in accordance with manufacturer's instructions (BGI Stereo-seq Transcriptomics set for Chip-on-a-slide user manual version B) with slight variations. 10□m sections of mouse tissue samples including E13.5 heads and 21PCW hearts, brain organoids, kidney organoids, lung organoids, and haematopoietic organoids were cryo-sectioned on CM3050S cryostat (Leica) and mounted on room temperature T-chips, then processed according to the STOmics user manual. 10□m cryo-sections of cartilage, heart muscle, and heart valve organoids were placed on room temperature poly-L-lysine coated Stereo-seq T-chips, then processed according to the STOmics user manual. All cryo-sections were fixed in ice cold methanol for 30 minutes at -30oC. Tissue section permeabilization times were as follows: lung, haematopoietic, transwell kidney, suspension kidney, and cartilage organoid sections were permeabilised for 12 minutes, mouse head and brain organoids for 15 minutes, heart organoids and engineered heart valves for 18 minutes, and mouse heart for 24 minutes. The cDNA purification and library preparation of each sample was constructed using the STOmics library preparation kit (STOmics Cat# 111KL114) in accordance with manufacturer's instructions (BGI STOmics Stereo-seq Transcriptomics set for Chip-on-a-

slide user manual version B). Quality of the final cDNA library size was assessed using

D1000 Tapestation Reagent (Agilent Technologies Cat# 5067-5602) and Screentape (Agilent

Technologies Cat# 5067-5582) and the concentration measured with Qubit dsDNA HS kit

(Thermo Fisher Scientific Cat# Q33231). The sequencing of the libraries was performed at

the BGI Oceanic headquarters using MGI DNBSEQ T7 sequencer, with 1260 million reads

targeted recovery per sample.

Bioinformatics Workflow

Preprocessing

The stitched .czi files taken with the Z2 Axio Imager were processed through ImageStudio v

2.1.1 using the ssDNA setting for staining to ensure they passed QC. They were then

transformed into .ipr . files to ensure compatibility with the SAW pipeline. The paired

FASTQ, mask and transformed .ipr files were then processed using the SAW pipeline v 6.12

using default parameters including cell binning. Raw FASTQ files generated from mouse

hearts were aligned using default SAW mapping parameters to the reference genome

GRCm39 while raw FASTQ files generated from organoids were aligned to the reference

genome GRCh38.

Analysis of organoids and reference tissues

All organoids and in vivo reference tissues was conducted using the Squidpy package v1.2.2

[32] and Stereopy package v1.2 [33] with Python v3.8.2.

Evaluation of quality metrics

Quality metrics were evaluated globally across the whole chip for each organoid type, and at

the individual-organoid level for organoids of interest. Number of counts and number of

genes per chip were computed using Stereopy, while number of mapped reads, mRNA-

captured DNA nanoballs under tissue and sequencing saturation values were extracted from

SAW output files and report. Per-organoid metrics were computed after filtering the data to

retain only a single organoid, using the coordinates extracted with the with the

plt.interact_spatial_scatter() function of Stereopy. This analysis was repeated for 3 different

bin sizes (50, 75 and 100). Before computing the number of bins, genes and counts, a filtering

was applied to remove signal outside of the organoid boundaries and bins with extremely low

counts or high mitochondrial gene percentage. This filtering ensured that we counted and

evaluated only bins representing true biological signal. The filtering criteria varied between
kinds of organoids and tissues as it was highly dependent on the specific counts' distribution.
In some case, poor-quality bins were not limited to clear outliers but represented a substantial
part of the distribution, making outlier-based filtering inadequate. The specific filtering steps
are documented in the code associated with the publication.

Organoid-specific regional analysis to uncover spatially-restricted transcriptomic
differences

Artifacts and outliers at the edge of the organoids were first filtered to retain biologically
relevant spatial information, as these frequently contained abnormal amount of UMIs counts
(Fig. 3A-3C, Fig. S4A-S4C). For each of the brain, cartilage and transwell kidney organoids
(which were all largely circular in shape), we first calculated the centroid coordinates, then
Euclidean distance of each point from the centroid was computed. We identified points with
distances greater than the 90th percentile of all distances as "distant points." Then we
proceeded to detect outliers, that are points located outside the organoid represented false
signals. These artifacts needed to be systematically removed to ensure that only biologically
relevant spatial information was retained. These artifacts were located at a distance from the
rest of the organoid cells, making it necessary to apply a targeted filtering approach. To
remove outliers, we applied a k-nearest neighbours (k-NN) approach using the
NearestNeighbors() function from sklearn v1.5.0 and ball_tree algorithm parameter. A k
parameter of 10 was used for brain organoids and 250 for kidney and cartilage. Since most of
the false signals were spatially distant from the organoid cells, k-NN was used to identify
points with abnormally high distances to their nearest neighbours. However, to prevent
central hollow regions from being misclassified as outliers, we restricted this analysis to the
distant bins. For each distant bin, the average distance to its nearest neighbours was computed
and standardized using a Z-score transformation. Bins with a Z-score greater than 2 (or
higher depending on the specific case) were classified as outliers and subsequently removed
from further analysis. (Fig. 3C, Fig. S4D-S4F).

Following the removal of outliers, we computed the convex hull of the remaining bins to
define the spatial boundary of the dataset. To further classify points into distinct spatial
regions of the 'edge', 'border' and 'core', we calculated the minimum distance of each point
to the convex hull boundary. Based on the distance distribution we classified each bin as
either belonging to the: 'edge' for bins within the 10th percentile of boundary distances,

'border' for bins with distances between the edge threshold and half of the maximum

distance, and 'core' for bins with distances exceeding the border threshold (Fig. 3D, Fig.

S4G-S4I). The 'edge' of the organoid often contained an abnormal number of counts and was

removed from further analysis. After spatial classification, we generated pseudo-bulk samples

using core and border regions of individual organoids placed on the same chip and performed

differential expression analysis between the core and border regions of the same organoid

types. Pseudobulk was performed using the pydeseq2 v1.6.0 and decoupler v0.4.11 packages.

The get_pseudobulk() function was used to generate pseudobulk samples from the anndata

object using the core and border regions from each kind of organoid. This was then built into

a DeSeq2 object and differential expression analysis performed. We treated the organoids on

the same chip as pseudo-replicates, as even if the organoids were not exactly biological

replicates but belonged to either different lines subjected to editing (brain organoids) or

different growing conditions (cartilage organoids), the aim of the analysis was to identify

regional differences in general within the inner and outer areas of the organoids, regardless of

the organoid conditions. Subsequently, we conducted Gene Set Enrichment Analysis (GSEA)

to determine enriched pathways and biological processes associated with differentially

expressed genes.

For the heart organoids which had an elliptic shape, to approximate the organoid boundary, a

convex hull was computed using the set of spatial coordinates. Specifically, the algorithm

Quickhull was used to identify the smallest convex polygon capable of enclosing all points.

Then each bin's distance to the convex hull boundary was computed by treating each bin's

coordinates as a point and measuring its distance to the perimeter of the convex polygon. This

metric provided an estimate of whether a bin was located near the tissue edge (small distance)

or more centrally (larger distance). To remove perimeter effects and focus on interior tissue

regions, bins in the lower 15th percentile of boundary distance were excluded. The remaining

data (with distances above this threshold) formed the "filtered" subset of bins, considered

representative of the tissue's interior.

The spatial coordinates of the filtered bins were then subjected to PCA. The first principal

component (PC1) was assumed to capture the tissue's major "long axis." This vector (PC1)

was used for subsequent segmentation of the tissue into poles and centre regions. Each

filtered bin was projected onto the long-axis vector by calculating a dot product between the

spot's spatial coordinates and the principal-axis vector. This yielded a one-dimensional scalar

value for each spot, referred to as its "projected distance." The minimum and maximum
projected distances defined the overall range (or "length") along this axis. The axis was then
divided into three segments of equal length, the first segment from the minimum projection to
one-third of the total range, the middle segment covering the next one-third of the range and
the final segment from two-thirds of the range to the maximum projection. Bins in the bottom
or top third of this range were labelled as "pole", whereas spots in the central third were
labelled as "centre." Differential expression analysis in pseudo-bulk was performed between
centre of the serum free and directed maturation heart organoids.

Figure Legends

Figure 1 – Overview of quality metrics across multiple organoid types and in vivo reference
tissues when captured using the Stereo-seq platform.

(A) Schematic of organoid models and in vivo reference tissues included on individual
Stereo-seq 1cm2 chips. Immunofluorescence images display regions of ssDNA capture
across the chip while accompanying areas of high-count expression are visualised as UMI
heatmaps. White dotted lines indicate organoid boundaries on UMI heatmaps and
immunofluorescence images.

(B) Schematic of experimental design. i) Steps included from sample preparation and
optimisation through to data analysis and investigation of regional patterns. ii) Outline of
datasets included in each experimental step. Protocol optimisation and cross-organoid
profiling was performed on a total of 10 Stereo-seq chips, divided across two in vivo
reference tissues and eight organoid types. After assessing chips that passed quality control
filters, both in vivo reference tissues and five organoid types were retained for analysis.
Finally, four organoid models were selected for region-specific analysis patterns of gene
expression.

(C) Number of UMI counts per CellBin across each chip.

(D) Number of genes per Cellbin across each chip.

Figure 2 – Effect of increasing bin size (50, 75, 100) on spatial patterning and data
visualisation across in vivo reference tissues and organoids.

(A) Immunofluorescence image of ssDNA capture and heatmaps visualising regions of high
counts and genes per bin across in vivo reference tissue E13.5 mouse head at increasing bin
sizes.

(B) Immunofluorescence image of ssDNA capture and heatmaps visualising regions of
counts and genes per bin across in vivo reference tissue 21 PCW mouse heart tissue at
increasing bin sizes.

(C) Immunofluorescence image of ssDNA capture and heatmaps visualising regions of
counts and genes per bin across a single brain organoid at increasing bin sizes. Brain
organoid 1 displayed in the heatmaps is highlighted in immunofluorescence image.

(D) Immunofluorescence image of ssDNA capture and heatmaps visualising regions of
counts and genes per bin across a single pseudo-replicate of cartilage organoid. Cartilage
organoid 1 displayed in the heatmaps is highlighted in immunofluorescence image.

(E) Bar plots of total number of bins captured across tissue at different bin sizes. (i) In vivo
reference E13.5 mouse head and 21 PCW mouse heart tissue. (ii) Brain organoid 1 and
Cartilage organoid 1.

(F) Distribution of number of genes captured in each bin at bin sizes 50, 75 and, 100 for in
vivo reference E13.5 mouse head, 21 PCW mouse heart, brain organoid 1 and cartilage
organoid 1.

Figure 3 – Analysis of spatially-restricted, differentially expressed genes across centre and
border regions.

(A) Schematic of representative workflow of region-specific analysis between core and
border regions demonstrated using cartilage organoids.

(B) Heatmap of regions of UMI counts across brain organoid 1 using CellBin.

(C) Outlier CellBins identified and visualised in magenta across brain organoid 1,
representing a false positive signal to be removed before region analysis.

(D) Edge (green), border (blue) and centre (red) regions highlighted for analysis in brain
organoid 1.

(E) Volcano plots of genes upregulated in core (visualised in red) and those upregulated in
border region (visualised in blue) for brain organoid pseudo-replicates 1-4.

(F) Gene set enrichment analysis output showing top 10 negatively enriched terms in core of
brain organoids. Each curve represents the enrichment profile the gene set. Legend
annotations include Normalised Enrichment Score, p-value, and FDR q-value.

(G) Gene set enrichment analysis output showing top 10 positively enriched terms in core of
brain organoids.

Figure 4 – Analysis of differentially expressed genes across centre regions of heart muscle
organoids of different conditions.

(A) Example of outlined centre (blue) and pole (orange) regions across DM and control heart
muscle organoids.

(B) Overview of immunofluorescent heart muscle organoids across Stereo-seq chip outlining
placement of six DM organoids on the top and six control organoids on the bottom side of the
chip.

(C) PCA plot of data from 3 centre regions of control organoids and 3 regions of DM
organoids.

(D) Volcano plots of genes upregulated in centre of 3 DM organoids (red) compared to genes
upregulated in centre of 3 control organoids (blue).

(E) Gene set enrichment analysis showing top enriched terms in the centre of control heart
muscle organoids.

(G) Gene set enrichment analysis showing top enriched terms in the centre of DM heart
muscle organoids.

Supplementary Figure 1- Overview of data capture across different organoids and in vivo
reference tissues.

(A) Total DNA nanoballs captured across each Stereo-seq chip for different organoids and in
vivo reference tissues.

(B) Total number of UMIs captured across each Stereo-seq chip for different organoids and
in vivo reference tissues.

(C) Sequence depth and sequence saturation curves for different organoids and reference
tissues. Vendor recommended 80% saturation cut-off displayed as blue dotted line.

(D) Correlation between total counts and number of CellBins across different organoids
passing QC and two in vivo reference tissues.

(E) Total number of CellBins across in vivo reference tissues and individual brain, kidney
transwell and cartilage organoids.

(F) Distribution of number of counts per CellBin across in vivo reference tissues and single
organoids from chips passing QC.

(G) Distribution of number of genes per CellBin across in vivo reference tissues and single
organoids from chips passing QC.

(H) Distribution of number of counts per bin size 50 across in vivo reference tissues and
single organoids from chips passing QC.

(I) Distribution of number of genes per bin size 50 across in vivo reference tissues and single
organoids from chips passing QC.

Supplementary Figure 2 - Overview of data capture across different organoids and in vivo
reference tissues.

(A) Bar plot of percentage of CellBins expression housekeeping genes GAPDH/Gapdh and
ACTB/Actb across in vivo reference tissues and single organoids from chips passing QC.

(B) Dot plot of relative intensity of expression of key marker genes across different organoid

types and in vivo reference tissues.

(C) Heatmap of gene expression housekeeping and mitochondrial genes across brain,

cartilage, heart muscle, kidney transwell and lung organoid chips compared in vivo reference

tissues E13.5 mouse head and 21PCW mouse heart.

(D) Heatmap of tissue-specific marker gene expression across brain, cartilage, heart muscle,

kidney transwell and lung organoid chips compared in vivo reference tissues E13.5 mouse

head and 21PCW mouse heart.

Supplementary Figure 3 – Effect of application of normalisation across chip or per organoid

on expression of housekeeping and marker genes for brain, cartilage and kidney transwell

organoids.

Supplementary Figure 4 - Analysis of spatially-restricted, differentially expressed genes

across centre and border regions for individual brain, cartilage and kidney transwell

organoids.

(A) Heatmap of UMI counts across brain organoid 2-3 using CellBin. Regions of high UMI

counts are displayed in dark purple.

(B) Heatmap of UMI counts across cartilage organoids 1-4 using CellBin. Regions of high

UMI counts are displayed in dark purple.

(C) Heatmap of UMI counts across kidney transwell organoid 1-2 using CellBin. Regions of

high UMI counts are displayed in dark purple.

(D) Outlier CellBins identified and visualised in magenta across brain organoid 2-3,

representing a false positive signal to be removed before region analysis.

(E) Outlier CellBins identified and visualised in magenta across cartilage organoid 1-4,

representing a false positive signal to be removed before region analysis.

(F) Outlier CellBins identified and visualised in magenta across kidney organoid replicates 1-

2, representing a false positive signal to be removed before region analysis.

(G) Edge (blue), border (green) and centre (red) regions highlighted for analysis in brain
organoid replicates 2-3.

(H) Edge (blue), border (green) and centre (red) regions highlighted for analysis in cartilage
organoids 1-4.

(I) Edge (blue), border (green) and centre (red) regions highlighted for analysis in kidney
organoids 1-2.

(J) Volcano plots of genes upregulated in core displayed in red and those upregulated in
border region visualised in blue for cartilage organoid samples 1-4. The analysis did not
return statistically significant differentially expressed genes.

(K) Heatmap of UMI counts 3 selected control heart muscle organoids and 3 selected DM
organoids.

(L) Visualisation of centre and poles for 3 selected control heart muscle organoids and 3
selected DM organoids.

Supplementary Table 1 – Overview of number of retained bins and corresponding tissue area
across bin size 50, 75 and 100 for in vivo reference tissues E13.5 mouse head and 21 PCW
mouse heart and Brain organoid 1 and Cartilage organoid .1

Resource Availability

Lead Contact

Further information and requests for resources should be directed to and will be fulfilled by
the lead contacts, Fernando Rossello (Fernando.rossello@mcri.edu.au) and Mirana
Ramialison (mirana.ramialison@mcri.edu.au).

Data and code availability

•   All original code is available on Github: https://github.com/Ramialison-
Lab/StereoseqOrganoids

Acknowledgements

This work was supported by the Novo Nordisk Foundation Center for Stem Cell Medicine, reNEW, supported by Novo Nordisk Foundation grant number NNF21CC0073729; a BGI STOmics Grant to NC and MR. E.R.P. is supported by an Investigator Grant from the National Health and Medical Research Council (NHMRC) of Australia (GNT2008376). S.V. is supported by an Investigator Grant from the NHMRC of Australia (GNT2025776) and a Fellowship from the Sarah and Lachlan Murdoch Foundation. H.K.V was supported by a Postdoctoral Fellowship (106645) from the National Heart Foundation of Australia. MR is funded by a Future Leader Fellowship (107328) from the Heart Foundation of Australia, a Human Frontiers Science Program Grant (RGP008/2024). MR and HTN are supported by an NHMRC Ideas Grant (APP1180905). Additional infrastructure funding to the Murdoch Children's Research Institute was provided by the Australian Government National Health and Medical Research Council Independent Research Institute Infrastructure Support Scheme. The Australian Regenerative Medicine Institute is supported by grants from the State Government of Victoria and the Australian Government.

Author contributions

Conceptualization: MN, NC, RFL, MR. Methodology: MN, NC, RFL, MR. Software: MN. Formal Analysis: MN. Investigation: MN, NC, RFL, AL, LR, RS, AM, MS, NT, VS, DT, MW, PXE, SKT, RS, FA, HV, HN. Resources: XS, BY, KLL. Data Curation: MN, NC, RFL. Writing Original Draft: MN, NC, RFL, FJR, MR. Writing review and editing: all authors. Supervision: SV, GB, MCF, DDE, FJR, MR, DAE, ERP, RJM. Funding acquisition: NC, MR, AGE, SRL, KTL, JMV, RJM, ESN, EGS, RBW, MHL, DAE, PER, HKV, MCF, DDE, SV.

Declarations of interest

H.K.V. is a co-inventor on a patent held by the Murdoch Children's Research Institute that relates to stem cell derivation of valve cells and tissues. E.R.P. and R.J.M. are co-inventors on patents relating to cardiac organoid maturation and cardiac therapeutics, and are co-founders, scientific advisors, and stockholders in Dynomics. F.J.R. receives institutional and salary support as a) a coinvestigator and subcontractor with the Peter MacCallum Cancer Centre for an investigator-initiated trial which receives funding support from Regeneron

719   Pharmaceuticals; and b) a co-investigator on a translational research project funded by a

720   Regeneron   Pharmaceuticals   grant.

Bibliography

[1]   D. Zhao, W. Lei, and S. Hu, "Cardiac organoid — a promising perspective of preclinical model," Dec. 01, 2021, *BioMed Central Ltd*. doi: 10.1186/s13287-021-02340-7.

[2]   M. Prasad, R. Kumar, L. Buragohain, A. Kumari, and M. Ghosh, "Organoid Technology: A Reliable Developmental Biology Tool for Organ-Specific Nanotoxicity Evaluation," Sep. 23, 2021, *Frontiers Media S.A.* doi: 10.3389/fcell.2021.696668.

[3]   Z. Zhao *et al.*, "Organoids," *Nature Reviews Methods Primers*, vol. 2, no. 1, Dec. 2022, doi: 10.1038/s43586-022-00174-y.

[4]   L. N. Waylen, H. T. Nim, L. G. Martelotto, and M. Ramialison, "From whole-mount to single-cell spatial assessment of gene expression in 3D," Dec. 01, 2020, *Nature Research*. doi: 10.1038/s42003-020-01341-1.

[5]   A. Chen *et al.*, "Spatiotemporal transcriptomic atlas of mouse organogenesis using DNA nanoball-patterned arrays," *Cell*, vol. 185, no. 10, pp. 1777-1792.e21, May 2022, doi: 10.1016/j.cell.2022.04.003.

[6]   P. Wahle *et al.*, "Multimodal spatiotemporal phenotyping of human retinal organoid development," *Nat Biotechnol*, vol. 41, no. 12, pp. 1765–1775, Dec. 2023, doi: 10.1038/s41587-023-01747-2.

[7]   I. Chiaradia *et al.*, "Tissue morphology influences the temporal program of human brain organoid development," *Cell Stem Cell*, vol. 30, no. 10, pp. 1351-1367.e10, Oct. 2023, doi: 10.1016/j.stem.2023.09.003.

[8]   S. Ma *et al.*, "Lamination-based organoid spatially resolved transcriptomics technique for primary lung and liver organoid characterization," *Proc Natl Acad Sci U S A*, vol. 121, no. 46, Nov. 2024, doi: 10.1073/pnas.2408939121.

[9]   Y. Lin *et al.*, "Evaluating stably expressed genes in single cells," *Gigascience*, vol. 8, no. 9, Sep. 2019, doi: 10.1093/gigascience/giz106.

[10]   I. Hautefort, M. Poletti, D. Papp, and T. Korcsmaros, "Everything You Always Wanted to Know About Organoid-Based Models (and Never Dared to Ask)," Jan. 01, 2022, *Elsevier Inc.* doi: 10.1016/j.jcmgh.2022.04.012.

[11]   S. P. Moss, E. Bakirci, and A. W. Feinberg, "Engineering the 3D structure of organoids," Jan. 14, 2025, *Cell Press*. doi: 10.1016/j.stemcr.2024.11.009.

[12]   S. R. Lamandé *et al.*, "Modeling human skeletal development using human pluripotent stem cells," *Proc Natl Acad Sci U S A*, vol. 120, no. 19, May 2023, doi: 10.1073/pnas.2211510120.

[13]   A. Uzquiano *et al.*, "Proper acquisition of cell class identity in organoids allows definition of fate specification programs of the human cerebral cortex," *Cell*, vol. 185, no. 20, pp. 3770-3788.e27, Sep. 2022, doi: 10.1016/j.cell.2022.09.010.

[14]   R. J. Mills *et al.*, "Functional screening in human cardiac organoids reveals a metabolic mechanism for cardiomyocyte cell cycle arrest," *Proc Natl Acad Sci U S A*, vol. 114, no. 40, pp. E8372–E8381, Oct. 2017, doi: 10.1073/pnas.1707316114.

[15]   H. K. Voges *et al.*, "Vascular cells improve functionality of human cardiac organoids," *Cell Rep*, vol. 42, no. 5, May 2023, doi: 10.1016/j.celrep.2023.112322.

[16]   M. Pocock *et al.*, "Maturation of human cardiac organoids enables complex disease modelling and drug discovery," Sep. 09, 2024. doi: 10.1101/2024.09.05.611336.

[17]   P. Xie *et al.*, "Digital reconstruction of full embryos during early mouse organogenesis," *Cell*, vol. 188, no. 17, pp. 4754-4772.e18, Aug. 2025, doi: 10.1016/j.cell.2025.05.035.

[18]   M. F. Oliveira *et al.*, "Characterization of immune cell populations in the tumor microenvironment of colorectal cancer using high definition spatial profiling," Jun. 05, 2024. doi: 10.1101/2024.06.04.597233.

[19]   K. Vlahos *et al.*, "Generation of iPSC lines from peripheral blood mononuclear cells from 5 healthy adults," *Stem Cell Res*, vol. 34, Jan. 2019, doi: 10.1016/j.scr.2018.101380.

[20]   J. M. Vanslambrouck *et al.*, "A toolbox to characterize human induced pluripotent stem cell-derived kidney cell types and organoids," *Journal of the American Society of Nephrology*, vol. 30, no. 10, pp. 1811–1823, 2019, doi: 10.1681/ASN.2019030303.

[21]   R. B. Werder *et al.*, "CRISPR interference interrogation of COPD GWAS genes reveals the functional significance of desmoplakin in iPSC-derived alveolar epithelial cells," Sci Adv 2022. doi: 10.1126/sciadv.abo6566

[22]   K. M. Yammine, S. Mirda Abularach, L. Sampurno, J. F. Bateman, S. R. Lamandé, and M. D. Shoulders, "Using CRISPR/Cas9 to generate a heterozygous COL2A1 p.R719C iPSC line (MCRIi019-A-6) model of human precocious osteoarthritis," *Stem Cell Res*, vol. 67, Mar. 2023, doi: 10.1016/j.scr.2023.103020.

[23]   K. M. Abo *et al.*, "Air-liquid interface culture promotes maturation and allows environmental exposure of pluripotent stem cell-derived alveolar epithelium," *Resource and Technical Advance* 2022, doi: 10.1172/jci. insight.155589.

[24]   F. J. Hawkins *et al.*, "Derivation of airway basal stem cells from human pluripotent stem cells," *Cell Stem Cell*, vol. 28, no. 1, pp. 79–95.e8, 2021, doi: 10.1016/j.stem.2020.09.017.

[25]   A. Jacob *et al.*, "Differentiation of Human Pluripotent Stem Cells into Functional Lung Alveolar Epithelial Cells," *Cell Stem Cell*, vol. 21, no. 4, pp. 472-488.e10, Oct. 2017, doi: 10.1016/j.stem.2017.08.014.

[26]   E. S. Ng *et al.*, "Long-term engrafting multilineage hematopoietic cells differentiated from human induced pluripotent stem cells," *Nat Biotechnol*, 2024, doi: 10.1038/s41587-024-02360-7.

[27]   H. K. Voges, R. J. Mills, E. R. Porrello, and J. E. Hudson, "Generation of vascularized human cardiac organoids for 3D in vitro modeling," *STAR Protoc*, vol. 4, no. 3, Sep. 2023, doi: 10.1016/j.xpro.2023.102371.

[28]   K. T. Lawlor *et al.*, "Cellular extrusion bioprinting improves kidney organoid reproducibility and conformation. HHS Public Access," *Nat Mater*, vol. 20, no. 2, pp. 260–271, 2021, doi: 10.6084/m9.figshare.12957122.

[29]   S. V. Kumar *et al.*, "Kidney micro-organoids in suspension culture as a scalable source of human pluripotent stem cell-derived kidney cells," *Development (Cambridge)*, vol. 146, no. 5, Mar. 2019, doi: 10.1242/dev.172361.

[30]   S. Velasco *et al.*, "Individual brain organoids reproducibly form cell diversity of the human cerebral cortex," *Nature*, vol. 570, no. 7762, pp. 523–527, Jun. 2019, doi: 10.1038/s41586-019-1289-x.

[31]   J. Huang *et al.*, "SARS-CoV-2 Infection of Pluripotent Stem Cell-Derived Human Lung Alveolar Type 2 Cells Elicits a Rapid Epithelial-Intrinsic Inflammatory Response," *Cell Stem Cell*, vol. 27, no. 6, pp. 962-973.e7, Dec. 2020, doi: 10.1016/j.stem.2020.09.013.

[32]   G. Palla *et al.*, "Squidpy: a scalable framework for spatial omics analysis," *Nat Methods*, vol. 19, no. 2, pp. 171–178, Feb. 2022, doi: 10.1038/s41592-021-01358-2.

[33]   S. Fang *et al.*, "Stereopy: modeling comparative and spatiotemporal cellular heterogeneity via multi-sample spatial transcriptomics," Dec. 05, 2023. doi: 10.1101/2023.12.04.569485.

A

Types of Human Organoids

Mouse reference tissue

| | Brain | Lungs | Haema-topoeitic | Transwell Kidney | Suspension Kidney | Heart Valves | Cartilage | Heart Muscle | 21 PCW Mouse Heart | E13.5 Mouse Head |

ssDNA

UMI

1 cm

1 cm

B

Workflow Schematic

Nucera, Charitakis, Leung et al Fig1

C

Number of counts per CellBin in the whole chip

Number of counts

21 PCW mouse heart · E13.5 mouse head · brain · lung · blood · kidney suspension protocol · kidney transwell protocol · valve · cartilage · cardiac

D

Number of genes per CellBin in the whole chip

Number of genes

21 PCW mouse heart · E13.5 mouse head · brain · lung · blood · kidney suspension protocol · kidney transwell protocol · valve · cartilage · cardiac

Nucera, Charitakis, Leung et al 2025 Fig2

Nucera, Chritakis, Leung et al Fig3