The complexity of multiple CRISPR arrays in strains with (co-occurring) CRISPR systems

Axel Fehrenbach[1,2]**, Alexander Mitrofanov**[3]**, Rolf Backofen**[3,4]**, and Franz Baumdicker**[1,2, ✉]

[1]Cluster of Excellence "Controlling Microbes to Fight Infections", Mathematical and Computational Population Genetics, University of Tübingen, 72076 Tübingen, Germany
[2]Institute for Bioinformatics and Medical Informatics (IBMI), University of Tübingen, 72076 Tübingen, Germany
[3]Bioinformatics group, Department of Computer Science, University of Freiburg, 79085 Freiburg, Germany
[4]Signalling Research Centres BIOSS and CIBSS, University of Freiburg, 79085 Freiburg, Germany

CRISPR and their associated Cas proteins provide adaptive immunity in prokaryotes, protecting against invading genetic elements. These systems are categorized into types and are highly diverse. Genomes often harbor multiple CRISPR arrays varying in length and distance from Cas loci. However, the ecological roles of multiple CRISPR arrays and their interactions with multiple Cas loci remain poorly understood.

We present a comprehensive analysis of CRISPR systems that uncovers variation between diverse Cas types regarding the occurrence of multiple arrays, the distribution of their lengths and positions relative to Cas loci, and the diversity of their repeat sequences. Some types tend to occurr as the sole Cas present, but typically comprise two or more arrays, especially for types I-E and I-F. Multiple Cas types are also common, with some systems showing a preference for specific co-occurrence. Distinct array distributions and orientations around Cas loci indicate substantial differences in functionality and transcriptional behavior among Cas types.

Our analysis suggests that arrays with identical repeats in the same genome acquire new spacers at comparable rates, irrespective of their proximity to the Cas locus. Furthermore, repeat similarities in our data set indicate that arrays of systems that often co-occur with other systems tend to have more diverse repeats than those mostly appearing alongside solitary systems within the genome. Our analysis suggests that co-occurring Cas type pairs might not only collaborate in spacer acquisition but also maintain independent and complementary functions and that CRISPR systems distribute their defensive spacer repertoire equally across multiple CRISPR arrays.

CRISPR-Cas | bacterial genomics | CRISPR co-occurrence | spacer acquisition | genome organization | CRISPR repeats
Correspondence: *franz.baumdicker@uni-tuebingen.de*

Introduction

CRISPR systems provide adaptive immunity in bacteria and archaea by incorporating short DNA fragments from invading genetic elements, such as phages and plasmids, into CRISPR arrays. These fragments, so called spacers, enable sequence-specific recognition and degradation of future invaders. CRISPR-Cas plays an important role in shaping microbial evolution and community dynamics [1]. The distribution of CRISPR systems and, in particular, the variability of their CRISPR arrays offer insights into the defense strategies and evolutionary pressures exerted by phages and mobile genetic elements [2, 3].

Prokaryotic genomes encode a remarkable variety of CRISPR systems that differ in their molecular components and organization and provide distinct functions [4]. CRISPR systems are widespread, can be encoded on the chromosome as well as plasmids [5], but exhibit a patchy distribution across bacterial genomes, which indicates that CRISPR systems also carry a cost [6]. Multiple types of CRISPR systems can coexist within a single strain [7, 8] which can lead to interactions between the systems and the associated arrays, such as spacer acquisition in trans [9–13].

In addition to the diversity of types and variety of Cas proteins, the variability of the associated CRISPR arrays are of interest. The spacer arrays provide a partial track record of spacer acquisitions that occurred, e.g. due to phage infection attempts and reflect the defensive capabilities of a cell. Since the frequency of transcription and the degree of immunity conferred by the spacers are influenced by the length of the array and the positions of the spacers [13–15], the composition of CRISPR arrays can provide insights into the arm race between bacteria and phages [3]. Arrays vary in their configuration; they may be located near the Cas genes or dispersed throughout the genome, and can differ in their length (i.e. the number of spacers), leader and repeat uniformity. The repeat sequences, which intersperse with spacer sequences within CRISPR arrays, are known to be type-specific to CRISPRs [8, 16]. The leader sequence plays an important role in regulating spacer integration and expression. In general, the leader sequence is located upstream of the CRISPR array and has been found to be very diverse, containing motifs that are type- but also taxonomy-specific [16, 17]. Together this variability suggests that besides the repeats and leaders, the structure and positioning of CRISPR arrays have important implications for the functional efficiency and adaptability of CRISPR systems.

When multiple CRISPR arrays exist within a bacterial genome, it remains unclear if and if so which arrays are preferentially used for new spacer acquisitions. One key factor may be their genomic proximity to Cas genes, as closer arrays might be more actively transcribed, while distant arrays could serve as archival records. Additionally, the presence of multiple arrays raises the question of whether they serve distinct functional roles or provide redundancy. While previous studies and simulations have explored spacer acquisition dynamics and array length distributions ([7, 8, 18]), a large-scale comparative analysis is still missing. Similarly, the interaction between multiple CRISPR-Cas systems is not yet fully understood. If different systems coexist, do their arrays exhibit

more similar repeats, potentially allowing cross-utilization of Cas proteins, are certain systems incompatible with each other, or do they function independently? Understanding these relationships can shed light on the evolutionary pressures shaping the diversity of CRISPR arrays and systems.

Here we systematically explore a database with thousands of prokaryotic genomes, their CRISPR systems and CRISPR arrays. We utilize the relationship between mean spacer acquisition rates and array lengths to investigate the relative activity of multiple CRISPR arrays and how the array position relative to Cas genes and the repeat sequence influences spacer acquisition. We also consider how multiple CRISPR systems within a genome interact, and find that all these patterns differ substantially between Cas types. We hope that the comprehensive analysis here provides the basis for a better understanding of these differences and can shed light on why certain array configurations are favored and how they impact bacterial immunity.

Methods

Dataset. We used a comprehensive dataset of CRISPR arrays obtained by extracting available whole genomes from GenBank (in November 2024) and running CRISPRidentify [19] to detect all high confidence (labeled by CRISPRidentify as "bona-fide" and "possible") CRISPR arrays within each genome. Cas genes were identified and labeled by CRISPRCasidentifier [20]. Cas types are assigned according to the closest Cas locus, if no Cas was found within the genome they are assigned "no cas type" and called "orphans". Arrays are labeled as "close" to their Cas, if they are less than 10000 bp from their respective Cas locus. Array orientations were determined using CRISPRstrand [21].

We obtained a total of 47789 genomes with 82981 CRISPR arrays, 44760 distinct Cas loci and 19946 unique consensus repeats; see Supplementary Tables S1, S2, S3 and Supplementary Fig. S1 - S5 for additional statistics.

Cas type co-occurrence analysis. We also considered the co-occurrence of Cas types relative to their total abundance. For each possible combination of Cas types, we compared the observed and expected numbers, where the expected number is given by the mean number of such genomes when the presence and absence of Cas types would be shuffled across the dataset. We did this (i) for the observed and expected number of times a system occurs as the only system within a genome, and (ii) the observed and expected number of times a pair of CRISPR systems co-occurrs. The resulting ratios are depicted in Figure 1 and the corresponding data and p-values are given in Supplementary Tables S4 and S5.

Pairwise distances between consensus repeats. We compute any distances between repeats solely for the respective consensus repeats of the arrays. For any two consensus repeats we compute the Levenshtein distance between the two. These distances are normalized by the maximum length of consensus repeats within the genome for Supplementary Fig. S13 or globally for Figure 4, respectively.

Visualization of repeat distances with multidimensional scaling (MDS). We computed the pairwise Levenshtein distances for all consensus repeats for all arrays in the dataset as described above. Then we performed metric MDS, using the obtained distance matrix as dissimilarity measure, to obtain a projection into two-dimensional space using the MDS function of the python package scikit-learn (v. 1.3.2). This MDS function uses the metric SMACOF algorithm and minimizes the stress (residual sum of squares of the original distances and the distances in the 2-d MDS space). The projection of each array's consensus repeat into the MDS space is shown in Figures 4 and 5. All subfigures show subsets of arrays projected using the *same* MDS projection for greater visual clarity.

In addition to the scatterplots of the repeats projected into the two-dimensional MDS space, we provide in Figure 5 kernel density estimates with Gaussian kernels of the marginal distributions for both MDS coordinates and the indicated (colored) subsets, respectively. To compute the kernel density estimates we use the kdeplot function of the python package seaborn (v. 0.12.2) with default parameters, which relies on SciPy (v. 1.9.1) for the Gaussian kernel density estimates.

Estimation of comparative insertion rate weights. To compute the insertion rate weights (Figure 3) we assume an independent deletion model (IDM) as described in [2] which evolves by independent acquisitions and deletions of spacers with insertion rate θ and deletion rate ρ.

In the IDM the array length is Poisson distributed with mean θ/ρ and thus the average length n is given by θ/ρ. As found in [2], deletion rates were found to be independent of Cas type and likely caused by processes, e.g. replication misalignment, which should be consistent for different arrays within a single genome. Thus, we assume the deletion rate to be constant and the insertion rate is likely the main driver of differing array lengths of arrays within the same genome.

Therefore, for a genome with k arrays we can roughly estimate the insertion rate of array $i \in \{1, \ldots, k\}$ as $\theta_i = n_i \rho$, where n_i is the length of the array. Furthermore, we define the total insertion rate for all arrays in the genome as $\theta = \theta_1 + \ldots + \theta_k$ for arrays $i \in \{1, \ldots, k\}$ within one genome with k spacer arrays. To compare the acquisition behavior of CRISPR arrays within the same genome we compute what we call (relative) *insertion rate weights* given by $w_i = \theta_i/\theta$. Clearly, by definition of θ and θ_i, and due to the constant deletion rate, w_i is solely determined by the proportions of array lengths, i.e. we find $w_i = n_i/n$, where $n = n_1 + \ldots + n_k$ is the combined total length of all arrays.

Results & Discussion

Diversity and co-occurrence of Cas types. We found that although the distribution of Cas types is dominated by a few abundant types such as I-E, I-F, III-A, V-A, VI-B and a substantial amount of orphan arrays (where no Cas locus was found in the genome), there is substantial diversity to explore (see Supplementary Tables S1, S2 and Supplementary Fig. S1, S2).

Fig. 1. Single and Co-occurring CRISPR Types For each pair of different Cas types we show the actually observed and the expected (if their occurrence would be reshuffled) number of joint occurrences (in the format observed/expected). For system-pairs where at least 50 genomes are observed or expected the color indicates the fold-change between the observed and the expected number. I.e. a -5 fold-change is shown in red and indicates that the number of observed co-occurrences is 5-fold smaller than expected if CRISPR systems would have been reshuffled, while blue colors indicate a more frequent co-occurrence. On the diagonal the color indicates a different fold-change. Namely, the tendency for each type to occur as the only CRISPR system (positive fold-change, orange) or to occur in combination with at least one other system (negative fold-change, purple). For example types I-E and I-F are alone 2.6-fold and 3.4-fold more often than expected and II-C occurs alone 1.7-fold less often than expected. On the diagonal only systems with at least 50 expected or observed solo ocurrences and with a significant fold-change (bionomial test p-value < 0.0001) got colored.

Moreover, although most genomes carry less than 7 distinct Cas loci with a respective CRISPR array (Supplementary Fig. S3) of less than 4 distinct types (Supplementary Fig. S4), a substantial number of genomes carry at least two different Cas types. Notably, certain combinations of Cas types are much more common than others, suggesting a potential interaction or synergy between some types [22]. Based on the considered dataset this might for example be the case for II-A and I-C, as well as II-C and III-A, and also III-A and V-F (see Figure 1) as those systems often occur together. On the other hand, some systems, including I-E and I-F, tend to occur as the only CRISPR system more often than expected, which could indicate that these systems either provide already sufficient defense or that the presence of other systems interferes with their function.

While other studies have already considered the co-occurrence of cas types [7, 8, 12], here we also consider the number and length of the associated spacer arrays and the similarity of repeat sequences to obtain a more comprehensive perspective.

Cas loci often support multiple arrays. Clearly, the number of arrays tends to exceed the number of Cas loci, indicating that many Cas loci can and do serve multiple arrays (Supplementary Fig. S5). Although many Cas loci have only one array, we found that many Cas loci have multiple close CRISPR arrays, as there is an overall average of 1.48 arrays per Cas locus within the dataset (Supplementary Table S1 and Supplementary Fig. S6). Larger amounts of around 5 arrays do occur quite often and larger numbers (up to around 10) are rare, but do appear (with a maximum of 23 for one V-A locus, see Supplementary Table S1).

Particular outliers are types I-E, I-F and III-A since they have a median of two arrays which we will discuss further below. On the other hand, Type II-A strongly favors only a single array (average 1.07). Similar observations have been made before based on the signature cas genes for type I and II [7].

CRISPR arrays are differently positioned and oriented for different Cas types. Most arrays tend to be close (< 10000 bp) to their respective Cas locus (Supplementary Fig. S2). However, there are huge differences between Cas types. Types I, II, and III tend to favor closeness with few exceptions, but arrays of type V and VI are almost always located farther away (> 500000 bp) from their Cas locus (Supplementary Fig. S7).

We analyze the position of CRISPR arrays in the genome with respect to their corresponding Cas locus by computing the distance between the closest array end and the respective

Fig. 2. Distance to Cas and array orientation. We show the distance of CRISPR arrays to their respective Cas locus. The distances and labels 'before' and 'after' were obtained by comparing the positions of the arrays and Cas genes on the genome, i.e. arrays are positioned 'before'/'after' their Cas cassette and assigned negative/positive "Distance to Cas", respectively. A small number of arrays are contained within their respective Cas locus and were labeled as 'overlap' (with distance 0). Additionally, we indicate the array orientation as predicted by CRISPRstrand. As expected, all types favor close CRISPR arrays, however the different Cas types show large differences in the distribution of distances and orientations with particular biases. In general, arrays close to the Cas locus tend to be forward oriented and more distantly located arrays tend to be reverse oriented. We identify multiple classes of distributions and show representative types for these behaviors here. Distributions for all Cas types with sufficient data can be found in Supplementary Fig. S8. In particular, type I-E has a bias for close arrays to be forward oriented and to be positioned after the Cas locus. Type I-E places close arrays on both sides of the Cas locus. I-E and II-A tend to favor to place the array before the Cas locus. However, II-A shows a significantly wider distribution. Lastly, type III-B shows significantly higher variance and places less importance of closeness to the Cas genes.

Cas locus end. Thereby we determine if they are positioned "before" or "after" the Cas. Moreover, we determined the orientation of the arrays. We show both the distance and orientation (for arrays with close (< 10000 bp distance) Cas) for a subselection of interesting Cas types in Figure 2. Distributions for all types with sufficient data are shown in Supplementary Fig. S8.

Closeness to the Cas locus tends to be favored for all types, and most arrays are less than 500 bp distant from a Cas locus. However, CRISPR types also show very different distributions of distances and distributions of orientations. Firstly, we found a tendency for all types that arrays close to Cas are forward oriented, whereas types further away from the Cas are reverse oriented. We grouped the different subtypes, with respect to their distance and orientation distributions, into multiple groups. For each group we show particularly interesting and representative types in Figure 2; marked **bold** in the text.

1. Type I-C, **I-E**, II-A, and IV-A tend to have arrays *after* and close to the Cas locus (in forward orientation), and arrays *before* the Cas tend to be more distant (in reverse orientation).

2. Type I-A, I-B, I-D, and **I-F** prefer close arrays *both before and after* the Cas locus (in forward orientation). More remote arrays are typically found *after* the Cas (in reverse orientation).

3. Type II-B, **II-C**, **III-A** tend to favor arrays arrays *before* and close to the Cas (in forward orientation). The more distant arrays are usually *after* the Cas (in reversed

orientation). For III-A there are a substantial number of arrays more than 2500 bp away from the Cas locus.

4. Type **III-B** and III-D have a behavior similar to 2. However, they have distributions with much higher variance and less favoritism for specific orientations. This resembles a distribution one might expect in a setting with lower selection pressures that act on the orientation and the distance of CRISPR arrays and the associated Cas locus.

5. Although types V-A and VI-B are very abundant, very few arrays of their arrays are close to their Cas. Where they are close, they show (similarly to 4) more random distributions (see Supplementary Fig. S8).

We further investigated the orientation and position of pairs of CRISPR arrays in genomes with only one Cas type and exactly two CRISPR arrays (Supplementary Fig. S9 and S10). Remarkably, we found that types I-D, I-E and I-F with two arrays almost exclusively position their arrays one on each side of the Cas locus. Possessing two arrays (or more) at the respective ends of the Cas locus is likely beneficial for various reasons. Multiple arrays provide redundancy, and their proximity to the Cas locus maintains high efficiency and enables joint regulation. Moreover, two arrays are likely to have more evenly distributed spacer expression compared to a single longer array, since spacers near their leader positions provide higher levels of immunity [7, 18].

Notably, I-E and I-F are very different in the distribution of array orientation. For I-E the pairs of arrays are almost

exclusively both in forward or both in reverse orientation, whereas for type I-F the distribution of forward and reverse orientation of the two arrays seems to be independent of each other, regardless of whether the arrays are on the same or different sides of the cas locus (Supplementary Fig. S10). I-D favors different orientations for pairs of arrays, but possesses a substantial number of array pairs with the same orientation (Supplementary Fig. S9). III-A behaves differently and typically places both arrays before the Cas and in the same (forward) orientation. The peaks around -2500 to -5000 in the distribution in Figure 2 for type III-A could thus indicate the beginning of a second array next to another array closer to the Cas locus.

Connecting the fact that I-E tends to have arrays in the same orientation with Figure 2, noting that most Cas loci of type I-E have one or two arrays, indicates that single arrays close to the Cas are most often after the Cas and forward oriented or before the Cas and reverse oriented. However, for array pairs one of the arrays is placed close before the Cas and reverse oriented or after the Cas and forward oriented and, quite frequently, the second array is much farther away and on the other side of the Cas (around 7500 bp) in the same orientation. In terms of colors in Figure 2: blue and green tend to form (forward oriented) pairs and red and orange tend to form (reverse oriented) pairs.

Arrays vary in length across Cas types and are typically longer when located near Cas genes . The spacer array length tends to depend heavily on the Cas type, although most arrays have less than 50 spacers (average of 15.11, median of 9) and there is a small number of arrays with between 50 and 100 spacers (Supplementary Table S2 and Supplementary Fig. S11 and S12). In particular, I-A and I-D have remarkably large arrays with a mean of 51 and 42 spacers respectively. Very large arrays are very rare, but e.g. up to 246/317 spacers for type I-E/I-F respectively and 587/587 for type I-B/I-U exist. Types I (excluding I-A/I-D) and type III tend to have averages between 15 and 30, while type II have slightly smaller arrays with averages between 12 (II-C) and 18 (II-B) and types V and VI have far smaller arrays with averages and medians of 10 or lower. If the hypothesis that the loss rate of spacers is comparable between different types and species holds [2] this could indicate that type V and VI CRISPR systems are less often triggered to acquire new spacers, compared to type I, II, and III.

Splitting the set of all arrays regardless of the repeat sequence into two groups depending on how close they are to the Cas and redoing the statistics yields additional insight; see Supplementary Table S3. We found that for most types, arrays close to their Cas are much longer than arrays further away. In particular, type II-C has a large difference between medians 12 (close) and 5 (not close). This holds for most types except I-A, III-B, III-D, V-F, VI-A and VI-C. Thus, one might be tempted to assume that distance to the Cas locus has a large impact on the spacer acquisition rate.

However, this distribution could also be the result of other effects. For example, leaderless arrays have been shown to tend to be shorter [16] and the predicted arrays that are further away could be more often leaderless. In addition, when considering all CRISPR arrays within a genome, this will also include arrays with different repeats and misclassified or other CRISPR-like elements, which could cause the shift towards shorter arrays. In line with this argument, we find that arrays that are not close to a cas locus tend to have diverse repeat sequences that differ from the repeats in arrays that are close (Supplementary Figure S15).

Relative spacer acquisition rates for multiple arrays. We further investigated the insertion behavior by comparing the array lengths of arrays within the same genome. In contrast to the global array length distribution, when looking at the lengths of multiple arrays within the same genome, the deletion rate of spacers should be consistent across different arrays. This allows us to compare array lengths as a proxy for spacer acquisition rates within cells, see Methods and [2]. We call these comparative weights *insertion rate weights* and show their distribution in Figure 3. We restricted our analysis to genomes with exactly 2, 3 or 4 arrays to reduce other biases. We would expect in the neutral case, when all arrays have a similar chance to acquire new spacers, and thus similar lengths, that the relative weight concentrates around the inverse of the number of arrays.

However, in Figure 3a) we see that the overall insertion rate weights are only distributed around 0.5/0.33 0.25, for a number of arrays 2/3, respectively, but not around 0.25 for 4 arrays. The distribution has an increasing weight towards the lower end indicating fewer large and more small arrays. For the arrays close to their Cas (light red/dark red) and the arrays far away from their Cas (blue/dark red) we noted that the distribution of relative lengths is shifted and arrays with close Cas tend to have larger weights than those with a more distant Cas locus. This can also be seen in the distribution of array lengths with respect to their proximity to Cas (Supplementary Table S3, Supplementary Fig. S11 and S12).

One might be tempted to conclude from this pattern that spacer acquisition is more efficient if the acquisition machinery is positioned closer to the array. However, when we further reduced the available arrays to arrays with the same consensus repeat we found that this shift of distributions vanishes. In Figure 3b), the distribution of arrays more distant to the Cas locus is entirely enveloped by the distribution for close to Cas arrays. Moreover, more weight of the distribution is around 0.5/0.33/0.25 as expected for equally distributed acquisitions among the 2/3/4 arrays. This indicates that although the distance to the Cas is correlated with array length, possessing the same (consensus) repeat is far more important for consistent acquisition of new spacers and arrays with the same repeat often maintain comparable lengths and are equally likely updated regardless of their distance to the cas locus. We are aware of only one experimental setup in which acquisitions split among two arrays are considered. While the authors also found that both arrays are used for acquisition the relative acquistion rates are hard to disentangle from a potential copy number bias towards the origin of replication in a replicating bacterial culture [23].

Fig. 3. Distribution of the estimated insertion rate weights of arrays. We show the count of insertion rate weights of arrays for fixed numbers of arrays (genomes with exactly 2, 3 or 4 arrays) colored blue, if the respective Cas genes are nearer than 10000 bp, or colored red otherwise (genomes with orphan arrays are excluded). The count plots are layered over each other (overlap in dark red). The total counts are shown in black. **a)** shows the weights across the whole dataset and **b)** only for arrays with the same consensus repeat. Note that in a), the distribution of arrays which are further away (dark and light red) dominate the the left side of the plots and the distribution of arrays closer to their Cas genes (blue and dark red) are further to the right. This indidates that arrays close to their Cas genes are generally larger than arrays further away. In b), most arrays are close to the Cas genes, but there is no bias of arrays away from the Cas locus to be longer. This indicates that, even though distance from the Cas genes is relevant for the acquisition rate of arrays, possessing the same consensus repeat is of even greater importance.

Repeat distances cluster according to Cas types. We computed the (normalized) Levenshtein distance between consensus repeats of the dataset. We found that, in general, the distribution of the normalized consensus repeat distances is bimodal (or even multimodal) with one focal point around zero and one or even more focal points in the distance range 0.4 - 0.6 for increasing number of arrays and types (Supplementary Fig. S13). Multiple arrays close to the same Cas locus frequently occur, and in this case it is likely beneficial to maintain similar repeats. Moreover, some types are known to use the acquisition machinery of other Cas loci and thus might benefit from similar repeats. Both cases result in the mode around 0. In general, repeat similarity might not be necessary or even detrimental when supporting multiple CRISPR-Cas systems with different tasks. This and an increasing number of co-occurring different types of Cas for which only a limited number are compatible likely leads to the additional modes.

To characterize and compare the repeats within Cas types and for different Cas types further, we performed MDS, with respect to the pairwise distances of the consensus repeats, to project the repeats of each array into a 2-d space; see Figure 4

and Supplementary Fig. S14, S15.

MDS is a method to translate the pairwise distances of a set of points (here the consensus repeats of each array) into a less-dimensional (here 2-d) space such that the pairwise distances are maintained as well as possible. Thus, MDS allows us to visualize the similarity of the consensus repeats losing as little information about their edit distance as possible; see Methods for details about the MDS. Note that the MDS was performed using all arrays, and all Figures are created using the same MDS projection, only different subsets of arrays are shown for visual clarity.

Most type I repeats form clearly distinguishable clusters with some outliers (which occur for all types) that are possibly misclassified arrays. In Supplementary Fig. S15 which shows the repeat MDS colored according to the distance from their Cas locus, it becomes clear that most repeat sequence outliers are also far away from their Cas locus. This further reinforces that these arrays might be misclassified as the wrong type, inactive, or false CRISPR detections [24, 25].

However, it is remarkable that types I-B, I-E and I-F, although they have more similar repeats, co-occur only rarely (Figure 1)

Fig. 4. Visualization of repeat similarity by projection into 2-d space with multidimensional scaling. The plot shows the projection of the consensus repeats (for each array) into 2-d space with multi dimensional scaling with respect to the pairwise consensus repeat edit distances (see Methods for details). The MDS is done globally across all arrays and types, i.e. all shown figures are using the same MDS projection and thus are in the same coordinate system. However, we separated the Cas types into subplots for visual clarity. Each dot corresponds to the consensus repeat of one array and is colored blue, if other Cas types were found within the same genome, and orange otherwise. Each single dot is semi-transparent, thus more intensive color indicates multiple arrays with the same consensus repeat. We show only a selection of representative Cas types here, see Supplementary Fig. S14 for plots of all Cas types. The types show large differences in their distributions. In particular, type I arrays tend to have very concentrated distinguishable distributions and have closely related repeats. However, type II shows much more variance, in particular, type II-C shows multiple clearly distinguishable clusters. Type V-A and VI-B are distributed across the whole space and seem to share similar repeats with all other types. Moreover, the distribution resembles the distribution of repeats of orphan arrays.

with I-E and I-F in particular avoiding most other Cas types, excluding some of the widely spread V and VI subtypes. Type V and VI show very wide distributions that populate almost the whole mds space which is very similar to the seemingly random distribution exhibited by the orphan arrays (Supplementary Fig. S14). This could suggest that they are very widely compatible with other Cas types, an idea that is supported by their relative frequent co-occurence with other types seen in Figure 1 and experimental studies [12, 26] who found that a VI-B array used the acquisition machinery of a II-C locus. However, it can also suggest that many type V and VI arrays might be misclassified or false CRISPR detections [24, 25].

Types II tend to fall somewhere in between types I and V/VI

showing wider distributions with some still distinguishable clusters. In particular, type II-C shows multiple clusters.

Type II-C repeats cluster into subgroups. In Figure 5 we show a more detailed breakdown of the MDS projections with marginal distributions for a subgroup of types. For II-C in a) there are clear differences between repeats of II-C that co-occur with other types and the two smaller clusters where no other Cas types is present. This is consistent with the classification by Makarova et al. [4] where they found two subtypes of II-C arrays. However, considering Supplementary Fig. S15, it becomes clear that arrays in the cluster on the right, which co-occurs with other Cas types, are mostly far away from their Cas locus and thus have remarkably small

Fig. 5. Repeat similarity MDS joint scatterplot and marginal distributions. In this plot, we show the MDS projections of the repeats shown in Figure 4 with additional kernel density estimates of the marginal distributions for both MDS coordinates for interesting subsets of arrays. See methods for details about the MDS and kernel density estimates. **a)** shows only Cas-II-C which clearly shows the existence of different subgroups of repeats. Moreover, we see that they can be separated into groups which occur more frequently and less frequently with other Cas types. **b) - e)** show only the repeats of arrays in genomes that posess arrays of both Cas types that have been found to significantly co-occur (see Figure 1), as noted in each title. **b)** shows that co-occurring II-C and III-B repeats are most often distant from each other. Most II-C repeats are found in the right cluster and are mostly far away from their Cas and shorter in length (see Supplementary Fig. S12, S15). In **c)**, repeats of types I-B and III-B are very similar and similarly distributed with almost complete overlap. In **d)**, some arrays of type III-A and V-F have similar repeats, but in general are relatively distant from each other, and V-F has a much wider distribution than III-A. I-C and II-A in **e)** clearly have distant distinct repeats.

arrays with often less than 5 spacers, see Supplementary Table S3 and Supplementary Fig. S12. It is possible that these are misclassifications, however, none of the other Cas types shows similar behavior, and their close clustering suggests that this might be a special case for type II-C.

In general, repeat similarity might be expected to be important for efficient spacer acquisition in trans. However, Hoikkala et al. [12] show that for II-C and VI-B – arrays with only limited repeat similarity – VI-B can acquire spacers using the II-C machinery.

Repeat similarity and CRISPR co-occurrence. In Figure 5b) - e) we show the repeat MDS projections for pairs of CRISPR types with high co-occurrence rate (Figure 1).

They show a differentiated picture, suggesting that beneficial co-occurrence of types can be efficient for very varied repeat agreement.

We see in Figure 5b) that type III-A has a high overlap with the middle II-C cluster. However, type II-C arrays with co-occuring types (and in particular III-A) are mostly found within the right cluster. Thus, II-C and III-A repeats that co-occur have, in most cases, different repeats.

I-B and III-B clearly have very similar almost exactly overlapping distributions for their repeats (Figure 5c)). Makarova et al. [4] found that I-B has spacer acquisition genes, like Cas1 and Cas2, and III-B does not. This suggests that III-B uses the acquisition machinery of I-B. Vink et al. found results consistent with this in the distribution of spacers and targets of arrays of both types [8].

III-A and V-F share similar repeats for only a subset of co-occurring arrays (Figure 5d)). According to Makarova et al. [4] both type III-A and some subtypes of V-F do not necessarily possess integral genes for spacer integration such as Cas1 and Cas2. This suggests that it might be beneficial for III-A and V-F to maintain similar repeats and pool their Cas machinery to allow for new spacer acquisition and to retain functionality. It is remarkable that, in both Figures 5b) and d), III-A is shown to have a very concentrated distribution of repeats, but co-occurs with much more universalist types in terms of repeat diversity and negative aloneness score (II-C, V-F, VI-B, Figure 1).

I-C and II-A are remarkable in the sense that their repeats are related, but they do not overlap in Figure 5e), moreover, they are the types with the highest co-occurrence ratio and according to Makarova et al. [4] both have functional genes for spacer acquisition. This suggests that these types fulfill complementary and beneficial functions but operate independently for spacer acquisition.

Conclusion

We show that the arrays of CRISPR-Cas systems have a remarkable diversity that can diverge wildly between types. However, some clear patterns emerge. Although array length decreases for arrays distant from the next Cas locus hinting at a low spacer acquisition rate, this effect almost completely vanishes, if the repeats of the arrays agree. This suggests that arrays with identical repeats are equally likely to acquire new spacers, regardless of their genomic position and distance to the cas locus. Moreover, types show clear distinct patterns of positioning and orienting their arrays around the Cas locus. The observed variations might be a result of the selective pressure to diversify the bacterial defense reportoire to avoid or compensate anti-CRISPR mechanisms and persist in the evolutionary arms race with phages.

The shown patterns in the frequency of co-occurrence of Cas types and the differences in repeat (dis)similarity and array orientation invite further exploration. Understanding if and how those CRISPR-Cas systems cooperate and interact for spacer acquisition and inference will help to assess the synergies and incompatibilities of prokaryotic defense systems.

ACKNOWLEDGEMENTS
AF and FB are funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under the Priority Program - SPP 2141 - Project number Ba-5529/1-1. FB is funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy – EXC number 2064/1 – Project number 390727645, and EXC 2124 – Project number 390838134.
This work was supported by German Research Foundation (DFG) [BA 2168/23-1 and BA 2168/23-2 SPP 2141]; Much more than Defence: the Multiple Functions and Facets of CRISPR–Cas; Baden-Wuerttemberg Ministry of Science, Research and Art; University of Freiburg.
This work was supported by the BMBF-funded de.NBI Cloud within the German Network for Bioinformatics Infrastructure (de.NBI) (031A532B, 031A533A, 031A533B, 031A534A, 031A535A, 031A537A, 031A537B, 031A537C, 031A537D, 031A538A).

Bibliography

1. Edze R. Westra, Andrea J. Dowling, Jenny M. Broniewski, and Stineke van Houte. Evolution and Ecology of CRISPR. *Annual Review of Ecology, Evolution, and Systematics*, 47(1): 307–331, 2016. doi: 10.1146/annurev-ecolsys-121415-032428.

2. Axel Fehrenbach, Alexander Mitrofanov, Omer S Alkhnbashi, Rolf Backofen, and Franz Baumdicker. SpacerPlacer: ancestral reconstruction of CRISPR arrays reveals the evolutionary dynamics of spacer deletions. *Nucleic Acids Research*, gkae772, 2024. doi: 10.1093/nar/gkae772.

3. Adrián López-Beltrán, João Botelho, and Jaime Iranzo. Dynamics of CRISPR-mediated virus-host interactions in the human gut microbiome. *bioRxiv*, 10.1101/2024.01.23.576851, 2024. doi: 10.1101/2024.01.23.576851.

4. Kira S. Makarova, Yuri I. Wolf, Jaime Iranzo, Sergey A. Shmakov, Omer S. Alkhnbashi, Stan J. J. Brouns, Emmanuelle Charpentier, David Cheng, Daniel H. Haft, Philippe Horvath, Sylvain Moineau, Francisco J. M. Mojica, David Scott, Shiraz A. Shah, Virginijus Siksnys, Michael P. Terns, Česlovas Venclovas, Malcolm F. White, Alexander F. Yakunin, Winston Yan, Feng Zhang, Roger A. Garrett, Rolf Backofen, John Van Der Oost, Rodolphe Barrangou, and Eugene V. Koonin. Evolutionary classification of CRISPR–Cas systems: a burst of class 2 and derived variants. *Nature Reviews Microbiology*, 18(2):67–83, 2020. doi: 10.1038/s41579-019-0299-x.

5. Rafael Pinilla-Redondo, Jakob Russel, David Mayo-Muñoz, Shiraz A Shah, Roger A Garrett, Joseph Nesme, Jonas S Madsen, Peter C Fineran, and Søren J Sørensen. CRISPR-Cas systems are widespread accessory elements across bacterial and archaeal plasmids. *Nucleic Acids Research*, 50(8):4315–4328, 2022. doi: 10.1093/nar/gkab859.

6. Michael Zaayman and Rachel M. Wheatley. Fitness costs of CRISPR-Cas systems in bacteria. *Microbiology*, 168(7), 2022. doi: 10.1099/mic.0.001209.

7. Jl. Weissman, William F. Fagan, and Philip L.F. Johnson. Selective Maintenance of Multiple CRISPR Arrays Across Prokaryotes. *The CRISPR Journal*, 1(6):405–413, 2018. doi: 10.1089/crispr.2018.0034.

8. Jochem N. A. Vink, Jan H. L. Baijens, and Stan J. J. Brouns. PAM-repeat associations and spacer selection preferences in single and co-occurring CRISPR-Cas systems. *Genome Biology*, 22(1):281, 2021. doi: 10.1186/s13059-021-02495-9.

9. Susanne Erdmann and Roger A. Garrett. Selective and hyperactive uptake of foreign DNA by adaptive immune systems of an archaeon via two distinct mechanisms. *Molecular Microbiology*, 85(6):1044–1056, 2012. doi: 10.1111/j.1365-2958.2012.08171.x.

10. Lisa Nickel, Weidenbach , Katrin, Jäger , Dominik, Backofen , Rolf, Lange , Sita J., Heidrich , Nadja, , and Ruth A. Schmitz. Two CRISPR-Cas systems inMethanosarcina mazeistrain Gö1 display common processing features despite belonging to different types I and III. *RNA Biology*, 10(5):779–791, 2013. doi: 10.4161/rna.23928.

11. Sukrit Silas, Patricia Lucas-Elio, Simon A. Jackson, Alejandra Aroca-Crevillén, Loren L. Hansen, Peter C. Fineran, Andrew Z. Fire, and Antonio Sánchez-Amat. Type III CRISPR-Cas systems can provide redundancy to counteract viral escape from type I systems. *eLife*, 6: e27601, 2017. doi: 10.7554/eLife.27601.

12. Ville Hoikkala, Janne Ravantti, César Díez-Villaseñor, Marja Tiirola, Rachel A. Conrad, Mark J. McBride, Sylvain Moineau, and Lotta-Riina Sundberg. Cooperation between Different CRISPR-Cas Types Enables Adaptation in an RNA-Targeting System. *mBio*, 12(2):e03338–20, 2021. doi: 10.1128/mBio.03338-20.

13. Shayna R Deecker and Alexander W Ensminger. Type I-F CRISPR-Cas Distribution and Array Dynamics in Legionella pneumophila. *G3 Genes|Genomes|Genetics*, 10(3):1039–1050, 2020. doi: 10.1534/g3.119.400813.

14. Jon McGinn and Luciano A. Marraffini. CRISPR-Cas Systems Optimize Their Immune Response by Specifying the Site of Spacer Integration. *Molecular Cell*, 64(3):616–623, 2016. doi: 10.1016/j.molcel.2016.08.038.

15. Chunyu Liao, Sahil Sharma, Sarah L. Svensson, Anuja Kibe, Zasha Weinberg, Omer S. Alkhnbashi, Thorsten Bischler, Rolf Backofen, Neva Caliskan, Cynthia M. Sharma, and Chase L. Beisel. Spacer prioritization in CRISPR-Cas9 immunity is enabled by the leader RNA. *Nature Microbiology*, 7(4):530–541, 2022. doi: 10.1038/s41564-022-01074-3.

16. Omer S Alkhnbashi, Shiraz A Shah, Roger A Garrett, Sita J Saunders, Fabrizio Costa, and Rolf Backofen. Characterizing leader sequences of CRISPR loci. *Bioinformatics*, 32(17): i576–i585, 2016. doi: 10.1093/bioinformatics/btw454.

17. Murat Buyukyoruk, Pushya Krishna, Andrew Santiago-Frangos, and Blake Wiedenheft. Discovery of Diverse CRISPR Leader Motifs, Putative Functions, and Applications for Enhanced CRISPR Detection and Subtype Annotation. *The CRISPR Journal*, crispr.2024.0093, 2025. doi: 10.1089/crispr.2024.0093.

18. Alexander Martynov, Konstantin Severinov, and Iaroslav Ispolatov. Optimal number of spacers in CRISPR arrays. *PLOS Computational Biology*, 13(12):e1005891, 2017. doi: 10.1371/journal.pcbi.1005891.

19. Alexander Mitrofanov, Omer S Alkhnbashi, Sergey A Shmakov, Kira S Makarova, Eugene V Koonin, and Rolf Backofen. CRISPRidentify: identification of CRISPR arrays using machine learning approach. *Nucleic Acids Research*, 49(4):e20–e20, 2021. doi: 10.1093/nar/gkaa1158.

20. Victor A Padilha, Omer S Alkhnbashi, Shiraz A Shah, André C P L F de Carvalho, and Rolf Backofen. CRISPRcasIdentifier: Machine learning for accurate identification and classification of CRISPR-Cas systems. *GigaScience*, 9(6):giaa062, 2020. doi: 10.1093/gigascience/giaa062.

21. Omer S. Alkhnbashi, Fabrizio Costa, Shiraz A. Shah, Roger A. Garrett, Sita J. Saunders, and Rolf Backofen. CRISPRstrand: predicting repeat orientations to determine the crRNA-encoding strand at CRISPR loci. *Bioinformatics*, 30(17):i489–i496, 2014. doi: 10.1093/bioinformatics/btu459.

22. Yi Wu, Sofya K. Garushyants, Anne van den Hurk, Cristian Aparicio-Maldonado, Simran Krishnakant Kushwaha, Claire M. King, Yaqing Ou, Thomas C. Todeschini, Martha R. J. Clokie, Andrew D. Millard, Yilmaz Emre Gençay, Eugene V. Koonin, and Franklin L. Nobrega. Bacterial defense systems exhibit synergistic anti-phage activity. *Cell Host & Microbe*, 32(4):557–572.e6, 2024. doi: 10.1016/j.chom.2024.01.015.

23. Luke J. Peach, Haoyun Zhang, Brian P. Weaver, and James Q. Boedicker. Assessing spacer acquisition rates in E. coli type I-E CRISPR arrays. *Frontiers in Microbiology*, 15, 2025. doi: 10.3389/fmicb.2024.1498959.

24. Quan Zhang and Yuzhen Ye. Not all predicted CRISPR–Cas systems are equal: isolated cas genes and classes of CRISPR like elements. *BMC Bioinformatics*, 18(1):92, 2017. doi: 10.1186/s12859-017-1512-4.

25. Murat Buyukyoruk, William S. Henriques, and Blake Wiedenheft. Clarifying CRISPR: Why Repeats Identified in the Human Genome Should Not Be Considered CRISPRs. *The CRISPR Journal*, 6(3):216–221, 2023. doi: 10.1089/crispr.2022.0106.

26. Elina Laanto, Ville Hoikkala, Janne Ravantti, and Lotta-Riina Sundberg. Long-term genomic coevolution of host-parasite interaction in the natural environment. *Nature Communications*, 8(1):111, 2017. doi: 10.1038/s41467-017-00158-7.