Disrupted priming within draining lymph nodes drives immune quiescence in gastric

cancer

Sohrab Salehi[1,2]*, Emily E. Stroobant[3]*, Hannah Lees[1]**, Ya-Hui Lin[4]**, Shoji Shimada[5]**,

Miseker Abate[3,6]**, Matthew Zatzman[1], Nicholas Ceglia[1], Sam Freeman[1], Monika Laszkowska[7],

Steven Maron[8], Andrew McPherson[1], Nicole Rusk[1], Eliyahu Havasov[1], Harrison Drebin[9], Ping

Gu[8], Laura H. Tang[10], Yelena Y. Janjigian[8], Ruslan Soldatov[1], Ronan Chaligne[11], Sohrab P.

Shah[1]***, Vivian E. Strong[3,6]***, Santosha A. Vardhana[4]***

*These authors contributed equally

** These authors contributed equally

***Co-corresponding

[1]The Halvorsen Center for Computational Oncology, Memorial Sloan Kettering Cancer Center,

New York, NY, USA

[2] Irving Institute for Cancer Dynamics, Columbia University, New York, NY, USA.

[3]Gastric and Mixed Tumor Service, Department of Surgery, Memorial Sloan Kettering Cancer

Center, New York, NY, USA.

[4]Human Oncology and Pathogenesis Program, Memorial Sloan Kettering Cancer Center, New

York, NY, USA.

[5]Digestive Disease Center, Showa University, Northern Yokohama Hospital, Yokohama, Japan.

[6]Department of Surgery, New York-Presbyterian Hospital/Weill Cornell Medicine, New York,

NY, USA.

[7]Gastroenterology, Hepatology, and Nutrition Service, Department of Medicine, Memorial Sloan

Kettering Cancer Center, New York, NY, USA.

[8]Gastrointestinal Oncology Service, Department of Medicine, Memorial Sloan Kettering Cancer

Center, New York, New York.

[9]Vagelos College of Physicians and Surgeons, Columbia University Irving Medical Center, New

York, NY, USA

[10]Department of Pathology, Memorial Sloan Kettering Cancer Center, New York, New York.

[11]Single-cell Analytics Innovation Laboratory, Memorial Sloan Kettering Cancer Center, New

York, NY, USA.

Correspondence: shahs3@mskcc.org, strongv@mskcc.org, vardhans@mskcc.org

Abstract

The gastric mucosa is characterized by continuous innate immune surveillance and inflammatory signaling, yet a high proportion of gastric carcinomas (GCs) are recalcitrant to immune-directed therapies. The mechanisms by which GCs evade adaptive immune surveillance within the highly antigenic microenvironment of the gastric mucosa remains unknown. To address this, we collected patient-matched tumor tissue, distant normal tissue, metastasis, and draining lymph nodes to generate a large-scale single-cell immune profiling dataset from 64 patients (n=179 samples, >150,000 cells). From single cell analysis, we identified two distinct sources of impaired tumor surveillance within tumor draining lymph nodes. First, we observed that a significant fraction of tumor draining lymph nodes had undergone cytokine-driven reprogramming, leading to reduced dendritic cell homing and limited T cell priming. Second, T cells undergoing successful activation exhibited limited expansion and constrained differentiation, marked by expression of the quiescence-associated transcription factor Kruppel-like Factor 2 (*KLF2*). Overexpression of *KLF2* in primary T cells limited both their differentiation and cytotoxic capacity. These findings implicate both impaired T cell priming and *KLF2*-dependent T cell quiescence in limiting T cell immunity in gastric adenocarcinoma. We suggest these findings represent an emerging model for immune silencing in tumors developing from tissues with chronic inflammation.

Introduction

Tumor growth and progression require evasion of immune surveillance; yet, paradoxically, chronic inflammation is a well-established risk factor for multiple cancers. This dichotomy is evident in gastric adenocarcinoma, a malignancy with rising global incidence and poor outcomes, particularly once lymphatic or distant spread occurs[1]. The role of chronic inflammation in gastric carcinogenesis was first established with the classification of *Helicobacter pylori* as a carcinogen in 1994[2]. Subsequent associations with microbiota and autoimmune conditions further positioned chronic inflammation as a central driver of gastric carcinogenesis[3-6]. Yet at clinical diagnosis, most gastric cancers, and particularly those without evidence of microsatellite instability, are thought to be immunologically silent with sparse infiltration of cytotoxic CD8+ T lymphocytes (TILs)[7]. How chronic inflammation disrupts CD8+ T cell-dependent immune surveillance remains unclear. We recently found that intra-tumoral T cell abundance in gastric cancer correlates with effective T cell priming in gastric tumor-draining lymph nodes[8]. This led us to hypothesize that chronic inflammation impairs adaptive immune surveillance by disrupting T cell priming within draining lymph nodes. To test this hypothesis, we developed a surgical collection protocol enabling paired profiling of immune cells from gastric tumors and matched tumor-draining lymph nodes through integrated single-cell RNA sequencing, T-cell receptor (TCR) sequencing and cell-surface epitope assessment (CITE-Seq)[9]. Our findings establish two complementary mechanisms by which sustained innate inflammation disrupts anti-tumor T cell immunity: by i) reducing the frequency of successful T cell priming events within draining lymph nodes and ii) stabilizing transcriptional programs that reinforce T cell quiescence.

Patient cohort, sample collection and single cell data generation

To characterize adaptive immune networks during both the priming and effector stages of the immune response, we developed a tissue collection protocol from patients with gastric adenocarcinoma, emphasizing tumor resections paired with matched draining lymph nodes

(dLN) (**Fig. 1a**). 64 patients with gastric adenocarcinoma were included with patient tumor

characteristics broadly distributed across gastric cancer stages, subtypes and characteristics

(**Extended Data Table 1, 2**). This included at least eight patients from each tumor subtype by

Lauren classification[10], and representation from all TCGA subtypes[11] (**Fig. 1b**). Importantly, a

broad distribution of PD-L1 expression by combined positive score (CPS)[12] was also observed

(0-100), indicating that analyzed tumors represented a broad range of immunogenicity by

conventional metrics. Median follow-up was 2.5 years after initial specimen collection.

In total, 179 fresh tissue samples (74 tumor samples, 56 distant normal samples, 36 draining

lymph nodes, and 13 metastasis samples) were collected from patients undergoing clinically

indicated diagnostic laparoscopy (DL) with esophagogastroduodenoscopy (EGD) or

gastrectomy for gastric adenocarcinoma by a specialist gastric cancer surgeon (Methods,

section Sample Collection and Processing) (**Fig. 1a,b**). All samples were dissociated into single

cell suspensions and viably cryopreserved within 2.5 hours of resection. Subsequently, batches

of no more than 30 samples were thawed, barcoded, and stained for viability. Sorted viable cells

were   multiplexed   and   profiled   by   CITE-Seq   and   single-cell   TCR-Seq[13].

Single cell immune profiling of gastric tumors and matched draining lymph nodes

We first analyzed immune cell subsets from 152,981 CD45+ cells passing quality control (QC)

filtering (**Methods, section** *Single-Cell Analysis***, Extended Data Table 3**). Unsupervised graph-

based clustering identified six main CD45+ cell types marked by canonical identifying genes

(**Fig. 2a, Extended Fig. 1a**): 98,274 T/NK cells (*CD2, CD3D, CD3E*), 23,560 B cells (*CD19,*

MS4A1), 19,774 plasma cells (*JCHAIN, CD38*), 8,876 myeloid cells (*CD14, CD68, S100A9*),

1,687 mast cells (*KIT, TPSAB1*), and 810 plasmacytoid dendritic cells (DCs) (*IL3RA, CLEC4C*).

We then further subclustered CD8+ T cells (**Fig. 2b**), CD4+ T cells (**Fig. 2c**), and myeloid cells/DCs (**Fig. 2d**) using a combination of known marker genes and surface proteins (**Extended Data Table 4**). Classical CD8+ T cell phenotypes including naïve (*SELL, LEF1*), early activated (*FOS, FOSB*), late activated (*HLA-DRA, HLA-DRB1*), effector (*CCL4, XCL1*), central memory (*GZMK, CD28*), proliferating (*MKI67, TYMS*), early exhausted (*TOX, CTLA4*), progenitor exhausted (*THEMIS, SLAMF6*), terminally exhausted (*LAG3, GZMB*), effector memory (*IL7R, ID2*), heat shock protein (HSP) associated (*HSPA1A, HSPA1B),* and effector memory with expression of CD45RA (EMRA) (*PRF1, NKG7*) subsets were identified, as well as a cluster of CD8+ T cells marked by expression of *NEAT1* (*NEAT1, CBLB*) (**Extended Fig. 1b**). CD4+ T cell phenotypes included naïve (*SELL, LEF1*), central memory (*CXCR4, CCR7*), effector memory (*CXCR3, GPR183*), follicular helper (*CXCR5, BCL6*), cytotoxic (*PRF1, GZMB*), HSP-associated (*HSPA1A, HSPA1B*), proliferating (*MKI67, TYMS*), and Th17 (*IL17A, RORC*) cells, as well as a *NEAT1*-expressing cluster (*NEAT1, CBLB*) and two subsets of regulatory T cells (*FOXP3, TIGIT*) (**Extended Fig. 1c**). A proliferative cluster was identified in both CD4+ and CD8+ T cells (*MKI67, TOP2A*) (**Extended Fig. 1d**).

Within the myeloid compartment, we identified mast cells (*KIT, TPSAB1*) as well as dendritic cell subsets including cDC1 (*CLEC9A, XCR1*), cDC2 (*CD1C, CLEC10A*), migratory DCs (*CCR7, LAMP3*), and plasmacytoid DCs (*CLEC4C, LILRA4, TCF4*). We observed two macrophage populations, distinguished by either NLRP3+ inflammasome activation (*NLRP3, IL1B, IL1RN*) or expression of complement proteins (*C1QA, C1QB, C1QC*) (**Extended Fig. 1e**). CD8+ T cells were the more abundant immune cell type within tumors, whereas CD4+ T cells dominated both tumors and dLN (**Fig. 2a, Extended Fig. 1f**). Amongst CD8+ T cells, dLN primarily contained naïve and central memory cells while tumors contained primarily activated, effector memory, and exhausted T cells (**Fig. 2b**). CD4+ T cells within dLN were primarily naïve while tumors contained a mix of regulatory, Th17, and cytotoxic CD4 T cells (**Fig. 2c**). Finally, within the

myeloid compartment, lymph nodes contained predominantly cDC2s and plasmacytoid DCs,

while tumors were marked by NLRP3 macrophages and mast cells (**Fig. 2d**). Heat maps for

normal and metastasis samples are in **Extended Fig. 2**. Lymph node and tumor samples were

primarily non-overlapping in all three cell compartments (CD8+, CD4+ and myeloid).

Patient-specific immune networks defined by cytokine-driven cell-cell communication

We set out to analyze how chronic inflammation influenced the distribution and interactions of

immune cells within gastric tumors and tumor-draining LN. Using a recently generated cytokine

dictionary of gene expression changes in individual immune cell types isolated from mice treated

with a library of 86 distinct cytokines[14] (**Fig. 3a**, **Methods,** section *Tensor factorization for*

Cytokine Activation using IREA), we quantified activation of gene expression programs

associated with cytokine exposure in each cell type from our cohort. Briefly, after mapping cell

types identified in **Fig. 2** to their murine counterparts, we computed cytokine enrichment in each

cell type using an over-representation analysis at the sample level (**Methods,** section *Tensor*

factorization for Cytokine Activation using IREA). We then applied a Bayesian sparse tensor

factorization approach, effectively modeling variation in cytokine-driven expression programs

across samples. This resulted in five robustly enumerated factors, with model weights in both

'receiver' cell-type and 'cytokine' activation dimensions (**Extended Fig. 3a**). Factor 0 was

strongly associated with cDC2 activation, and moderately with pDCs. Factor 1 was strongly

associated with Treg activation, and (γδ) T cells. Factor 2 was associated with CD4 T cells and

Tregs, Factor 3 with macrophages, and Factor 4 with CD8+ T cells, NK cells, and B cells. Factors

0 and 2 were enriched in the dLN, while Factors 3 and 4 were enriched in the tumor; this is

consistent with dendritic cell and CD4+ T cell abundance within lymph nodes, and macrophage

and CD8+ T cell abundance within tumors. Regarding cytokines, IL1a was enriched in Factor 2

while IL15 and TNFa were enriched in Factor 3. IL1 alarmin was the highest activated cytokine

family in Factors 1, 2, and 3, while Factors 0 and 4 were more balanced with the highest activity

from the IL2 family. Growth factors and receptor antagonists were the least activated cytokine

families across all Factors.

We next stratified tumor and dLN samples by factor enrichment. This process yielded two

clusters of cytokine-driven immune networks in dLN, and 3 clusters in tumors (**Methods**, section

Clustering of samples based on IREA scores) (**Fig. 3b**). One dLN cluster ("canonical" LN) was

defined by crosstalk between classical dendritic cells and T cells. Consistent with intact LN

priming events, central memory CD8+ T cells, follicular helper CD4+ T cells, and type 2 and

plasmacytoid dendritic cells were prevalent in this cluster (**Fig. 3c**, **Extended Fig. 3b**). The

second dLN cluster ("disrupted" LN) showed a decrease in cytokine-driven transcriptional activity,

with macrophages as the dominant cytokine-responsive cell type (**Fig. 3c**, **Extended Fig. 3b**).

Notably this cluster harbored an abundance of CD8+ and CD4+ T cells with a naïve phenotype

(**Extended Fig. 3b**) and macrophages with an NLRP3+ phenotype (**Fig. 3c**). Given the relative

dominance of macrophage-driven crosstalk and the absence of dendritic cells needed for

classical T cell priming, we labeled LN within this cluster as "disrupted". Tumor clusters were

similarly interpreted based on cytokine-driven immune cell interactions, with "T cell engaged"

dominated by high-strength T cell interactions, activated and exhausted CD8+ T cells, and Th17

CD4+ T cells, "inflamed myeloid" with primarily NLRP3+ macrophage driven interactions, and

"immune silent" with minimal immune cell infiltration or cytokine-driven interactions (**Fig. 3c,**

Extended Fig. 3b).

We next asked which ligand-receptor interactions contributed most strongly to these cytokine-

driven phenotypes. We inferred intercellular ligand-receptor (L-R) interactions, with a focus on

cytokine-driven L-R interactions[15,16] (**Methods,** section *Receptor Ligand activation analysis*).

Inference of interactions was consistent with dendritic cells as the dominant target of cytokine-

driven L-R interactions within canonical LN, integrating signals from several immune cell subsets.

Analogously, macrophages were the dominant source of cytokine-driven L-R interactions, broadly targeting immune cell subsets within disrupted LN (**Fig. 3d**). Transforming growth factor beta (*TGFB1*) was the dominant source of cytokine-mediated crosstalk within canonical LN, whereas several innate inflammatory cytokines, including IL-1 (*IL1A, IL1B*) and TNF, dominated cytokine-mediated crosstalk in disrupted LN samples (**Fig. 3d**).

We then explored whether paired tumors and their dLN showed co-varying cytokine-driven reprogramming. The two dLN types led to significantly different tumor type distributions (p-value = 0.04, chi-squared test); canonical dLN were more often paired with T-cell engaged tumors, while disrupted dLN were more often paired with either immune silent or inflamed myeloid tumors (**Fig. 3e**). Individual patient tumor and dLN types are listed in **Extended Data Table 5**. There was no clear correlation of genomic features with either tumor or dLN type (**Extended Fig. 3c**). Interestingly, there was an uneven distribution amongst traditional tumor types, with diffuse-type tumors, which carry the poorest prognosis amongst gastric tumor subtypes[17], pairing exclusively with disrupted dLN (**Fig. 3f**). As T cell priming occurs in the lymph node, these results suggest that altered priming within dLN contributes to the lack of intratumoral T cell surveillance in these tumor subtypes.

Reduced T cell priming in disrupted tumor draining lymph nodes

Linked cytokine-driven reprogramming between tumor and matched dLN suggests that chronic inflammation-driven loss of dendritic cell homing to disrupted dLN might impair adaptive immune responses. To investigate this, we next focused on the impact of canonical and disrupted dLN on CD8+ T cell phenotypic trajectories. We generated a CD8+ T cell trajectory manifold (**Methods**, section *CD8+ T cell Trajectory Analysis*) and mapped T cells according to their transcriptional phenotypes (**Fig. 4a**). T cells from dLN and tumors had limited phenotypic overlap (**Fig. 4b**). A computational fate analysis (**Methods**, section *CD8+ T cell Trajectory Analysis*) revealed 6 primary terminal states for naïve CD8+ T cells, including activated, effector, EMRA,

Neat1+, and proliferating fates, as well as a cell fate in which T cells remained in a naïve-like

state (**Fig. 4c**). Using single-cell TCR analysis, we confirmed that clonally expanded T cells were

mostly, though not exclusively in the tumor, with clonally expanded T cells within the tumor

occupying a wide distribution of effector and exhausted states and expanded T cells within the

LN occupying primarily a central memory state (**Extended Fig. 4a**).

We next asked whether T cell trajectories were altered by canonical versus disrupted lymph

node priming by mapping T cell trajectories between canonical and disrupted LN and their most

frequent paired tumor states. In canonical LN-tumor pairs, dLN CD8+ T cells were primarily of

a central memory phenotype and could be found within all six terminal states within the tumor.

In sharp contrast, T cells from disrupted LN-tumor pairs were primarily phenotypically naïve

within the dLN, and their intratumoral trajectories were substantially more restricted (**Fig. 4d**).

This constrained differentiation could be directly attributable to altered priming within the dLN,

as TCR analysis revealed that CD8+ T cells within disrupted LN displayed reduced clonal

expansion and a reduced phenotypic entropy (**Fig. 4e**), with the majority of CD8+ T cells

occupying either a naïve or EMRA phenotypic state (**Fig. 4f**).

Curiously, a fraction of phenotypically naïve CD8+ T cells from disrupted dLN downregulated

surface expression of CD45RA, which typically occurs in the context of antigen experience[18,19]

(**Fig. 4g**), despite being transcriptionally similar to naïve cells (**Fig. 4h**). We therefore stratified

phenotypically naïve T cells based on CD45 isoform expression (CD45RA vs. CD45RO,

included in our CITE-seq antibody-derived tags (ADT) panel). Phenotypically naïve T cells

negative for CD45RA and positive for CD45RO ('RA⁻RO⁺') showed signs of antigen experience

on the protein level, including downregulation of L-selectin and CD3[20,21], as well as upregulation

of Fas[22], consistent with being recently activated. In contrast, T cells negative for CD45RA and

CD45RO ('RA⁻RO⁻') largely overlapped with CD45RA+, CD45RO- canonically naïve T cells, with

sustained expression of L-selectin and CD3 (**Fig. 4i**).   Trajectory analysis of these 3

phenotypically naïve T cell subsets along with early activated T cells confirmed that RA⁻RO⁻ T

cells were similar to RA⁺RO⁻naïve T cells, whereas RA⁻RO⁺ cells were more similar to early

activated T cells (**Fig 4j, Extended Fig. 4b**). We therefore combined double negative RA⁻RO⁻ T

cells with RA⁺RO⁻ true naïve cells and RA⁻RO⁺ cells with early activated T cells for downstream

analysis. We found that compared to canonical dLN, T cells within disrupted dLN were enriched

in either a naïve or EMRA phenotype, indicating that inflammatory reprogramming of disrupted

dLN is sufficient to limit T cell priming and downstream tumor surveillance (**Extended Fig. 4c**).

KLF2 suppresses effector T cell differentiation in gastric tumor-draining LN

Despite showing increased clonal expansion relative to T cells from disrupted LN, T cells from

canonical LN also showed evidence of phenotypic constraint, with the majority of clonally

expanded T cells within dLN remaining in a central memory state (**Extended Fig. 5a**). To identify

transcriptional regulators that might restrict differentiation and egress of CD8+ T cells from dLN,

we analyzed expression of transcription factors in clonally expanded CD8+ T cells compared

with clonally expanded T cells from tumors. We found that Kruppel-like Factor-2 (*KLF2*)[23] was

the most differentially upregulated transcription factor in clonally expanded T cells retained within

dLN (**Fig. 5a**). Within both dLN and tumors, *KLF2* expression correlated strongly with expression

of genes linked to quiescence (*LEF1, TCF7*) and LN retention (*S1PR1, S1PR4, CCR7, SELL*)

and inversely with genes linked to tissue infiltration (*ITGAE*, *CXCR6*) and effector function (*IRF4,*

GZMB) (**Fig 5b**). Moreover, *KLF2* expression inversely correlated with clone size amongst

expanded clones within the LN and tumor, suggesting that *KLF2* expression may reinforce

quiescence and LN retention while restricting T cell expansion, differentiation, and LN egress

(**Fig. 5c**).

To directly assess the consequences of *KLF2* stabilization on T cell activation and differentiation,

we transduced activated primary human T cells with either lentivirus encoding *KLF2* or a vector

control (**Extended Fig. 5b**) and interrogated cells using a combination of flow cytometry and

combined single-cell RNA and ATAC sequencing (DOGMA-Seq)[24]. *In vitro* activated T cells

differentiated into several T cell states marked by distinct transcriptional modules, including

proliferating, effector, effector memory, and central memory T cells, with a minority of T cells

occupying innate-like, activated, or intermediate states (**Fig 5d, Extended Fig. 5c**). Naïve cells

were not present as T cells had been previously activated to facilitate lentiviral transduction.

Consistent with *KLF2* expression and accessibility marking a central memory phenotype[25], we

found that overexpression of *KLF2* in primary T cells led to an increased abundance of CD8+ T

cells with a central memory phenotype and a decreased abundance of T cells with an effector

memory phenotype (**Fig. 5e,f**). To determine the mechanism by which *KLF2* alters T cell

differentiation, we identified differentially accessible motifs in central memory versus effector

memory T cells using ChromVar[26]. We found that CCCTC-binding factor (*CTCF*) and Interferon

Regulatory Factor (*IRF*) motif accessibility were associated with effector memory T cells,

whereas *FOXO1* and T-box (*TBX*) family members were associated with central memory T cells

(**Fig. 5g**). Measurement of promoter-centered chromatin accessibility confirmed that promoter-

associated accessibility of both *CTCF* and *IRF* transcription factors was reduced in *KLF2*-

overexpressing cells (**Fig. 5h**). As both *CTCF* and *IRF* transcription factors are known to be

required for effector CD8+ T cell differentiation[27-32], we measured both effector T cell

differentiation and cytotoxic function and found them to be reduced in *KLF2* overexpressing T

cells (**Fig 5i, Extended Fig. 5d**).

Finally, to identify potential mechanisms by which *KLF2* might limit T cell differentiation and

tumor infiltration, we identified putative *KLF2* target genes by asking which genes containing a

KLF-binding motif[33] (CCACGCCC) were differentially expressed in *KLF2* versus vector

overexpressing CD8+ T cells. *KLF2* overexpression was associated with reduced expression of

target genes associated with effector memory T cells, such as *TOX*, *TOX2*, and *ITGAE* and

increased expression of the GTPase activating proteins (GAP) *RAPGAP2* and *RASA3* (**Fig. 5j**).

RAPGAP2 and *RASA3* bind and stimulate the catalytic activity of the Rap1 GTPase[34-36]. Rap1

has been shown to be required for the TCR-dependent conformational change in Lymphocyte

function-associated antigen-1 (LFA-1) that enables interaction with its ligands, otherwise known

as 'inside out' signaling. This led us to hypothesize that *KLF2*-driven expression of Rap1 GAPs

may limit T cell priming by disrupting LFA-1 activation. Consistent with this hypothesis, surface

expression of both components of the LFA-1 heterodimer (CD11a and CD18) was reduced in

naïve and central memory T cells from disrupted LN; moreover, *KLF2* expression *in vitro* was

sufficient to lower expression levels of CD11a and CD18 when compared to our empty vector

control (**Fig. 5k**). To directly test the impact of LFA-1 activity on T cell priming and differentiation,

we activated OT-I TCR transgenic T cells with cognate peptide across a range of antigen dose

and affinities using altered peptide ligands of the canonical OT-I TCR cognate peptide SIINFEKL

(**Extended Fig. 6a**). Addition of LFA-1 blocking antibodies during T cell priming by peptide-

pulsed DCs increased the threshold for activation, proliferation, and effector function across

peptide doses and affinities, providing a mechanism by which *KLF2* stabilization influences both

T cell priming and effector differentiation in GC (**Extended Fig. 6b,c**).

Cytokine driven dLN types are not fixed

Lymph nodes are typically only surgically resected during gastrectomy, as a fundamental part of

staging. However, for one specific patient (GC028) a biopsy of a visible draining lymph node

was collected during diagnostic laparoscopy (DL) for evaluation of metastatic gastric cancer as

well as during surgical resection after achieving a near complete response to neoadjuvant

chemotherapy (FOLFOX)[37] (**Fig. 6a**). This unique pairing of tumor and draining lymph node both

before and after neoadjuvant chemotherapy allowed us to investigate whether cytokine driven

dLN types remain fixed. Classification of both pre-treatment and post-treatment samples based

on cytokine networks revealed a disrupted dLN prior to treatment and a canonical dLN at the

time of resection (**Fig. 6b**). Accordingly, the post-treatment dLN showed a decreased abundance

of C1QC+ macrophages and an increased abundance of plasmacytoid DCs (**Fig. 6c, Extended**

Fig. 7a).

Clonotype tracking between the dLN and tumor pairs revealed that shared clones were present

between all 4 samples (**Extended Fig. 7b**). Phenotypes of shared CD8+ T cell TCR clonotypes

changed post-treatment, with an increase in the fraction of progenitor Texh in the tumor and CM

in the dLN (**Extended Fig. 7c**). The post-treatment canonical dLN also had both a larger number

of expanded clones and a greater diversity of expanded T cell phenotypes from the dLN to its

paired tumor (**Fig. 6d**). We examined overall *KLF2* expression in CD8+ T cells in the pre-therapy

disrupted dLN and the post-therapy canonical dLN and found that *KLF2* expression was

significantly higher in the disrupted dLN (pre) than the canonical dLN (post) (**Fig. 6e**). *KLF2* was

also the second most transcriptionally decreased gene between these pre-treatment and post-

treatment dLN (**Extended Fig. 7d**). In line with this evidence, expression of both LFA-1

components (CD11a and CD18) was increased in the canonical dLN (post-treatment)

(**Extended Fig. 7e**). The ability for tumor draining lymph nodes to change from disrupted to

canonical types, paired with changes in *KLF2* expression, reveals that cytokine driven dLN types

can respond to therapeutic intervention and may have prognostic implications, as this patient

remains in complete remission 2.7 years after diagnosis despite presenting with metastatic

disease.

Discussion

The paradoxical role of the immune responses in tumorigenesis is evidenced by decades of

epidemiologic studies demonstrating an increased risk of cancers in both immunocompromised

patients and patients with autoimmune conditions[38-41]. This archetype is exemplified by gastric

cancer, where chronic inflammation is a well-established driver of carcinogenesis, yet most

gastric tumors show limited evidence of T cell surveillance. Here we provide a novel mechanistic

explanation for how persistent innate inflammation can impair anti-tumor T cell immunity.

Chronic inflammation limits dendritic cell accumulation within tumor draining lymph nodes,

preventing exposure of putatively tumor-reactive naïve T cells within the lymph node to tumor-

derived neoantigens. Whether targeting innate inflammatory signaling can enhance DC homing

and T cell priming represents an important area for future investigation.

We also identify a second, previously unappreciated checkpoint on T cell surveillance in gastric

cancer. Activated T cells within the draining LN largely fail to differentiate into effectors, instead

becoming central memory T cells that remain in the LN and express high levels of *KLF2*. This

may be in part due to reduced integrin activity increasing the threshold for T cell activation during

naïve T cell priming events within dLN. These findings are consistent with recent evidence

demonstrating a role for *KLF2* in preventing T cell exhaustion[42]. By suppressing the exhaustion

program*, KLF2* may paradoxically reduce the clinical efficacy of immune checkpoint inhibitors

that target inhibitory receptors preferentially expressed on exhausted T cells, including PD-1,

CTLA-4, and LAG-3. Future studies to determine if *KLF2* expression in T cells is associated

with lack of immune checkpoint blockade (ICB) response or whether targeting either *KLF2* or

KLF2-dependent integrin activation can enhance sensitivity to ICB would have immediate clinical

relevance.

Our findings provide insights into how reciprocal balances between innate and adaptive immune

responses maintain homeostasis in an inflammatory microenvironment, and how an imbalance

between these responses contributes to human disease. The extent to which the T cell

phenotypes observed in this study apply to other tumor types remains unknown. However, this

unique dataset, derived from prospective surgical sampling of paired tumors and draining LN,

may serve as a valuable reference to understand whether lymph node priming is disrupted in

other cancers. Blunting of the adaptive immune response during chronic inflammation may

represent a physiologic mechanism to preserve mucosal tolerance and barrier integrity;

consistent with this hypothesis, T cell specific *KLF2* expression is required to maintain

commensal tolerance and prevent inflammatory bowel disease[43]. However, adaptive immune

suppression has been reported in other chronic inflammatory conditions, including sepsis-

associated immune paralysis[44]. Potential targets such as *KLF2* may enable T cells to escape

from this maladaptive state, and the ability for dLN to change with intervention further supports

tumor draining lymph nodes as a viable focus for therapeutic development. Anecdotally, our

report of a patient with pre- and post-treatment tissue collections demonstrating phenotypic T

cell state transitions linked to a remarkable clinical response indicates that the non-canonical

immune states we describe in this study are potentially reversible with therapeutic intervention.

Characterizing the prevalence and determinants of such immune state transitions will be central

to future translational efforts aimed at enhancing lymph node priming in gastric cancer.

Figure Legends

Figure 1. **Patient cohort and sample collection.** (a) Schematic of surgical sample collection from gastric cancer patients and processing for scRNA-seq. (b) Swimmer's plot showing individual patients with samples collected, tumor characteristics, and simplified patient timelines.

Figure 2: Single cell characterization of immune cells from primary tumor, distal normal, draining lymph nodes, and metastases in GC. Uniform manifold approximation and projections **(**UMAP) depicting (a) broad CD45+ cell subsets (left), density heatmap of CD45+ cells from dLN (center) and tumors (right), (b) CD8+ T cell subsets (left), density heatmap of CD8+ cells from dLN (center) and tumors (right), (c) CD4+ T cell subsets (left), density heatmap of CD4+ cells from dLN (center) and tumors (right), and (d) myeloid cell subsets (left), density heatmap of myeloid cells from dLN (center) and tumors (right).

Figure 3. **Patient stratification by distinct cytokine driven immune networks in tumors and their draining lymph nodes.** (a) Workflow of sample level Bayesian sparse tensor decomposition. (b) Clustering of individual patient samples from dLN (left) and tumors (right) based on sparse tensor decomposition from (a). (c) Summary of enriched cytokine-cell type interactions in dLN and tumor subtypes with UMAPs showing the fraction of myeloid cell type content and paired myeloid cell density. (d) Dominant cell type-specific receptor-ligand interactions identified using CellChat in canonical and disrupted dLN subtypes. (e) Correlation of cytokine interaction-dependent dLN and tumor subtypes in patients with paired samples demonstrates a significant association between dLN and tumor types as indicated (p-value = 0.04, chi-squared test) (f) Distribution of Lauren and molecular subtypes across paired dLN-Tumor networks.

Figure 4. **Reduced T cell priming in disrupted tumor draining lymph nodes.** (a-d) Trajectory

manifold of CD8+ T cell subsets across dLN and tumor, showing distribution of (a) CD8+ T cell

subsets, (b) distribution within dLN and tumors, (c) fate probabilities, and (d) density mapping of

cells from 'disrupted-immune silent' pairs and 'canonical-T cell engaged' pairs. (e) Fraction of

shared expanded (clonotypes >20) CD8+ T cell clones between dLN and Tumor (left) and

average phenotype entropy in CD8+ T cells (right) (**Methods**, section *Phenotype entropy*),

stratified by canonical dLN versus disrupted dLN. (f) Difference in mean cell fractions of CD8+

T cell subsets between disrupted dLN and canonical dLN. (g) Surface CD45 isoform expression

distribution on CD8+ T cells from disrupted dLN with a transcriptionally naïve phenotype. (h)

Cells from (g) projected on the trajectory manifold from (a), coded by CD45 isoform expression.

(i) Expression of genes and proteins associated with T cell activation on phenotypically naïve T

cells from (g,h), stratified by surface expression of CD45 isoforms. (j) Density heatmaps of

subtypes of naïve cells from (g-i), projected onto the CD8+ trajectory manifold from (a).

Figure 5. **KLF2 suppresses effector T cell differentiation.** (a) Differentially expressed

transcription factors expressed in non-naïve CD8+ T cells from tumors (up) vs. dLN (down). (b)

Genes correlated with the KLF2 expression in non-naïve CD8+ T cells across tumors and dLN

(c) Correlation of KLF2 expression on non-naïve CD8+ T cells across tumors and dLN with

clonotype size. (d) UMAP of activated human CD8+ T cells transduced with either empty vector

or a KLF2 overexpression vector. (e) Density plot showing relative distribution of CD8+ T cell

phenotypes in primary human CD8+ T cells transduced with either empty vector (left) or KLF2

overexpressing vector (right). (f) Mean fraction of T cell subsets in primary human CD8+ T cells

transduced with either empty vector or KLF2 overexpressing vector as indicated. (g)

Transcription factor motifs with top accessibility scores in central memory (right) compared with

effector memory (left) T cell subsets. (h) Volcano plot demonstrating accessibility of CTCF/L

(left) or IRF4/8 (right) motifs, centered around the transcriptional start site, in human CD8+ T

cells transduced with empty vector or KLF2 overexpression vector. (i) Expression of CXCR6 and

PD-1 (left) and intracellular accumulation of IFN-g and TNF following restimulation with PMA and

ionomycin in CD4+ and CD8+ T cells transduced with empty vector or KLF2 overexpression

vector. *ADT* antibody-derived tags (j) Relative expression of genes (KLF2 overexpression vector

vs. Empty vector) with an overlapping KLF2 motif (k) Differential surface expression of the LFA-

1 subunits CD11a and CD18 in indicated CD8+ T cell subsets from dLN, stratified by canonical

vs. disrupted dLN phenotype (left) and in primary human T cells transduced with empty vector

or KLF2 overexpression vector.

Figure 6. **Cytokine driven dLN types are not fixed.** (a) Patient timeline with tumor and LN

collection pre- and post-neoadjuvant chemotherapy treatment. (b) UMAP projections of myeloid

cells (left) and CD8+ T cells (right) for patient GC028 across both pre-treatment and post-

treatment conditions, colored by either cell subset or distribution in tumors and dLN. (c) Heat-

encoded density embeddings of myeloid and CD8+ T cell compartments prior to and following

neoadjuvant treatment, stratified by timepoint and anatomical site. (d) Alluvial plots showing

CD8+ T cells with shared TCRs across dLN and tumor, both pre- and post-treatment (e) KLF2

expression in CD8+ T cells within dLN of patient GC028, pre- and post-treatment.

Extended Figure Legends

Extended Figure 1. **Marker genes for major CD45+ cell compartments.** The expression of

marker genes in each major CD45+ cell compartment (a), CD8+ T cells (b), CD4+ T cells (c),

cycling cells in CD4+ and CD8+ T cells (d) and myeloid cells (e). (f) Fraction of cells in the major

CD45+ cell compartments within tumors and dLN as indicated.

Extended Figure 2. **Single cell characterization of immune cells from normal tissue and**

metastatic sites in GC. UMAPs depicting (a) broad CD45+ cell subsets across tissue sites (left),

density heatmap of CD45+ cells from distant normal tissue (center) and metastases (right), (b)

CD8+ T cell subsets across tissue sites (left), density heatmap of CD8+ cells from distant normal

tissue (center) and metastases (right), (c) CD4+ T cell subsets across tissues (left), density

heatmap of CD4+ cells from distant normal tissue (center) and metastases (right), and (d)

myeloid cell subsets across tissue sites (left), density heatmap of myeloid cells from distant

normal tissue (center) and tumors (right).

Extended Figure 3. **Patient stratification by distinct cytokine driven immune networks in**

tumors and their draining lymph nodes. (a) Posterior mean for latent sample by factor (left),

cell-type by factor (center), and cytokine by factor (right) tensor slices. For visualization purposes,

the cytokine by factor slide heatmap includes only cytokines with an average factor activation

score of more than 0.2. (b) UMAPs showing the fraction of CD8+ T cell type content and paired

CD8+ T cell density (left) and fraction of CD4 and myeloid cell subsets in each cytokine

interaction-driven immune network cluster. (c) Oncoplot of tumor mutation⬜⬜, separated by either

dLN or tumor subtype.

Extended Figure 4. **CD8+ T cell expansion.** (a) CD8+ T cell clonotype size distribution in dLN

and Tumor (left) and cell phenotypes for expanded TCR clonotypes (right). (b) Average distance

of naïve like T cells from non-naïve T cell phenotypes indicating closer proximity of RA-RO+

naïve T cells to early activated T cells (**Methods**, section *Distance between T Cell Phenotypes*).

(c) Difference in mean cell fractions of CD8+ T cell subsets between disrupted dLN and

canonical dLN following recategorization of naïve-like CD8+ T cells (RA$^+$RO$^-$ naïve and RA$^-$RO$^-$

naïve into 'naïve merged', RA$^-$RO$^+$ naïve and early activated into 'early activated merged').

Extended Figure 5. **KLF2 suppresses effector T cell differentiation.** (a) Fraction of CD8+ T

cell phenotypes in expanded TCR clonotypes in each cytokine-driven immune network cluster.

(b) Western blot showing KLF2 and Histone H3 (loading control) protein abundance in primary

activated human T cells transduced with either empty vector control or KLF2 overexpression

vector. (c) Marker genes and their expression in CD8+ T cell phenotypes reported in empty

vector and KLF2 overexpression vector-transduced cells from **Fig. 5d**. (d) Flow cytometry gating

for **Fig. 5i**.

Extended Figure 6. LFA-1 blockade attenuates T cell priming and effector differentiation.

(a) Experimental design to assess the contribution of LFA-1 to DC-mediated T cell priming and

effector differentiation. (b) Flow cytometry scatter plots and gating of live, B220$^-$NK1.1$^-$

TCRb$^+$CD8$^+$ T cells to assess T cell activation (CD69$^+$CD25$^+$), proliferation (CD44$^+$CellTrace

Violet^{low}), effector differentiation (CX3CR1$^-$CXCR6$^+$) and cytokine production (TNF$^+$IFN-g$^+$). (c)

Quantitation of impact of peptide dose and affinity on CD8+ T cell activation, proliferation,

effector differentiation, and cytokine production as described in (b). Each **table entry** represents

the **average** of three independent biological replicates for each peptide affinity and dose.

Extended Figure 7. **Cytokine driven dLN types are not fixed** (a) Fraction of CD8+ T (left) and

myeloid cell type subsets (right) in patient GC0028, stratified by tissue (dLN and tumor) as well

as timepoint (pre- and post-treatment) (b) Abundance of TCR clonotypes that are shared across

time and anatomical sites (left) and their projection onto the patient GC0028-specific CD8+ T

cell UMAP (right) (c) Alluvial plots showing phenotypes of CD8+ T cells with shared TCR pre-

and post-treatment in both dLN (left) and tumor (right) samples. (d) Differentially expressed

genes in the dLN pre-treatment (up) vs post-treatment (down). (e) Expression of CD11a and

CD18 in CD8+ T cells from patient GC0028 dLN pre- and post-treatment.

EXTENDED DATA TABLES

Extended Data Table 1: Individual patient and tumor characteristics

GC#	pT-stage	pN-stage	pM-stage	Combined Tumor Subtype	*H. Pylori* History	Histological Grade	TRG	PDL1 CPS	HER2 Status	Therapy Prior to First Collection	Any Chemotherapy	Any Immunotherapy	Any Trastuzumab
GC001	ypT3	ypN2	M0	Mixed	Yes	poorly differentiated	3	1	negative	neoadj chemo	Yes	No	No
GC002	ypT3	ypN0	M0	Intestinal	No	moderately differentiated	3	50	negative	neoadj chemo	Yes	No	No
GC003	ypT3	ypN1	M0	Mixed	No	moderately to poorly differentiated		15	positive	No	Yes	Yes	No
GC004	pT1b	pN2	M0	Mixed	No	moderately differentiated		3	negative	No	Yes	No	No
GC005	Tx	Nx	M1	Diffuse	No	poorly differentiated		2	negative	No	Yes	Yes	No
GC006	Tx	Nx	M1	NA	No	poorly differentiated		5	NP	No	Yes	Yes	No
GC007	ypT0	ypN0	M0	NA	No	moderately differentiated		2	negative	No	Yes	Yes	No
GC008	Tx	Nx	M1	Diffuse	No	poorly differentiated		2	negative	chemo	Yes	Yes	No

GC009	ypT1a	ypN0	M0	Intestinal	No	moderately differentiated	2	10	negative	neoadj chemo	Yes	No	No
GC010	ypT3	ypN3	M0	Mixed	Yes	poorly differentiated		2	positive	neoadj chemo	Yes	No	No
GC011	ypT3	ypN0	M0	MSI-H	No	moderately differentiated	2	15	negative	neoadj chemo	Yes	No	No
GC012	Tx	Nx	M1	Diffuse	Yes	poorly differentiated		1	negative	neoadj chemo, adj chemo	Yes	No	No
GC013	Tx	Nx	M1	NA	No	poorly differentiated		20	negative	No	Yes	Yes	No
GC014	ypT3	ypN0	M0	Intestinal	No	poorly differentiated	3	0	negative	neoadj chemo	Yes	No	No
GC015	ypT4a	ypN3	M0	MSI-H	No	poorly differentiated		0	negative	No	Yes	No	No
GC016	ypT3	ypN0	M0	Mixed	Yes	poorly differentiated	2	0	negative	neoadj chemo	Yes	No	No
GC017	pT3	pN1	M0	Diffuse	No	poorly differentiated		1	negative	No	No	No	No
GC018	Tx	Nx	M1	MSI-H	No	poorly differentiated		25	negative	immuno	No	Yes	No
GC019	Tx	Nx	M1	NA	No	poorly differentiated		0	negative	No	Yes	No	No
GC020	Tx	Nx	M0	MSI-H	No	moderately to poorly differentiated		15	negative	No	Yes	Yes	No

GC021	Tx	Nx	M1	Intestinal	Yes	NA		0	positive	No	Yes	Yes	Yes
GC022	pT1b	pN0	M0	Intestinal	No	moderately differentiated		5	negative	No	No	No	No
GC023	Tx	Nx	M1	Diffuse	No	poorly differentiated		1	negative	No	Yes	Yes	No
GC024	pT1b	pN1	M0	Intestinal	Yes	moderately differentiated		35	negative	No	No	No	No
GC025	Tx	Nx	M1	NA	No	poorly differentiated		1	negative	No	Yes	No	No
GC026	pT1b	pN0	M0	NA	No	moderately differentiated		0	negative	No	Yes	No	No
GC027	ypT4b	ypN3	M0	EBV	No	poorly differentiated	3	15	negative	neoadj chemo	Yes	Yes	No
GC028	ypT3	ypN1	M1	Intestinal	Yes	NA		0	negative	No	Yes	No	No
GC029	pT2	pN0	M0	Mixed	No	poorly differentiated		1	negative	No	No	No	No
GC030	pT2	pN1	M0	MSI-H	Yes	poorly differentiated		1	negative	No	No	No	No
GC031	pT1a	pN0	M0	Intestinal	No	moderately differentiated		2	negative	No	No	No	No
GC032	Tx	Nx	M1	Diffuse	No	poorly differentiated		0	negative	No	Yes	Yes	No
GC033	ypT4a	ypN3	M0	Diffuse	No	poorly differentiated	3	1	negative	neoadj chemo	Yes	Yes	No

ID	T	N	M	Subtype		Differentiation				Treatment				
GC034	pT4a	pN2	M0	Mixed	Yes	poorly differentiated		0	negative	No	Yes	No	No
GC035	pT4a	pN0	M0	Diffuse	Yes	poorly differentiated		0	negative	No	Yes	No	No
GC036	ypT2	ypN0	M0	EBV	No	poorly differentiated	2	100	negative	neoadj chemo	Yes	No	No
GC037	ypT1b	ypN0	M0	Intestinal	No	moderately differentiated	1	1	positive	neoadj chemo	Yes	No	No
GC038	pT1a	pN0	M0	Intestinal	No	well differentiated		0	NP	No	No	No	No
GC039	ypT3	ypN0	M0	NA	No	poorly differentiated	2	0	negative	neoadj chemo	Yes	No	No
GC040	ypT1b	ypN0	M0	Intestinal	Yes	moderately to poorly differentiated	3	2	negative	neoadj chemo	Yes	No	No
GC041	pT1a	pN0	M0	Diffuse	Yes	poorly differentiated		0	negative	No	No	No	No
GC042	Tx	Nx	M0	MSI-H	No	NA		20	negative	No	No	Yes	No
GC043	Tx	Nx	M1	NA	No	moderately differentiated		30	negative	No	No	No	No
GC044	pT1b	pN0	M0	MSI-H	No	moderately to poorly differentiated		5	negative	No	No	No	No
GC045	pT3	pN0	M0	Diffuse	No	poorly differentiated		0	negative	No	Yes	No	No

GC046	Tx	Nx	M1	NA	No	moderately to poorly differentiated		0	negative	neoadj chemo	Yes	No	No
GC047	ypT3	ypN2	M0	NA	No	NA		1	negative	No	Yes	Yes	No
GC048	Tx	Nx	M1	NA	No	poorly differentiated		10	negative	chemo+immuno	Yes	Yes	No
GC049	Tx	Nx	M0	MSI-H	No	poorly differentiated		80	negative	No	Yes	Yes	No
GC050	pT2	pN0	M0	MSI-H	No	moderately differentiated		2	negative	No	Yes	Yes	No
GC051	pT1b	pN2	M0	Intestinal	No	poorly differentiated		5	negative	No	Yes	No	No
GC052	ypT3	ypN3	M0	Intestinal	Yes	NA		23	negative	No	Yes	No	No
GC053	pT4a	pN2	M0	Diffuse	Yes	poorly differentiated		0	negative	No	Yes	No	No
GC054	ypT2	ypN0	M0	Diffuse	No	poorly differentiated	2	10	negative	neoadj chemo	Yes	No	No
GC055	ypT4a	ypN3	M1	Diffuse	No	poorly differentiated		1	negative	No	Yes	No	No
GC056	ypT3	ypN1	M1	Intestinal	No	NA		10	negative	No	Yes	Yes	No
GC057	Tx	Nx	M0	MSI-H	No	moderately differentiated		0	negative	No	No	Yes	No
GC058	Tx	Nx	M1	NA	No	poorly differentiated		15	negative	No	No	No	No

GC059	ypT3	ypN0	M0	EBV	Yes	moderately to poorly differentiated		30	negative	No	Yes	Yes	No
GC060	ypT4b	ypN1	M0	Intestinal	No	NA		0	positive	neoadj chemo	Yes	No	Yes
GC061	Tx	Nx	M0	MSI-H	Yes	poorly differentiated		90	negative	No	No		No
GC062	ypT1a	ypN2	M0	Mixed	No	poorly differentiated		2	negative	No	Yes	No	No
GC063	ypT1b	ypN0	M0	Intestinal	Yes	moderately to poorly differentiated		70	negative	No	Yes	Yes	No
GC064	Tx	Nx	M0	MSI-H	No	moderately differentiated		30	negative	No	No		No

512 *Tx, Nx* pathological staging unknown; *NA* not available; *MSI-H* microsatellite instability high; *EBV*

513 Epstein-Barr Virus associated; *TRG* tumor regression grade; *NP* not performed; *neoadj chemo*

514 neoadjuvant chemotherapy; *chemo* chemotherapy; *immuno* immunotherapy

515 **Extended Data Table 2:** Patient demographics

Age	65.1 (11.0 - 90.0)
Stage	
Early	10 (15.6%)
Locally advanced	36 (56.2%)
Metastatic	18 (28.1%)
Gender	
Male	39 (60.9%)

Female	25 (39.1%)
Race	
White	44 (68.8%)
Asian	11 (17.2%)
Black	6 (9.4%)
NA	3 (4.7%)
Combined tumor subtype	
Intestinal	16 (25.0%)
Diffuse	13 (20.3%)
MSI-H	12 (18.8%)
Mixed	8 (12.5%)
EBV	3 (4.7%)
NA	12 (18.8%)
H. pylori **history**	
No	47 (73.4%)
Yes	17 (26.6%)
T-stage	
ypT4b	2 (3.1%)
ypT4a	3 (4.7%)
ypT3	3 (4.7%)
ypT3	10 (15.6%)
ypT2	2 (3.1%)
ypT1b	3 (4.7%)
ypT1a	2 (3.1%)

ypT0	1 (1.6%)
pT4a	3 (4.7%)
pT3	2 (3.1%)
pT2	3 (4.7%)
pT1b	6 (9.4%)
pT1a	3 (4.7%)
NA	21 (32.8%)
N-stage	
ypN3	6 (9.4%)
ypN2	3 (4.7%)
ypN1	4 (6.2%)
ypN0	13 (20.3%)
pN2	4 (6.2%)
pN1	3 (4.7%)
pN0	10 (15.6%)
NA	21 (32.8%)
M-stage	
M1	18 (28.1%)
M0	42 (65.6%)
NA	4 (6.2%)
HER2 status	
Negative	57 (89.1%)
Positive	5 (7.8%)
NA	2 (3.1%)

PD-L1 CPS	
<1	18 (28.1%)
1-4	20 (31.2%)
5-9	4 (6.2%)
>=10	22 (34.4%)
Any Chemotherapy	
Yes	48 (75.0%)
No	16 (25.0%)
Any Immunotherapy	
No	41 (64.1%)
Yes	22 (34.4%)
NA	1 (1.6%)
Histological grade	
Well differentiated	1 (1.6%)
Moderately differentiated	17 (26.6%)
Moderately to poorly differentiated	7 (10.9%)
Poorly differentiated	38 (59.4%)
NA	1 (1.6%)

516 *NA* not available; *MSI-H* microsatellite instability high; *EBV* Epstein-Barr Virus associated

517 **Extended Data Table 3**: Data subsets and cell counts

Object	Cell count	Sample count	Gene count
Anndata object: all cell types	296,231	192	36601

Anndata QCed, cells with <200 genes and samples with <100 cells removed; doublets removed, cells with >40% mitochondrial genes removed). 13 samples removed due to failing QC/insufficient cell count.	251,167	179	36601
Subset to CD45+ cells. Cells with >10% mitochondrial genes removed. Cells double positive and double negative for CD4/CD8 removed.	152,981	179	36601

518 *Anndata* structed data format used in Scanpy eco-system; *QCed* after quality control

519 **Extended Data Table 4:** Top 50 DEGs for cell type cluster annotation for CD45+, CD8, CD4

520 and Myeloid subsets

521 (see attached file)

522 **Extended Data Table 5**: Patients with dLN and tumor IREA types

GC Patient ID	Collection	Lymph Node cytokine group	Tumor cytokine group
GC001		NaN	Tumor Immune Silent
GC002		LN Canonical	NaN
GC003	First collection	NaN	Tumor Inflamed Myeloid
GC003	Second collection	NaN	Tumor T Cell Engaged
GC004		NaN	NaN
GC005		NaN	Tumor T Cell Engaged
GC006		NaN	Tumor T Cell Engaged
GC007		NaN	Tumor Inflamed Myeloid
GC008		NaN	Tumor Inflamed Myeloid
GC009		LN Disrupted	Tumor Inflamed Myeloid
GC010		LN Disrupted	Tumor Immune Silent
GC011		LN Canonical	Tumor Inflamed Myeloid
GC012		NaN	Tumor T Cell Engaged
GC013	First collection	NaN	Tumor T Cell Engaged
GC013	Second collection	NaN	Tumor T Cell Engaged
GC014		LN Canonical	Tumor T Cell Engaged
GC015	First collection	NaN	Tumor T Cell Engaged
GC015	Second collection	LN Canonical	Tumor T Cell Engaged
GC016		LN Canonical	Tumor Immune Silent
GC017		NaN	Tumor T Cell Engaged
GC019		NaN	Tumor Immune Silent

GC020		NaN	Tumor Immune Silent
GC021	First collection	NaN	Tumor T Cell Engaged
GC021	Second collection	NaN	Tumor T Cell Engaged
GC022		LN Disrupted	Tumor Inflamed Myeloid
GC023		LN Disrupted	Tumor Immune Silent
GC024		LN Disrupted	Tumor Inflamed Myeloid
GC025		NaN	Tumor Inflamed Myeloid
GC026		NaN	Tumor T Cell Engaged
GC027		LN Canonical	Tumor Inflamed Myeloid
GC028	First collection	LN Disrupted	Tumor T Cell Engaged
GC028	Second collection	LN Canonical	Tumor T Cell Engaged
GC029		NaN	Tumor Inflamed Myeloid
GC030		LN Disrupted	Tumor Immune Silent
GC031		NaN	Tumor Immune Silent
GC032		NaN	Tumor T Cell Engaged
GC033		LN Disrupted	Tumor Inflamed Myeloid
GC034		LN Disrupted	Tumor Inflamed Myeloid
GC035		NaN	Tumor Immune Silent
GC036		LN Disrupted	Tumor T Cell Engaged
GC037		NaN	Tumor Immune Silent
GC038		LN Disrupted	Tumor T Cell Engaged
GC039		LN Disrupted	Tumor Immune Silent
GC040		LN Canonical	Tumor T Cell Engaged
GC041		LN Disrupted	Tumor Immune Silent
GC042		NaN	Tumor Immune Silent
GC043		NaN	Tumor Immune Silent
GC044		NaN	NaN
GC045		LN Disrupted	Tumor Immune Silent
GC046		NaN	Tumor Immune Silent
GC047	First collection	NaN	Tumor T Cell Engaged
GC047	Second collection	LN Canonical	NaN
GC048		NaN	Tumor T Cell Engaged
GC049		NaN	Tumor T Cell Engaged
GC050		LN Disrupted	Tumor Inflamed Myeloid
GC051		LN Canonical	Tumor T Cell Engaged
GC052	First collection	NaN	Tumor T Cell Engaged
GC052	Second collection	LN Canonical	Tumor T Cell Engaged

GC053		LN Disrupted	Tumor T Cell Engaged
GC054		LN Disrupted	Tumor Immune Silent
GC055	First collection	NaN	Tumor Immune Silent
GC055	Second collection	LN Disrupted	Tumor T Cell Engaged
GC056	First collection	NaN	Tumor Immune Silent
GC056	Second collection	LN Disrupted	Tumor Immune Silent
GC057		NaN	Tumor T Cell Engaged
GC058		LN Canonical	Tumor T Cell Engaged
GC059	First collection	NaN	Tumor T Cell Engaged
GC059	Second collection	LN Canonical	Tumor T Cell Engaged
GC060		LN Canonical	Tumor T Cell Engaged
GC061		NaN	Tumor T Cell Engaged
GC062	First collection	NaN	Tumor T Cell Engaged
GC062	Second collection	LN Disrupted	NaN
GC063	First collection	NaN	Tumor T Cell Engaged
GC063	Second collection	LN Canonical	Tumor Inflamed Myeloid
GC064		NaN	Tumor T Cell Engaged

LN lymph node; *NaN* IREA group not able to be calculated

END NOTES

Acknowledgements

This research was supported in part by the institution's Cancer Center Support Grant (P30

CA008748) from the NIH, and the Cycle for Survival-Equinox Grant (MA, EES, VES, SAV). SAV

was supported by a Damon Runyon Clinical Investigator Award. This work was additionally

supported by research funding from Bristol Myers Squibb. Figure 1a was made using

Biorender.com. We thank the MSKCC Single Cell Analysis and Innovation Lab (SAIL) for their

thoughtful insights and assistance and Anthony Verdone for assistance with data management.

Author Contributions

EES, SS, SS, MA, HL, VES, SPS, and SAV conceived the project and designed all experiments.

SS, MA, HD, VES and EES performed all sample acquisition. EES and SS assembled all

metadata and reviewed clinical characteristics. YHL, SS and EES performed all tissue

processing. YHL and EES performed all spectral cytometry assays. YHL and EES performed all laboratory experiments. SS, HL, MZ, and NC performed all computational and statistical analyses. EES, SS, HL, SPS, VES and SAV wrote the manuscript; all authors reviewed the manuscript.

Competing Interests Statement

VES has received speaking honoraria from Merck Pharmaceuticals and Astra Zeneca. SAV is an advisor for Generate Bio and for the Institute for Follicular Lymphoma Innovation (I-FLI), previously served as a consultant for Koch Disruptive Technologies and has received research funding from Bristol Meyers Squibb. SPS reports grants/contracts from AstraZeneca and Bristol Meyers Squibb and a leadership role in the Genome Canada Scientific Advisory Council. ML ML is supported by a grant from the NIDDK of the NIH (K08 DK125876). No other authors have relationships with outside entities to disclose.

Data Availability

Supplementary Information is available for this paper.

Data is available at ***.

Correspondence and requests for materials should be addressed to shahs3@mskcc.org, strongv@mskcc.org, or vardhans@mskcc.org.

Reprints and permissions information is available at www.nature.com/reprints.

REFERENCES

SEER*Explorer: An interactive website for SEER cancer statistics [Internet]. <https://seer.cancer.gov/statistics-network/explorer/> (2025 Apr 16).

Schistosomes, liver flukes and Helicobacter pylori. *IARC Monogr Eval Carcinog Risks Hum* **61**, 1-241 (1994).

Fu, K. *et al.* Streptococcus anginosus promotes gastric inflammation, atrophy, and tumorigenesis in mice. *Cell* **187**, 882-896 e817 (2024). https://doi.org/10.1016/j.cell.2024.01.004

Norcross, J. W., Monroe, S. E. & Griffin, B. G. The development of gastric carcinoma in pernicious anemia. *Ann Intern Med* **37**, 338-343 (1952). https://doi.org/10.7326/0003-4819-37-2-338

Sasaki, H. *et al.* Presence of Streptococcus anginosus DNA in esophageal cancer, dysplasia of esophagus, and gastric cancer. *Cancer Res* **58**, 2991-2995 (1998).

Zamcheck, N., Grable, E., Ley, A. & Norman, L. Occurrence of gastric cancer among patients with pernicious anemia at the Boston City Hospital. *N Engl J Med* **252**, 1103-1110 (1955). https://doi.org/10.1056/NEJM195506302522601

Rooney, M. S., Shukla, S. A., Wu, C. J., Getz, G. & Hacohen, N. Molecular and genetic properties of tumors associated with local immune cytolytic activity. *Cell* **160**, 48-61 (2015). https://doi.org/10.1016/j.cell.2014.12.033

Abate, M. *et al.* Host Tissue Factors Predict Immune Surveillance and Therapeutic Outcomes in Gastric Cancer. *Cancer Immunol Res* **13**, 591-601 (2025). https://doi.org/10.1158/2326-6066.CIR-23-0563

Stoeckius, M. *et al.* Simultaneous epitope and transcriptome measurement in single cells. *Nat Methods* **14**, 865-868 (2017). https://doi.org/10.1038/nmeth.4380

Lauren, P. The Two Histological Main Types of Gastric Carcinoma: Diffuse and So-Called Intestinal-Type Carcinoma. An Attempt at a Histo-Clinical Classification. *Acta Pathol Microbiol Scand* **64**, 31-49 (1965). https://doi.org/10.1111/apm.1965.64.1.31

Cancer Genome Atlas Research, N. Comprehensive molecular characterization of gastric adenocarcinoma. *Nature* **513**, 202-209 (2014). https://doi.org/10.1038/nature13480

Bellmunt, J. *et al.* Pembrolizumab as Second-Line Therapy for Advanced Urothelial Carcinoma. *N Engl J Med* **376**, 1015-1026 (2017). https://doi.org/10.1056/NEJMoa1613683

Pai, J. A. & Satpathy, A. T. High-throughput and single-cell T cell receptor sequencing technologies. *Nat Methods* **18**, 881-892 (2021). https://doi.org/10.1038/s41592-021-01201-8

Cui, A. *et al.* Dictionary of immune responses to cytokines at single-cell resolution. *Nature* **625**, 377-384 (2024). https://doi.org/10.1038/s41586-023-06816-9

Jin, S. *et al.* Inference and analysis of cell-cell communication using CellChat. *Nat Commun* **12**, 1088 (2021). https://doi.org/10.1038/s41467-021-21246-9

Jin, S., Plikus, M. V. & Nie, Q. CellChat for systematic analysis of cell-cell communication from single-cell transcriptomics. *Nat Protoc* **20**, 180-219 (2025). https://doi.org/10.1038/s41596-024-01045-4

Nakauchi, M. *et al.* Outcomes of Neoadjuvant Chemotherapy for Clinical Stages 2 and 3 Gastric Cancer Patients: Analysis of Timing and Site of Recurrence. *Ann Surg Oncol* **28**, 4829-4838 (2021). https://doi.org/10.1245/s10434-021-09624-5

18 Akbar, A. N., Terry, L., Timms, A., Beverley, P. C. & Janossy, G. Loss of CD45R and gain
of UCHL1 reactivity is a feature of primed T cells. *J Immunol* **140**, 2171-2178 (1988).
19 Serra, H. M., Krowka, J. F., Ledbetter, J. A. & Pilarski, L. M. Loss of CD45R (Lp220)
represents a post-thymic T cell differentiation event. *J Immunol* **140**, 1435-1441 (1988).
20 Tedder, T. F., Penta, A. C., Levine, H. B. & Freedman, A. S. Expression of the human
leukocyte adhesion molecule, LAM1. Identity with the TQ1 and Leu-8 differentiation
antigens. *J Immunol* **144**, 532-540 (1990).
21 Telerman, A. *et al.* Internalization of human T lymphocyte receptors. *Eur J Immunol* **17**,
991-997 (1987). https://doi.org/10.1002/eji.1830170715
22 Miyawaki, T. *et al.* Differential expression of apoptosis-related Fas antigen on lymphocyte
subpopulations in human peripheral blood. *J Immunol* **149**, 3753-3758 (1992).
23 Kuo, C. T., Veselits, M. L. & Leiden, J. M. LKLF: A transcriptional regulator of single-
positive T cell quiescence and survival. *Science* **277**, 1986-1990 (1997).
https://doi.org/10.1126/science.277.5334.1986
24 Mimitou, E. P. *et al.* Scalable, multimodal profiling of chromatin accessibility, gene
expression and protein levels in single cells. *Nat Biotechnol* **39**, 1246-1258 (2021).
https://doi.org/10.1038/s41587-021-00927-2
25 Schober, S. L. *et al.* Expression of the transcription factor lung Kruppel-like factor is
regulated by cytokines and correlates with survival of memory T cells in vitro and in vivo.
J Immunol **163**, 3662-3667 (1999).
26 Schep, A. N., Wu, B., Buenrostro, J. D. & Greenleaf, W. J. chromVAR: inferring
transcription-factor-associated accessibility from single-cell epigenomic data. *Nat
Methods* **14**, 975-978 (2017). https://doi.org/10.1038/nmeth.4401
27 Liu, J. *et al.* CTCF mediates CD8+ effector differentiation through dynamic redistribution
and genomic reorganization. *J Exp Med* **220** (2023).
https://doi.org/10.1084/jem.20221288
28 Raczkowski, F. *et al.* The transcription factor Interferon Regulatory Factor 4 is required for
the generation of protective effector CD8+ T cells. *Proc Natl Acad Sci U S A* **110**, 15019-
15024 (2013). https://doi.org/10.1073/pnas.1309378110
29 Yao, S. *et al.* Interferon regulatory factor 4 sustains CD8(+) T cell expansion and effector
differentiation. *Immunity* **39**, 833-845 (2013).
https://doi.org/10.1016/j.immuni.2013.10.007
30 Man, K. *et al.* Transcription Factor IRF4 Promotes CD8(+) T Cell Exhaustion and Limits
the Development of Memory-like T Cells during Chronic Infection. *Immunity* **47**, 1129-
1141 e1125 (2017). https://doi.org/10.1016/j.immuni.2017.11.021
31 Harberts, A. *et al.* Interferon regulatory factor 4 controls effector functions of CD8(+)
memory T cells. *Proc Natl Acad Sci U S A* **118** (2021).
https://doi.org/10.1073/pnas.2014553118
32 Miyagawa, F. *et al.* Interferon regulatory factor 8 integrates T-cell receptor and cytokine-
signaling pathways and drives effector differentiation of CD8 T cells. *Proc Natl Acad Sci
U S A* **109**, 12123-12128 (2012). https://doi.org/10.1073/pnas.1201453109
33 Miller, I. J. & Bieker, J. J. A novel, erythroid cell-specific murine transcription factor that
binds to the CACCC element and is related to the Kruppel family of nuclear proteins. *Mol
Cell Biol* **13**, 2776-2786 (1993). https://doi.org/10.1128/mcb.13.5.2776-2786.1993
34 Schultess, J., Danielewski, O. & Smolenski, A. P. Rap1GAP2 is a new GTPase-activating
protein of Rap1 expressed in human platelets. *Blood* **105**, 3185-3192 (2005).
https://doi.org/10.1182/blood-2004-09-3605

35   Molina-Ortiz, P. *et al.* Rasa3 controls megakaryocyte Rap1 activation, integrin signaling
and differentiation into proplatelet. *PLoS Genet* **10**, e1004420 (2014).
https://doi.org/10.1371/journal.pgen.1004420
36   Stefanini, L. *et al.* RASA3 is a critical inhibitor of RAP1-dependent platelet activation. *J*
Clin Invest **125**, 1419-1432 (2015). https://doi.org/10.1172/JCI77993
37   Wang, X. *et al.* A phase II study of a modified FOLFOX6 regimen as neoadjuvant
chemotherapy for locally advanced gastric cancer. *Br J Cancer* **114**, 1326-1333 (2016).
https://doi.org/10.1038/bjc.2016.126
38   Man, S. M. Inflammasomes in the gastrointestinal tract: infection, cancer and gut
microbiota homeostasis. *Nat Rev Gastroenterol Hepatol* **15**, 721-737 (2018).
https://doi.org/10.1038/s41575-018-0054-1
39   Hussain, S. P., Hofseth, L. J. & Harris, C. C. Radical causes of cancer. *Nat Rev Cancer* **3**,
276-285 (2003). https://doi.org/10.1038/nrc1046
40   Good, R. A. Relations between immunity and malignancy. *Proc Natl Acad Sci U S A* **69**,
1026-1032 (1972). https://doi.org/10.1073/pnas.69.4.1026
41   Immunodeficiency disease and malignancy. Various immunologic deficiencies of man and
the role of immune processes in the control of malignant disease. *Ann Intern Med* **77**,
605-628 (1972).
42   Fagerberg, E. *et al.* KLF2 maintains lineage fidelity and suppresses CD8 T cell exhaustion
during acute LCMV infection. *Science* **387**, eadn2337 (2025).
https://doi.org/10.1126/science.adn2337
43   Shao, T. Y. *et al.* Kruppel-like factor 2+ CD4 T cells avert microbiota-induced intestinal
inflammation. *Cell Rep* **42**, 113323 (2023). https://doi.org/10.1016/j.celrep.2023.113323
44   Schwandt, T. *et al.* Expression of type I interferon by splenic macrophages suppresses
adaptive immunity during sepsis. *EMBO J* **31**, 201-213 (2012).
https://doi.org/10.1038/emboj.2011.380
45   Zehir, A. *et al.* Mutational landscape of metastatic cancer revealed from prospective
clinical sequencing of 10,000 patients. *Nat Med* **23**, 703-713 (2017).
https://doi.org/10.1038/nm.4333

METHODS

Patient Inclusion

Under Institutional Review Board approval (IRB 20-025) fresh tissue was collected from gastric tumors, distant normal gastric mucosa, draining gastric lymph nodes (dLN), and metastases from patients with gastric adenocarcinoma at Memorial Sloan Kettering Cancer Center (MSK). Patients had either institutional 6-107 and/or 12-245 tissue consent or specific IRB approval for tissue use after patient death. IRB approval was obtained for patients under 18. Tissues were collected from patients undergoing either gastric resection or diagnostic laparoscopy (DL) with esophagogastroduodenoscopy (EGD) as part of treatment or staging/diagnosis. Gastric resections were either open or robotic assisted. Biopsies taken during EGD were removed using cold endoscopic forceps, and biopsies taken during DL were removed using cold laparoscopic graspers. Pathology was reviewed by institutional gastrointestinal pathologists and all tumors were pathologically confirmed to be gastric adenocarcinoma. The 8th edition of the AJCC staging system was used by convention for pathologic TNM staging.[1] Clinical information for patients was collected prospectively in a clinical gastric database. Lauren classification[2] and TCGA classifications[3] were combined to describe patients with intestinal, diffuse, mixed, MSI-H and EBV tumors. MSI-H or EBV designation superseded Lauren classification. Lauren classification was determined by pathology report. MSI-H status was determined by immunohistochemistry (IHC) of MLH1, MSH2, MSH6, and PMS2 as reported by pathology or by tumor mutation burden on institutional genomic testing (MSK IMPACT). EBV positivity was determined by IHC on pathology report. HER2 status was reported as negative, positive, or NA (not available) as a summary of HER2 IHC and fluorescence in situ hybridization (FISH) analysis for equivocal IHC results from pathology reports. PD-L1 scores were reported as CPS from 0-100, with any scores over 100 reduced to 100. For patients who received systemic therapy prior to resection, tumor response grade (TRG) was reported as called in the institutional pathology report from 0-3

according to the NCCN system.[4] *H. pylori* history was determined by a previous positive IHC

test or a documentation of previous *H. pylori* infection or treatment in a patient's chart.

Treatments were grouped as chemotherapy, immunotherapy, trastuzumab, radiation, or other

and were categorized as prior to or following specimen collection. Survival, follow-up, and

recurrence were collected up to January 17, 2025. Outcomes for two patients were not included

due to their enrollment in a clinical trial after specimen collection. Disaggregated data could not

be publicly shared for **7** patients due to IRB restrictions.

Sample Collection and Processing

As mentioned above, tissue was collected in the operating room from clinically indicated gastric

cancer resections or DLs with EGDs. During robotic gastric resections, Indocyanin Green (ICG)

was injected intra-operatively to mark the tumor bed and highlight tumor-draining lymph nodes

as previously described.[5] During open operations, tumor draining lymph nodes were determined

by the operating surgeon according to tumor location and lymph node station. For gastrectomies,

within 3 minutes of resection, 0.5-1.0 cm^3 of tumor, distant normal gastric mucosa (defined as

macroscopically normal appearing to the surgeon and not directly adjacent to the tumor), and

LNs (identified to be draining nodes by ICG or station by operating surgeon) were isolated, cut

into ~1 mm^3 pieces and placed into 15mL conical tubes containing phosphate buffered saline

(PBS). These samples were placed on ice and immediately transported to the laboratory. During

DL with EGD tumor and distant normal samples were collected during EGD with endoscopic

cold forceps, 4 bites of tumor were collected then immediately suspended in PBS, as were 3

bites of distant normal gastric mucosa with the forceps cleaned in-between collections. If

peritoneal or liver metastases were seen on DL with sufficient tissue for pathologic diagnosis, a

biopsy of 0.5 cm^3 of metastasis was collected and immediately transferred to a 15mL conical

tube with PBS. These samples were placed on ice and immediately transported to the laboratory

in the same manner as for gastric resections. All samples were digested using a GentleMACS

Dissociator in 2.2 ml or 4.7mL (based on tissue size) of R-10 solution (RPMI, 10% fetal bovine

serum, 10mM HEPES, 1% penicillin-streptomycin, 1% L-glutamine) and Miltenyi enzymes

(Miltenyi Biotec Tumor Dissociation Kit, Human) at 37°C using GentleMACS program TK3. The

digested tissue was then filtered through a 100-μm cell strainer and washed with cold R-10.

Samples were centrifuged at 500 g x 5 minutes at 4°C and supernatant was aspirated. Pelleted

cells were resuspended in 1mL of ammonium-chloride-potassium (ACK) lysing buffer for 3

minutes at room temperature and then washed with 5 mL of R-10. Samples were centrifuged

again at 500 g x 5 minutes at 4°C and supernatant was aspirated, the cell pellet was

resuspended in 1mL of RPMI plus 2uL of 20mg/mL DNase and incubated at room temperature

for 5 minutes. After incubation 4mL of R10 was added and cells were counted. Cells were spun

again at 500 g x 5 minutes at 4°C and resuspended in 1 million/mL aliquots in BAMBANKER

Serum-Free Cell Freezing Medium (Wako Chemicals) for cryopreservation at -80°C.

Single cell immune profiling

Samples were batched for multiplexed combined 3' single-cell RNA, 5' T-cell receptor (TCR),

and cell-surface epitope assessment (CITE-seq) (10x Genomics.) Selected samples were

thawed from -80°C in a 37°C water bath with gentle agitation, transferred to a 15mL conical tube

and then slowly diluted with 9mL of RPMI. Samples were then centrifuged at 200 rpm for 10 min

at 4°C and the supernatant was aspirated. Each sample was resuspended in 50uL of 1:400

dilution in PBS of Ghost Dye 510 (Tonbo) and transferred to a 96-well V-bottom plate, then

incubated at room temperature in the dark for 10 minutes. Samples were washed with 100uL of

PBS, centrifuged at 700g for 2 minutes at 4°C, and supernatant was discarded. Each sample

was then resuspended in 25 uL of Human TruStain FcX (Biolegand) diluted 1:20 in PBS and

incubated for 10 minutes in the dark on ice. Then an antibody pool of TotalSeq-C Human

Universal Cocktail V1.0 (Biolegend), TotalSeqC anti-human Hashtag Antibodies (Biolegend)

was added, and samples were incubated for 30 minutes on ice. Samples were washed with

100uL of PBS, centrifuged at 700g for 2 minutes at 4°C and supernatant was discarded. Cells

were then resuspended in 200uL of 0.5% bovine serum albumin (BSA) in PBS and transferred

to a flow tube with strainer cap. Cells were flow sorted for live cells and then pooled and counted.

All samples were concentrated to a volume of 50uL and then delivered to the Single Cell

Analytics Innovation Lab (SAIL) for library construction and sequencing.

Single-cell RNA-Seq of FACS-sorted cell suspensions was performed on Chromium instrument

(10x genomics) following the user manual for 5′ v2 chemistry. In brief, FACS-sorted cells were

washed once with PBS containing 2% bovine serum albumin (BSA) and resuspended in PBS

containing 2% BSA to a final concentration of 700–1,200 cells per μL. The viability of cells was

usually above 70%, as confirmed with 0.2% (w/v) Trypan Blue staining (Countess II). Between

10,000 and 30,000 cells were targeted for each sample, where several samples were

multiplexed together on one lane of 10x Chromium and specific extracellular protein epitope

characterized (using Antibody Derived Tag - ADT) according to the cell hashing protocol.[6]

The scRNA-seq and scTCR-seq libraries were prepared using the 10x Single Cell Immune

Profiling Solution Kit, according to the manufacturer's instructions. Briefly, amplified cDNA was

used for both 5′ gene expression library construction and TCR enrichment. For gene expression

library construction, amplified cDNA was fragmented and end-repaired, double-sided size-

selected with SPRIselect beads, PCR-amplified with sample indexing primers (98 °C for 45 s;

14–16 cycles of 98 °C for 20 s, 54 °C for 30 s, 72 °C for 20 s; 72 °C for 1 min), and double-sided

size-selected with SPRIselect beads.

For TCR library construction, TCR transcripts were enriched from 2 µl of amplified cDNA by PCR

(primer sets 1 and 2: 98 °C for 45 s; 10 cycles of 98 °C for 20 s, 67 °C for 30 s, 72 °C for 1 min;

72 °C for 1 min). Following TCR enrichment, enriched PCR product was fragmented and end-

repaired, size-selected with SPRIselect beads, PCR-amplified with sample-indexing primers

(98 °C for 45 s; 9 cycles of 98 °C for 20 s, 54 °C for 30 s, 72 °C for 20 s; 72 °C for 1 min), and

size-selected with SPRIselect beads.

Final libraries (GEX, TCR and HTO/CITE) were sequenced on Illumina NovaSeq S4 or X+

platform (R1 – 26 cycles, i7 – 10 cycles, i5 – 10 cycles, R2 – 90 cycles).

Demultiplexing

HTO sequencing data were aligned to the HTO barcodes, and UMIs were counted for each cell

using CITE-seq-Count. Using a two-component K-means algorithm, we partitioned logged HTO

counts into two distributions: background noise (lower mean) and positive tags (larger mean).

Each droplet was then assigned to its source sample based on tags in the positive signal

component. We classified droplets with multiple assignments as doublets and those with a single

assignment   as   singlets.   This   analysis   was   performed   using   SHARP   v0.1.1

(https://github.com/hisplan/sharp).

Single-Cell Analysis

We used a hierarchical supervised approach to assign phenotypes to cells as follows. After QC

and pre-processing, we used unsupervised graph-based clustering to define the main CD45+

cell types. This yielded 6 separate clusters, namely T/NK, B, Plasma, Mast, Myeloid, and

Plasmacytoid DC (Figure 2a), annotated using known marker genes (Extended Fig. 1,

Supplementary Table 4).

Preprocessing

Raw binary base call (BCL) files were demultiplexed into FASTQ files using Cell Ranger. Gene expression libraries were aligned to the GRCh38 reference genome and quantified using `cellranger count` (v7.1.0). TCR V(D)J libraries were processed with `cellranger vdj` (v6.1.2). Cell barcodes were selected using Cell Ranger's cell-calling algorithm EmptyDrops, and the resulting whitelist was used to filter ADT data. The ADT data was then quantified using the SHARP pipeline, which builds on CITE-seq-Count to count cell barcodes and UMIs. Finally, these matrices—gene expression, ADT, and TCR—were loaded into individual AnnData objects for downstream single-cell analysis.

Cells with fewer than 200 genes and more than 40% mitochondrial content were removed and samples with fewer than 100 cells were removed. Doublet cells were identified using scDblFinder[7] and excluded. Raw counts were normalized by library size and log-transformed with a pseudo-count of 1 using Scanpy.[8] The top 2000 highly variable genes were selected following the removal of ribosomal, mitochondrial and T cell receptor genes using the `highly_variable_genes` function with the `flavor` option set to `seurat` and sample as `batch_key`. Principal component analysis (PCA) was performed retaining the top 50 PCs. `Harmonypy`[9] (v0.0.9) was used to perform batch correction to control for inter-patient variation. The batch corrected embeddings were used for leiden clustering and creating the UMAP[10] projection for visualization. Clusters for the broad cell lineages were annotated using marker genes, derived from Wilcoxon rank-sum tests comparing each cluster to all others. The 8 annotated clusters were: B (MS4A1, CD19, CD79A), T/NK (CD3D, CD3E, NCAM1), epithelial (EPCAM, KRT19, KRT8), endothelial (PECAM1, VWF, CDH5), fibroblasts (COL1A1, FAP, PDGFRA), myeloid (CD14, LYZ, CSF1R), mast (KIT, TPSB2, FCER1A, CPA3, HDC, MS4A2), and plasma (MZB1, SDC1, JCHAIN). Using these annotated cell type clusters, the data were subset to CD45+ cells and cells with more than 10% mitochondrial content were removed. Highly

155 variable gene selection, PCA, batch correction, leiden clustering and UMAP computation were

156 performed as outlined above.

Cell-type annotations

CD45+ cells were annotated using a combination of gene expression and protein marker

intensity. Cells were grouped into major compartments, namely T/NK, myeloid, B, plasma, mast

and plasmacytoid DC cells (Supplementary Table 4). T cells were annotated using cite-seq

protein CD3D, with custom intensity thresholds to filter for CD4+ and CD8+ T cells using CD8A

and CD4 proteins, respectively. For CD4, CD8 and myeloid cells, clustering analysis was

performed and the UMAP re-computed for each subset. Differentially expressed genes between

clusters were evaluated alongside protein expression (e.g. CD45RA and CD45RO) to annotate

13 CD8, 11 CD4 and 7 myeloid subclusters (Supplementary Table 4).

Gaussian Kernel Density Estimates

169 We estimated cell density across the UMAP embedding using Scanpy's

170 `sc.tl.embedding_density` with default parameters, grouping cells by the respective

171 covariate. For each group, we visualized the resulting Gaussian kernel density using

172 `sc.pl.embedding_density`.

Tensor factorization for Cytokine Activation using IREA

We applied cytokine networks to identify intercellular cytokine driven bridging tumors and

tumor-draining lymph nodes. To do so, we computed an enrichment score for each of the 86

cytokines in each cell type per sample. To account for the variation of cell types and cytokines

in our samples, we used a Bayesian Poisson tensor factorization algorithm shown to be robust

to sparse datasets. The algorithm infers sample specific representations in terms of latent

factors. For each anatomical site of origin, we grouped samples based on their factor activation

scores, using a nearest-neighbor graph community detection algorithm.

Concretely, top up-regulated genes in each cell type for each cytokine was acquired from a

recently published cytokine dictionary (hereafter called IREA genes.)[11] For each sample, and

each cytokine, the overlap between the IREA genes and the DE genes in each cell type was

used to compute an overrepresentation analysis and p-values were computed via the

hypergeometric test and corrected for multiple testing using false discovery rate (FDR). An

enrichment score was computed as the negative log10 of the adjusted p-values per (sample,

cell-type, cytokine) tuple. A sample by cytokine score was computed as the sum of all

cytokine-cell type interactions. The sample by cell type by cytokine enrichment score,

computed thus, was used as the input to the tensor factorization algorithm. Let x_{ijc} denote the

enrichment of cytokine c , in cell type j , in sample i , then we assume the following

decomposition: $x_{ijc} \approx \widehat{x_{ijc}} = \sum_{k=1}^{K} A_{ik} B_{jk} C_{ck}$ where $A \in R_+^{N \times K}$, $B \in R_+^{J \times K}$, $C \in R_+^{D \times K}$ are latent

non-negative matrices, representing sample by factor, cell type by factor, and cytokine by

factor scores respectively. To encourage sparsity, we use a Poisson likelihood $x_{ijc} \sim$

Poisson$(x_{ijc}; \widehat{x_{ijc}})$. The target of inference is the posterior distribution on the latent matrices A ,

B , C , that is, $P(A, B, C \mid X, \lambda)$. We use variational inference to infer the latent variables.

Cell type-Cytokine Activation Graph Construction

Let ψ_{ij} denote the activity of cytokine i in cell type j . We use $\psi \in R_+^{J \times C}$ as a weighted adjacency

matrix for a bipartite graph where nodes are the set of cytokines and cell types, and edges

denote the inferred activity level of a source cytokine in a receiver cell type. To compute ψ, we

average over factors to compute sample-specific (Cell-Type, Cytokine) pairings. For each

sample, per Cell Type per Cytokine $\psi_{ij} = \sum_{k=1}^{K} A_{ik} B_{jk} C_{:K} \in R^M$. For each group of samples G ,

we can compute a cell type-cytokine activation matrix as $\psi_{Gj} := \sum_{i \in G} \psi_{ij}$. Similarly, for a set of

cytokines H, we report a summary ψ_{GH} as $\psi_{GH} := \sum_{i \in G, j \in H} \psi_{ij}$.

Cytokine type groupings

We grouped into biologically relevant cytokine families as below:

• Class 2 cytokines: IL-10 and all the interferons (k, g, e, b, a1)

• Common beta-chain cytokines: IL3, IL5, GM-CSF

• Colony stimulating factors: G-CSF, M-CSF, Flt3l

• Hormones: Prolactin, Adiponectin

• IL-1/Alarmin family: TSLP, IL1a, Il1b, IL18, IL33, IL36a

• IL-12 family: IL12, IL23, IL27

• IL-17 family: IL17E

• IL-2 family: IL2, IL7, IL15, IL21

• IL-6 family: IL11, IL31, LIF, Cardiotrophin, Neuropoietin, Oncostatin M

• Th2-like: IL4, IL9, IL13

• TNF family: TNF, TL1A, CD40L, BAFF

Clustering of samples based on IREA scores

We grouped samples that belong to each anatomical site separately. Briefly, given the sample

by factor matrix A, we selected rows corresponding to samples from the dLN (tumor), formed

the nearest neighbor graph and computed the leiden clustering with resolution 0.5. For tumor

samples, this initially yielded 4 clusters. For interpretability, we merged the two that were high in

T cell interactions.

Receptor Ligand activation analysis

We inferred cell-cell interactions using Cellchat,[12] calculating communication probabilities for

each cell type pair and associated significance using permutation testing for disrupted and

canonical lymph node samples separately. We merged these Cellchat objects to perform

comparisons between the two groups and to identify significantly differing ligand-receptor

interaction probabilities. We then filtered these results to cytokine interactions. To further

evaluate the evidence for these predicted interactions, we compared the gene expression for

each significant ligand or receptor highlighted by Cellchat. This was performed by pooling all

cells from each lymph node group and comparing gene expression using a Wilcoxon rank-sum

test with Benjamini–Hochberg correction for multiple comparisons. This strategy allowed us to

include all samples (particularly valuable for myeloid cell types due to the relatively low cell

counts for this cell lineage). Gene expression after pseudobulking by sample was assessed and

only genes that showed a consistent trend between the pseudobulked and pooled data were

retained in our results.

CD8+ T cell Trajectory Analysis

For trajectory analysis, we used Palantir and CellRank. First, we selected CD8+ T cells derived

from dLN and tumor samples based on prior annotations. We then identified the top 2,000 highly

variable genes using the Scanpy function `sc.pp.highly_variable_genes`. Next, we

computed the top 50 principal components using `sc.tl.pca` and performed batch correction

via Harmony integration across individual patient identifiers using

`sc.external.pp.harmony_integrate`. We computed a diffusion map based on these

Harmony-corrected PCA embeddings using `sc.tl.diffmap` to capture global structure and

lineage relationships. We initialized the pseudotime trajectory by selecting a root cell

representative of a biologically meaningful starting point—specifically, a cluster of phenotypically

naïve T cells identified using our annotations. We then used Palantir's

`palantir.core.run_palantir` function to infer cell fate potential and generate pseudotime

values, employing 1,000 waypoints. Finally, we use CellRank's

`cellrank.tl.terminal_states` and `cellrank.tl.lineages` functions, leveraging

Palantir-derived pseudotime and diffusion-based transition probabilities to identify terminal

states and major cellular lineages. We executed CellRank with 1,000 permutations to identify

lineage drivers and associated genes.

Phenotype entropy

We first compute the fraction of CD8+ T cells per sample. We then compute the Shannon entropy

as $H = \sum_{k=1}^{K} p_k \log(p_k)$.

Identifying Naïve-cell subpopulations

We used Gaussian Mixture Models (GMM) to identify subpopulations of naïve cells based on

the protein expression of CD45 isoforms. Briefly, we selected all cells that had a naïve phenotype

based on their transcription profile. We used Bayesian information criterion to select the number

of components for the GMM. We used the from `sklearn.mixture.GaussianMixture` with

a full covariance, `max_iter=10000`, and 100 random restarts.

Distance between T Cell Phenotypes

To compute the distance between different T cell phenotype clusters, we used the geometric

median computed Weiszfeld's algorithm.[13] Geometric median of a set of n points $\{X_i\}$, is defined

as the point y that minimizes the sum of distances from all points in the set, i.e., $y :=$

$\text{argmin}_{y \in R^D} \sum_{i=1}^{n} D(X_i - y)$, where D is the L_2 norm (Euclidean distance). We used the

diffusion map representation of cells as input. We computed the geometric median for each

phenotype group the computed the pairwise distance of geometric medians of each phenotype

cluster from all others.

Tumor vs dLN Transcription Factor Comparison

To find which transcription factors (TF) affect expansion in CD8+ T cells, we compared the

expression of TFs in CD8+ T cells, excluding the naïve phenotype, derived from Tumor to

draining lymph node. We obtained a list of human TFs[14] and ranked their expression using the

Wilcoxon rank-sum test.

KLF2 Correlated Genes

We used regularized linear regression to find genes predictive of KLF2's expression in a sample

specific manner. Gene expression counts were normalized for library-size, log-transformed, and

standardized such that the mean and the variance of each gene are zero and one respectively.

Using linear regression with an L2 penalty (i.e., ridge regression), we estimated a per sample

coefficient vector of per gene association with KLF2 expression. We used the collection of these

coefficients to compute a mean coefficient and Z-score per gene.

KLF2 Gene Cloning

The plasmid PpyCAG-KLF2-IB[15] was a gift from Austin Smith (Addgene plasmid # 60441;

http://n2t.net/addgene:60441; RRID:Addgene 60441), which contained human *KLF2*. PCR

amplification of *KLF2* was performed, using primers (in table below) and Q5® High-Fidelity DNA

Polymerase (New England Biolabs) according to manufacturer's instructions. A restriction digest

was performed of both source plasmid PpyCAG-KLF2-IB and destination plasmid pTRIP-SFFV-

mTagBFP-P2A (Addgene 102585), using BamHI-HF (New England Biolabs) and SalI-HF (New

England Biolabs) with rCutSmart™ Buffer (New England Biolabs) according to manufacturer's

instructions, gel isolated and purified. Construct assembly was then performed using Gibson

Assembly® Master Mix (New England Biolabs) according to manufacturer's instructions. The

constructed plasmid was then transformed into NEB 5-alpha competent *E. coli* cells (New

England Biolabs) using 1uL of plasmid DNA. Mini-prep using Qiagen Mini Kit (Qiagen) was

performed for 3 individual clones per manufacturer's instructions, and then Sanger sequencing

was used to confirm the appropriate sequence for clone 2 which was subsequently used for all

experiments.

Cloning primer sequences

Primer Experiment	Primer Name	Primer Sequence
KLF2 Overexpression Plasmid Primers	NL65_KLF2.F	gagaaccctggacctgcgatggcgctgagtgaacccatcc
	NL65_KLF2.R	cgaggtcgacactagtgctacatgtgccgtttcatgtgc
KLF2 Plasmid Sequencing Primer	SFFV-mTagBFP-F	caaacgccaagaccacatatag

KLF2 Overexpression

Lentivirus was produced using 293T cells as follows: briefly, 293T cells were transiently co-

transfected with viral packaging (pCG-gag-pol), and envelope (pCG-VSVg) constructs along with

the expression vector (pTRIP-SFFV-mTagBFP-P2A-Empty or pTRIP-SFFV-mTagBFP-P2A-

KLF2) using 100 mg/mL polyethylenimine (Polysciences Inc). Virus-containing supernatant was

collected beginning 48 hours after transfection and filtered with a 0.45µm syringe, then aliquoted

and stored at -80°C until use. Primary human T-cell activation was performed as follows: Buffy

coats containing PBMCs were obtained from New York Blood Center. PBMCs were isolated

using Lymphoprep density gradient-based centrifugation. Subsequently, T cells were isolated

using a Dynabeads Untouched Human T Cell kit according to manufacturer's instructions. T cells

were activated in T cell media (RPMI-1640 supplemented with 10% heat-inactivated FBS, 2 mM

L-glutamine, and 100 U/mL Penicillin-Streptomycin) supplemented with recombinant IL-2

(Peprotech, 5 ng/mL). Cells were plated onto non-tissue culture-treated 6-well plates that had

been coated with anti-CD3 (Biolegend, 5 µg/mL) and anti-CD28 antibody (Biolegend, 2 µg/mL)

in PBS. 24 hours after activation, 100x concentrated virus was diluted 1:50 in T cell media (40uL

virus in 160uL T cell media, see above), then 200uL was added dropwise to each well and plates

were shaken gently. 24 hours later, cells were passaged into tissue culture-treated 6-well plates

containing fresh T cell media supplemented with IL-2. 5 days following initial stimulation, cells

were either harvested directly for spectral cytometry-based phenotyping, DOGMA-Seq,

assessment of KLF2 by Western blot, or restimulated to interrogate intracellular cytokine

production as described below.

Protein extraction and Western Blot

Cell lysis was performed using RIPA buffer (Cell Signaling) containing 1% SDS as well as

protease and phosphatase inhibitors. Following lysis, cells were sonicated (Bioruptor) and

centrifuged and supernatants were analyzed for protein concentration using BCA. 40 ug of each

lysate were run on a 4-12% Bis-Tris Gel (Thermo Fisher) and transferred to nitrocellulose

membranes. Membranes were blocked using 5% milk and then blotted with the following

antibodies: anti-KLF2 (see table below) and anti-H3 (see table below) in 3% BSA overnight at

4C on a rocker. Membranes were washed and then incubated with HRP-conjugated secondary

antibodies at 1:5000 dilution for 1 hour at room temperature in 5% milk. Membranes were then

washed and imaged using a Bio-Rad ChemiDoc Imaging System.

Target Name	Host	Company	Catalog #	Dilution
KLF2	Mouse	Novus Biologicals	NBP2-61812-0.025ml	1:1000
Histone H3	Rabbit	Cell Signaling	9715	1:1000
Anti-mouse IgG, HRP-linked Antibody	Horse	Cell Signaling	7076	1:5000
Amersham ECL Rabbit IgG, HRP-linked whole Ab	Donkey	Cytiva	NA934	1:5000

Spectral Cytometry

For phenotyping, cells were incubated with Fc block (Human TruStain FcX, Biolegend, cat.

422302) at a 1:20 dilution for 30 minutes at 4°C. Viability staining was performed for 30 minutes

at 4°C using Ghost Dye 510 Violet (cat. 59863) diluted 1:800 in PBS. Cells were subsequently

washed and stained for surface markers in FACS buffer containing 1X Brilliant Buffer (cat.

566349) for 30 minutes at 4°C, then washed twice and resuspended in 1% BSA in PBS. Data

was acquired using a Cytek Aurora spectral cytometer.

Staining type	Marker	Fluorophore	Catalog #	Clone	Dilution
Phenotype, Surface	PD-1	APC	BioLegend, 329908	EH12.2H7	1:100
	CCR7	BB700	BD Biosciences, 566437	3D12	1:50

CD45RA	BUV395	BD Biosciences, 740298	HI100	1:400
CD8	BUV496	BD Biosciences, 612943	RPA-T8	1:800
CD38	BUV661	BD Biosciences, 612969	HIT2	1:200
HLA-DR	BV786	BD Biosciences, 564041	G46-6	1:200
CD4	BV570	BioLegend, 300534	RPA-T4	1:800
CD28	BUV805	BD Biosciences, 742037	CD28.2 (RUO)	1:100
CD69	BV605	BioLegend, 310937	FN50	1:200
Tim-3	BV650	BioLegend, 345028	F38-2E2	1:50
CD103	PE-594	BioLegend, 350223	Ber-ACT8	1:100
CD39	PE-810	BioLegend, 328245	A1	1:100
CXCR3	A700	BD Biosciences, 561320	1C6/CXCR3	1:100
CXCR6	BV421	BioLegend, 356013	K041E5	1:100

Phenotype, Intracellular	TCF-1	A488	Cell Signaling, 6444S	C63D9	1:100
	TOX	PE	MACS Miltenyi Biotec, 130-120-716	REA473	1: 25
	CTLA4	BV711	BioLegend, 369632	BNI3	1: 200

For interrogation of cytokine production, PMA and ionomycin were added to wells to a final

concentration of 50 ng/mL and 500 ng/mL, respectively. 1 hour after addition of PMA and

ionomycin, Brefeldin A was added at a 1:1000 dilution and cells were incubated for another 3

hours, after which they were harvested and stained with surface antibodies as described above

using the panel below. Fixation was performed using an eBioscience FoxP3

Fixation/Permeabilization kit and stained overnight with intracellular antibodies diluted in

permeabilization buffer overnight at 4C. Data was acquired using a Cytek Aurora spectral

cytometer.

Staining type	Marker	Fluorophore	Catalog #	Clone	Dilution
Restimulation, Surface	PD-1	APC	BioLegend, 329908	EH12.2H7	1: 100
	CCR7	BB700	BD Biosciences, 566437	3D12	1: 50

CD45RA	BUV395	BD Biosciences, 740298	HI100	1: 400
CD8	BUV496	BD Biosciences, 612943	RPA-T8	1: 800
CD38	BUV661	BD Biosciences, 612969	HIT2	1: 200
HLA-DR	BV786	BD Biosciences, 564041	G46-6	1: 200
CD4	BV570	BioLegend, 300534	RPA-T4	1: 800
CD28	BUV805	BD Biosciences, 742037	CD28.2 (RUO)	1: 100
CD69	BV605	BioLegend, 310937	FN50	1: 200
Tim-3	BV650	BioLegend, 345028	F38-2E2	1: 50
CD103	PE-594	BioLegend, 350223	Ber-ACT8	1: 100
CD39	PE-810	BioLegend, 328245	A1	1: 100

Restimulation, Intracellular	GzmB	A700	BD Biosciences, 560213	GB11	1: 1600
	GzmK	FITC	BioLegend, 370508	GM26E7	1: 800
	IL-2	APC-F750	BioLegend, 500351	MQ1-17H12	1: 400
	IFN-g	BV421	BioLegend, 502531	4S.B3	1: 800
	TNF	PE	BioLegend, 502908	MAb11	1: 800

DOGMA-Seq

Primary human T cells were isolated and cultured as described above ('**KLF2 Overexpression**').

On day 5 following initial activation, T cells were collected and stained with a viability dye

followed by surface staining with anti-CD4 (RPA-T4, BioLegend 300534, 1:100), anti-CD8 (RPA-

T8, BD Biosciences 612914, 1:1200), TotalSeq-A Human Universal Cocktail, and TSA-

compatible hashtag oligos for multiplexing. Live BFP+ CD4$^+$ and CD8$^+$ T cells from both KLF2

overexpressing and empty vector conditions were flow sorted and delivered to SAIL for

sequencing. Single Cell Multiome ATAC + Gene Expression was performed with 10x genomics

system using Chromium Next GEM Single Cell Multiome Reagent Kit. We followed User Guide

with the difference that single cell suspension was prepared as described in Mimitou et al

(DOGMA-seq protocol with Digitonin buffer).[16] Around 10,000 cells were targeted for each

sample. Samples were multiplexed together on one lane of 10x Chromium (using Hash Tag

Oligonucleotides - HTO) and specific extracellular protein epitope characterized (using Antibody

Derived Tag - ADT) following previously published protocol.[6] Next, cells were treated with a

digitonin lysis buffer (20 mM Tris-HCl pH 7.4, 150 mM NaCl, 3 mM MgCl2, 0.01% DIG and 2 U

µl−1 of RNase inhibitor) for 5 min on ice, followed by adding 1 ml of chilled DIG wash buffer

(20 mM Tris-HCl pH 7.4, 150 mM NaCl, 3 mM MgCl2 and 1 U µl−1 of RNAse inhibitor) and

inversion before centrifugation at 500g for 5 min at 4 °C. The supernatant was discarded, and

cells were resuspended in DIG wash buffer, followed by counting using trypan blue and DAPI

on a Countess II FL Automated Cell Counter. Then cells were processed according to the

Chromium Next GEM Single Cell Multiome ATAC + Gene Expression user guide. Final libraries

were sequenced on Illumina NovaSeq X+ platform following 10x genomics recommendations.

For sample demultiplexing, HTO sequencing data were aligned to the HTO barcodes, and UMIs

were counted for each cell using CITE-seq-Count. Using a two-component K-means algorithm,

we partitioned logged HTO counts into two distributions: background noise (lower mean) and

positive tags (larger mean). Each droplet was then assigned to its source sample based on tags

in the positive signal component. We classified droplets with multiple assignments as doublets

and those with a single assignment as singlets. This analysis was performed using SHARP

v0.1.1 (https://github.com/hisplan/sharp).

Cell typing of DOGMA-seq

The KLF2 over-expression and empty vector were demultiplexed as above in '**DOGMA-Seq**'.

We processed single-cell RNA-, ATAC- and CITE-seq (DOGMA-seq) data using the Seurat (v.

5.0.3) and Signac packages (v.1.13.0). A total of 9,616 cells were sequenced, 6,487 in the KLF2

over-expression and 3,129 in the empty vector conditions. We loaded RNA and ATAC-seq data

into a single Seurat object. We computed quality control metrics—including nucleosome signal

and transcription start site (TSS) enrichment and filtered cells based on RNA/ATAC count

thresholds, nucleosome signal, and TSS enrichment to exclude low-quality cells. Next, we established a unified peak set by merging and filtering called peaks, removing non-standard chromosomes and blacklist regions. We quantified ATAC counts within these peaks using `FeatureMatrix` and added them to a new chromatin assay in the Seurat object. We then split the CD4+ and CD8+ T cells based on the CD4 and CD8A protein expression. To annotate cell phenotypes, we normalized the RNA expression counts with `SCTransform` and reduced dimensions using `RunPCA`. We processed ATAC data with term frequency-inverse document frequency (`RunTFIDF`) and `RunSVD`. We then integrated RNA and ATAC modalities using `FindMultiModalNeighbors` and `FindClusters` on the resulting joint weighted nearest-neighbor graph to identify clusters. We used `FindAllMarkers` on the SCT assay and separately on the ADT assay to report marker genes and proteins per cluster. We used these markers to annotate the T cell phenotypes.

Motif Accessibility Analysis

We used the `RunChromVAR` function with default parameters from the Signac package to compute transcription factor (motif activities from single-cell chromatin accessibility data. Briefly, it quantifies how accessible each cell is at genomic regions associated with specific TF motifs by comparing observed accessibility to background expectations. ChromVAR generates bias-corrected, normalized activity scores (z-scores) for each motif in each cell, enabling the identification of TFs that are potentially driving differences in chromatin states across cell populations or experimental conditions. With the accessibility scores at hand, we used the `FindMarkers` function to compute the differentially accessible markers per cell type (with parameters `mean.fxn = rowMeans` and `fc.name = "avg_diff"`).

KLF2 Targets

We first extracted KLF2's motif's binding sites that overlapped with any identified peak. We used

the motif MA1515.2.[17] We then filtered for regions that were within 1000bp of a protein coding

gene's promoter to identify putative target genes. Among these genes, we reported those with a

log-fold-change (between the KLF2 over expression versus the empty vector conditions) of

above 1.0 or below –1.0 and a gene expression value of 0.1 in at least one of the conditions.

LFA-1 dependent T cell priming assay

Cell Isolation

OT-I T cells and dendritic cells were isolated as follows: for DC isolation, spleens from C57/Bl6

mice were minced and incubated in digestion buffer (200X DNase, 50X Liberase, RPMI) for 30

minutes at 37 degrees. During this incubation period, spleens from OT-I mice were isolated,

mashed over a 40 µm strainer. Subsequently, spleens were washed with RPMI and treated with

hypotonic ACK lysis buffer to lyse red blood cells. RBC lysis was quenched with 10x excess

PBS and centrifuged. OT-I T cells and DCs were then enriched using T-cell isolation

(Dynabeads™ Untouched™ Mouse T Cells Kit) and DC enrichment kits (Dynabeads™ Mouse

DC Enrichment kit), respectively.   These are both negative selection kits in which cell

suspensions were first incubated with antibodies for 20 minutes at 4 degrees, subsequently

incubated with magnetic beads for 15 minutes with gentle tilting and rotation, and finally placed

on a magnet after which suspensions containing OT-I T cell and dendritic cell-enriched

populations were transferred to a fresh tube for downstream assays.

T cell-DC coculture

After isolation, DCs were resuspended in complete RPMI-1640 media containing 10% FBS, 2

mM L-glutamine and Pen/Strep and incubated at 37 degrees for 1 hour with gentle rotation in

the presence of escalating doses of SIINFEKL (N4), SIIQFEKL (Q4), SIITFEKL (T4), or

SIIVFEKL (V4) ranging from 0.1, 1, 10, and 100 nM. Cells were subsequently washed and

resuspended at 1 million cells/mL.

During this period, OT-I T cells were labeled with CellTrace Violet by resuspending in PBS at 1

million cells/mL, warming to 37 degrees, mixing 1:1 with CellTrace Violet diluted 1:1000 in pre-

warmed PBS and incubating in a 37 degree water bath for 20 minutes. Labeling was quenched

by adding 5x volume of RPMI+++. Labeled T cells were counted, centrifuged and resuspended

in complete T cell media (RPMI+++ containing 50 uM BME and 10 ng/mL IL-2) at 200K/mL, with

either isotype control (rat IgG2a kappa, BioXCell) or anti-LFA-1 (M17/4, BioXCell) at a final

concentration of 5 ug/mL. 500 uL containing 100K T cells were added to each well of a 96 well

Polypropylene Deep Well Plate (Corning) into which 5 uL DCs (5,000 DC total) were added. The

plate was then incubated at 37 degrees for 4 days.

For interrogation of cytokine production, PMA and ionomycin were added to wells to a final

concentration of 50 ng/mL and 500 ng/mL, respectively. 1 hour after addition of PMA and

ionomycin, Brefeldin A was added at a 1:1000 dilution and cells were incubated for another 3

hours, after which they were harvested and stained with surface antibodies as described above

using the panel below.   Fixation was performed using an eBioscience FoxP3

Fixation/Permeabilization kit and stained overnight with intracellular antibodies diluted in

permeabilization buffer overnight at 4C. Data was acquired using a Cytek Aurora spectral

cytometer.

Staining type	Marker	Fluorophore	Catalog #	Clone	Dilution
Restimulation, Surface	B220	BUV496	BD Biosciences 612950	RA3-6B2	1:800
	TCRb	BUV805	BD Biosciences 748405	H57-597	1:100
	NK1.1	BUV615	BD Biosciences 751111	PK136	1:50

	CD4	BUV563	BD Biosciences 612923	GK1.5	1:800
	CD8	BV650	Biolegend 100741	53-6.7	1:800
	CD62L	BV570	Biolegend 104433	ME-14	1:200
	CD44	BUV737	BD Biosciences 612799	IM7	1:400
	PD-1	APC-Cy7	Biolegend 135224	29F.1A12	1:200
	CD39	BUV395	BD Biosciences 567264	Y23-1185	1:800
	CD25	PE	Biolegend 102008	PC61	1:200
	CD127	PE-Cy5	Biolegend 135015	A7R34	1:200
	CD69	PerCP-Cy5.5	Biolegend 104522	H1.2F3	1:200
	CX3CR1	APC	Biolegend 149008	SA011F11	1:400
	CXCR6	PE-Dazzle594	Biolegend 151117	SA051D1	1:200
Restimulation, Intracellular	IL-2	BV711	Biolegend 503837	JES6-5H4	1:800
	TNF	FITC	Biolegend 506304	MP6-XT22	1:800
	IFN-g	PE-Cy7	Biolegend 505826	XMG1.2	1:1600
	Granzyme B	Alexa 700	Biolegend 372222	QA16A02	1:3200

Methods Reference Section:

1   Amin, M., Edge, S., Greene, F. & et al. *AJCC Cancer Staging Manual*. 8th edn, (Springer,
2017).
2   Lauren, P. The Two Histological Main Types of Gastric Carcinoma: Diffuse and So-Called
Intestinal-Type Carcinoma. An Attempt at a Histo-Clinical Classification. *Acta Pathol*
Microbiol Scand **64**, 31-49 (1965). https://doi.org/10.1111/apm.1965.64.1.31
3   Cancer Genome Atlas Research, N. Comprehensive molecular characterization of gastric
adenocarcinoma. *Nature* **513**, 202-209 (2014). https://doi.org/10.1038/nature13480
4   NCCN®.   *NCCN   Guidelines®   Gastric   Cancer   2.2023*,
<https://www.nccn.org/guidelines/guidelines-detail?category=1&id=1434> (2023).
5   Herrera-Almario, G., Patane, M., Sarkaria, I. & Strong, V. E. Initial report of near-infrared
fluorescence imaging as an intraoperative adjunct for lymph node harvesting during

robot-assisted laparoscopic gastrectomy. *Journal of Surgical Oncology* **113**, 768-770 (2016). https://doi.org/10.1002/jso.24226

Stoeckius, M. *et al.* Cell Hashing with barcoded antibodies enables multiplexing and doublet detection for single cell genomics. *Genome Biology* **19** (2018). https://doi.org/10.1186/s13059-018-1603-1

Germain, P.-L., Lun, A., Garcia Meixide, C., Macnair, W. & Robinson, M. D. Doublet identification in single-cell sequencing data using scDblFinder. *F1000Research* **10**, 979 (2022). https://doi.org/10.12688/f1000research.73600.2

Wolf, F. A., Angerer, P. & Theis, F. J. SCANPY: large-scale single-cell gene expression data analysis. *Genome Biology* **19** (2018). https://doi.org/10.1186/s13059-017-1382-0

Korsunsky, I. *et al.* Fast, sensitive and accurate integration of single-cell data with Harmony. *Nature Methods* **16**, 1289-1296 (2019). https://doi.org/10.1038/s41592-019-0619-0

Leland McInnes, J. H., James Melville. UMAP: Uniform Manifold Approximation and Projection for Dimension Reduction.   https://doi.org/10.48550/arxiv.1802.03426

Cui, A. *et al.* Dictionary of immune responses to cytokines at single-cell resolution. *Nature* (2023). https://doi.org/10.1038/s41586-023-06816-9

Jin, S. *et al.* Inference and analysis of cell-cell communication using CellChat. *Nature Communications* **12** (2021). https://doi.org/10.1038/s41467-021-21246-9

Chandrasekaran, R. & Tamir, A. Open questions concerning Weiszfeld's algorithm for the Fermat-Weber location problem. *Mathematical Programming* **44**, 293-295 (1989). https://doi.org/10.1007/bf01587094

Lambert, S. A. *et al.* The Human Transcription Factors. *Cell* **172**, 650-665 (2018). https://doi.org/10.1016/j.cell.2018.01.029

Takashima, Y. *et al.* Resetting Transcription Factor Control Circuitry toward Ground-State Pluripotency in Human. *Cell* **158**, 1254-1269 (2014). https://doi.org/10.1016/j.cell.2014.08.029

Mimitou, E. P. *et al.* Scalable, multimodal profiling of chromatin accessibility, gene expression and protein levels in single cells. *Nature Biotechnology* **39**, 1246-1258 (2021). https://doi.org/10.1038/s41587-021-00927-2

Rauluseviciute, I. *et al.* JASPAR 2024: 20th anniversary of the open-access database of transcription factor binding profiles. *Nucleic Acids Research* **52**, D174-D182 (2024). https://doi.org/10.1093/nar/gkad1059

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

a GC028   Time from diagnosis (days)   **Cytokine-driven Immune Network**
- LN Disrupted
- LN Canonical
- Tumor T-cell Engaged

b Myeloid *n = 228*   CD8 *n = 1,208*

CD8:
- Central memory
- Naïve
- Neat1
- Early activated
- Late activated
- Progenitor Texh
- Effector memory
- HSP
- Temra
- Terminal Texh
- Early Texh
- Effector
- Proliferating

Site:
- Lymph Node
- Tumor

Myeloid:
- Mast
- cDC1
- Migratory/cDC3
- Plasmacytoid DC
- cDC2
- Macrophage C1QC
- Macrophage NLRP3

c LN   Tumor

Myeloid / CD8:
- Pre (Disrupted)
- Post (Canonical)
- Pre (T cell engaged)
- Post (T cell engaged)

d Pre   Post

e KLF2 expression in dLNs (p-value < 1e-10)