Sex-dependent neural plasticity in response to damage

Marina Recatalà-Martinez[1], Manel Bosch[2], Pedro Gaspar[3], Alessandro Mineo[3], Santiago

Rios[1], Irene Miguel-Aliaga[3], Marta Morey[1,4,*]

1.Departament de Genètica, Microbiologia i Estadística, Facultat de Biologia, Universitat

de Barcelona, Barcelona, Spain

2. Unitat de Microscòpia Òptica Avançada, Centres Científics i Tecnològics de la

Universitat de Barcelona (CCiTUB), Barcelona, Spain

3. The Francis Crick Institute, London, United Kingdom.

4. Human Genetics and Disease, Institut de Biomedicina de la Universitat de Barcelona

(IBUB), Barcelona, Spain

*Corresponding author: mmorey@ub.edu

ABSTRACT

Plasticity of intact neural tissue in the vicinity of neural damage serves to restoring

functionality of circuits. Much remains to be learned about the mechanisms regulating

this process and the reported sex differences in recovery outcomes. Here we present the

fly gut and its innervation as simplified model to address these questions. We show that

gut damage caused by ingestion of toxic agents resulted in plasticity of the enteric

neuronal network, manifested as a Reactive Oxygen Species (ROS)-dependent increase

in neural tissue, which was reversible after a recovery period. Interestingly, males did not

display neural plasticity, and masculinization of neurons in females suppressed the

damage-dependent neural growth. Together, these findings position the fly gut as a

system to investigate the cellular, molecular and sex-specific underpinnings of neural

plasticity, with implications for therapeutic advancements in neural circuit recovery.

INTRODUCTION

Neural plasticity encompasses two crucial mechanisms for functional recovery after

injury: axonal regeneration and structural plasticity of intact neural tissue. While

significant research has focused on understanding and overcoming limitations to axonal

regeneration (Mahar and Cavalli, 2018; Tedeschi and Bradke, 2017), modulating the
plasticity of remaining healthy neural tissue offers an alternative approach to restoring
functionality (Gao et al., 2022; Petersen et al., 2022). Indeed, extensive clinical evidence
demonstrates that structural plasticity of intact neural tissue plays a vital role in
functional recovery and circuit rewiring after acute damage such as stroke (Cirillo et al.,
2020; Murphy and Corbett, 2009; Nudo, 2003; Sampaio-Baptista et al., 2018) or multiple
sclerosis episodes (Ksiazek-Winiarek et al., 2015; Prosperini et al., 2015). Interestingly,
stroke recovery outcomes differ between sexes (Turtzo and McCullough, 2008; Yu et al.,
2015), potentially indicating a connection to variations in neural plasticity capacity (Hyer
et al., 2018; Kirby et al., 2024).

Despite these observations, insight into the cellular and molecular mechanisms
regulating the plasticity capacity of adult intact neural tissue is very limited. An obstacle
to advancing our understanding in this area is the intricate biological context and
associated technical challenges. Specifically, the presence of healthy neural tissue in
close proximity to, or even intermingled with, damaged or dying neural tissue
complicates research efforts, making it difficult to isolate and study specific structural
plasticity processes. We developed a simplified paradigm that focuses on the plasticity
of intact neural tissue in response to non-neural tissue damage. Our approach leverages
the fact that organs are innervated by the peripheral nervous system. This allows us to
use the organ as a platform for inducing diverse types of damage, exposing the intact
nervous system that innervates it to different noxious stimuli and analyzing the plasticity
response. This approach offers several advantages: it provides a less complex
environment compared to direct neural injury models; it allows for precise control over
the type and extend of non-neural tissue damage; and it enables the observation of intact
neural tissue responses in a physiologically relevant context.

We used *Drosophila melanogaster*'s digestive tract as a model system. Similar to
mammals the fly gut is structurally and physiologically compartmentalized (Buchon et
al., 2013; Lemaitre and Miguel-Aliaga, 2013; Marianes and Spradling, 2013). The
epithelium contains similar cell types: intestinal stem cells (ISCs), enteroblast
progenitors (EBs) that differentiate into absorptive enterocytes (ECs) and progenitors that

give rise to secretory enteroendocrine (EEs) cells. It is surrounded by muscle, oxygenated

by a vasculature-like respiratory tracheal system and innervated by neurons located in

the central nervous system, the corpora cardiaca (neurosecretory structure), and the

hypocerebral ganglion (HCG, one of the stomatogastric ganglia) (Kuraishi et al., 2015;

Miguel-Aliaga et al., 2018). While most neurites terminate on the visceral muscles, some

appear to reach the underlying epithelium (Cognigni et al., 2011; Cui et al., 2024;

Kenmoku et al., 2016; Petsakou et al., 2023). Contrary to the extensive innervation along

the length of the tract in mammals, innervation in the fly gut is still complex but restricted

to specific regions facilitating its analysis (Kuraishi et al., 2015; Miguel-Aliaga et al.,

2018). Recent studies in mammals and flies show the effects of microbiota, nutrient, sex,

ageing, reproductive stage, and recovery after injury on the morphology and/or function

of gastrointestinal innervation, showcasing its plasticity capacity (Ameku et al., 2020;

Hadjieconomou et al., 2020; Petsakou et al., 2023). Here we analyzed the structural

plasticity capacity of intact neural tissue innervating the anterior midgut in response to

damage.

RESULTS

Intestinal damage induces reversible neural plasticity.

In order to quantify and characterize the innervation in the anterior part of the midgut we

proceeded to select a neural driver and reporter that would specifically label neuronal

processes with native GFP fluorescence. This prevented background signal from

secondary antibodies and provided a high signal to noise ratio which facilitated faithful

neural process quantification as GFP signal. General neuronal markers, such as Elav

(Chen et al., 2016; Holsopple et al., 2022; Titos et al., 2023) or n-Synaptobrevin (Weaver

et al., 2020), also label EEs. However, we used the *GMR51F12-GAL4* driver line from the

Janelia FlyLight collection (Jenett et al., 2012; Min et al., 2021; Olds and Xu, 2014) and a

specific *nSyb.S-GAL4* driver (Weaver et al., 2020), neither of which labels EEs.

Morphological landmarks in whole-mount preparations, combined with phalloidin co-

staining to visualize the surrounding musculature, enabled consistent imaging of the

same digestive tract region and its associated neural innervation (Figure 1A). Using 3D

reconstruction and image analysis we obtained volume quantifications for both the

digestive tube and neural tissue (see Material and Methods). Scaling growth is a prevalent

phenomenon in the peripheral nervous system where sensory and motor neurons must adjust the size of their arborizations according to the area of their target tissues to maintain their functionality (Bentley and Toroian-Raymond, 1981; Bucher and Pflüger, 2000; Lee and Stevens, 2007; Menon et al., 2013). Measuring the volume of the digestive tube enabled us to use statistical analysis methods that accounted for and removed any potential influence of digestive tube size variation in adults (driven by treatment or genetic background) on neural tissue quantification comparisons (see Material and Methods).

To damage the gut we fed flies with DSS. DSS is a polymer that induces colitis in mammals (Yang and Merlin, 2024) and has been used in *Drosophila* to study the cellular mechanisms and molecular pathways that regulate ISC proliferation during regeneration (Jiang et al., 2016; Tian et al., 2022). When flies ingest DSS it crosses the epithelium expanding the basement membrane sheet and altering muscle morphology (Howard et al., 2019). The resulting change in biomechanical forces in the niche is thought to activate ISC division (Howard et al., 2019). DSS feeding also results in trachea sprouting (Perochon et al., 2021), which is necessary for ISC proliferation (Perochon et al., 2021; Tamamouna et al., 2021). Interestingly, basement membrane and muscle defects, as well as trachea remodeling, are reversible after a recovery period (Howard et al., 2019; Perochon et al., 2021; Tamamouna et al., 2021). We wondered whether adult neurons were plastic and could also respond to gut damage.

To analyze the effect of gut damage on neural plasticity we fed adult mated females with DSS for 4 days and compared them to controls fed with the carrier (sucrose). We confirmed the effectiveness of DSS by replicating previously published observations, including a reduction in lifespan (Amcheslavsky et al., 2009) (data not shown) and an increase in the proliferative response of ISCs in the midgut (Amcheslavsky et al., 2009) (see later in Figure 2A). Over 90% of the flies that survived the feeding protocol tested negative for the SMURF assay (Rera et al., 2012), indicating that in the majority of animals (93,3%±SD4.5) that were dissected gut permeability was not compromised. Thus, during DSS feeding, neurons were largely shielded from direct exposure to gut contents that could potentially cause damage. When we dissected the DSS-fed females and compared

them to control flies we observed a significant increase in neural tissue (Figure 1B). To rule out neurogenesis as the source of the new neural tissue, we quantified the number of neural nuclei in the HCG, the primary source of neurons innervating the anterior midgut (Figure 1A). No differences were observed between control and DSS-fed animals (Figure 1C), suggesting that the increase in neural tissue was due to neural growth. This neural growth was not exclusive to DSS damage since we also observed it when feeding the flies with H_2O_2 (Figure 1D).

Tracheal terminal cells (TTCs) are known to sprout in response to DSS and H_2O_2 (Perochon et al., 2021; Tamamouna et al., 2021). Hence, we investigated their response in the anterior midgut, where we observed the increase in neural tissue. Our findings revealed that this region was poorly tracheated, with very few TTCs (Supplementary Figure 1A) and minimal branch coverage (Supplementary Figure 1B). This pattern remained unchanged in DSS-fed flies, indicating that TTCs in this area were unresponsive to gut damage (Supplementary Figure 1A, B). Furthermore, their morphology did not account for the observed extent of neural growth, which far exceeded the coverage provided by TTCs. These findings suggested that in the anterior midgut TTCs did not serve as a physical scaffold for neural tissue growth.

We next explored the stability of this neural plasticity and conducted recovery experiments (Figure 1E). To carry out comparisons within the same cohort of flies, a subset of flies was dissected after the DSS treatment and the rest after the recovery period. While DSS feeding increased the neural tissue (C versus DSS), a 4-day recovery period (DSS + Rec) reverted the amount of neural tissue to levels as those of control flies (C + Rec). Consistent with this observation, a clear reversal of neural growth was observed when comparing flies dissected after the DSS treatment (DSS) and those dissected after the recovery phase (DSS + Rec). No differences were observed between control flies (C versus C + Rec).

In summary, these experiments indicate that the intact neural tissue innervating the digestive tube is plastic and responds to damage with growth. This neural growth is reversible after a recovery period, as the gut returns to homeostasis after injury.

161   Importantly, these observations were validated with two independent neural drivers

162   (*GMR51F12-GAL4,* Figure 1 and *nSyb.S-GAL4,* Supplementary Figure 1C).

Figure 1

Figure 1. Neural plasticity induced by gut damage resolves after recovery. A) Drawing

of the adult gut with two areas of interest highlighted and corresponding z-stack

projected confocal images. The square box region corresponds to the area above the

proventriculus (pv) and the image shows neuronal nuclei (magenta, anti-Elav) and

neuronal membrane (grey, GFP) of the HCG. Autofluorescence of the peritrophic

membrane lining the gut in this region is observed. The rectangular box region

corresponds to the area of the anterior midgut imaged in subsequent analysis and the

images show neuronal processes (grey, GFP) and muscle fibers (grey, Phalloidin). B)

Schematic outlining the feeding protocol and dissection time. Z-stack projected

confocal images of neuronal processes innervating the anterior midgut of control (C) and

DSS-fed (DSS) females and quantification of neural tissue. Red outline in the box plots or

bar graphs indicates female. C) Z-stack projected confocal images of the HCG of control

and DSS-fed females stained with Hoechst (grey, which also accumulates in some

tracheal branches) and showing neuronal nuclei (magenta, anti-Elav). Quantification of

neuronal nuclei in the HCG of C and DSS-fed females. D) Quantification of neural tissue

in control and H_2O_2-fed females. E) Schematic outlining the feeding protocol as well as

the feeding protocol followed by the recovery period. Quantification of neural tissue after

the feeding protocol and after the recovery period (Rec). Scale bar 50μm. Statistical

significance is indicated as follows: *$p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, n.s = not

significant. Gut schematic from *BioRender.com*.

ISC proliferation does not mediate neural plasticity.

Since DSS -feeding results in an increase in ISC proliferation, we set out to test if ISC cell

division was signaling neurons to grow. DSS-mediated ISC proliferation has been

described as an increase in histone 3 phosphorylation (H3P) signal along the midgut

(Amcheslavsky et al., 2009), mostly concentrated in its posterior part at the midgut to

hindgut junction. Although H3P signal in the anterior part is quite scarce if not absent in

control animals a significant increase in signal was detected in DSS-fed flies (Figure 2A).

To better monitor the extend of ISC divisions exclusively taking place during the feeding

protocol in the anterior part of the gut we used lineage tracing (ReDDM) (Antonello et al.,

2015) and observed substantial ISC proliferation in DSS-fed versus control animals

(Figure 2B). To test the effect of this extensive ISC proliferation on neural growth we

sought to block ISC cell division in adult animals at the same time they were feeding on

DSS to assess if neural growth would be suppressed. We blocked ISC proliferation using

a combination of a null allele (*Cdk1^null^*) and a temperature-sensitive allele of *Cdk1* (*Cdk1^1E1–*

*24^*) (Rodríguez et al., 2024), which rendered flies mutant for *Cdk1* during DSS feeding

(Figure 2C). We observed no difference in neural growth between DSS-fed flies with and

without proliferation blockage (Figure 2D), indicating that ISC cell division is not

responsible for the DSS induced neural growth.

Gut-derived ROS stimulates neural plasticity.

DSS feeding results in epithelial damage, causing cell death and an increase in ROS

levels (Wu et al., 2012). To test the possible contribution of ROS levels to neural growth,

and to overcome the limitation that using the GAL4/UAS system to visualize neurons

imposes on the genetic manipulation of the gut with other heterologous systems, we took

advantage of antioxidant drugs. To investigate if ROS could signal neurons to grow, we fed

flies with DSS plus the ROS inhibitor N-acetyl cysteine (NAC) and compared them to flies

fed with DSS only. In this case we observed a reduction of neural growth in the NAC-fed

condition (Figure 2E). Moreover, the neural tissue values obtained for the DSS + NAC

feeding protocol were not significantly different from the values observed for the Control

+ NAC condition, arguing for a strong suppression of the neural growth (Figure 2E). Thus,

DSS-dependent ROS promotes neural plasticity.

Reversal of neural growth as the gut returns to homeostasis is not caspase

dependent.

We were intrigued about the mechanisms regulating the reversal of the neural growth

after the recovery period. Axon pruning can take place either through axon retraction or

degeneration (Luo and O'Leary, 2005). Examples of the latter during developmental

remodeling include the pruning of mushroom body gamma neurons (Watts et al., 2003)

as well as dendritic arborizing (da) sensory neurons ddaC (Williams and Truman, 2005).

While deletion of apoptotic activators or expression of a caspase inhibitor in MB neurons

had no effect on axon pruning (Watts et al., 2003), da neurons use local caspase

activation to direct dendrite engulfment (Williams et al., 2006). To test if there was a role

of caspases in reversal of the neural growth in the enteric nervous system, we inhibited

apoptosis specifically in neurons by overexpressing the caspase inhibitor Diap1. We

observed that reversal of neural growth still took place (Supplementary Figure 2). This

result suggests the observed reversal of neural growth could potentially be mediated by

axon retraction, rather than caspase-dependent mechanisms.

Figure 2

Figure 2. Neural plasticity does not depend on ISC proliferation but on gut-derived ROS. A) Quantification of H3P/Hoechst positive cells in the anterior midgut and the whole midgut in C and DSS-fed females. B) Schematic showing activation of ReDDM lineage tracing during the feeding protocol. Z-stack projected confocal images of the anterior midgut stained with Hoechst (grey) and showing the cell membrane of ISCs/EBs (magenta, anti-GFP) in C and DSS-fed females. Quantification of the percentage of the ISC/EB area (GFP signal) from the total area of the anterior mid gut in C and DSS-fed females. C-D) Schematic showing temperature dependent knock out of Cdk1 function during the feeding protocol. C) Quantification of H3P/Hoechst positive cells in the anterior midgut and the whole midgut in wild type (heterozygous) and *Cdk1* null females after DSS feeding. D) Quantification of neural tissue in wild type and *Cdk1* null females after DSS feeding. E) Z-stack projected confocal images of neuronal processes in the anterior midgut of DSS, DSS + NAC and C +NAC -fed females and quantification of neural tissue. Scale bar 50μm. Statistical significance is indicated as follows: * *p < 0.05,* ** *p < 0.01,* *** *p < 0.001,* n.s = not significant.

Neural plasticity is exclusive to females and is actively suppressed when female neurons are masculinized.

Interestingly, males did not exhibit neural growth following DSS treatment (Figure 3A). To investigate whether this neural growth was exclusive to female neurons, we aimed to create female individuals with masculinized neurons. Most sex differences in *Drosophila* arise from the female specific expression of the *Sex lethal* gene (*Sxl*) which codes for an RNA binding protein (Bell et al., 1988). Sxl induces the alternative splicing of the *transformer* (*tra*) gene generating the female fate determining TraF protein only in females (Belote et al., 1989; Boggs et al., 1987; Inoue et al., 1990; Sosnowski et al., 1989). We first confirmed that neurons in the HCG expressed the female-specific alternative splicing variant of *tra* while males didn't (Figure 3B,C) (Hérault et al., 2024). Subsequently, we proceeded to masculinize neurons in females by means of *tra* downregulation with RNAi during development (Figure 3D). We confirmed that the presence of the *UAS-Dcr2* transgene did not affect the response of wild type females to DSS, and thus, the expected increase in neural tissue was observed. Interestingly, DSS feeding did not have any effect in masculinized females and the amount of neural tissue was not significantly different

269 between control and DSS-fed flies. In addition, DSS-fed masculinized females displayed

270 significantly less neural tissue that DSS-fed control flies, aligning with the absence of

271 neural growth in masculinized females.

Figure 3

Figure 3. **Neural plasticity is not observed in males and masculinization of female**

neurons abolishes neural growth. A) Z-stack projected confocal images of neuronal

processes in the anterior midgut of C and DSS-fed males, and quantification of neural

tissue. Blue outline in box plots indicates male. B-C) Z-stack projected confocal images

at the level of the HCG showing female-specific *tra* alternative splicing in neurons (grey,

GFP; magenta, Elav) in females (B) and absence in males (C). D-E) Schematic showing

transgene expression in neurons in relation to the feeding protocol. D) Quantification and

comparison of neural tissue in females and females with masculinized neurons both in

C versus DSS-fed conditions. E) Quantification and comparison of neural tissue in males and males with feminized neurons both in C versus DSS-fed conditions. Scale bar 50μm. Statistical significance is indicated as follows: * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, n.s = not significant.

We also carried out the reverse experiment and attempted to confer plasticity to male neurons by feminizing them through developmental *Sxl* overexpression (Figure 3E) (Bell et al., 1991). In this case, the DSS treatment had no effect in feminized males, and we did not observe the neural growth we expected to see for a successfully feminized male. Moreover, both control and DSS-fed feminized males showed less amount of neural tissue than wild type males. These results indicate that neuronal Sxl overexpression in males is not sufficient to induce damage-responsive plasticity, and may even impair neural tissue growth in our context.

The lack of neural growth in males following DSS-induced gut damage, along with the suppression of this response in masculinized female neurons, suggests that neural plasticity is limited in males compared to females.

DISCUSSION

Our findings establish the fly gut and its innervation as a tractable model to dissect the mechanisms of neural plasticity in response to acute nearby tissue damage. This system enables the study of both structural growth and retraction in intact adult neurons. We show that DSS-induced ROS promotes neural growth, consistent with growing evidence that ROS supports structural plasticity (Oswald et al., 2018). In *Drosophila*, regrowth after developmental pruning and regeneration involves TOR signaling and interactions with the InR/PI3K/Akt pathway (Sanal et al., 2023; Yaniv and Schuldiner, 2016). Mammals also rely on mTOR for neuronal regeneration (Lund-Ricard et al., 2020). Given that low, transient ROS can activate both pathways (Yoshida et al., 2011; Zhang et al., 2020), these are candidates to mediate the observed DSS-dependent neural growth. Conversely, axon retraction upon recovery may involve cytoskeletal remodeling via RhoA signaling. In *Drosophila* there is evidence for a RhoA-mediated retraction pathway actively suppressed by a GTPase-activating protein (GAP) (Billuart et al., 2001). Additionally, Rho

guanine nucleotide exchange factors (GEFs) have also been identified (Hakeda-Suzuki et

al., 2002). Thus, extracellular cues may modulate retraction via GAP/GEF balance. Given

its parallels with repulsive axon guidance, cell surface molecules like Ephrins, Slits, and

Semaphorins—which signal through RhoA (Bashaw and Klein, 2010; Kania and Klein,

2016; Pasterkamp, 2012)—could be expressed by the gut or muscle during recovery to

help coordinate neural plasticity.

In addition, we uncover a clear sexual dimorphism: neural growth is absent in males and

suppressed in masculinized female neurons. However, feminizing male neurons was not

sufficient to induce plasticity. One possibility is that signals from other feminized

surrounding tissues could be necessary. Indeed, sex-specific differences in morphology,

physiology, pathology, and cell behavior—including intestinal stem cells and

enterocytes—have been documented in the fly digestive system (Blackie et al., 2024;

Hadjieconomou et al., 2020; Hudry et al., 2019, 2016; Regan et al., 2022, 2016; Reiff et

al., 2015). Interestingly, although we did not detect structural changes in TTCs in the

anterior midgut, plasticity in the tracheal system has been reported in other regions of

the gut (Linneweber et al., 2014; Perochon et al., 2021; Tamamouna et al., 2021). TTC's

behavior is strikingly similar to that of neurons: they respond to DSS and H_2O_2 feeding

with ROS-dependent sprouting (Perochon et al., 2021; Tamamouna et al., 2021),

although sex specific differences remain to be tested. Given the recent link between the

dimorphic tracheal system and sex-specific gut morphology (Blackie et al., 2024), it is

tempting to speculate that the functional and/or structural plasticity of subsets of TTCs

could also be dimorphic. It is well established that sex differences influence organ

formation, function and physiology (Mank and Rideout, 2021). The results presented here

showcase the fly gut and its innervation as a model to unravel how biological sex

influences neural plasticity. Understanding these mechanisms could guide the

development of sex-specific therapeutic strategies for neural repair.

MATERIALS AND METHODS

Fly stocks

The following fly lines were used: *GMR51F12-GAL4* (BDSC:58685), *nSyb-GAL4.S* (BDSC:

51635), *UAS-Flybow.1.1B* (used as 10xUAS-CD8::GFP, BDSC: 56802, 56803), *DSRF.Term-*

GAL4 (BDSC: 25753), *esg-GAL4* (a gift from M. Milán, unknown insertion), *UAS-*
mCD8::GFP (BDSC:5130), *UAS-H2B::RFP* (a gift from J. Morante), *tubP-GAL80ts* (BDSC:
7017), *Cdk1null* (BDSC: 6643), *Cdk1E1–24* (BDSC: 6641), *UAS-Diap1* (BDSC: 15310), *traF-*
GAL4 (a gift from B. Hudry), *UAS-Dcr-2* (BDSC: 24650), *UAS-traRNAi.TRiPJF03132* (*UAS-*
traRNAi, BDSC: 28512), *UASp-Sxl.alt5-C8* (*UAS-Sxl*, BDSC: 58484). The list of complete
genotypes is available in Supplementary Information.

Feeding protocols
For feeding assays, virgin females were collected and aged on standard media at 25°C for
3-4 days before undergoing mating for 24hrs. Then same procedure was carried out when
experiments were performed with males. Flies were recovered from the mating and
transferred in groups of 10 to an empty vial containing 5 pieces of 2cm × 2cm of bench
paper soaked with 500ml of the desired feeding solution. Flies were transferred to a new
vial with fresh feeding paper every day for the specified number of days prior to
dissection. The feeding protocol was always done at 29°C. For DSS (Merck, 42867)
induced damage, flies were fed with 5% sucrose (Control) or with 6% DSS in 5% sucrose
(DSS) for 4 days. For the SMURF assay, the last control and DSS feeding solutions prior to
the day of dissection were laced with Brilliant Blue FCF (80717, Merck) at 2.5% w/v. For
H_2O_2 (1072100250, Merck) induced damage, flies were fed with 5% Sucrose (Control) or
1% H_2O_2 in 5% sucrose (H_2O_2) for 4 days. For NAC treatment, flies were fed with 1,2mM
NAC in 5% sucrose (Control + NAC) or 6% DSS + 1,2mM NAC in 5% sucrose (DSS + NAC)
for 4 days. For recovery protocols, after the DSS feeding protocol control and DSS-fed
flies were each transferred to vials with 5% sucrose for 4 days (Control + Recovery; DSS
+ Recovery).

Temperature sensitive assays
To visualize the amount of proliferation taking place exclusively during the four days of
the feeding protocol we set ReDDM-based crosses (Antonello et al., 2015). This lineage
tracing approach uses the Gal80 temperature sensitive allele (*Gal80ts*), which inhibits
GAL4 transcription at permissive temperature (19°C) and allows it at the restrictive one
(29°C) (McGuire et al., 2003), and *esg-GAL4* to label ISCs and EBs. Crosses were grown
and flies were kept at 17°C to ensure Gal80ts repression until adult mated female flies

were shifted to 29°C to inactive the Gal80 and carry out the feeding protocol. To block ISC

proliferation during the feeding protocol, *Cdk1^{null}/ Cdk1$^{1E1–24}$* flies were grown, aged and

mated at 17°C, Cdk1 function was abrogated once the flies were transferred to 29°C

(Rodriguez et al., 2024) during the DSS feeding protocol.

Dissection and immunostaining

To obtain the gut flies were dissected in PBS. Adult animals had their heads cut off with

scissors by the neck, their abdomens were pulled apart from the thorax and opened to

extract the gut. With this procedure the digestive tube was severed well above the crop

duct. Once fat and ovaries were removed, the gut was handled by its posterior end and

transferred to PBS in a well drawn onto a poly-l-lysine-coated slide (P1524, Merck) using

hydrophobic silicone (Flowsil, Intek Adhesives). In cases were the anterior part of the

midgut had to be repositioned, we manipulated the gut from the crop to avoid touching

the digestive tube. Guts were fixed at room temperature for 30min in 4% PFA in PBS.

Washes were done with PBS-T (PBS, 0.25% Triton X-100). To label the muscle or nuclei,

guts were incubated at room temperature for 30 min with Alexa Fluor™ 635 Phalloidin

(1/400, A34054, Invitrogen) or Hoechst 33258 (1/1000, H3569, Invitrogen) respectively in

PBS-T. A final wash with PBS was done prior to mounting. When antibodies were used,

primaries were incubated overnight at 4°C, and secondaries were incubated at room

temperature for 2–3 h. The following antibodies were used: rat-anti-Elav (1/100, 7E8A10,

DSHB), rabbit-anti-H3P (1/1000, 06-570, EDM Milipore), rabbit-anti-DsRed (1/200,

632496, Clontech), chicken-anti-GFP (1/800, ab13970, Abcam), Alexa Fluor™ 568 goat-

anti-rat, Alexa Fluor™ 488 goat-anti-rabbit, Alexa Fluor™ 568 goat-anti-rabbit, Alexa

Fluor™ 488 goat-anti-chicken (1/500, A11077, A11008, A11011, A11039, Life

Technologies). Samples were mounted in VECTASHIELD® PLUS Antifade Mounting (H-

1900-10, Vector Laboratories).

Microscopy

Samples were imaged using a Zeiss LSM 880 confocal microscope (Zeiss, Germany)

equipped with a 25X 0.8 glycerol immersion media objective and an Argon and a

HeNe633 lasers. For neural network analysis, to consistently image the same region in

each gut, we used the end of the proventriculus as a positional landmark and captured

the gut section that fit within the visual field of the 25X objective at 0.8 magnification,

which in our set up generated images of 566,79 μm^2. As quality control for dissection,

only guts with an intact HCG were considered since this ganglion contains the cell bodies

of many of the neurons innervating the anterior midgut. Stacks of images were acquired

with a pixel size of 0.55x0.55x1um (xyz) and a pixel dwell time of 2.05us.

Image processing and quantifications based on images

Images were processed with Fiji (version ImageJ 1.53c, (Schindelin et al., 2012) and

figures assembled using Adobe Illustrator (Adobe, San Jose, CA).

For quantification of neural tissue and gut parameters: The channel with the neuron

staining was filtered in 3D (Ollion et al., 2013) first with the maximum filter and then with

the minimum one, with a radius of 3.3um in x and y and 2um in z, in both filters. Then the

filtered stack was processed using the Tubeness plugin (sigma = 0.66) (Sato et al., 1998).

Finally, images were binarized using the Bernsen's AutoLocal threshold method (radius =

2) (Bernsen, 1986). The resulting binary stack was analyzed first with the 3D Object

Counter function to obtain the total neuronal network volume (mm^3). The channel with

the gut staining was processed by background subtraction and then thresholder using

the Li's method (Li and Lee, 1993). The resulting binary stack was analyzed with the 3D

Object Counter function to obtain the volume of the gut.

For other quantifications: The number of Elav/Hoechst positive nuclei in the HCG was

counted from the confocal stack using the 3D Object Counter plugin. The number of TTC

nuclei was counted by hand from the confocal stack maximum projection. TTCs volume

was obtained using the same procedure as for neural tissue quantification. The number

of H3P/ Hoechst positive nuclei in the midgut was counted directly under the confocal

microscope. For quantification of cumulative proliferation using ReDDM, xyz stacks of

half the gut were taken and z-projected. The Hoechst signal was manually outlined to

measure the gut area. To quantify the area of GFP signal from mCD8::GFP-labeled cells

(representing ISCs and EBs resulting from ISCs divisions) a Gaussian Blur filter (radius =

2.0) was applied. The image was then thresholded using the "MaxEntropy" algorithm

(Kapur et al., 1985) and finally, the GFP-positive regions larger than 10μm^2 were

quantified. The GFP area was represented as a percentage of the total area.

Other quantifications

For the SMURF assay (Rera et al., 2012), on the 4th day of the DSS feeding protocol, we assessed the proportion of surviving flies that were positive for the SMURF phenotype.

Statistics

Data analysis was carried out using Prism 6 (GraphPad Software Inc., San Diego, CA) and Statgraphics Centurion version 18 (Statgraphics Technologies Inc., The Plains, VA). Experiments were performed with three biological replicates yielding similar results, and a representative replicate is shown. A minimum of 15 guts were analyzed for each condition. For the SMURF assay ten tubes with 10 flies each were used. For data shown in box plots, the median is given between the first and third quartiles (ends of the box). Whiskers represent the maximum and minimum values of the data. Bar graphs display the mean, with error bars representing the standard deviation.

For the amount of neural tissue and TTCs tissue: To analyze the effect of the treatment on neural tissue volume and TTCs tissue volume, analysis of covariance (ANCOVA) was performed. The dependent variable was neural tissue volume or TTCs tissue volume, the fixed factor was the treatment, and the covariate was gut volume (Figure 1B, D, E; Figure 2D, E; Figure 3A, D, E; Supplementary Figure 1B, C; Supplementary Figure 2).

For other quantifications: For pairwise comparisons of data sets following a normal distribution (Figure 1C) the unpaired t-test was performed. For pairwise comparisons of data sets not following a normal distribution we used the non-parametric Mann Whitney test (Figure 2A-C; Supplementary Figure 1A).

FUNDING

This work was funded by the Spanish Ministerio de Ciencia e Innovación (grant PID2019-107340GB-I00) and Agència de Gestió d'Ajuts Universitaris i de Recerca of the Generalitat de Catalunya (2021SGR00293) to M.M. Work in I.M-A. was funded by an ERC Advanced Grant (ERCAdG 787470 "IntraGutSex"), a UKRI Horizon Europe guarantee grant (EPSRC-ERC EP/Y03298/1), MRC intramural funding and the Francis Crick Institute, which receives its core funding from Cancer Research UK (FC001317 and FC001175), the UK Medical Research Council (FC001317 and FC001175), and the Wellcome Trust (FC001317 and FC001175).

ACKNOWLEDGEMENTS

We thank C. González, M. Milán, C. Estella, J. Morante, B. Hudry, M. Corominas, F. Serras, BDSC and DSHB for reagents and members of the Corominas' and Serras' laboratories for insightful discussions. We are grateful to Q. Zhu and O. Ramirez for their assistance with early pilot experiments that helped shape this research. We are indebted to our colleagues F. Cebrià, M. Milán, M. Corominas, F. Serras, B. Payer and the CCiTUB technological platform for their invaluable support and encouragement during this project.

CONFLICT OF INTERESTS

The authors declare that they have no conflict of interest.

CONTRIBUTIONS

Conceptualization: M.M.; Methodology: M.R-M., M.B., P.G., A.M., S.R., I.M-A., M.M.; Investigation: M.R-M., M.M.; Formal Analysis: M.R-M., P.G., A.M., S.R., I.M-A., M.M.; Validation: M.R-M., M.M.; Visualization: M.R-M., M.M.; Data Curation: M.R-M., M.M.; Resources: P.G., A.M., I.M-A.; Supervision: M.M.; Project Administration: M.M.; Funding Acquisition: M.M.; Writing – Original Draft: M.M. with contributions from all the authors.

REFERENCES

Amcheslavsky, A., Jiang, J., Ip, Y.T., 2009. Tissue Damage-Induced Intestinal Stem Cell Division in Drosophila. Cell Stem Cell 4, 49–61. https://doi.org/10.1016/j.stem.2008.10.016

Ameku, T., Beckwith, H., Blackie, L., Miguel-Aliaga, I., 2020. Food, microbes, sex and old age: on the plasticity of gastrointestinal innervation. Curr. Opin. Neurobiol. 62, 83–91. https://doi.org/10.1016/j.conb.2019.12.004

Antonello, Z.A., Reiff, T., Ballesta-Illan, E., Dominguez, M., 2015. Robust intestinal homeostasis relies on cellular plasticity in enteroblasts mediated by miR-8– Escargot switch. EMBO J. 34, 2025–2041. https://doi.org/10.15252/embj.201591517

Bashaw, G.J., Klein, R., 2010. Signaling from Axon Guidance Receptors. Cold Spring Harb. Perspect. Biol. 2, a001941–a001941. https://doi.org/10.1101/cshperspect.a001941

Bell, L.R., Horabin, J.I., Schedl, P., Cline, T.W., 1991. Positive autoregulation of Sex-lethal by alternative splicing maintains the female determined state in Drosophila. Cell 65, 229–239. https://doi.org/10.1016/0092-8674(91)90157-T

Bell, L.R., Maine, E.M., Schedl, P., Cline, T.W., 1988. Sex-lethal, a Drosophila sex
determination switch gene, exhibits sex-specific RNA splicing and sequence
similarity to RNA binding proteins. Cell 55, 1037–1046.
https://doi.org/10.1016/0092-8674(88)90248-6
Belote, J.M., McKeown, M., Boggs, R.T., Ohkawa, R., Sosnowski, B.A., 1989. Molecular
genetics of *transformer* , a genetic switch controlling sexual differentiation in
Drosophila. Dev. Genet. 10, 143–154. https://doi.org/10.1002/dvg.1020100304
Bentley, D., Toroian-Raymond, A., 1981. Embryonic and postembryonic morphogenesis
of a grasshopper interneuron. J. Comp. Neurol. 201, 507–518.
https://doi.org/10.1002/cne.902010404
Bernsen, J., 1986. Dynamic Thresholding of Gray Level Image. Proc. Int. Conf. Pattern
Recognit. Berl. ICPR 86 1251–1255.
Billuart, P., Winter, C.G., Maresh, A., Zhao, X., Luo, L., 2001. Regulating Axon Branch
Stability. Cell 107, 195–207. https://doi.org/10.1016/S0092-8674(01)00522-0
Blackie, L., Gaspar, P., Mosleh, S., Lushchak, O., Kong, L., Jin, Y., Zielinska, A.P., Cao, B.,
Mineo, A., Silva, B., Ameku, T., Lim, S.E., Mao, Y., Prieto-Godino, L., Schoborg, T.,
Varela, M., Mahadevan, L., Miguel-Aliaga, I., 2024. The sex of organ geometry.
Nature 630, 392–400. https://doi.org/10.1038/s41586-024-07463-4
Boggs, R.T., Gregor, P., Idriss, S., Belote, J.M., McKeown, M., 1987. Regulation of sexual
differentiation in D. melanogaster via alternative splicing of RNA from the
transformer gene. Cell 50, 739–747. https://doi.org/10.1016/0092-
8674(87)90332-1
Bucher, D., Pflüger, H.-J., 2000. Directional sensitivity of an identified wind-sensitive
interneuron during the postembryonic development of the locust. J. Insect
Physiol. 46, 1545–1556. https://doi.org/10.1016/S0022-1910(00)00078-0
Buchon, N., Osman, D., David, F.P.A., Yu Fang, H., Boquete, J.-P., Deplancke, B.,
Lemaitre, B., 2013. Morphological and Molecular Characterization of Adult
Midgut Compartmentalization in Drosophila. Cell Rep. 3, 1725–1738.
https://doi.org/10.1016/j.celrep.2013.04.001
Chen, J., Reiher, W., Hermann-Luibl, C., Sellami, A., Cognigni, P., Kondo, S., Helfrich-
Förster, C., Veenstra, J.A., Wegener, C., 2016. Allatostatin A Signalling in
Drosophila Regulates Feeding and Sleep and Is Modulated by PDF. PLOS Genet.
12, e1006346. https://doi.org/10.1371/journal.pgen.1006346
Cirillo, C., Brihmat, N., Castel-Lacanal, E., Le Friec, A., Barbieux-Guillot, M., Raposo, N.,
Pariente, J., Viguier, A., Simonetta-Moreau, M., Albucher, J.-F., Olivot, J.-M.,
Desmoulin, F., Marque, P., Chollet, F., Loubinoux, I., 2020. Post-stroke
remodeling processes in animal models and humans. J. Cereb. Blood Flow
Metab. 40, 3–22. https://doi.org/10.1177/0271678X19882788
Cognigni, P., Bailey, A.P., Miguel-Aliaga, I., 2011. Enteric Neurons and Systemic Signals
Couple Nutritional and Reproductive Status with Intestinal Homeostasis. Cell
Metab. 13, 92–104. https://doi.org/10.1016/j.cmet.2010.12.010
Cui, X., Meiselman, M.R., Thornton, S.N., Yapici, N., 2024. A gut-brain-gut interoceptive
circuit loop gates sugar ingestion in Drosophila.
https://doi.org/10.1101/2024.09.02.610892
Gao, Z., Pang, Z., Chen, Y., Lei, G., Zhu, S., Li, G., Shen, Y., Xu, W., 2022. Restoring After
Central Nervous System Injuries: Neural Mechanisms and Translational

Applications of Motor Recovery. Neurosci. Bull. 38, 1569–1587.
https://doi.org/10.1007/s12264-022-00959-x

Hadjieconomou, D., King, G., Gaspar, P., Mineo, A., Blackie, L., Ameku, T., Studd, C., De
Mendoza, A., Diao, F., White, B.H., Brown, A.E.X., Plaçais, P.-Y., Préat, T., Miguel-
Aliaga, I., 2020. Enteric neurons increase maternal food intake during
reproduction. Nature 587, 455–459. https://doi.org/10.1038/s41586-020-2866-8

Hakeda-Suzuki, S., Ng, J., Tzu, J., Dietzl, G., Sun, Y., Harms, M., Nardine, T., Luo, L.,
Dickson, B.J., 2002. Rac function and regulation during Drosophila development.
Nature 416, 438–442. https://doi.org/10.1038/416438a

Hérault, C., Pihl, T., Hudry, B., 2024. Cellular sex throughout the organism underlies
somatic sexual differentiation. Nat. Commun. 15, 6925.
https://doi.org/10.1038/s41467-024-51228-6

Holsopple, J.M., Cook, K.R., Popodi, E.M., 2022. Enteroendocrine cell expression of
split-GAL4 drivers bearing regulatory sequences associated with panneuronally
expressed genes in Drosophila melanogaster. Open Access.

Howard, A.M., LaFever, K.S., Fenix, A.M., Scurrah, C.R., Lau, K.S., Burnette, D.T., Bhave,
G., Ferrell, N., Page-McCaw, A., 2019. DSS-induced damage to basement
membranes is repaired by matrix replacement and crosslinking. J. Cell Sci. 132,
jcs226860. https://doi.org/10.1242/jcs.226860

Hudry, B., De Goeij, E., Mineo, A., Gaspar, P., Hadjieconomou, D., Studd, C.,
Mokochinski, J.B., Kramer, H.B., Plaçais, P.-Y., Preat, T., Miguel-Aliaga, I., 2019.
Sex Differences in Intestinal Carbohydrate Metabolism Promote Food Intake and
Sperm Maturation. Cell 178, 901-918.e16.
https://doi.org/10.1016/j.cell.2019.07.029

Hudry, B., Khadayate, S., Miguel-Aliaga, I., 2016. The sexual identity of adult intestinal
stem cells controls organ size and plasticity. Nature 530, 344–348.
https://doi.org/10.1038/nature16953

Hyer, M.M., Phillips, L.L., Neigh, G.N., 2018. Sex Differences in Synaptic Plasticity:
Hormones and Beyond. Front. Mol. Neurosci. 11, 266.
https://doi.org/10.3389/fnmol.2018.00266

Inoue, K., Hoshijima, K., Sakamoto, H., Shimura, Y., 1990. Binding of the Drosophila Sex-
lethal gene product to the alternative splice site of transformer primary
transcript. Nature 344, 461–463. https://doi.org/10.1038/344461a0

Jenett, A., Rubin, G.M., Ngo, T.-T.B., Shepherd, D., Murphy, C., Dionne, H., Pfeiffer, B.D.,
Cavallaro, A., Hall, D., Jeter, J., Iyer, N., Fetter, D., Hausenfluck, J.H., Peng, H.,
Trautman, E.T., Svirskas, R.R., Myers, E.W., Iwinski, Z.R., Aso, Y., DePasquale,
G.M., Enos, A., Hulamm, P., Lam, S.C.B., Li, H.-H., Laverty, T.R., Long, F., Qu, L.,
Murphy, S.D., Rokicki, K., Safford, T., Shaw, K., Simpson, J.H., Sowell, A., Tae, S.,
Yu, Y., Zugates, C.T., 2012. A GAL4-Driver Line Resource for Drosophila
Neurobiology. Cell Rep. 2, 991–1001.
https://doi.org/10.1016/j.celrep.2012.09.011

Jiang, H., Tian, A., Jiang, J., 2016. Intestinal stem cell response to injury: lessons from
Drosophila. Cell. Mol. Life Sci. 73, 3337–3349. https://doi.org/10.1007/s00018-
016-2235-9

Kania, A., Klein, R., 2016. Mechanisms of ephrin–Eph signalling in development,
physiology and disease. Nat. Rev. Mol. Cell Biol. 17, 240–256.
https://doi.org/10.1038/nrm.2015.16

Kapur, J.N., Sahoo, P.K., Wong, A.K.C., 1985. A new method for gray-level picture thresholding using the entropy of the histogram. Comput. Vis. Graph. Image Process. 29, 273–285. https://doi.org/10.1016/0734-189X(85)90125-2

Kenmoku, H., Ishikawa, H., Ote, M., Kuraishi, T., Kurata, S., 2016. A subset of neurons controls the permeability of the peritrophic matrix and midgut structure in *Drosophila* adults. J. Exp. Biol. jeb.122960. https://doi.org/10.1242/jeb.122960

Kirby, E.D., Andrushko, J.W., Rinat, S., D'Arcy, R.C.N., Boyd, L.A., 2024. Investigating female versus male differences in white matter neuroplasticity associated with complex visuo-motor learning. Sci. Rep. 14, 5951. https://doi.org/10.1038/s41598-024-56453-z

Ksiazek-Winiarek, D.J., Szpakowski, P., Glabinski, A., 2015. Neural Plasticity in Multiple Sclerosis: The Functional and Molecular Background. Neural Plast. 2015, 1–11. https://doi.org/10.1155/2015/307175

Kuraishi, T., Kenmoku, H., Kurata, S., 2015. From mouth to anus: Functional and structural relevance of enteric neurons in the Drosophila melanogaster gut. Insect Biochem. Mol. Biol. 67, 21–26. https://doi.org/10.1016/j.ibmb.2015.07.003

Lee, S., Stevens, C.F., 2007. General design principle for scalable neural circuits in a vertebrate retina. Proc. Natl. Acad. Sci. 104, 12931–12935. https://doi.org/10.1073/pnas.0705469104

Lemaitre, B., Miguel-Aliaga, I., 2013. The Digestive Tract of *Drosophila melanogaster*. Annu. Rev. Genet. 47, 377–404. https://doi.org/10.1146/annurev-genet-111212-133343

Li, C.H., Lee, C.K., 1993. Minimum cross entropy thresholding. Pattern Recognit. 26, 617–625. https://doi.org/10.1016/0031-3203(93)90115-D

Linneweber, G.A., Jacobson, J., Busch, K.E., Hudry, B., Christov, C.P., Dormann, D., Yuan, M., Otani, T., Knust, E., de Bono, M., Miguel-Aliaga, I., 2014. Neuronal Control of Metabolism through Nutrient-Dependent Modulation of Tracheal Branching. Cell 156, 69–83. https://doi.org/10.1016/j.cell.2013.12.008

Lund-Ricard, Y., Cormier, P., Morales, J., Boutet, A., 2020. mTOR Signaling at the Crossroad between Metazoan Regeneration and Human Diseases. Int. J. Mol. Sci. 21, 2718. https://doi.org/10.3390/ijms21082718

Luo, L., O'Leary, D.D.M., 2005. AXON RETRACTION AND DEGENERATION IN DEVELOPMENT AND DISEASE. Annu. Rev. Neurosci. 28, 127–156. https://doi.org/10.1146/annurev.neuro.28.061604.135632

Mahar, M., Cavalli, V., 2018. Intrinsic mechanisms of neuronal axon regeneration. Nat. Rev. Neurosci. 19, 323–337. https://doi.org/10.1038/s41583-018-0001-8

Mank, J.E., Rideout, E.J., 2021. Developmental mechanisms of sex differences: from cells to organisms. Development 148. https://doi.org/10.1242/dev.199750

Marianes, A., Spradling, A.C., 2013. Physiological and stem cell compartmentalization within the Drosophila midgut. eLife 2, e00886. https://doi.org/10.7554/eLife.00886

McGuire, S.E., Le, P.T., Osborn, A.J., Matsumoto, K., Davis, R.L., 2003. Spatiotemporal Rescue of Memory Dysfunction in *Drosophila*. Science 302, 1765–1768. https://doi.org/10.1126/science.1089035

Menon, K.P., Carrillo, R.A., Zinn, K., 2013. Development and plasticity of the *Drosophila*
larval neuromuscular junction. WIREs Dev. Biol. 2, 647–670.
https://doi.org/10.1002/wdev.108
Miguel-Aliaga, I., Jasper, H., Lemaitre, B., 2018. Anatomy and Physiology of the Digestive
Tract of *Drosophila melanogaster*. Genetics 210, 357–396.
https://doi.org/10.1534/genetics.118.300224
Min, S., Oh, Y., Verma, P., Whitehead, S.C., Yapici, N., Van Vactor, D., Suh, G.S., Liberles,
S., 2021. Control of feeding by Piezo-mediated gut mechanosensation in
Drosophila. eLife 10, e63049. https://doi.org/10.7554/eLife.63049
Murphy, T.H., Corbett, D., 2009. Plasticity during stroke recovery: from synapse to
behaviour. Nat. Rev. Neurosci. 10, 861–872. https://doi.org/10.1038/nrn2735
Nudo, R.J., 2003. Functional and structural plasticity in motor cortex: implications for
stroke recovery. Phys. Med. Rehabil. Clin. N. Am. 14, S57–S76.
https://doi.org/10.1016/S1047-9651(02)00054-2
Olds, W.H., Xu, T., 2014. Regulation of food intake by mechanosensory ion channels in
enteric neurons. eLife 3, e04402. https://doi.org/10.7554/eLife.04402
Ollion, J., Cochennec, J., Loll, F., Escudé, C., Boudier, T., 2013. TANGO: a generic tool for
high-throughput 3D image analysis for studying nuclear organization.
Bioinformatics 29, 1840–1841. https://doi.org/10.1093/bioinformatics/btt276
Oswald, M.C.W., Garnham, N., Sweeney, S.T., Landgraf, M., 2018. Regulation of
neuronal development and function by ROS. FEBS Lett. 592, 679–691.
https://doi.org/10.1002/1873-3468.12972
Pasterkamp, R.J., 2012. Getting neural circuits into shape with semaphorins. Nat. Rev.
Neurosci. 13, 605–618. https://doi.org/10.1038/nrn3302
Perochon, J., Yu, Y., Aughey, G.N., Medina, A.B., Southall, T.D., Cordero, J.B., 2021.
Dynamic adult tracheal plasticity drives stem cell adaptation to changes in
intestinal homeostasis in Drosophila. Nat. Cell Biol. 23, 485–496.
https://doi.org/10.1038/s41556-021-00676-z
Petersen, E.D., Sharkey, E.D., Pal, A., Shafau, L.O., Zenchak-Petersen, J., Peña, A.J.,
Aggarwal, A., Prakash, M., Hochgeschwender, U., 2022. Restoring Function After
Severe Spinal Cord Injury Through BioLuminescent-OptoGenetics. Front. Neurol.
12, 792643. https://doi.org/10.3389/fneur.2021.792643
Petsakou, A., Liu, Yifang, Liu, Ying, Comjean, A., Hu, Y., Perrimon, N., 2023. Cholinergic
neurons trigger epithelial Ca2+ currents to heal the gut. Nature 623, 122–131.
https://doi.org/10.1038/s41586-023-06627-y
Prosperini, L., Piattella, M.C., Giannì, C., Pantano, P., 2015. Functional and Structural
Brain Plasticity Enhanced by Motor and Cognitive Rehabilitation in Multiple
Sclerosis. Neural Plast. 2015, 1–12. https://doi.org/10.1155/2015/481574
Regan, J.C., Khericha, M., Dobson, A.J., Bolukbasi, E., Rattanavirotkul, N., Partridge, L.,
2016. Sex difference in pathology of the ageing gut mediates the greater
response of female lifespan to dietary restriction. eLife 5.
https://doi.org/10.7554/elife.10956
Regan, J.C., Lu, Y.-X., Ureña, E., Meilenbrock, R.L., Catterson, J.H., Kißler, D., Fröhlich, J.,
Funk, E., Partridge, L., 2022. Sexual identity of enterocytes regulates autophagy
to determine intestinal health, lifespan and responses to rapamycin. Nat. Aging
2, 1145–1158. https://doi.org/10.1038/s43587-022-00308-7

Reiff, T., Jacobson, J., Cognigni, P., Antonello, Z., Ballesta, E., Tan, K.J., Yew, J.Y.,
Dominguez, M., Miguel-Aliaga, I., 2015. Endocrine remodelling of the adult
intestine sustains reproduction in Drosophila. eLife 4.
https://doi.org/10.7554/elife.06930
Rera, M., Clark, R.I., Walker, D.W., 2012. Intestinal barrier dysfunction links metabolic
and inflammatory markers of aging to death in Drosophila. Proc. Natl. Acad. Sci.
U. S. A. 109, 21528–21533. https://doi.org/10.1073/pnas.1215849110
Rodríguez, A., Foronda, D., Córdoba, S., Felipe-Cordero, D., Baonza, A., Miguez, D.G.,
Estella, C., 2024. Cell proliferation and Notch signaling coordinate the formation
of epithelial folds in the *Drosophila* leg. Development 151, dev202384.
https://doi.org/10.1242/dev.202384
Sampaio-Baptista, C., Sanders, Z.-B., Johansen-Berg, H., 2018. Structural Plasticity in
Adulthood with Motor Learning and Stroke Rehabilitation. Annu. Rev. Neurosci.
41, 25–40. https://doi.org/10.1146/annurev-neuro-080317-062015
Sanal, N., Keding, L., Gigengack, U., Michalke, E., Rumpf, S., 2023. TORC1 regulation of
dendrite regrowth after pruning is linked to actin and exocytosis. PLOS Genet. 19,
e1010526. https://doi.org/10.1371/journal.pgen.1010526
Sato, Y., Nakajima, S., Shiraga, N., Atsumi, H., Yoshida, S., Koller, T., Gerig, G., Kikinis,
R., 1998. Three-dimensional multi-scale line filter for segmentation and
visualization of curvilinear structures in medical images. Med. Image Anal. 2,
143–168. https://doi.org/10.1016/S1361-8415(98)80009-1
Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D.J.,
Hartenstein, V., Eliceiri, K., Tomancak, P., Cardona, A., 2012. Fiji: an open-source
platform for biological-image analysis. Nat. Methods 9, 676–682.
https://doi.org/10.1038/nmeth.2019
Sosnowski, B.A., Belote, J.M., McKeown, M., 1989. Sex-specific alternative splicing of
RNA from the transformer gene results from sequence-dependent splice site
blockage. Cell 58, 449–459. https://doi.org/10.1016/0092-8674(89)90426-1
Tamamouna, V., Rahman, M.M., Petersson, M., Charalambous, I., Kux, K., Mainor, H.,
Bolender, V., Isbilir, B., Edgar, B.A., Pitsouli, C., 2021. Remodelling of oxygen-
transporting tracheoles drives intestinal regeneration and tumorigenesis in
Drosophila. Nat. Cell Biol. 23, 497–510. https://doi.org/10.1038/s41556-021-
00674-1
Tedeschi, A., Bradke, F., 2017. Spatial and temporal arrangement of neuronal intrinsic
and extrinsic mechanisms controlling axon regeneration. Curr. Opin. Neurobiol.
42, 118–127. https://doi.org/10.1016/j.conb.2016.12.005
Tian, A., Morejon, V., Kohoutek, S., Huang, Y., Deng, W., Jiang, J., 2022. Damage-induced
regeneration of the intestinal stem cell pool through enteroblast mitosis in the
Drosophila midgut. EMBO J. 41, e110834.
https://doi.org/10.15252/embj.2022110834
Titos, I., Juginović, A., Vaccaro, A., Nambara, K., Gorelik, P., Mazor, O., Rogulja, D., 2023.
A gut-secreted peptide suppresses arousability from sleep. Cell 186, 1382-
1397.e21. https://doi.org/10.1016/j.cell.2023.02.022
Turtzo, L.C., McCullough, L.D., 2008. Sex Differences in Stroke. Cerebrovasc. Dis. 26,
462–474. https://doi.org/10.1159/000155983

Watts, R.J., Hoopfer, E.D., Luo, L., 2003. Axon Pruning during Drosophila
Metamorphosis. Neuron 38, 871–885. https://doi.org/10.1016/S0896-
6273(03)00295-2
Weaver, L.N., Ma, T., Drummond-Barbosa, D., 2020. Analysis of Gal4 Expression
Patterns in Adult *Drosophila* Females. G3 GenesGenomesGenetics 10, 4147–
4158. https://doi.org/10.1534/g3.120.401676
Williams, D.W., Kondo, S., Krzyzanowska, A., Hiromi, Y., Truman, J.W., 2006. Local
caspase activity directs engulfment of dendrites during pruning. Nat. Neurosci.
9, 1234–1236. https://doi.org/10.1038/nn1774
Williams, D.W., Truman, J.W., 2005. Cellular mechanisms of dendrite pruning in
Drosophila : insights from in vivo time-lapse of remodeling dendritic arborizing
sensory neurons. Development 132, 3631–3642.
https://doi.org/10.1242/dev.01928
Wu, S.-C., Liao, C.-W., Pan, R.-L., Juang, J.-L., 2012. Infection-Induced Intestinal
Oxidative Stress Triggers Organ-to-Organ Immunological Communication in
Drosophila. Cell Host Microbe 11, 410–417.
https://doi.org/10.1016/j.chom.2012.03.004
Yang, C., Merlin, D., 2024. Unveiling Colitis: A Journey through the Dextran Sodium
Sulfate-induced Model. Inflamm. Bowel Dis. 30, 844–853.
https://doi.org/10.1093/ibd/izad312
Yaniv, S.P., Schuldiner, O., 2016. A fly's view of neuronal remodeling. WIREs Dev. Biol. 5,
618–635. https://doi.org/10.1002/wdev.241
Yoshida, S., Hong, S., Suzuki, T., Nada, S., Mannan, A.M., Wang, J., Okada, M., Guan, K.-
L., Inoki, K., 2011. Redox Regulates Mammalian Target of Rapamycin Complex 1
(mTORC1) Activity by Modulating the TSC1/TSC2-Rheb GTPase Pathway. J. Biol.
Chem. 286, 32651–32660. https://doi.org/10.1074/jbc.M111.238014
Yu, C., An, Z., Zhao, W., Wang, W., Gao, C., Liu, S., Wang, J., Wu, J., 2015. Sex
Differences in Stroke Subtypes, Severity, Risk Factors, and Outcomes among
Elderly Patients with Acute Ischemic Stroke. Front. Aging Neurosci. 7.
https://doi.org/10.3389/fnagi.2015.00174
Zhang, Y., Park, J., Han, S.-J., Yang, S.Y., Yoon, H.J., Park, I., Woo, H.A., Lee, S.-R., 2020.
Redox regulation of tumor suppressor PTEN in cell signaling. Redox Biol. 34,
101553. https://doi.org/10.1016/j.redox.2020.101553
775
776