Genetic modification of *Clostridium kluyveri* for heterologous *n*-butanol and *n*-hexanol

production

Caroline Schlaiß[a,b], **Saskia T. Baur**[b], **James W. Marsh**[c], **Kurt Gemeinhardt**[b],

Largus T. Angenent[a,b,d,e,f,,#]

[a]AG Angenent, Max Planck Institute for Biology, Max-Planck-Ring 5, 72076Tübingen,

Germany

[b]Environmental Biotechnology Group, Department of Geosciences, University of

Tübingen, Schnarrenbergstraße 94-96, 72076 Tübingen, Germany

[c]Department of Microbiome Science, Max Planck Institute for Biology, Max-Planck-Ring

5, 72076 Tübingen, Germany

[d]The Novo Nordisk Foundation CO_2 Research Center (CORC), Aarhus University,

Gustav Wieds Vej 10C, 8000 Aarhus C, Denmark

[e]Department of Biological and Chemical Engineering, Aarhus University, Gustav Wieds

Vej 10D, 8000 Aarhus C, Denmark

[f]Cluster of Excellence – Controlling Microbes to Fight Infections, University of Tübingen,

Auf der Morgenstelle 28, 72074 Tübingen, Germany

Running Head: Heterologous *n*-butanol and *n*-hexanol production of *C. kluyveri*

#Address correspondence to Largus T. Angenent, l.angenent@uni-tuebingen.de

*Present address: Largus T. Angenent, Department of Geosciences, University of

Tübingen, Schnarrenbergstraße 94-96, 72076 Tübingen, Germany

Abstract: 218 words

ABSTRACT

The mesophilic microbe *Clostridium kluyveri* serves as the most commonly used model microbe to elucidate the physiology and biochemistry of ethanol-based chain elongation *via* reverse ß-oxidation. In this pathway, ethanol and acetate are converted into short- and medium-chain carboxylates. However, to date, no genetic system has been published in a peer-reviewed publication. Here, we report the development of versatile genetic tools for *C. kluyveri*, utilizing the pMTL *Clostridia* shuttle vector system and thiamphenicol as a selective marker. We identified the native restriction-modification system of *C. kluyveri* as a critical barrier to DNA transfer and overcame it by identifying and characterizing the crucial methyltransferase. To mimic the native DNA methylation pattern of *C. kluyveri*, we performed *in-vivo* methylation of the shuttle vector plasmid by expressing the methyltransferase in *Escherichia coli*, followed by DNA transfer *via* conjugation. After validating the genetic system, we demonstrated heterologous expression of different combinations of both NADH and NADPH-dependent alcohol dehydrogenases from *Clostridium acetobutylicum*. The expression of these genes was controlled by the P_{thl} promoter, which is commonly used in *Clostridia,* and the P_{adhE2} promoter, leading to *n*-butanol and *n*-hexanol production of the mutant strains. This genetic system for *C. kluyveri* will not only enable further research on the metabolism of this microbe but also enable more profound insights into ethanol-based chain elongation in general.

IMPORTANCE

Medium-chain carboxylates are required in various everyday products, including cosmetics, pharmaceuticals, and fragrances, and show a natural antimicrobial property. Further, they represent food additives and serve as chemical building blocks for several other compounds. Traditionally, these carboxylates are produced from fossil resources, contributing to increased greenhouse gas emissions. Alternatively, they are derived from animal- or plant-based fat (*e.g.*, coconut oil), which competes with agricultural land that is needed for food production. However, microbial chain elongation, which is a biotechnological approach relying on microbes, such as *Clostridium kluyveri*, is sustainable and a promising alternative to the conventional production of medium-chain carboxylates. Notably, it enables the use of industrial waste streams (*e.g.*, off-gases, carbohydrate-rich industrial waste) as substrates, making the process more environmentally friendly. By applying our genetic system for *C. kluyveri*, a better understanding of microbial chain elongation can be achieved, and potentially even an extension of its product portfolio.

INTRODUCTION

Despite advances in renewable energy technologies, modern society remains heavily dependent on fossil fuels. This continued reliance contributes considerably to greenhouse gas emissions, which in turn accelerate climate change and the depletion of natural resources (1). Therefore, moving toward a more circular economy represents a critical challenge for sustainability to ensure long-term resilience for future generations. A key aspect of this transformation is the development of sustainable alternatives for essential chemical building blocks, which are conventionally assembled using fossil resources.

The development of sustainable alternatives represents a central challenge for modern environmental biotechnology. In particular, chemical building blocks that are in high demand from industry include short- and medium-chain carboxylates (SCCs and MCCs, respectively) (2). Both SCCs and MCCs contain a carboxylic group; while SCCs have carbon chains with a length of two to five carbon atoms, MCCs consist of six to twelve carbons (3) and their monetary value increases with an increasing chain length (4). MCCs serve as versatile chemical compounds with numerous industrial applications. Beyond functioning as essential building blocks for bioplastics (5, 6) and potential biofuel precursors through Kolbe electrolysis (7-9), these compounds have considerable utility across multiple sectors. They are widely incorporated into cosmetics, pharmaceuticals (2), fragrances (10), and rubber production (11). Additionally, MCCs feature prominently in food products as flavor additives (11, 12) and in animal feed formulations, where they provide valuable antimicrobial properties (13, 14). Also, they serve as precursors for alcohols such as *n*-butanol and *n*-hexanol (15).

Conventionally, MCCs are produced from petrochemicals (16), leading to

environmental devastation and mobilization and release of carbon dioxide (CO_2). A

potential bio-alternative that is already applied on an industrial scale is the production

of MCCs from animal or plant fat, for example, coconut oil, palm kernel oil, and castor

oil (17, 18). However, the sustainability of this approach is contested because it

requires agricultural land, which is indispensable for growing food, especially given

the steadily increasing global population (19). Another promising biotechnological

alternative is microbial chain elongation to produce MCCs.

By using the reverse ß-oxidation pathway, microbes fuse a C2 compound to short-

chain carboxylates, resulting in a stepwise elongation of the carbon chain by two

carbon atoms at a time, respectively, up to a maximum chain length of six carbon

atoms (20). This elongation process can be achieved by reactor microbiomes using

an open culture of microbial consortia. The open cultures do not require sterile

treatment and are resilient to disturbances (21, 22). Another promising attempt, which

was demonstrated in several studies, is to couple syngas fermentation with chain

elongation by cultivating an acetogenic and a chain-elongating microbe in a co-

culture (23-26). First, the acetogenic microbe produces the intermediates ethanol and

acetate from a mixture of hydrogen, CO_2, and carbon monoxide. Second, the chain-

elongating microbe further elongates the intermediates to medium-chain carboxylates

via reverse ß-oxidation. Third, the acetogenic microbe reduces SCCs and MCCs into

their corresponding alcohol (*n*-butanol and *n*-hexanol). Besides open- and co-

cultures, medium-chain carboxylate production performed by pure cultures offers

higher production rates (27).

Clostridium kluyveri has been extensively used to study the physiology and

biochemistry of ethanol-based chain elongation *via* the reverse ß-oxidation pathway

(28-31). Therefore, it serves as a model microbe for microbial chain elongation. To

date, no genetic system for this microbe has been published in a peer-reviewed

study. Here, we developed a genetic system for *C. kluyveri* by mimicking its native

methylation pattern on a shuttle vector plasmid to circumvent its restriction-

modification defense system. As a proof of principle, we introduced and

heterologously expressed alcohol dehydrogenases from *Clostridium acetobutylicum*

to enable *n*-butanol and *n*-hexanol production. By applying the findings from this

study, metabolic engineering of *C. kluyveri* can be used to increase its productivity,

expand its product portfolio, and obtain more profound insights into the reverse ß-

oxidation metabolism in general.

RESULTS

Plating optimization

To enable successful genetic modification, our initial focus was to optimize plating conditions for the reliable growth of single colonies, because a high plating efficiency increases the likelihood of isolating the correct clone. We systematically adjusted each variable to determine the optimal plating conditions. Of these, supplementing the DSMZ52 medium with a 10-fold concentration of vitamins and using cells in their early exponential growth phase had the most substantial positive impact on colony formation (**Table S1**). Other modifications, such as pour-plating with 0.8% Bacto agar, increasing the atmospheric pressure in the storage tanks to 0.6 bar, and placing a sterile filter paper (1 cm × 1 cm) soaked with ethanol in the lid of the Petri dish to ensure sufficient substrate supply, showed moderate positive effects (**Table S1**).

In contrast, using complex modified Reinforced Clostridial Medium (RCM) (32), which was supplemented with 20 mL/L ethanol and 2.5 g/L $NaHCO_3$ instead of DSMZ52 medium, and adding ethanol to the external plate atmosphere affected the plating efficiency negatively (**Table S1**). Further, using paper clips to increase ventilation resulted in a drastic reduction in colony formation (**Table S1**), most likely due to ethanol evaporation and plate desiccation from the enhanced airflow. Because hydrogen production of *C. kluyveri* resulted in large bubbles of the pour-plated medium, which complicated post-treatment, we decided to perform spread-plating whenever a minor reduction in plating efficiency was acceptable. For routine plating, we chose DSMZ52 medium with increased vitamin supplementation over RCM to reduce contamination risk and improve consistency. However, RCM medium was necessary during conjugation to support the growth of *E. coli*.

Antibiotic testing

Selection of cells carrying the desired plasmid requires two key factors: **1)** the sensitivity of the wild-type strain to the respective antibiotic, and **2)** the expression of an enzyme that ensures resistance to the antibiotic to which the wild-type strain is sensitive. Thiamphenicol resistance is a well-established marker in many clostridial strains, including strains closely related to *C. kluyveri* (33, 34). Therefore, we tested the sensitivity of the cells to thiamphenicol in liquid DSMZ52 medium and on plates and determined the minimum inhibitory concentration (MIC) after one week, which was 3 µg/mL (**Figure S1**). However, we chose to use a higher concentration of 5 µg/mL thiamphenicol in liquid medium to prevent the growth of wild-type cells due to spontaneous resistance or a reduction in antibiotic concentration caused by instability of the antibiotic at 37°C after prolonged incubation. Additionally, we used this concentration to select plasmid-carrying cells on solid media.

Identification of the crucial methylation pattern

A restriction assay of the pMTL83151 shuttle vector plasmid (35), using the cell lysate of *C. kluyveri*, showed digestion of the foreign DNA, producing fragment sizes of approximately 2 kbp and 0.5 kbp (**Figure 1A**). This result indicated that the native restriction and modification defense system of *C. kluyveri* (**Figure 1B**) is a critical barrier that must be overcome for successful genetic modification. PacBio sequencing of the native methylation pattern of the genomic DNA of *C. kluyveri* DSM555 revealed methylation at the following DNA sites: GATC, GWTAAT, CCAAG, CAAAAAT, and CCGG, suggesting the presence of multiple restriction-modification systems. *In-silico* analysis revealed that the observed restriction pattern of the unmethylated pMTL83151 corresponded to the CCGG as the critical methylation site, corresponding with the findings of Schneider, 2024 (36).

We identified a native methyltransferase in the REBASE database (37) that

methylates CCGG sites and is encoded by the genome of *C. kluyveri* (locus tag

CKL_2671). To counteract the restriction, we expressed this methyltransferase under

the control of the Isopropyl-ß-D-thiogalactopyranoside (IPTG)-inducible P*lac* promoter

on the plasmid pMeth (**Figure 2**) to mimic *C. kluyveri*-specific methylation of

pMTL83151. As a result, the restriction assay testing CCGG-methylated pMTL83151

showed reduced plasmid digestion. Next, we optimized the methyltransferase

expression by testing different IPTG concentrations (10-1000 mM) for induction in

E. coli TOP10. In a final restriction digest (**Figure 1C**), we only observed minimal

digestion of the shuttle vector after 24 h, indicating that the restriction sites of the

foreign DNA were protected by methylation (**Figure 1D**). The lowest restriction

activity occurred at IPTG induction concentrations between 200 mM and 500 mM

right after inoculation. Based on these results, we established 500 mM IPTG as the

standard concentration for inducing the expression of the methyltransferase.

Triparental conjugational DNA transfer

The *E. coli* TOP10 strain, harboring both pMTL83151 and pMeth plasmids, exhibited

reduced growth rates. This growth inhibition was likely attributable to the metabolic

burden that was imposed by maintaining resistance to two antibiotics simultaneously

(chloramphenicol for pMTL83151 and ampicillin for pMeth), combined with the

additional cellular stress of protein expression that is induced by IPTG treatment. To

avoid additional burden on this strain by introducing a third plasmid to enable

conjugation, we instead performed triparental mating for shuttle vector transfer into

the *C. kluyveri* recipient using a modified transformation protocol (38). We retained

E. coli TOP10, containing both pMeth and the shuttle vector plasmid, as the donor

strain and *E. coli* HB101, carrying pRK2013, as the helper strain (39). To support the

growth of the *E. coli* strains and *C. kluyveri* on the same plate, we tested the growth

of the two *E. coli* strains on RCM plates supplemented with varying ethanol

concentrations (86-342 mM, 0.5-2.0% [v/v]) under a N_2/CO_2 atmosphere. Both *E. coli*

strains exhibited normal growth up to 257 mM ethanol. We decided to use the highest

ethanol concentration because it is the closest to the ethanol level that we use to

cultivate *C. kluyveri* (342 mM), making it the optimum condition for co-cultivating

these three strains.

We evaluated different mating durations (24, 48, 72, 96, and 120 h) by selecting

potentially transformed cells on plates containing thiamphenicol and trimethoprim.

After 24 h, individual colonies were visible (**Figure S2A**). However, when extending

mating times from 48 h to 120 h, individual colonies became indistinguishable on

selective plates, requiring re-streaking to isolate single colonies (**Figure S2B**). In the

next step, we successfully verified the presence of the shuttle vector plasmid inside

C. kluyveri cells *via* colony PCR (**Figure 3A**), and we sequenced the 16S rRNA gene

to verify pure cultures (**Figure 3B**). From these findings, we conclude that longer

mating times between 48 and 120 h enhance or at least do not adversely affect the

transformation efficiency of the cells.

Heterologous gene expression as proof of concept

To expand the product portfolio of *C. kluyveri*, we aimed to enable the conversion of

n-butyrate and *n*-caproate into *n*-butanol and *n*-hexanol. This was achieved by

introducing the *adhE2* gene from *C. acetobutylicum* into the shuttle vector plasmid

pMTL83151 under the control of its native promoter, which is P_{adhE2}. *adhE2* encodes

a bifunctional alcohol dehydrogenase, which catalyzes both the conversion of acyl-

CoA to the corresponding aldehyde in the first step and the reduction of the aldehyde

to the respective alcohol in the second step (40). After successful transformation of

the plasmid into *C. kluyveri* cells, we observed the production of alcohols

corresponding to the chain length of the carboxylates produced. However, preliminary

measurements of the alcohols produced by *C. kluyveri* pPadhE2_adhE2 revealed

relatively low alcohol concentrations (*n*-butanol: 1.87 mM, *n*-hexanol: 3.15 mM). To

increase alcohol production, we tested two conditions: **1)** increasing the expression of

the *adhE2* gene by using a stronger promoter (41); and **2)** introducing an NADPH-

dependent alcohol dehydrogenase because NADPH represents a more powerful co-

factor than NADH (42) and the use of NADH by AdhE2 potentially lowers the

NADH/NAD+ ratio too much. Therefore, using NADPH as an alternative co-factor

could be advantageous for *n*-butanol and *n*-hexanol production

To address these limitations, we enhanced *adhE2* expression by replacing the native

promoter with the reportedly stronger P$_{thl}$ promoter from *C. acetobutylicum* (43, 44),

which resulted in higher *n*-butanol and *n*-hexanol concentrations (*n*-butanol:

2.73 mM, *n*-hexanol: 3.25 mM) compared to the P$_{adhE2}$ promoter. To further increase

alcohol production, we introduced the *bdhB* gene from *C. acetobutylicum*, encoding

butanol dehydrogenase B. Unlike AdhE2, BdhB uses NADPH as a co-factor for

converting aldehydes to alcohols (45, 46). Since BdhB exclusively catalyzes the

second step from the aldehyde to the alcohol (47), we retained AdhE2 to ensure acyl-

CoA conversion to the corresponding aldehyde. Thus, we tested three constructs on

the shuttle vector plasmid pMTL83151: **1)** pPadhE2_adhE2; **2)** pPthl_adhE2; and

3) pPthl_adhE2_bdhB and used the pMTL83151 vector as the negative control in a

growth experiment (**Figure 4**). We successfully transformed *C. kluyveri* cells with the

three constructs and the negative control and evaluated the resulting phenotypes.

Notably, during the conjugation process, we observed that DNA transfer for the

plasmid encoding both *adhE2* and *bdhB* was only successful after 72 h of mating.

n-Butanol and *n*-hexanol production of the different strains

In bottle experiments, *C. kluyveri* pPadhE2_adhE2, *C. kluyveri* pPthl_adhE2_bdhB, and the negative control exhibited comparable growth with doubling times between 4.5 and 6 h (**Figure 4A**). In comparison, the three replicates of *C. kluyveri* pPthl_adhE2 started to grow at different times after inoculation, thus, leading to higher standard deviations, not only for the OD_{600} but also for substrate and product concentrations for this strain. However, the growth rates were similar to the other strains tested (data not shown). Unfortunately, several attempts to repeat the growth experiment failed due to a high mutation rate in the promoter region of *adhE2* in *C. kluyveri* pPthl_adhE2. Average *n*-butanol and *n*-hexanol production was first observed after 33 h for *C. kluyveri* pPthl_adhE2_bdhB and after 38 h for *C. kluyveri* pPadhE2_adhE2 (**Figures 4B,C**). Due to the extended lag phase and high standard deviations (up to 54% for *n*-butanol and 173% for *n*-hexanol), *C. kluyveri* pPthl_adhE2 was the last strain starting to produce *n*-butanol and *n*-hexanol (first production observed between 61 h and 81 h, data not shown), resulting in a slower average alcohol production than the other replicates.

Regarding *n*-butanol production, *C. kluyveri* pPthl_adhE2_bdhB reached a maximum concentration of 4.2 ± 0.0 mM, whereas *C. kluyveri* pPadhE2_adhE2 and *C. kluyveri* pPthl_adhE2 exhibited similar *n*-butanol production levels (**Figure 4B**). In line with this observation, *C. kluyveri* pPthl_adhE2_bdhB also achieved the highest *n*-hexanol concentration (3.7 ± 0.6 mM), followed by *C. kluyveri* pPthl_adhE2 (2.3 ± 0.4 mM). *C. kluyveri* pPadhE2_adhE2 produced the lowest amount of *n*-hexanol (2.2 ± 0.4 mM) (**Figure 4C**). The difference in *n*-hexanol production between *C. kluyveri* pPthl_adhE2 and *C. kluyveri* pPadhE2_adhE2 likely would have been notably higher if *C. kluyveri* pPthl_adhE2 had exhibited more consistent growth

across the replicates. Also, a longer experiment duration might have further

highlighted a potential disparity.

Interestingly, both strains carrying P_{thl} started producing the corresponding alcohols

at lower n-butyrate and n-caproate concentrations (28.6 ± 10.3 mM and

15.6 ± 12.1 mM, and 34.2 ± 1.4 mM and 21.7 ± 2.3 mM, respectively)

(**Figures 4B,C)**. In contrast, *C. kluyveri* pPadhE2_adhE2 started n-butanol and n-

hexanol production at 42.6 ± 2.6 mM n-butyrate and 38.7 ± 5.3 mM n-caproate

(**Figures 4B,C** and **Figures S3C,D)**. Remarkably, *C. kluyveri* pPthl_adhE2_bdhB

converted most acyl-CoA into the corresponding alcohol, formed more n-butyrate

than the other strains, and reached the highest OD_{600} (**Figures 4A** and

Figures S3C,D). It produced 30% less n-caproate than the other alcohol-producing

strains and approximately 35% less than the negative control (**Figure S3D)**. Due to

the lower n-caproate production, its conversion rate for substrate carbon into product

carbon (n-butyrate, n-caproate, n-butanol, and n-hexanol) only reached 53%,

whereas the other strains converted 62-67% of it (**Figures 4B,C and Figure S3**).

Further, *C. kluyveri* pPthl_adhE2_bdhB showed a higher acetate-to-ethanol ratio at

the end of the experiment (0.16) than the other strains (0.10-0.12) (**Figures S3A,B)**.

DISCUSSION

Our results show that: **1)** adapting the growth conditions for *C. kluyveri* on solid DSMZ52 medium; **2)** determining the MIC of thiamphenicol; and **3)** counteracting restriction modification barriers enabled successful DNA transfer of a shuttle vector plasmid into *C. kluyveri* cells *via* conjugation after 24 h of mating. However, extending the mating time to 48 h improved the transformation efficiency. Notably, transferring the plasmid pPthl_adhE2_bdhB (8.234 kbp) was only successful after 72 h, suggesting that shuttle vectors with critical genes or an increased size may require extended mating times. This extension has been shown to improve conjugation efficiencies overall (48).

All generated strains, but the negative control, showed plasmid-based gene expression of the *adhE2* and *bdhB* genes from *C. acetobutylicum*. This led to phenotypically modified *C. kluyveri* strains that were capable of producing *n*-butanol and *n*-hexanol. In line with our expectations, both replacing P_{adhE2} with the reportedly stronger P_{thl} promoter and introducing a second, NADPH-dependent butanol dehydrogenase resulted in enhanced *n*-butanol and *n*-hexanol production. The clear variability observed in the lag phase duration among the replicates of *C. kluyveri* pPthl_adhE2 distorted the calculated mean values. This statistical distortion resulted in an artificial lowering of the estimated growth rate because the extended lag phases of some replicates disproportionately influenced the overall growth calculations. This bias not only lowers the slope of the OD_{600} but also needs to be considered for the graphs representing the substrates and products. A potential explanation for the extended lag phase is cell stress caused by the P_{thl} promoter, which is stronger than P_{adhE2}. Because AdhE2 catalyzes both reactions of acyl-CoA to aldehyde and aldehyde to alcohol, it may have a higher affinity towards the first reaction in

C. kluyveri. Hence, the promoter exchange could lead to aldehyde accumulation,

which is known to be toxic (49, 50), causing cell stress and impaired cells.

The batch experiment provided compelling evidence supporting the hypothesis that

the combination of the P$_{thl}$ promoter and solely the *adhE2* gene exhibits toxicity. This

toxicity was demonstrated conclusively by metabolic impairment: across three

independent growth experiments, none of the *C. kluyveri* pPthl_adhE2 replicates

produced detectable levels of either *n*-butanol or *n*-hexanol. In contrast, only one

experiment resulted in the successful production of the alcohols. Sequencing of the

non-alcohol-producing cultures revealed that all contained a deletion in the P$_{thl}$ region

(**Figure S4A**), suggesting a selection pressure against the expression of *adhE2*.

Additionally, sequencing *C. kluyveri* pPthl_adhE2 cultures, which produced *n*-butanol

and *n*-hexanol, revealed a less defined baseline in the promoter region (**Figure S4B**),

indicating that a fraction of the cells had already mutated. Given these findings, we

cannot rule out that an unmutated pure culture of *C. kluyveri* pPthl_adhE2 could have

performed better. Nonetheless, while using a stronger promoter enhanced the

production of *n*-hexanol, adding the butanol dehydrogenase presumably helped

accelerate aldehyde removal. This prevented toxic accumulation and resulted in a

stable, mutation-free strain.

Furthermore, we observed reduced formation of *n*-caproate by *C. kluyveri*

pPthl_adhE2_bdhB despite reaching the highest OD$_{600}$ of all strains. This might be

due to a more balanced use of the co-factors NADH and NADPH. Both co-factors are

utilized to prevent excessive depletion of NADH, which would result in a reduced

NADH/NAD$^+$ ratio. This results in a higher NADH reducing capability and increased

alcohol production (51). Additionally, improved recycling of the co-factors could

explain the lower final concentration of *n*-caproate. Since NADH and NADPH are

used as electron donors for alcohol production, they are less available for other

processes, such as the formation of carboxylates (28). As a result, these reactions

occur less frequently, leading to a decrease in *n*-caproate production.

The genetic system demonstrated in this study enabled plasmid-based gene

expression, leading to heterologous production of *n*-butanol and *n*-hexanol by

C. kluyveri. Besides expanding the product portfolio of this microbe, our research

represents the first step towards developing a broader toolkit for *C. kluyveri*, including

genome editing techniques (*e.g.*, homologous recombination, CRISPR-based

methods). Consequently, the genetic accessibility of the model microbe for chain

elongation is a precondition for further research on the reverse ß-oxidation pathway,

which holds strong industrial potential. Therefore, gaining a deeper understanding of

this pathway, both on a genetic level and in terms of the enzymes involved in their

central energy metabolism, is crucial to enhancing its productivity.

MATERIALS AND METHODS

Microbial strains, media, and cultivation

C. kluyveri DSM555 was obtained from the DSMZ (Braunschweig, Germany) and used as the wild-type strain. Unless stated otherwise, all *C. kluyveri* strains in this study were cultivated in DSMZ52 medium for which 2.5 g/L $NaHCO_3$ was used to buffer the medium instead of Na_2CO_3. Cultures were grown in 100-mL medium with a headspace of 140 mL under an N_2/CO_2 (80/20) atmosphere. Both liquid cultures and plates (containing 0.8% [v/v] Bacto agar) were incubated at 37°C until visible growth was observed. When required, the medium was supplemented with 5 µg/mL thiamphenicol and 10 µg/mL trimethoprim for selection. All genetic manipulations of *C. kluyveri* were conducted under strict anoxic conditions.

Unless otherwise noted, *E. coli* HB101 pRK2013 and *E. coli* TOP10 strains were grown in liquid Luria-Bertani (LB) medium or on LB plates containing 1.5% agar. For plasmid selection and maintenance, the medium was supplemented with 100 µg/mL ampicillin (for liquid cultures), 100 µg/mL carbenicillin (for plates), 30 µg/mL chloramphenicol, or 30 µg/mL kanamycin. IPTG was added at a concentration of 500 mM to induce the expression of the methyltransferase encoded on pMeth.

Optimization of the plating efficiency

The variables (**Table S1**) were systematically adjusted one at a time to determine the optimal plating conditions, while all other conditions remained identical. The resulting outcomes were compared to the previously identified best conditions. Both the modified condition and the previously best-performing combination of conditions were tested using the same culture conditions and inoculation volume (4% [v/v]) with simultaneous incubation. The inoculum was serially diluted by increasing the dilution

factor by a magnitude of 10, reaching a final dilution of 1:100,000, and each dilution

was plated in duplicates for both conditions. Colony counts were performed by

capturing images using the UVP GelStudio system (Analytic Jena, Jena, Germany)

and analyzed with the integrated colony-count software. The plating conditions

yielding the highest colony count were determined to be the updated optimal

combination, and the following variable was tested accordingly.

Restriction digest

10 mL of a late exponential phase *C. kluyveri* culture were harvested, washed twice

with phosphate-buffered saline (PBS) buffer (137 mM sodium chloride, 2.7 mM

potassium chloride, 10 mM sodium phosphate dibasic, and p1.8 mM potassium

phosphate monobasic), and resuspended in 300 μL PBS buffer. Cells were lysed

using bead-beating (30 sec, 6.5 m/s; FastPrep-24™, MP BioMedicals, Santa Ana,

USA). The lysing procedure was performed four times, and in between, the cells

were kept on ice for 3 min. The lysis was centrifuged at 2000 × *g* for 20 min at 4°C.

To assess the sensitivity of the respective plasmid to the native restriction enzymes of

C. kluyveri, 5 μL of the supernatant were added to 13 μL plasmid (200 ng) and 2 μL

rCutSmart™ buffer (New England Biolabs, Ipswich, Massachusetts, USA). The

prepared reaction mix was stored at 37°C for 1 to 24 h. Results were analyzed *via*

gel electrophoresis and potential corresponding restriction sites were analyzed *in*

silico with SnapGene (GSL Biotech LLC, San Diego, USA).

PacBio sequencing

High molecular weight DNA (25 ng/μL) was isolated with the NucleoSpin Microbial

DNA Mini kit for DNA (Macherey-Nagel, Düren, Germany) and sheared using 35 kb

settings with a Megaruptor 2 instrument (Diagenode, SA, Lüttich, Belgium). Average

insert sizes were approximately 10 kb. Sheared fragments were used to prepare

libraries with the HiFi SMRTbell Express Template Prep Kit 2.0 (Pacific Biosciences,

Menlo Park, California, USA). The libraries were size-selected with a BluePippin

system (SageScience, Beverly, USA) with 17 kb cutoff in a 0.75% DF Marker

S1 High-Pass 6-10 kb vs3 gel cassette (Biozym, Hamburg, Germany). The library

was sequenced with sequencing primer v2 (Pacific Biosciences, #101-847-900) and

4 h of pre-extension time on a single SMRT Cell with the Sequel II system using

Binding Kit 2.0.

Methylation profiling

Reads were demultiplexed using Lima (52) and converted into High-Fidelity (HiFi)

reads retaining kinetic information with Circular Consensus Sequencing (53). BAM

files were converted to FASTQ with bam2fastx (54). Reads shorter than 1000 bp and

the lowest 5% of reads based on quality scores were discarded using Filtlong (55).

Trycycler (56) was used to subsample reads and generate a consensus sequence

from 12 assemblies independently generated with Flye (57), Miniasm (58), and

Raven (59). HiFi reads were then aligned with the consensus assembly using pbmm2

(60), and ipdSummary (61) was used to detect and classify DNA base-modifications

from the kinetic signatures. MotifMaker (62) was used to identify statistically

significant methylated motifs, with a minimum score threshold of 60. Motifs were

retained if they were identified for 90% or more possible sites in the genome. The

native methyltransferase of *C. kluyveri,* responsible for methylating the critical CCGG

sites, was identified using the REBASE database (37).

Plasmid construction

Plasmids (**Table S2**) were constructed with *E. coli* TOP10 and the respective primers (**Table S3)**. The pUC19 vector (63) served as the backbone for the methylation plasmid pMeth. The gene encoding the critical methyltransferase (CKL_2671) from *C. kluyveri* was PCR amplified (Q5 polymerase, New England Biolabs (NEB), Ipswich, MA, USA) from genomic DNA (NucleoSpin, Microbial DNA, Macherey-Nagel) and inserted into the multiple cloning site *via* Gibson Assembly (NEBuilder, NEB). The lambda t1 terminator was introduced downstream of the CKL_2671 gene *via* PCR, amplifying the plasmid with overlapping primers and transferring the linear DNA inside *E. coli*. The same method was applied to remove the nucleotide sequence between the promoter and the start codon of CKL_2671 to utilize the promoter present in the backbone (P_{lac}). In the final step, the origin of replication (ORI) was exchanged from the pSC101 ORI to the p15a ORI (64) by Gibson Assembly (NEBuilder, NEB) to ensure a low copy number and compatibility with the shuttle vector plasmids for *C. kluyveri*. For *n*-butanol and *n*-hexanol production, pMTL83151 (35) served as the backbone for all shuttle vector plasmids. The DNA fragments *adhE2*, *bdhB*, P_{adhE2}, and P_{thl} from *C. acetobutylicum* were PCR amplified (Q5 polymerase, NEB) from genomic DNA, and the respective fragments were inserted into the plasmids *via* Gibson Assembly (NEBuilder, NEB).

Triparental conjugational DNA transfer

We inoculated LB medium supplemented with ampicillin, chloramphenicol, and IPTG (to induce the expression of the methyltransferase) with the donor strain *E. coli* TOP10, carrying the desired plasmid and incubated the culture overnight. When both the donor strain and the helper strain *E. coli* HB101 pRK2013, which was grown in LB medium supplemented with 30 µL kanamycin, reached an OD_{600} of 0.6-0.8, 1 mL

of each culture was harvested by centrifugation (17,900 $\times$ g, 3 min). The resulting

pellets were suspended and mixed in PBS buffer. We again centrifuged the *E. coli*

mixture (17,900 $\times$ g, 3 min) and transferred the received cell pellet into the anoxic

chamber. Subsequently, 2 mL of mid-exponential *C. kluyveri* wild-type culture

(OD_{600}: 0.4-0.7) was harvested *via* centrifugation (9.800 $\times$ g, 4 min) and resuspended

in 50 µL RCM under an N_2 atmosphere. This cell suspension was used to resuspend

the *E. coli* cell pellet. The cell mixture was then spot-plated on RCM plates (1.5%

[v/v] agar) containing 500 mM IPTG and 257 mM ethanol (1.5% [v/v]). Plates were

incubated at 37°C in an N_2/CO_2 (80/20) atmosphere for up to 120 h. After mating,

cells from the mating spots were transferred *via* striking onto DSMZ52 plates

containing thiamphenicol to select for the respective plasmid. After cells had grown

on the first selective plate, they were streaked onto a thiamphenicol and

trimethoprim-containing DSMZ52 plate to select against *E. coli* strains. The resulting

colonies were then further analyzed.

Molecular methods for analysis of genetically modified *C. kluyveri*

Direct colony PCR was conducted on both colonies and liquid cultures to confirm the

presence of the respective plasmid in *C. kluyveri* cells. Individual colonies were

picked and resuspended in 25 µL of 10 mM NaOH to extract the DNA template. For

PCR performed on liquid cultures, 100 µL of culture was sampled. In both cases, the

cells were lysed at 98°C for 10 min using a ThermoMixerC (Eppendorf, Hamburg,

Germany). For the PCR reactions, 1 µL of the lysate was utilized in two separate

10 µL PCR reactions (PhirePlant (ThermoScientific™, Waltham, USA)) – one using

plasmid-specific primers and the other using *C. kluyveri*-specific primers. The results

of the reactions were analyzed *via* gel electrophoresis. Additionally, a 30 µL PCR

reaction targeting the 16S rRNA gene was performed using 3 µL of the lysate as a

template. The resulting PCR product was purified using the QIAquick PCR

Purification Kit (QIAGEN, Venlo, Netherlands) and subsequently analyzed *via* Sanger

sequencing (GENEWIZ, Leipzig, Germany) to verify a pure culture of *C. kluyveri*.

Analysis of compound concentration

To determine the concentrations of carboxylates (*e.g.*, acetate, *n*-butyrate, *n*-

caproate) and alcohols (*n*-butanol and *n*-hexanol), an Agilent 7890B gas

chromatograph (Agilent Technologies, Inc., Santa Clara, CA, USA) that were

equipped with a DB-Fatwax UI capillary column (30 m x 0.25 mm x 0.25 μm) and

flame ionization detector (FID) was used. Hydrogen served as the carrier gas, and

the temperature program consisted of an initial temperature of 40°C for 3 min,

ramped at 12°C/min to 160°C, followed by a further ramp of 20°C/min to 230°C with a

5-min hold. Ethyllactate was used as an internal standard at a concentration of 2 mM

for quantifying carboxylates. The sampling volume was 1 mL of culture that was

centrifuged (17,900 *g*, 3 min), and from which we stored the supernatant at -20°C

until further analysis. Before measuring, the samples were thawed and filtered

through a 0.2 μm polyvinylidene difluoride (PVDF) sterilefilter. For the analysis,

samples were diluted at ratios of 1:2 for *n*-butanol and *n*-hexanol and 1:50 for MCCs

measurements.

For ethanol concentration measurements, undiluted samples were analyzed using

the LC-20 high-performance liquid chromatography (HPLC) system (Shimadzu,

Nishinokyo Kuwabara-cho, Japan). The system was equipped with an Aminex-HPX-

87H column, and 5 mM sulfuric acid was utilized as the mobile phase at a flow rate of

0.6 mL/min (LC-20AD). The oven temperature was maintained at 65°C (CTO-20AC),

and the samples were cooled at 15°C in the autosampler unit.

ACKNOWLEDGEMENTS

We want to express our sincere gratitude to Silke Dauser and Heike Budde for performing the library preparation and PacBio sequencing. Further, the authors want to thank José Antonio Velázquez Gómez for supporting the HPLC analysis. Funding sources were the Max Planck Institute to L.T.A. as a Max Planck Fellow, the Alexander von Humboldt Foundation through the Alexander von Humboldt Professorship (L.T.A.), the Novo Nordisk Foundation through the CO_2 Research Center (CORC) with grant number NNF21SA0072700 (L.T.A.), the Deutsche Forschungsgesellschaft (DFG, German Research Foundation) through the Leibniz Prize (L.T.A.), and the Deutsche Forschungsgemeinschaft under Germany's Excellence Strategy – EXC 2124 – 390838134 (L.T.A.).

L.T.A. initiated the work. C.S., S.T.B., and L.T.A. designed the experiments. C.S., S.T.B., J.W.M., and K.G. performed laboratory experiments and analyzed the data. L.T.A. and S.T.B. supervised the project. C.S., L.T.A., J.W.M., and S.T.B. wrote the manuscript, while all edited the paper and approved the final version.

We declare no conflict of interest.

REFERENCES

1. Masson-Delmotte V, Zhai P, Pirani A, Connors SL, Péan C, Berger S, Caud N, Chen Y, Goldfarb L, Gomis MI, Huang M, Leitzell K, Lonnoy E, Matthews JBR, Waterfield T, Yelekci O, R Y, Zhou B. 2022. Climate change 2021: the physical science basis, contribution of working group I to the sixth assessment report of the interfovernmetal panel on climate change. Cambridge University Press.

2. Angenent LT, Richter H, Buckel W, Spirito CM, Steinbusch KJ, Plugge CM, Strik DP, Grootscholten TI, Buisman CJ, Hamelers HV. 2016. Chain elongation with reactor microbiomes: open-culture biotechnology to produce biochemicals. Environmental Science & Technology 50:2796-2810.

3. Steinbusch KJ, Hamelers HV, Plugge CM, Buisman CJ. 2011. Biological formation of caproate and caprylate from acetate: fuel and chemical production from low grade biomass. Energy & Environmental Science 4:216-224.

4. Wang J, Yin Y. 2022. Biological production of medium-chain carboxylates through chain elongation: an overview. Biotechnology Advances 55:107882.

5. Marang L, Jiang Y, van Loosdrecht MC, Kleerebezem R. 2013. Butyrate as preferred substrate for polyhydroxybutyrate production. Bioresource Technology 142:232-239.

6. Szacherska K, Oleskowicz-Popiel P, Ciesielski S, Mozejko-Ciesielska J. 2021. Volatile fatty acids as carbon sources for polyhydroxyalkanoates production. Polymers 13:321.

7. de Araújo Cavalcante W, Leitão RC, Gehring TA, Angenent LT, Santaella ST. 2017. Anaerobic fermentation for *n*-caproic acid production: A review. Process Biochemistry 54:106-119.

8. Palkovits S, Palkovits R. 2019. The role of electrochemistry in future dynamic bio‐refineries: a focus on (non‐) kolbe electrolysis. Chemie Ingenieur Technik 91:699-706.

9. Harvey BG, Meylemans HA. 2014. 1-Hexene: a renewable C6 platform for full-performance jet and diesel fuels. Green Chemistry 16:770-776.

10. Gupta C, Prakash D, Gupta S. 2015. A biotechnological approach to microbial based perfumes and flavours. Journal of Microbiological Methods. doi.org/10.15406/jmen.20150100034.

11. Jeon BS, Moon C, Kim B-C, Kim H, Um Y, Sang B-I. 2013. *In situ* extractive fermentation for the production of hexanoic acid from galactitol by *Clostridium* sp. BS-1. Enzyme and Microbial Technology 53:143-151.

12. Serhan M, Mattar J, Debs L. 2016. Concentrated yogurt (labneh) made of a mixture of goats' and cows' milk: Physicochemical, microbiological and sensory analysis. Small Ruminant Research 138:46-52.

13. Van Immerseel F, De Buck J, Boyen F, Bohez L, Pasmans F, Volf J, Sevcik M, Rychlik I, Haesebrouck F, Ducatelle R. 2004. Medium-chain fatty acids decrease colonization and invasion through *hilA* suppression shortly after infection of chickens with *Salmonella Enterica* serovar Enteritidis. Applied and Environmental Microbiology 70:3582-3587.

14. Huang CB, Alimova Y, Myers TM, Ebersole JL. 2011. Short-and medium-chain fatty acids exhibit antimicrobial activity for oral microorganisms. Archives of Oral Biology 56:650-654.

15. Zhang J, Taylor S, Wang Y. 2016. Effects of end products on fermentation
profiles in *Clostridium carboxidivorans* P7 for syngas fermentation.
Bioresource Technology 218:1055-1063.
16. Liu P, Jarboe LR. 2012. Metabolic engineering of biocatalysts for carboxylic
acids production. Computational and Structural Biotechnology Journal
doi.org/10.5936/csbj.201210011
17. Roghair M, Hoogstad T, Strik DP, Plugge CM, Timmers PH, Weusthuis RA,
Bruins ME, Buisman CJ. 2018. Controlling ethanol use in chain elongation by
CO_2 loading rate. Environmental Science & Technology 52:1496-1505.
18. Gervajio GC. 2005. Fatty acids and derivatives from coconut oil. In Bailey's
industrial oil and fat products. John Wiley & Sons, Inc: Hooboken, NJ, USA,
2005. ISBN 9780471678496 6:1-56.
19. Schmitz LM, Kreitli N, Obermaier L, Weber N, Rychlik M, Angenent LT. 2024.
Power-to-vitamins: producing folate (vitamin B9) from renewable electric
power and CO_2 with a microbial protein system. Trends in Biotechnology
42:1691-1714.
20. Fernández-Blanco C, Robles-Iglesias R, Naveira-Pazos C, Veiga MC, Kennes
C. 2023. Production of biofuels from C1-gases with *Clostridium* and related
bacteria—recent advances. Microbial Biotechnology 16:726-741.
21. Werner JJ, Knights D, Garcia ML, Scalfone NB, Smith S, Yarasheski K,
Cummings TA, Beers AR, Knight R, Angenent LT. 2011. Bacterial community
structures are unique and resilient in full-scale bioenergy systems.
Proceedings of the National Academy of Sciences 108:4158-4163.
22. Agler MT, Spirito CM, Usack JG, Werner JJ, Angenent LT. 2012. Chain
elongation with reactor microbiomes: upgrading dilute ethanol to medium-
chain carboxylates. Energy & Environmental Science 5:8189-8192.
23. Richter H, Molitor B, Diender M, Sousa DZ, Angenent LT. 2016. A narrow pH
range supports butanol, hexanol, and octanol production from syngas in a
continuous co-culture of *Clostridium ljungdahlii* and *Clostridium kluyveri* with
in-line product extraction. Frontiers in Microbiology 7:1773.
24. Diender M, Stams AJ, Sousa DZ. 2016. Production of medium-chain fatty
acids and higher alcohols by a synthetic co-culture grown on carbon monoxide
or syngas. Biotechnology for Biofuels 9:1-11.
25. Schneider M, Bäumler M, Lee NM, Weuster-Botz D, Ehrenreich A, Liebl W.
2021. Monitoring co-cultures of *Clostridium carboxidivorans* and *Clostridium*
kluyveri by fluorescence *in situ* hybridization with specific 23S rRNA
oligonucleotide probes. Systematic and Applied Microbiology 44:126271.
26. Bäumler M, Schneider M, Ehrenreich A, Liebl W, Weuster-Botz D. 2021.
Synthetic co-culture of autotrophic *Clostridium carboxidivorans* and chain
elongating *Clostridium kluyveri* monitored by flow cytometry. Microbial
Biotechnology 15:1471-1485.
27. Gildemyn S, Molitor B, Usack JG, Nguyen M, Rabaey K, Angenent LT. 2017.
Upgrading syngas fermentation effluent using *Clostridium kluyveri* in a
continuous fermentation. Biotechnology for Biofuels 10:1-15.
28. Thauer RK, Jungermann K, Decker K. 1977. Energy conservation in
chemotrophic anaerobic bacteria. Bacteriological Reviews 41:100-180.
29. Thauer R, Jungermann K, Henninger H, Wenning J, Decker K. 1968. The
energy metabolism of *Clostridium kluyveri*. European Journal of Biochemistry
4:173-180.
30. Li F, Hinderberger J, Seedorf H, Zhang J, Buckel W, Thauer RK. 2008.
Coupled ferredoxin and crotonyl coenzyme A (CoA) reduction with NADH

catalyzed by the butyryl-CoA dehydrogenase/Etf complex from *Clostridium*
kluyveri. Journal of Bacteriology 190:843-850.

31. Scherf U, Söhling B, Gottschalk G, Linder D, Buckel W. 1994. Succinate-
    ethanol fermentation in *Clostridium kluyveri*: purification and characterisation
    of 4-hydroxybutyryl-CoA dehydratase/vinylacetyl-CoA Δ3-Δ2-isomerase.
    Archives of Microbiology 161:239-245.

32. Molitor B, Kirchner K, Henrich AW, Schmitz S, Rosenbaum MA. 2016.
    Expanding the molecular toolkit for the homoacetogen *Clostridium ljungdahlii*.
    Scientific Reports 6:1-10.

33. Kuehne SA, Heap JT, Cooksley CM, Cartman ST, Minton NP. 2011. ClosTron-
    mediated engineering of *Clostridium*, p 389-407, Strain Engineering. Springer.

34. Klask C-M, Jäger B, Casini I, Angenent LT, Molitor B. 2022. Genetic evidence
    reveals the indispensable role of the *rseC* gene for autotrophy and the
    importance of a functional electron balance for nitrate reduction in *Clostridium*
    ljungdahlii. Frontiers in Microbiology 13:887578.

35. Heap JT, Pennington OJ, Cartman ST, Minton NP. 2009. A modular system for
    Clostridium shuttle plasmids. Journal of Microbiological Methods 78:79-85.

36. Schneider MI. 2024. Entwicklung von molekularbiologischen Werkzeugen für
    den sekundären Gärer *Clostridium kluyveri*. Technische Universität München.

37. Roberts RJ, Vincze T, Posfai J, Macelis D. 2015. REBASE—a database for
    DNA restriction and modification: enzymes, genes and genomes. Nucleic
    Acids Research 43:D298-D299.

38. Lesiak JM, Liebl W, Ehrenreich A. 2014. Development of an *in-vivo*
    methylation system for the solventogen *Clostridium saccharobutylicum* NCP
    262 and analysis of two endonuclease mutants. Journal of Biotechnology
    188:97-99.

39. Ditta G, Stanfield S, Corbin D, Helinski DR. 1980. Broad host range DNA
    cloning system for gram-negative bacteria: construction of a gene bank of
    Rhizobium meliloti. Proceedings of the National Academy of Sciences
    77:7347-7351.

40. Fontaine L, Meynial-Salles I, Girbal L, Yang X, Croux C, Soucaille P. 2002.
    Molecular characterization and transcriptional analysis of *adhE2*, the gene
    encoding the NADH-dependent aldehyde/alcohol dehydrogenase responsible
    for butanol production in alcohologenic cultures of *Clostridium acetobutylicum*
    ATCC 824. Journal of Bacteriology 184:821-830.

41. Nevoigt E, Kohnke J, Fischer CR, Alper H, Stahl U, Stephanopoulos G. 2006.
    Engineering of promoter replacement cassettes for fine-tuning of gene
    expression in *Saccharomyces cerevisiae*. Applied and Environmental
    Microbiology 72:5266-5273.

42. Zhou Z, Jing Y, Wei S, Zhang Q, Peng S, An X, Li H. 2023. Enhancement of
    butanol production in *Clostridium acetobutylicum* SE25 through oxidation-
    reduction potential regulation and analysis of its metabolic mechanisms. Fuel
    331:125708.

43. Huang H, Chai C, Li N, Rowe P, Minton NP, Yang S, Jiang W, Gu Y. 2016.
    CRISPR/Cas9-based efficient genome editing in *Clostridium ljungdahlii*, an
    autotrophic gas-fermenting bacterium. ACS Synthetic Biology 5:1355-1361.

44. Hoffmeister S, Gerdom M, Bengelsdorf FR, Linder S, Flüchter S, Öztürk H,
    Blümke W, May A, Fischer R-J, Bahl H. 2016. Acetone production with
    metabolically engineered strains of *Acetobacterium woodii*. Metabolic
    Engineering 36:37-47.

45. Yoo M, Bestel-Corre G, Croux C, Riviere A, Meynial-Salles I, Soucaille P.
2015. A quantitative system-scale characterization of the metabolism of
Clostridium acetobutylicum. MBio 6:1-12.
46. Lee SY, Park JH, Jang SH, Nielsen LK, Kim J, Jung KS. 2008. Fermentative
butanol production by Clostridia. Biotechnology and Bioengineering 101:209-
228.
47. Foulquier C, Rivière A, Heulot M, Dos Reis S, Perdu C, Girbal L, Pinault M,
Dusséaux S, Yoo M, Soucaille P. 2022. Molecular characterization of the
missing electron pathways for butanol synthesis in *Clostridium acetobutylicum*.
Nature Communications 13:4691.
48. Huisman JS, Benz F, Duxbury SJ, de Visser JAG, Hall AR, Fischer EA,
Bonhoeffer S. 2022. Estimating plasmid conjugation rates: a new
computational tool and a critical comparison of methods. Plasmid 121:102627.
49. Kunjapur AM, Prather KL. 2015. Microbial engineering for aldehyde synthesis.
Applied and Environmental Microbiology 81:1892-1901.
50. Heldt D, Frank S, Seyedarabi A, Ladikis D, Parsons JB, Warren MJ, Pickersgill
RW. 2009. Structure of a trimeric bacterial microcompartment shell protein,
EtuB, associated with ethanol utilization in *Clostridium kluyveri*. Biochemical
Journal 423:199-207.
51. Cheng C, Bao T, Yang S-T. 2019. Engineering *Clostridium* for improved
solvent production: recent progress and perspective. Applied Microbiology and
Biotechnology 103:5549-5566.
52. Armin Töpfer JM, Mitchell Robert Volger, Chan Ye, K L Weber. Lima -
Demultiplex Barcoded PacBio Samples, v2.9.0. GitHub,
https://github.com/PacificBiosciences/barcoding.
53. Töpfer A. CCS: Generate Highly Accurate Single-Molecule Consensus Reads
(HiFi Reads), v6.4.0. GitHub, https://github.com/PacificBiosciences/ccs.
54. Armin Töpfer JM. PacBio BAM toolkit, v3.0.0. GitHub,
https://github.com/pacificbiosciences/pbtk/.
55. Ryan Wick PM. Quality filtering tool for long reads, v0.2.1. GitHub,
https://github.com/rrwick/Filtlong.
56. Wick RR, Judd LM, Cerdeira LT, Hawkey J, Méric G, Vezina B, Wyres KL, Holt
KE. 2021. Trycycler: consensus long-read assemblies for bacterial genomes.
Genome biology 22:1-17.
57. Kolmogorov M, Yuan J, Lin Y, Pevzner PA. 2019. Assembly of long, error-
prone reads using repeat graphs. Nature Biotechnology 37:540-546.
58. Heng Li SM, Riku Walve, Ryan Wick. Ultrafast de novo assembly for long
noisy reads (though having no censensus step), v0.3. GitHub,
https://github.com/lh3/miniasm.
59. Robert Vaser IV, Andrew Zhang, Thanh Lee. De novo genome assembler for
long uncorrected reads, v1.8.3. GitHub, https://github.com/lbcb-sci/raven.
60. Nat Echols DA, David Seifert, Lance Hepler, Derek W. Barnett, James Drake.
Tools for detecting DNA modifications from single molecule, real-time
sequencing data, v1.13.1. GitHub,
https://github.com/PacificBiosciences/kineticsTools.
61. Patrick Marks NE, David Alexander, Mengjuei Hsieh, David Seifert, Lance
Hepler, James Drake, Derek W. Barnett, Yuri D'Elia. Tools for detecting DNA
modifications from single molecule, real-time sequencing data, v3.0.0. GitHub,
https://github.com/PacificBiosciences/kineticsTools.
62. Maeder H. Find methylated motifs in prokaryotic genomes, v12. GitHub,
https://github.com/wangdi2014/MotifMaker.

63. Yanisch-Perron C, Vieira J, Messing J. 1985. Improved M13 phage cloning
vectors and host strains: nucleotide sequences of the M13mpl8 and pUC19
vectors. Gene 33:103-119.
64. Chang AC, Cohen SN. 1978. Construction and characterization of amplifiable
multicopy DNA cloning vehicles derived from the P15A cryptic miniplasmid.
Journal of Bacteriology 134:1141-1156.
730

Figure 1. Restriction modification system of C. kluyveri. *A) Restriction digest of the unmethylated pMTL83151 plasmid after 4 h. The solid-lined box in the 2nd lane shows the unrestricted, circular plasmid (~4.5 kbp), whereas the dashed-lined box in the 4th lane shows the vector digested by* C. kluyveri *cell lysate (fragment sizes: ~2 kbp, ~0.5 kbp). The linearized pMTL83151 vector in the 3rd lane (~4.5 kbp) and the mixture of circular plasmids pMTL83151 and pMeth (5th lane) serve as controls. B) In bacterial cells, DNA restriction serves as a defense mechanism against foreign DNA that either lacks methylation at the critical sites or shows a different methylation pattern from the native DNA of the respective bacterium. C) Restriction digest of* C. kluyveri-*specific methylated pMTL83151 after 24 h. Because the expression of the methyltransferase (CKL_2671) of* C. kluyveri *was controlled by the P$_{lac}$ promoter, we tested different IPTG*

levels for induction (10 mM to 1 M). All induction levels result in minor restriction of pMTL83151 highlighted with dashed boxes (fragment sizes: ~2 kbp and ~0.5 kbp), however, the restriction was clearly reduced and minimized. Best results with the least restriction were obtained for the induction with 500 mM IPTG. D) As a result of the same methylation pattern as the native DNA, the foreign DNA is protected from restriction by the bacterial restriction modification system. Created with BioRender.com.

Figure 2. *Plasmid map of pMeth based on pUC19. CKL_2671 was amplified from C. kluyveri genomic DNA and cloned into the multiple cloning site of pUC19. The T7 terminator was cloned into the plasmid and the ORI of puC19 was exchanged to the p15A ORI. Created with BioRender.com.*

Figure 3. *Verification of pMTL83151 inside* C. kluyveri *cells. A) We screened 18* C. kluyveri *colonies (lanes 1-18) for the presence of the plasmid pMTL83151 and verified its presence* via *colony PCR of the respective clones. Negative and positive (fragment size: ~1 kbp) controls are shown on the right side. The band corresponding to a longer fragment in the positive control results from the circular pMTL83151 plasmid, which served as template. B) Sanger Sequencing results of the 16S rRNA gene from the screened clones. All clones showed clear signals, confirming pure* C. kluyveri *pMTL83151 cultures. We show only one of the sequencing results here, which represents all tested clones. Created with BioRender.com.*

Figure 4. Results of the growth experiment. A) The growth behavior was measured by OD$_{600}$, showing that all strains but C. kluyveri *pPthl_adhE2 showed comparable growth. B)* C. kluyveri *pPthl_adhE2_bdhB reached the highest* n-*butanol concentration, whereas* C. kluyveri *pPadhE2_adhE2 and* C. kluyveri *pPthl_adhE2 showed comparable production levels. C) We observed the highest* n-*hexanol levels for* C kluyveri *pPthl_adhE2_bdhB, followed by* C. kluyveri *pPthl_adhE2. Created with MATLAB2024b.*

SUPPORTING INFORMATION

Plating efficiency

The plating efficiency was optimized by stepwise exchanging one condition while using the same pre-culture for each comparison (**Table S1**). Colony forming units (CFU) were counted, and the best-performing combination of the two combinations was defined as the current best condition. The current best condition was then used as a control for the next round of the experiment when, again, one factor/condition was changed. If the following combination showed better results than the control, it was updated to be the new current best condition. As a result, we achieved an iterative improvement of the plating efficiency in each round. We observed the biggest positive impact on the plating efficiency by supplementing a 10x concentration of the seven-vitamin solution, compared to the concentration used for cultivating *C. kluyveri* in liquid culture. We refer to this modified medium as DSMZ52*. The different conditions that we tested and their respective outcome are shown in **Table S1**.

Table S1. *Optimization of the plating efficiency. Each line represents an independent and individual experiment in which two conditions were tested, respectively. Experiments are listed chronologically, meaning that the plating efficiency stays constant with every line or is improved. DSMZ52* means that DSMZ52 medium with 10x vitamin supplementation was used. From all conditions tested, pour plating a culture in its early exponential phase in DSMZ52* with 0.8% agar, 0.6 bar overpressure (N_2/CO_2), and a filter paper soaked with ethanol showed the best plating result.*

Dilution factor	Current best combination	CFU per plate	Adapted condition	CFU per plate
1:100	Spread plating on DSMZ52	563	Spread plating on DSMZ52 with 10x vitamin supplementation	1,095

1:1,000	Spread plating on DSMZ52*	707	Spread plating on DSMZ52*, with paperclips for increased ventilation of CO2	148
1:100	Spread plating on DSMZ52*	1,001	Spread plating on RCM	402
1:1,000	Spread plating on DSMZ52*	980	Pour plating in DSMZ52*in 0.8% agar	1,056
1:100	Pour plating in DSMZ52* in 0.8% agar	944	Pour plating in DSMZ52* in 0.8% agar, additional EtOH in the headspace	141
1:10,000	Pour plating in DSMZ52* 0.8% agar, early exponential phase	678	Pour plating in DSMZ52 * in 0.8% agar, late exponential phase	216
1:10,000	Pour plating in DSMZ52* in 0.8% agar, early exponential phase	577	Pour plating in DSMZ52* in 0.8% agar, early exponential phase, at 0.6 bar overpressure	748
1:10,00	Pour plating in DSMZ52* in 0.8% agar, early exponential phase, at 0.6 bar overpressure	551	Pour plating in DSMZ52* in 0.8% agar, early exponential phase, at 0.6 bar overpressure, with a filter paper soaked with EtOH inside the petri dish	579

*CFU: colony forming units per plate

DSMZ52: DSMZ52 medium with 10x seven-vitamin solution

Determination of the minimal inhibitory concentration of thiamphenicol for

C. kluyveri

Wild-type cells of *C. kluyveri* were grown in liquid DSMZ52 medium at different thiamphenicol concentrations to determine the minimum inhibitory concentration (MIC). After one week, we observed growth up to a thiamphenicol concentration of 2 µg/mL, suggesting that 3 µg/mL was the MIC (**Figure S1**). However, to ensure no wild-type cells would grow, for example, due to spontaneous resistance, we decided to use a thiamphenicol concentration of 5 µg/mL to select the plasmid-carrying cells in liquid medium and 3 µg/mL on plates.

Figure S1. *Determination of the MIC of thiamphenicol for* C. kluyveri.

Selection for plasmids

To identify the ideal mating time to obtain *C. kluyveri* mutants, we restreaked the potentially pMTL83151-carrying cells after 24, 48, 72, 96, and 120 h of mating. After 24 h, we observed individual colonies (**Figure S2A**), whereas, after 48 h of mating, the plates showed a lawn of cells carrying the plasmid (**Figure S2B**), indicating that more of the transferred cells carried the plasmid pMTL83151. Because colonies could not be distinguished after 48 h for the empty vector pMTL83151, no increased efficiency could be observed after more than 48 h of mating. However, longer mating times might be beneficial for larger or more toxic plasmids.

Figure S2. *Growth of* C. kluyveri *colonies on selective plates after 24 and 48 h mating time.*

Growth experiment for *n*-butanol and *n*-hexanol production

We measured the substrates, ethanol (**Figure S3A**) and acetate (**Figure S3B**), and

the products, *n*-butyrate (**Figure S3C**) and *n*-caproate (**Figure S3D**), for the different

mutant strains (*C. kluyveri* pMTL83151, *C. kluyveri* pPadhE2_adhE2, *C. kluyveri*

pPthl_adhE2, and *C. kluyveri* pPthl_adhE2_bdhB producing *n*-butanol and *n*-hexanol

and the negative control carrying the empty pMTL83151 vector throughout time.

Figure S3. *Concentration of the substrates ethanol A) and acetate B) and the products*

n-*butyrate C) and* n-*caproate D).*

Sanger Sequencing of P*thl* in *C. kluyveri* pPthl_adhE2 cells

Sanger Sequencing of *C. kluyveri* pPthl_adhE2 revealed mutations in the promoter region P*thl* of the *C. kluyveri* pPthl_adhE2, which did not produce any longer-chain alcohols than ethanol (**Figure S4A**). Also, the replicates of *C. kluyveri* pPthl_adhE2 seemed to have a fraction of mutated cells (marked with the red box) at the end of the experiment (**Figure S4B**).

Figure S4. *Sequencing results of the promoter region of* C. kluyveri *pPthl_adhE2. A) Deletion in the promoter region (*P*thl*) *of* adhE2 *in* C. kluyveri *pPthl_adhE2 cultures, which did not produce* n-*butanol and* n-*hexanol. B) Shows the same, not deleted, promoter region of* C. kluyveri *pPthl_adhE2 cultures, which produced* n-*butanol and* n-*hexanol, at the end of the experiment. However, sequencing these cultures revealed mixed signals (especially inside the region marked with the red box), indicating that a fraction of the cells had mutated during the growth experiment. Created with BioRender.com.*

Plasmid list

Table S2. *List of plasmids used in this study.*

Name	Information	Reference
pMTL83151	Shuttle vector for *Clostridia*	Heap et al. 2009
pMeth	Methylation plasmid, encoding a methyltransferase of *C. kluyveri* DSM555 (CKL_2671) for methylation of plasmid DNA matching the methylation pattern of *C. kluyveri* (CCGG site)	This study
pUC19	Standard cloning vector	Yanisch-Perron et al. 1985
pPadhE2_adhE2	Alcohol production *via* bifunctional aldehyde/alcohol dehydrogenase AdhE2 from *C. acetobutylicum* (CA_P0035) controlled by the native promoter P_{adhE2}	This study
pPthl_adhE2	Alcohol production *via* bifunctional aldehyde/alcohol dehydrogenase AdhE2 from *C. acetobutylicum* (CA_P0035) controlled by the promoter P_{thl}	This study

| pPthl_adhE2_bdhB | Alcohol production *via* bifunctional aldehyde/alcohol dehydrogenase AdhE2 from *C. acetobutylicum* (CA_P0035) and the butanol dehydrogenase BdhB from *C. acetobutylicum* (CA_C3298) controlled by the promoter P*thl* | This study |

Primer list

Table S3. *List of oligonucleotides used in this study.*

Name	Sequence	Function
8F	AGAGTGTGATCCTGGCTCAG	Sequencing of the 16S rRNA gene region
1492r	GGTTTTTTTACGACTT	
CS-089	CTAGTCGACATGCGTGGTGGAAAT CG	Amplification of CKL_2671
CS-090	CATCTAGATTATAAAGCTATAGTTT TTATTTTTTCAATATCCTG	
CS-095	CATTAGCGCAAGGTGATTTTTGTCT TCTTGCGCTAATTTTTTTCTAGAGG ATCCCCGGG	T1 terminator exchange
CS-096	CCTTGCGCTAATGCTCTGTTACAGT TATAAAGCTATAGTTTTTATTTTTTC AATATCCTG	
CS-141	GAAACAGCTATGCGTGGTGGAAAT CGAC	Gibson Assembly for p15a ORI
CS-142	CACCACGCATAGCTGTTTCCTGTG TGAAATTG	
CS-143	CCCGTAGAAAAGATCAAAGGATCT TCAAGATGATCTTCTTGAGATCGTT TTG	
CS-144	CGCGTTGCTGGCGTTCGTTTTTCC ATAGGCTCCG	
CS-145	CGGAGCCTATGGAAAAACGAACGC CAGCAACGCG	Removal of remaining *lacZ* gene
CS-146	CAAAACGATCTCAAGAAGATCATCT TGAAGATCCTTTGATCTTTTCTACG GG	
CS-112	CAAAACCTTCACGCTTGCCTATATT G	Sequencing pMeth
CS-147	GAGTCAGGCAACTATGGATGAAC	
CS-148	GAGTCAGTGAGCGAGGAAG	
CS-159	TTTTTATTGCCAAATACTAGCATAA AAATAAGAAGCCTGCAT	Gibson Assembly for P$_{adhE2}$_*adhE2*_terminator insert into pMTL83151
CS-160	GCTTTTTACTCCTGCTTGACATGGA TACAGCGGCC	
CS-161	ACCGCGGCCGCTGTATCCATGTCA AGCAGGAGTAAAAAGCG	
CS-162	GCAGGCTTCTTATTTTTATGCTAGT ATTTGGCAATAAAAATAGTTATAAT CATTAATTATTG	

CS-240	TTTTGATTTGTAACTTTCATATGAAC TAACCTCCTAAATTTTGATACGGGG	
CS-241	TGCAGGCTTCTTATTTTTATTTTTTA ACAAAATATATTGATAAAAATAATAA TAGTGGGTATAATTAAG	Gibson Assembly for P$_{thl}$ insert
CS-242	TCAATATATTTTGTTAAAAAATAAAA ATAAGAAGCCTGCATTTGCAGG	
CS-243	AATTTAGGAGGTTAGTTCATATGAA AGTTACAAATCAAAAGAACTAAAA CAAAAGCTAAATG	
CS-236	TTTATCTACAATTTTTTTATACTCCC CAAGATATTAATGCTTT	
CS-237	TAAAATCATTTTAAAAAATAGTGGT TGATTTCGAATATTCAATACC	
CS-238	GAATATTCGAAATCAACCACTATTT TTTAAAATGATTTTATATAGATATCC TTAAGTTCACTTATAAG	Gibson Assembly for bdhB_terminator insert
CS-239	GCATTAATATCTTGGGGAGTATAAA AAAATTGTAGATAAAACTATTTTATA AAATTTATCTACAAT	
EB00003594	GTACTGCTTCCAATACACAAAGGC	
CS-162	GCAGGCTTCTTATTTTTATGCTAGT ATTTGGCAATAAAAATAGTTATAAT CATTAATTATTG	
CS-227	CTGGATAGATGAGCCATCAATAG	Sequencing P$_{adhE2}$_adhE2, P$_{thl}$_adhE2, and P$_{thl}$_adhE2_bdhB
CS-228	GTGTTGCTGAAAGAAGGGAAAATA TG	
CS-229	GATTATACTGATGAATTAGCCTTAA GAGC	
CS-237	TAAAATCATTTTAAAAAATAGTGGT TGATTTCGAATATTCAATACC	
MRS-040	CAAGGCAAGACCGATCG	
EB00002844	GCGGCCGCTCTTTCCTGCGTTATC CCCTG	Verification of pMTL83151-based plasmids
EB00002845	ACATGTGGTCATAGCTGTTTCCTGA TAAAAAAATTG	

Bacterial strains

Table S4. *List of bacterial strains used in this study.*

Name	Source
C. acetobutylicum ATCC824	ATCC
C. kluyveri DSM555	DSMZ
C. kluyveri pMTL83151	This study
C. kluyveri pPadhE2_adhE2	This study
C. kluyveri pPthl_adhE2	This study
C. kluyveri pPthl_adhE2_bdhB	This study
E. coli TOP10 pMeth	This study
E. coli TOP10 pMeth pMTL83151	This study
E. coli TOP10 pPadhE2_adhE2	This study
E. coli TOP10 pMeth pPadhE2_adhE2	This study
E. coli TOP10 pPthl_adhE2	This study
E. coli TOP10 pMeth pPthl_adhE2	This study
E. coli TOP10 pPthl_adhE2_bdhB	This study
E. coli TOP10 pMeth pPthl_adhE2_bdhB	This study
E. coli HB101 pKR2013	DSMZ