Crossing effects in the tactile temporal order judgment task:
A meta-analysis

Nina Schmelter[1], Benedikt Langenberg[2], Axel Mayer[1], Tobias Heed[3,4]

[1] Department of Psychology, Faculty of Psychology and Sports Science, Bielefeld University, Bielefeld, Germany

[2] Department of Methodology and Statistics, Maastricht University, Netherlands

[3] Department of Psychology, University of Salzburg, Salzburg, Austria

[4] Centre for Cognitive Neuroscience, University of Salzburg, Salzburg, Austria

Correspondence

T.H., tobias.heed@plus.ac.at

Acknowledgments

We thank Tine-Marie Wujciak for helpful discussion during conceptualization, data curation, and formal analysis.

Author Contributions

Contributions according to https://credit.niso.org

NS: conceptualization, data curation, formal analysis, investigation, methodology, validation, visualization, writing – original draft

BL: formal analysis

AM: conceptualization, methodology, formal analysis, project administration, supervision, validation, writing – original draft

TH: conceptualization, project administration, supervision, writing – original draft

1 Abstract

The tactile temporal order judgment (TOJ) task is widely used in multisensory
neuroscience. Participants judge which of two tactile stimuli, one on each hand,
came first. A key finding is that TOJ performance declines when the arms are
crossed, likely due to interactions between tactile, proprioceptive, and visual
information. The TOJ crossing effect has been widely reported, but studies have
employed various analysis methods, leaving open whether the choice of method
influences the effect's magnitude. Moreover, some studies have reported
modulations by the availability of visual information, response modality, or speeded
response requirements. Through an exhaustive, systematic literature search, we
identified 37 experiments that investigated the TOJ crossing effect. Meta-analysis
estimated the effect size at approximately 1.4. The moderators did not significantly
affect this estimate, though this is likely due to too few studies being available to
obtain sufficient statistical power. The large effect size supports the TOJ task's use in
other, e.g. developmental and clinical, research. However, the lack of statistically
significant effect modulation by the moderators calls for caution when applying it for
research questions beyond the crossing effect itself.

18 Keywords

19 meta-analysis; touch; remapping; multisensory; tactile temporal order judgement

Introduction

Crossing the arms over the body midline impairs tactile choice tasks, a phenomenon known as the (tactile) crossing effect[1]. A prominent experimental paradigm for its investigation is the tactile Temporal Order Judgment (TOJ) task, in which participants report which of two consecutive stimuli, applied one to each hand, occurred first (or second), typically by pressing a button with the respective hand[2,3]. Intuitively, it would seem irrelevant for choosing a hand for a button press whether the arms are held uncrossed or crossed. However, when participants cross their arms, their TOJ are strongly impaired and sometimes even systematically reversed. Therefore, the TOJ task has become a prime example of a tactile crossing effect, and it has been used extensively over the last two decades to investigate tactile spatial processing.

The origin of crossing effects such as that observed in the TOJ task has been debated. The most prominent idea has been that crossing effects arise from a conflict between different spatial codes: a skin-based, or anatomical, code that signals where on the skin the touch occurred; and a spatiotopic, or external, code that signals where the touch was in space[1,4]. With crossed arms, these two codes carry opposing information; the right hand, for instance, is then located in left space, and it may be this kind of conflict that impairs tactile judgments. Newer studies have cast doubt on this theory and have suggested that crossing effects, instead, stem from confusion of where the touched limb is usually located – its default posture – and where it is placed at the moment of touch[5–7].

Irrespective of its theoretical underpinnings, the TOJ crossing effect is often large, and it is probably this feature that has contributed the most to the task's popularity. However, there are multiple versions of the TOJ paradigm, and the task has been analyzed in multiple ways. This variability in how the task is run and analyzed has made it difficult to compare studies directly, beyond assessing qualitative similarities. Moreover, no attempt has been made to estimate a statistical effect size based on evidence from multiple studies.

An effect size estimate would aid future research, for instance, for planning sample sizes based on statistical power or for defining priors in Bayesian analysis. The TOJ task has been employed, among others, in developmental and clinical studies, including research on mental disorders, physical illnesses, and brain processes[8–11].

Such studies typically aim at optimizing experimental efficiency and determining the
minimum number of required study participants. One option to reduce experimental
time may be to apply yet another version of the TOJ task which applies adaptive
testing[12]. Yet, a proper estimate of the expected effect size is necessary to
determine an appropriate sample size. In practice, therefore, experiments are often
either inefficient or underpowered[13].
The TOJ task has been employed for a wide range of scientific questions, and
numerous manipulations have been applied next to limb crossing for this purpose.
Some of them have been scrutinized across multiple studies, potentially allowing
investigating their effect sizes as well. One key variable that has differed across
studies is the response modality. Often, participants have responded with the hands,
that is, with the same limbs that received stimulation. Some studies, however, have
required button responses with the feet. With this response modality, it is necessary
to instruct participants how to match stimulus hand and response foot: under
anatomical instructions, responses are given with the foot that belongs to the same
body side as the hand that received the first stimulus. In contrast, under external
instructions, responses are given with the foot that is on the same side of external
space as the stimulated hand. These two instructions require different responses
when the hands are crossed: for instance, when the participant chooses the right
hand, (s)he must press the right foot under anatomical instructions (stimulus on right
body side), but the left foot under external instructions (stimulus on left side of space
due to crossed posture). Several studies have reported a larger TOJ crossing effect
for external than anatomical instructions[14,15]. In yet other studies, participants
responded verbally[11,16], but, to our knowledge, no direct comparison of verbal vs.
button press responses has been published so far. However, the apparent
differences between these response modalities have not been systematically
explored.
A second manipulation that has been investigated in multiple studies is the
availability of visual information. The crossing effect was reported to be diminished
when the hands were crossed behind the back[17], when the eyes were closed[18], and
when uncrossed rubber arms were placed above the hidden, crossed arms[19]. While
most studies have reported that the absence of visual input reduces the crossing
effect, some findings have been inconsistent[20].

A third aspect which may affect TOJ are task instructions regarding speed and
accuracy. The TOJ has sometimes been run as an unspeeded task that emphasized
correct responding[2], whereas many studies have asked participants to respond both
as fast and as correct as possible[3,21–23]. Emphasizing speed may compel participants
to respond before they have fully processed the stimuli, leading to more frequent
guessing and errors in the crossed posture, which may, in turn, amplify the crossing
effect. We are, however, not aware of any study systematically addressing this detail
for the TOJ task.
Finally, the reported size of experimental effects may systematically depend on the
analysis method[1]. TOJ performance has often been assessed using a range of
stimulus onset asynchronies (SOAs) between the two tactile stimuli. By convention,
SOAs are marked as negative when the left hand stimulation occurred first and
plotted as "percentage of right-first" responses (y-axis) against the SOA (x-axis).
With uncrossed hands, performance follows a typical psychophysical, S-shaped
curve. In contrast, with crossed hands, the curve is often N-shaped, with responses
for long SOAs correct but responses for short SOAs reversed[3]. This N-shaped result
pattern is unusual – in fact, we are not aware that it has been observed in any other
psychophysical task to date. Reaction times in the TOJ task are longer for shorter
SOAs, and are significantly prolonged in the crossed compared to the uncrossed
posture[3,23].
Whether speeded or not, most reports have focused on analyzing TOJ accuracy
rather than RT. One way to analyze accuracy has been to probit-transform
percentage of right-first responses to linearize the response curve's S-shape, and to
derive the slope of the resulting straight line[2]. A second measure has been to derive
the just noticeable difference as half the distance between the SOA required for 25%
and 75% correct right-first responses after probit transformation[2]. A third measure
has been to simply sum up accuracy across all SOAs[21]. With all three methods, the
crossing effect was defined as the difference of the respective measure between the
uncrossed and crossed postures. Finally, a fourth analysis method has been to fit the
N-shaped response curves of the crossed posture. This can be done by adding up
three Gaussians[3]. Two of the three Gaussians fit the N's bulks; their amplitudes,
termed A_{right} and A_{left}, are then interpreted as a measure of the crossing effect,[3,24,25].
Note, the two "flip Gaussians" are derived solely from the crossed condition; this

distinguishes the flip fitting procedure from all other crossing effect measures, which

all rely on subtracting performance between uncrossed and crossed conditions.

There is no consensus on which analysis method is most appropriate, nor does any

method exhibit generally greater sensitivity or consistently yield a larger crossing

effect[23]. In addition to these analysis choices, the choice of SOAs used in the

experiment, as well as some further analysis options, may result in differences

across studies[26].

In sum, both the need for an effect size estimate to aid future study planning and the

lack of consensus about the formal analysis for TOJ regarding the crossing effect,

and the so far unsystematic exploration of the four discussed potential moderators,

all call for a systematic review of the research, which we provide by conducting a

meta-analysis for the TOJ crossing effect and its moderation by response mode,

visual input, speed and accuracy instructions, and analysis method.

Methods

Search Strategy

We searched the databases BASE, EMBASE, Pubmed, and Web of Science,

accessed through Bielefeld University on April 28, 2023. We chose these databases

because they have demonstrated high levels of precision, recall, and

reproducibility[27]. We employed the search string "(tactile OR hand OR arm) temporal

order judg*" for all databases with the respective, appropriate syntax (see Appendix

A, Table A1). We included only English articles when this option was available.

Selection Criteria

Studies were included in the meta-analysis if they: (1) conducted the tactile TOJ task

with a crossed and an uncrossed condition, (2) presented stimuli sequentially on

both hands, (3) asked participants to judge the temporal order of both stimuli, (4)

collected responses from participants using their hands, feet, or verbally, (5) included

participants aged 18 and above, (6) were published peer-reviewed articles, (7) were

in English. Studies were excluded if they: (1) reported mental or health impairment of

participants and no unaffected control group was available, (2) did not provide

sufficient data within the published articles to calculate a crossing effect. This

criterion led us to exclude studies that reported only flip parameters obtained by

fitting Gaussians (see Introduction, fourth analysis method). As flip parameters are unique to the crossed condition, they cannot be tested against the uncrossed condition, making them formally incomparable to the remaining measures which all compare crossed and uncrossed conditions. Qualitatively, however, the respective studies support the conclusions we draw from our formal analysis (see Discussion).

Data Extraction and Derived Variables

All studies were collected in Zotero (Corporation for Digital Scholarship, Vienna, Virginia, USA, https://www.zotero.org). We employed automated procedures to eliminate duplicates and removed undetected duplicates manually. The assessment of the remaining studies regarding the fulfillment of the inclusion criteria was performed by one coder (N.S.). The same coder extracted data from each study, including statistical data for calculating effect sizes and standard errors. All results were double-checked by the same coder to ensure accuracy.

 Additionally, potential moderator variables were operationalized and coded for each study in the following manner:

       (1) outcome measure

           a. "accuracy" – the difference in performance between the crossed and uncrossed condition was calculated for each SOA and then pooled over all SOAs, as introduced in *(21)*

           b. "slope" – the S-shaped curve of the percentage of right-first responses was probit, logit, or Weibull transformed to linearize the psychometric curve, and the resulting straight line displayed the performance in the TOJ task as a function of the SOA

           c. "JND 75%" – the JND was obtained from linearized response data (see b) and calculated as the time window between the two applied stimuli in which 75% of the judgments were correct)

           d. "JND 84%" – response data were pooled across participants and fitted with a cumulative four-parametric cumulative Gaussian function, as introduced in *(3)*. Subsequently, the standard deviation (σ) was calculated, reflecting the smallest time window between the two applied stimuli in which 84% of the judgments were correct)

182 If multiple outcome measures were available, we entered only one into our analysis,

183 using the following order: accuracy, slope, JND 75%, JND 84%.

184   (2) eyes

185   a. "open" – participants' eyes were open; we did not differentiate

186   whether fixation was required or not, or whether participants'

187   hands were hidden or not

188   b. "closed" – participants were blindfolded

189   (3) response modality and response demands

190   a. "hands" participants responded with the hands or a finger of the

191   hand that was stimulated first [or second]

192   b. "feet external" – participants responded with the foot on the

193   same side of space as the hand stimulated first [or second]

194   c. "feet anatomical" – participants responded with the foot on the

195   same body side as the hand stimulated first [or second]

196   d. "verbal" participants responded verbally

197   (4) task instruction regarding speed and accuracy

198   a. "unspeeded" – participants responded without any time

199   restriction and/or were instructed to prioritize accuracy

200   b. "time-restricted" – participants had to respond within a specific

201   time interval or were instructed to respond as accurately and as

202   quickly as possible; the latter were usually combined)

Statistical Analysis

Effect Measures

All included studies used within-subject designs, and all studies except two

aggregated the dichotomous dependent variable (correct or incorrect) into a

percentage correct score that can be treated as continuous for the range of values

that is not too close to the limits. Therefore, we calculated a version of Cohen's d

known as Cohen's d_z or d_D (hereafter referred to as Cohen's d_z), which is appropriate

for within subject-designs with a continuous outcome measure. Using Cohen's d_z

allowed us to pool primary studies with different outcome measures, such as slope

versus percentage correct. One possibility to calculate Cohen's d_z is to compute the

mean difference between crossed and uncrossed conditions and then use the

standard deviation of these difference scores as the denominator[28]:

$$Cohen's\ d_z = \frac{M_{crossed} - M_{uncrossed}}{SD_{difference}} \quad (1)$$

215 The mean difference and its standard deviation were only available in studies that
216 used accuracy as an outcome measure. Alternative derivations are available for
217 Cohen's d_z depending on the available information:

218   a. calculation based on the t-value and the sample size[29]:

$$Cohen's\ d_z = \frac{t}{\sqrt{n}} \quad (2)$$

219   b. calculation based on the F-value (with one numerator degree of freedom)
220   and the sample size[30]:

$$F = t^2,\ \text{consequently}\ Cohen's\ d_z = \frac{\sqrt{F}}{\sqrt{n}} \quad (3)$$

221   c. from the χ^2- value and the sample size, with the χ^2- value for categorical
222   outcomes treated as F-value to obtain comparability with the analysis of
223   continuous variables as described in b:

$$Cohen's\ d_z = \frac{\sqrt{\chi^2}}{\sqrt{n}} \quad (4)$$

224   d. from the regression coefficient (β) in a generalized linear mixed model
225   (GLMM) and the sample size, employing the additonal assupmtions that
226   the β-value is unstandardized, and the z-value is approximately equivalent
227   to the t-value:

$$z = \frac{\beta}{SE},\ Cohen's\ d_z = \frac{z}{\sqrt{n}} \quad (5)$$

228   e. from the means of the crossed and uncrossed conditions, their standard
229   deviations, and the correlation between the two conditions[28]; we estimated
230   the required correlation from three openly available datasets (see
231   Supplementary Information for details):

$$Cohen's\ d_z$$
$$= \frac{M_{crossed} - M_{uncrossed}}{\sqrt{SD^2_{crossed} + SD^2_{uncrossed} - 2 * r * SD_{crossed} * SD_{uncrossed}}} \quad (6)$$

Equations (4) and (5) are approximations based on the properties of F and t, which
converge against the χ^2 and the normal distribution for large n.
If multiple statistical metrics were available to calculate the effect size, formulas were
chosen in the order reported above.

Standard Error

The standard error of each effect size was calculated as shown in equation (7). The

derivation of this formula is shown in Appendix B.

$$SE(d_z)$$

$$= \sqrt{\frac{1}{n} * \frac{n-1}{(n-1)-2} + d_z^2 * \left(\frac{n-1}{(n-1)-2} - \frac{1}{\left(1 - \frac{3}{4*(n-1)-1} \right)^2} \right)} \quad (7)$$

Meta-Analysis

We conducted statistical analyses with R[31] and R Studio[32]. We employed the

packages meta[33] to conduct the meta-analysis, dmetar[34] for outlier and influence

analysis, metasens[35] for Rücker's Limit Meta-Analysis Method, and tidyverse[36] for

data handling.

We used a random-effects model to pool effect sizes, because we anticipated

considerable between-study heterogeneity. We calculated the heterogeneity

variance (τ^2) using the restricted maximum likelihood (REML) estimator[37] and the

confidence interval around the pooled effect using Knapp-Hartung adjustments[38].

$$Cohen's\ d_z = \frac{\sum_{k=1}^{K} Cohen's\ d_{z\ k} w_k^*}{\sum_{k=1}^{K} w_k^*} \quad (8)$$

where:

w_k^* is the weight assigned to each study k

K is the number of studies included in the meta-analysis

with:

$$w_k^* = \frac{1}{se_k^2 * n_k + \tau^2} \quad (9)$$

where:

n_k is the sample size of each study k

We used Cochran's Q to test if between-study variation exceeded sampling error

and estimated the proportion of variability due to true differences using Higgins and

Thompson's I^2 [39]. Additionally, we calculated prediction intervals to estimate the

expected range of effects in future studies[40]. To detect divergent studies, that is,

studies that differ markedly from the study pool, we set as outlier criterion that a

study's 95% confidence interval was beyond the 95% confidence interval of the

pooled effect. Additionally, we identified studies with a high influence on the pooled

effect size using the leave-one-out method. To this end, we calculated, for each

study, the difference in Cohen's d_z when the study was included in the analysis

compared to when it is not included. The resulting differences were plotted in a

Baujat plot on the y-axis, with overall heterogeneity, measured by Cochran's Q, on

the x-axis[41].

Analysis of moderator effects

To explore the impact of the four potential moderators of the crossing effect's effect

size – analysis method, visual input, response mode, and speed and accuracy

instructions – we conducted separate moderator analyses, each over those articles

that reported the respective moderator characteristics. Ten studies lacked

information on one moderator each; therefore, we opted for separate analyses, so

that we retained as many studies as possible for each individual analysis[42]. We

applied a random-effects (plural) model for these analyses[43]. We used a pooled

version of the between-study heterogeneity variance τ^2 across all studies subsets

when any subset consisted of fewer than 6 studies[44].

Data availability statement

All variables created for our report, as well as the R code used for analysis, are

freely available at strategy https://osf.io/xtd9g/. Moreover, the data contain several

variables we extracted during data extraction but did not use for our report, such as

the reported *p* and *df* values for *t*, *F*, and χ^2 tests.

Results

Search Results

The database search yielded 973 studies (Fig. 1). After removing duplicates, 346
eligible studies remained. 301 of these were removed during abstract screening.
Forty-five studies underwent full-text review; they comprised a total of 85
experiments. Of these, 18 studies/51 experiments were excluded, 18 experiments
due to insufficient data and 33 experiments due to not meeting the inclusion criteria
(see Supplementary Table B1 for references and reasons for exclusion of fully
screened studies).

The remaining 27 studies comprised a total of 37 experiments and comprised 636
participants (see Table 1 for study citations and characteristics). The difference
between studies and experiments results from our including between-subject
experimental groups, such as separate groups that conducted the experiment with
their eyes open vs. closed, as well as different experiments within a single study if
they comprised different individuals.

Figure 1. Flowchart of study selection. Design adapted from the PRISMA 2020 statement[45].

295 Seven studies comprised 2-4 experiments each (see Supplementary Table B2). As
296 the experiments were conducted with different persons, we treat them as
297 independent experiments. For further information and limitations regarding the
298 coding of subgroup analysis variables outcome measure, response modality, and
299 task instructions about speed see Supplementary Tables B3, B4, and B5.

Table 1. Included studies and characteristics used for the meta-analysis.

Author	n	Statistical characteristics	Outcome measure	Eyes	Response modality	Speed
Azañón & Soto-Faraco (2007)[19] - exp. 1	19	*M, SE*	JND 84%	open	hands	unspeeded
Azañón & Soto-Faraco (2007)[19] - exp. 2	20	*M, SE*	JND 84%	open	hands	unspeeded
Azañón et al. (2015)[46]	12	*F*	JND 75%	open	hands	unspeeded
Azañón, Mihaljevic et al. (2016)[47]	16	*t*	JND 75%	closed	hands	unspeeded
Azañón, Radulova et al. (2016)[48]	11	*t*	slope	NA	hands	unspeeded
Badde et al. (2014)[49] - exp. 1	17	*F*	Accuracy	open	hands	time restriction
Badde et al. (2014)[49] – exp. 2	17	*F*	Accuracy	open	hands	time restriction
Badde et al. (2015)[50] - exp. 1	13	*F*	Accuracy	open	hands	unspeeded
Badde et al. (2015)[50] - exp. 2	14	*F*	Accuracy	open	hands	unspeeded
Badde et al. (2017)[51]	16	χ^2	Accuracy	NA	hands	time restriction
Cadieux & Shore (2013)[14] - eyes closed, feet external	11	*t*	Accuracy	closed	feet external	time restriction
Cadieux & Shore (2013)[14] - eyes closed, hands	20	*t*	Accuracy	closed	hands	time restriction
Cadieux & Shore (2013)[14] - eyes open, feet external	11	*t*	Accuracy	open	feet external	time restriction
Cadieux & Shore (2013)[14] - eyes open, hands	20	*t*	Accuracy	open	hands	time restriction
Cadieux et al. (2010)[21]	48	*t*	slope	open	feet external	time restriction
Crollen, Albouy, et al. (2017)[52]	11	*t*	slope	closed	hands	time restriction
Crollen, Lazzouni, et al. (2017)[22]	11	*t*	slope	closed	hands	time restriction
Crollen et al. (2019)[53]	15	*t*	slope	closed	feet external	time restriction
Heed et al. (2012)[23] - exp. 1	15	*F*	Accuracy	open	hands	unspeeded
Heed et al. (2012)[23] - exp. 2	14	*t*	Accuracy	open	hands	unspeeded
Heed et al. (2016)[4]	24	*β*	Accuracy	open	hands	NA

Study	n	Statistic	Measure	Eyes	Posture	Speed
Kóbor et al. (2006)[17]	18	*F*	JND 75%	closed	feet anatomical	unspeeded
Landry & Champoux (2018)[54]	20	*MD, SD~within~*	Accuracy	NA	feet external	unspeeded
Moseley et al. (2012)[11]	10	*t*	JND 75%	open	verbal	NA
Roberts & Humphreys (2008)[55] – exp. 1	9	*t*	slope	closed	feet external	unspeeded
Röder et al. (2004)[20] - eyes closed	13	*F*	JND 75%	closed	hands	unspeeded
Röder et al. (2004)[20] - eyes open	12	*F*	JND 75%	open	hands	unspeeded
Sambo et al. (2013)[56]	12	*t*	JND 75%	closed	hands	time restriction
Schicke & Röder (2006)[26]	10	*F*	JND 75%	closed	hands	unspeeded
Sharp et al. (2018)[57]	13	*MD, SD~within~*	Accuracy	NA	feet external	unspeeded
Shore et al. (2002)[2]	9	*F*	slope	open	hands	unspeeded
Soto-Faraco & Azañón (2013)[10]	32	*F*	JND 84%	open	hands	NA
Unwalla et al. (2020)[58]	20	*t*	Accuracy	NA	hands	time restriction
Unwalla, Goldreich, et al. (2021)[15] – exp. 1	20	*t*	Accuracy	NA	feet anatomical	time restriction
Unwalla, Goldreich, et al. (2021)[15] - exp. 2	43	*t*	Accuracy	NA	feet anatomical	time restriction
Unwalla, Cadieux, et al. (2021)[18] - eyes closed	20	*t*	Accuracy	closed	hands	time restriction
Unwalla, Cadieux, et al. (2021)[18] - eyes open	20	*t*	Accuracy	open	hands	time restriction

Note. Exp. = experiment, *n* = simple size, *M* = mean, *SE* = standard error, *MD* = mean difference, *SD* = standard deviation, NA = not available

Our main analysis involved the data from all identified 27 studies with their 37

experiments. The estimated effect size of the crossing effect, quantified as Cohen's

d_z, was 1.42 (t(36) = 19.44, p < .001, 95% CI [1.27, 1.56]). This effect size is very

large, based on Cohen's classification of effects as large when d_z >= 0.8[59]. The

estimated between-study heterogeneity variance, τ^2, was 0.06 (95%CI [0.00, 0.15]),

with Q(36) = 47.67, p = .09, and an I^2 value of 25% (95%CI [0%, 50%]). The

prediction interval ranged from Cohen's d_z = 0.87 to 1.96, indicating that the TOJ

crossing effect is large and should be reliably observable in future studies.

Figure 2. Forest plot of the effect sizes of included studies. *SE* = standard error,

CI = confidence interval, *df* = degrees of freedom. Red squares represent the point

estimates for each study and black lines are 95% confidence intervals. The size of

the squares reflects the study weight with which a study contributes to the pooled

effect size. The black diamond symbolizes the pooled effect estimate, while the red

line below denotes the prediction interval.

The outlier analysis identified two studies: Cadieux et al. (2010)[21] and Soto-Faraco

and Azanon (2013)[10]. The same two studies also contributed strongly to the overall

heterogeneity measured by Cochran's Q and exhibited a strong influence on the

pooled result measured by the leave-one-out method (see Fig. 3). Notably, the effect

size in Cadieux et al. (2010) was higher than the pooled effect size, while it was

lower in Soto-Faraco & Azañón (2013). As a result, excluding these two studies

affected the numerical results only slightly (Cohen's d_z, = 1.38, p < .001, 95% CI

[1.25, 1.51], 95% PI [1.18, 1.59; I^2 value of 0%, 95%CI [0%, 38%]).

Figure 3: Baujat Plot. Illustration of the heterogeneity with which each study

included in the main meta-analysis contributes to the effect size estimate. X-axis:

Cochran's Q, indicating the contribution to heterogeneity; Y-axis: Cohen`s d_z,

representing the difference in the overall effect when a study is included or not

assessed with the leave-one-out method. Note, the Y-axis only displays positive

values, meaning that both positive and negative deviations from the pooled effect are

represented with positive values. Circle size indicates the study weight with which a

study contributes to the pooled effect size. See main text for details.

Influence of Moderators on Effect Size Estimate

Table 2 summarizes influence of the four potential moderators – analysis method,

visual input, response mode, and speed and accuracy instructions – in the same

order as above.

Regarding the analysis method, the largest descriptive crossing effects was present
for slope and accuracy, a slightly smaller crossing effect for JND 75%, and the
smallest for JND 84%. Nevertheless, all effect size estimates were still very large,
ranging from 1.04-1.53, and these differences did not reach statistical significance.
For the moderator visual input, the crossing effect was slightly larger when
participants had their eyes open than when they had them closed. However, this
descriptive difference was not statistically significant, and again, the effect size was
very large for both cases.
For response modality, the crossing effect was smaller for foot responses under
anatomical instructions and for hand responses. The effect was larger when
participants responded verbally and when they gave foot responses under external-
spatial instructions. Note, however, that only a single study employed verbal
responses, and three studies employed anatomical foot responses. Only the
difference between the two most extreme effect size values – hand responses and
external foot responses – was statistically significant.
The crossing effect was smaller when participants gave unspeeded than speeded
responses, but again this descriptive difference did not reach statistical significance.
354
355   **Table 2. Effect size estimates when different potential moderators were**
356   **considered.**

Moderator	k	Cohen's d_z	95% CI	p	I^2	$p_{subgroup}$
			Analysis method			
JND 84%	3	1.04	0.2 - 1.88	.03	29%	
JND 75%	8	1.23	0.9 - 1.57	< .001	0%	
Slope	7	1.5	1.03 – 1.98	< .001	35%	
Accuracy	19	1.53	1.33 – 1.73	< .001	13%	
Total	37					.07
			Visual input			
Open	18	1.5	1.28 - 1.72	< .001	35%	
Closed	12	1.27	1.07 – 1.46	< .001	0%	
Total	30					.1
			Response modality			
Hands	26	1.3	1.14 – 1.46	< .001	1%	

Feet external	7	1.95	1.56 – 2.33	< .001	0%
Feet anatomical	3	1.25	0.29 – 2.22	.03	40%
Verbal	1	1.8	0.5 – 3.11	.007	-
Total	37				.002
Task instruction of speed					
Unspeeded	17	1.41	1.17 – 1.64	< .001	1%
Time-restricted	17	1.46	1.24 – 1.68	< .001	29%
Total	34				.72

357 *Note.* k = number of studies, CI = confidence interval. *p*-values for each moderator
358 characteristic indicate whether the crossing effect is significant when only including
359 studies possessing that characteristic. *p*-value for the total indicates whether the
360 moderator has a significant effect.

Discussion

Our meta-analysis over 27 studies, which comprised 37 experiments, provides robust evidence for a significant crossing effect in tactile TOJ tasks. Specifically, arm crossing leads to a pronounced performance decline, as evidenced by a pooled effect size of Cohen's $d_z = 1.42$. The prediction interval ranged from 0.87 to 1.96, suggesting that we can be certain that the effect size is large (> 0.8). Moreover, between-study heterogeneity was low. This formal result confirms the colloquial perception of the TOJ crossing effect in the tactile research community, encouraging its further use in research into tactile spatial processing. It is noteworthy that the task has also been attested high reliability[58], further supporting this conclusion.

The effect size was not consistently affected by the analysis method. This finding is in line with a report that analyzed three experiments with several analysis approaches[23]: All approaches yielded qualitatively comparable results, and none of them resulted in consistent biases, such as always identifying larger effects than others. This suggests that methodological variations in analysis should not be a major concern when comparing studies in this field. Notably, the comparison included the use of the flip gaussian fitting procedure, which we excluded in the present meta-analysis because it does not involve a direct comparison of uncrossed and crossed performance. The result pattern of the studies that used this method is well in line with all others. One could consider that the $A_{r/l}$ parameter be converted into an effect size (e.g. by assessing its t-value tested against zero) and including flip fitting studies in the meta-analysis. Our data and code are freely available to implement this strategy (https://osf.io/xtd9g/), and we have listed flip fitting studies in all lists provided in the Supplementary Information.

Our moderator analysis addressing the response mode suggest that the spatial content of the instructions affects the TOJ crossing effect. Specifically, the crossing effect was larger when foot responses had to be given with external instructions (i.e., "on which side of space did the first stimulus occur") than with anatomical instructions (i.e., "respond with the foot of the same body side as the hand that was stimulated first"). This difference was the numerically largest effect in our analysis; we consider it likely that it was not significant because the low number of studies that have implemented foot responses leads to a lack of statistical power. Furthermore, the difference between external foot responses and hand responses was almost as

large, with a significantly greater crossing effect for external foot responses. Note,

that hand responses can be considered implicitly anatomically instructed, because

participants respond with the limb that was touched first, without reference to the

limbs' spatial location. We are not aware that these two response modes have been

directly tested against each other. Together, these results suggest that TOJ are

impaired by external response instructions, independent of which limbs are used.

Descriptively, the crossing effect was smaller when participants were blindfolded

than when they could see. While this effect did not reach statistical significance, its

numerical direction aligns with the hypotheses that have been formulated regarding

availability of visual information[20]. Notably, some prior studies have demonstrated a

strong influence of visual input[14], and the relatively small effect observed here may

have been affected by our grouping criteria. We categorized studies only based on

whether participants' eyes were open or closed, without accounting for factors such

as room illumination and fixation control, which may play a moderating role[4,14].

Future studies could adopt more precise classifications to clarify the impact of visual

input on the crossing effect, though currently the number of available studies is too

low for such a meta-analysis.

We observed a slight tendency for a more pronounced crossing effect when studies

imposed time constraints than when they did not. However, this difference was

minimal and did not reach statistical significance. We found this experimental detail

surprisingly difficult to extract from the original papers and, therefore, had to apply

coarse categorization. More detailed reporting in primary studies would enable more

fine-grained analyses in future research and may help resolve whether time pressure

amplifies the crossing effect.

Limitations

In contrast to the large crossing effect, moderation effects are much smaller. Larger

number of studies would be needed to reliably detect them, but these are not

currently available. Therefore, our subset analyses addressing potential moderators

of the crossing effect suffered from a lack of power, so that even apparently large

effect size differences were not statistically significant. On the one hand, future

studies can help such endeavors by diligently reporting on all experimental details,

even if they appear irrelevant at the time of writing up the study. This may be even

more relevant as the TOJ paradigm is often employed to test for additional

moderators, rather than crossing per se. On the other hand, our review may

encourage further dedicated research into these moderators. It is our perception

that, colloquially, researchers in the field already assume that some of the effects we

tested for do exist – for instance roles of vision availability and spatial response

mode – but our results show that the evidence is not yet firm.

An important aspect of meta-analyses is whether a publication bias exists. We did

not control for this for two reasons. First, given that there is a considerable number of

tactile TOJ publications, and they consistently report a large effect size, it is unlikely

that even studies with small sample sizes would have non-significant results. In fact,

in our own work, we have seen highly significant results with as few as five

participants. Second, commonly used small-study effect methods might produce

biased results when the standard error depends on the effect size[60], as is the case

here, and p-curve analyses only provide accurate results when the specific p-value

analyzed is directly related to the hypothesis[61], which is rarely the case in the TOJ

literature. Notably, whereas we are confident about the validity of the crossing effect

size, the same cannot be said about the moderators we have addressed: there are

markedly fewer publications for them, and accordingly we cannot say with

confidence that some studies have not landed in the drawer.

Conclusion

In summary, we have confirmed that the tactile TOJ crossing effect is a very large

effect, irrespective of the specific criteria one uses to include a particular study or

not. The obtained effect size value of approximately 1.4 may be of help in planning

future studies, especially if they target the crossing effect per se, as may be the

case, for instance, in clinical and developmental studies. The picture is much less

clear regarding the moderators we identified as prevalent in the tactile TOJ literature.

If we are truly interested in them, more research and diligent reporting will be

necessary.

Competing interests

The author(s) declare no competing interests.

References

1. Heed, T. & Azanon, E. Using time to investigate space: a review of tactile temporal order judgments as a window onto spatial processing in touch. *Frontiers in Psychology* **5**, (2014).

2. Shore, D., Spry, E. & Spence, C. Confusing the mind by crossing the hands. *Cognitive Brain Research* **14**, 153–163 (2002).

3. Yamamoto, S. & Kitazawa, S. Reversal of subjective temporal order due to arm crossing. *Nat. Neurosci.* **4**, 759–765 (2001).

4. Heed, T., Backhaus, J., Röder, B. & Badde, S. Disentangling the external reference frames relevant to tactile localization. *PLoS ONE* **11**, (2016).

5. Badde, S., Röder, B. & Heed, T. Feeling a touch to the hand on the foot. *Current Biology* **29**, 1491-1497.e4 (2019).

6. Heed, T., Burbach, J., Rödenbeck, N., Habets, B. & Fuchs, X. Limb, not touch location, is coded in 3D space. 2024.02.01.578364 Preprint at https://doi.org/10.1101/2024.02.01.578364 (2024).

7. Maij, F., Seegelke, C., Medendorp, W. P. & Heed, T. External location of touch is constructed post-hoc based on limb choice. *eLife* **9**, e57804 (2020).

8. Hense, M., Badde, S., Köhne, S., Dziobek, I. & Röder, B. Visual and proprioceptive influences on tactile spatial processing in adults with autism spectrum disorders. *Autism Res.* **12**, 1745–1757 (2019).

9. Espenhahn, S. *et al.* Atypical Tactile Perception in Early Childhood Autism. *J Autism Dev Disord* (2022) doi:10.1007/s10803-022-05570-7.

10. Soto-Faraco, S., 1970- & Azañón, E. Electrophysiological correlates of tactile remapping. *Neuropsychologia* **51**, 1584–1594 (2013).

11. Moseley, G. L., Gallace, A. & Iannetti, G. D. Spatially defined modulation of skin

temperature and hand ownership of both hands in patients with unilateral complex

regional pain syndrome. *Brain* **135**, 3676–3686 (2012).

12. Vanderclausen, C., Filbrich, L., De Volder, A. & Legrain, V. Measuring the sensitivity of

tactile temporal order judgments in sighted and blind participants using the adaptive psi

method. *Atten Percept Psychophys* **83**, 2995–3007 (2021).

13. Lakens, D. Sample Size Justification. *Collabra: Psychology* **8**, 33267 (2022).

14. Cadieux, M. L. & Shore, D. I. Response demands and blindfolding in the crossed-hands

deficit: an exploration of reference frame conflict. *Multisens Res* **26**, 465–482 (2013).

15. Unwalla, K., Goldreich, D. & Shore, D. I. Exploring reference frame integration using

response demands in a tactile temporal-order judgement task. *Multisens Res* **34**, 1–32

(2021).

16. Ritterband-Rosenbaum, A., Hermosillo, R., Kroliczak, G. & van Donkelaar, P. Hand

position-dependent modulation of errors in vibrotactile temporal order judgments:the

effects of transcranial magnetic stimulation to the human posterior parietal cortex.

Experimental Brain Research **232**, 1689–1698 (2014).

17. Kóbor, I., Füredi, L., Kovács, G., Spence, C. & Vidnyánszky, Z. Back-to-front: Improved

tactile discrimination performance in the space you cannot see. *Neurosci. Lett.* **400**, 163–

167 (2006).

18. Unwalla, K., Cadieux, M. L. & Shore, D. I. Haptic awareness changes when lying down.

Scientific reports **11**, 13479 (2021).

19. Azañón, E. & Soto-Faraco, S. Alleviating the 'crossed-hands' deficit by seeing uncrossed

rubber hands. *Exp. Brain Res.* **182**, 537–548 (2007).

20. Röder, B., Rösler, F. & Spence, C. Early vision impairs tactile perception in the blind.

Curr. Biol. **14**, 121–124 (2004).

21. Cadieux, M. L., Barnett-Cowan, M. & Shore, D. Crossing the hands is more confusing

for females than males. *Experimental brain research,* **204**, 431–446 (2010).

22. Crollen, V. *et al.* Visual experience shapes the neural networks remapping touch into

external space. *Journal of neuroscience* **37**, 10097–10103 (2017).

23. Heed, T., Backhaus, J. & Roder, B. Integration of hand and finger location in external

spatial coordinates for tactile localization. *Journal of Experimental Psychology: Human*

Perception and Performance **38**, 386–401 (2012).

24. Nishikawa, N., Shimo, Y., Wada, M., Hattori, N. & Kitazawa, S. Effects of aging and

idiopathic Parkinson's disease on tactile temporal order judgment. *PLoS ONE* **10**, (2015).

25. Wada, M., Yamamoto, S. & Kitazawa, S. Effects of handedness on tactile temporal order

judgment. *Neuropsychologia* **42**, 1887–1895 (2004).

26. Schicke, T. & Röder, B. Spatial remapping of touch: confusion of perceived stimulus

order across hand and foot. *Proc Natl Acad Sci U S A* **103**, 11808–11813 (2006).

27. Gusenbauer, M. & Haddaway, N. R. Which academic search systems are suitable for

systematic reviews or meta-analyses? Evaluating retrieval qualities of Google Scholar,

PubMed, and 26 other resources. *Research Synthesis Methods* **11**, 181–217 (2020).

28. Goulet-Pelletier, J.-C. & Cousineau, D. A review of effect sizes and their confidence

intervals, Part I: The Cohen's d family. *TQMP* **14**, 242–265 (2018).

29. Lakens, D. Calculating and reporting effect sizes to facilitate cumulative science: a

practical primer for t-tests and ANOVAs. *Frontiers in Psychology* **4**, (2013).

30. Langenberg, B., Janczyk, M., Koob, V., Kliegl, R. & Mayer, A. A tutorial on using the

paired t test for power calculations in repeated measures ANOVA with interactions.

Behav Res **55**, 2467–2484 (2022).

31. R Core Team. R: A Language and Environment for Statistical Computing. R Foundation

for Statistical Computing (2023).

32. Posit team. RStudio: Integrated Development Environment for R. Posit Software, PBC
(2023).

33. Balduzzi, S., Rücker, G. & Schwarzer, G. How to perform a meta-analysis with R: A
practical tutorial. **Evidence-Based Mental Health**, 153–160 (2019).

34. Harrer, M., Cuijpers, P., Furukawa, T. & Ebert, D. D. dmetar: Companion R package for
the guide 'Doing meta-analysis in R'. (2019).

35. Schwarzer, G., Carpenter, J. R. & Rücker, G. metasens: Statistical methods for sensitivity
analysis in meta-analysis. (2023).

36. Wickham, H. *et al.* Welcome to the {tidyverse}. *Journal of Open Source Software* **4**,
1686 (2019).

37. Veroniki, A. A. *et al.* Methods to estimate the between-study variance and its uncertainty
in meta-analysis. *Research Synthesis Methods* **7**, 55–79 (2016).

38. Knapp, G. & Hartung, J. Improved tests for a random effects meta-regression with a
single covariate. *Statistics in Medicine* **22**, 2693–2710 (2003).

39. Higgins, J. P. T. & Thompson, S. G. Quantifying heterogeneity in a meta-analysis.
Statistics in Medicine **21**, 1539–1558 (2002).

40. IntHout, J., Ioannidis, J. P. A., Rovers, M. M. & Goeman, J. J. Plea for routinely
presenting prediction intervals in meta-analysis. *BMJ Open* **6**, e010247 (2016).

41. Baujat, B., Mahé, C., Pignon, J.-P. & Hill, C. A graphical method for exploring
heterogeneity in meta-analyses: application to a meta-analysis of 65 trials. *Stat Med* **21**,
2641–2652 (2002).

42. Thompson, S. G. & Higgins, J. P. T. How should meta-regression analyses be undertaken
and interpreted? *Statistics in Medicine* **21**, 1559–1573 (2002).

43. Borenstein, M. & Higgins, J. P. T. Meta-analysis and subgroups. *Prev Sci* **14**, 134–143
(2013).

44. Borenstein, M., Hedges, L. V., Higgins, J. & Rothstein, H. R. Subgroup Analyses. in

Introduction to Meta-Analysis 149–186 (John Wiley & Sons, Ltd, 2009).

doi:10.1002/9780470743386.ch19.

45. Page, M. J. *et al.* The PRISMA 2020 statement: an updated guideline for reporting

systematic reviews. *BMJ* n71 (2021) doi:10.1136/bmj.n71.

46. Azañón, E., Stenner, M., Cardini, F. & Haggard, P. Dynamic tuning of tactile localization

to body posture. *Current Biology* **25**, 512–517 (2015).

47. Azañón, E., Mihaljevic, K. & Longo, M. R. A three-dimensional spatial characterization

of the crossed-hands deficit. *Cognition* **157**, 289–295 (2016).

48. Azañón, E., Radulova, S., Haggard, P. & Longo, M. R. Does the crossed-limb deficit

affect the uncrossed portions of limbs? *Journal of Experimental Psychology: Human*

Perception and Performance **42**, 1320–1331 (2016).

49. Badde, S., Heed, T. & Röder, B. Processing load impairs coordinate integration for the

localization of touch. *Atten Percept Psychophys* **76**, 1136–1150 (2014).

50. Badde, S., Röder, B. & Heed, T. Flexibly weighted integration of tactile reference frames.

Neuropsychologia **70**, 367–374 (2015).

51. Badde, S., Röder, B. & Bruns, P. Task-irrelevant sounds influence both temporal order

and apparent-motion judgments about tactile stimuli applied to crossed and uncrossed

hands. *Attention, Perception, & Psychophysics* **80**, 773–783 (2017).

52. Crollen, V., Albouy, F., Lepore, F. & Collignon, O. How visual experience impacts the

internal and external spatial mapping of sensorimotor functions. *Scientific reports* **7**, 1022

(2017).

53. Crollen, V., Spruyt, T., Mahau, P., Bottini, R. & Collignon, O. How visual experience

and task context modulate the use of internal and external spatial coordinate for

perception and action. *Journal of Experimental Psychology: Human Perception and*

Performance **45**, 354–362 (2019).

54. Landry, S. P. & Champoux, F. Long-term musical training alters tactile temporal-order

judgment. *Multisens Res* **31**, 373–389 (2018).

55. Roberts, R. D. & Humphreys, G. W. Task effects on tactile temporal order judgments:

when space does and does not matter. *J. Exp. Psychol. Hum. Percept. Perform.* **34**, 592–

604 (2008).

56. Sambo, C. F., Gallace, A., Liang, M., Moseley, G. L. & Iannetti, G. D. The temporal

order judgement of tactile and nociceptive stimuli is impaired by crossing the hands over

the body midline. *Pain* **154**, 242–247 (2013).

57. Sharp, A., Landry, S. P., Maheu, M. & Champoux, F. Deafness alters the spatial mapping

of touch. *PLoS ONE* **13**, e0192993 (2018).

58. Unwalla, K., Kearney, H. & Shore, D. I. Reliability of the crossed-hands deficit in tactile

temporal order judgements. *Multisens Res* **34**, 387–421 (2020).

59. Cohen, J. *Statistical Power Analysis for the Behavioral Sciences*. (Erlbaum, Hillsdale, NJ

u.a., 1988).

60. Zwetsloot, P.-P. *et al.* Standardized mean differences cause funnel plot distortion in

publication bias assessments. *eLife* **6**, e24260 (2017).

61. Simonsohn, U., Nelson, L. D. & Simmons, J. P. P-curve: A key to the file-drawer.

Journal of Experimental Psychology: General **143**, 534–547 (2014).

62. Hogben, D., Pinkham, R. S. & Wilk, M. B. The Moments of the Non-Central t-

Distribution. *Biometrika* **48**, 465–468 (1961).

63. Hedges, L. V. Distribution Theory for Glass's Estimator of Effect size and Related

Estimators. *Journal of Educational Statistics* **6**, 107–128 (1981).

64. Tothfalusi, L. & Endrenyi, L. An Exact Procedure for the Evaluation of Reference-Scaled

Average Bioequivalence. *AAPS J* **18**, 476–489 (2016).

Appendix A

Table A1. Search String for Each Database

Database	Search string
BASE	(tactile OR hand OR arm) temporal order judg* doctype:121; english manually selected
EMBASE	('tactile temporal order' OR ((hand OR tactile OR arm) AND temporal AND order AND judg*)) AND ([article]/lim OR [article in press]/lim) AND [english]/lim
PUBMED	(tactile OR hand OR arm) AND temporal order judg* AND (english[Filter]) NOT(review pt])
Web of Science	ALL=((hand OR tactile OR arm) temporal order judg*); articles manually selected

Appendix B

Transformation of noncentral t variance to variance of Cohen's d_z

To calculate the standard error of Cohen's d_z, typically either the standard deviation

of the mean difference or the correlation between the two measurements (crossed

and uncrossed conditions) is needed[28,29]. However, these data were not available in

all studies. Therefore, to avoid using an estimated correlation for the standard error

calculation, the relationship between t and Cohen's d_z was utilized to calculate the

standard errors of the effect sizes:

1. Variance of the random variable t which follows a non-central t-

618   distribution[62] (using an approximation[63,64]):

$$Var(t) = \frac{v(1 + \mu^2)}{v - 2} - \frac{\mu^2 * v}{2} * \left(1 - \frac{3}{4 * v - 1}\right)^{-2} \tag{B1}$$

619   where v is the degrees of freedom and μ is the noncentrality parameter.

620   2. Given the relation between t and Cohen's d_z[29], t can be calculated from d_z,

621   and the sample size:

$$Cohen's\ d_z = \frac{t}{\sqrt{n}} \text{ and therefore } t = d_z * \sqrt{n} \tag{B2}$$

622   3. Substitution of v and μ for within-subject designs: where v = n-1 and

623           $\mu = t = d_z * \sqrt{n}$:

$$Var(t) = \frac{(n - 1)\left(1 + (d_z * \sqrt{n})^2\right)}{(n - 1) - 2} - \frac{(d_z * \sqrt{n})^2 * (n - 1)}{2}$$
$$* \left(1 - \frac{3}{4 * (n - 1) - 1}\right)^{-2} \tag{B3}$$

624   4. Given equation (B2) the relationship between $Var(t)$ and $Var(d_z)$ can be

625   expressed as:

$$Var(t) = n * Var(d_z) \tag{B4}$$

626   therefore:

$$Var(d_z) = \frac{Var(t)}{n} \tag{B5}$$

628   5. Substitution of $Var(t)$ into the equation for $Var(d_z)$:

$$Var(d_z) = \frac{1}{n} * \left(\frac{(n-1)\left(1 + (d_z * \sqrt{n})^2\right)}{(n-1) - 2} - \frac{(d_z * \sqrt{n})^2 * (n-1)}{2} \right.$$
$$\left. * \left(1 - \frac{3}{4*(n-1) - 1}\right)^{-2} \right)$$

(B6)

629   6. Simplified to:

$$Var(d_z) = \frac{1}{n} * \frac{n-1}{(n-1) - 2} + d_z^2$$
$$* \left(\frac{n-1}{(n-1) - 2} - \frac{1}{\left(1 - \frac{3}{4*(n-1) - 1}\right)^2} \right)$$

(B7)

630   7. Derive the standard error SE(d_z), which in this case equals to the standard
631   deviation and is therefore defined as the square root of the variance:

$$SE(d_z)$$
$$= \sqrt{\frac{1}{n} * \frac{n-1}{(n-1) - 2} + d_z^2 * \left(\frac{n-1}{(n-1) - 2} - \frac{1}{\left(1 - \frac{3}{4*(n-1) - 1}\right)^2} \right)}$$

(B8)

632