circEPSTI1 Enables Dengue Innate Immune Evasion by Sponging miR-942-5p, a Repressor of AKT1 and the Viral Genome

Nilanjana Das [a1], Pandikannan Krishnamoorthy [a1], Kushal Ramrao Junghare [a], Athira S Raj [a],

Priyanka Saha [b], Kuhu Pal [b], Shouvanik Adhya [c], Atul Garg [d], Dheeraj Khetan [e], Ashok Kumar [f],

Himanshu Kumar [a, g*]

Affiliations:

[a] Department of Biological Sciences, Laboratory of Immunology and Infectious Disease Biology,

Indian Institute of Science Education and Research (IISER) Bhopal, Bhopal-462066, MP, India

[b] Department of Microbiology, College of Medicine and JNM Hospital, Kalyani, West Bengal.

[c] Department of Forensic Medicine and Toxicology, College of Medicine and JNM Hospital,

Kalyani, West Bengal.

[d] Department of Microbiology, Sanjay Gandhi post graduate Institute of Medical sciences,

Lucknow

[e] Department of Transfusion Medicine, Sanjay Gandhi Postgraduate Institute of Medical

Sciences, Lucknow, India.

[f] Department of Biochemistry, All India Institute of Medical Sciences (AIIMS), Bhopal, Madhya

Pradesh, India

[g] Laboratory of Host Defense, WPI Immunology, Frontier Research Centre, Osaka University,

Osaka 5650871, Japan

1 These authors contributed equally.

*Corresponding author: H Kumar, Department of Biological Sciences, Laboratory of

Immunology and Infectious Disease Biology, Indian Institute of Science Education and Research

24   (IISER) Bhopal, AB-3, Room No. 220, Bhopal By-pass Road, Bhauri, Bhopal 462066, MP,

25   India. Tel: +91 755 6691413; Fax: +91 755 669 2392; E-mail: hkumar@iiserb.ac.in

26   Running head: **circEPSTI1 regulates antiviral miR-942-5p during DENV infection**

ABSTRACT

Circular RNAs (circRNAs) are recently emerging transcripts that are gaining attention for their potential role in host-pathogen interactions. However, the role of circRNAs in Dengue virus (DENV) infection remains unexplored. In this study, circEPSTI1 induction was observed in DENV-infected peripheral blood mononuclear cells (PBMCs) and plasma of Dengue patients. circEPSTI1 induction is interferon-dependent, as IRF3-knockout cells completely abolished its induction. circEPSTI1 enhances DENV infection by sponging the expression of hsa-miR-942-5p. The antiviral role of hsa-miR-942-5p is mediated by bidirectional inhibition of conserved genomic sequences of the DENV genome across NS1, NS3 and NS5 and the host AKT1 signaling pathway. circEPSTI1 sponges hsa-miR-942-5p and relieves its inhibition over DENV genomic RNA and the host AKT1 signaling pathway, thus leading to increased AKT1 activation and DENV replication. These findings highlight the diagnostic and therapeutic potential of circEPSTI1 and the novel circEPSTI1-miR-942-5p-AKT1 axis that regulates the host immune response against DENV infection.

Keywords: circRNA; Dengue; Innate immunity; microRNA; Host-pathogen interaction

Introduction:

Dengue is an emerging global health threat due to the increased outbreaks and its distribution to over half of the global population (1). Dengue is linked to symptoms ranging from moderate fever in the case of Dengue fever (DF) to thrombocytopenia, vascular leakage in the case of Dengue with Warning signs (DWS), and excessive production of cytokines and organ dysfunction in the case of Severe Dengue (SD) (2). Different serotypes of the Dengue virus (DENV-1, DENV-2, DENV-3, and DENV-4) are transmitted through *Aedes* aegypti. The DENV genome is an 11kb positive-sense RNA genome consisting of a highly conserved 5'-UTR, a lengthy ORF coding polyprotein, and a very organized, unstable 3'-UTR. Secondary infection with a different serotype can lead to severe clinical outcomes through the process known as antibody-dependent enhancement, which limits the DENV vaccine development (3).

Host innate immunity plays a crucial antiviral role during DENV infection (4). DENV is sensed by Pathogen recognition receptors (PRRs) such as Toll-like receptors (TLR-3/7) and Retinoic-acid-like receptors (RLRs), which initiate type-I Interferons (IFN-α, IFN-β) production to limit DENV replication. RIG-I senses the DENV RNA and recruits a sole adaptor protein, IPS-1 (also known as MAVS, VISA, and CARDIF), subsequently activating transcription factor IFN regulatory factor 3 (IRF3) and NFκB to produce Type-I, Type-III IFN and inflammatory cytokines, respectively (5). However, DENV utilizes multiple strategies to evade the antiviral innate immunity and establish the infection, skew physiology, and develop a range of clinical symptoms (6).

Non-coding RNAs play an important role in regulating genes and various signaling pathways

that are part of the antiviral immune responses (3). Non-coding RNAs can be categorized into

small non-coding RNAs, such as microRNAs (length less than 200 nucleotides), and long non-

coding RNAs, such as lncRNAs (length more than 200 nucleotides). An emerging class of

transcripts, circRNAs, single-stranded, covalently-closed RNAs without free ends, formed

through back-splicing, are shown to have higher diagnostic and therapeutic potential due to their

excellent stability and presence in diverse cell types and extracellular fluids. CircRNAs were

reported to regulate various biological processes in diseases like cancer, neurological disorders,

and autoimmune diseases (7).

CircRNAs exert their mechanistic role by binding to RNA-binding proteins (RBPs) and are also

well known to sponge microRNAs (miRNAs) (8). miRNAs are small non-coding RNAs that

post-transcriptionally regulate the expression of genes by binding to the 3'-UTR of mRNAs.

miRNAs can either bind to the mRNAs of the innate immune genes or bind to the viral genomes

of RNA viruses and regulate the viral replication (9). The regulatory functions of miRNAs are

further regulated by circRNAs and thus form a tight layer of competing endogenous RNA

network to regulate various signaling pathways under different physiological conditions. The

role of circRNAs in regulating innate antiviral immune response against viral infection is not

well-studied. Most circRNA studies are related to DNA viruses, and no previous studies have

explored the mechanistic role of circRNAs in DENV infection (10).

In this study, after the comprehensive computational analysis of the high-depth clinical RNA-seq

dataset, circEPSTI1 was identified to be highly expressed and upregulated in the PBMCs of the

Dengue patients. The upregulation was validated in the plasma of Indian Dengue patients

recruited for this study. Further, the characterization of circEPSTI1 revealed that the

upregulation of circEPSTI1 is interferon-dependent, and the knockdown and overexpression of

circEPSTI1 suppress and enhance the DENV replication, respectively. Mechanistically,

circEPSTI1 exerts its pro-viral role by sponging hsa-miR-942-5p, a miRNA that suppresses

DENV replication by targeting the DENV genomic RNA at NS1, NS3 and NS5 loci positions

and host AKT1 signaling pathway. The sponging of hsa-miR-942-5p by circEPSTI1 relieves its

suppression on AKT1, a key protein in cellular survival that DENV hijacks for replication, thus

facilitating a conducive microenvironment for viral propagation. Thus, circEPSTI1 functions as a

competing endogenous RNA (ceRNA) that modulates the host antiviral response by regulating

the activity of miR-942-5p during DENV infection.

RESULTS:

Transcriptome-wide analysis of circRNAs dysregulation in Dengue-infected patients
identifies circEPSTI1 upregulation:

The circRNAs are known to be dysregulated in the blood components upon different
pathological conditions such as cancer and viral infections. However, the circRNAs dysregulated
in Dengue patients compared to uninfected cases have not yet been identified. Quantifying
circRNAs from RNA-seq datasets is possible from total-RNA libraries but not polyA-enriched
libraries. Upon mining public transcriptome datasets for the DENV-infected clinical samples,
only the dataset, GSE94892, was found to be derived from the total RNA library without polyA-
enrichment (11). Identification of circRNAs was highly influenced by the depth of sequencing
and the number of reads per sample. The sequencing depth was higher (more than 80 million
reads per sample) and not poly-A enriched in this dataset. As illustrated in **(Fig. 1A)**, the
distribution of mapped, unmapped, and circular reads supported accurate quantification of
circular transcripts. Principal Component Analysis (PCA) revealed a clear segregation of Control
and DENV-infected groups based on circRNA expression profiles **(Supplementary Fig. 1A)**,
indicating distinct transcriptomic alterations during infection. Differential expression analysis
identified 180 significantly dysregulated circRNAs (padj < 0.05, $|\log FC| > 1$), including 142
upregulated and 36 downregulated circRNAs, denoted by the volcano plot **(Fig. 1B)**. To identify
clinically relevant circRNAs for translational application in clinical settings, it is important to
find highly expressed circRNAs. The expression of the top circRNAs in each sample was
visualized through the violin plot **(Supplementary Fig 1B**) and the heatmap **(Supplementary**
Fig. **1C).** The dot plot **(Figure 1C)** representing the normalized counts of the top significant

circRNAs reveals hsa_circEPSTI1_001(circEPSTI1) with consistently higher expression across

every sample (above the red dotted line denoting the normalized expression value of more than

5) and also significantly upregulated upon DENV-infected patients (triangle datapoints)

compared to the control group (circle datapoints). circEPSTI1 (hsa_circ_0000479, in circBase) is

derived by the back-splicing between the exon 2 and exon 6 of EPSTI1 pre-mRNA, leading to

the 375 bp transcript (**Supplementary Fig. 2A**). Circular RNAs are not expected to be amplified

from genomic DNA (gDNA) as well as the cDNA synthesized through the oligo-dT method, as

they lack poly-A ends. To ensure the circRNA is amplified but not the linear one, the divergent

primer flanking the backsplice junction was designed to amplify the circEPSTI1, and the

convergent primer was designed to amplify the linear ESPTI1 **(Fig. 1D)**. The genomic DNA and

cDNA generated from human PBMCs through random hexamers were amplified with the

divergent and convergent primers. circEPSTI1 was amplified only in the cDNA amplified

through the random hexamer method but not gDNA and cDNA generated through the oligo-dT

method. However, linear EPSTI1 was amplified in both the gDNA and cDNA from PBMCs

(Fig. 1D), suggesting the amplified product by divergent primer belongs to circEPSTI1 but not

the linear counterpart. Similar results were obtained in the A549 cell line **(Supplementary Fig. 2**

B). Sanger sequencing of the amplified bands revealed the presence of the sequence of the back-

splice junction of circEPSTI1 **(Fig. 1E)**. CircRNAs generally have higher stability than linear

RNA. Actinomycin D (ActD) inhibits transcription by binding to DNA and inhibits the activity

of RNA polymerase to synthesize new RNA, and hence can be utilized for measuring the

stability of circRNA, compared to the linear RNA. Upon ActD treatment in PBMCs **(Fig.1F)** and

A549 **(Supplementary Fig. 2C)**, linear EPSTI1 decreases, whereas circEPSTI1 remains stable,

suggesting the amplified product is a circRNA. RNase R is a 3'-5' exoribonuclease that can

selectively degrade linear RNAs but not circular RNAs. RNAse R treatment in PBMC **(Fig. 1G)**

and A549 **(Supplementary Fig. 2D)** showed that the circEPSTI1 level remained unchanged,

whereas linear EPSTI1 almost completely degraded. To gain insight of circEPSTI1 cellular

localization, the nuclear and cytoplasmic fractions were isolated, and the level of enrichment of

circEPSTI1 were measured. circEPSTI1 is predominantly distributed in the cytoplasm of the

PBMCs **(Fig. 1H)** and A549 **(Supplementary Fig. 2E)**. To strengthen our biochemical

observations, cell imaging was performed through Fluorescence in situ Hybridization (FISH)

utilizing a Cy3-labelled probe targeting circEPSTI1. The enrichment in the cytoplasm of the

A549 cells was observed as visualized through confocal microscopy **(Fig. 1I)**. Overall,

computational analysis and multiple experiments suggested that the circEPSTI1 is one of the

highly expressed circRNAs in DENV-infected patients and possesses the properties of the

circRNAs and is widely distributed in the cytoplasm.

DENV infection upregulates circEPSTI1, and induction is interferon-dependent

To validate the prediction from the RNA-seq results, clinical recruitment of DENV patients was

performed. RNA from the plasma of healthy controls and Dengue patients was isolated, and the

level of circEPSTI1 was measured by qPCR. circEPSTI1 level was found to be upregulated

significantly in the DENV patients compared to the healthy controls **(Fig. 2A)**.

PBMCs from the healthy volunteers were isolated and infected with DENV ex vivo, and

circEPSTI1 level was found to be upregulated **(Fig. 2B)**, as observed in the plasma. Next, the

copy number of circRNA in A549 cell was determined because it is crucial for therapeutic

manipulation. To this end a highly sensitive, digital-PCR method was utilized to measure the

copy number of circEPSTI1, and it is observed that the copy number of circEPSTI1 is very low

in uninfected cells, and upon infection, it increases to 750 copies (more than 7 times) per ul **(Fig.**

2C). This suggests that the expression and copy number of circEPSTI1 are increased upon

DENV infection. Further, to understand the dynamics of the induction of circEPSTI1, different

cell lines were infected with DENV, and the circEPSTI1 level was measured post 12, 24, 36, and

48 hours of infection. circEPSTI1 levels were not upregulated till 12 hours post-infection, and

subsequently, the expression was increased in PBMCs, A549, MRC5, and HEPG2 cell lines

(Fig. 2D). Overall, circEPSTI1 induction upon DENV infection was observed in diverse cell

types, and the magnitude and extent of expression vary between different cell types and DENV

infectivity (**Supplementary Fig. 3**). Upon DENV infection, the primary innate antiviral response

is initiated by TLR-3 signaling and RLR signaling that induces type-I interferons and

inflammatory cytokines. Upon TLR-3 ligand, Poly(I:C) stimulation, circEPSTI1 was induced as

observed in the case of viral infection **(Fig. 2E)**, and treatment of cells with type-I interferons,

IFN-α and IFN-β, also induced circEPSTI1 expression, suggesting the role of interferon-

mediated induction, rather than DENV-specific induction **(Fig.2F, 2G)**. To explore the

involvement of type-I interferon signaling for circEPSTI1 induction and whether it is RLR-

dependent, the innate immune RLR sensor, RIG-I, RLR signaling adaptor IPS-1 (also known as

MAVS, VISA, CARDIF), and the downstream interferon-inducing transcription factor, IRF3,

were knocked out using the CRISPR-Cas9 technique in A549 cells. DENV infection was

performed in the WT, RIG-I KO, IPS-1 KO, and IRF3 KO cells at different time points.

circEPSTI1 was induced in WT cells but not induced in the cells deficient in RIG-I, IPS-1, and

IRF3, suggesting the interferon-dependent induction of circEPSTI1 **(Fig. 2H)**. In the previous

section, cellular localization of circEPSTI1 was shown to be predominantly present in the

cytoplasm. Therefore, cellular localization of circEPSTI1 was imaged upon DENV infection by

confocal microscopy as described previously. The results revealed the significant enrichment of

circEPSTI1 in the cytoplasm of DENV-infected cells **(Fig. 2I)**. Taken together, circEPSTI1

induction upon DENV infection is observed to be interferon-dependent and is predominantly

expressed in the cytosolic compartment.

circEPSTI1 regulates DENV infection

The effect of induction of circEPSTI1 over DENV replication was explored by the siRNA-

mediated knockdown study. siRNA was designed to target the back-splice junction of

circEPSTI1, and the knockdown of circEPSTI1 was verified by RT-PCR in both human PBMC

and A549 cells, as shown in **Fig. 3A**. The siRNA reduced 45% to 60% expression in these cells.

Knockdown of circEPSTI1 led to a reduction in the viral load in both PBMC and A549 cells and

culture supernatant **(Fig. 3B & C)**. Similar effects were observed in THP1 and MRC5 cell lines

(**Supplementary Fig. 4A and 4B**). To further investigate the effect of circEPSTI1 knockdown

on DENV viral load, western blot, Flow cytometry, and confocal microscopy were performed

using DENV NS5-specific antibodies. At the protein level, the dengue NS5 level was suppressed

upon knockdown of circEPSTI1 **(Fig. 3D)**. Flow cytometry analysis revealed a reduction of

DENV in the circEPSTI1 knockdown **(Fig.3E)**, and confocal microscopy also showed a

reduction in the intensity of DENV in circEPSTI1 knockdown cells **(Fig. 3F)**. Collectively, our

results with multiple techniques demonstrate that circEPSTI1 knockdown was found to suppress

DENV replication. To strengthen the findings, circEPSTI1 overexpression was performed by

cloning circEPSTI1 in the mc2mNeonV2 vector, and the overexpression was verified by RT-

PCR, which shows a more than 1400-fold increase in circEPSTI1 **(Fig. 3G)**. DENV viral load

was increased to a higher extent upon circEPSTI1 overexpression in A549 cells and in culture

supernatant **(Fig. 3H & I).** The finding was further validated through western blot **(Fig. 3J)**, flow

cytometry **(Fig. 3K)**, and confocal microscopy **(Fig. 3L)**. Altogether, the DENV viral load

increased upon circEPSTI1 overexpression, as demonstrated by various assays. Next, the innate

antiviral responses were measured, interferon-induced genes, such as IFN-β and OAS2, were

significantly increased and suppressed upon knockdown **(Fig. 3M)** and overexpression **(Fig. 3N)**

of circEPSTI1, respectively. The levels of IFN-inducible protein, IP-10 and inflammatory

cytokine IL-6, in culture supernatant upon circEPSTI1 knockdown, were significantly enhanced

(Fig. 3O). Similarly, IP-10 and IL-6, in culture supernatant upon circEPSTI1 overexpression,

were significantly reduced **(Fig. 3P)**. Overall, knockdown of circEPSTI1 reduces the viral load

and increases IFNβ and inflammatory cytokines, whereas overexpression of circEPSTI1

increases the viral load and decreases the level of IFNβ and inflammatory cytokines, suggesting

the pro-viral role of circEPSTI1 facilitating DENV replication.

circEPSTI1 sponges miR-942-5p during DENV infection

Circular RNAs exert their regulatory functions mainly by sponging miRNAs and stabilizing the

target transcripts, which subsequently result in sustained protein expression. The miRNAs

binding to circEPSTI1 are predicted by three widely utilized algorithms- miRanda, PITA, and

TargetScan. miRanda predicts the sequence complementarity between the circRNA and miRNA

and the thermodynamic stability of the interaction, and calculates the binding energy and score.

hsa-miR-942-5p was the top miRNA with the lowest free energy and the highest miRanda score.

PITA considers the accessibility and miRNA-target pair and computes free energy, and Target

Scan considers seed region complementarity, conservation, and number of sites. miR-942-5p was

the common miRNA predicted by all three tools with a stringent cutoff **(Fig. 4A)**. To validate the

predictions, the circRNA sequence was cloned downstream of luciferase under the CMV

promoter as shown in schematic of **Fig. 4B**. Upon overexpression of hsa-miR-942-5p, the

luciferase activity of the circEPSTI1-luc vector was reduced significantly, indicating that hsa-

miR-942-5p binds to circEPSTI1 and suppresses its expression. In contrast, when the hsa-miR-

942 inhibitor was introduced which sequester endogenous hsa-miR-942-5p, the luciferase

activity was not reduced, suggesting the suppression is not due to non-specific binding **(Fig. 4B)**.

To further validate the finding, the predicted binding sites of miR-942 in the circ-EPSTI1

sequence were mutated, and this mutated construct failed to reduce the luciferase activity upon

hsa-miR-942 overexpression, suggesting that miR-942 binds to circEPST1 with its seed

sequence at the predicted sites **(Fig. 4C)**. The miRNA targets the transcript through a

multiprotein complex known as RNA-induced silencing complex (RISC) and Argounate-2

protein (AGO2) is one of essential component of RISC for miRNA-mediated silencing. To

explore the involvement of AGO2 in circEPSTI1 and hsa-miR-942-5p interaction, the AGO2

pulldown assay was performed using a FLAG-specific antibody. The cells transfected with hsa-

miR-942 showed greater than four fold enhancement of circEPSTI1 levels in the pull-down

fraction compared to the NC-transfected cells **(Fig. 4D)**. Conversely, to further validate the

findings, circEPSTI1 was pulled down using biotin probes and hsa-miR-942-5p levels were

enriched in the pull-down fraction in circEPSTI1 overexpressed cells compared to the control

vector transfected cells, suggesting circEPSTI1 and hsa-miR-942-5p bind with each other **(Fig.**

4E). To assess the functional consequences of this interaction, upon knockdown of circEPSTI1,

hsa-miR-942-5p levels were quantified. hsa-miR-942-5p levels were significantly upregulated

upon circEPSTI1 knockdown in PBMCs and A549 cells compared to the negative control

transfected cells **(Fig. 4F)** suggesting the inverse regulatory relationship between circEPSTI1

and hsa-miR-942-5p. Interestingly, simultaneous siRNA-mediated knockdown of circEPSTI1

and inhibition of hsa-miR-942-5p through the inhibitor restored the DENV viral load achieved

only by knocking down circEPSTI1. **(Fig. 4G)**. This implies that the pro-viral activity of

circEPSTI1 is mediated through the regulation of hsa-miR-942-5p expression. Further, miR-942-

5p expression was suppressed upon DENV infection at different time points in PBMC, A549,

MRC5, and HEPG2 cell lines, negatively correlated with the induction of circEPSTI1

expression. **(Fig. 4H).**

hsa-miR-942-5p targets the DENV genome and inhibits DENV replication

MicroRNAs bind to the genome of RNA viruses such as Influenza and regulate their replication

(9, 12). The miRanda tool was utilized to find whether miR-942-5p can bind to the DENV

genomic RNA across all four serotypes **(Fig. 5A)**. Based on the conservation of the site across

more than one DENV serotype and the free energy, 8 sites in the DENV genome were identified.

NS5 bears 4 sites, NS1 bears 3 sites, NS3 bears 2 sites, and E bears a single site

(Supplementary Fig. 5A). To validate the prediction, NS5, NS3, and NS1 were cloned

downstream to the luciferase, and the luciferase activity upon miR-942-5p overexpression was

measured. NS1 luciferase activity was suppressed the most, followed by NS3 and NS5. When

NS2B with no predicted binding site was cloned downstream of luciferase, luciferase activity

suppression was not observed, suggesting the predictions are not random and non-specific **(Fig.**

5B). To determine the involvement of AGO2-mediated miR-942-5p DENV viral RNA

interaction, AGO2 pulldown was performed and DENV RNA was enriched in the pull-down

fraction compared to the control uninfected cells **(Fig. 5C)**. miR-942-5p overexpression in A549

and PBMC also reduced the DENV levels significantly. The same effect was not observed when

the miR-942-5p inhibitor was used **(Fig. 5D).** The suppression of DENV load was also observed

in the supernatant in A549 cells **(Fig. 5E)**. The level of NS1 and NS5 at the protein level was

evaluated through the western blot. DENV protein levels were reduced upon miR-942-5p

overexpression, suggesting the anti-viral role of miR-942-5p upon DENV infection **(Fig. 5F)**.

Further, the reduction of DENV NS1 and NS5 was validated by FACS **(Fig. 5G)** and

Immunofluorescence through confocal microscopy **(Fig. 5H)** using NS1 and NS5-specific

antibodies. Further, the levels of interferon-inducible protein, IP-10 and inflammation-associated

cytokines such as IL-6 was measured through ELISA, and it was observed that upon miR-942-5p

overexpression, IL-6 and IP-10 levels were significantly induced **(Fig. 5I)**. Overall, miR-942-5p

suppresses DENV infection by primarily targeting the DENV NS1 and NS5 and to some extent

NS3 to modulate the host innate immune response.

miR-942-5p targets the host Akt-signaling pathway and regulates DENV infection:

To identify the host targets of miR-942-5p, miRabel tool was utilized, which reports the top host

genes based on different miRNA-binding tools (13). Upon performing the pathway enrichment

analysis of miR-942-5p targets, we found AKT1 signaling pathway as one of the enriched

pathways, next to metabolic and cancer pathways **(Fig. 6A)**. Furthermore, NS1, NS3, and NS5

were found to interact with various proteins involved in PI3K/AKT signaling (**Supplementary**

Fig. 5B). Various factors activate AKT1 signaling, and the phosphorylation of AKT1 (S473)

indicates one of the activation mechanisms of the pathway (14). The level of p-AKT1 (S473)

decreases in initial time points and increases until 48 hours post-DENV infection **(Fig. 6B)**,

suggesting the activation of the AKT1 signaling pathway upon DENV infection. Among the

target genes predicted to be bound by miR-942-5p, AKT1 was the central mediator of this

pathway. The 3'UTR of AKT1 was cloned downstream to luciferase, and the luciferase activity

was suppressed when miR-942-5p was overexpressed in HEK293T cells. However, miR-942-5p

inhibitor did not suppress the luciferase activity **(Fig. 6C)**. Further mutation of the miR-942-5p

binding site in AKT1 UTR could not reduce the luciferase activity, proving the specific

interaction of miR-942-5p with the 3'-UTR of miR-942-5p **(Fig. 6D)**. AGO2-pulldown upon

miR-942-5p showed enrichment of AKT1 level. The enrichment was not observed when the

miR-942-5p inhibitor was used **(Fig. 6E)**. At the protein level, AKT1 was suppressed upon miR-

942-5p overexpression during DENV infection **(Fig. 6F).** Overall, miR-942-5p was found to

target viral genomic RNA as well as the host AKT1 RNA and modulate the AKT1 signaling

pathway. To know the role of AKT1 in DENV infection, siRNA-mediated knockdown of AKT1

was performed. The knockdown of AKT1 reduced the DENV viral load, suggesting the pro-viral

role of AKT1 **(Fig. 6G)**. At the protein level, the reduction in the protein level of NS5 was

observed upon AKT1 knockdown **(Fig. 6H)**. The reduction in viral load upon siAKT1

knockdown was further validated by FACS **(Fig. 6I)** and confocal microscopy **(Fig. 6J)**. IFN-β

and interferon-induced gene OAS2, were significantly increased upon siAKT1 knockdown **(Fig.**

6K). The levels of interferon induced gene IP-10 and inflammatory cytokine IL-6 were

quantified through ELISA and were found to be induced significantly upon AKT1 knockdown

(Fig. 6L). To know the role of circEPSTI1 in miR-942-5p-mediated AKT1 reduction,

sicircEPSTI1 was transfected at increasing concentrations, and it was observed that AKT1 at the

protein level was decreased upon the increased dose of sicircEPSTI1. Upon DENV infection,

when the siRNA-mediated suppression of circEPSTI1 was performed, its sponging activity on

miR-942-5p was reduced, leading to more availability of miR-942-5p and hence leading to

higher AKT1 suppression **(Fig. 6M)**. Overall, circEPSTI1 regulation of antiviral miR-942-5p

that interacts with both viral RNA and host AKT1 can be the immune evasion strategy of DENV,

utilizing the AKT1 signaling pathway. The pharmacological inhibition of AKT1 signaling was

performed by Wortmannin treatment at two different doses. The reduction in viral load upon

Wortmannin treatment was confirmed by Western blot analysis **(Fig. 6N)** and confocal

microscopy **(Fig. 6O)**, suggesting a promising target molecule to target this axis and a promising

DENV therapeutic target.

Discussion

CircRNAs have recently gained attention due to their diagnostic and therapeutic potential across different pathological conditions (8). In recent years, thousands of circRNAs have been identified and characterized across several tissues and many diseases (7). The role of circRNAs in regulating immune responses against the virus infection has not been well studied. circRNAs play an important role in the regulation of the miRNAs that, in turn, can regulate the genes, subsequently cellular signaling pathways, and finally disease outcome (7). The regulation by non-coding RNAs impacts cell signaling and shapes the complex host-pathogen interactions and the disease pathology (3). CircRNA detection methods from the high-throughput RNA-sequencing methods demand non-polyA enriched libraries and higher read depth to detect a significant number of circRNAs. Upon DENV infection, the role of the circRNAs has not been previously studied. Using microarray, two circRNAs were identified to be dysregulated between pretreatment and post-treatment whole blood samples of non-severe dengue patients (15). This study also did not explore the functional role of the identified circRNAs (15).

In this study, using computational analysis of RNA-sequencing datasets, circEPSTI1 was found to be significantly upregulated in the PBMCs of dengue patients compared to the control group and also bears higher expression in the blood (**Fig. 1C**). To validate the findings, the dengue patients were recruited from India, and using RT-PCR, the circEPSTI1 was found to be upregulated in the plasma of severe and non-severe dengue patients compared to the healthy controls, revealing its diagnostic potential (**Fig. 2A**).

circEPSTI1 was reported to be upregulated in several cancers, including cervical cancer (16, 17),

osteosarcoma (18), breast cancer (19), non-small cell lung cancer (20, 21), ovarian cancer (22),

and oral cancer (23). Increased expression of circEPSTI1 was reported to correlate with

increased cell proliferation, migration, invasion, and overall tumor progression (16, 17, 20).

Combination of circEPSTI1 and MIF (Macrophage Migration Inhibitory Factor) was reported to

have diagnostic value in endometrial cancer (24). Apart from cancer, circEPSTI1 was also found

to be a potential biomarker candidate in SLE (25). However, the expression of circEPSTI1 in

virus infections and its role have not been previously studied. Hence, this study aimed to explore

the functional role of circEPSTI1 in regulating the host immune response during DENV

infection.

Host pathogen recognition receptor (PRRs) plays an important role in recognizing the viral

elements to initiate host innate immune responses (26). Upon DENV infection, TLRs such as

TLR3 (27) and TLR7 (28), RLRs such as RIG-I (29), were reported to recognize DENV RNA

and initiate downstream interferon signaling. circEPSTI1 levels were upregulated upon PolyIC

stimulation (a TLR3 Ligand), and recombinant IFN-α and β treatment. Interestingly, circEPSTI1

induction upon DENV infection was not observed in RLR sensor, RIG-I, RLR signaling adaptor,

IPS-1, and the transcription factor IRF-3 knockout, suggesting interferon-dependent induction

(**Fig. 2H**). Further functional analysis of circEPSTI1 by siRNA-mediated knockdown and

overexpression revealed its pro-viral role in DENV replication **(Fig. 3)**.

Circular RNAs were reported to exert their function by binding to RNA-binding proteins (RBPs)

and acting as a sponge for miRNAs (3, 8). To further elucidate the mechanism by which

circEPSTI1 regulates DENV infection, the miRNA sponging activity of circEPSTI1 was

explored. Through three different algorithms, the top host miRNA predicted to be targeted by

circEPSTI1 was found to be hsa-miR-942-5p. The cytoplasmic localization of circEPSTI1

further supports its functional relevance in regulating miRNA and protein interactions during

DENV infection (**Fig. 2I**). Since circRNAs are known to interact with RNA-binding proteins

(RBPs) and components of the RNA interference machinery, the cytoplasmic presence of

circEPSTI1 may facilitate efficient sequestration of miR-942-5p, thereby modulating the

downstream immune and signaling responses. This prediction is validated using the luciferase

assay and AGO2-pull-down experiments (**Fig. 4**). hsa-miR-942-5p has been previously reported

to play a role in antiviral responses and immune regulation upon HCV infection (30). However,

its role in DENV infection has not yet been explored. The level of hsa-miR-942-5p was

suppressed upon DENV infection (**Fig.4I**).

miRNAs exert their functions by binding to the 3'-UTR of the genes and to the viral genome of

RNA viruses and regulating the host immune responses (9, 12). To explore whether hsa-miR-

942-5p can bind to the DENV genome, the genomic sequences of all four serotypes of DENV

were scanned for the presence of miR-942-5p binding sites. A total of 8 binding sites were

identified and conserved in more than two serotypes (**Fig. 5A**). Out of which, NS5, NS3, and

NS1 bear more sites. Prediction was further validated using luciferase assay and AGO2-

pulldown experiments. hsa-miR-942-5p overexpression reduced DENV replication by binding to

DENV NS5, NS3, and NS1. However, the host genes and the pathways that interplay with this

interaction were unclear.

To identify the host hsa-miR-942-5p targets, the miRabel tool that reports the top gene

candidates based on the results of multiple algorithms was utilized. The enrichment analysis of

the top hits identified the genes belonging to broader cancer and metabolic pathways, followed

by the specific AKT1 signaling pathway (**Fig. 6A**). DENV proteins were reported to interact

with the host AKT1 signaling proteins and modulate the pathway at the protein level (31). The

suppression of miR-942-5p upon DENV infection, mediated by circEPSTI1, likely results in the

dysregulation of downstream signaling pathways that promote viral replication. Specifically,

circEPSTI1-mediated miR-942-5p sponging was linked to the activation of the AKT1 signaling

pathway, which has been implicated in supporting viral replication and immune evasion. This

highlights a novel mechanism by which DENV manipulates host cellular signaling through

circRNA-mediated miRNA regulation.

Among the AKT1 signaling-related targets, the effector protein AKT1 was predicted to be

targeted by hsa-miR-942-5p and validated using functional experiments. AKT1 is a

serine/threonine kinase that plays a crucial role in cell survival, metabolism, and innate immune

regulation. The AKT1 signaling activation is mediated through the phosphorylation of AKT1 at

Serine 473 residue (p-AKT1). Multiple viruses utilize this pathway to promote their replication

(14, 32). However, the interplay of AKT1 signaling and virus immunity is complex and was

reported to be proviral as well as antiviral in different virus infections. A previous study reported

the pro-viral role of AKT1 signaling upon DENV infection (33).

In this study, upon DENV infection, p-AKT1 level was increased, indicating the activation of

AKT1 signaling. Both siRNA-mediated knockdown and pharmacological inhibition by

wortmannin increase the DENV replication. Overexpression of hsa-miR-942-5p and siRNA-

mediated knockdown of circEPSTI1 reduced the AKT1 level, suggesting the novel regulatory

axis of circEPSTI1-hsa-miR-942-5p-AKT1 that modulates the host-virus interactions.

CONCLUSION

circEPSTI1, found to be upregulated in the plasma of Dengue patients, the correlation of

circEPSTI1 level with the different severity is not yet known. Although circEPSTI1 is

conserved, the divergent primers designed at the back-splice junction of the circEPSTI1 could

not amplify the desired sequence in the mouse macrophages. Future studies are needed to

explore the correlation between circEPSTI1 levels and the dengue severity and the mechanistic

role in in-vivo models. Since DENV naturally doesn't infect mice, adapted mouse models

susceptible to DENV infection can be utilized to explore the mechanistic role of circEPSTI1

upon DENV infection in vivo. Despite these shortcomings, to our knowledge, this is the first

study that explored the role of any circRNA upon DENV infection and also shows a high

potential for the clinical diagnosis of Dengue. Further research focusing on the clinical validation

of circEPSTI1-based diagnostics and the development of therapeutic strategies targeting

circEPSTI1 could provide new avenues for the management and treatment of dengue.

Materials and Methods

Reanalysis of publicly available data from the NCBI-GEO database

Publicly available RNA-sequencing datasets related to Dengue virus infection clinical samples in which Total RNA-sequencing was performed were retrieved from the NCBI-GEO database. Only one dataset (GSE94892) was found eligible for the study(11, 34). The raw sequencing reads were downloaded and quality-checked. Adapter trimming and low-quality read filtering were performed using fastp (35). The clean reads were aligned to the human reference genome (GRCh38) using the BWA aligner (36). The unmapped reads were further processed for circRNA detection using the CIRI2 algorithm (37). The quantification of circRNAs was performed through CIRIquant (38). The circRNA expression counts were pre-filtered with rowsums more than 5 in at least 10 samples to avoid the least-expressing circRNAs with zero counts. Differential expression analysis between DENV-infected and control samples was performed using DESeq2 (39). The top differently expressed circRNAs (DEcircRNAs) were identified based on the adjusted p-value and the logFC. The top DECircRNAs were visualized through a heatmap, a violin plot, and a dot plot using the ggplot2 R package.

Cells and transfection

A549 human alveolar basal epithelial cells (ATCC CCL-185), HEK293T human embryonic kidney cells (ATCC CRL-3216), HepG2 human hepatoblastoma cell line (HB-8065) and MRC5 Medical Research Council cell strain 5 (ATCC CCL-171) were grown in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% fetal bovine serum (FBS) and 1% penicillin-streptomycin. DNA, siRNA, or poly I:C transfection was carried out using Lipofectamine 3000 (Invitrogen). For electroporation of human PBMCs 1×10^6 cells were resuspended in Opti-

470 MEM (Invitrogen) containing 50 nM miRNA mimics (ThermoFisher Scientific). The cells were

471 subjected to two pulses of 1000 V for 0.5 ms with a 5-second interval between pulses using the

472 Gene Pulser Xcell electroporation system. Following electroporation, the cells were transferred

473 to RPMI supplemented with 10% FBS.

Generation of the DENV-2 virus

Dengue virus (strain DENV-2, New Guinea C) virus was used in this study. Dengue virus was

rescued from the reverse genetic system gifted by Prof. Andrew D Davidson, University of

Bristol (40). Briefly, the plasmid was cloned into a low-copy-number plasmid pWSK29.

Transformation of the vector pWSK29 was done into E. coli (DH5α) followed by overnight

incubation and plasmid DNA isolation. Xba I linearized pDVWS601 plasmid was transcribed in

vitro using a T7 RNA transcription kit followed by determination of the concentration of the

RNA. For virus recovery, BHK-21 cells were transfected using DENV-2 transcripts, and the

virus was harvested 48 hours after transfection. The cells were fixed, over-dried, and then

assayed for evidence of viral replication by immunofluorescence assay using an antibody

recognizing the DENV E protein. The remaining supernatant was used for the setup of passage 1

on C6/36 cells and for RT-PCR analysis.

Viruses and infection

487 Cells were washed with PBS and infected with 0.5 MOI DENV-2 virus dissolved in SF-DMEM.

488 After 1.5 hrs, cells were washed with PBS again and incubated with SF-DMEM supplemented

489 with 1% FBS.

PBMC isolation

Human PBMCs were collected from healthy donors using EDTA blood tubes and isolated using Histopaque-1077 (Sigma) as per manufacturer's guidelines. The cells were cultured in RPMI 1640 media supplemented with 5% FBS. The cells were infected with MOI=1.

Quantitative real-time reverse transcription-PCR

Intracellular RNA was extracted using TRIzol reagent (Invitrogen) while RNA from supernatant and patient samples were isolated by TRIzol LS reagent (Invitrogen) and then converted into cDNA with the High-Capacity cDNA synthesis kit (ThermoFisher Scientific) as per the manufacturer's guidelines. Gene expression was evaluated through quantitative real-time PCR using gene-specific primers and PowerUp SYBR Green Master Mix (ThermoFisher Scientific).

RNA Fluorescence in situ hybridization

A549 cells were seeded in 24-well plate and infected with DENV virus at an MOI=0.5 for 36h. The cells were fixed with 4% formaldehyde for 20min and permeabilized with 0.1% Triton X-100 for 5 min. The cells were treated with prehybridization buffer (2X saline sodium citrate buffer, 10% formamide) for 20min at room temperature. The cells were then hybridized in hybridization buffer (10% Dextran sulfate, 4X SSC, 40% formamide, 10×10^{-3} M DDT, 100 µg/ml yeast tRNA, 100 µg/ml sheared salmon sperm DNA, 1X Denhardt's solution) with 2 µl of Cy3-labeled circEPSTI1 probe at 55°C overnight. The cells were washed with PBS and the nuclei was stained with DAPI. The images were taken with LSM 780 confocal laser microscope.

Digital PCR

RNA was extracted with Trizol and cDNA synthesis was done using High-Capacity cDNA synthesis kit (ThermoFisher Scientific) as per the manufacturer's guidelines. dPCR reactions were performed using A549 cell line as template and circEPSTI1 primer using the kit QIAcuity EG PCR Kit (250111, QIAGEN).

Luciferase reporter assay

HEK 293T cells were seeded in 24-well plates and transfected at 60–70% confluency with 30 ng of the control plasmid pRL-TK, 100 ng of the luciferase reporter plasmid, and 25 nM of the control or miR-942 mimic. Cells were collected and lysed 36 hours post transfection, and luciferase activity was measured using Dual-Luciferase® Reporter Assay System (Promega, Cat# E1960) according to the manufacturer's instructions and the reading were recorded in the GloMax® Multi+ system (Promega).

RNA immunoprecipitation

HEK293T cells were lysed using ice-cold lysis buffer supplemented with 1X protease inhibitor cocktail (11836145001, Roche). The lysate was incubated with FLAG M2 affinity beads overnight at 4°C. The lysate was washed with washing buffer (300 mM NaCl, 50 mM tris-glycine (pH 7.5), 5 mM MgCl2, and 0.05% NP-40). The RNA was extracted using TRIZOL reagent.

Biotin Pull Down Assay

A549 cells were transfected with empty vector and circEPSTI1 plasmid. The cells were lysed

post 24 hours after transfection and was incubated with control and circEPSTI1 biotin probe

overnight at 4°C. The lysate was incubated with Pierce™ Streptavidin Magnetic Beads (88817,

ThermoFisher Scientific) at 4°C for 2 hours. The beads were then washed thrice with lysis buffer

and the RNA was extracted using TRIZOL.

Enzyme-linked immunosorbent assay

Culture supernatants from A549 cell lines and PBMC were collected and analyzed by Enzyme-

linked immunosorbent assay following the manufacturer's protocols to determine the levels of

IP-10 (550926, BD OptEIA™ Human IP-10 ELISA Set) and IL-6 (555220, BD OptEIA™

Human IL-6 ELISA Set).

Immunoblotting analysis

Cells were harvested 36 hours post infection, washed with PBS and lysed with ice-cold lysis

buffer containing 1x protease inhibitor cocktail (11836145001, Roche) and 1mM of sodium

orthovanadate (S6508, Sigma-Aldrich). The lysate was centrifuged at 15000g for 10 minutes at

4°C and the supernatant was collected. Protein was quantified by Bradford assay and around 10

µg of total protein was loaded. Immunoblotting was done using anti-AKT1 (9Q7, Invitrogen),

anti-phospho-Akt (9271, Ser473, Cell Signaling Technology) anti-NS5 (MA5-17295,

Invitrogen), anti-NS1 (SAB2700022, Sigma-Aldrich), anti-β actin (A1978, Sigma-Aldrich)

antibody. Secondary anti-mouse IgG and anti-rabbit IgG antibodies were purchased from LI-

COR. The immunoblotted PVDF membrane was visualized with LI-COR system.

Confocal microscopy

A549 cells were grown under cover slips in 12 well plate. The cells were fixed with acetone: methanol in 1:1 ratio for 10 mins at -20°C. The fixed cells were blocked with BSA supplemented with 0.1% Triton X-100 for permeabilization for 2 hours at room temperature and incubated with primary antibody for 1 hour. The cells were washed three times with PBST and incubated with secondary antibody for 30 minutes at room temperature. The cells were washed three times with PBST, the nucleus was stained with DAPI, and the coverslip was mounted on a glass slide. The cells were then analyzed with LSM 780 confocal laser microscope. ImageJ software was used to analyse the images.

Flow cytometry

Cells were harvested post 36 hours of infection, trypsinized and fixed with 4% formaldehyde for 5 minutes. The cells were permeabilized with 0.1% Triton X-100 for 10 minutes, blocked by 5% FBS for 40 minutes, incubated with primary antibodies for 40 minutes and secondary antibody for 30 minutes. The cells were analyzed using flow cytometer FACSAria III (Becton Dickinson) and the data was analyzed using FlowJo software.

Statistical analysis

All experiments were conducted with appropriate control or mock-transfected samples. Each experiment was repeated independently two or three times. Data were analyzed for statistical significance using GraphPad Prism software. Differences between two groups were assessed using an unpaired, two-tailed Student's t-test, while differences between three or more groups were evaluated using ANOVA followed by the Newman-Keuls test. Differences were regarded

573   as statistically significant when P < 0.05. Statistical significance in the figures is represented as

574   follows: ***P < 0.001, **P < 0.01, *P < 0.05; ns, not significant.

REFERENCES:

1. J. P. Messina *et al.*, The current and future global distribution and population at risk of dengue. *Nat Microbiol* **4**, 1508-1515 (2019).
2. A. Srikiatkhachorn *et al.*, Dengue--how best to classify it. *Clin Infect Dis* **53**, 563-567 (2011).
3. P. Krishnamoorthy, A. S. Raj, P. Kumar, N. Das, H. Kumar, Host and viral non-coding RNAs in dengue pathogenesis. *Rev Med Virol* **32**, e2360 (2022).
4. H. Kumar, T. Kawai, S. Akira, Toll-like receptors and innate immunity. *Biochem Biophys Res Commun* **388**, 621-625 (2009).
5. J. Sprokholt, L. C. Helgers, T. B. Geijtenbeek, Innate immune receptors drive dengue virus immune activation and disease. *Future Virol* **13**, 287-305 (2017).
6. Y. T. Kao, M. M. C. Lai, C. Y. Yu, How Dengue Virus Circumvents Innate Immunity. *Front Immunol* **9**, 2860 (2018).
7. Y. Wang *et al.*, Circular RNAs in human diseases. *MedComm (2020)* **5**, e699 (2024).
8. A. Gu *et al.*, Functions of Circular RNA in Human Diseases and Illnesses. *Noncoding RNA* **9** (2023).
9. H. Ingle *et al.*, The microRNA miR-485 targets host and influenza virus transcripts to regulate antiviral immunity and restrict viral replication. *Sci Signal* **8**, ra126 (2015).
10. T. Tagawa *et al.*, Discovery of Kaposi's sarcoma herpesvirus-encoded circular RNAs and a human antiviral circular RNA. *Proc Natl Acad Sci U S A* **115**, 12805-12810 (2018).
11. A. D. Pandey *et al.*, Correlation of altered expression of a long non-coding RNA, NEAT1, in peripheral blood mononuclear cells with dengue disease progression. *J Infect* **75**, 541-554 (2017).
12. A. Kumar *et al.*, MicroRNA hsa-miR-324-5p Suppresses H5N1 Virus Replication by Targeting the Viral PB1 and Host CUEDC2. *J Virol* **92** (2018).
13. A. Quillet *et al.*, Improving Bioinformatics Prediction of microRNA Targets by Ranks Aggregation. *Front Genet* **10**, 1330 (2019).
14. N. Diehl, H. Schaal, Make yourself at home: viral hijacking of the PI3K/Akt signaling pathway. *Viruses* **5**, 3192-3212 (2013).
15. J. He *et al.*, hsa_circ_0006459 and hsa_circ_0015962 affect prognosis of Dengue fever. *Sci Rep* **9**, 19425 (2019).
16. P. Wu *et al.*, circEPSTI1 promotes tumor progression and cisplatin resistance via upregulating MSH2 in cervical cancer. *Aging (Albany NY)* **14**, 5406-5416 (2022).
17. P. Wu *et al.*, Circular RNA circEPSTI1 accelerates cervical cancer progression via miR-375/409-3P/515-5p-SLC7A11 axis. *Aging (Albany NY)* **13**, 4663-4673 (2021).
18. X. Tan *et al.*, circEPSTI1 Acts as a ceRNA to Regulate the Progression of Osteosarcoma. *Curr Cancer Drug Targets* **20**, 288-294 (2020).
19. B. Chen *et al.*, circEPSTI1 as a Prognostic Marker and Mediator of Triple-Negative Breast Cancer Progression. *Theranostics* **8**, 4003-4015 (2018).
20. Y. Xie, L. Wang, D. Yang, CircEPSTI1 Promotes the Progression of Non-Small Cell Lung Cancer Through miR-145/HMGB3 Axis. *Cancer Manag Res* **12**, 6827-6836 (2020).
21. T. Yang *et al.*, Downregulation of circEPSTI1 represses the proliferation and invasion of non-small cell lung cancer by inhibiting TRIM24 via miR-1248 upregulation. *Biochem Biophys Res Commun* **530**, 348-354 (2020).
22. J. Xie *et al.*, circEPSTI1 regulates ovarian cancer progression via decoying miR-942. *J Cell Mol Med* **23**, 3597-3602 (2019).
23. J. Wang *et al.*, The circEPSTI1/mir-942-5p/LTBP2 axis regulates the progression of OSCC in the background of OSF via EMT and the PI3K/Akt/mTOR pathway. *Cell Death Dis* **11**, 682 (2020).

24. Z. Cui *et al.*, The Combination of circEPSTI1 and MIF Offers Diagnostic Value for
Endometrial Cancer. *Int J Gen Med* **17**, 1395-1403 (2024).
25. X. Wang *et al.*, CircEPSTI1 in peripheral blood as a novel potential biomarker for
childhood-onset systemic lupus erythematosus. *Rheumatology & Autoimmunity* **4**, 20-26
(2024).
26. T. Kawai, S. Akira, Toll-like receptor and RIG-I-like receptor signaling. *Ann N Y Acad Sci*
1143, 1-20 (2008).
27. Z. Liang *et al.*, Activation of Toll-like receptor 3 impairs the dengue virus serotype 2
replication through induction of IFN-beta in cultured hepatoma cells. *PLoS One* **6**,
e23346 (2011).
28. J. P. Wang *et al.*, Flavivirus activation of plasmacytoid dendritic cells delineates key
elements of TLR7 signaling beyond endosomal recognition. *J Immunol* **177**, 7114-7121
(2006).
29. J. K. Sprokholt *et al.*, RIG-I-like Receptor Triggering by Dengue Virus Drives Dendritic
Cell Immune Activation and T(H)1 Differentiation. *J Immunol* **198**, 4764-4771 (2017).
30. D. Yang *et al.*, MiR-942 mediates hepatitis C virus-induced apoptosis via regulation of
ISG12a. *PLoS One* **9**, e94501 (2014).
31. P. Karyala *et al.*, DenHunt - A Comprehensive Database of the Intricate Network of
Dengue-Human Interactions. *PLoS Negl Trop Dis* **10**, e0004965 (2016).
32. V. S. Lekshmi *et al.*, PI3K/Akt/Nrf2 mediated cellular signaling and virus-host
interactions: latest updates on the potential therapeutic management of SARS-CoV-2
infection. *Front Mol Biosci* **10**, 1158133 (2023).
33. A. Lahon, R. P. Arya, A. C. Banerjea, Dengue Virus Dysregulates Master Transcription
Factors and PI3K/AKT/mTOR Signaling Pathway in Megakaryocytes. *Front Cell Infect*
Microbiol **11**, 715208 (2021).
34. A. Banerjee *et al.*, RNA-Seq analysis of peripheral blood mononuclear cells reveals
unique transcriptional signatures associated with disease progression in dengue
patients. *Transl Res* **186**, 62-78 e69 (2017).
35. S. Chen, Y. Zhou, Y. Chen, J. Gu, fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics **34**, i884-i890 (2018).
36. H. Li, R. Durbin, Fast and accurate short read alignment with Burrows-Wheeler
transform. *Bioinformatics* **25**, 1754-1760 (2009).
37. Y. Gao, J. Zhang, F. Zhao, Circular RNA identification based on multiple seed matching.
Brief Bioinform **19**, 803-810 (2018).
38. J. Zhang, S. Chen, J. Yang, F. Zhao, Accurate quantification of circular RNAs identifies
extensive circular isoform switching events. *Nat Commun* **11**, 90 (2020).
39. M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).
40. A. D. Davidson, Development and application of dengue virus reverse genetic systems.
Methods Mol Biol **1138**, 113-130 (2014).
664
665

Acknowledgement:

This work was supported by the intramural research grant from the Indian Institute of Science Education and Research, Bhopal (IISER Bhopal). The funders had no role in study design, data collection, interpretation, or the decision to submit the work for publication. The pIRESneo-Flag/HA Ago2 plasmid was a gift from T. Tuschl (Addgene; plasmid 10822). The DENV-2 reverse genetics system was a kind gift from Prof. Davidson. We thank IISER Bhopal for providing access to the Central Instrumentation Facility.

Contributions:

Nilanjana Das – Conceptualization, Methodology, Validation, Formal Analysis, Investigation, Data curation, Writing – Original Draft, Writing- Review & Editing, Visualization.

Pandikannan Krishnamoorthy - Conceptualization, Methodology, Software, Formal Analysis, Investigation, Data curation, Writing – Original Draft, Writing- Review & Editing, Visualization.

Kushal Ramrao Junghare- Investigation, Validation, Visualization, Writing- Review & Editing.

Athira S Raj- Investigation, Visualization, Data curation, Writing- Review & Editing.

Priyanka De- Resources, Writing- Review & Editing

Shouvanik Adhya - Resources, Writing- Review & Editing

Atul Garg – Resources, Writing- Review & Editing

Dheeraj Kheten – Resources, Writing- Review & Editing

Ashok Kumar- Resources, Writing- Review & Editing

Himanshu Kumar- Conceptualization, Methodology, Formal Analysis, Data curation, Writing –

Original Draft, Writing- Review & Editing, Project administration, Funding Acquisition

Visualization, Supervision

Ethics declarations

Experiments were performed after approval from the Institutional Ethical Committee (IEC),

Indian Institute of Science Education and Research (IISER) Bhopal

(IISERB/IEC/Certificate/20l7-II/01). The Dengue patient samples were obtained from the

College of Medicine and JNM Hospital, Kalyani, West Bengal. The IEC approval number for the

patient samples is F-24/PR/COMJNMH/IEC/24/2/42.

Declaration of interests

The authors declare no competing interests.

 Figure Legends

Figure 1:

Computational analysis of DENV-clinical samples identifies the upregulation of

circEPSTI1

(A) Distribution of sequencing reads mapped to the human genome: mapped reads (blue),

unmapped reads (green), both in millions, and circular reads (orange), in thousands across

samples from dataset GSE94892, supporting high-depth circRNA profiling in DENV-infected

human PBMCs. (B) Volcano plot showing differentially expressed circRNAs between control

and DENV-infected groups. Red points indicate statistically significant circRNAs based on log☐

fold change and adjusted p-values. The top three circRNAs are annotated with arrows. (C) Dot

plot depicting normalized expression levels of the top five circRNAs. The red dotted line is the

threshold with the normalized value of 5, above which the circRNAs are expressed at a higher

level. (D) Schematic of circEPSTI1 biogenesis from EPSTI1 pre-mRNA and the primer design.

circEPSTI1 and EPSTI1 levels were measured from the cDNA synthesized with the random

hexamer method, the oligo-dT primers method, and the genomic DNA and cDNA as templates

from the PBMCs for the amplification. (E) Sanger sequencing of the amplified bands using

specific divergent primers spanning the back-splice junction. (F) Pre-treatment of Actinomycin

D (10µg/ml) was performed for the indicated times, and the expression of linear EPSTI1 and

circEPSTI1 was measured. (G) DENV (48hr) infected cellular RNA from PBMCs was treated

with and without RNase R (2U/µl), and linear EPSTI1 and circEPSTI1 were quantified by

qPCR. (H) DENV (48hr) infected cellular RNA from PBMCs isolated from cytoplasmic and

nuclear cell fractionation was used to quantify circEPSTI1 by qPCR (I) Confocal microscopy

images of A549 cells using cy3-labeled probes targeting circEPSTI1 for localization of

circEPSTI1. Data are mean ± SEMs from triplicate samples of a single experiment and are

representative of results from three independent experiments.

Figure 2:

DENV infection induced circEPSTI1 depends on the RLR pathway, in vitro and ex vivo.

(A) Schematic of clinical recruitment of DENV patients' plasma from India, and circEPSTI1

levels were measured using qPCR in healthy controls and DENV patients. (B) Human PBMCs

(PBMC) of healthy volunteers were infected with DENV (MOI=1) for 36h ex vivo, and

circEPSTI1 levels were measured in cells using qPCR. (C) The copy number of circEPSTI1 was

measured using digital-PCR in the DENV-infected (MOI=0.5) and mock-infected A549 cells at

36h. (D) Different cell lines, such as A549 (Lung), MRC5 (Lung) (MOI=0.5), and HepG2

(Liver) (MOI=0.5), and primary cells, PBMCs, were infected with DENV and the circEPSTI1

level was measured at 12, 24, 36, and 48h post-infection. A549 cells were treated with Poly I:C.

(E), recombinant IFN-α (F), and recombinant IFN-β (G), and the level of circEPSTI1 was

measured at 6, 12, and 18h. (H) DENV infection (MOI=0.5) was performed in wild-type (WT),

RIG-I (Sensor) Knock-out (KO), IPS-1 (Adaptor) KO, or IRF-3 (transcription factor) KO A549

cells at 12, 24, 36, and 48 h, and the expression of circEPSTI1 was measured. (I) Mock and

DENV-2 infected A549 cells were fixed and hybridized with cy3-labelled circEPSTI1 probe and

subjected to confocal microscopy. Data are mean ± SEMs from triplicate samples of a single

experiment and are representative of results from three independent experiments. ∗∗∗∗, $p <$

0.0001, ∗∗∗$p < 0.001$ ∗∗, $p < 0.01$, and ∗$p < 0.05$, by two-tailed unpaired t test (B). ns,

nonsignificant.

 Figure 3:

DENV establishes infection through circEPSTI1

(A) siRNA-mediated knockdown of circEPSTI1 was performed by designing the siRNA against

the back-splice junction as shown in the schematic. PBMC and A549 cells were transfected with

circEPSTI1 siRNA for 24h, and subsequently, the cells were infected with DENV (MOI=0.5) for

36h. Knockdown efficiency was tested by qPCR. (B) Intracellular viral load was measured by

qPCR. (C) Viral load in the culture supernatant was measured upon circEPSTI1 knockdown by

qPCR. A549 cells transfected with circEPSTI1 siRNA were infected with DENV (MOI=0.5) for

36h. DENV viral load at protein level (NS5) upon circEPSTI1 knockdown was quantified using

the (D) western blot, (E) FACS analysis, and (F) confocal microscopy using DENV NS5-specific

antibody. Nuclei were stained with DAPI. (G) circEPSTI1 was cloned in Circminivector mneon

v2 vector, as shown in the schematic. A549 cells were transfected with the overexpression vector

and infected with DENV (MOI=0.5) for 36h, and overexpression was tested by q-PCR. A549

cells transfected with the circEPSTI1 vector were infected with DENV (MOI=0.5) for 36h. Viral

load in cells (H) and culture supernatant (I) and was tested by qPCR. DENV NS5 protein level in

circEPSTI1 was quantified through western blot analysis. (J) FACS analysis and confocal

microscopy (L) using DENV NS5 antibody. Nuclei were stained with DAPI. IFNβ and OAS2

levels were tested by qPCR upon (M) circEPSTI1 knockdown and (N) circEPSTI1

overexpressed A549 cells. The levels of interferon induced protein IP-10 and inflammatory

cytokine IL-6 were quantified using ELISA upon (O) circEPSTI1 knockdown and (P)

circEPSTI1 overexpression. Data are mean ± SEMs from triplicate samples of a single

experiment and are representative of results from three independent experiments. ∗∗∗∗, p <

0.0001, ∗∗∗, p < 0.001, ∗∗, p < 0.01, and ∗, p < 0.05, by two-tailed unpaired t test (B). ns,

nonsignificant.

Figure 4:

CircEPSTI1 sponges hsa-miR-942-5p

(A) Three RNA interaction prediction algorithms, miRanda, PITA, and TargetScan, were

employed to predict the interaction of miRNAs with circEPSTI1 with a stringent cutoff:

miRanda - Binding energy cutoff -40 kcal/mol and minimum miRanda Score of 300; PITA -

PITA score of minimum -10 and number of binding sites to be minimum 6; TargetScan - more

than one binding site. (B & C) circEPSTI1 wild type (WT) and mutant (M) sequences were

cloned downstream of luciferase HEK293T cells were transfected with indicated, 100ng of

circEPSTI1-luc (WT) or mut circEPSTI1 (M), 25nM miR-942-5p mimic or 25nM miR-942-5p

inhibitor, and 30ng pRL-TK vector. Luciferase assay was performed after 48h. (D) HEK293T

cells were transfected with 2µg of plasmid encoding Flag-Ago2 and 25nM mir-942 and were

infected with DENV virus (MOI=2) for 36h. RNA immunoprecipitation was done using FLAG-

M2 beads, and the enrichment of miR-942 was tested by qPCR. (E) A biotin-labeled circEPSTI1

probe was used to pull down the RNA fractions binding to the circEPSTI1, and the level of miR-

942 was measured by qPCR. (F) Upon knockdown of circEPSTI1, miR-942-5p expression was

quantified in PBMC and A549 cells by qPCR. (G) A549 cells were transfected with sicirEPSTI1

and sicircEPSTI1 along with miR-942 inhibitor. miR-942 level and viral load were quantified by

qPCR. (H) miR-942-5p levels upon DENV infection at different time points were quantified in

PBMC (MOI =1), A549 (MOI=0.5), MRC5(MOI=0.5), and HEPG2(MOI=0.5) by qPCR. Data

are mean ± SEMs from triplicate samples of a single experiment and are representative of results

from three independent experiments. ****, p < 0.0001, ***p < 0.001 **, p < 0.01, and *p < 0.05,

by two-tailed unpaired t test (B). ns, nonsignificant.

Figure 5:

miR-942-5p targets DENV genomic RNA and suppresses viral replication

(A) Predicted binding sites of miR-942-5p across the genomes of four different DENV

serotypes. (B) NS1, NS5, NS3, and NS2B sequences were cloned downstream of the pmiR-

luciferase vector. HEK293T cells were transfected with 100ng of NS1, NS3, NS5, and NS2B

luciferase constructs, 25nM of miR-942 mimic, and 30ng of PRL-Tk plasmid. Luciferase assay

was performed after 48h. (C) A549 cells were transfected with 2µg of FLAG-Ago2 plasmid and

50 nM of miR-942 mimic, followed by DENV infection for 36h. RNA immunoprecipitation was

done using FLAG-M2 beads, and the enrichment of virus was tested by qPCR. (D) A549 cells

and PBMCs were transfected with 25nM of miR-942 mimic and 25nM of miR-942 inhibitor for

24h and infected with DENV for 36h. The viral load was quantified by qPCR. (E) Viral load in

the supernatant of A549 cells was quantified by qPCR. (F) DENV NS1 and NS5 protein level in

miR-942 overexpressed A549 cells was quantified through western blot analysis. (G) FACS

analysis was performed with the DENV NS5-specific antibody upon miR-942 overexpression in

A549 cells. (H) A549 cells transfected with miR-942 mimic were infected with DENV for 36h

and were subjected to confocal microscopy using DENV NS1 and NS5 antibody. Nuclei were

stained with DAPI. (I) The levels of interferon induced protein IP-10 and inflammatory cytokine

IL-6 were quantified using ELISA upon miR-942 overexpression in A549 cells. A549 cells and

PBMCs were infected with DENV at an MOI=0.5 and MOI=1 respectively. Data are mean ±

SEMs from triplicate samples of a single experiment and are representative of results from three

Figure 6:

circ-ESPTI1-regulated miR-942-5p affects host AKT-signaling upon DENV infection

(A) The host targets of miR-942-5p were tested using miRabeL. (B) The expression of AKT1

and p-AKT1 at the protein level was quantified using western blot analysis at 6, 12, 24, 36, and

48h in DENV-infected A549 cells. (C) AKT1-3' UTR sequence was cloned downstream of the

pmiR-luciferase vector. HEK293T cells were transfected with 100ng of AKT1-3'UTR luciferase

construct, 25nM of miR-942 mimic, 25nM of miR-942 inhibitor and 30ng of PRL-Tk plasmid.

Luciferase assay was performed after 48h. (D) HEK293T cells were transfected with 100ng of

WT AKT1-luc, 100ng of Mut AKT1-luc, 25nM miR-942-5p mimic, and 30ng pRL-TK vector.

Luciferase assay was performed after 48h. (E) A549 cells were transfected with 2µg of FLAG-

Ago2 plasmid and 50 nM of miR-942 mimic, followed by DENV infection for 36h. RNA

immunoprecipitation was done using FLAG-M2 beads and the enrichment of AKT1 was tested

by qPCR. (F) A549 cells were transfected with 25nM of miR-942 mimic for 24h and infected

with DENV for 36h. DENV NS5 and AKT1 protein level was quantified through western blot

analysis. (G) A549 cells were transfected with 25nM of AKT1 siRNA for 24h and infected with

DENV for 36h. Efficiency knockdown of AKT1 and viral load was quantified by qPCR. (H)

DENV NS5 protein level in siAKT1 transfected A549 cells was quantified through western blot

analysis (I) FACS analysis was performed with the DENV NS5 specific antibody upon

knockdown of AKT1 in A549 cells. (J) A549 cells transfected with siAKT1 was infected with

DENV for 36h and was subjected to confocal microscopy using DENV NS5 antibody. Nuclei

were stained with DAPI. (K) IFNβ and OAS2 levels were checked by qPCR upon AKT1

knockdown. An increased dose of siRNA targeting circEPSTI1 reduced AKT1 level, in turn, led

to the proviral activity of circEPSTI1. (L) The levels of interferon induced protein IP-10 and

inflammatory cytokine IL-6 were quantified using ELISA upon AKT1 knockdown (M) A549

cells were transfected with increased doses of sicircEPSTI1 for 24h and infected with DENV for

36h. AKT1 protein level was quantified through western blot analysis. (N) A549 cells were

infected with DENV and treated with 20uM and 40uM of Wortmannin (WM), an AKT1

pharmacological inhibitor for 36h. pAKT1 and DENV NS5 protein level was quantified through

western blot analysis. (O) A549 cells were infected with DENV and treated with 20uM WM for

36h and was subjected to confocal microscopy using DENV NS5 antibody. Nuclei were stained

with DAPI. A549 cells were infected with DENV at an MOI=0.5.

Data are mean ± SEMs from triplicate samples of a single experiment and are representative of

results from three independent experiments. ****, $p < 0.0001$, ***, $p < 0.001$, **, $p < 0.01$, and

*, $p < 0.05$, by two-tailed unpaired t test (B). ns, nonsignificant.

Fig 1.

Fig 5.

A miR-942-5p Binding sites Across Dengue Viral Genome