GNAQ Induces Melanomagenesis in Mitfa-Independent Melanocyte Progenitors in a Zebrafish Model of Uveal Melanoma

Julius I. Yevdash,[1,2] Delaney Robinson,[1] Rachel Moore,[1] Zhijie Li,[1] Katelyn R. Campbell-Hanson,[1] Danielle Gutelius,[3] Stephen P. G. Moore,[3] Dylan Friend,[1] Isaac O'Toole,[1] Brynnon E Harman,[6,7] Collin Montgomery,[1] Robert A. Cornell,[4] Alexander Birbrair,[5] Jesse D. Riordan,[2,8] Adam J. Dupuy,[2,8] Elaine M. Binkley,[2,6,7] Robert F. Mullins,[6,7] Deborah Lang,[3] Ronald J. Weigel,[1,2,8] Colin Kenny[1,2,4,8]

1. Department of Surgery, College of Medicine, University of Iowa, Iowa City, IA 52242, USA.
2. Holden Comprehensive Cancer Center, University of Iowa, Iowa City, IA 52242, USA.
3. Department of Dermatology, Boston University School of Medicine, Boston, MA, 02118, USA.
4. Department of Oral Health Sciences, University of Washington, School of Dentistry, Seattle, WA, 98195
5. Department of Dermatology, University of Wisconsin-Madison, School of Medicine and Public Health.
6. Department of Ophthalmology and Visual Sciences, University of Iowa, Iowa City, IA 52242, Iowa
7. Institute for Vision Research, University of Iowa, Iowa City, IA 52242, Iowa
8. Department of Anatomy & Cell Biology, University of Iowa, Iowa City, IA, 52242, USA.

Running title: GNAQ transforms melanocyte progenitors in zebrafish choroid

* Corresponding author: colin-kenny@uiowa.edu

The authors declare no potential conflicts of interest

Abstract

Melanocytes reside in diverse microenvironments that influence their susceptibility to oncogenic transformation; however, investigation of rare melanoma subsets has been limited by the lack of suitable pre-clinical animal models. Here, we developed a primary, immune-competent zebrafish model to study uveal melanoma (UM) using choroidal melanocyte-targeted injection and electroporation of plasmids encoding human GNAQ^{Q209L} together with CRISPR/Cas9 cassettes for somatic tumor suppressor gene deletion. Single-cell transcriptional profiling of primary melanocytes and melanoma derived from the eye and skin revealed distinct transcriptional programs, with epithelial-to-mesenchymal transition pathways enriched in ocular tumors. In addition, choroidal fibroblasts from tumor-bearing eyes exhibited marked transcriptional changes, including increased fibronectin and collagen expression, consistent with stromal remodeling. Given prior associations between *mitfa* loss and accelerated GNAQ^{Q209L} tumor onset, the model was applied to determine whether melanocyte differentiation state contributes to the emergence of GNAQ-driven tumors. The increased susceptibility resulted from expansion of Mitfa-independent melanocyte progenitor populations in germline *mitfa* mutant zebrafish, rather than somatic *mitfa* loss in differentiated melanocytes, as conditional, melanocyte-specific *mitfa* deletion in adult zebrafish did not accelerate tumor growth. Furthermore, *pax3a*-positive melanocyte progenitor cells in *mitfa*-deficient zebrafish embryos and adult eyes and skin were highly susceptible to transformation induced by GNAQ^{Q209L} but not BRAF^{V600E}. Analogous *PAX3* positive populations were also identified in mouse and human single-cell transcriptomic datasets. Collectively, these findings establish a critical role for Mitfa-independent melanocyte progenitors in UM pathogenesis.

54   **Statement of Significance**

56   Choroid-targeted GNAQ^{Q209L} expression induces anatomically correct uveal melanoma in adult

57   zebrafish, with germline *mitfa* deletion expanding mitfa-independent melanocyte progenitors

58   with enhanced susceptibility that are transcriptionally distinct from the subpopulation

59   transformed by BRAF^{V600E}.

Introduction

Uveal melanoma (UM) is a rare cancer of the melanocytes in the uveal tract. The majority of UM's (~90%) occur in the choroid, the vascular structure supplying the outer retina, with the remainder arising from the ciliary body or the iris.[1] Oncogenic mutations in *GNAQ* or its paralog *GNA11* are observed in up to 90% of UMs but are rare in cutaneous melanoma (CM).[2] Such mutations reduce GTPase activity of the heterotrimeric G protein alpha subunit, resulting in constitutive activation that drives cellular proliferation through downstream signaling pathways, including MAPK, YAP/TAZ, and PI3K.[3,4] The transcriptional mechanisms of UM onset and progression remain understudied, in part, because existing preclinical models inadequately capture the earliest events of ocular melanocyte transformation within the correct microenvironment.

A recent study by Phelps et al.[5] showed that loss of *mitfa,* which is considered the master regulator of the melanocyte lineage, accelerates tumorigenesis in transgenic *Tg(mitfa-GNAQ^{Q209L}); mitfa^{-/-}; tp53^{-/-}* zebrafish compared to *Tg(mitfa-GNAQ^{Q209L}); tp53^{-/-}* controls. These findings indicate that, in the context of tumor-suppressor loss, oncogenic GNAQ can drive tumor formation independently of differentiated melanocytes. This contrasts with *BRAF^{V600E}*-induced melanoma, where Mitfa is required for tumor formation.[6] Such observations imply that melanocyte progenitor cells, which persist in *mitfa*-mutant zebrafish, may serve as the cells of origin for GNAQ-induced transformation. While transgenic models have been incredibly informative for studying spontaneous tumor formation, germline loss of *mitfa* or tumor suppressor genes, together with melanocyte-lineage expression of oncogenes throughout development, may unintentionally perturb melanocyte lineage specification. Consequently, these models may capture tumorigenesis in a lineage already altered during development rather than melanoma initiated by somatic mutations in normal melanocytes. Conditional knockout

approaches as well as lineage restricted expression of oncogenes in the appropriate

microenvironment will improve our understanding of UM pathogenesis.

Zebrafish are a well-established model for studying melanocytes and melanoma due to

conserved melanocyte differentiation pathways, efficient genetic manipulation, and the

availability of transparent lines that facilitate high-resolution live-cell imaging. Zebrafish have

three types of pigment cells; dark melanocytes, iridescent iridophores, and yellow xanthophores,

that arise from multipotent, neural crest-derived pigment progenitors.[7] To study early events in

UM, we developed a novel zebrafish model that utilizes transgene electroporation of adult

zebrafish (TEAZ)[8] to express the human $GNAQ^{Q209L}$ oncogene and conditionally delete tumor

suppressor genes in cells of the melanocyte lineage within the skin and eye. Choroidal targeted

injection and electroporation drive primary UM onset from the same anatomical site as human

UM. In this study, we demonstrate the existence of a melanocyte progenitor population in

zebrafish eyes with increased sensitivity to transformation by $GNAQ^{Q209L}$. Such progenitor cells

are expanded in *mitfa* deficient zebrafish and result in reduced $GNAQ^{Q209L}$-induced tumor

latency compared to wild-type animals.

Materials and Methods

Plasmid constructs and zebrafish generated in this study may be obtained by a request to the

lead contact.

Data and code availability

Datasets generated in this article are available from NCBI GEO accession number GEO:

GSE307227. This study does not report original code. Any additional information required to

reanalyze the data reported in this work paper is available from the lead contact upon request.

Human donor eye procurement and histology

115 Human donor eyes were obtained from the Iowa Lions Eye Bank with full informed consent from

116 the next of kin for use in research. All procedures adhered to the tenets of the Declaration of

117 Helsinki and institutional guidelines. Globes were processed for histological analysis in the

118 laboratory of Dr. Mullins at the University of Iowa. Tissues were fixed, paraffin embedded,

119 sectioned, and stained with hematoxylin and eosin (H&E) using standard protocols.

Zebrafish husbandry

All zebrafish used in this study were bred and maintained at the University of Iowa Animal Care

Facility. The zebrafish are kept consistently at 28°C, 7.4 pH, and controlled salt concentrations

and cyclically receive 14 h of light followed by 10 hours of darkness. All zebrafish experiments

were performed in compliance with the ethical regulations of the Institutional Animal Care and

Use Committee at the University of Iowa and in compliance with NIH guidelines (protocol

#3022523). Zebrafish embryos were maintained at 28.5°C and staged by hours or days post-

fertilization (hpf or dpf).

Zebrafish mutant lines

Wild-type zebrafish in this study include the *AB* line (RRID:ZFIN_ZDBGENO-960809-7) and the

WIK line (RRID:ZIRC_ZL84). Transgenic lines include *casper* ($mitfa^{w2/w2}$; $mpv17^{-/-}$)[9,10]

(RRID:ZIRC_ZL1714), *nacre* ($mitfa^{w2/w2}$)[11] (RRID:ZIRC_ZL2104), *Tg*(*mitfa*:GFP)[18], and

Tg(*mitfa*:GFP); $mitfa^{w2/w2}$. TEAZ-Eye and TEAZ-Skin was performed on adult zebrafish

(inclusion criteria: greater than 4 months postfertilization and less than 1.5 years old). Similar

proportions of male and female zebrafish were used within each experiment.

Induction of melanoma using Strategies A and B.

To model GNAQ-driven melanoma, we generated *mitfa* promoter–driven plasmid systems

expressing human oncogenic GNAQ^{Q209L}, CRISPR/Cas9, and guide RNAs (gRNAs) targeting

tp53, ptena, and ptenb. Two delivery strategies were used. In Strategy A, a single plasmid

contained two expression cassettes under the mitfa promoter: one encoding GNAQ^{Q209L} fused to

a T2A self-cleaving peptide followed by eGFP, and a second encoding Cas9, together with three

U6-driven gRNAs targeting *tp53* and *ptena/b*. Because T2A-mediated cleavage leaves residual

C-terminal amino acids that may alter GNAQ oncogenic activity, we also employed an

alternative multi-plasmid approach (Strategy B). This system consisted of five separate

plasmids: mitfa:*GNAQ^{Q209L}*, mitfa:GFP, mitfa:Cas9, U6:gRNA-*tp53*, U6:gRNA-*ptena*, and

U6:gRNA-*ptenb*. Plasmids expressing mitfa:GNAQ^{Q209L} and mitfa:GFP were generously

provided by Dr. Jacqueline Lees (MIT), and plasmids encoding Cas9 and gRNAs were provided

by Dr. Richard White (University of Oxford). Strategy B was designed to avoid unintended C-

terminal modification of GNAQ and to improve electroporation efficiency due to reduced plasmid

size. Fish were monitored weekly for tumor formation at the electroporation site. Tumor onset

was defined as the first appearance of a visible, expanding pigmented or translucent mass.

Tumor latency was calculated as days post-electroporation to first detection. Kaplan–Meier

analyses were used to compare tumor onset between experimental groups.

Cloning

The MiniCoopR[6] vector (MiniCoopR 2x*U6*:gRNA, *mitfa*:Cas9 was a gift from Leonard Zon

Addgene plasmid # 118844) was modified to replace the *mitfa* coding sequence with *GNAQ*^{Q209L}

by removing *mitfa* using StuI and SmaI restriction enzymes (NEB). Gibson assembly was used

to insert *GNAQ*^{Q209L}:T2A:GFP DNA as a gBlock hifi (IDT) with sequence on the 3'

(GAAGCTAACACATAGTTGAAC) and 5' (CAATGCCAACTAAATTTCATG) as overhangs.

Gibson assemblies were transformed using NEB 5-alpha competent *E. coli* cells via heat shock.

GNAQ^{Q209L}:T2A:GFP insertion was confirmed using Sanger sequencing (Primers:

CAAGGAAGCCCGGCGGATCAACGA, CATACTTGTATGGGATCTTGAGTGTGTCCA,

GTGGAGTCAGACAATGAGAACCGAATGGAG, CAAGCTGGAGTACAACTACAA).

mitfa sgRNA was designed using IDT's CRISPR-Cas9 guide RNA design tool (sequence:

ATGGACAAAGCTGGACCATG). *U6*:sgRNA-*ptena* was linearized by PCR (forward primer:

GTTTAAGAGCTATGCTGGAAACAGCAT, reverse primer: GAACAAAGAGCTGGAGGGAG).

Linear plasmid was purified from a 2% agarose gel using the Monarch GNA Gel Extraction Kit

(#T1020S). ssDNA sequences for sgRNA-*mitfa* were annealed together in NE buffer r2.1.

Gibson assembly was used to insert sgRNA-*mitfa* into linearized U6 plasmid. Assembled

product was transformed into NEB 5-alpha competent *E. coli* cells via heat shock and selected

with 50 ug/mL streptomycin. Insertion of sgRNA-*mitfa* was confirmed using PCR amplification of

lysed bacteria from single colonies (forward primer: ATGGACAAAGCTGGACCATG, reverse

primer: CTTAGCTGGATAACGCCAC). Positive products were isolated using the QIAGEN

Plasmid Maxi Kit. Entire plasmid was sequenced using Plasmidsaurus sequencing service.

Transgene Electroporation of Adult Zebrafish Skin (TEAZ-Skin)

Adult zebrafish were anesthetized in 0.2% Tricaine prior to upright placement on a prewetted

Kimwipe. Each injection consisted of 1000 ng total DNA: 100 ng of Tol2 plasmid and an equal

division of the remaining 900 ng for the remaining constructs. Injection stocks contained a small

volume (<1uL, 1:50 dilution) of green food coloring to visualize localization of injection. Injection

volumes ranged from 1 to 1.5 uL depending on the injection mix, keeping 1000 ng total

constant. Needles were pulled from 10 cm long filamented borosilicate glass capillaries at a

pressure of 500, heat of 650, pull strength of 100, velocity of 200, and time delay of 40 in a

Model P-97 Flaming/Brown micropipette puller from Sutter Instrument Co. Following injection,

zebrafish were immediately electroporated (<1 min) by placing the cathode probe of ultrasound

gel-coated Platnium Tweezertrode, 3MM, on the side of the injection site and the anode probe

on the other side of the fish body. Electroporation was achieved with 5 60-ms pulses of 40 volts

with a 1s pulse interval from a ECM 830 Electro Square Porator from BTX Harvard Apparatus.

Following electroporation, zebrafish are placed into a recovery tank with fresh, gently stirred

system water and are monitored until normal swimming resumes. Electroporated zebrafish are

imaged weekly. *mitfa*:$GNAQ^{Q209L}$; *mitfa*:GFP plasmid constructs were kindly provided by Dr.

Jacqueline A. Lees (Massachusetts Institute of Technology, Cambridge, USA). *mitfa*:Cas9,

mitfa:tdTomato, PCS2FA Tol2, *U6*:gRNA-*tp53, U6*:gRNA-*ptena,* and *U6*:gRNA-*ptenb* plasmids

were kindly provided by Dr. Richard White (Sloan Kettering Institute, New York, USA).

Transgene Electroporation of Adult Zebrafish Eye (TEAZ-Eye)

Pre-injection preparation for eye injections is the same as for skin injections including: 0.2%

Tricaine anesthesia, needle size, and injection concentrations. To inject the eye, a blunt

spherical head of a pin was gently pressed against the bottom half of the zebrafish eyeball to

reveal the dorsal posterior regions of the eye. The needle was gently pressed into this location,

where the cornea is softer, just through the sclera, keeping the needle as superficial as possible

as to not damage the back of the eye. Injection volumes were around 0.25 uL and were

terminated with a slight visual swelling of the eye. Following injection zebrafish were

immediately electroporated (<1 min) by the cathode probe of ultrasound gel-coated Platnium

Tweezertrode, 1MM, on the superior side of the eye nearest the injection and the anode probe

on the inferior side of the eye. Electroporation was achieved with 3 50-ms pulses of 75 volts with

a 1s pulse interval from a ECM 830 Electro Square Porator from BTX Harvard Apparatus.

Recovery and imaging procedures for these fish did not differ from the skin injections.

Imaging and image processing

Injected zebrafish were imaged using an upright Leica M205 FCA microscope with brightfield

and GFP filters. Zebrafish were briefly anesthetized with 0.2% Tricaine and positioned on a

Kimwipe mound for upright orientation. Images were taken and processed through the LAS X

software. All histology slides were imaged using brightfield, DAPI, and GFP filters on an EVOS

M5000 microscope from Invitrogen for individual images or on a Leica DMi8 microscope and

stitched together by LAS X Thunder Imaging software for full tissue images.

H&E and IF – zebrafish tissue

Whole zebrafish or zebrafish eyes were fixed in 4% paraformaldehyde for 48 hours at 4°C and

paraffin embedded in the University of Iowa Pathology Core. Fish were sectioned at 7 µM

thickness and placed on Superfrost Plus Microscope slides from Fisher Scientific, baked at

60°C, stained with hematoxylin and eosin, and mounted with Cytoseal 60 from Epredia. For

immunofluorescence, sections were deparaffinized and hydrated followed by heat-induced

antigen retrieval with 10 mM Na-citrate buffer and blocking with 3% milk in 0.1% Tween 20.

slides were stained with primary antibodies against GFP (NB600-308 Rabbit anti-GFP

polyclonal from Novus Biologicals, RRID:AB_10003058, at 1:250) overnight at 4°C and then

fluorescent secondary antibodies against rabbit IgG (Alexa Fluor 488 goat anti-rabbit IgG,

RRID:AB_2576217, at 1:500 from a 2 mg/mL stock) for 1 hr at room temperature. RPE

autofluorescence was minimized using the Vector TrueVIEW Autofluorescence Quenching Kit.

Sections were counterstained with DAPI, mounted using VECTASHIELD Vibrance Antifade

Mounting Medium, and imaged the same day.

IF - mouse eye tissue

C57BL/6 mice (RRID:MGI:2159769) eyes were collected, formalin-fixed, paraffin-embedded,

and sectioned into 7 μm-thick sections. Sections were deparaffinized and hydrated followed by

heat-induced antigen retrieval with citrate buffer (Vector Laboratories, Inc.), permeabilized with

1% horse serum in 0.2% Triton-X, blocked with 5% Horse serum in 0.2% Triton-X, and

incubated with PAX3 primary antibody (Invitrogen 38-1801 Rabbit anti-PAX3 polyclonal,

RRID:AB_2533359, at 1:200) overnight at 4°C. Samples were incubated with DyLightTM 488-

labeled secondary antibody (Vector Laboratories, DI-1088-1.5, horse anti-rabbit,

RRID:AB_2336403, at 1:2000) for 1 hr at room temperature, treated with autofluorescence

quenching kit (Vector Laboratories, Inc. SP-8400-15), and cover slips fitted using Vectashield

Antifade Mounting Medium with DAPI (Vector Laboratories, Inc. H-1000-10).

Kaplan-Meier analysis

Zebrafish were followed for up to 70 weeks and tumor-free survival was analyzed with the

Kaplan-Meier method. Tumors were defined as the first presence of a GFP-positive nodular

mass at the injection site. Zebrafish were excluded from curves if they perished or were

euthanized without any visible signs of tumors during the study. Differences between injection

groups were analyzed using log-rank statistics.

Single cell suspension and single-cell RNA Seq analysis

Zebrafish were euthanized with 0.4% Tricaine for 10 mins followed by immersion in ice water

until no opercular movement was observed for at least 30 min as detailed in our IACUC

protocol. Tumors were harvested and minced into small pieces. Minced tumors were digested

for 1 hour at 37°C with collagenase and hyaluronidase in HBSS with 2% FBS (HF).

Suspensions were washed with 1:4 HF:NH$_4$Cl and then further digested for 5 min with pre-

warmed trypsin. Following trypsin inactivation, cell suspensions were washed twice with 10%

FBS in HBSS.

Cellular suspensions were loaded on a 10x Genomics Chromium instrument to generate single-

cell gel beads in emulsion (GEMs). Approximately 10,000-20,000 cells were loaded per channel

depending on the 10x kit used. See figures S5 and S12 for kit information per sample.

Next GEM kit: targeted cell number = 10,000. Single-cell RNA-Seq libraries were prepared

using Single Cell 3′ Reagent Kits v2: Chromium Single Cell 3′ Library & Gel Bead Kit v2, PN-

120237; Single Cell 3′ Chip Kit v2, PN-120236; and i7 Multiplex Kit, PN-120262 (10x Genomics)

and following the Single Cell 3′ Reagent Kits v2 User Guide (Manual Part # CG00052 Rev A).

GEM-X kit: targeted cell number = 20,000. Single-cell RNA-Seq libraries were prepared using

Chromium GEM-X Single Cell 3′ Reagent Kits v4: Library Construction Kit C, PN-1000694;

Single Cell 3' GEM Kit v4, PN-1000693; Single Cell 3' Gel Bead Kit v4, PN-2001128; and Dual

Index Kit TT Set A, PN-3000431; and following the Chromium GEM-X Single Cell 3′ Reagent

Kits v4 User Guide (Manual Part # CG000731 Rev B). Libraries were sequenced on an Illumina

HiSeq 4000 as 2□×□150 paired-end reads. Sequencing results were demultiplexed and

converted to FASTQ format using Illumina bcl2fastq software.

A custom reference genome for the Zebrafish was constructed with GRCz11 primary assembly

(Ensembl, RRID:SCR_002344) using Cell Ranger (Cell Ranger mkref function). 10X Genomics

scRNA-Seq reads were then processed and aligned to this reference with Cell Ranger (Cell

Ranger Count function). Further analysis and visualization was performed using Seurat

(v4.1.0)[12], cells with fewer than 200 RNA feature counts and greater than 5% mitochondrial

contamination were removed with filtering (greater than 15% for wildtype eye samples). RNA

counts were normalized, and FindVariableFeatures was run with the following parameters:

selection.method = "vst", nfeatures = 2,000. The cells were originally clustered in a UMAP by

RunPCA then RunUMAP with dims 1:30. FindNeighbors was run with dims 1:30 followed by

FindClusters using a resolution of 0.4. These methods identified up to 20 clusters in individual

samples.

For comparative analysis, RNA-seq datasets were integrated together using IntegrateData with

dims 1:30. New clusters were generated as before. Clusters were annotated using

FindAllMarkers and the top 25 enriched genes in each cluster were compared to the Daniocell

dataset[13] and Spectacle[14]. Cluster names were given according to the closest matched cellular

population between these two datasets. To determine enriched pathways, *danio rerio* gene

names were converted into human Ensembl gene codes with g:Profiler[15] and then data were

analyzed with the use of QIAGEN IPA (QIAGEN Inc)[16] and GSEA.[17] Pseudotime was performed

using Monocle3 (v0.2.3.0).[18]

Verification of sgRNAs

Zebrafish were euthanized with 0.4% Tricaine for 10 mins followed by immersion in ice water

until no opercular movement was observed for at least 10 min as detailed in the IACUC

protocol. Tumors were harvested and minced into small pieces. DNA was harvested according

to the DNeasy Blood and Tissue Kit from Qiagen. Regions surrounding cut sites were PCR-

amplified and the purity of the PCR amplicon was confirmed with agarose gel electrophoresis.

Samples were cleaned up with the QIAquick PCR Purification Kit from Qiagen and sent for

Sanger sequencing at the Iowa Institute of Human Genetics. Gene editing was confirmed with

DECODR INDEL analysis by comparing to wildtype sequence[19]. PCR primer sequences follow:

mitfa PCR F: CCGGTATGTATTCACATTGTCTTG

mitfa PCR R: TCTTGCTTAGGATGCCTATGTATT

ptena PCR F: CATCCCACCAAGTGAGGTTAAAC

ptena PCR R: CACATACACAGTCAAGGGTGAG

ptenb PCR F: CAGTTCTGTTGCACCCAATAAG

ptenb PCR R: CTGGTGGTGTTGAGGCTATAAAG

tp53 PCR F: AAGTATTCAGCCCCCAGGTG

tp53 PCR R: CGCTTTTGACTCACAGTGCAAG

Results

Generation of GNAQ-driven melanoma in adult zebrafish

To model GNAQ-driven melanoma in adult zebrafish, we generated *mitfa* promoter-driven

plasmid systems expressing oncogenic GNAQ^{Q209L}, Cas9, and gRNAs targeting *tp53* and

ptena/b (**Figure 1A**). We tested two delivery strategies: a single-plasmid GNAQ-T2A-GFP

construct incorporating all expression cassettes (**Strategy A**), and a multi-plasmid approach

designed to avoid C-terminal modification of GNAQ by the T2A cleavage site while improving

electroporation efficiency through the use of smaller individual plasmids (**Strategy B; see**

Methods).TEAZ applied to adult zebrafish skin (TEAZ-Skin) induced melanoma in wild-type

animals using either strategy, with Strategy B significantly reducing tumor latency (**Figure 1C–**

D, Figure S1A). Melanomas developed exclusively from clonal expansion of cells harboring

edits introduced by all gRNAs. Neither oncogenic GNAQ expression nor individual tumor

suppressor loss alone was sufficient to drive tumorigenesis, demonstrating that combined

oncogenic signaling and inactivation of all gRNA-targeted tumor suppressors were required for

transformation (**Figure S1A**). Consistent with prior reports, loss of *mitfa* markedly accelerated

tumor onset following TEAZ-Skin, with tumors arising within ~30 days in *nacre* ($mitfa^{w2/w2}$) and

casper ($mitfa^{w2/w2}$, $mpv17^{a9/a9}$) zebrafish (**Figure 1C–D, Figure S1B–D**). In these backgrounds

(**see Methods**), oncogenic GNAQ alone was sufficient to drive melanoma, consistent with

increased susceptibility of Mitfa-independent cells. Tumor identity was distinguished from

xanthomas by gross morphology and pigmentation (**Figure S1E**), and CRISPR-mediated editing

of target genes was confirmed by Sanger sequencing (**Figure S2**). Together, these data

demonstrate that TEAZ-Skin efficiently induces GNAQ-driven melanoma in adult zebrafish, with

Strategy B providing faster tumor onset than Strategy A and *mitfa* loss substantially accelerating

disease compared to TEAZ-Skin in wild-type zebrafish (**Figure 1C–D**).

The *mitfa* promoter is active in choroidal melanocytes

We next sought to adapt the TEAZ approach to induce melanocyte transformation in the

zebrafish eye (TEAZ-Eye)**.** The zebrafish and human eye share highly conserved overall

organization and cellular composition (reviewed in [20]), including distinct retinal layers, retinal

pigmented epithelium, and a melanocyte-containing choroidal layer. Most major human ocular

compartments have clear zebrafish counterparts, although the zebrafish lacks a fovea and has

a proportionally thinner choroid (**Figure 1E**). We first assessed specificity of the *mitfa* promoter

in driving transgene expression in choroidal melanocytes by performing immunofluorescent

imaging with an anti-GFP antibody in eyes harvested from transgenic *Tg(mitfa:GFP)* zebrafish.

Choroidal melanocytes exhibited strong GFP positivity compared to the secondary antibody-

only control (**Figure 1F**). Quenching autofluorescence signal from RPE improved the signal-to-

noise ratio (**Figure S3A-B**). We next applied TEAZ-Eye by injecting and electroporating

plasmids containing *mitfa* promoter-driven *GFP* expression (*mitfa:GFP*) into the choroid. Eyes

were harvested two weeks post-electroporation to assess GFP expression by

immunofluorescence. Cells at the injection site exhibited strong GFP signal compared to the

secondary antibody-only control (**Figure S3C**), demonstrating that the *mitfa* promoter is

sufficient to drive transgene expression in choroidal melanocytes using TEAZ-Eye.

TEAZ-Eye induces UM in the choroid of wild-type and *casper* zebrafish

A major limitation in the study of UM onset and progression is the lack of anatomically accurate

in vivo models. To address this gap, we used the TEAZ-Eye technique to introduce oncogenic

GNAQ and tumor suppressor gRNAs into the choroidal space of wild-type zebrafish (**Figure 1G-**

H). Remarkably, TEAZ-Eye in wild-type zebrafish promoted transformation of cells within the

choroid. Live *in vivo* imaging of TEAZ-Eye enabled identification of individual GFP+ cells at the

injection site, serving as a proxy for $GNAQ^{Q209L}$ expression (**Figure 1H-H'**). We next applied

TEAZ-Eye to *mitfa*-deficient *casper* zebrafish (**Figure 1I-I'**). As with TEAZ-Skin, tumor latency of

TEAZ-Eye injected *casper* zebrafish (n=13) was significantly reduced compared to wild-type

(n=5) (**Figure 1J**). H&E analysis confirmed that TEAZ-Eye induced UM within the choroid,

progressing into surrounding ocular structures (**Figure 1K**). Notably, 100% (n=10) of tumors

formed between the sclera and RPE, highlighting the model's ability to induce anatomically

accurate UM. Finally, anti-GFP immunofluorescent analysis confirmed expression of plasmid

constructs following TEAZ-Eye (**Figure 1K'**).

UM transformation involves activation of neural crest transcriptional programs

During embryonic development, choroidal melanocytes are derived from the cranial neural crest

(CNC).[21] Since our model can detect the earliest events in UM transformation, we asked

whether reactivation of CNC programs is one of the initial steps in UM development. To address

this point, tumors were derived in wild-type zebrafish using TEAZ-Eye with Strategy B plasmids.

At 50-90 dpe, when gross nodular tumors were identified, surgical enucleation was performed,

and dissociated cells from the entire eye were subjected to single-cell RNA sequencing (scRNA-

Seq). We compared pooled wild-type tumors (2 sequencing replicates) with tumor-free paired

eyes or sibling-matched control eyes (2 and 4 pooled eyes, respectively, 2 sequencing

replicates) (**Figure 2A, Figure S4**). Given that whole eyes were used as control samples and

melanocytes are most abundant in the choroid, melanocytes identified under control conditions

are hereafter referred to as choroidal melanocytes. Tumor cells were identified by expression of

GFP and oncogenic *GNAQ* (**Figure 2B**) whereas primary choroidal melanocytes were marked

by expression of *mlana* (**Figure S5**). Annotation of additional cell types within the TME was

accomplished by comparing to previously published human choroid and retina scRNA-Seq

datasets,[14,22] the Zebrafish Information Network (ZFIN),[23] and from the zebrafish embryo single

cell atlas "Daniocell"[13] (**Figure 2A**). Interestingly, tumor cells aligned most closely with choroidal

melanocytes, ruling out the retinal pigmented epithelium (RPE) as an origin of our TEAZ-Eye

induced tumors (**Figure 2C**). Additionally, profiling *GFP/GNAQ*^{Q209L}+ UM cells revealed that

tumors exhibited strong expression of neural crest markers (*tfap2a, tfap2c, pax7b, sox10, foxd3*)

and melanocytic markers (*mitfa, tfap2e, kita, mlana, pmela, slc22a7a, mlpha, mtbl*) (**Figure 2D**).

Unexpectedly, CNC marker genes *twist1* or *dlx1*[24,25] were not enriched in UM tumors compared

to TME. Pathway analysis showed that genes enriched in TEAZ-Eye-derived tumors were

strongly associated with MITF-M-dependent gene expression, RAF/MAP kinase, mTOR,

PI3K/AKT signaling, NF-kB, RHO GTPase, and NOTCH activity, as well as fatty acid oxidation

pathways (**Figure S6A-B**).

Re-clustering of choroidal cells from control and tumor conditions revealed a marked expansion

of tumor cells relative to melanocytes, accompanied by increases in immune cells and

fibroblasts (**Figure 2E-F**). Immune cell populations within the TME included B-cells (*igic1s1,*

pax5, cd79a, zgc:153659), T-cells (*traf1, si:ch211-67e16.3, cd27, zap70, cd8a, sla2*), a

population of cytokine producing cells, likely representing T-cells (*il4, il13, il11, ca2, il34, il6*) as

well as macrophages (*marco, ccl34a.4, mrc1b, c1qa/b/c, grn1*) and neutrophils (*cpa5, lect2l,*

mmp9, npsn, scpp8). Additional cell types within the choroidal TME included fibroblasts (*fn1a,*

col1a1a, col1a1b, col1a2, col5a1, dcn, serpinf1), pericytes (*pdgfrb, notch3, foxf2b, cd248a*),

Schwann cells (*dlx5a, apoda.1, mpz, plp1b, cd59, s100b, mbap*) and vasculature endothelial

cells (*vwf, flt4, epas1b, ecscr, fabp11a*) (**Figure 2D-F**, **Table S1**).

We next compared choroidal melanocytes and UM tumors, revealing a significant overlap in

gene expression (n=644; hypergeometric p<0.0001) alongside a large set of tumor-enriched

genes (n=1177) (**Figure 2G**). Shared genes included melanocytic regulators (*ednrb, sox10,*

foxd3, mitfa, mlpha, kita, mlana, pax7b), supporting a melanocytic origin, with many transcripts

further upregulated in UM tumors compared to choroidal melanocytes (**Figure 2H**). HOMER

promoter motif enrichment analysis of genes enriched in UM tumors revealed binding motifs for

the MiT/TFE family of transcription factors, as well as PAX3/7 and HOXB13, suggesting that

these regulators may contribute to UM-specific transcriptional programs (**Figure 2G'**).

To assess the relevance of our model to human disease, we examined whether genes enriched

in zebrafish UM tumors were associated with clinical outcomes in patients. Differential

expression analysis identified a gene signature, including *LGALS2, FABP3/5, GCH, MET*, and

KIT, whose high expression was associated with reduced overall and disease-free survival and

with TCGA molecular subtypes 3 and 4,[26] corresponding to aggressive UM (**Figure 2G, 2I-K,**

Figure S7-8, Table S2).

Cancer associated fibroblasts within the uveal melanoma TME upregulate components of the extracellular matrix (ECM)

Recent studies have significantly advanced our understanding of the TME as an active driver of

melanoma progression.[27,28] While it is well established that components of the TME, particularly

cancer-associated fibroblasts (CAFs), play a key role in malignancy,[29-31] *in vivo* models to study

such interactions in UM remain limited. Our model allows for direct comparison of the

microenvironment in TEAZ-Eye and control injected eyes within the same animals (i.e., left and

right eye). We compared the transcriptome of CAFs and normal fibroblasts in wild-type

zebrafish. Upon re-clustering and differential gene expression analysis, we found that several

ECM proteins were upregulated in CAFs, indicating ECM remodeling in the presence of UM.

Enriched genes broadly fell into four categories: ECM, ECM regulators, TNFs, and collagens.

Notably, fibronectin (*fn1a*) was among the most significantly upregulated genes in CAFs

compared to normal fibroblasts (q<0.0001). Several collagens were also found to have higher

expression in CAFs, including Collagen I (*col1a1b, col1a2),* Collagen V (*col5a1, col5a2a,*

col5a3a) and Collagen XII (*col12a1a)* (**Figure 2L, Table S3**)**.** Consistently, cell–cell

communication analysis using CellChat[32] identified FN1 and collagen signaling from CAFs to

tumor cells as significantly enriched pathways and predicted the transmembrane heparan

sulfate proteoglycan SDC4 as a mediator of these interactions in UM cells (**Figure S9**).

To control for injection-related fibrosis potentially causing fibroblast activation and upregulation

of ECM components, we mock-injected three wild-type eyes with plasmids containing *mitfa*:GFP

and *mitfa*:Cas9 cassettes followed by electroporation (mock-TEAZ) and performed scRNA-Seq.

Fibroblasts from mock-injected sibling controls (three pooled eyes), non-injected sibling controls

(four pooled eyes), and non-injected tumor-matched control eyes (two pooled eyes) were

equally distributed on the UMAP, with no alteration in *fn1a* or *lgals2a* expression or other

changes to extracellular matrix (ECM) genes upon mock-TEAZ (**Figure S10**). These findings

indicate that TEAZ-Eye tumor induction depends on genetic modification of target cells, not

simply physical injection, triggering a CAF response that may be critical to melanoma

progression.

Skin and eye primary melanocytes, and their resulting tumors, have unique gene

expression profiles

Having established a system to induce genetically identical tumors in the eye and skin of adult

zebrafish, we asked whether tumor transcriptional programs differed by anatomical site. Such

differences could reflect site-specific tumor–stromal interactions or intrinsic differences between

uveal and cutaneous melanocytes. We generated TEAZ-Eye and TEAZ-Skin tumors using

Strategy B plasmids and performed scRNA-Seq with matched control eye and skin tissues

(**Figure 3A**). We compared tumors to a combined TME and performed a parallel analysis of eye

and skin primary melanocytes relative to the combined microenvironment (ME) (**Figure 3B,**

Table S4). $GNAQ^{Q209L}$ expression levels were comparable between eye and skin wild-type

tumors (**Figure S11A**) and DEGs showed substantial overlap between sites, with 41% of 2,192

genes shared, consistent with common transcriptional programs driven by oncogenic GNAQ

during transformation (**Figure 3B**). Notably, only 25.8% of DEGs overlapped between eye and

skin melanocytes, indicating substantial transcriptional differences in melanocytes originating

from distinct anatomical regions (**Figure 3B**).

We hypothesized that DEGs shared by only the two tumor sites and not the primary

melanocytes would represent essential mechanisms for UM growth and survival. Consistent

with this hypothesis, pathway analysis revealed enrichment of pathways involved in cancer

proliferation and metabolism (**Figure S12A**). Additional enriched pathways included WNT

signaling, PTEN regulation, and RAF/MAPK signaling, which have been previously implicated in

UM pathogenesis.[33-41] In contrast, IPA analysis of gene sets shared between eye and skin

melanocytes represented homeostatic pathways including enrichment of MITF-M–dependent

programs, as well as pathways related to nucleotide biosynthesis and mTORC1-regulated

amino acid metabolism (**Figure S12B**).

We next re-clustered normal melanocytes from eye and skin for direct comparison, revealing

distinct UMAP clusters (**Figure 3C**). We identified 186 DEGs enriched in eye melanocytes and

229 DEGs enriched in skin melanocytes (log2FC > |0.5|, FDR< 0.01), while 3,387 genes were

not differentially expressed (FDR > 0.01, expressed in ≥10% of cells, **Table S5**). GSEA revealed

that skin melanocytes were enriched for DNA repair pathways and UV response-associated

genes (**Figure 3D**). In contrast, eye melanocytes showed enrichment for pathways involved in

UV response downregulation and oxidative phosphorylation. We next re-clustered and directly

compared eye and skin tumors, which also separated by tissue of origin on the UMAP (**Figure**

3E). We identified 230 DEGs enriched in eye tumors and 375 DEGs enriched in skin tumors

(log2FC > 0.5, FDR < 0.01), while 5,754 genes were not differentially expressed between tumor

types. Eye tumors were enriched for epithelial-to-mesenchymal transition (EMT) and

selenoamino acid metabolism pathways (**Figure 3F**). In contrast, skin tumors were enriched for

PI3K–AKT–mTOR signaling and RHO GTPase–mediated activation and YAP1 activity. Together,

these analyses demonstrate that both primary melanocytes and GNAQ^{Q209L}-driven melanomas

exhibit site-specific transcriptional programs.

Melanocyte progenitor cells are highly susceptible to oncogenic GNAQ transformation in

***mitfa*-deficient zebrafish.**

H&E analysis of *nacre* and *casper* zebrafish eyes showed loss of choroidal melanocytes

(**Figure 4A**), whereas melanin in the RPE remained less affected, as previously demonstrated.[42]

These observations imply that an *mitfa*-independent melanocyte progenitor cell persists in *nacre*

and *casper* eyes and that they are susceptible to oncogenic GNAQ-driven transformation. To

identify such cells, we crossed *nacre* (*mitfa$^{w2/w2}$*) with transgenic *Tg(mitfa:GFP)* lines to create

Tg(mitfa:GFP); mitfa$^{w2/w2}$ zebrafish. We then harvested adult eyes at 5 months and performed

anti-GFP immunofluorescence. While GFP+ melanocytes were readily detected in the choroid of

Tg(mitfa:GFP) zebrafish (**Figure 1F, Figure S3**), staining of *Tg(mitfa:GFP); mitfa$^{w2/w2}$* eyes

revealed rare cells with GFP-positivity (**Figure 4B**). Such GFP+ cells were primarily located in

the ciliary body, with isolated cells also found in the choroid, suggesting the persistence of an

mitfa-independent progenitor population in adult *mitfa*-deficient zebrafish eyes (**Figure 4B**). We

next asked whether such cells retain the ability to undergo melanocyte differentiation by using

TEAZ-Eye to deliver plasmids containing the *mitfa*-promoter driving expression of *mitfa* (i.e., to

overexpress *mitfa* mRNA). Interestingly, re-expression of *mitfa* transcript resulted in expansion

of choroidal melanocytes in wild-type animals and rescue of choroidal melanocytes in *mitfa*-

deficient (*nacre*) zebrafish as shown by H&E analysis (**Figure S13A-B'**). To assess whether

GFP-positive cells are sensitive to oncogenic transformation by GNAQ, we expressed

mitfa:GNAQ^{Q209L} in *Tg(mitfa:GFP); mitfa$^{w2/w2}$* zebrafish using TEAZ-Eye. In all injected fish (n=3),

UM tumors were GFP-positive, suggesting they originated from GFP+ progenitor cells within the

eye of *Tg(mitfa:GFP); mitfa$^{w2/w2}$* zebrafish (**Figure S13C**).

Melanocyte progenitor cells express neural crest marker genes in *mitfa*-deficient

zebrafish.

We performed scRNA-Seq and pseudotime analysis on 3 pooled UM tumors from the eyes of

casper zebrafish and compared them to the pooled tissue from 3 normal *casper* eyes (**Figure**

4C, Figure S14). Within the *casper* eyes, we identified a population of cells expressing the

mitfa$^{w2/w2}$ allele that clustered closely with UM tumors under TEAZ-Eye conditions, termed

progenitors (**Figure S15A-B and Tables S6-8**)**.** We re-clustered the progenitor and tumor cells

(**Figure 4D**), revealing seven distinct clusters consisting of neural crest-like (NC-like),

melanoblast-like, proliferating, progenitor, angiogenic, stress-like, as well as an unidentified

cluster marked by *tmsb1, krt91* and *vim* (**Figure 4D, Table S9**). Unsupervised pseudotime

analysis supports a lineage trajectory from the progenitor cells to heterogeneous UM tumor cells

(**Figure S15C**). Notably, the progenitor cell population in normal *casper* eyes expressed neural

crest and melanocyte stem cell marker genes, including *tfap2a, foxd3, sox10, pax3a, vim,* and

yap1 (**Figure 4E-F**). UM tumor cells retained expression of these genes but activated additional

genes associated with pigment progenitor cell function, including *tfec* and *kita* (**Figure 4F**).

Notably, *mitfa* expression in tumors and progenitor cells in *casper* zebrafish reflects the

expression of the mutant *mitfa*$^{w2/w2}$ allele, which functions as a lineage tracer to mark cells that

would normally express *mitfa* under wild-type conditions (**Figure S15B**). Interestingly, despite

the lack of Mitfa activity in *casper* zebrafish, GNAQ^{Q209L}-positive UM tumors included a

melanoblast-like cluster that exhibited expression of pigmentation genes, including *pmela,*

mlpha, and *mlana,* likely explaining the occasional emergence of melanin within tumors as they

progress in *casper* zebrafish.

Conditional deletion of *mitfa* in adult zebrafish fails to recapitulate germline loss-of-

function effects on tumor-free survival in GNAQ^{Q209L}-driven melanoma.

We next asked whether the accelerated growth of GNAQ^{Q209L}-driven tumors in *mitfa*-deficient

zebrafish was attributable to increased oncogene expression or cell-intrinsic loss of Mitfa in

adult melanocytes. scRNA-Seq analysis showed higher GNAQ^{Q209L} expression in wild-type

tumors than in *casper* tumors (**Figure S11B**), thus excluding oncogene overexpression as the

basis for the accelerated phenotype. To directly test cell-intrinsic effects of *mitfa* loss in adult

melanocytes, we performed conditional somatic deletion of *mitfa* using Strategy B plasmids with

either non-targeting gRNAs or gRNAs targeting exon 5 of *mitfa*. TEAZ-Skin was conducted in

three groups: (1) *casper* zebrafish with *mitfa*-targeting gRNAs (n=9; which controls for off-target

effects of the gRNA and for potential function of the *mitfa*$^{w2/w2}$ allele); (2) wild-type zebrafish with

non-targeting gRNAs (n=9); and (3) wild-type zebrafish with *mitfa*-targeting gRNAs (n=9). GFP

expression was confirmed in all injected animals (**Table S11)**. As expected, by 100 dpe tumors

developed in 100% (9 of 9) of *casper* zebrafish. Interestingly, there was no significant difference

in tumor latency between wild-type and *mitfa* conditional-KO, with both indicating significantly

longer latency than in *casper* animals (**Figure 4G–H**). Thus, somatic loss of *mitfa* in adult

melanocytes is insufficient to reproduce the reduced tumor-free survival observed with germline

mitfa deficiency. Notably, *mitfa*-deficient tumors arising in either background developed distal

metastases, a feature not observed in *mitfa* wild-type TEAZ-Skin tumors (**Figure 4H, Figure**

S16). Together, these findings argue against Mitfa as a repressor of GNAQ-induced

transformation in adult melanocytes and indicate that somatic loss in the adult is insufficient to

drive tumor initiation.

***mitfa*-deficient zebrafish exhibit expanded *pax3a* and *tfec* positive melanocyte**

progenitors during embryogenesis.

We next asked whether melanocyte progenitor cells are expanded in *mitfa* mutant zebrafish

relative to wild-type zebrafish. To test this prediction, we performed scRNA-Seq on GFP-positive

cells sorted from *Tg(mitfa:GFP);mitfa*$^{w2/w2}$ and *mitfa*$^{+/w2}$ sibling-matched transgenic zebrafish

embryos at 28 hours post fertilization (hpf), when melanocytes (zebrafish melanophores) begin

to differentiate (**Figure 4I, Figure S14**). We assigned cell-type annotations based on our

previously annotated GFP-positive cells from *Tg(mitfa:GFP)* embryos.[43] The 10 clusters

included six main cell types: neural crest cells (*sox10, foxd3*), a tripotent precursor of

melanoblasts (M), iridoblasts (I), and xanthoblasts (X) (*tfap2a, cdkn1ca, slc15a2, ino80e, id3,*

mycn, tfec), termed MIX cells, a cluster that expressed high levels of melanoblast/xanthoblast

markers (*mitfa, erbb3b, impdh1b, gch2, id3*), termed MX cells, a melanoblast cluster (*mitfa, dct,*

pmel, tyr), as well as two additional clusters corresponding to xanthoblasts and xanthophores

(**Table S10**). For this analysis we referred to MIX and MX cells as melanophore progenitors

(**Figure 4I**). As expected, melanophore cells were reduced in *Tg(mitfa:GFP): mitfa$^{w2/w2}$* zebrafish

(**Figure 4I'**). However, melanophore progenitors were expanded in *Tg(mitfa:GFP); mitfa$^{w2/w2}$*,

while neural crest cells remained unaffected (**Figure 4I-I'**). Melanophore markers were

downregulated in GFP-positive cells sorted from *Tg(mitfa:GFP); mitfa$^{w2/w2}$,* whereas melanocyte

progenitor markers (*pax3a, sox10, foxd3*), as we identified in adult zebrafish eyes, were

upregulated in embryonic progenitors from *Tg(mitfa:GFP); mitfa$^{w2/w2}$* embryos (**Figure 4J, Table**

S12). A role for *pax3a* in adult fish eye tissue has not been previously characterized; however,

recent studies have identified *PAX3* expression in ocular surface melanocytes,[44] and another

study reports an association between *PAX3* and stem cell markers in UM progression.[45]

Transcriptional programs enriched in oncogenic BRAF- versus GNAQ-driven melanoma

overlap with specific melanocyte progenitor populations.

Oncogenic BRAF and NRAS are unable to promote transformation in the absence of *mitfa,*[6,46]

suggesting that distinct transcriptional programs, potentially reflecting unique cellular origins, are

required for UM and CM transformation. To address this hypothesis, we compared previously

published scRNA-Seq datasets in transgenic *Tg(mitfa:BRAF^{V600E}): tp53$^{-/-}$: mitfa$^{w2/w2}$* zebrafish

where TEAZ-Skin was used to rescue *mitfa* and induce melanoma,[46] to our GNAQ^{Q209L}-induced

TEAZ-Skin tumors (**Figure 5A)**. We predicted that by analyzing primary tumors, the

transcriptional signature of the origin cell will be retained and can be identified by overlapping

gene signatures identified in the embryonic melanocyte lineage.[43] Interestingly, while the

expression of *mitfa* was significantly higher in GNAQ-positive tumors, the expression of

pigmentation genes; including *tyrp1b, dct, scl45a2, sox10* and *pmela* were significantly higher in

BRAF-positive melanoma (**Figure S17**). Genes enriched in GNAQ-UM and BRAF-CM were

defined as those with a Log2FC > 1 and Adj p-value < 0.0001 between the two tumor types

(**Figure 5B, Table S13**). We next overlapped this differential gene signature with genes

expressed by cells within the melanocyte lineage isolated from zebrafish embryos (**Figure 5C**).

As expected, BRAF-driven tumors strongly mapped to melanoblast and melanophore clusters

(**Figure 5D**), supporting recent findings that BRAF-driven melanoma originates from

melanoblasts, a melanocyte progenitor population[46,47]. When comparing GNAQ-enriched genes

to the melanocyte lineage, we found that signatures were not confined to a single cluster but

were distributed among neural crest cells, melanocyte progenitors, and xanthophore clusters

(**Figure 5D; Figure S18**). Such results support the hypothesis that GNAQ- and BRAF-driven

melanomas depend on distinct transcriptional programs, reflecting less and more differentiated

melanocytic states, respectively.

We next asked whether less differentiated melanocytes, similar to the *mitfa*-deficient progenitors

in *casper* eyes, are also present in adult wild-type eyes. Since melanocyte stem cells are known

to reside in the zebrafish skin,[48] we included skin tissue as a control and re-clustered

melanocytes from wild-type skin and eyes with progenitor cells from *casper* eyes. Interestingly,

a subset of wild-type melanocytes from both eye and skin samples clustered closely with *casper*

progenitors (**Figure 5E, Figure S19**) and expressed pigment cell progenitor marker genes,

including *pax3a,*[49] *tfec,*[42] *sox10,*[50] *sox4a,*[51,52] *foxd3,*[53] and *yap1* (**Figure 5F**). These findings

suggest either melanocyte plasticity within the eye or the presence of an eye-resident

melanocyte progenitor cell population in adult wild-type zebrafish. In support of this notion,

reanalysis of published scRNA-Seq datasets from human[22] and mouse[54] eyes identified *PAX3-*

positive melanocyte subpopulations (**Figure S20**) Moreover, we confirmed the presence of

PAX3-positive cell populations in adult C57BL/6 mouse eyes by immunohistochemistry (**Figure**

S21). Together, these data support the existence of a *PAX3*-expressing melanocyte precursor

population in both fish and mammalian eyes, which may mirror *PAX3* expression in skin

melanocyte stem cells.[49,55] Such cells may be more susceptible to GNAQ^{Q209L}-driven

transformation and represent candidate cells of origin for UM in this context (**Figure 5G**).

Discussion

Our UM model enables analysis of the earliest events in UM initiation and progression.

Compared with transgenic approaches, this system produces anatomically correct tumors with

shorter latency. Importantly, genes identified in this model are associated with survival

differences in human uveal melanoma, offering an efficient and clinically relevant platform for

studying UM pathogenesis.

We found that fatty acid binding proteins, particularly *fabp3*, are enriched in UM tumors. While

lipid droplets have been identified as vulnerable targets in CM and regulators of melanoma

plasticity,[46,56] lipid signaling in UM has received limited attention.[57-60] Interestingly, high

expression of *FABP3* and *FABP5* are associated with worse disease-free survival in UM but not

CM patients. Our analysis also identified *LGALS2* and *GCH*, two fatty acid metabolism-related

genes whose expression was strongly associated with worse overall survival in UM but showed

an inverse relationship in CM. We additionally found increased *kita* expression in GNAQ-driven

tumors compared to choroidal melanocytes, which is particularly notable given recent studies

linking KIT-positive UM tumors with poor prognosis and chromosome 3 monosomy.[61] In addition,

met expression was elevated in UM tumors relative to choroidal melanocytes. MET signaling is

a key pathway in UM pathogenesis, with established roles in tumor invasion, metastatic

progression, and responsiveness to hepatocyte growth factor (HGF), particularly in the context

of liver metastasis.[62,63] Notably, MET activation has also been described in other cancer types

through ligand independent fibronectin-integrin α5β1-mediated signaling.[64] Given the marked

increase of *fn1a* in choroidal CAFs, it is plausible that fibroblast-driven ECM remodeling

enhances MET signaling in UM. Furthermore, expression of these genes was highest in

molecular subclasses 3 and 4 of the UM TCGA dataset which represents the subset of patients

with the highest rates of BAP1 loss and metastasis.[26] Such observations demonstrate the utility

of our model in uncovering key genes with disease-specific roles in UM whose function could be

further explored through genetic perturbations in zebrafish.

To explore how anatomical context influences transcriptional states, we compared tumors

induced by identical genetic perturbations in the skin and eye. We find that transcriptional

profiles of UM tumors arising from melanocytes in the choroid and skin are influenced by more

than the intrinsic differences of their cells of origin. For example, eye tumors were enriched for

EMT programs compared with skin tumors. In contrast, eye melanocytes did not exhibit these

signatures relative to skin melanocytes, suggesting that signals from the ocular

microenvironment may promote a more mesenchymal state. Alternatively, these differences may

represent a unique cellular origin of UM in the eye and skin. Further investigation is needed to

determine whether these differences are driven by the microenvironment or by intrinsic

properties of melanocytes or precursors cells from distinct anatomical sites.

An interesting distinction between GNAQ and BRAF oncogenes is their differential dependency

on Mitfa activity for tumorigenesis. While BRAF^{V600E} requires Mitfa activity to induce melanoma,[6]

GNAQ^{Q209L}-driven tumor latency was significantly reduced in *mitfa*-deficient zebrafish compared

to wild-type controls. Our results are consistent with prior studies showing that *mitfa*-deficient

zebrafish activate distinct signaling pathways, favoring YAP over MAPK signaling, and give rise

to tumors with reduced latency compared to wild type tumors.[5] We observed *yap1* expression in

melanocyte progenitor cells of *mitfa*-deficient and wild type zebrafish, suggesting that Yap1

activity may be carried over from the progenitor cell during transformation. While germline *mitfa*

loss accelerates tumor onset, our data suggest this effect is not solely due to the deletion of

Mitfa from differentiated melanocytes. Instead, we propose that germline *mitfa* deficiency

expands a population of *pax3a*-positive melanocyte progenitor cells, which are more susceptible

to GNAQ-induced transformation. Alternatively, *mitfa* loss may impair the differentiation of

melanocyte progenitors, preventing the formation of GNAQ^{Q209L}-positive nevi and promoting

direct progression to malignancy. Another explanation for decreased tumor latency in *casper*

zebrafish could be that germline loss of *mitfa* has effects on other cell types, such as immune

cells, which may influence tumor behavior.

Although Mitfa-independent melanocytic cells are unlikely to be the cellular origin of CM in

zebrafish models, studies have shown that MITF-independent BRAF-positive melanoma cells

are central to disease recurrence[65]. GNAQ-driven transformation may be facilitated by Mitfa

paralogs such as Tfec and Tfeb. The Mitfa paralog Tfec has been shown to activate

pigmentation genes in both the retinal pigment epithelium and melanocyte progenitor cells. Tfec

can also rescue ectopic melanocytes in *nacre* zebrafish embryos.[42,66] Moreover, we have

recently shown that the MITF paralog TFE3 promotes cellular plasticity in MITF-low melanoma

cells,[67] suggesting that MITF paralogs are compelling candidate transcription factors to drive

UM onset in *casper* zebrafish.

In our zebrafish study, melanocyte progenitors inherently express neural crest-associated

programs, which are retained during GNAQ^{Q209L}-driven transformation. Re-analysis of mouse

and human single-cell datasets reveals a subpopulation of *PAX3*-positive melanocytes,

suggesting that similar progenitor-like states may be conserved in mammals. Together, these

findings highlight how lineage plasticity contribute to UM initiation and progression across

species. Such observations are supported by an elegant study by Xu, Karreth and colleagues,

who established an immune-competent mouse UM model showing that transcriptional plasticity

from melanocytic to neural crest-like states drives tumor progression.[68]

Identifying the transcriptional and signaling mechanisms within *mitfa*-independent melanocyte

progenitor cells that facilitate UM onset and progression will have significant clinical

implications, particularly for the differential diagnosis of high-risk lesions and the design of

targeted therapies.

Limitation of study

The use of *tp53* and *ptena/b* mutations in our UM model does not fully recapitulate the genetics

of human UM. Mutations in *TP53* and *PTEN* are rarely observed in UM, where loss-of-function

mutations in *BAP1* represent the most frequent tumor suppressor alteration.[39,69,70] However,

both *TP53* and *PTEN* signaling pathways have been shown to be altered in UM pathogenesis.

MDM2 overexpression, an upstream negative regulator of p53, and downregulation of *PERP*, a

downstream effector of p53, have been demonstrated in many UM patients and are associated

with worse outcome.[71-75] Studies have also shown that over half of UM tumors have decreased

PTEN immunostaining and its loss is correlated with decreased survival.[39-41] Whether genetic

knockout of these tumor suppressors recapitulate the dysregulation of these pathways in human

UM remains an open question. Another limitation to our approach when examining the

differences between genetically identical skin and eye-derived tumors is the inability to

distinguish between microenvironmental effects and differences in cell of origin (melanocytes in

the skin versus in the eye). To address this question, a syngeneic or immunodeficient model

could be applied to transplant cells from a primary UM tumor into both anatomical sites.

Additionally, our use of the *mitfa* promoter to drive oncogene expression presents a challenge in

defining the exact cell of origin, as this promoter is active in both melanocyte progenitors and

differentiated melanocytes. As such, it remains unclear whether tumors in *mitfa* wild-type fish (or

those with conditional *mitfa* deletion) arise from a progenitor or a mature melanocyte. Lineage

tracing studies will be essential to determine whether differentiated melanocytes possess the

capacity to initiate primary UM in *mitfa*-competent or conditional *mitfa*-KO zebrafish.

Acknowledgments

This work was supported by the Holden Comprehensive Cancer Center (P30 CA086862).

Additional support to C.K. was provided by an institutional research grant from the American

Cancer Society administered through the HCCC (IRG-18-164-43), the Melanoma Research

Alliance (ID#1426690) and Outrun the Sun in honor of Doug Stickney. J.Y. and D.R. were

supported by NIH predoctoral fellowships (T32 GM144636 and T90 DE023520, respectively).

Additional NIH support was provided to C.K. (R03 CA297549-01), R.A.C. (R01 AR062457),

R.F.M. (P30 EY025580), and D.L. (R03 CA288281-01). Further funding was provided by the

Leo Foundation (LF-OC-21-000888) to D.L., the Wisconsin Partnership Program (AAN4246) to

A.B., and the U.S. Department of Defense (ME240045) to A.B. The funders had no role in study

design; data collection, analysis, or interpretation; manuscript preparation; or the decision to

publish.

Figure 1: $GNAQ^{Q209L}$-induced melanoma by TEAZ-Skin and TEAZ-Eye.

(**A**) TEAZ-Skin was used to induce GNAQ-positive melanoma using two strategies. Strategy A involves a single plasmid construct containing expression cassettes driven by the *mitfa* promoter for $GNAQ^{Q209L}$-T2A-eGFP and Cas9, as well as a U6 promoter driving the expression of three separate gRNAs (targeting *tp53*, *ptena*, and *ptenb*, or a non-targeting control). Strategy B uses a multi-plasmid approach with five plasmids for TEAZ-Skin. Expression cassettes include: (a) *mitfa*:$GNAQ^{Q209L}$-*mitfa*:eGFP, (b) *mitfa*:Cas9, (c) *U6*:gRNA (*tp53*), (d) *U6*:gRNA (*ptena*), and (e) *U6*:gRNA (*ptenb*).

(**B**) Schematic of TEAZ-Skin using wild type (*AB*), *nacre* (*mitfa$^{w2/w2}$*) and *casper* (*mitfa$^{w2/w2}$*: *mpv17$^{a9/a9}$*) zebrafish lines.

(**C**) TEAZ-Skin using Strategy B plasmids in adult wild-type (*AB*) and *casper* zebrafish. Representative brightfield and fluorescent images shown at 4, 8, and 12 weeks post electroporation for WT and at 2, 4, and 6 weeks for *casper*. Scale bar = 0.5 cm.

(**D**) Kaplan-Meier curves were used to estimate tumor-free survival, defined as the time from electroporation until the first visible appearance of GFP+ nodular tumors, in adult zebrafish via TEAZ-Skin. Strategy B wild-type (n=9), *nacre* (n=5), and *casper* (n=24) curves and TEAZ-Skin ($GNAQ^{Q209L}$:GFP) curves (n=3 per condition) compared pairwise with logrank (Mantel-Cox) test. ns = not significant, * p-value<0.05; ** p-value<0.01; ***p-value<0.001; ****p-value<0.0001.

(**E**) Schematic of the zebrafish eye with dashed box indicating the retina and choroid. Comparative H&E of the retina and choroid from a wild-type zebrafish (left) and human donor (right) (20×). Choroidal melanocytes indicated with red arrows. VIT, vitreous; ILM, internal limiting membrane, GCL, ganglion cell layer; IPL, inner plexiform layer, INL, inner nuclear layer; ONL, outer nuclear layer; POS, photoreceptor outer segments; RPE, retinal pigmented epithelium; CH, choroid; SCL, Sclera. Scale bars = 20 µm.

(**F**) Immunofluorescent analysis of transgenic *Tg(mitfa:GFP)* zebrafish using an anti-GFP antibody. Sections were counterstained with DAPI to visualize nuclei. Representative image

showing GFP expression in *mitfa*-positive cells. Secondary-only antibody was used as

background control. Scale bar = 50 µm.

(**G**) Schematic of TEAZ-Eye using choroidal targeted injection.

(**H-H'**) Live imaging of adult wild-type zebrafish following TEAZ-Eye with Strategy B plasmids to

induce uveal melanoma. Brightfield imaging and the corresponding GFP fluorescence images

highlighting tumor progression. Timepoints in day post-electroporation (dpe) as labeled. (H)

Scale bar = 250 µm. (H') Scale bar = 0.5 cm.

(**I-I'**) Live imaging of adult *casper* zebrafish over 21 days post-electroporation (dpe) using TEAZ-

Eye with Strategy B plasmids to induce uveal melanoma. Brightfield imaging and the

corresponding GFP fluorescence images highlighting tumor progression. (I) Scale bar = 250

µm. (I') Scale bar = 0.5 cm.

(**J**) Kaplan-Meier curves comparing tumor-free survival among wild-type (n=5) and *casper*

(n=13) zebrafish following TEAZ-Eye induction using Strategy B plasmids. Curves compared

pairwise with logrank (Mantel-Cox) test. ** p-value<0.01.

(**K, K'**) H&E analysis of uveal melanoma in *casper* zebrafish induced by TEAZ-Eye. Images

were acquired at 20× magnification and stitched from 85 fields using Thunder imaging software.

Normal eye tissue and uveal melanoma as labeled. Scale bar = 100 µm.

(**K'**) Representative 40x magnification image showing GFP expression in tumors cells after anti-

GFP immunofluorescent analysis of TEAZ-Eye induced uveal melanoma in *casper* zebrafish.

Figure 2: Single cell profiling of uveal melanoma and the TME in adult zebrafish.

(**A**) UMAP representation of single-cell RNA-Seq (scRNA-Seq) from three dissociated wild-type

eyes harboring uveal melanoma (UM). UM tumors were generated by TEAZ-Eye (sequencing

replicates, n=2). Control eyes represent pools of two, tumor-free paired wild-type eyes and four

sibling-matched wild-type eyes (sequencing replicates, n=2). Annotated cell clusters as labeled.

(**B**) UMAP and feature plot showing *eGFP* and *GNAQ*^Q209L expression in control eyes and UM

tumor. Red dotted line highlights the melanocyte and UM clusters.

(**C**) Comparison of UM tumor marker genes with the markers of cell types within the control eye

microenvironment. Only genes with a log2FC > 1 and FDR < 0.01 were included in the analysis.

(**D**) Dot plot representing enriched genes (log2FC > 1; q-value < 0.01) in choroidal cell

populations, UM tumors cells, immune cell populations, fibroblasts, Schwann cells, pericytes,

and vascular endothelial cells as shown. Dot size indicates the percentage of cells expressing

each gene; color intensity (grey to blue) reflects normalized average expression levels (low to

high).

(**E**) UMAP clustering of cells in the choroidal microenvironment. Annotated cell clusters as

labelled.

(**F**) Comparison of relative cluster sizes between the two conditions, control and TEAZ-Eye,

shown in (E). Schwann cells serve as a control cluster, displaying no change in frequency

between conditions. Gray = TEAZ-Eye, Blue = Control.

(**G**) Venn diagram representing the overlap of genes between UM tumors and choroidal

melanocytes compared to the cells in their respective microenvironments. Enriched genes for

each condition were identified by differential expression analysis comparing melanocytes to the

microenvironment (ME) or tumor cells to the tumor microenvironment (TME), using a threshold

of log2FC > 1 and FDR < 0.01. Genes of interest have been labelled in their respective subsets,

bolded genes represent genes associated with worse overall survival or disease-free survival in

human UM TCGA datasets. (G') HOMER promoter motif enrichment analysis of genes enriched

in UM tumors vs TME. The top ranked motif for "known" and "*de novo*" motifs are shown with

their corresponding p-values.

(**H**) Dot plot showing enriched genes (log2FC > 1; FDR < 0.01) in UM tumor cells, choroidal

melanocytes, or the tumor microenvironment (TME). Dot size indicates the percentage of cells

expressing each gene; color intensity (grey to blue) reflects normalized average expression

levels (low to high).

(**I-K**) Kaplan–Meier disease-free survival and overall survival curves generated using uveal

melanoma TCGA data. Patients stratified by relative expression levels of *KIT*, *FABP3*, or

LGALS2. Gene expression thresholds were defined by median expression within each cohort.

Log rank p-values as shown. Dotted lines represent 95% confidence interval.

(**L**) Dot plot showing enriched genes (log2FC > 1; FDR < 0.01) in cancer-associated fibroblasts

(CAF), normal fibroblasts (NF), or the tumor microenvironment (TME) without CAF. Dot size

indicates the percentage of cells expressing each gene; color intensity (grey to blue) reflects

normalized average expression levels (low to high). Genes have been grouped according to

their labelled functions.

Figure 3: Comparison of GNAQ-driven tumors and primary melanocytes in the skin and

eyes of adult zebrafish.

(**A**) UMAP representation of GNAQ-positive tumors generated using TEAZ-Eye and TEAZ-Skin,

along with their respective control tissues using scRNA-Seq. Tumor samples include three

dissociated wild-type eyes harboring UM tumors generated via TEAZ-Eye and two dissociated

wild-type skin UM tumors generated via TEAZ-Skin (n=2 sequencing replicates each). Control

eye samples consist of a pool of two uninjected tumor paired wild-type eyes, three mock-

injected sibling-matched wild-type eyes, and four uninjected sibling-matched wild-type eyes

(n=3 sequencing replicates). Control skin samples consist of dissociated normal skin from two

wild-type zebrafish (n=2 sequencing replicates). Cell clusters are annotated as tumor cells (red

= Eye; Blue = Skin), tumor microenvironment (TME) cells shared by both eye and skin samples

(light gray), and TME cells specific to the eye samples (dark gray).

(**B**) Venn diagram showing overlapping genes between TEAZ-Eye, TEAZ-Skin, choroidal

melanocytes, and skin melanocytes compared to the cells in their respective

microenvironments. Enriched genes for each condition were identified by differential expression

analysis comparing melanocytes to the microenvironment (ME) or tumor cells to the tumor

microenvironment (TME), using a threshold of log2FC > 0.5 and FDR < 0.01. Bar chart

displaying overlaps between enriched gene sets from (B). Percentages represented as the

number of overlapping genes over the total unique genes in both gene sets.

(**C**) UMAP representation of primary melanocytes from both eye (salmon) and skin (teal).

Volcano plot of differentially expressed genes (DEGs) between eye and skin primary

melanocytes (log2FC > |0.5| and FDR < 0.01).

(**D**) GSEA enrichment plots of pathways that were enriched in skin melanocytes and of

pathways enriched in eye melanocytes from (C). P-values and FDRs as labelled.

(**E**) UMAP representation of GNAQ^{Q209L}-induced melanoma from both eye (salmon) and skin

(teal). Volcano plot of differentially expressed genes (DEGs) between eye and skin tumors

(log2FC > |0.5| and FDR < 0.01).

(**F**) GSEA enrichment plots of pathways that were enriched in skin melanoma and of pathways

enriched in eye melanoma from (E) P-values and FDRs as labelled.

Figure 4: *mitf*-independent melanocyte progenitor cells are expanded in *mitfa*-deficient

zebrafish and are susceptible to *GNAQ*^{Q209L} transformation.

(**A**) H&E analysis representation of the retina and choroid in wild-type (left), *nacre* (middle), and

casper (right) zebrafish (20x magnification). Note the loss of melanocytes (red arrows) in the

choroid of *nacre* and *casper* eyes. "Vit" = vitreous, "GCL" = ganglion cell layer, "INL" = inner

nuclear layer, "ONL" = outer nuclear layer, "POS" = photoreceptor outer segment, "RPE" =

retinal pigmented epithelium, "CH" = choroid. Scale bar = 50 μm.

(**B**) Immunofluorescent analysis of transgenic *Tg(mitfa:GFP); mitfa*$^{w2/w2}$ (*nacre*) zebrafish using

an anti-GFP antibody. Sections were counterstained with DAPI to visualize nuclei.

Representative images showing GFP-positive cells in the ciliary body and choroid are shown by

yellow arrow heads. Secondary-only antibody was used as background control. Scale bar = 20

μm.

(**C**) UMAP representation of single-cell RNA sequencing (scRNA-Seq) data from three paired

dissociated *casper* eyes and from three *casper* eyes harboring uveal melanoma (UM) tumors

induced using Strategy B via TEAZ-Eye (n=1, sequencing replicate). Annotated cell clusters are

labeled. Progenitor cells and UM tumor clusters are outlined with dashed red lines.

Representative brightfield images of a normal eye and a UM tumor in *casper* zebrafish are

shown in the top left corner of the UMAP. Scale bar = 250 μm.

(**D**) Re-clustered UMAP representing progenitor cells from control eyes and uveal melanoma

(UM) tumor cells from TEAZ-Eye–injected *casper* zebrafish. Heterogeneous cell populations as

labeled.

(**E**) Re-clustered UMAP from (D) showing melanocyte progenitor cells from control eyes in red

and UM tumor cells from TEAZ-Eye in blue. Feature plots for *pax3a, foxd3* and *sox10*

expression are shown, dashed line marks the progenitor cell population.

(**F**) Dot plot illustrating differentiational expressed genes (log2FC > 0.5; FDR < 0.05) between

heterogeneous cell populations in (D). Dot size indicates the percentage of cells expressing

each gene; color intensity (blue to red) reflects normalized average expression levels (low to

high).

(**G**) Kaplan–Meier curves comparing tumor-free survival in wild-type and *casper* zebrafish

following TEAZ-Eye injection using Strategy B plasmids for *mitfa* conditional knockout. Wild-type

zebrafish were injected with either Strategy B plasmids plus non-targeting gRNAs (n=9) or

Strategy B plasmids plus *mitfa*-targeting gRNAs (n=9). *Casper* zebrafish were injected with

Strategy B plasmids and *mitfa*-targeting gRNAs (n=9).

(**H**) Representative brightfield and GFP-overlay images of tumors that formed in *casper* (9 of 9)

and wild-type zebrafish injected with *mitfa*-targeting gRNAs (8 of 9). Metastatic cells are

indicated by red arrows. Scale bar = 0.5 cm.

(**I, I'**) UMAP obtained after clustering GFP-positive cells sorted from *Tg(mitfa:GFP)* and

Tg(mitfa:GFP); mitfa$^{w2/w2}$ (*nacre*) zebrafish embryos at 28 hours post fertilization (hpf).

Annotated cell clusters as labelled. (**I'**) Comparison of relative cluster size for the two

genotypes; *Tg(mitfa:GFP)* and *Tg(mitfa:GFP); mitfa$^{w2/w2}$* (*nacre*) in (I). Scale bar = 0.5 mm.

(**J**) Dot plot illustrating DEGs (log2FC > 0.5; FDR < 0.05) in melanophores and progenitor cells

in *mitfa$^{w2/w2}$* and sibling embryos at 28hpf. Dot size indicates the percentage of cells expressing

each gene; color intensity (grey to red) reflects normalized average expression levels (low to

high).

Figure 5: Oncogenic GNAQ and BRAF transform transcriptionally distinct cells within the

melanocyte lineage.

(**A**) UMAP obtained after clustering tumor cells from *Tg(mitfa:BRAF^{V600E}): tp53$^{-/-}$: mitfa$^{w2/w2}$*

zebrafish where TEAZ-Skin was used to rescue *mitfa* expression and knockout *ptena* and *ptenb*

to induce cutaneous melanoma,[46] along with our GNAQ-induced tumors by TEAZ-Skin (Strategy

B: *mitfa:Cas9, U6*:gRNA-*tp53, U6*:gRNA-*ptena, U6*:gRNA-*ptenb, mitfa:GNAQ^{Q209L}-mitfa:GFP*).

BRAF-induced tumors in red and GNAQ-induced tumors in blue.

(**B**) Violin plot showing differentially expressed genes (FDR < 0.05) between BRAF-induced and

GNAQ-induced tumors from (A). The plot displays the distribution of gene expression across

individual cells, with the width indicating cell density at specific expression levels and the height

representing the range of expression values.

(**C**) Integrated UMAP after clustering GFP-positive cells sorted from *Tg(mitfa:GFP)* and

Tg(mitfa:GFP); mitfa^{w2/w2} (*nacre*) zebrafish embryos at 28 hours post fertilization (hpf).

Annotated cell clusters as labelled.

(**D**) Average enrichment of signature genes from BRAF- and GNAQ-driven tumors projected

onto the UMAP in (C) using UCell with 'FeaturePlot' function. Signature genes represent the top

50 enriched genes from each tumor genotype that are also expressed in the embryonic

melanocyte lineage.

(**E**) UMAP of primary melanocytes and melanocyte progenitors from 9 control wild-type eyes, 2

control wild-type skin sections, and 3 control *casper* eyes. Cells are labeled according to their

genotype (*casper* and wild-type) and tissue origin (eye and skin).

(**F**) Feature plots representing the relative mRNA expression of *pax3a*, *tfec, sox4a, yap1, sox10*,

foxd3 and *ednrba* in melanocytes described in (E). Color intensity (grey to red) reflects

normalized average expression levels (low to high). The red dotted line indicates cells that are

clustering closely with the melanocyte progenitors identified in *casper* eyes.

(**G**) Schematic illustrating GNAQ^{Q209L}-driven uveal melanoma (UM) in the context of tumor

suppressor loss. Melanocyte progenitor cells (*pax3a/b, tfec, sox4a*) can directly give rise to UM

when oncogenic GNAQ^{Q209L} is present (black arrow), or differentiate into MITF^{high} melanocytes

that preferentially form nevi (black arrow). These nevi may subsequently progress to UM

following reduced MITF activity (gray dashed arrow). In contrast, germline *mitfa* loss blocks

melanocyte differentiation and expands melanocyte progenitors, enabling GNAQ^{Q209L} to drive

UM directly from this progenitor population.

1.   Jager, M.J., Shields, C.L., Cebulla, C.M., Abdel-Rahman, M.H., Grossniklaus, H.E., Stern,
M.H., Carvajal, R.D., Belfort, R.N., Jia, R., Shields, J.A., and Damato, B.E. (2020). Uveal
melanoma. Nat Rev Dis Primers *6*, 24. 10.1038/s41572-020-0158-0.

2.   Van Raamsdonk, C.D., Bezrookove, V., Green, G., Bauer, J., Gaugler, L., O'Brien, J.M.,
Simpson, E.M., Barsh, G.S., and Bastian, B.C. (2009). Frequent somatic mutations of
GNAQ in uveal melanoma and blue naevi. Nature *457*, 599-602. 10.1038/nature07586.

3.   Kaliki, S., and Shields, C.L. (2017). Uveal melanoma: relatively rare but deadly cancer. Eye
(Lond) *31*, 241-257. 10.1038/eye.2016.275.

4.   Yu, F.X., Luo, J., Mo, J.S., Liu, G., Kim, Y.C., Meng, Z., Zhao, L., Peyman, G., Ouyang, H.,
Jiang, W., et al. (2014). Mutant Gq/11 promote uveal melanoma tumorigenesis by
activating YAP. Cancer Cell *25*, 822-830. 10.1016/j.ccr.2014.04.017.

5.   Phelps, G.B., Amsterdam, A., Hagen, H.R., García, N.Z., and Lees, J.A. (2022). MITF
deficiency and oncogenic GNAQ each promote proliferation programs in zebrafish
melanocyte lineage cells. Pigment Cell Melanoma Res *35*, 539-547. 10.1111/pcmr.13057.

6.   Ceol, C.J., Houvras, Y., Jane-Valbuena, J., Bilodeau, S., Orlando, D.A., Battisti, V., Fritsch,
L., Lin, W.M., Hollmann, T.J., Ferré, F., et al. (2011). The histone methyltransferase
SETDB1 is recurrently amplified in melanoma and accelerates its onset. Nature *471*, 513-
517. 10.1038/nature09806.

7.   Singh, A.P., Dinwiddie, A., Mahalwar, P., Schach, U., Linker, C., Irion, U., and Nüsslein-
Volhard, C. (2016). Pigment Cell Progenitors in Zebrafish Remain Multipotent through
Metamorphosis. Dev Cell *38*, 316-330. 10.1016/j.devcel.2016.06.020.

8.   Callahan, S.J., Tepan, S., Zhang, Y.M., Lindsay, H., Burger, A., Campbell, N.R., Kim, I.S.,
Hollmann, T.J., Studer, L., Mosimann, C., and White, R.M. (2018). Cancer modeling by
Transgene Electroporation in Adult Zebrafish (TEAZ). Dis Model Mech *11*.
10.1242/dmm.034561.

9.   D'Agati, G., Beltre, R., Sessa, A., Burger, A., Zhou, Y., Mosimann, C., and White, R.M.
(2017). A defect in the mitochondrial protein Mpv17 underlies the transparent casper
zebrafish. Dev Biol *430*, 11-17. 10.1016/j.ydbio.2017.07.017.

10.   White, R.M., Sessa, A., Burke, C., Bowman, T., LeBlanc, J., Ceol, C., Bourque, C., Dovey,
M., Goessling, W., Burns, C.E., and Zon, L.I. (2008). Transparent adult zebrafish as a tool
for in vivo transplantation analysis. Cell Stem Cell *2*, 183-189.
10.1016/j.stem.2007.11.002.

11.   Lister, J.A., Robertson, C.P., Lepage, T., Johnson, S.L., and Raible, D.W. (1999). nacre
encodes a zebrafish microphthalmia-related protein that regulates neural-crest-derived
pigment cell fate. Development *126*, 3757-3767. 10.1242/dev.126.17.3757.

12.   Butler, A., Hoffman, P., Smibert, P., Papalexi, E., and Satija, R. (2018). Integrating single-
cell transcriptomic data across different conditions, technologies, and species. Nat
Biotechnol *36*, 411-420. 10.1038/nbt.4096.

13.   Sur, A., Wang, Y., Capar, P., Margolin, G., Prochaska, M.K., and Farrell, J.A. (2023). Single-
cell analysis of shared signatures and transcriptional diversity during zebrafish
development. Dev Cell *58*, 3028-3047.e3012. 10.1016/j.devcel.2023.11.001.

14.   Voigt, A.P., Whitmore, S.S., Lessing, N.D., DeLuca, A.P., Tucker, B.A., Stone, E.M., Mullins,
R.F., and Scheetz, T.E. (2020). Spectacle: An interactive resource for ocular single-cell
RNA sequencing data analysis. Exp Eye Res *200*, 108204. 10.1016/j.exer.2020.108204.

15.   Kolberg, L., Raudvere, U., Kuzmin, I., Adler, P., Vilo, J., and Peterson, H. (2023). g:Profiler-
interoperable web service for functional enrichment analysis and gene identifier
mapping (2023 update). Nucleic Acids Res *51*, W207-w212. 10.1093/nar/gkad347.
16.   Krämer, A., Green, J., Pollard, J., Jr., and Tugendreich, S. (2014). Causal analysis
approaches in Ingenuity Pathway Analysis. Bioinformatics *30*, 523-530.
10.1093/bioinformatics/btt703.
17.   Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A.,
Paulovich, A., Pomeroy, S.L., Golub, T.R., Lander, E.S., and Mesirov, J.P. (2005). Gene set
enrichment analysis: a knowledge-based approach for interpreting genome-wide
expression profiles. Proc Natl Acad Sci U S A *102*, 15545-15550.
10.1073/pnas.0506580102.
18.   Trapnell, C., Cacchiarelli, D., Grimsby, J., Pokharel, P., Li, S., Morse, M., Lennon, N.J.,
Livak, K.J., Mikkelsen, T.S., and Rinn, J.L. (2014). The dynamics and regulators of cell fate
decisions are revealed by pseudotemporal ordering of single cells. Nat Biotechnol *32*,
381-386. 10.1038/nbt.2859.
19.   Bloh, K., Kanchana, R., Bialk, P., Banas, K., Zhang, Z., Yoo, B.C., and Kmiec, E.B. (2021).
Deconvolution of Complex DNA Repair (DECODR): Establishing a Novel Deconvolution
Algorithm for Comprehensive Analysis of CRISPR-Edited Sanger Sequencing Data. Crispr j
4, 120-131. 10.1089/crispr.2020.0022.
20.   Richardson, R., Tracey-White, D., Webster, A., and Moosajee, M. (2017). The zebrafish
eye-a paradigm for investigating human ocular genetics. Eye (Lond) *31*, 68-86.
10.1038/eye.2016.198.
21.   McMenamin, P.G., Shields, G.T., Seyed-Razavi, Y., Kalirai, H., Insall, R.H., Machesky, L.M.,
and Coupland, S.E. (2020). Melanoblasts Populate the Mouse Choroid Earlier in
Development Than Previously Described. Invest Ophthalmol Vis Sci *61*, 33.
10.1167/iovs.61.10.33.
22.   Voigt, A.P., Mulfaul, K., Mullin, N.K., Flamme-Wiese, M.J., Giacalone, J.C., Stone, E.M.,
Tucker, B.A., Scheetz, T.E., and Mullins, R.F. (2019). Single-cell transcriptomics of the
human retinal pigment epithelium and choroid in health and macular degeneration. Proc
Natl Acad Sci U S A *116*, 24100-24107. 10.1073/pnas.1914143116.
23.   Howe, D.G., Bradford, Y.M., Conlin, T., Eagle, A.E., Fashena, D., Frazer, K., Knight, J., Mani,
P., Martin, R., Moxon, S.A., et al. (2013). ZFIN, the Zebrafish Model Organism Database:
increased support for mutants and transgenics. Nucleic Acids Res *41*, D854-860.
10.1093/nar/gks938.
24.   Tatarakis, D., Cang, Z., Wu, X., Sharma, P.P., Karikomi, M., MacLean, A.L., Nie, Q., and
Schilling, T.F. (2021). Single-cell transcriptomic analysis of zebrafish cranial neural crest
reveals spatiotemporal regulation of lineage decisions during development. Cell Rep *37*,
110140. 10.1016/j.celrep.2021.110140.
25.   Simões-Costa, M., and Bronner, M.E. (2015). Establishing neural crest identity: a gene
regulatory recipe. Development *142*, 242-257. 10.1242/dev.105445.
26.   Robertson, A.G., Shih, J., Yau, C., Gibb, E.A., Oba, J., Mungall, K.L., Hess, J.M.,
Uzunangelov, V., Walter, V., Danilova, L., et al. (2017). Integrative Analysis Identifies Four
Molecular and Clinical Subsets in Uveal Melanoma. Cancer Cell *32*, 204-220.e215.
10.1016/j.ccell.2017.07.003.

27. Hunter, M.V., Joshi, E., Bowker, S., Montal, E., Ma, Y., Kim, Y.H., Yang, Z., Tuffery, L., Li, Z.,
Rosiek, E., et al. (2025). Mechanical confinement governs phenotypic plasticity in
melanoma. Nature. 10.1038/s41586-025-09445-6.

28. Hunter, M.V., Moncada, R., Weiss, J.M., Yanai, I., and White, R.M. (2021). Spatially
resolved transcriptomics reveals the architecture of the tumor-microenvironment
interface. Nat Commun 12, 6278. 10.1038/s41467-021-26614-z.

29. Liu, T., Zhou, L., Xiao, Y., Andl, T., and Zhang, Y. (2022). BRAF Inhibitors Reprogram
Cancer-Associated Fibroblasts to Drive Matrix Remodeling and Therapeutic Escape in
Melanoma. Cancer Res 82, 419-432. 10.1158/0008-5472.Can-21-0614.

30. Mao, X., Xu, J., Wang, W., Liang, C., Hua, J., Liu, J., Zhang, B., Meng, Q., Yu, X., and Shi, S.
(2021). Crosstalk between cancer-associated fibroblasts and immune cells in the tumor
microenvironment: new findings and future perspectives. Mol Cancer 20, 131.
10.1186/s12943-021-01428-1.

31. Wang, J., Song, Y., Li, Z., Gao, T., Shen, W., Kang, Z., and Xu, C. (2025). Combining single-
cell and bulk RNA sequencing to identify CAF-related signature for prognostic prediction
and treatment response in patients with melanoma. Sci Rep 15, 29082. 10.1038/s41598-
025-14979-w.

32. Jin, S., Guerrero-Juarez, C.F., Zhang, L., Chang, I., Ramos, R., Kuan, C.H., Myung, P., Plikus,
M.V., and Nie, Q. (2021). Inference and analysis of cell-cell communication using
CellChat. Nat Commun 12, 1088. 10.1038/s41467-021-21246-9.

33. Kaochar, S., Dong, J., Torres, M., Rajapakshe, K., Nikolos, F., Davis, C.M., Ehli, E.A., Coarfa,
C., Mitsiades, N., and Poulaki, V. (2018). ICG-001 Exerts Potent Anticancer Activity
Against Uveal Melanoma Cells. Invest Ophthalmol Vis Sci 59, 132-143. 10.1167/iovs.17-
22454.

34. Zheng, L., Liu, Y., and Pan, J. (2017). Inhibitory effect of pyrvinium pamoate on uveal
melanoma cells involves blocking of Wnt/β-catenin pathway. Acta Biochim Biophys Sin
(Shanghai) 49, 890-898. 10.1093/abbs/gmx089.

35. Wu, S., Han, M., and Zhang, C. (2022). Overexpression of microRNA-130a represses
uveal melanoma cell migration and invasion through inactivation of the Wnt/β-catenin
signaling pathway by downregulating USP6. Cancer Gene Ther 29, 930-939.
10.1038/s41417-021-00377-7.

36. Zuidervaart, W., van Nieuwpoort, F., Stark, M., Dijkman, R., Packer, L., Borgstein, A.M.,
Pavey, S., van der Velden, P., Out, C., Jager, M.J., et al. (2005). Activation of the MAPK
pathway is a common event in uveal melanomas although it rarely occurs through
mutation of BRAF or RAS. Br J Cancer 92, 2032-2038. 10.1038/sj.bjc.6602598.

37. Chen, X., Wu, Q., Depeille, P., Chen, P., Thornton, S., Kalirai, H., Coupland, S.E., Roose,
J.P., and Bastian, B.C. (2017). RasGRP3 Mediates MAPK Pathway Activation in GNAQ
Mutant Uveal Melanoma. Cancer Cell 31, 685-696.e686. 10.1016/j.ccell.2017.04.002.

38. Rimoldi, D., Salvi, S., Liénard, D., Lejeune, F.J., Speiser, D., Zografos, L., and Cerottini, J.C.
(2003). Lack of BRAF mutations in uveal melanoma. Cancer Res 63, 5712-5715.

39. Abdel-Rahman, M.H., Yang, Y., Zhou, X.P., Craig, E.L., Davidorf, F.H., and Eng, C. (2006).
High frequency of submicroscopic hemizygous deletion is a major mechanism of loss of
expression of PTEN in uveal melanoma. J Clin Oncol 24, 288-295.
10.1200/jco.2005.02.2418.

40.   Pópulo, H., Soares, P., Rocha, A.S., Silva, P., and Lopes, J.M. (2010). Evaluation of the
1110   mTOR pathway in ocular (uvea and conjunctiva) melanoma. Melanoma Res *20*, 107-117.
1111   10.1097/CMR.0b013e32832ccd09.

1112   41.   Woodman, S.E. (2012). Metastatic uveal melanoma: biology and emerging treatments.
1113   Cancer J *18*, 148-152. 10.1097/PPO.0b013e31824bd256.

1114   42.   Petratou, K., Spencer, S.A., Kelsh, R.N., and Lister, J.A. (2021). The MITF paralog tfec is
1115   required in neural crest development for fate specification of the iridophore lineage
1116   from a multipotent pigment cell progenitor. PLoS One *16*, e0244794.
1117   10.1371/journal.pone.0244794.

1118   43.   Kenny, C., Dilshat, R., Seberg, H.E., Van Otterloo, E., Bonde, G., Helverson, A., Franke,
1119   C.M., Steingrímsson, E., and Cornell, R.A. (2022). TFAP2 paralogs facilitate chromatin
1120   access for MITF at pigmentation and cell proliferation genes. PLoS Genet *18*, e1010207.
1121   10.1371/journal.pgen.1010207.

1122   44.   Ulrich, E., Kistenmacher, S., Martin, G., Schlötzer-Schrehardt, U., Seitz, B., Auw-Hädrich,
1123   C., Schlunck, G., Reinhard, T., and Polisetti, N. (2025). PAX3 expression patterns in ocular
1124   surface melanocytes. Sci Rep *15*, 12472. 10.1038/s41598-025-90318-3.

1125   45.   Kumar, N., Singh, M.K., Singh, L., Jha, J., Pushker, N., Lomi, N., Meel, R., Sen, S., and
1126   Kashyap, S. (2025). Prognostic significance of pigmentation and stem cell markers in
1127   Indian population of uveal melanoma. Br J Ophthalmol *109*, 799-808. 10.1136/bjo-2024-
1128   325886.

1129   46.   Lumaquin-Yin, D., Montal, E., Johns, E., Baggiolini, A., Huang, T.H., Ma, Y., LaPlante, C.,
1130   Suresh, S., Studer, L., and White, R.M. (2023). Lipid droplets are a metabolic vulnerability
1131   in melanoma. Nat Commun *14*, 3192. 10.1038/s41467-023-38831-9.

1132   47.   Baggiolini, A., Callahan, S.J., Montal, E., Weiss, J.M., Trieu, T., Tagore, M.M., Tischfield,
1133   S.E., Walsh, R.M., Suresh, S., Fan, Y., et al. (2021). Developmental chromatin programs
1134   determine oncogenic competence in melanoma. Science *373*, eabc1048.
1135   10.1126/science.abc1048.

1136   48.   Brombin, A., Simpson, D.J., Travnickova, J., Brunsdon, H., Zeng, Z., Lu, Y., Young, A.I.J.,
1137   Chandra, T., and Patton, E.E. (2022). Tfap2b specifies an embryonic melanocyte stem cell
1138   that retains adult multifate potential. Cell Rep *38*, 110234.
1139   10.1016/j.celrep.2021.110234.

1140   49.   Kubic, J.D., Young, K.P., Plummer, R.S., Ludvik, A.E., and Lang, D. (2008). Pigmentation
1141   PAX-ways: the role of Pax3 in melanogenesis, melanocyte stem cell maintenance, and
1142   disease. Pigment Cell Melanoma Res *21*, 627-645. 10.1111/j.1755-148X.2008.00514.x.

1143   50.   Dutton, K.A., Pauliny, A., Lopes, S.S., Elworthy, S., Carney, T.J., Rauch, J., Geisler, R.,
1144   Haffter, P., and Kelsh, R.N. (2001). Zebrafish colourless encodes sox10 and specifies non-
1145   ectomesenchymal neural crest fates. Development *128*, 4113-4125.
1146   10.1242/dev.128.21.4113.

1147   51.   Brombin, A., MacMaster, S., Travnickova, J., Wyatt, C., Brunsdon, H., Ramsey, E., Vu, H.N.,
1148   Steingrimsson, E., Chandra, T., and Patton, E.E. (2025). Loss of MITF activity leads to
1149   emergent cell states from the melanocyte stem cell lineage. bioRxiv,
1150   2025.2012.2023.695681. 10.64898/2025.12.23.695681.

52. Frantz, W.T., Iyengar, S., Neiswender, J., Cousineau, A., Maehr, R., and Ceol, C.J. (2023).
Pigment cell progenitor heterogeneity and reiteration of developmental signaling
underlie melanocyte regeneration in zebrafish. Elife *12*. 10.7554/eLife.78942.

53. Thomas, A.J., and Erickson, C.A. (2009). FOXD3 regulates the lineage switch between
neural crest-derived glial cells and pigment cells by repressing MITF through a non-
canonical mechanism. Development *136*, 1849-1858. 10.1242/dev.031989.

54. Youkilis, J.C., and Bassnett, S. (2021). Single-cell RNA-sequencing analysis of the ciliary
epithelium and contiguous tissues in the mouse eye. Exp Eye Res *213*, 108811.
10.1016/j.exer.2021.108811.

55. Lang, D., Lu, M.M., Huang, L., Engleka, K.A., Zhang, M., Chu, E.Y., Lipner, S., Skoultchi, A.,
Millar, S.E., and Epstein, J.A. (2005). Pax3 functions at a nodal point in melanocyte stem
cell differentiation. Nature *433*, 884-887. 10.1038/nature03292.

56. Johns, E., Ma, Y., Louphrasitthiphol, P., Peralta, C., Hunter, M.V., Raymond, J.H., Molina,
H., Goding, C.R., and White, R.M. (2025). The Lipid Droplet Protein DHRS3 Is a Regulator
of Melanoma Cell State. Pigment Cell Melanoma Res *38*, e13208. 10.1111/pcmr.13208.

57. Xu, Y., Xu, W.H., Yang, X.L., Zhang, H.L., and Zhang, X.F. (2020). Fatty acid-binding protein
5 predicts poor prognosis in patients with uveal melanoma. Oncol Lett *19*, 1771-1780.
10.3892/ol.2020.11301.

58. Van Ly, D., Wang, D., Conway, R.M., Giblin, M., Liang, S., Lukeis, R., Lim, L.A., Hesson, L.,
and Cherepanoff, S. (2020). Lipid-Producing Ciliochoroidal Melanoma with Expression of
HMG-CoA Reductase. Ocul Oncol Pathol *6*, 416-421. 10.1159/000510393.

59. Yavuzyigitoglu, S., Kilic, E., Vaarwater, J., de Klein, A., Paridaens, D., and Verdijk, R.M.
(2016). Lipomatous Change in Uveal Melanoma: Histopathological,
Immunohistochemical and Cytogenetic Analysis. Ocul Oncol Pathol *2*, 133-135.
10.1159/000440981.

60. Fiorentzis, M., Kalirai, H., Katopodis, P., and Coupland, S.E. (2017). Adipophilin
expression in primary and metastatic uveal melanoma: a pilot study. Graefes Arch Clin
Exp Ophthalmol *255*, 1049-1051. 10.1007/s00417-017-3630-0.

61. Horvathova Kajabova, V., Soltysova, A., Demkova, L., Plesnikova, P., Lyskova, D., Furdova,
A., Babal, P., and Smolkova, B. (2021). KIT Expression Is Regulated by DNA Methylation in
Uveal Melanoma Tumors. Int J Mol Sci *22*. 10.3390/ijms221910748.

62. Surriga, O., Rajasekhar, V.K., Ambrosini, G., Dogan, Y., Huang, R., and Schwartz, G.K.
(2013). Crizotinib, a c-Met inhibitor, prevents metastasis in a metastatic uveal melanoma
model. Mol Cancer Ther *12*, 2817-2826. 10.1158/1535-7163.Mct-13-0499.

63. Cheng, H., Terai, M., Kageyama, K., Ozaki, S., McCue, P.A., Sato, T., and Aplin, A.E. (2015).
Paracrine Effect of NRG1 and HGF Drives Resistance to MEK Inhibitors in Metastatic
Uveal Melanoma. Cancer Res *75*, 2737-2748. 10.1158/0008-5472.Can-15-0370.

64. Mitra, A.K., Sawada, K., Tiwari, P., Mui, K., Gwin, K., and Lengyel, E. (2011). Ligand-
independent activation of c-Met by fibronectin and $\alpha(5)\beta(1)$-integrin regulates ovarian
cancer invasion and metastasis. Oncogene *30*, 1566-1576. 10.1038/onc.2010.532.

65. Travnickova, J., Wojciechowska, S., Khamseh, A., Gautier, P., Brown, D.V., Lefevre, T.,
Brombin, A., Ewing, A., Capper, A., Spitzer, M., et al. (2019). Zebrafish MITF-Low
Melanoma Subtype Models Reveal Transcriptional Subclusters and MITF-Independent
Residual Disease. Cancer Res *79*, 5769-5784. 10.1158/0008-5472.Can-19-0037.

66.   Lister, J.A., Lane, B.M., Nguyen, A., and Lunney, K. (2011). Embryonic expression of
zebrafish MiT family genes tfe3b, tfeb, and tfec. Dev Dyn *240*, 2529-2538.
10.1002/dvdy.22743.
67.   Chang, J., Campbell-Hanson, K.R., Vanneste, M., Bartschat, N.I., Nagel, R., Arnadottir,
A.K., Vu, H.N., Montgomery, C., Yevdash, J., Jiang, J., et al. (2025). Antagonistic roles for
MITF and TFE3 in melanoma plasticity. Cell Rep *44*, 115474.
10.1016/j.celrep.2025.115474.
68.   Xu, X., Liu, X., Jarajapu, V., Neelature Sriramareddy, S., Konecny, F., Posorske, B., Dollar,
J.J., Liu, X., Jasani, N., Wang, K., et al. (2026). A Multi-Step Immune-Competent
Genetically Engineered Mouse Model Reveals Phenotypic Plasticity in Uveal Melanoma.
Cancer Res. 10.1158/0008-5472.Can-25-2684.
69.   Hajkova, N., Hojny, J., Nemejcova, K., Dundr, P., Ulrych, J., Jirsova, K., Glezgova, J., and
Ticha, I. (2018). Germline mutation in the TP53 gene in uveal melanoma. Sci Rep *8*, 7618.
10.1038/s41598-018-26040-0.
70.   Harbour, J.W., Onken, M.D., Roberson, E.D., Duan, S., Cao, L., Worley, L.A., Council, M.L.,
Matatall, K.A., Helms, C., and Bowcock, A.M. (2010). Frequent mutation of BAP1 in
metastasizing uveal melanomas. Science *330*, 1410-1413. 10.1126/science.1194472.
71.   Brantley, M.A., Jr., and Harbour, J.W. (2000). Deregulation of the Rb and p53 pathways in
uveal melanoma. Am J Pathol *157*, 1795-1801. 10.1016/s0002-9440(10)64817-1.
72.   Coupland, S.E., Anastassiou, G., Stang, A., Schilling, H., Anagnostopoulos, I., Bornfeld, N.,
and Stein, H. (2000). The prognostic value of cyclin D1, p53, and MDM2 protein
expression in uveal melanoma. J Pathol *191*, 120-126. 10.1002/(sici)1096-
9896(200006)191:2<120::Aid-path591>3.0.Co;2-p.
73.   Paraoan, L., Gray, D., Hiscott, P., Ebrahimi, B., Damato, B., and Grierson, I. (2006).
Expression of p53-induced apoptosis effector PERP in primary uveal melanomas:
downregulation is associated with aggressive type. Exp Eye Res *83*, 911-919.
10.1016/j.exer.2006.04.016.
74.   Davies, L., Spiller, D., White, M.R., Grierson, I., and Paraoan, L. (2011). PERP expression
stabilizes active p53 via modulation of p53-MDM2 interaction in uveal melanoma cells.
Cell Death Dis *2*, e136. 10.1038/cddis.2011.19.
75.   de Lange, J., Ly, L.V., Lodder, K., Verlaan-de Vries, M., Teunisse, A.F., Jager, M.J., and
Jochemsen, A.G. (2012). Synergistic growth inhibition based on small-molecule p53
activation as treatment for intraocular melanoma. Oncogene *31*, 1105-1116.
10.1038/onc.2011.309.
1229

FIGURE 1

FIGURE 2

FIGURE 3

A Control eye (n=3); Control skin (n=2);
TEAZ-Eye (n=2), TEAZ-Skin (n=2)

Common TME
Eye TME
Tumor_Skin
Tumor_Eye

B Comparison of Primary Melanocytes and GNAQ-induced Melanoma Across Eye and Skin

Eye Mel vs. ME
Eye Tumor vs. TME
Skin Tumor vs. TME
Skin Mel vs. ME

Total unique gene overlap

552/ 2350   386/ 1823   898/ 2192   414/ 1607

Skin melanocytes
Skin tumor
Eye melanocytes
Eye tumor

C DEGs: <u>Primary Melanocytes</u> Eye vs. Skin
(FDR<0.01, Log2FC >|0.5|)

Primary Melanocytes — Eye, Skin

E DEGs: <u>GNAQ-induced Melanoma</u> Eye vs. Skin
(FDR<0.01, Log2FC >|0.5|)

GNAQ-induced Melanoma — Eye, Skin

D
Up in Skin Melanocytes
DNA Repair
P-value = 0.002
FDR = 0.016

Up in Eye Melanocytes
Oxidative phosphorylation
P-value = 0.00
FDR<0.01

UV Response Up
P-value = 0.00
FDR = 0.1

UV Response Down
P-value = 0.00
FDR<0.1

F
Up in Skin Melanoma
PI3K/AKT/MTOR
P-value <0.009
FDR<0.1

Up in Eye Melanoma
EMT transition
P-value = 0.00
FDR<0.02

RHO GTPase
P-value <0.002
FDR<0.05

Selenoamino acid metabolism
P-value = 0.00
FDR = 0.00

YAP1 activity
P-value <0.002
FDR<0.03

FIGURE 4

FIGURE 5

FIGURE S1

A Tumor Free Survival Following Electroporation

Tumor free survival — *Wild type*

Days

- *ptena/b* CRISPR
- *tp53* CRISPR
- GNAQ: GFP
- **Strategy A**
- **Strategy B**

B Tumor Free Survival Following Electroporation

Tumor free survival — *Nacre*

Days

- *ptena/b* CRISPR
- *tp53* CRISPR
- GNAQ: GFP
- **Strategy A**
- **Strategy B**

B' *Strategy B*

nacre — Week 2

Week 4

Week 6 — GFP

C Tumor Free Survival Following Electroporation

Tumor free survival — *Casper*

Days

- *ptena/b* CRISPR
- *tp53* CRISPR
- GNAQ: GFP
- **Strategy A**
- **Strategy B**

D Tumor Free Survival Following Electroporation

Tumor free survival

Days

- GNAQ: GFP (WT)
- GNAQ: GFP (Nacre)
- GNAQ: GFP (Casper)
- Strategy B (WT)
- Strategy B (Nacre)
- Strategy B (Casper)

E TEAZ-Skin in wild-type

n=10

TEAZ-Skin in *casper*

n=20

Xanthoma in *Tg(mitfa:BRAF^{V600E}); mitfa$^{w2/w2}$; tp53$^{-/-}$)*

n=20

FIGURE S2

Tumor #	ptena	ptenb	tp53	# of edited genes
1	🟩	🟩	🟩	3
2	🟩	🟩	🟩	3
3	🟥	🟩	🟩	2
4	🟩	🟩	🟩	3
5	🟩	🟩	🟩	3
6	🟩	🟩	🟩	3
7	🟩	🟩	🟩	3
8	🟩	🟩	🟩	3
9	🟩	🟩	🟩	3
10	🟩	🟩	🟩	3
11	🟩	🟩	🟩	3
Germline	**ptena**	**ptenb**	**tp53**	
WT1	🟥	🟥	🟥	0
WT2	🟥	🟥	🟥	0

🟩 CRISPR editing

🟥 No editing

A

Tg(mitfa-GFP) Zebrafish Line

B *Tg(mitfa-GFP)* Zebrafish Line

C TEAZ-Eye using *mitfa:GFP* plasmids into WT Zebrafish

Anti-GFP IF

Secondary-only Control

FIGURE S4

Single-cell RNA seq Replicates

Tissue	Genotype	Replicate 1 (pooled tissue, tumors)	Replicate 2 (pooled tissue, tumors)
Skin	Wild-type	1 skin sample	1 skin sample
Skin Tumor	Wild-type	1 tumor sample	1 tumor sample
Eye Paired (tumor matched bilateral eye)	Wild-type	2 pooled eyes	
Eye sibling matched	Wild-type	4 pooled eyes	
Eye mock injected, sibling matched	Wild-type	3 pooled eyes	
Eye Tumor	Wild-type	1 tumor sample	2 pooled tumor samples

FIGURE S5

FIGURE S6

FIGURE S7

FIGURE S8

FABP5 Expression in
4 TCGA Molecular Subclass of UM

FABP3 Expression in
4 TCGA Molecular Subclass of UM

LGALS1 Expression in
4 TCGA Molecular Subclass of UM

GCH Expression in
4 TCGA Molecular Subclass of UM

LGALS2 Expression in
4 TCGA Molecular Subclass of UM

KIT Expression in
4 TCGA Molecular Subclass of UM

MET Expression in
4 TCGA Molecular Subclass of UM

HGF Expression in
4 TCGA Molecular Subclass of UM

FIGURE S9

A

Number of interactions

Interaction weights/strength

B

FN1 Signaling from CAFs

C

Collagen Signaling from CAFs

D

FIGURE S11

A

GNAQ

0.93

1.1

p-value < 0.001

B

GNAQ

1.1

0.69

p-value < 0.001

FIGURE S13

A   Control injected: Wild type

A' TEAZ-Eye (miniCoopR) : Wild type

B   Control injected: *nacre*

B' TEAZ-Eye (miniCoopR): *nacre*

C   TEAZ-Eye: (*mitfa:GNAQ^{Q209L}*)

Tg(mitfa:GFP): mitfa^{w2/w2}

GFP

FIGURE S14
Single-cell RNA seq Replicates

Tissue	Genotype	Replicate 1 (pooled tissue, tumors)
Eye	Casper	3 pooled eyes
Eye Tumor	Casper	3 pooled tumors
Whole embryo 28hpf	Wild-type Tg(mitfa-GFP); mitfa$^{+/w2}$	Pooled embryos 25 total
Whole embryo 28hpf	Nacre Tg(mitfa-GFP); mitfa$^{w2/w2}$	Pooled embryos 25 total

FIGURE S15

FIGURE S16

A TEAZ-Skin: *Strategy B* (Wild type)

16 weeks

Wild type   Lateral

Dorsal

Bright overlay GFP

GFP

GFP

B TEAZ-Skin: *Strategy B + gRNA-mitfa* (Wild type)

8 weeks

Wild type   Lateral

GFP

metastasis

Bright overlay GFP

C TEAZ-Skin: *Strategy B* (Casper)

8 weeks

Lateral

Dorsal

metastasis   primary

Bright overlay GFP

metastasis   primary

Bright overlay GFP

FIGURE S17

FIGURE S18
TEAZ-Eye Strategy B (Wild-type)

FIGURE S19

A Mouse (Eye)

Tyr *Pax3*

Ednrb *Kit*

Melanocytes

RPE

Log2 Exp
3.0

0.0

B Human (Eye)

SCHWANN-MYE

MAST-CELLS SCHWANN-NON

TNK-CELL

ENDOTHELIAL

FIBROBLAST

MACROPHAGE

B-CELL

PERICYTE RPE

Melanocytes

MELANOCYTE

umap-1

umap-1

PAX3 FOXD3 SOX10 EDNRB KIT TFE3 MITF

Schwann-mye

Schwann-non

Melanocyte

Endothelial

Pericyte

Fibroblast

RPE

B-cell

T/NK-cell

macrophage

mast-cells

FIGURE S21