Mitfa-Independent Melanocyte Progenitors are Highly Susceptible to GNAQ-induced Uveal Melanoma in Adult Zebrafish

Julius I. Yevdash,[1] Delaney Robinson,[1] Rachel Moore,[1] Zhijie Li,[1] Katelyn R. Campbell-Hanson,[1] Danielle Gutelius,[2] Stephen P. G. Moore,[2] Dylan Friend,[1] Isaac O'Toole,[1] Collin Montgomery,[1] Jesse D. Riordan,[3,4] Adam J. Dupuy,[3,4] Robert A. Cornell,[5] Elaine M. Binkley,[4,6,7] Deborah Lang,[2] Ronald J. Weigel,[1,3,4] Colin Kenny[1,3,4,5,8]

1. Department of Surgery, College of Medicine, University of Iowa, Iowa City, IA 52242, USA.
2. Department of Dermatology, Boston University School of Medicine, Boston, MA, 02118, USA.
3. Department of Anatomy & Cell Biology, University of Iowa, Iowa City, IA, 52242, USA.
4. Holden Comprehensive Cancer Center, University of Iowa, Iowa City, IA 52242, USA.
5. Department of Oral Health Sciences, University of Washington, School of Dentistry, Seattle, WA, 98195
6. Department of Ophthalmology and Visual Sciences, University of Iowa, Iowa City, IA 52242, Iowa
7. University of Iowa, Institute for Vision Research, Iowa City, IA 52242, Iowa
8. Lead Contact.

* Corresponding author: colin-kenny@uiowa.edu

Summary

Melanocytes reside in diverse microenvironments that influence their susceptibility to oncogenic transformation, however, studying rare melanoma subsets has been hindered by the lack of suitable animal models. We developed a primary, immune-competent zebrafish model to study uveal melanoma (UM), utilizing choroidal-targeted injection and electroporation of plasmids containing human $GNAQ^{Q209L}$ and CRISPR/Cas9 cassettes for tumor suppressor gene deletion. Single-cell transcriptional profiling of genetically identical eye- and skin-derived tumors revealed distinct oncogenic pathways, highlighting the importance of studying melanoma subtypes in their correct anatomical context. Additionally, we identified a population of *tfec*- and *pax3a*-expressing melanocyte progenitor cells in *mitfa*-deficient embryos and adult zebrafish eyes, which were highly susceptible to GNAQ-driven transformation. While previous studies have linked *mitfa* deficiency to accelerated UM onset, our findings suggest that an expanded progenitor population in *mitfa*-deficient animals drives this susceptibility. Our study establishes a critical role for Mitfa-independent melanocyte progenitors in UM pathogenesis

Highlights

- Choroid-targeted electroporation of oncogenic GNAQ induces anatomically correct UM in adult zebrafish.

- Mitfa-independent melanocyte progenitor cells are highly susceptible to GNAQ-driven transformation.

- Germline loss of *mitfa* accelerates UM onset via progenitor enrichment, a phenotype not recapitulated by conditional *mitfa*-KO in adult zebrafish.

- Oncogenic BRAF and GNAQ transform molecularly and developmentally distinct subpopulations of cells within the melanocyte lineage.

Introduction

Uveal melanoma (UM) is a rare cancer of the melanocytes in the eye that is both vision threatening and deadly.[1-4] The majority of UM's (~90%) occur in the choroid, the vascular structure supplying the outer retina, with the remainder arising from the ciliary body or the iris.[5-9] Oncogenic mutations in *GNAQ* or its paralog *GNA11* are observed in up to 90% of UMs, but are rare in cutaneous melanoma (CM).[4,10] Such mutations reduce GTPase activity of the heterotrimeric G protein alpha subunit, resulting in constitutive activation that drives cellular proliferation through downstream signaling pathways, including MAPK, YAP/TAZ, and PI3K signaling.[5,11-13]

UM and CM display distinct metastatic progression. Up to 50% of UM tumors metastasize to the liver via hematogenous spread,[3,14,15] and survival after metastasis is generally poor [16] There remain few FDA approved options for patients with metastatic UM. Tebentafusp was the first systemic therapy shown to improve overall survival in metastatic UM at both 1 year and 3 years; however, this is only approved in HLA-A02:01-positive patients and, despite being an improvement, progression free survival at 6 months was only 31%, and median overall survival at 3 years was only 21.6 months in treated patients [17,18] Chemotherapy and immune checkpoint inhibitors, which are highly effective in CM, have very limited efficacy in UM.[19,20] Some promising targeted therapies are currently under investigation, including Darovasertib, a PKC inhibitor targeting GNAQ/GNA11-driven MAPK signaling,[21] and more invasive options, such as melphalan-based liver treatment, have demonstrated some benefit in the absence of other effective systemic options.[22]

While *GNAQ/11* oncogenic mutations are uncommon in CM, the majority of CMs harbor activating mutations in *BRAF* or *NRAS*.[5,23-25] This distinction suggests that unique properties of

melanocytes in different tissues, or perhaps their tumor microenvironments (TME), may drive

differential oncogene sensitivity. Melanocytes in the choroid interact with fibroblasts and

vasculature endothelial cells, whereas melanocytes in the skin are in direct contact with

keratinocytes. In transgenic experiments, BRAF[V600E]-positive melanocytes exhibit significantly

greater survival in the epidermis compared to GNAQ[Q209L]-positive melanocytes. However,

GNAQ[Q209L]-positive melanocytes preferentially survived in the dermis, where fibroblasts are

abundant.[26] This suggests that oncogene-specific interactions with the microenvironment

influence melanocyte survival and provide a basis for the absence of *GNAQ* mutations in CM,

except for dermal blue nevi which interact closely with fibroblasts.[10,26] Interestingly, while UM

and CM are unique in their metastatic patterns, GNAQ/11-mutant non-uveal melanoma

demonstrates a lymphatic metastatic progression that resembles CM, thus further indicating the

importance of studying UM in the correct anatomical context.[24]

Zebrafish are a well-established model for studying melanocytes and melanoma due to

conserved melanocyte differentiation pathways, efficient genetic manipulation, and the use of

transparent lines that facilitate high-resolution live-cell imaging.[24,27-35] Zebrafish have three types

of pigment cells; dark melanocytes, iridescent iridophores, and yellow xanthophores, that arise

from multipotent, neural crest-derived pigment progenitors.[24,33,36-39] Melanomas have been

shown to upregulate neural crest transcriptional programs, which are associated with worse

clinical outcomes.[29,40-42] Similarly, *Tg(mitfa:GNAQ[Q209L]); mitfa[-/-]* zebrafish, which retain

melanocyte progenitor cells but lack differentiated melanocytes,[43] have earlier occurrences of

UM than wild-type zebrafish.[34] By contrast, BRAF-driven CM displays a reliance on Mitfa activity

to progress.[44,45] To study the early events of UM, we developed a novel zebrafish model that

utilizes the transgene electroporation of adult zebrafish (TEAZ) approach to express the human

GNAQ[Q209L] oncogene in cells of the melanocyte lineage within the skin and eye. By targeting

injections to the choroid, primary UM tumors arise from the same anatomical location as human

UM, recapitulating similar TME and initiating cell types. In this study, we analyze expression

differences between genetically identical skin and eye-derived GNAQ[Q209L] tumors and

demonstrate the existence of a melanocyte precursor population in the eyes zebrafish that have

an increased sensitivity to transformation by GNAQ[Q209L].

Results

Generation of plasmid constructs to drive GNAQ[Q209L]-induced melanoma.

To study GNAQ-driven melanoma in adult zebrafish, we generated expression plasmids

containing two cassettes under the control of the *mitfa* promoter. The first cassette encoded

human oncogenic *GNAQ[Q209L]* fused to a T2A self-cleaving peptide followed by eGFP, while the

second cassette expressed CRISPR/Cas9. Additionally, three zebrafish U6 promoters were

used to drive gRNAs targeting *tp53* and *ptena/b* (**Figure 1A** – strategy A). Recognizing that T2A

cleavage results in residual amino acids remaining on the C-terminus of GNAQ, which may alter

its oncogenic activity, we employed an alternative delivery strategy. This system utilized five

separate plasmids: (1) *mitfa:GNAQ[Q209L]; mitfa:eGFP* (a generous gift from Dr. Jacqueline Lees

at the KOCH institute Massachusetts Institute of Technology), (2) *mitfa:Cas9*, (3) *U6:*gRNA-

tp53, (4) *U6:*gRNA-*ptena* and (5) *U6:*gRNA-*ptenb* (**Figure 1A** – strategy B) (plasmids 2-5 were a

generous gift from Dr. Richard White at the University of Oxford, UK). This design aimed to

improve the oncogenicity of GNAQ[Q209L] by avoiding unintended C-terminal modifications by T2A

cleavage. Moreover, the plasmids used in strategy B are smaller in size, which may enhance

the efficiency of uptake during electroporation.

We applied TEAZ to the skin (TEAZ-Skin) of adult wild-type (*AB*), *mitfa[w2/w2]* (*nacre*

[43]), and *mitfa[w2/w2]; mpv17[a9/a9]* (*casper[46,47]*) zebrafish to validate the oncogenic potential of

GNAQ[Q209L] (**Figure 1B**). TEAZ-Skin using Strategy A and Strategy B induced melanoma in wild-

type zebrafish within an average of 147- and 104-days, respectively (**Figure 1C-D, Figure**

S1A). TEAZ-Skin using plasmids to express oncogenic *GNAQ* alone or Cas9 with gRNAs to

delete *tp53, ptena* or *ptenb* alone did not induce melanoma, suggesting that both oncogenic

GNAQ and tumor suppressor deletion are required for transformation (**Figure S1A,** black line,

n=15). TEAZ-Skin was subsequently applied to zebrafish lacking functional *mitfa* (*nacre* and

casper), which lack differentiated melanocytes. Consistent with previous observations in

transgenic *Tg(mitfa-GNAQ^{Q209L}); mitfa$^{w2/w2}$; tp53$^{-/-}$* zebrafish,[34] *mitfa* loss resulted in decreased

tumor latency compared to wild-type zebrafish (**Figure 1C-D, Figure S1B,B'-C**). Tumors were

detected within an average of 30- and 28-days post-electroporation following Strategy B in

nacre and *casper*, respectively. Moreover, unlike wild-type zebrafish and consistent with

previous reports,[40] expression of oncogenic GNAQ alone was sufficient to induce melanoma in

nacre (**Figure 1D, Figure S1B**) and *casper* zebrafish (**Figure 1D, Figure S1C-D**). We ruled out

xanthomas as the cell of origin for TEAZ-Skin tumors, given the distinct visual differences

between the tumors and the xanthomas observed in *Tg(mitfa$^{w2/w2}$, mitfa:BRAF^{V600E}, tp53$^{-/-}$)*

zebrafish. The xanthomas were bright orange, opaque, and clearly distinguishable from the

darkly pigmented UM tumors in wild-type zebrafish and the beige, translucent UM tumors in

casper zebrafish (**Figure S1E**). CRISPR editing in TEAZ-Skin tumors was confirmed by Sanger

sequencing. All but one tumor (10 of 11) exhibited edits in all three targeted tumor suppressors

(*tp53*, *ptena*, and *ptenb*), with the remaining tumor lacking *ptena* mutation. As expected,

uninjected tissue showed no evidence of CRISPR editing (**Figure S2**).

Together, our findings demonstrate that both TEAZ-Skin strategies are sufficient to induce

GNAQ-driven melanoma, with strategy B demonstrating reduced tumor latency compared to

strategy A. Notably, TEAZ-Skin accelerates melanoma onset relative to transgenic models;[34]

while tumors in transgenic *Tg(mitfa:GNAQ^{Q209L}); tp53$^{-/-}$* zebrafish typically arise 200-400 days

post-fertilization,[34] TEAZ-Skin induces UM tumors by approximately 100 days in wild-type

zebrafish.

The *mitfa* promoter is active in choroidal melanocytes

Having confirmed the oncogenicity of our $GNAQ^{Q209L}$ expression plasmids using TEAZ-Skin, we next sought to adapt this approach to induce melanocyte transformation in the zebrafish eye (TEAZ-Eye), specifically targeting the choroid, where the majority of ocular melanocytes reside (**Figure 1E**). To determine whether TEAZ-Eye could serve as a model for UM in adult zebrafish, we first assessed the specificity of the *mitfa* promoter in driving transgene expression in choroidal melanocytes. To this end, we harvested eyes from transgenic *Tg(mitfa:GFP)* zebrafish and performed immunofluorescent imaging using an anti-GFP antibody. Choroidal melanocytes exhibited strong anti-GFP positivity compared to the secondary-only control (**Figure 1F**). Of note, the retinal pigment epithelium (RPE), located apically to the choroidal melanocytes, contains autofluorescent granules. Quenching the autofluorescence signal from the RPE improved the anti-GFP signal-to-noise ratio (**Figure S3A-B**). We next applied TEAZ-Eye by injecting and electroporating plasmids containing the *mitfa* promoter driving *eGFP* expression (*mitfa:eGFP*) into the choroid. Eyes were harvested two weeks post-electroporation, and immunofluorescence using an anti-GFP antibody was performed. Remarkably, cells at the injection site exhibited a strong eGFP signal compared to the secondary-only control (**Figure S3C**). These results demonstrate that the *mitfa* promoter is sufficient to drive transgene expression in choroidal melanocytes using TEAZ-Eye.

TEAZ-Eye induces UM in the choroid of wild-type and *casper* zebrafish

A major limitation in the study of UM onset and progression is the lack of anatomically accurate *in vivo* models. For instance, UM tumors derived in transgenic *Tg(GNAQ^{Q209L}): tp53^{-/-}* zebrafish predominantly form in the skin, with ~1-2% of tumors forming in the eye.[34] To address this gap, we applied TEAZ-Eye using a plasmid pool containing *mitfa:GNAQ^{Q209L}, mitfa:Cas9, U6:*gRNA-

*tp53, U6:*gRNA-*ptena, and U6:*gRNA-*ptenb* (i.e., Strategy B) into the choroidal space of wild-

type zebrafish, followed by electroporation (**Figure 1G-H**). Remarkably, TEAZ-Eye in wild-type

zebrafish resulted in the transformation of cells within the choroid. Live *in vivo* imaging of TEAZ-

Eye enabled the identification of single GFP+ cells at the injection site, serving as a proxy for

GNAQ[Q209L] expression (**Figure 1H-H'**). We next applied TEAZ-Eye to *mitfa*-deficient *casper*

zebrafish (**Figure 1I-I'**), and like with TEAZ-Skin, tumor latency of TEAZ-Eye injected *casper*

zebrafish (n=10) was significantly reduced compared to wild-type (n=5) (**Figure 1J**). H&E

analysis confirmed that TEAZ-Eye induced UM within the choroid, progressing into surrounding

ocular structures (**Figure 1K**). Notably, 100% (n=10) of tumors formed between the sclera and

RPE, highlighting the model's ability to induce anatomically accurate UM. Finally,

immunofluorescent analysis using anti-GFP antibodies confirmed the expression of plasmid

constructs following TEAZ-Eye (**Figure 1K'**).

UM transformation involves activation of neural crest transcriptional programs

During embryonic development, choroidal melanocytes are derived from the cranial neural crest

(CNC)[48]. Since our model can detect the earliest events in UM transformation, we asked

whether the reactivation of CNC programs is one of the initial steps in UM development. To

achieve this goal, tumors were derived in wild-type zebrafish using TEAZ-Eye with Strategy B

plasmids. At 50-90 dpe, when gross nodular tumors were identified, surgical enucleation was

performed, and dissociated cells from the entire eye were subjected to single-cell RNA

sequencing (scRNA-seq). We compared pooled wild-type tumors (2 sequencing replicates) with

tumor paired bilateral eyes or sibling-matched control eyes (2 and 4 pooled eyes, respectively, 2

sequencing replicates). Given that whole eyes were used as control samples and melanocytes

are most abundant in the choroid, melanocytes identified under control conditions are hereafter

referred to as choroidal melanocytes. Tumor cells were identified by expression of *GFP* and

oncogenic *GNAQ* (**Figure 2B**) whereas primary choroidal melanocytes were marked by

expression of *mlana* (**Figure S4**). Annotation of additional cell types within the TME was

accomplished by comparing to previously published scRNA-seq datasets using human choroid

and retina,[49,50] the Zebrafish Information Network (ZFIN),[51] and from the zebrafish embryo single

cell atlas "Daniocell"[52] (**Figure 2A**). Interestingly, tumor cells aligned most closely with choroidal

melanocytes, ruling out the retinal pigmented epithelium (RPE) as an origin of our TEAZ-Eye

induced tumors (**Figure 2C**). Additionally, profiling *GFP/GNAQ^{Q209L}+* UM cells revealed that

tumors exhibited strong expression of neural crest markers (*tfap2a, tfap2c, pax7b, sox10, foxd3*)

and melanocytic markers (*mitfa, tfap2e, mlana, pmela, slc22a7a, mlpha, mtbl*) (**Figure 2D**).

Unexpectedly, the CNC marker genes *twist1* or *dlx1*[53,54] were not enriched in UM tumors

compared to the TME. Pathway analysis showed that the genes enriched in TEAZ-Eye-derived

tumors were strongly associated with MITF-M-dependent gene expression, RAF/MAP kinase,

mTOR, PI3K/AKT signaling, NF-kB, RHO GTPase, and NOTCH activity, as well as fatty acid

oxidation pathways (**Figure S6A-B**).

Re-clustering of choroidal cells from control and tumor conditions revealed a marked expansion

of tumor cells relative to melanocytes which was accompanied by increases in immune cell

populations and fibroblasts (**Figure 2E-F**). Immune cell populations within the TME included B-

cells (*igic1s1, pax5, cd79a, zgc:153659*), T-cells (*traf1, si:ch211-67e16.3, cd27, zap70, cd8a,*

sla2), a population of cytokine producing cells, likely representing T-cells (*il4, il13, il11, ca2, il34,*

il6) as well as macrophages (*marco, ccl34a.4, mrc1b, c1qa/b/c, grn1*) and neutrophils (*cpa5,*

lect2l, mmp9, npsn, scpp8). Additional cell types within the choroidal TME included fibroblasts

(*col1a1a, col1a1b, col1a2, col5a1, dcn, serpinf1*), pericytes (*pdgfrb, notch3, foxf2b, cd248a*),

Schwann cells (*dlx5a, apoda.1, mpz, plp1b, cd59, s100b, mbap*) and vasculature endothelial

cells (*vwf, flt4, epas1b, ecscr, fabp11a*) (**Figure 2D-F**, **Table S1**).

We next compared choroidal melanocytes and UM tumors, revealing a significant overlap in

gene expression (n=644; hypergeometric p < 0.0001) alongside a large set of tumor-enriched

genes (n=1177) (**Figure 2G**). Shared genes included melanocytic regulators (*ednrb, sox10,*

foxd3, mitfa, mlpha, kita, mlana, pax7b), supporting a melanocytic origin, and many of these

transcripts were further upregulated in UM tumors compared to choroidal melanocytes (**Figure**

2H). HOMER promoter motif enrichment analysis of genes enriched in UM tumors revealed

binding motifs for the MiT/TFE family of transcription factors, as well as PAX7 and HOXB13,

suggesting that these regulators may contribute to UM-specific transcriptional programs (**Figure**

2G') Differential expression analysis further identified several genes enriched in UM tumors and

associated with patient survival, including *lgals2a, fabp3, gch2*, and the proto-oncogene *kita*

(**Figure 3H**). High expression of *FABP3, FABP5, KIT, GCH,* and *LGALS2* correlated with worse

overall survival (OS) and disease-free survival (DFS) in UM (**Figure 3I-K, Figure S7**), and these

genes were preferentially associated with TCGA molecular subtypes 3 and 4, representing the

more aggressive forms of UM (**Figure S8, Table S2**).[55]

Cancer associated fibroblasts within the uveal TME upregulate components of the

extracellular matrix (ECM)

Recent studies have significantly advanced our understanding of the TME as an active driver of

melanoma progression.[56,57] While it is well established that components of the TME, particularly

cancer-associated fibroblasts (CAFs), play a key role in malignancy,[58-60] *in vivo* models to study

such interactions in UM remain limited. Our model allows for direct comparison of the TME in

TEAZ-Eye and control injected eyes within the same animals. We compared the transcriptome

of CAFs and normal fibroblasts in wild-type zebrafish. Upon reclustering and differential gene

expression analysis of normal eye fibroblasts and CAFs, we found that several ECM proteins

were upregulated, indicating remodeling of the ECM in the presence of UM. Enriched genes

broadly fell into four categories: ECM, ECM regulators, TNFs, and collagens. Notably,

fibronectin (*fn1a*) was among the most significantly upregulated gene in CAFs compared to

normal fibroblasts (q-value < 0.0001). Several collagens were also found to have higher

expression in CAFs, including Collagen I (*col1a1b, col1a2),* Collagen V (*col5a1, col5a2a,*

col5a3a) and Collagen XII (*col12a1a)* (**Figure 2L, Table S3**).

To control for injection-related fibrosis potentially causing fibroblast activation and upregulation

of ECM components, we mock-injected three wild-type eyes with plasmids containing *mitfa*:GFP

and *mitfa*:Cas9 cassettes followed by electroporation (mock-TEAZ) and performed scRNA-seq.

Fibroblasts from mock-injected sibling control eyes (three pooled eyes), non-injected sibling

control eyes (four pooled eyes), and tumor-matched control non-injected eyes (three pooled

eyes) were equally distributed on the UMAP, with no alteration in *fn1a* or *lgals2a* expression or

other changes to extracellular matrix (ECM) genes upon mock-TEAZ injection (**Figure S5,**

Figure S9). These findings suggest that TEAZ-Eye tumor induction, and not the physical

injection, in zebrafish triggers a CAF response that may be critical to melanoma progression.

Skin and eye melanocytes, and their resulting tumors, have unique gene expression

profiles

With the ability to induce genetically identical tumors in both the eye and the skin of adult

zebrafish, we next asked whether tumors arising in these sites were transcriptionally distinct.

Any such differences could be due to the relationship between the unique microenvironments of

the skin and eye with their respective tumors or they could arise from inherent differences in the

untransformed melanocytic cells that reside in the skin or eye. To this end, we generated tumors

using TEAZ-Eye and TEAZ-Skin with strategy B plasmids in wild-type zebrafish and compared

them with control eye and skin tissue, respectively, using scRNA-seq (**Figure S5**). UMAP

analysis revealed that while the tumors and the majority of microenvironment clusters overlap,

the eye tissue expectedly had unique cell types corresponding to retinal bipolar cells, glial cells,

photoreceptors, and retinal interneurons (**Figure 3A**). To identify transcriptional differences

between TEAZ-Eye and TEAZ-Skin tumors, we compared the gene expression profiles of each

tumor type relative to the combined TME from both sites. A similar analysis was performed for

eye and skin melanocytes relative to the combined uninjected microenvironment (**Figure 3B,**

Table S4). Notably, the strongest overlap was observed between genes upregulated in skin

tumors versus TME and those upregulated in eye tumors versus TME, with 41% of the 2,192

unique differentially expressed genes (DEGs) shared, highlighting the broad transcriptional

programs activated by $GNAQ^{Q209L}$ expression, $tp53$ loss, and $ptena/b$ loss (**Figure 3C**). We

hypothesized that the DEGs shared by only the two tumors and not the primary melanocytes

would represent mechanisms that are essential for UM growth and survival. In line with this

prediction, IPA analysis of this gene subset revealed that pathways pertaining to cancer

proliferation and metabolism were enriched (**Figure S10A**). Furthermore, we noticed an

enrichment of WNT signaling, PTEN regulation, and RAF/MAPK pathways which have been

previously implicated as important signaling cascades in UM.[61-69] Interestingly, only 5% of DEGs

overlapped across all datasets (eye and skin melanocytes and tumors), and we predicted these

genes to be associated with melanocyte identity and survival (**Figure 3C**). IPA analysis

confirmed this prediction with the most enriched pathway being MITF-M dependent gene

expression. Other enriched terms were signaling events that are downstream of GPCRs, such

as GNAQ, and pathways that are essential for survival such as nucleotide biosynthesis and

amino acid regulation of mTORC1 (**Figure S10B**). Notably, only 25.8% of DEGs overlapped

between eye and skin melanocytes, indicating substantial transcriptional differences in

melanocytes originating from distinct anatomical regions (**Figure 3C**). We next asked whether

transcriptional differences in primary melanocytes would be reflected in the corresponding

tumors, that is, whether eye melanocytes would be more similar to eye tumors than skin

melanocytes. Interestingly, the overlap of unique DEGs between eye melanocytes and eye

tumors relative to their respective microenvironments (386/1823) was not significantly different

than the overlap between skin melanocytes and eye tumors (489/1689) (permutation test, p-

value = 1, 10,000 permutations) (**Figure 3C**). In contrast, the overlap of unique DEGs between

skin melanocytes and skin tumors (552/2350) was significantly greater than the overlap

between eye melanocytes and skin tumors (424/2509) (permutation test, p-value = 0, 10,000

permutations).

While this analysis allowed for consideration of genes that were differentially expressed in the

same direction and magnitude versus the microenvironment, it would miss the genes that are

differentially regulated between the two tissue types directly. Therefore, we reclustered just the

normal melanocytes from both eye and skin for direct comparison and found that skin and eye

melanocytes clustered separately on the UMAP (**Figure 3D**). Our analysis returned 186 DEGs

enriched in eye melanocytes (log2FC > |0.5|, q-value < 0.01) and 229 DEGs enriched in skin

melanocytes (log2FC > |0.5|, q-value < 0.01) with 3387 genes not differentially regulated (q-

value > 0.01, expressed in at least 10% of cells), displaying modest transcriptional variance

between melanocytes of the skin and the eye (**Figure 3E, Table S5**). GSEA analysis of these

DEGs between eye and skin melanocytes revealed that skin melanocytes were enriched for

DNA repair pathways (p-value < 0.002, FDR < 0.016) and for genes that are upregulated during

UV response (p-value < 0.009, FDR < 0.081) (**Figure 3G**). On the other hand, DEGs in eye

melanocytes showed an enrichment for the genes involved in downregulating the UV response

pathway (p-value < 0.000, FDR < 0.090) and for the gene ontology oxidative phosphorylation

pathway (p-value < 0.000, FDR < 0.006). We also reclustered and directly compared skin and

eye tumors, with a similar tissue-specific grouping observed (**Figure 3D**). 230 DEGs were

enriched in the eye tumors (log2FC > 0.5, q-value < 0.01) and 375 DEGs were enriched in the

skin tumors (log2FC > 0.5, q-value < 0.01), while 5,754 genes remained not differentially

regulated between the two tumor types (q-value > 0.01, expressed in at least 10% of cells)

(**Figure 3F**). Interestingly, there was an enrichment of the epithelial to mesenchymal transition

dataset (p-value < 0.000, FDR < 0.020) in the eye tumor as well as an enrichment for the

metabolism of selenoamino acids (p-value < 0.000, FDR < 0.000) (**Figure 3H**). The skin tumor

was instead enriched for PI3K-AKT-MTOR signaling (p-value < 0.009, FDR < 0.086) and

pathways for RHO GTPases activating formins (p-value < 0.002, FDR < 0.058). Therefore, after

both types of analyses we conclude that both the primary melanocytes and the GNAQ-driven

melanomas are significantly different between the eye and the skin emphasizing the importance

of studying UM, and melanocytes in general, in their proper anatomical context.

Melanocyte progenitor cells are highly susceptible to oncogenic GNAQ transformation in

***mitfa*-deficient zebrafish.**

H&E analysis of *nacre* and *casper* zebrafish eyes showed loss of choroidal melanocytes

(**Figure 4A**), whereas melanin in the RPE remains less affected, as previously demonstrated.[33]

These observations imply that an *mitfa*-independent melanocyte progenitor cell persists in *nacre*

and *casper* eyes and that they are susceptible to oncogenic GNAQ transformation. To identify

such cells, we crossed *nacre* (*mitfa$^{w2/w2}$*) with transgenic *Tg(mitfa:GFP)* lines to create

Tg(mitfa:GFP); mitfa$^{w2/w2}$ zebrafish. We then harvested adult eyes at 5 months and performed

anti-GFP immunofluorescence analysis, with the assumption that loss of *mitfa* does not alter

expression of the *mitfa* promoter. While GFP+ melanocytes were readily detected in the choroid

of *Tg(mitfa:GFP)* zebrafish (**Figure 1F, Figure S3**), staining of *Tg(mitfa:GFP); mitfa$^{w2/w2}$* eyes

revealed rare cells with GFP-positivity (**Figure 4B**). Such GFP+ cells were primarily located in

the ciliary body, with isolated cells also found in the choroid, suggesting the persistence of a

mitfa-independent progenitor population in adult *mitfa*-deficient zebrafish (**Figure 4B**). We next

asked whether such cells retain the ability to undergo melanocyte differentiation by using TEAZ-

Eye to deliver plasmids containing the *mitfa*-promoter driving expression of *mitfa* (i.e., to

overexpress *mitfa* mRNA). Interestingly, overexpression of *mitfa* transcript levels resulted in

expansion of choroidal melanocytes in wild-type animals and the rescue of choroidal

melanocytes in *mitfa*-deficient (*nacre*) zebrafish as shown by H&E analysis (**Figure S11A-B'**).

To assess whether GFP-positive cells are sensitive to oncogenic transformation by GNAQ, we

expressed *mitfa:GNAQ^{Q209L}* in *Tg(mitfa:GFP); mitfa$^{w2/w2}$* zebrafish using TEAZ-Eye. In all

injected fish (n=3), UM tumors were GFP-positive, suggesting they originated from GFP+

progenitor cells within the eye of *Tg(mitfa:GFP); mitfa$^{w2/w2}$* zebrafish (**Figure S11C**).

Melanocyte progenitor cells express neural crest marker genes in *mitfa*-deficient

zebrafish.

We performed scRNA-seq and pseudotime analysis on 3 pooled UM tumors from the eyes of

casper zebrafish and compared them to the pooled tissue from 3 normal *casper* eyes (**Figure**

4C, Figure S12). Within the *casper* eyes, we identified a population of cells expressing the

nacre allele that clustered closely with UM tumors under TEAZ-Eye conditions, termed

progenitors (**Figure S13A-B and Tables S6-8**)**.** We re-clustered the progenitor and tumor cells

(**Figure 4D**), revealing seven distinct clusters consisting of neural crest-like (NC-like),

melanoblast-like, proliferating, progenitor, angiogenic, stress-like, as well as an unidentified

cluster marked by *tmsb1, krt91* and *vim* (**Figure 4D, Table S9**). Unsupervised pseudotime

analysis supports a lineage trajectory from the progenitor cells to heterogeneous UM tumor cells

(**Figure S13C**). Notably, the progenitor cell population in normal *casper* eyes expressed neural

crest and melanocyte stem cell marker genes, including *tfap2a, foxd3, sox10*, *pax3a, vim,* and

yap1 (**Figure 4E-F**). UM tumor cells retained expression of these genes but activated additional

genes associated with pigment progenitor cell function, including *tfec* and *kita* (**Figure 4F**).

Notably, *mitfa* expression in tumors and progenitor cells in *casper* zebrafish reflects the

expression of the mutant *mitfa$^{w2/w2}$* allele, which functions as a lineage tracer to mark cells that

would normally express *mitfa* under wild-type conditions (**Figure S13B**). Interestingly, despite

the lack of Mitfa activity in *casper* zebrafish, GNAQ-positive UM tumors included a melanoblast-

like cluster that exhibited expression of pigmentation genes, including *pmela, mlpha*, and *mlana,*

likely explaining the occasional emergence of melanin within tumors as they progress in *casper*

zebrafish.

***mitfa*-deficient zebrafish exhibit expanded *pax3a* and *tfec* positive melanocyte**

progenitors during embryogenesis.

Having established that an Mitfa-independent progenitor population gives rise to UM in *mitfa-*

deficient zebrafish, we next examined whether this population is expanded relative to wild-type

zebrafish. To test this prediction, we performed scRNA-seq on GFP-positive cells sorted from

Tg(mitfa:GFP);mitfa$^{w2/w2}$ and *mitfa$^{+/w2}$* sibling-matched transgenic zebrafish embryos at 28 hours

post fertilization (hpf), when melanocytes (zebrafish melanophores) begin to differentiate

(**Figure 4G, Figure S12**). We assigned cell-type annotations based on our previously annotated

GFP-positive cells from *Tg(mitfa:GFP)* embryos.[30] The 10 clusters included six main cell types:

neural crest cells (*sox10, foxd3*), a tripotent precursor of melanoblasts (M), iridoblasts (I), and

xanthoblasts (X) (*tfap2a*, *cdkn1ca, slc15a2, ino80e, id3, mycn, tfec*), termed MIX cells, a cluster

that expressed high levels of melanoblast/xanthoblast markers (*mitfa, erbb3b, impdh1b, gch2,*

id3), termed MX cells, a melanoblast cluster (*mitfa, dct, pmel, tyr*), as well as two additional

clusters corresponding to xanthoblasts and xanthophores (**Table S10**). For this analysis we

referred to MIX and MX cells as melanophore progenitors (**Figure 4G**). As expected,

melanophore cells were reduced in *Tg(mitfa:GFP): mitfa$^{w2/w2}$* zebrafish (**Figure 4G-G'**).

However, melanophore progenitors were expanded in *Tg(mitfa:GFP); mitfa$^{w2/w2}$*, while neural

crest cells remained unaffected (**Figure 4G-G'**). Melanophore markers were downregulated in

GFP-positive cells sorted from *Tg(mitfa:GFP); mitfa$^{w2/w2}$,* whereas melanocyte progenitor

markers (*pax3a, sox10, foxd3*), as we identified in adult zebrafish eyes, were upregulated in

embryonic progenitors from *Tg(mitfa:GFP); mitfa$^{w2/w2}$* embryos (**Figure 4H, Table S11**). A role

for *pax3a* in adult fish eye tissue had not been previously characterized; however, recent studies

have identified PAX3 expression in ocular surface melanocytes,[70] and another study reports an

association between PAX3 and stem cell markers in UM progression.[71] Consistent with these

observations, we identified a *pax3* expressing cell population in adult C57BL/6 mouse eyes in

the limbal region, and rarely in the choroid (**Figure S14**). These findings support a rare *pax3-*

expressing population in both zebrafish and mouse eyes and may mirror *pax3* expression in the

melanocyte stem cell in the skin.[72,73]

Conditional deletion of *mitfa* in adult zebrafish fails to recapitulate germline loss-of-

function effects on tumor-free survival in GNAQ-driven melanoma.

We next investigated whether the accelerated growth of GNAQ-driven tumors in *mitfa*-deficient

zebrafish, compared to wild-type animals, was due to an expansion of progenitor cells or the

cell-intrinsic loss of Mitfa activity. To this end, we performed conditional *mitfa* knockout

experiments using Strategy B plasmids in combination with either (1) non-targeting control

gRNAs or (2) gRNAs targeting exon 5 of *mitfa*. TEAZ-Skin was performed in three experimental

groups: (1) *casper* zebrafish (n=9) with gRNAs targeting *mitfa* (which controls for off-target

effects of the gRNA and for potential function of the *mitfa$^{w2/w2}$* allele), (2) wild-type zebrafish

(n=9) with non-targeting (NT) gRNAs, and (3) wild-type zebrafish (n=9) with *U6*:gRNA-*mitfa*.

GFP expression was confirmed in all injected zebrafish (**Table S12**). Within 100 days post-

electroporation (dpe), tumors developed in 9/9 casper, 2/9 wild-type (+U6:gRNA-mitfa), and 2/9

wild-type (+U6:gRNA-NT) zebrafish (**Figure 4I-J**). Of note, *mitfa*-deficient tumors that formed in

wild-type or *casper* zebrafish developed distal metastases, an observation not seen in *mitfa*

wild-type TEAZ-Skin injected zebrafish (**Figure 4J, Figure S15**). Our findings support a model

in which failed differentiation as well as the expansion of melanocyte progenitor cells in *mitfa-*

deficient animals creates a permissive cellular environment for GNAQ-induced transformation

and reduced tumor latency.

Transcriptional programs enriched in BRAF- versus GNAQ-driven melanoma overlap with

specific melanocyte progenitor populations.

Oncogenic BRAF and NRAS are unable to promote transformation in the absence of *mitfa,*[29,32]

suggesting that distinct transcriptional programs, potentially reflecting unique cellular origins, are

required for UM and CM transformation. To address this hypothesis, we analyzed previously

published scRNA-seq datasets in transgenic *Tg(mitfa:BRAF^{V600E}): tp53$^{-/-}$: mitfa$^{w2/w2}$* zebrafish

where TEAZ-Skin was used to rescue *mitfa* and induce melanoma,[32] to our GNAQ-induced

tumors by TEAZ-Skin (**Figure 5A)**. We predicted that by analyzing primary tumors, the

transcriptional signature of the origin cell will be retained and can be identified by overlapping

gene signatures identified in the embryonic melanocyte lineage.[30] Interestingly, while the

expression of *mitfa* was significantly higher in GNAQ-positive tumors, the expression of

pigmentation genes; including *tyrp1b, dct, scl45a2, sox10* and *pmela* were significantly higher in

BRAF-positive melanoma (**Figure S16**). Genes enriched in GNAQ-UM and BRAF-CM were

identified by those with a Log2FC > 1 and Adj p-value < 0.0001 between the two tumor types

(**Figure 5B, Table S13**). We next overlapped this differential gene signature with genes

expressed by cells within the melanocyte lineage isolated from zebrafish embryos (**Figure 5C**).

As expected, BRAF-driven tumors strongly mapped to the melanoblast and melanophore

clusters (**Figure 5D**), supporting recent findings that BRAF-driven melanoma originates from

melanoblasts, a melanocyte progenitor population[32,74]. When comparing GNAQ-enriched genes

to the melanocyte lineage, we found that the signatures were not confined to a single cluster but

were distributed among neural crest cells, melanocyte progenitors, and xanthophore clusters

(**Figure 5D; Figure S17**). Such results support the hypothesis that GNAQ- and BRAF-driven

melanomas depend on distinct transcriptional programs, reflecting less and more differentiated

melanocytic states, respectively.

We next asked whether less differentiated melanocytes, similar to the *mitfa*-deficient progenitors

in *casper* eyes, are also present in adult wild-type eyes. Since melanocyte stem cells are known

to reside in the zebrafish skin,[28] we included skin tissue as a control and reclustered

melanocytes from wild-type skin and eyes with progenitor cells from *casper* eyes. Interestingly,

a subset of wild-type melanocytes from both eye and skin samples clustered closely with *casper*

progenitors (**Figure 5E, Figure S18**) and expressed pigment cell progenitor marker genes,

including *pax3a, sox10, foxd3,* and *tfec* (**Figure 5F**). These findings suggest either plasticity

among melanocytes within the eye or the existence of an eye melanocyte stem cell population

in adult wild-type zebrafish. Moreover, such cells may be more susceptible to transformation by

GNAQ[Q209L] and could represent the cells of origin for UM tumors in this context.

Discussion

Herein, we present an immunocompetent zebrafish model of primary UM where tumors arise

within the choroid and progressively invade surrounding ocular structures. This model enables

the study of the earliest events of UM pathogenesis and how anatomical context impacts

disease progression. Notably, genes identified through our analysis correlate with survival

differences in human UM, highlighting the model's potential for mechanistic studies on gene

function in UM pathogenesis. Compared to transgenic models, our system produces

anatomically correct tumors at an accelerated rate, offering an efficient platform for studying UM

progression and therapeutic responses.

By generating UM tumors from the same anatomical location as the majority of human UM, we

were able to conduct key analyses such as the comparison of the tumor to untransformed

primary choroidal melanocytes and the tumor microenvironment. This method allowed for the

identification of genes that are upregulated in the tumor and therefore important for disease

onset and progression. Specifically, we found that fatty acid binding proteins, particularly *fabp3*,

are enriched in UM tumors. While lipid droplets have been identified as vulnerable targets in CM

and regulators of melanoma plasticity,[32,75] lipid signaling in UM has received limited attention.[76-

79] Interestingly, high expression of *FABP3* as well as *FABP5* associated with worse disease-free

survival in patients with UM but not in CM. Interestingly, our analysis also identified *LGALS2* and

GCH, two genes whose expression was strongly associated with worse overall survival in UM

but intriguingly showed the opposite in CM. Finally, we found greater *kita* expression in GNAQ-

driven tumors compared to choroidal melanocytes, a result that is particularly notable given

recent studies linking KIT-positive UM tumors with poor prognosis and monosomy of

chromosome 3.[80] Furthermore, expression of these genes was highest in TCGA molecular

classes 3 and 4 of the UM dataset, the subset of patients with the highest rates of metastatic

incidence and BAP1 loss.[55] Such observations show that this model can uncover genes that are

likely to have key, disease-specific roles in UM whose function could be further explored through

genetic perturbations in zebrafish. These genes could also be developed into new biomarkers of

disease risk to refine the identification of patients likely to progress to metastatic disease.

Although UM and CM both arise from cells within the melanocyte lineage, they differ

significantly in their biology and clinical outcomes. To explore how anatomical context influences

transcriptional states, we compared genetically identical tumors derived in the skin and eye and

found that eye-derived tumors display distinct expression profiles. To explore whether the

differences in tumors were due to inherent differences in their primary cells, we also compared

normal melanocytes from the eye and the skin. While we did find some transcriptional

distinctions between eye and skin melanocytes, these variances only represented a small

fraction of the differentially expressed genes between eye and skin melanomas. These results

suggest that the transcriptional profiles of genetically identical tumors in distinct anatomical

locations are influenced by more than the intrinsic differences in their origin cells, perhaps

varying due to signals from their unique microenvironments. Of the expression differences

between normal eye and skin melanocytes, GSEA pathway analysis revealed that skin

melanocytes are enriched for UV responses and DNA damage repair. Zebrafish, like humans,

have been shown to alter their melanin production in response to UV exposure.[81] Our results

suggest that skin melanocytes in the zebrafish may be more primed to enact UV-protective

functions than eye melanocytes, even with no UV exposure. Furthermore, the enrichment of

oxidative phosphorylation in eye melanocytes and of selenoamino acid metabolism in eye

tumors suggests metabolic differences between melanocytes derived from the eye or skin.

Interestingly, disruptions in selenoamino acid metabolism have been shown to adjust the TME

and growth of primary tumors in a number of cancers, including melanoma.[82] Pathway analysis

between skin and eye tumors has also revealed potential differences in the progression of these

tumors. Eye tumors were enriched for EMT genetic programs, suggesting that eye melanocytes

may exist in a more mesenchymal state than skin melanocytes or that greater expression of

EMT programs is necessary for metastatic spread out of the eye and towards distant sites like

the liver. On the other hand, skin tumors were enriched for the certain arms of the GNAQ

signaling cascade,[5,11-13] namely PI3K/AKT/MTOR signaling and RHO GTPases activating

formins. Further investigation is needed to confirm whether microenvironment differences or

anatomical origin of melanocytes can influence the relative importance of oncogenic pathways

in the setting of $GNAQ^{Q209L}$ expression. Using our animal model to compare genetically identical

tumors, and their primary cells, across different anatomical sites has revealed distinct biological

behaviors, highlighting the model's potential for further investigations into site-specific tumor

biology and therapeutic strategies.

An interesting distinction between GNAQ and BRAF oncogenes is their differential dependency

on Mitfa activity for tumorigenesis. While BRAF[V600E] requires Mitfa activity to induce

melanoma,[27,29] GNAQ[Q209L]-driven tumor latency was significantly reduced in *mitfa*-deficient

zebrafish compared to wild-type controls. Our results are consistent with prior studies showing

that *mitfa*-deficient zebrafish activate distinct signaling pathways, favoring YAP over MAPK

signaling, and give rise to tumors with reduced latency compared to wild type tumors. Moreover,

manipulating YAP activity in wild-type animals was shown to recapitulate the enhanced growth

seen in *mitfa*-deficient tumors.[40] We observed high *yap1* expression in melanocyte progenitor

cells of *mitfa*-deficient zebrafish, suggesting that elevated Yap1 activity may be carried over

from the progenitor cell during transformation. While germline *mitfa* loss accelerates tumor

onset, our data suggest this effect is not solely due to the deletion of Mitfa from differentiated

melanocytes. Instead, we propose that germline *mitfa* deficiency expands a population of

pax3a- and *tfec*-positive melanocyte progenitor cells, which are more susceptible to GNAQ-

induced transformation. Notably, conditional loss of *mitfa* in differentiated melanocytes may

activate progenitor-like transcriptional programs, increasing their vulnerability to oncogenic

transformation. Alternatively, *mitfa* loss may impair the differentiation of melanocyte progenitors,

preventing the formation of GNAQ-positive nevi and promoting direct progression to malignancy.

Interestingly, these progenitor cells appear to maintain a neural crest signature, as evidenced by

the relatively high expression of additional markers such as *foxd3, pax3a, sox10,* and *tfec*.

[30,31,83-85] Together, these findings suggest that germline loss of *mitfa* accelerates GNAQ-driven

transformation of melanocyte progenitor cells by promoting their expansion and maintaining a

neural crest-like transcriptional profile. Identifying the transcriptional and signaling mechanisms

within *mitfa*-independent melanocyte progenitor cells that facilitate UM onset and progression

will have significant clinical implications, particularly for the differential diagnosis of high-risk

lesions and the rational design of targeted therapies.

Studies in human UM have shown that less pigmented tumor regions are more metastatic and

that BAP1-negative tumors exhibit a stem-like phenotype.[1,2,86,87] Our findings raise intriguing

questions about whether the progenitor cells observed in *mitfa*-deficient zebrafish are similarly

predisposed to a more undifferentiated, stem-like phenotype, contributing to their enhanced

transformation potential and aggressive tumor behavior. Interestingly, the direct comparison of

melanocytes and tumors from the eye or skin against their microenvironments revealed that skin

melanocytes overlap with the skin tumor greater than the overlap of eye melanocytes with the

skin tumor. However, this was not true in reverse with eye melanocytes being no more similar to

the eye tumor than skin melanocytes are. These findings suggest that eye tumors may originate

from a less differentiated cell compared to the melanocytic cell type that CM arises from. This

progenitor cell may be underrepresented in the normal eye melanocyte population, leading to

the observed lower transcriptional overlap between total enriched genes in eye tumors and

normal eye melanocytes. Furthermore, re-clustering of normal WT melanocytes and melanocyte

progenitor cells from *mitfa*-deficient zebrafish demonstrates that both the eye and skin have

melanocyte populations that cluster closely with melanocyte progenitors. These melanocyte

populations may be less differentiated as evidenced by increased expression of *pax3a*, *tfec*, and

foxd3. Whether this subset of melanocytes represents bona fide melanocyte stem cells or

melanocytes that can oscillate between differentiation states within the eye remains to be

uncovered.

Furthermore, differences in *mitfa* dependency between BRAF-driven CM and GNAQ-driven UM

suggest fundamental distinctions in their cellular origins. Although MITF-independent

melanocytic cells are unlikely to be the cellular origin of CM in zebrafish models, studies have

shown that MITF-independent BRAF-positive melanoma cells are central to disease

recurrence[45]. Our results highlight that *mitfa* expression does not directly reflect Mitfa activity, as

mitfa-target genes are more highly expressed in BRAF-positive melanoma despite lower *mitfa*

expression compared to GNAQ-positive melanoma. GNAQ-driven transformation may be

facilitated by Mitfa paralogs such as Tfec and Tfeb. The Mitfa paralog Tfec has been shown to

activate pigmentation genes in both the retinal pigment epithelium and melanocyte progenitor

cells. Tfec can also rescue ectopic melanocytes in *nacre* zebrafish embryos.[31,33] Moreover, we

have recently shown that the MITF paralog TFE3 promotes cellular plasticity and stemness

genes in MITF-low melanoma cells,[88] suggesting that MITF paralogs are compelling candidate

transcription factors in UM pathogenesis.

Although understudied, the choroidal stroma is abundant in fibroblasts that produce various

ECM proteins, including collagen, elastin, laminin, fibrin and fibronectin.[89-92] This

microenvironment differs from that of the keratinocyte-rich epidermis where CM arises. The

differences in TME between UM and CM are likely to play an important role in the phenotypic

differences observed in these tumors. Recent studies show that increased fibronectin promotes

a less differentiated and more mobile phenotype in murine melanocytes *in vitro*.[93] Our finding

that fibronectin (*fn1a* and *fn1b)* was upregulated in CAFs compared to normal fibroblasts

provides evidence that fibronectin may promote invasiveness or dedifferentiation in a malignant,

in vivo context. Similar to other studies finding an accumulation of collagen I in the TME of

various cancer types, we found upregulation of *col1a1b* and *col1a2* in CAFs relative to normal

fibroblasts, indicating increased collagen I deposition.[59,60] The precise mechanism of how

different subclasses of CAFs regulate UM differentiation state and propensity to metastasize

through ECM remodeling is still to be defined.

Another explanation for decreased tumor latency in *casper* zebrafish is that the global loss of

mitfa could have secondary effects on other cell types, such as immune cells, which may

influence tumor behavior. Furthermore, the presence of xanthophore, iridophore, and

melanophore markers within UM tumor cells raises the possibility of a pluripotent progenitor

population. This model also enables us to examine whether differences in differentiation

potential contribute to tumor aggressiveness, as *casper* fish may exhibit more pronounced

tumor acceleration due to the simultaneous blockade of two chromatophore differentiation

pathways compared to single-pathway inhibition in *nacre* fish.

Our findings open numerous avenues for future research. This model can be used to study the

effects of *BAP1* loss and other secondary mutations commonly associated with UM, such as

EIF1AX and *SF3B1*. It also provides a platform for investigating metastatic events, as tumors

emerge in defined primary sites. Establishing zebrafish UM cell lines from different anatomical

locations and genetic backgrounds could offer further insights into the role of microenvironment

and *mitfa* dependency in UM progression. Additionally, characterizing the melanocyte progenitor

population in the eye and determining if a similar population exists in mammals could provide

critical insights into uveal stem cell biology. While we identified a population of Pax3-positive

cells in the mouse limbus, it remains to be determined if these cells are sensitive to oncogenic

GNAQ transformation.

Limitation of study

The use of *tp53* and *ptena/b* mutations in our UM model does not fully recapitulate the genetics

of human UM. Mutations in *TP53* and *PTEN* are rarely observed in UM, where loss-of-function

mutations in *BAP1* represent the most frequent tumor suppressor alteration.[67,94,95] However,

both *TP53* and *PTEN* signaling pathways have been shown to be altered in UM pathogenesis.

MDM2 overexpression, an upstream negative regulator of p53, and downregulation of *PERP*, a

downstream effector of p53, have been demonstrated in many UM patients and are associated

with worse outcome.[96-100] Studies have also shown that over half of UM tumors have decreased

PTEN immunostaining and its loss is correlated with decreased survival.[67-69] Nevertheless,

whether genetic knockout of these tumor suppressors recapitulate the dysregulation of these

pathways in human UM remains an open question. Another limitation to our approach when

examining the differences between genetically identical skin and eye-derived tumors is the

inability to distinguish between purely microenvironmental effects and differences in cell of origin

(melanocytes in the skin versus in the eye). To address this question, a syngeneic or

immunodeficient model could be applied to transplant cells from a primary UM tumor into both

anatomical sites. Additionally, our use of the *mitfa* promoter to drive oncogene expression

presents a challenge in defining the exact cell of origin, as this promoter is active in both

melanocyte progenitors and differentiated melanocytes. As such, it remains unclear whether

tumors in *mitfa* wild-type fish (or those with conditional *mitfa* deletion) arise from a progenitor or

a mature melanocyte. Lineage tracing studies will be essential to determine whether

differentiated melanocytes possess the capacity to initiate primary UM in *mitfa*-competent or

conditional *mitfa*-KO zebrafish.

Resource availability

Lead contact

Requests for further information and resources should be directed to, and will be fulfilled by, the

lead contact, Colin Kenny, PhD (colin-kenny@uiowa.edu).

Materials availability

Plasmid constructs and zebrafish generated in this study may be obtained by a request to the

lead contact.

Data and code availability

Datasets generated in this article are available from NCBI GEO accession number GEO:

GSE307227. This study does not report original code. Any additional information required to

reanalyze the data reported in this work paper is available from the lead contact upon request.

Acknowledgments

This project was supported by grants from the Holden Comprehensive Cancer Center (HCCC;

P30- CA086862) at the University of Iowa. Additional support came from an institutional

research grant from the American Cancer Society, administered through the HCCC, to C.K.

(IRG-18-164-43) and from the Melanoma Research Alliance (ID#1426690) to C.K. Additional

support came from the National Institutes of Health (NIH) to R.A.C. (R01-AR062457). JY and

DR were supported by predoctoral fellowships from the National Institutes of Health (T32

GM144636 and T90 DE023520, respectively). Additional support came from the Leo Foundation

(LF-OC-21-000888) to DL. The funders had no role in study design, data collection and

analysis, decision to publish, or preparation of the manuscript. We gratefully acknowledge Dr.

Richard White from Oxford University, UK. for providing mitfa-Cas9, *U6*:gRNA-*ptena*, *U6*:gRNA-

ptenb and *U6*:gRNA-*tp53* plasmids constructs and for providing critical comments to

successfully preform TEAZ experiments. We additional acknowledge Dr. Jacqueline Lees and

Dr. Adam Amsterdam from the Koch Institute for Integrative Cancer Research at MIT for

generously providing *mitfa*:$GNAQ^{Q209L}$; *mitfa*:GFP plasmid constructs to induce UM tumors.

Author contributions

Conceptualization: C.K. Methodology: J.I.Y., D.R., R.M., Z.L., K.R.C-H., D.G., J.D.R., S.P.G.M.,

D.F., I.O.T. Investigation/data analysis: J.I.Y., D.R., R.M., Z.L., K.R.C-H., D.G., J.D.R., A,J,D.,

S.P.G.M., D.F., I.O.T., C.K. Visualization and data interpretation: J.I.Y., D.R., R.M., Z.L., K.R.C-

H., D.G., S.P.G.M., D.F., I.O.T. D.L., R.J.W., and C.K. Funding acquisition: R.A.C., D.L., C.K.

Project administration: C.K. Supervision: C.K. Writing—original draft: J.I.Y., and C.K. Writing—

review and editing: J.I.Y., D.R., C.M., E.M.B., D.L., R.J.W., and C.K. Approval of submission:

J.I.Y., R.M., Z.L., K.R.C-H., D.R., D.G., S.P.G.M., C.M., D.F., J.D.R., A,J,D., I.O.T. R.A.C.,

E.M.B., D.L., R.J.W., and C.K. Accountable for contributions: J.I.Y., D.R., R.M., Z.L., K.R.C-H.,

D.G., C.M., S.P.G.M., D.F., J.D.R., A,J,D., I.O.T. R.A.C., E.M.B., D.L., R.J.W., and C.K.

Declaration of interests

The authors declare no competing interests.

Methods

Experimental model and study participant details

Zebrafish husbandry

All zebrafish used in this study were bred and maintained at the University of Iowa Animal Care

Facility. The zebrafish are kept consistently at 28°C, 7.4 pH, and controlled salt concentrations

and cyclically receive 14 h of light followed by 10 hours of darkness. All zebrafish experiments

were performed in compliance with the ethical regulations of the Institutional Animal Care and

Use Committee at the University of Iowa and in compliance with NIH guidelines (protocol

#3022523). Zebrafish embryos were maintained at 28.5°C and staged by hours or days post-

fertilization (hpf or dpf).

Zebrafish mutant lines

Wild-type zebrafish in this study include the *AB* line and the *WIK* line. Transgenic lines include

casper (*mitfa*$^{w2/w2}$; *mpv17*$^{-/-}$)[46,47], *nacre* (*mitfa*$^{w2/w2}$)[43], *Tg*(*mitfa*:GFP)[18], and *Tg*(*mitfa*:GFP;

mitfa$^{w2/w2}$. TEAZ-Eye and TEAZ-Skin was performed on adult zebrafish (greater than 4 months

postfertilization).

Cloning

The MiniCoopR[29] vector (MiniCoopR 2x*U6*:gRNA, *mitfa*:Cas9 was a gift from Leonard Zon

Addgene plasmid # 118844) was modified to replace the *mitfa* coding sequence with *GNAQ*^{Q209L}

by removing *mitfa* using StuI and SmaI restriction enzymes (NEB). Gibson assembly was used

to insert *GNAQ*^{Q209L}:T2A:GFP DNA as a gBlock hifi (IDT) with sequence on the 3'

(GAAGCTAACACATAGTTGAAC) and 5' (CAATGCCAACTAAATTTCATG) as overhangs.

Gibson assemblies were transformed using NEB 5-alpha competent *E. coli* cells via heat shock.

GNAQ^Q209L:T2A:GFP insertion was confirmed using Sanger sequencing (Primers:

CAAGGAAGCCCGGCGGATCAACGA, CATACTTGTATGGGATCTTGAGTGTGTCCA,

GTGGAGTCAGACAATGAGAACCGAATGGAG, CAAGCTGGAGTACAACTACAA).

mitfa sgRNA was designed using IDT's CRISPR-Cas9 guide RNA design tool (sequence:

ATGGACAAAGCTGGACCATG). *U6*:sgRNA-*ptena* was linearized by PCR (forward primer:

GTTTAAGAGCTATGCTGGAAACAGCAT, reverse primer: GAACAAAGAGCTGGAGGGAG).

Linear plasmid was purified from a 2% agarose gel using the Monarch GNA Gel Extraction Kit

(#T1020S). ssDNA sequences for sgRNA-*mitfa* were annealed together in NE buffer r2.1.

Gibson assembly was used to insert sgRNA-*mitfa* into linearized U6 plasmid. Assembled

product was transformed into NEB 5-alpha competent *E. coli* cells via heat shock and selected

with 50 ug/mL streptomycin. Insertion of sgRNA-*mitfa* was confirmed using PCR amplification of

lysed bacteria from single colonies (forward primer: ATGGACAAAGCTGGACCATG, reverse

primer: CTTAGCTGGATAACGCCAC). Positive products were isolated using the QIAGEN

Plasmid Maxi Kit. Entire plasmid was sequenced using Plasmidsaurus sequencing service.

Transgene Electroporation of Adult Zebrafish Skin (TEAZ-Skin)

Adult zebrafish were anesthetized in 0.2% Tricaine prior to upright placement on a prewetted

Kimwipe. Each injection consisted of 1000 ng total DNA: 100 ng of Tol2 plasmid and an equal

division of the remaining 900 ng for the remaining constructs. Injection stocks contained a small

volume (<1uL, 1:50 dilution) of green food coloring to visualize localization of injection. Injection

volumes ranged from 1 to 1.5 uL depending on the injection mix, keeping 1000 ng total

constant. Needles were pulled from 10 cm long filamented borosilicate glass capillaries at a

pressure of 500, heat of 650, pull strength of 100, velocity of 200, and time delay of 40 in a

Model P-97 Flaming/Brown micropipette puller from Sutter Instrument Co. Following injection,

zebrafish were immediately electroporated (<1 min) by placing the cathode probe of ultrasound

gel-coated Platnium Tweezertrode, 3MM, on the side of the injection site and the anode probe

on the other side of the fish body. Electroporation was achieved with 5 60-ms pulses of 40 volts

with a 1s pulse interval from a ECM 830 Electro Square Porator from BTX Harvard Apparatus.

Following electroporation, zebrafish are placed into a recovery tank with fresh, gently stirred

system water and are monitored until normal swimming resumes. Electroporated zebrafish are

imaged weekly. *mitfa*:*GNAQ*^{Q209L}; *mitfa*:GFP plasmid constructs were kindly provided by Dr.

Jacqueline A. Lees (Massachusetts Institute of Technology, Cambridge, USA). *mitfa*:Cas9,

mitfa:tdTomato, PCS2FA Tol2, *U6*:gRNA-*tp53, U6*:gRNA-*ptena,* and *U6*:gRNA-*ptenb* plasmids

were kindly provided by Dr. Richard White (Sloan Kettering Institute, New York, USA).

Transgene Electroporation of Adult Zebrafish Eye (TEAZ-Eye)

Pre-injection preparation for eye injections is the same as for skin injections including: 0.2%

Tricaine anesthesia, needle size, and injection concentrations. To inject the eye, a blunt

spherical head of a pin was gently pressed against the bottom half of the zebrafish eyeball to

reveal the dorsal posterior regions of the eye. The needle was gently pressed into this location,

where the cornea is softer, just through the sclera, keeping the needle as superficial as possible

as to not damage the back of the eye. Injection volumes were around 0.25 uL and were

terminated with a slight visual swelling of the eye. Following injection zebrafish were

immediately electroporated (<1 min) by the cathode probe of ultrasound gel-coated Platnium

Tweezertrode, 1MM, on the superior side of the eye nearest the injection and the anode probe

on the inferior side of the eye. Electroporation was achieved with 3 50-ms pulses of 75 volts with

a 1s pulse interval from a ECM 830 Electro Square Porator from BTX Harvard Apparatus.

Recovery and imaging procedures for these fish did not differ from the skin injections.

Imaging and image processing

Injected zebrafish were imaged using an upright Leica M205 FCA microscope with brightfield

and GFP filters. Zebrafish were briefly anesthetized with 0.2% Tricaine and positioned on a

Kimwipe mound for upright orientation. Images were taken and processed through the LAS X

software. All histology slides were imaged using brightfield, DAPI, and GFP filters on an EVOS

M5000 microscope from Invitrogen for individual images or on a Leica DMi8 microscope and

stitched together by LAS X Thunder Imaging software for full tissue images.

H&E and IF – zebrafish tissue

Whole zebrafish or zebrafish eyes were fixed in 4% paraformaldehyde for 48 hours at 4°C and

paraffin embedded in the University of Iowa Pathology Core. Fish were sectioned at 7 µM

thickness and placed on Superfrost Plus Microscope slides from Fisher Scientific, baked at

60°C, stained with hematoxylin and eosin, and mounted with Cytoseal 60 from Epredia. For

immunofluorescence, sections were deparaffinized and hydrated followed by heat-induced

antigen retrieval with 10 mM Na-citrate buffer and blocking with 3% milk in 0.1% Tween 20.

slides were stained with primary antibodies against GFP (NB600-308 Rabbit anti-GFP

polyclonal from Novus Biologicals at 1:250) overnight at 4°C and then fluorescent secondary

antibodies against rabbit IgG (Alexa Fluor 488 goat anti-rabbit IgG at 1:500 from a 2 mg/mL

stock) for 1 hr at room temperature. RPE autofluorescence was minimized using the Vector

TrueVIEW Autofluorescence Quenching Kit. Sections were counterstained with DAPI, mounted

using VECTASHIELD Vibrance Antifade Mounting Medium, and imaged the same day.

IF - mouse eye tissue

C57BL/6 mice eyes were collected, formalin-fixed, paraffin-embedded, and sectioned into 7 μm-

thick sections. Sections were deparaffinized and hydrated followed by heat-induced antigen

retrieval with citrate buffer (Vector Laboratories, Inc.), permeabilized with 1% horse serum in

0.2% Triton-X, blocked with 5% Horse serum in 0.2% Triton-X, and incubated with PAX3 primary

antibody (1:200 anti-rabbit, Invitrogen, 38-1801) overnight at 4°C. Samples were incubated with

DyLightTM 488-labeled secondary antibody (1:2000, horse anti-rabbit. Vector Laboratories, Inc.

DI-1088-1.5) for 1 hr at room temperature, treated with autofluorescence quenching kit (Vector

Laboratories, Inc. SP-8400-15), and cover slips fitted using Vectashield Antifade Mounting

Medium with DAPI (Vector Laboratories, Inc. H-1000-10).

Kaplan-Meier analysis

Zebrafish were followed for up to 70 weeks and tumor-free survival was analyzed with the

Kaplan-Meier method. Tumors were defined as the first presence of a GFP-positive nodular

mass at the injection site. Differences between injection groups were analyzed using log-rank

statistics.

Single cell suspension and single-cell RNA Seq analysis

Zebrafish were euthanized with 0.4% Tricaine for 10 mins followed by immersion in ice water

until no opercular movement was observed for at least 30 min as detailed in our IACUC

protocol. Tumors were harvested and minced into small pieces. Minced tumors were digested

for 1 hour at 37°C with collagenase and hyaluronidase in HBSS with 2% FBS (HF).

Suspensions were washed with 1:4 HF:NH₄Cl and then further digested for 5 min with pre-

warmed trypsin. Following trypsin inactivation, cell suspensions were washed twice with 10%

FBS in HBSS.

Cellular suspensions were loaded on a 10x Genomics Chromium instrument to generate single-

cell gel beads in emulsion (GEMs). Approximately 10,000-20,000 cells were loaded per channel

depending on the 10x kit used. See figures S5 and S12 for kit information per sample.

Next GEM kit: targeted cell number = 10,000. Single-cell RNA-Seq libraries were prepared

using Single Cell 3′ Reagent Kits v2: Chromium Single Cell 3′ Library & Gel Bead Kit v2, PN-

120237; Single Cell 3′ Chip Kit v2, PN-120236; and i7 Multiplex Kit, PN-120262 (10x Genomics)

and following the Single Cell 3′ Reagent Kits v2 User Guide (Manual Part # CG00052 Rev A).

GEM-X kit: targeted cell number = 20,000. Single-cell RNA-Seq libraries were prepared using

Chromium GEM-X Single Cell 3′ Reagent Kits v4: Library Construction Kit C, PN-1000694;

Single Cell 3' GEM Kit v4, PN-1000693; Single Cell 3' Gel Bead Kit v4, PN-2001128; and Dual

Index Kit TT Set A, PN-3000431; and following the Chromium GEM-X Single Cell 3′ Reagent

Kits v4 User Guide (Manual Part # CG000731 Rev B). Libraries were sequenced on an Illumina

HiSeq 4000 as 2 × 150 paired-end reads. Sequencing results were demultiplexed and converted

to FASTQ format using Illumina bcl2fastq software.

A custom reference genome for the Zebrafish was constructed with GRCz11 primary assembly

(Ensembl) using Cell Ranger (Cell Ranger mkref function). 10X Genomics scRNA-seq reads

were then processed and aligned to this reference with Cell Ranger (Cell Ranger Count

function). Further analysis and visualization was performed using Seurat (v4.1.0)[101], cells with

fewer than 200 RNA feature counts and greater than 5% mitochondrial contamination were

removed with filtering (greater than 15% for wildtype eye samples). RNA counts were

normalized, and FindVariableFeatures was run with the following parameters: selection.method

= "vst", nfeatures = 2,000. The cells were originally clustered in a UMAP by RunPCA then

RunUMAP with dims 1:30. FindNeighbors was run with dims 1:30 followed by FindClusters

using a resolution of 0.4. These methods identified up to 20 clusters in individual samples.

For comparative analysis, RNA-seq datasets were integrated together using IntegrateData with

dims 1:30. New clusters were generated as before. Clusters were annotated using

FindAllMarkers and the top 25 enriched genes in each cluster were compared to the Daniocell

dataset[52] and Spectacle[49]. Cluster names were given according to the closest matched cellular

population between these two datasets. To determine enriched pathways, *danio rerio* gene

names were converted into human Ensembl gene codes with g:Profiler[102] and then data were

analyzed with the use of QIAGEN IPA (QIAGEN Inc)[103] and GSEA[104,105]. Pseudotime was

performed using Monocle3 (v0.2.3.0)[106].

Verification of sgRNAs

Zebrafish were euthanized with 0.4% Tricaine for 10 mins followed by immersion in ice water

until no opercular movement was observed for at least 10 min as detailed in the IACUC

protocol. Tumors were harvested and minced into small pieces. DNA was harvested according

to the DNeasy Blood and Tissue Kit from Qiagen. Regions surrounding cut sites were PCR-

amplified and the purity of the PCR amplicon was confirmed with agarose gel electrophoresis.

Samples were cleaned up with the QIAquick PCR Purification Kit from Qiagen and sent for

Sanger sequencing at the Iowa Institute of Human Genetics. Gene editing was confirmed with

DECODR INDEL analysis by comparing to wildtype sequence[107]. PCR primer sequences follow:

mitfa PCR F: CCGGTATGTATTCACATTGTCTTG

mitfa PCR R: TCTTGCTTAGGATGCCTATGTATT

ptena PCR F: CATCCCACCAAGTGAGGTTAAAC

ptena PCR R: CACATACACAGTCAAGGGTGAG

ptenb PCR F: CAGTTCTGTTGCACCCAATAAG

ptenb PCR R: CTGGTGGTGTTGAGGCTATAAAG

tp53 PCR F: AAGTATTCAGCCCCCAGGTG

tp53 PCR R: CGCTTTTGACTCACAGTGCAAG

Figure 1: *GNAQ*^{Q209L}-induced melanoma by TEAZ-Skin and TEAZ-Eye.

(**A**) TEAZ-Skin was used to induce GNAQ-positive melanoma using two strategies. <u>Strategy A</u> involves a single plasmid construct containing expression cassettes driven by the *mitfa* promoter for GNAQ^{Q209L}-T2A-eGFP and Cas9, as well as a U6 promoter driving the expression of three separate gRNAs (targeting *tp53*, *ptena*, and *ptenb*, or a non-targeting control). <u>Strategy B</u> uses a multi-plasmid approach with five plasmids for TEAZ-Skin. Expression cassettes include: (a) *mitfa*:*GNAQ*^{Q209L}-*mitfa*:eGFP, (b) *mitfa*:Cas9, (c) *U6*:gRNA (*tp53*), (d) *U6*:gRNA (*ptena*), and (e) *U6*:gRNA (*ptenb*).

(**B**) Schematic of TEAZ-Skin using wild type (*AB*), *nacre* (*mitfa*$^{w2/w2}$) and *casper* (*mitfa*$^{w2/w2}$: *mpv17*$^{a9/a9}$) zebrafish lines.

(**C**) TEAZ-Skin using Strategy B plasmids in adult wild-type (*AB*) and *casper* zebrafish. Representative brightfield and fluorescent images shown at 4, 8, and 12 weeks post electroporation for WT and at 2, 4, and 8 weeks for *casper*.

(**D**) Kaplan-Meier curves were used to estimate tumor-free survival, defined as the time from electroporation until the first visible appearance of GFP+ nodular tumors, in adult zebrafish via TEAZ-Skin. Strategy B wild-type (n=7), *nacre* (n=5), and *casper* (n=17) curves and TEAZ-Skin (*GNAQ*^{Q209L}:GFP) curves (n=3 per condition) compared pairwise with logrank (Mantel-Cox) test. ns = not significant, * p-value<0.05; ** p-value<0.01; ***p-value<0.001; ****p-value<0.0001.

(**E**) Schematic of the zebrafish eye, dashed box indicates the retina and choroid. Hematoxylin and eosin (H&E) staining of a wild-type zebrafish eye at 20× magnification, showing the retina and choroid. Eye structures are labeled, including choroidal melanocytes.

(**F**) Immunofluorescent analysis of transgenic *Tg(mitfa:GFP)* zebrafish using an anti-GFP antibody. Sections were counterstained with DAPI to visualize nuclei. Representative image showing GFP expression in *mitfa*-positive cells. Secondary-only antibody was used as background control.

(**G**) Schematic of TEAZ-Eye using choroidal targeted injection.

(**H-H'**) Live imaging of adult wild-type zebrafish following TEAZ-Eye with Strategy B plasmids to

induce uveal melanoma. Brightfield imaging and the corresponding GFP fluorescence images

highlighting tumor progression. Timepoints in day post-electroporation (dpe) as labeled.

(**I-I'**) Live imaging of adult *casper* zebrafish over 21 days post-electroporation (dpe) using TEAZ-

Eye with Strategy B plasmids to induce uveal melanoma. Brightfield imaging and the

corresponding GFP fluorescence images highlighting tumor progression.

(**J**) Kaplan-Meier curves comparing tumor-free survival among wild-type (n=5) and *casper*

(n=10) zebrafish following TEAZ-Eye induction using Strategy B plasmids. Curves compared

pairwise with logrank (Mantel-Cox) test. ** p-value<0.01.

(**K, K'**) H&E analysis of uveal melanoma in *casper* zebrafish induced by TEAZ-Eye. Images

were acquired at 20× magnification and stitched from 85 fields using Thunder imaging software.

Normal eye tissue and uveal melanoma as labeled.

(**K'**) Representative 40x magnification image showing GFP expression in tumors cells after anti-

GFP immunofluorescent analysis of TEAZ-Eye induced uveal melanoma in *casper* zebrafish.

Figure 2: Single cell profiling of uveal melanoma and the TME in adult zebrafish.

(**A**) UMAP representation of single-cell RNA-Seq (scRNA-seq) data from three dissociated wild-

type eyes harboring uveal melanoma (UM) that were generated by TEAZ-Eye (sequencing

replicates, n=2), and control eyes consisting of four pooled sibling-matched wild-type eyes, and

two tumor paired control eyes (sequencing replicates, n=2). Annotated cell clusters as labeled.

(**B**) UMAP and feature plot showing *eGFP* and *GNAQ*^{Q209L} expression in control eyes and UM

tumor. Red dotted line highlights the melanocyte and UM clusters.

(**C**) Comparison of UM tumor marker genes with the markers of cell types within the control eye

microenvironment. Only genes with a log2FC > 1 and q-value < 0.01 were included in the

analysis.

(**D**) Dot plot representing enriched genes (log2FC > 1; q-value < 0.01) in choroidal cell

populations, UM tumors cells, immune cell populations, fibroblasts, Schwann cells, pericytes,

and vascular endothelial cells as shown. Dot size indicates the percentage of cells expressing

each gene; color intensity (grey to blue) reflects normalized average expression levels (low to

high).

(**E**) UMAP clustering of cells in the choroidal microenvironment. Annotated cell clusters as

labelled.

(**F**) Comparison of relative cluster sizes between the two conditions, control and TEAZ-Eye,

shown in (E). Schwann cells serve as a control cluster, displaying no change in frequency

between conditions. Gray = TEAZ-Eye, Blue = Control.

(**G**) Venn diagram representing the overlap of genes between UM tumors and choroidal

melanocytes compared to the cells in their respective microenvironments. Enriched genes for

each condition were identified by differential expression analysis comparing melanocytes to the

microenvironment (ME) or tumor cells to the tumor microenvironment (TME), using a threshold

of log2FC > 1 and q-value < 0.01. Genes of interest have been labelled in their respective

subsets. (**G'**) HOMER promoter motif enrichment analysis of genes enriched in UM tumors vs

TME. The top ranked motif for "known" and "*de novo*" motifs are shown with their corresponding

p-values.

(**H**) Dot plot showing enriched genes (log2FC > 1; q-value < 0.01) in UM tumor cells, choroidal

melanocytes, or the tumor microenvironment (TME). Dot size indicates the percentage of cells

expressing each gene; color intensity (grey to blue) reflects normalized average expression

levels (low to high).

(**I-K**) Kaplan–Meier disease-free survival and overall survival curves generated using uveal

melanoma TCGA data. Patients stratified by relative expression levels of *KIT*, *FABP3*, or

LGALS2. Gene expression thresholds were defined by median expression within each cohort.

Log rank p-values as shown. Dotted lines represent 95% confidence interval.

(**L**) Dot plot showing enriched genes (log2FC > 1; q-value < 0.01) in cancer-associated

fibroblasts (CAF), normal fibroblasts (NF), or the tumor microenvironment (TME) without CAF.

Dot size indicates the percentage of cells expressing each gene; color intensity (grey to blue)

reflects normalized average expression levels (low to high). Genes have been grouped

according to their labelled functions.

Figure 3: Comparison of GNAQ-driven tumors and primary melanocytes in the skin and

eyes of adult zebrafish.

(**A**) UMAP representation of GNAQ-positive tumors generated using TEAZ-Eye and TEAZ-Skin,

along with their respective control tissues using scRNA-seq. Tumor samples include three

dissociated wild-type eyes harboring UM tumors generated via TEAZ-Eye and two dissociated

wild-type skin UM tumors generated via TEAZ-Skin (n=2 sequencing replicates). Control eye

samples consist of a pool of two uninjected tumor paired wild-type eyes, three mock-injected

sibling-matched wild-type eyes, and four uninjected sibling-matched wild-type eyes (n=3

sequencing replicates). Control skin samples consist of dissociated normal skin from two wild-

type zebrafish (n=2 sequencing replicates). Cell clusters are annotated as tumor cells (red),

tumor microenvironment (TME) cells shared by both eye and skin samples (light gray), and TME

cells specific to the eye samples (dark gray).

(**B**) Live imaging of representative samples used for single-cell RNA-Seq analysis and Venn

diagram showing overlapping genes between TEAZ-Eye, TEAZ-Skin, choroidal melanocytes,

and skin melanocytes compared to the cells in their respective microenvironments. Enriched

genes for each condition were identified by differential expression analysis comparing

melanocytes to the microenvironment (ME) or tumor cells to the tumor microenvironment (TME),

using a threshold of log2FC > 0.5 and q-value < 0.01.

(**C**) Table displaying overlaps between enriched gene sets from (B). Percentages represented

as the number of overlapping genes over the total unique genes of both gene sets.

(**D**) UMAP representation of primary melanocytes and UM tumor cells from both eye (salmon)

and skin (teal).

(**E**) Volcano plot of differentially expressed genes (DEGs) between eye and skin primary

melanocytes (log2FC > |0.5| and q-value < 0.01).

(**F**) Volcano plot of differentially expressed genes (DEGs) between eye and skin UM tumors

(log2FC > |0.5| and q-value < 0.01).

(**G**) GSEA enrichment plots of pathways that were enriched in skin melanocytes and of

pathways enriched in eye melanocytes from (E). P-values and FDRs as labelled.

(**H**) GSEA enrichment plots of pathways that were enriched in UM skin tumors and of pathways

enriched in UM eye tumors from (F). P-values and FDRs as labelled..

1010

1011

Figure 4: *mitf*-independent melanocyte progenitor cells are expanded in *mitfa*-deficient

zebrafish and are susceptible to *GNAQ*^{Q209L} transformation.

(**A**) H&E analysis representation of the retina and choroid in wild-type (left), *nacre* (middle), and

casper (right) zebrafish (20x magnification). Note the loss of melanocytes (red arrows) in the

choroid of *nacre* and *casper* eyes. "Vit" = vitreous, "GCL" = ganglion cell layer, "INL" = inner

nuclear layer, "ONL" = outer nuclear layer, "POS" = photoreceptor outer segment, "RPE" =

retinal pigmented epithelium, "CH" = choroid.

(**B**) Immunofluorescent analysis of transgenic *Tg(mitfa:GFP); mitfa*$^{w2/w2}$ (*nacre*) zebrafish using

an anti-GFP antibody. Sections were counterstained with DAPI to visualize nuclei.

Representative images showing GFP-positive cells in the ciliary body and choroid are shown by

yellow arrow heads. Secondary-only antibody was used as background control.

(**C**) UMAP representation of single-cell RNA sequencing (scRNA-seq) data from three paired

dissociated *casper* eyes and from three *casper* eyes harboring uveal melanoma (UM) tumors

induced using Strategy B via TEAZ-Eye (n=1 sequencing replicate for each). Annotated cell

clusters are labeled. Progenitor cells and UM tumor clusters are outlined with dashed red lines.

Representative brightfield images of a normal eye and a UM tumor in *casper* zebrafish are

shown in the top left corner of the UMAP.

(**D**) Re-clustered UMAP representing progenitor cells from control eyes and uveal melanoma

(UM) tumor cells from TEAZ-Eye–injected *casper* zebrafish. Heterogeneous cell populations as

labeled.

(**E**) Re-clustered UMAP from (D) showing melanocyte progenitor cells from control eyes in red

and UM tumor cells from TEAZ-Eye in blue. Feature plots for *pax3a, foxd3* and *sox10*

expression are shown, dashed line marks the progenitor cell population.

(**F**) Dot plot illustrating differentiational expressed genes (log2FC > 0.5; q-value < 0.05)

between heterogeneous cell populations in (D). Dot size indicates the percentage of cells

expressing each gene; color intensity (blue to red) reflects normalized average expression

levels (low to high).

(**G, G'**) UMAP obtained after clustering GFP-positive cells sorted from *Tg(mitfa:GFP)* and

Tg(mitfa:GFP); mitfa$^{w2/w2}$ (*nacre*) zebrafish embryos at 28 hours post fertilization (hpf).

Annotated cell clusters as labelled. (**G'**) Comparison of relative cluster size for the two

genotypes; *Tg(mitfa:GFP)* and *Tg(mitfa:GFP); mitfa$^{w2/w2}$* (*nacre*) in (G).

(**H**) Dot plot illustrating differentiational expressed genes (log2FC > 0.5; q-value < 0.05) in

melanophores and progenitor cells in *mitfa$^{w2/w2}$* and sibling embryos at 28hpf. Dot size indicates

the percentage of cells expressing each gene; color intensity (grey to red) reflects normalized

average expression levels (low to high).

(**I**) Kaplan–Meier curves comparing tumor-free survival in wild-type and *casper* zebrafish

following TEAZ-Eye injection using Strategy B plasmids for *mitfa* conditional knockout. Wild-type

zebrafish were injected with either Strategy B plasmids plus non-targeting gRNAs (n=9) or

Strategy B plasmids plus *mitfa*-targeting gRNAs (n=9). *Casper* zebrafish were injected with

Strategy B plasmids and *mitfa*-targeting gRNAs (n=9).

(**J**) Representative brightfield and GFP-overlay images of tumors that formed in *casper* (9 of 9)

and wild-type zebrafish injected with *mitfa*-targeting gRNAs (2 of 9). Metastatic cells are

indicated by red arrows.

1055

Figure 5: Oncogenic GNAQ and BRAF transform transcriptionally distinct cells within the

melanocyte lineage.

(**A**) UMAP obtained after clustering tumor cells from $Tg(mitfa:BRAF^{V600E}): tp53^{-/-}: mitfa^{w2/w2}$

zebrafish where TEAZ-Skin was used to rescue *mitfa* expression and knockout *ptena* and *ptenb*

to induce cutaneous melanoma[32], along with our GNAQ-induced tumors by TEAZ-Skin (Strategy

B: *mitfa:Cas9, U6:*gRNA-*tp53, U6:*gRNA-*ptena, U6:*gRNA-*ptenb, mitfa:GNAQ^{Q209L}-mitfa:GFP*).

BRAF-induced tumors in red and GNAQ-induced tumors in blue.

(**B**) Violin plot showing differentially expressed genes (log2FC > 0.5, q-value < 0.05) between

BRAF-induced and GNAQ-induced tumors from (A). The plot displays the distribution of gene

expression across individual cells, with the width indicating cell density at specific expression

levels and the height representing the range of expression values.

(**C**) Integrated UMAP after clustering GFP-positive cells sorted from *Tg(mitfa:GFP)* and

$Tg(mitfa:GFP); mitfa^{w2/w2}$ (*nacre*) zebrafish embryos at 28 hours post fertilization (hpf).

Annotated cell clusters as labelled.

(**D**) Average enrichment of signature genes from BRAF- and GNAQ-driven tumors projected

onto the UMAP in (C) using UCell with 'FeaturePlot' function. Signature genes represent the top

50 enriched genes from each tumor genotype that are also expressed in the embryonic

melanocyte lineage.

(**E**) UMAP of primary melanocytes and melanocyte progenitors from 9 control wild-type eyes, 2

control wild-type skin sections, and 3 control *casper* eyes. Cells are labeled according to their

genotype (*casper* and wild-type) and tissue origin (eye and skin).

(**F**) Feature plots representing the relative mRNA expression of *pax3a*, *foxd3*, *sox10*, and *tfec* in

melanocytes described in (E). Color intensity (grey to red) reflects normalized average

expression levels (low to high). The red dotted line indicates cells that are clustering closely with

the melanocyte progenitors identified in *casper* eyes.

1081

1082

Figure S1: Mitfa-deficient zebrafish have decreased tumor latency compared to wild-types by

TEAZ-Skin. (**A**) Kaplan-Meier curves estimating tumor-free survival, defined as the time from

electroporation until the first visible appearance of GFP+ nodular tumors, in adult wild-type

zebrafish via TEAZ-Skin. Injections contained *mitfa*:Cas9 with *U6*:gRNA-*ptena* + *U6*:gRNA-

ptenb (*ptena/b* CRISPR), *mitfa*:Cas9 with *U6*:gRNA-*tp53* (*tp53* CRISPR), *mitfa*:$GNAQ^{Q209L}$-

mitfa:GFP (GNAQ: GFP), strategy A plasmids, or strategy B plasmids (n=3 per condition). (**B**)

Kaplan-Meier curves estimating tumor-free survival in adult *nacre* (*mitfa*$^{w2/w2}$) zebrafish via

TEAZ-Skin. Same injection mixes as in (A). (n=5 for Strategy B, n=3 for rest) (**B'**)

Representative brightfield and fluorescent images shown at 2, 4, and 6 weeks for strategy B

injection in *nacre*. (**C**) Kaplan-Meier curves estimating tumor-free survival in adult *casper*

(*mitfa*$^{w2/w2}$, *mpv17*$^{a9/a9}$) zebrafish via TEAZ-Skin. Same injection mixes as in (A). (n=5 for

Strategies A and B, n=3 for rest) (**D**) Kaplan-Meier curves estimating tumor-free survival in adult

wild-type, *nacre*, and *casper* zebrafish via TEAZ-Skin. Oncogene alone (GNAQ: GFP) is

compared to strategy B plasmids for each genotype. (n=5 for strategy B in *nacre* and *casper*,

n=3 for rest). (**E**) Representative images of Strategy B TEAZ-Skin in wild-type and *casper* and a

xanthoma that formed in a *Tg(mitfa:BRAF*^{V600E}*); mitfa*$^{w2/w2}$*; tp53*$^{-/-}$ zebrafish. Note the bright

orange color of the xanthoma, indicated by the black arrow.

1100

1101 **Figure S2:** Targeted Sanger sequencing to confirm CRISPR gene editing following TEAZ-Skin

1102 injection using Strategy B plasmids. 11 tumors and 2 wild-type tissue controls were used for

1103   INDEL analysis to verify gene editing in *tp53, ptena,* and *ptenb.* Green box indicates successful

1104   gene editing; red box indicates no gene editing. Total number of edited genes per sample are

1105   indicated in the right-most column.

1106

1107   **Figure S3:** Immunofluorescence of the choroid using an anti-GFP antibody in (**A**) *Tg*(*mitfa*:GFP)

1108   with 10x magnification and in (**B**) *Tg*(*mitfa*:GFP) following quenching of autofluorescence. (**C**)

1109   TEAZ-Eye–injected zebrafish expressing an eGFP construct driven by the *mitfa* promoter

1110   (*mitfa:GFP*) with 40x magnification. Sections were counterstained with DAPI to visualize nuclei.

1111

1112   **Figure S4:** UMAP of normal eyes and TEAZ-Eye uveal melanoma in wild-type zebrafish

1113   (Strategy B). (**A**) Integrated object is split by sequencing replicates (top) and feature plots

1114   showing *mlana* RNA expression to identify normal choroidal melanocytes and uveal melanoma

1115   cells (bottom). Color intensity (grey to red) reflects normalized average expression levels (low to

1116   high). The red dotted line indicates the melanocyte and uveal melanoma clusters.

1117

1118   **Figure S5:** Gross images of tissue and tumors used for single cell RNA-Seq in wild-type

1119   zebrafish.

1120

1121   **Figure S6:** Ingenuity Pathway Analysis of genes enriched in TEAZ-Eye compared to the tumor

1122   microenvironment. (**A**) The top 49 pathways enriched in TEAZ-Eye. (**B**) Select pathways

1123   enriched in TEAZ-Eye pertaining to fatty acid oxidation and signaling. Color intensity indicates

1124   the sign and magnitude of the z-score from blue (negative) to orange (positive). Significance is

1125   indicated by the -log(p-value).

1126

1127   **Figure S7:** Overall survival of patients with uveal melanoma (UM) and cutaneous melanoma

1128   (CM) based on expression levels of *FABP3*, *KIT, FABP5, LGALS2,* and *GCH*. Disease-free

1129   survival in UM and CM patients based on *FABP3, KIT, and FABP5* expression. Dotted lines

1130   represent 95% confidence intervals. Data derived from TCGA datasets.

1131

1132   **Figure S8:** Normalized RNA expression of *FABP5, FABP3, LGALS1, GCH, LGALS2, KIT*

1133   across the 4 TCGA molecular subclasses. [55] Groups were compared by ANOVA with post-hoc

1134   Tukey test. ns = not significant, * p-value<0.05; ** p-value<0.01; ***p-value<0.001; ****p-

1135   value<0.0001.

1136

Figure S9: Mock injected wild-type zebrafish eyes do not show an increase in activated

fibroblast markers as compared to uninjected controls. (**A**) UMAP representation of wild-type

normal eyes from three different sources: mock injected, sibling-matched (3 pooled eyes),

uninjected tumor-paired (2 pooled eyes), and uninjected sibling-matched (4 pooled eyes). The

red dotted line indicates the fibroblast clusters. (**B**) Feature plots showing *fn1a* RNA expression

across wild-type eye control groups. Color intensity (grey to red) reflects normalized average

expression levels (low to high). (**C**) Dot plot of fibroblast activation marker genes in the fibroblast

clusters of wild-type eye control groups. Dot size indicates the percentage of cells expressing

each gene; color intensity (grey to blue) reflects normalized average expression levels (low to

high).

1147

Figure S10: Ingenuity Pathway Analysis of enriched genes in TEAZ-Eye and TEAZ-Skin

compared to the combined tumor microenvironment (eye and skin). (**A**) Enriched pathways from

genes that were common to only TEAZ-Eye and TEAZ-Skin compared to the combined tumor

microenvironment. Families of pathways have been identified with a colored box and colored

label including: cancer proliferation (green), WNT signaling (blue), metabolism (red), PTEN

regulation (black), and RAF/MAPK signaling (black). (**B**) Enriched pathways from genes that

were common to all four gene subsets (TEAZ-Eye, TEAZ-Skin, choroidal melanocytes, skin

melanocytes) versus their respective combined microenvironments (normal or tumor). Families

of pathways have been identified with a colored box and colored label including: melanocyte

identity (black), signals downstream of GNAQ/GNA11 (blue), and survival/proliferation (green).

Color intensity indicates the sign and magnitude of the z-score from blue (negative) to orange

(positive). Significance is indicated by the -log(p-value).

1160

Figure S11: Melanocyte rescue experiments in *nacre* zebrafish by TEAZ-Eye using plasmids

containing *mitfa*:*mitfa*; *mitfa*:GFP (miniCoopR[29]). (**A-A'**) H&E images (10x and 40x

magnification) of wild-type (*AB*) eyes following TEAZ-Eye injection with (**A**) control *mitfa*-GFP or

(**A'**) miniCoopR plasmids. (**B-B'**) H&E images (10x and 40x magnification) of *mitfa$^{w2/w2}$* (*nacre*)

eyes following TEAZ-Eye injection with (**B**) control *mitfa*-GFP or (**B'**) miniCoopR plasmids,

rescued melanocytes in the choroid are indicated by a red asterisk. (**C**) Representative dorsal

image of a transgenic *Tg(mitfa:GFP); mitfa$^{w2/w2}$* injected with *mitfa*:*GNAQ^{Q209L}* following TEAZ-

Eye injection.

1169

1170   **Figure S12:** Gross images of tissue and tumors used for single cell RNA-Seq in *casper*

1171   zebrafish and transgenic *Tg*(*mitfa*:GFP); *mitfa$^{+/w2}$* and *Tg*(*mitfa*:GFP); *mitfa$^{w2/w2}$* whole embryos.

1172

Figure S13: UMAP representation of UM tumors derived from *casper* zebrafish, with

accompanying pseudotime analysis. (**A**) UMAP representation following scRNA-seq on normal

casper control eyes (3 pooled eyes) and UM tumors (3 pooled tumors) derived in *casper* (TEAZ-

Eye) (**B**) Feature plot showing *eGFP* expression and *mitfa$^{w2/w2}$* expression (**C**) UMAP following

re-clustering of *mitfa$^{w2/w2}$* positive cells from (B), with overlayed unsupervised pseudotime

analysis using Monocle analysis.

1179

1180 **Figure S14:** Immunofluorescent staining for Pax3 in the mouse limbus. (**A**) Representative 20x

1181 image of the mouse eye showing Pax3 positive staining in the limbus region. Sections were

1182 counterstained with DAPI to visualize nuclei. (**B**) Post-image processing showing the retina

1183 (red), cornea (orange) and limbus (purple), Pax3-positive cells in green.

1184

1185 **Figure S15:** Representative images of TEAZ-Skin induced tumors (*Strategy B*) in (**A**) wild-type

1186 at 16 weeks, (**B**) conditional *mitfa*-KO (wild-type) at 8 weeks and (**C**) *casper* zebrafish at 8

1187 weeks post TEAZ-Skin. Brightfield and GFP-overlaid images are shown to highlight primary and

1188 metastatic cells in (B-C), and the absence of metastasis in (A).

1189

1190 **Figure S16:** Violin plots representing *mitfa* expression as well as pigmentation genes in

1191 $BRAF^{V600E}$- and $GNAQ^{Q209L}$-positive tumors by single cell RNA-Seq analysis.

1192

1193 **Figure S17:** Dot plot of genes enriched in chromatoblasts and in uveal melanoma tumors

1194 derived in wild-type zebrafish following TEAZ-Eye compared to the tumor microenvironment

1195 (TME). Genes enriched in a particular pigment cell type are labelled, including: iridophore,

1196 xanthophore, and melanocyte.

1197

1198 **Figure S18:** UMAP representation of melanocytes and melanocyte progenitors in *casper*

1199 normal eyes and in the three sequencing replicates of wild-type normal eyes. The red dotted

1200 line indicates melanocyte progenitors in the *casper* sample and melanocytes that are clustering

1201 closely with such progenitors in wild-type samples.

1202

1203

1204

Supplemental Table 1 – Findallmarkers of the integrated object between wild-type control eyes

and wild-type TEAZ-Eye.

Supplemental Table 2 – RNA expression data of *FABP3*, *FABP5*, *LGALS2*, *LGALS1*, *BSCL2*,

GCH1, and *KIT* across the 4 TCGA molecular subclasses in uveal melanoma with

corresponding sample IDs.

Supplemental Table 3 – Differentially expressed genes between cancer associated fibroblasts

from wild-type TEAZ-Eye and normal fibroblasts from wild-type normal eyes.

Supplemental Table 4 – Findmarkers for choroidal melanocytes, skin melanocytes, TEAZ-Eye,

and TEAZ-Skin versus their respective microenvironments. Table also includes gene names in

every intersection of the Venn Diagram comparison of all four groups.

Supplemental Table 5 – Differentially expressed genes between eye and skin melanocytes and

between TEAZ-Eye and TEAZ-Skin, respectively.

Supplemental Table 6 – Findallmarkers of the *casper* TEAZ-Eye single-cell object.

Supplemental Table 7 – Findallmarkers of the *casper* normal eye single-cell object.

Supplemental Table 8 – Findallmarkers of the integrated object between *casper* TEAZ-Eye and

casper normal eye with cluster names labelled.

Supplemental Table 9 – Findallmarkers of the subset of GFP-positive clusters from the

integrated object between *casper* TEAZ-Eye and *casper* normal eye.

Supplemental Table 10 – Findallmarkers of the integrated object of neural crest and neural

crest-derived cells from *Tg*(*mitfa*:GFP) *mitfa*$^{+/w2}$ and *Tg*(mitfa:GFP) *mitfa*$^{w2/w2}$ embryos.

Supplemental Table 11 – Differentially expressed genes between neural crest and neural crest-

derived clusters from *Tg*(*mitfa*:GFP) *mitfa*$^{+/w2}$ and *Tg*(mitfa:GFP) *mitfa*$^{w2/w2}$ embryos.

Supplemental Table 12 – Injection and tumor penetrance of wild-type and *casper* zebrafish

injected with Strategy B plasmids and either *U6*:gRNA-nontargeting or *U6*:gRNA-*mitfa*.

Supplemental Table 13 – Differentially expressed genes between *BRAF*^{V600E}-driven tumors and

GNAQ^{Q209L}-driven tumors in adult zebrafish.

1.   Gentien, D., Saberi-Ansari, E., Servant, N., Jolly, A., de la Grange, P., Némati, F., Liot, G.,
Saule, S., Teissandier, A., Bourc'his, D., et al. (2023). Multi-omics comparison of
malignant and normal uveal melanocytes reveals molecular features of uveal melanoma.
Cell Rep *42*, 113132. 10.1016/j.celrep.2023.113132.
2.   Rodrigues, M., Mobuchon, L., Houy, A., Alsafadi, S., Baulande, S., Mariani, O., Marande,
B., Ait Rais, K., Van der Kooij, M.K., Kapiteijn, E., et al. (2019). Evolutionary Routes in
Metastatic Uveal Melanomas Depend on MBD4 Alterations. Clin Cancer Res *25*, 5513-
5524. 10.1158/1078-0432.Ccr-19-1215.
3.   Kolandjian, N.A., Wei, C., Patel, S.P., Richard, J.L., Dett, T., Papadopoulos, N.E., and
Bedikian, A.Y. (2013). Delayed systemic recurrence of uveal melanoma. Am J Clin Oncol
36, 443-449. 10.1097/COC.0b013e3182546a6b.
4.   Jager, M.J., Shields, C.L., Cebulla, C.M., Abdel-Rahman, M.H., Grossniklaus, H.E., Stern,
M.H., Carvajal, R.D., Belfort, R.N., Jia, R., Shields, J.A., and Damato, B.E. (2020). Uveal
melanoma. Nat Rev Dis Primers *6*, 24. 10.1038/s41572-020-0158-0.
5.   Lamas, N.J., Martel, A., Nahon-Estève, S., Goffinet, S., Macocco, A., Bertolotto, C.,
Lassalle, S., and Hofman, P. (2021). Prognostic Biomarkers in Uveal Melanoma: The
Status Quo, Recent Advances and Future Directions. Cancers (Basel) *14*.
10.3390/cancers14010096.
6.   Kaliki, S., and Shields, C.L. (2017). Uveal melanoma: relatively rare but deadly cancer. Eye
(Lond) *31*, 241-257. 10.1038/eye.2016.275.
7.   Yu, F.X., Luo, J., Mo, J.S., Liu, G., Kim, Y.C., Meng, Z., Zhao, L., Peyman, G., Ouyang, H.,
Jiang, W., et al. (2014). Mutant Gq/11 promote uveal melanoma tumorigenesis by
activating YAP. Cancer Cell *25*, 822-830. 10.1016/j.ccr.2014.04.017.
8.   Singh, A.D., Bergman, L., and Seregard, S. (2005). Uveal melanoma: epidemiologic
aspects. Ophthalmol Clin North Am *18*, 75-84, viii. 10.1016/j.ohc.2004.07.002.
9.   Arnesen, K. (1985). The neural crest origin of uveal melanomas. Int Ophthalmol *7*, 143-
147. 10.1007/bf00128360.
10. Van Raamsdonk, C.D., Bezrookove, V., Green, G., Bauer, J., Gaugler, L., O'Brien, J.M.,
Simpson, E.M., Barsh, G.S., and Bastian, B.C. (2009). Frequent somatic mutations of
GNAQ in uveal melanoma and blue naevi. Nature *457*, 599-602. 10.1038/nature07586.
11. Shain, A.H., Bagger, M.M., Yu, R., Chang, D., Liu, S., Vemula, S., Weier, J.F., Wadt, K.,
Heegaard, S., Bastian, B.C., and Kiilgaard, J.F. (2019). The genetic evolution of metastatic
uveal melanoma. Nat Genet *51*, 1123-1130. 10.1038/s41588-019-0440-9.
12. Silva-Rodríguez, P., Fernández-Díaz, D., Bande, M., Pardo, M., Loidi, L., and Blanco-
Teijeiro, M.J. (2022). GNAQ and GNA11 Genes: A Comprehensive Review on
Oncogenesis, Prognosis and Therapeutic Opportunities in Uveal Melanoma. Cancers
(Basel) *14*. 10.3390/cancers14133066.
13. Li, Y., Shi, J., Yang, J., Ge, S., Zhang, J., Jia, R., and Fan, X. (2020). Uveal melanoma:
progress in molecular biology and therapeutics. Ther Adv Med Oncol *12*,
1758835920965852. 10.1177/1758835920965852.
14. Gragoudas, E.S., Egan, K.M., Seddon, J.M., Glynn, R.J., Walsh, S.M., Finn, S.M.,
Munzenrider, J.E., and Spar, M.D. (1991). Survival of patients with metastases from uveal
melanoma. Ophthalmology *98*, 383-389; discussion 390. 10.1016/s0161-6420(91)32285-
1.

15. Rantala, E.S., Hernberg, M., and Kivelä, T.T. (2019). Overall survival after treatment for
metastatic uveal melanoma: a systematic review and meta-analysis. Melanoma Res *29*,
561-568. 10.1097/cmr.0000000000000575.
16. Lane, A.M., Kim, I.K., and Gragoudas, E.S. (2018). Survival Rates in Patients After
Treatment for Metastasis From Uveal Melanoma. JAMA Ophthalmol *136*, 981-986.
10.1001/jamaophthalmol.2018.2466.
17. Nathan, P., Hassel, J.C., Rutkowski, P., Baurain, J.F., Butler, M.O., Schlaak, M., Sullivan,
R.J., Ochsenreither, S., Dummer, R., Kirkwood, J.M., et al. (2021). Overall Survival Benefit
with Tebentafusp in Metastatic Uveal Melanoma. N Engl J Med *385*, 1196-1206.
10.1056/NEJMoa2103485.
18. Hassel, J.C., Piperno-Neumann, S., Rutkowski, P., Baurain, J.F., Schlaak, M., Butler, M.O.,
Sullivan, R.J., Dummer, R., Kirkwood, J.M., Orloff, M., et al. (2023). Three-Year Overall
Survival with Tebentafusp in Metastatic Uveal Melanoma. N Engl J Med *389*, 2256-2266.
10.1056/NEJMoa2304753.
19. Wespiser, M., Neidhardt, E., and Negrier, S. (2023). Uveal melanoma: In the era of new
treatments. Cancer Treat Rev *119*, 102599. 10.1016/j.ctrv.2023.102599.
20. Kaštelan, S., Antunica, A.G., Oresković, L.B., Pelčić, G., Kasun, E., and Hat, K. (2020).
Immunotherapy for Uveal Melanoma - Current Knowledge and Perspectives. Curr Med
Chem *27*, 1350-1366. 10.2174/0929867326666190704141444.
21. Wdowiak, K., Dolar-Szczasny, J., Rejdak, R., Drab, A., and Maciocha, A. (2025). A Brief
Overview of Uveal Melanoma Treatment Methods with a Focus on the Latest Advances.
J Clin Med *14*. 10.3390/jcm14124058.
22. Souto, E.B., Zielinska, A., Luis, M., Carbone, C., Martins-Gomes, C., Souto, S.B., and Silva,
A.M. (2019). Uveal melanoma: physiopathology and new in situ-specific therapies.
Cancer Chemother Pharmacol *84*, 15-32. 10.1007/s00280-019-03860-z.
23. Ma, J., Weng, L., Bastian, B.C., and Chen, X. (2021). Functional characterization of uveal
melanoma oncogenes. Oncogene *40*, 806-820. 10.1038/s41388-020-01569-5.
24. Livingstone, E., Zaremba, A., Horn, S., Ugurel, S., Casalini, B., Schlaak, M., Hassel, J.C.,
Herbst, R., Utikal, J.S., Weide, B., et al. (2020). GNAQ and GNA11 mutant nonuveal
melanoma: a subtype distinct from both cutaneous and uveal melanoma. Br J Dermatol
183, 928-939. 10.1111/bjd.18947.
25. Lee, J.H., Choi, J.W., and Kim, Y.S. (2011). Frequencies of BRAF and NRAS mutations are
different in histological types and sites of origin of cutaneous melanoma: a meta-
analysis. Br J Dermatol *164*, 776-784. 10.1111/j.1365-2133.2010.10185.x.
26. Urtatiz, O., Haage, A., Tanentzapf, G., and Van Raamsdonk, C.D. (2021). Crosstalk with
keratinocytes causes GNAQ oncogene specificity in melanoma. Elife *10*.
10.7554/eLife.71825.
27. Callahan, S.J., Tepan, S., Zhang, Y.M., Lindsay, H., Burger, A., Campbell, N.R., Kim, I.S.,
Hollmann, T.J., Studer, L., Mosimann, C., and White, R.M. (2018). Cancer modeling by
Transgene Electroporation in Adult Zebrafish (TEAZ). Dis Model Mech *11*.
10.1242/dmm.034561.
28. Brombin, A., Simpson, D.J., Travnickova, J., Brunsdon, H., Zeng, Z., Lu, Y., Young, A.I.J.,
Chandra, T., and Patton, E.E. (2022). Tfap2b specifies an embryonic melanocyte stem cell

that retains adult multifate potential. Cell Rep *38*, 110234.
10.1016/j.celrep.2021.110234.

29. Ceol, C.J., Houvras, Y., Jane-Valbuena, J., Bilodeau, S., Orlando, D.A., Battisti, V., Fritsch,
L., Lin, W.M., Hollmann, T.J., Ferré, F., et al. (2011). The histone methyltransferase
SETDB1 is recurrently amplified in melanoma and accelerates its onset. Nature *471*, 513-
517. 10.1038/nature09806.

30. Kenny, C., Dilshat, R., Seberg, H.E., Van Otterloo, E., Bonde, G., Helverson, A., Franke,
C.M., Steingrímsson, E., and Cornell, R.A. (2022). TFAP2 paralogs facilitate chromatin
access for MITF at pigmentation and cell proliferation genes. PLoS Genet *18*, e1010207.
10.1371/journal.pgen.1010207.

31. Lister, J.A., Lane, B.M., Nguyen, A., and Lunney, K. (2011). Embryonic expression of
zebrafish MiT family genes tfe3b, tfeb, and tfec. Dev Dyn *240*, 2529-2538.
10.1002/dvdy.22743.

32. Lumaquin-Yin, D., Montal, E., Johns, E., Baggiolini, A., Huang, T.H., Ma, Y., LaPlante, C.,
Suresh, S., Studer, L., and White, R.M. (2023). Lipid droplets are a metabolic vulnerability
in melanoma. Nat Commun *14*, 3192. 10.1038/s41467-023-38831-9.

33. Petratou, K., Spencer, S.A., Kelsh, R.N., and Lister, J.A. (2021). The MITF paralog tfec is
required in neural crest development for fate specification of the iridophore lineage
from a multipotent pigment cell progenitor. PLoS One *16*, e0244794.
10.1371/journal.pone.0244794.

34. Phelps, G.B., Hagen, H.R., Amsterdam, A., and Lees, J.A. (2022). MITF deficiency
accelerates GNAQ-driven uveal melanoma. Proc Natl Acad Sci U S A *119*, e2107006119.
10.1073/pnas.2107006119.

35. Seberg, H.E., Van Otterloo, E., Loftus, S.K., Liu, H., Bonde, G., Sompallae, R., Gildea, D.E.,
Santana, J.F., Manak, J.R., Pavan, W.J., et al. (2017). TFAP2 paralogs regulate melanocyte
differentiation in parallel with MITF. PLoS Genet *13*, e1006636.
10.1371/journal.pgen.1006636.

36. Kelsh, R.N., Brand, M., Jiang, Y.J., Heisenberg, C.P., Lin, S., Haffter, P., Odenthal, J.,
Mullins, M.C., van Eeden, F.J., Furutani-Seiki, M., et al. (1996). Zebrafish pigmentation
mutations and the processes of neural crest development. Development *123*, 369-389.
10.1242/dev.123.1.369.

37. Dutton, K.A., Pauliny, A., Lopes, S.S., Elworthy, S., Carney, T.J., Rauch, J., Geisler, R.,
Haffter, P., and Kelsh, R.N. (2001). Zebrafish colourless encodes sox10 and specifies non-
ectomesenchymal neural crest fates. Development *128*, 4113-4125.
10.1242/dev.128.21.4113.

38. McGowan, K.A., and Barsh, G.S. (2016). Evolution: How the zebrafish got its stripes. Elife
5. 10.7554/eLife.14239.

39. Parichy, D.M., and Spiewak, J.E. (2015). Origins of adult pigmentation: diversity in
pigment stem cell lineages and implications for pattern evolution. Pigment Cell
Melanoma Res *28*, 31-50. 10.1111/pcmr.12332.

40. Phelps, G.B., Amsterdam, A., Hagen, H.R., García, N.Z., and Lees, J.A. (2022). MITF
deficiency and oncogenic GNAQ each promote proliferation programs in zebrafish
melanocyte lineage cells. Pigment Cell Melanoma Res *35*, 539-547. 10.1111/pcmr.13057.

41.   Wessely, A., Steeb, T., Berking, C., and Heppt, M.V. (2021). How Neural Crest
Transcription Factors Contribute to Melanoma Heterogeneity, Cellular Plasticity, and
Treatment Resistance. Int J Mol Sci *22*. 10.3390/ijms22115761.

42.   Kaufman, C.K., Mosimann, C., Fan, Z.P., Yang, S., Thomas, A.J., Ablain, J., Tan, J.L., Fogley,
R.D., van Rooijen, E., Hagedorn, E.J., et al. (2016). A zebrafish melanoma model reveals
emergence of neural crest identity during melanoma initiation. Science *351*, aad2197.
10.1126/science.aad2197.

43.   Lister, J.A., Robertson, C.P., Lepage, T., Johnson, S.L., and Raible, D.W. (1999). nacre
encodes a zebrafish microphthalmia-related protein that regulates neural-crest-derived
pigment cell fate. Development *126*, 3757-3767. 10.1242/dev.126.17.3757.

44.   Lister, J.A., Capper, A., Zeng, Z., Mathers, M.E., Richardson, J., Paranthaman, K., Jackson,
I.J., and Elizabeth Patton, E. (2014). A conditional zebrafish MITF mutation reveals MITF
levels are critical for melanoma promotion vs. regression in vivo. J Invest Dermatol *134*,
133-140. 10.1038/jid.2013.293.

45.   Travnickova, J., Wojciechowska, S., Khamseh, A., Gautier, P., Brown, D.V., Lefevre, T.,
Brombin, A., Ewing, A., Capper, A., Spitzer, M., et al. (2019). Zebrafish MITF-Low
Melanoma Subtype Models Reveal Transcriptional Subclusters and MITF-Independent
Residual Disease. Cancer Res *79*, 5769-5784. 10.1158/0008-5472.Can-19-0037.

46.   D'Agati, G., Beltre, R., Sessa, A., Burger, A., Zhou, Y., Mosimann, C., and White, R.M.
(2017). A defect in the mitochondrial protein Mpv17 underlies the transparent casper
zebrafish. Dev Biol *430*, 11-17. 10.1016/j.ydbio.2017.07.017.

47.   White, R.M., Sessa, A., Burke, C., Bowman, T., LeBlanc, J., Ceol, C., Bourque, C., Dovey,
M., Goessling, W., Burns, C.E., and Zon, L.I. (2008). Transparent adult zebrafish as a tool
for in vivo transplantation analysis. Cell Stem Cell *2*, 183-189.
10.1016/j.stem.2007.11.002.

48.   McMenamin, P.G., Shields, G.T., Seyed-Razavi, Y., Kalirai, H., Insall, R.H., Machesky, L.M.,
and Coupland, S.E. (2020). Melanoblasts Populate the Mouse Choroid Earlier in
Development Than Previously Described. Invest Ophthalmol Vis Sci *61*, 33.
10.1167/iovs.61.10.33.

49.   Voigt, A.P., Whitmore, S.S., Lessing, N.D., DeLuca, A.P., Tucker, B.A., Stone, E.M., Mullins,
R.F., and Scheetz, T.E. (2020). Spectacle: An interactive resource for ocular single-cell
RNA sequencing data analysis. Exp Eye Res *200*, 108204. 10.1016/j.exer.2020.108204.

50.   Voigt, A.P., Mulfaul, K., Mullin, N.K., Flamme-Wiese, M.J., Giacalone, J.C., Stone, E.M.,
Tucker, B.A., Scheetz, T.E., and Mullins, R.F. (2019). Single-cell transcriptomics of the
human retinal pigment epithelium and choroid in health and macular degeneration. Proc
Natl Acad Sci U S A *116*, 24100-24107. 10.1073/pnas.1914143116.

51.   Howe, D.G., Bradford, Y.M., Conlin, T., Eagle, A.E., Fashena, D., Frazer, K., Knight, J., Mani,
P., Martin, R., Moxon, S.A., et al. (2013). ZFIN, the Zebrafish Model Organism Database:
increased support for mutants and transgenics. Nucleic Acids Res *41*, D854-860.
10.1093/nar/gks938.

52.   Sur, A., Wang, Y., Capar, P., Margolin, G., Prochaska, M.K., and Farrell, J.A. (2023). Single-
cell analysis of shared signatures and transcriptional diversity during zebrafish
development. Dev Cell *58*, 3028-3047.e3012. 10.1016/j.devcel.2023.11.001.

53.   Tatarakis, D., Cang, Z., Wu, X., Sharma, P.P., Karikomi, M., MacLean, A.L., Nie, Q., and
Schilling, T.F. (2021). Single-cell transcriptomic analysis of zebrafish cranial neural crest
reveals spatiotemporal regulation of lineage decisions during development. Cell Rep *37*,
110140. 10.1016/j.celrep.2021.110140.
54.   Simões-Costa, M., and Bronner, M.E. (2015). Establishing neural crest identity: a gene
regulatory recipe. Development *142*, 242-257. 10.1242/dev.105445.
55.   Robertson, A.G., Shih, J., Yau, C., Gibb, E.A., Oba, J., Mungall, K.L., Hess, J.M.,
Uzunangelov, V., Walter, V., Danilova, L., et al. (2017). Integrative Analysis Identifies Four
Molecular and Clinical Subsets in Uveal Melanoma. Cancer Cell *32*, 204-220.e215.
10.1016/j.ccell.2017.07.003.
56.   Hunter, M.V., Joshi, E., Bowker, S., Montal, E., Ma, Y., Kim, Y.H., Yang, Z., Tuffery, L., Li, Z.,
Rosiek, E., et al. (2025). Mechanical confinement governs phenotypic plasticity in
melanoma. Nature. 10.1038/s41586-025-09445-6.
57.   Hunter, M.V., Moncada, R., Weiss, J.M., Yanai, I., and White, R.M. (2021). Spatially
resolved transcriptomics reveals the architecture of the tumor-microenvironment
interface. Nat Commun *12*, 6278. 10.1038/s41467-021-26614-z.
58.   Liu, T., Zhou, L., Xiao, Y., Andl, T., and Zhang, Y. (2022). BRAF Inhibitors Reprogram
Cancer-Associated Fibroblasts to Drive Matrix Remodeling and Therapeutic Escape in
Melanoma. Cancer Res *82*, 419-432. 10.1158/0008-5472.Can-21-0614.
59.   Mao, X., Xu, J., Wang, W., Liang, C., Hua, J., Liu, J., Zhang, B., Meng, Q., Yu, X., and Shi, S.
(2021). Crosstalk between cancer-associated fibroblasts and immune cells in the tumor
microenvironment: new findings and future perspectives. Mol Cancer *20*, 131.
10.1186/s12943-021-01428-1.
60.   Wang, J., Song, Y., Li, Z., Gao, T., Shen, W., Kang, Z., and Xu, C. (2025). Combining single-
cell and bulk RNA sequencing to identify CAF-related signature for prognostic prediction
and treatment response in patients with melanoma. Sci Rep *15*, 29082. 10.1038/s41598-
025-14979-w.
61.   Kaochar, S., Dong, J., Torres, M., Rajapakshe, K., Nikolos, F., Davis, C.M., Ehli, E.A., Coarfa,
C., Mitsiades, N., and Poulaki, V. (2018). ICG-001 Exerts Potent Anticancer Activity
Against Uveal Melanoma Cells. Invest Ophthalmol Vis Sci *59*, 132-143. 10.1167/iovs.17-
22454.
62.   Zheng, L., Liu, Y., and Pan, J. (2017). Inhibitory effect of pyrvinium pamoate on uveal
melanoma cells involves blocking of Wnt/β-catenin pathway. Acta Biochim Biophys Sin
(Shanghai) *49*, 890-898. 10.1093/abbs/gmx089.
63.   Wu, S., Han, M., and Zhang, C. (2022). Overexpression of microRNA-130a represses uveal
melanoma cell migration and invasion through inactivation of the Wnt/β-catenin
signaling pathway by downregulating USP6. Cancer Gene Ther *29*, 930-939.
10.1038/s41417-021-00377-7.
64.   Zuidervaart, W., van Nieuwpoort, F., Stark, M., Dijkman, R., Packer, L., Borgstein, A.M.,
Pavey, S., van der Velden, P., Out, C., Jager, M.J., et al. (2005). Activation of the MAPK
pathway is a common event in uveal melanomas although it rarely occurs through
mutation of BRAF or RAS. Br J Cancer *92*, 2032-2038. 10.1038/sj.bjc.6602598.

65. Chen, X., Wu, Q., Depeille, P., Chen, P., Thornton, S., Kalirai, H., Coupland, S.E., Roose,
J.P., and Bastian, B.C. (2017). RasGRP3 Mediates MAPK Pathway Activation in GNAQ
Mutant Uveal Melanoma. Cancer Cell *31*, 685-696.e686. 10.1016/j.ccell.2017.04.002.
66. Rimoldi, D., Salvi, S., Liénard, D., Lejeune, F.J., Speiser, D., Zografos, L., and Cerottini, J.C.
(2003). Lack of BRAF mutations in uveal melanoma. Cancer Res *63*, 5712-5715.
67. Abdel-Rahman, M.H., Yang, Y., Zhou, X.P., Craig, E.L., Davidorf, F.H., and Eng, C. (2006).
High frequency of submicroscopic hemizygous deletion is a major mechanism of loss of
expression of PTEN in uveal melanoma. J Clin Oncol *24*, 288-295.
10.1200/jco.2005.02.2418.
68. Pópulo, H., Soares, P., Rocha, A.S., Silva, P., and Lopes, J.M. (2010). Evaluation of the
mTOR pathway in ocular (uvea and conjunctiva) melanoma. Melanoma Res *20*, 107-117.
10.1097/CMR.0b013e32832ccd09.
69. Woodman, S.E. (2012). Metastatic uveal melanoma: biology and emerging treatments.
Cancer J *18*, 148-152. 10.1097/PPO.0b013e31824bd256.
70. Ulrich, E., Kistenmacher, S., Martin, G., Schlötzer-Schrehardt, U., Seitz, B., Auw-Hädrich,
C., Schlunck, G., Reinhard, T., and Polisetti, N. (2025). PAX3 expression patterns in ocular
surface melanocytes. Sci Rep *15*, 12472. 10.1038/s41598-025-90318-3.
71. Kumar, N., Singh, M.K., Singh, L., Jha, J., Pushker, N., Lomi, N., Meel, R., Sen, S., and
Kashyap, S. (2025). Prognostic significance of pigmentation and stem cell markers in
Indian population of uveal melanoma. Br J Ophthalmol *109*, 799-808. 10.1136/bjo-2024-
325886.
72. Kubic, J.D., Young, K.P., Plummer, R.S., Ludvik, A.E., and Lang, D. (2008). Pigmentation
PAX-ways: the role of Pax3 in melanogenesis, melanocyte stem cell maintenance, and
disease. Pigment Cell Melanoma Res *21*, 627-645. 10.1111/j.1755-148X.2008.00514.x.
73. Lang, D., Lu, M.M., Huang, L., Engleka, K.A., Zhang, M., Chu, E.Y., Lipner, S., Skoultchi, A.,
Millar, S.E., and Epstein, J.A. (2005). Pax3 functions at a nodal point in melanocyte stem
cell differentiation. Nature *433*, 884-887. 10.1038/nature03292.
74. Baggiolini, A., Callahan, S.J., Montal, E., Weiss, J.M., Trieu, T., Tagore, M.M., Tischfield,
S.E., Walsh, R.M., Suresh, S., Fan, Y., et al. (2021). Developmental chromatin programs
determine oncogenic competence in melanoma. Science *373*, eabc1048.
10.1126/science.abc1048.
75. Johns, E., Ma, Y., Louphrasitthiphol, P., Peralta, C., Hunter, M.V., Raymond, J.H., Molina,
H., Goding, C.R., and White, R.M. (2025). The Lipid Droplet Protein DHRS3 Is a Regulator
of Melanoma Cell State. Pigment Cell Melanoma Res *38*, e13208. 10.1111/pcmr.13208.
76. Xu, Y., Xu, W.H., Yang, X.L., Zhang, H.L., and Zhang, X.F. (2020). Fatty acid-binding protein
5 predicts poor prognosis in patients with uveal melanoma. Oncol Lett *19*, 1771-1780.
10.3892/ol.2020.11301.
77. Van Ly, D., Wang, D., Conway, R.M., Giblin, M., Liang, S., Lukeis, R., Lim, L.A., Hesson, L.,
and Cherepanoff, S. (2020). Lipid-Producing Ciliochoroidal Melanoma with Expression of
HMG-CoA Reductase. Ocul Oncol Pathol *6*, 416-421. 10.1159/000510393.
78. Yavuzyigitoglu, S., Kilic, E., Vaarwater, J., de Klein, A., Paridaens, D., and Verdijk, R.M.
(2016). Lipomatous Change in Uveal Melanoma: Histopathological,
Immunohistochemical and Cytogenetic Analysis. Ocul Oncol Pathol *2*, 133-135.
10.1159/000440981.

79.   Fiorentzis, M., Kalirai, H., Katopodis, P., and Coupland, S.E. (2017). Adipophilin
1491   expression in primary and metastatic uveal melanoma: a pilot study. Graefes Arch Clin
1492   Exp Ophthalmol *255*, 1049-1051. 10.1007/s00417-017-3630-0.
1493   80.   Horvathova Kajabova, V., Soltysova, A., Demkova, L., Plesnikova, P., Lyskova, D., Furdova,
1494   A., Babal, P., and Smolkova, B. (2021). KIT Expression Is Regulated by DNA Methylation in
1495   Uveal Melanoma Tumors. Int J Mol Sci *22*. 10.3390/ijms221910748.
1496   81.   Hu, S., Huang, J., Pei, S., Ouyang, Y., Ding, Y., Jiang, L., Lu, J., Kang, L., Huang, L., Xiang, H.,
1497   et al. (2019). Ganoderma lucidum polysaccharide inhibits UVB-induced melanogenesis
1498   by antagonizing cAMP/PKA and ROS/MAPK signaling pathways. J Cell Physiol *234*, 7330-
1499   7340. 10.1002/jcp.27492.
1500   82.   Liang, R., Cheng, A., Lu, S., Zhang, X., Ren, M., Lin, J., Wu, Y., Zhang, W., and Luan, X.
1501   (2024). Seleno-amino Acid Metabolism Reshapes the Tumor Microenvironment: from
1502   Cytotoxicity to Immunotherapy. Int J Biol Sci *20*, 2779-2789. 10.7150/ijbs.95484.
1503   83.   Li, W., and Cornell, R.A. (2007). Redundant activities of Tfap2a and Tfap2c are required
1504   for neural crest induction and development of other non-neural ectoderm derivatives in
1505   zebrafish embryos. Dev Biol *304*, 338-354. 10.1016/j.ydbio.2006.12.042.
1506   84.   Lewis, J.L., Bonner, J., Modrell, M., Ragland, J.W., Moon, R.T., Dorsky, R.I., and Raible,
1507   D.W. (2004). Reiterated Wnt signaling during zebrafish neural crest development.
1508   Development *131*, 1299-1308. 10.1242/dev.01007.
1509   85.   Monsoro-Burq, A.H., Wang, E., and Harland, R. (2005). Msx1 and Pax3 cooperate to
1510   mediate FGF8 and WNT signals during Xenopus neural crest induction. Dev Cell *8*, 167-
1511   178. 10.1016/j.devcel.2004.12.017.
1512   86.   Han, A., Purwin, T.J., Bechtel, N., Liao, C., Chua, V., Seifert, E., Sato, T., Schug, Z.T.,
1513   Speicher, D.W., Harbour, J.W., and Aplin, A.E. (2021). BAP1 mutant uveal melanoma is
1514   stratified by metabolic phenotypes with distinct vulnerability to metabolic inhibitors.
1515   Oncogene *40*, 618-632. 10.1038/s41388-020-01554-y.
1516   87.   Matatall, K.A., Agapova, O.A., Onken, M.D., Worley, L.A., Bowcock, A.M., and Harbour,
1517   J.W. (2013). BAP1 deficiency causes loss of melanocytic cell identity in uveal melanoma.
1518   BMC Cancer *13*, 371. 10.1186/1471-2407-13-371.
1519   88.   Chang, J., Campbell-Hanson, K.R., Vanneste, M., Bartschat, N.I., Nagel, R., Arnadottir,
1520   A.K., Vu, H.N., Montgomery, C., Yevdash, J., Jiang, J., et al. (2025). Antagonistic roles for
1521   MITF and TFE3 in melanoma plasticity. Cell Rep *44*, 115474.
1522   10.1016/j.celrep.2025.115474.
1523   89.   Pouw, A.E., Greiner, M.A., Coussa, R.G., Jiao, C., Han, I.C., Skeie, J.M., Fingert, J.H.,
1524   Mullins, R.F., and Sohn, E.H. (2021). Cell-Matrix Interactions in the Eye: From Cornea to
1525   Choroid. Cells *10*. 10.3390/cells10030687.
1526   90.   Chirco, K.R., Worthington, K.S., Flamme-Wiese, M.J., Riker, M.J., Andrade, J.D.,
1527   Ueberheide, B.M., Stone, E.M., Tucker, B.A., and Mullins, R.F. (2017). Preparation and
1528   evaluation of human choroid extracellular matrix scaffolds for the study of cell
1529   replacement strategies. Acta Biomater *57*, 293-303. 10.1016/j.actbio.2017.05.011.
1530   91.   Angi, M., Kalirai, H., Prendergast, S., Simpson, D., Hammond, D.E., Madigan, M.C.,
1531   Beynon, R.J., and Coupland, S.E. (2016). In-depth proteomic profiling of the uveal
1532   melanoma secretome. Oncotarget *7*, 49623-49635. 10.18632/oncotarget.10418.

92. Aughton, K., Hattersley, J., Coupland, S.E., and Kalirai, H. (2024). Revealing the structural
microenvironment of high metastatic risk uveal melanomas following decellularisation.
Sci Rep *14*, 26811. 10.1038/s41598-024-78171-2.

93. Luthold, C., Didion, M., Benedum, E., Burkhart, A.-K., Demmerle, N., Wirth, E.,
Gunaratnam, G., Racz, V.S., Thangamurugan, S., Helms, V., et al. (2025). Melanocyte
differentiation and mechanosensation are differentially modulated by distinct
extracellular matrix proteins. bioRxiv, 2024.2010.2004.616635.
10.1101/2024.10.04.616635.

94. Hajkova, N., Hojny, J., Nemejcova, K., Dundr, P., Ulrych, J., Jirsova, K., Glezgova, J., and
Ticha, I. (2018). Germline mutation in the TP53 gene in uveal melanoma. Sci Rep *8*, 7618.
10.1038/s41598-018-26040-0.

95. Harbour, J.W., Onken, M.D., Roberson, E.D., Duan, S., Cao, L., Worley, L.A., Council, M.L.,
Matatall, K.A., Helms, C., and Bowcock, A.M. (2010). Frequent mutation of BAP1 in
metastasizing uveal melanomas. Science *330*, 1410-1413. 10.1126/science.1194472.

96. Brantley, M.A., Jr., and Harbour, J.W. (2000). Deregulation of the Rb and p53 pathways in
uveal melanoma. Am J Pathol *157*, 1795-1801. 10.1016/s0002-9440(10)64817-1.

97. Coupland, S.E., Anastassiou, G., Stang, A., Schilling, H., Anagnostopoulos, I., Bornfeld, N.,
and Stein, H. (2000). The prognostic value of cyclin D1, p53, and MDM2 protein
expression in uveal melanoma. J Pathol *191*, 120-126. 10.1002/(sici)1096-
9896(200006)191:2<120::Aid-path591>3.0.Co;2-p.

98. Paraoan, L., Gray, D., Hiscott, P., Ebrahimi, B., Damato, B., and Grierson, I. (2006).
Expression of p53-induced apoptosis effector PERP in primary uveal melanomas:
downregulation is associated with aggressive type. Exp Eye Res *83*, 911-919.
10.1016/j.exer.2006.04.016.

99. Davies, L., Spiller, D., White, M.R., Grierson, I., and Paraoan, L. (2011). PERP expression
stabilizes active p53 via modulation of p53-MDM2 interaction in uveal melanoma cells.
Cell Death Dis *2*, e136. 10.1038/cddis.2011.19.

100. de Lange, J., Ly, L.V., Lodder, K., Verlaan-de Vries, M., Teunisse, A.F., Jager, M.J., and
Jochemsen, A.G. (2012). Synergistic growth inhibition based on small-molecule p53
activation as treatment for intraocular melanoma. Oncogene *31*, 1105-1116.
10.1038/onc.2011.309.

101. Butler, A., Hoffman, P., Smibert, P., Papalexi, E., and Satija, R. (2018). Integrating single-
cell transcriptomic data across different conditions, technologies, and species. Nat
Biotechnol *36*, 411-420. 10.1038/nbt.4096.

102. Kolberg, L., Raudvere, U., Kuzmin, I., Adler, P., Vilo, J., and Peterson, H. (2023). g:Profiler-
interoperable web service for functional enrichment analysis and gene identifier
mapping (2023 update). Nucleic Acids Res *51*, W207-w212. 10.1093/nar/gkad347.

103. Krämer, A., Green, J., Pollard, J., Jr., and Tugendreich, S. (2014). Causal analysis
approaches in Ingenuity Pathway Analysis. Bioinformatics *30*, 523-530.
10.1093/bioinformatics/btt703.

104. Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A.,
Paulovich, A., Pomeroy, S.L., Golub, T.R., Lander, E.S., and Mesirov, J.P. (2005). Gene set
enrichment analysis: a knowledge-based approach for interpreting genome-wide

expression profiles. Proc Natl Acad Sci U S A *102*, 15545-15550.
10.1073/pnas.0506580102.
105.   Mootha, V.K., Lindgren, C.M., Eriksson, K.F., Subramanian, A., Sihag, S., Lehar, J.,
Puigserver, P., Carlsson, E., Ridderstråle, M., Laurila, E., et al. (2003). PGC-1alpha-
responsive genes involved in oxidative phosphorylation are coordinately downregulated
in human diabetes. Nat Genet *34*, 267-273. 10.1038/ng1180.
106.   Trapnell, C., Cacchiarelli, D., Grimsby, J., Pokharel, P., Li, S., Morse, M., Lennon, N.J.,
Livak, K.J., Mikkelsen, T.S., and Rinn, J.L. (2014). The dynamics and regulators of cell fate
decisions are revealed by pseudotemporal ordering of single cells. Nat Biotechnol *32*,
381-386. 10.1038/nbt.2859.
107.   Bloh, K., Kanchana, R., Bialk, P., Banas, K., Zhang, Z., Yoo, B.C., and Kmiec, E.B. (2021).
Deconvolution of Complex DNA Repair (DECODR): Establishing a Novel Deconvolution
Algorithm for Comprehensive Analysis of CRISPR-Edited Sanger Sequencing Data. Crispr j
4, 120-131. 10.1089/crispr.2020.0022.
1590

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

FIGURE S1

Tumor #	ptena	ptenb	tp53	# of edited genes
1				3
2				3
3				2
4				3
5				3
6				3
7				3
8				3
9				3
10				3
11				3
Germline	ptena	ptenb	tp53	
WT1				0
WT2				0

CRISPR editing

No editing

FIGURE S2

A
Tg(mitfa-GFP) Zebrafish Line

B
Tg(mitfa-GFP) Zebrafish Line

C
TEAZ-Eye using *mitfa:GFP* plasmids into WT Zebrafish

FIGURE S3

A

FIGURE S4

Single-cell RNA seq Replicates

Tissue	Genotype	Replicate 1 (pooled tissue, tumors)	Replicate 2 (pooled tissue, tumors)
Skin	Wild-type	1 skin sample	1 skin sample
Skin Tumor	Wild-type	1 tumor sample	1 tumor sample
Eye Paired (tumor matched bilateral eye)	Wild-type	2 pooled eyes	
Eye sibling matched	Wild-type	4 pooled eyes	
Eye mock injected, sibling matched	Wild-type	3 pooled eyes	
Eye Tumor	Wild-type	1 tumor sample	2 pooled tumor samples

FIGURE S5

FIGURE S6

FIGURE S7

FIGURE S8

FIGURE S9

FIGURE S10

A Control injected: Wild type

RPE

Choroid

A' TEAZ-Eye (miniCoopR) : Wild type

RPE

Choroid

B Control injected: *nacre*

RPE

Choroid

B' TEAZ-Eye (miniCoopR): *nacre*

RPE

Choroid

C

TEAZ-Eye: (*mitfa:GNAQ^{Q209L}*)

Tg(mitfa:GFP): mitfa^{w2/w2}

GFP

Figure S11

Single-cell RNA seq Replicates

Tissue	Genotype	Replicate 1 (pooled tissue, tumors)
Eye	Casper	3 pooled eyes
Eye Tumor	Casper	3 pooled tumors
Whole embryo 28hpf	Wild-type Tg(mitfa:GFP); mitfa$^{+/w2}$	Pooled embryos 25 total
Whole embryo 28hpf	Nacre Tg(mitfa:GFP); mitfa$^{w2/w2}$	Pooled embryos 25 total

Figure S12

Figure S13

PAX3 DAPI

Retina

Limbus   Cornea

FIGURE S14

A TEAZ-Skin: *Strategy B* (Wild type)

16 weeks

Wild type · Lateral · Dorsal

Bright overlay GFP

GFP · GFP

B TEAZ-Skin: *Strategy B + gRNA-mitfa* (Wild type)

8 weeks

Wild type · Lateral

GFP · metastasis · Bright overlay GFP

C TEAZ-Skin: *Strategy B* (Casper)

8 weeks

Lateral · Dorsal

metastasis · primary · metastasis · primary

Bright overlay GFP · Bright overlay GFP

Figure S15

Figure S16

TEAZ-Eye Strategy B (Wild-type)

Figure S17

Figure S18