1 **Targeting *Leishmania donovani* Sphingosine Kinase 1 using PF-543 enhances immune**
2 **response and limits parasite load**
3
4 **Short Title: Targeting Sphingosine-1-phosphate kinase to limit *Leishmania* growth**
5
6

7 **Evanka Madan**[1,2]**, Ruby Bansal**[1]**, Jhalak Singhal**[1]**, Monica Saini**[3]**, Sadat Shafi**[1,4]**, Nishant**
8 **Joshi**[5]**, Shailja Singh**[1*]
9

[1]Special Centre for Molecular Medicine, Jawaharlal Nehru University, New Delhi
110067, India
[2]Department of Biotechnology and Research, Sir Ganga Ram Hospital, New Delhi-110008,
India
[3]Biology Centre Czech Academy of Sciences, Institute of Parasitology, Laboratory of RNA
Biology of Protists, Czech Republic
[4]Department of Pharmacology, School of Pharmaceutical Education and Research, Jamia
Hamdard, New Delhi, India.
[5] Eli Lilly Services India Private Limited, Bengaluru, Karnataka 560103
19

email addresses of authors:
Evanka Madan: evanka.madan@gmail.com
Ruby Bansal: rubybansal1242@gmail.com
Jhalak Singhal: jhalakbiotech@gmail.com
Monica Saini: monikabcas@gmail.com
Sadat Shafi: sadatshafi2016@gmail.com
Nishant Joshi: nj633@snu.edu.in
27
28

29 Correspondence:
30 ∗Shailja Singh; Special Centre for Molecular Medicine, Jawaharlal Nehru University, Delhi,
31 110067, India; shailja.jnu@gmail.com
32
33

Abstract

Background

Sphingosine-1-phosphate (S1P) is a bioactive lipid mediator regulating apoptosis,

proliferation, and immune responses. While S1Ps presence in *Leishmania donovani*

phagolysosomes has been reported, the role of sphingosine kinases, especially SphK1, in

parasite survival and host immune modulation remains underexplored. This study

investigates the molecular and functional role of *L. donovani* SphK1 (*Ld*SphK1) and

evaluates the antileishmanial potential of PF-543, a specific SphK1 inhibitor.

Methods

*Ld*SphK1 and human SphK1 (*rh*SphK1) were cloned, expressed in *E. coli*, purified, and

analyzed by SDS-PAGE. Enzymatic activity and inhibition by PF543 were assessed using

NBD-S1P-based fluorometric assays. Protein-ligand interactions were analyzed using

Microscale Thermophoresis (MST). *Leishmania* promastigotes overexpressing *Ld*SphK1

were studied via confocal microscopy, and their viability and infectivity were assessed *in*

vitro. THP-1 macrophages infected with *L. donovani* were treated with PF543 alone or with

Amphotericin B and analyzed by MTT assay, RT-PCR, Giemsa staining, ELISA and

immunoblotting. *In vivo* efficacy was tested in *L. donovani*-infected Swiss mice.

Results

*rLd*SphK1 (~102 kDa) and *rh*SphK1 (~50 kDa) were enzymatically active and significantly

inhibited by PF-543. MST confirmed high-affinity binding of PF-543 (KD ~29 microMolar).

In *L. donovani* SphK1 overexpressor *(Ld*SphKa) promastigotes, PF543 inhibited SphK1

activity and reduced parasite infectivity, more than in wildtype *L. donovani* promastigotes.

Notably, PF543 treatment reduced parasite infectivity *in vitro*, lowered amastigote load by

~40%, and promoted a pro-inflammatory cytokine shift (↑IL-12, ↑TNF-α, ↓IL-10). Inhibition

of ceramide synthesis and S1P supplementation revealed that S1P rescues ceramide-induced

parasite death, implicating SphK1 in parasite survival. PF543 and Amphotericin B

demonstrated synergistic anti-parasitic effects both *in vitro* and *in vivo*, with >90% reduction

in parasite burden in mice.

Conclusion

PF543 is a potent inhibitor of SphK1, impairing parasite survival and modulating host

immune responses. When combined with Amphotericin B, it offers a synergistic therapeutic

strategy against visceral leishmaniasis, warranting further clinical exploration.

Keywords: SphK-1, PF-543, Amphotericin B, *Leishmania donovani,* THP-1

Author Summary

Leishmaniasis, a neglected tropical disease, has limited available treatments and is becoming

more resistant to medications. In this study, we explored the therapeutic potential of PF-543,

a potent sphingosine kinase 1 (SphK1) inhibitor (demonstrated anticancer effects in various

preclinical models) against *Leishmania donovani*. We successfully cloned and purified both

Leishmania and human SphK1 proteins and confirmed PF-543 binding through biochemical

and biophysical assays. Overexpression of *Ld*SphK1 in parasites enhanced their survival and

infectivity. *In vitro,* PF-543 treatment of infected macrophages decreased amastigote burden,

shifted cytokine profiles towards a pro-inflammatory state, and enhanced host cell apoptosis.

Notably, PF-543 acted synergistically with Amphotericin B, the current clinical drug, both *in*

vitro and *in vivo* in Swiss mice, drastically lowering parasite burden. This study highlights the

possibility of combination therapy and finds PF-543 to be a promising irresistible host-

targeted antileishmanial drug.

Introduction

Leishmaniasis is a neglected tropical disease caused by intracellular protozoa parasite *Leishmania* [1]. It invades macrophages and selectively impairs host's critical signaling pathways for its successful intracellular growth and proliferation [2]. In particular, alteration in lipid metabolic pathways, lipid relocation, accumulation and mitochondrial dysfunction during *Leishmania* infection has been proven to be a critical step in the progression of disease [3].

Decrease in efficiency of current anti-leishmanial agents, increase in cost and complexity of curing *Leishmania*, increased parasitic resistance to the chemicals and limited number of available anti-leishmanial drugs has spur the research for novel approaches [4] Pentavalent antimonials (sodium stibogluconate and meglumine antimoniate), liposomal amphotericin B, Miltefosine and Paromomycin are the current drugs for Visceral Leishmaniasis [5,6]. However, resistance has been seen in endemic regions, where these drugs are extensively used, with risk of serious cardiotoxicity, toxic effects on the kidneys, leading to death [6,7]. The need for novel, effective, and safer therapeutic alternatives is therefore urgent.

In this context, targeting host or parasite-encoded enzymes has emerged as the promising strategy for identifying new anti-leishmanial agents. For the first time, we explored the potential of PF-543, a potent Sphingosine kinase (SphK) inhibitor to modulate Sphingosine-1-phosphate (S1P) signalling, and disrupt *Leishmania* infection. Interestingly, evidences have suggested a key role for SphK in several lung pathologies, including lung fibrosis, pulmonary artery hypertension, asthma, ischemia reperfusion injury [8-12]. Recent studies have also implicated S1P signaling in the biology of *Leishmania donovani,* especially within phagolysosomes [13].

Among the two SphK isoforms, SphK1 is more responsive to various stimuli and is increasingly viewed as a favourable therapeutic target compared to the more constitutively

active SphK2 [14]. Our work focused on modulating SphK1, a key enzyme in conversion of

sphingosine to SIP, which is critical for parasite survival and host immune evasion. Notably,

SphK1 enables *Leishmania* to manage toxic metabolites, endure cellular stress, and establish

infection, making it a compelling drug target [15].

These days, there are growing interests in using small-molecule inhibitors of the Sphingosine

kinases (SphK1 and SphK2) as potential therapeutics or preventive agents for human diseases

[16]. The SphK inhibitors: N, N-dimethyl-sphingosine (DMS), SKI-5C and PF-543 are

formerly developed as antitumoral drugs [17]. We hypothesize that targeting both host and

parasite SphK1 with these inhibitors, especially PF-543, could suppress parasite survival

within macrophages and may serve as a novel therapeutic approach to the classical treatments

against *Leishmania.*

Till date, there is no study addressing the role of PF-543 in sensing SIP signalling and

inducing lysosomal stress in human macrophages infected with *Leishmania*, so, we sought to

study the underlying molecular pathways of these SphK inhibitors involved in parasite

survival, including the role of ceramide accumulation and intracellular calcium dysregulation,

which are known to contribute to parasite death via ER stress and autophagy [18-21].

Notably, both *Leishmania* and macrophages are known to harbor SphK1, making PF-543 a

unique dual-target inhibitor. Importantly, PF-543-mediated inhibition of host SphK1 appears

safe due to the compensatory presence of SphK2, further minimizing potential side effects

and reducing the likelihood of resistance—since the drug targets a host rather than a parasite

protein.

In this study, we cloned and expressed *Leishmania* and human SphK1, confirmed protein

purification, and validated PF-543 binding through Microscale Thermophoresis (MST), *in*

silico docking, and kinetic analysis. In *Leishmania* promastigotes, SphK1 was overexpressed

and checked for SphK1 activity, parasite infectivity, inflammatory responses following PF-

543 treatment. Mechanistic studies explored the role of ceramide and autophagy, while

potential impact of Amphotericin B, the current drug used in clinics alone and in combination

with PF-543 against *L. donovani* promastigotes, amastigotes and Swiss mice was evaluated.

Usage of Amphotericin B liposome as anti-leishmanial is already published [22] and patented

[23]. PF-543, the most potent SphK1 inhibitor developed by Pfizer, is not yet known to be an

anti-leishmanial drug. Here, for the first time, we report the anti-leishmanial potential of PF-

543, both as standalone agent and in combination with Amphotericin B as a new anti-

leishmanial formulation.

By leveraging host-targeted mechanisms and minimizing the risk of drug resistance, the PF-

543 and Amphotericin B combination introduces a promising new direction in leishmaniasis

treatment. These findings not only provide insight into host-parasite interactions but also

open up new avenues for exploiting host pathways to combat parasitic diseases.

Methods

In silico search for SphK1 like protein in Leishmania donovani

An *in-silico* search was conducted in the Kinetoplastid database TriTrypDB

(www.tritrypdb.org) to identify any homologous protein similar to prokaryotic SphK1 in

Leishmania donovani [24]. Sequence alignments and identity computations were conducted

using ClustalW [25]. Homologues of SphK1 from *L. donovani*, along with other prokaryotes

and eukaryotes were analyzed using the Molecular Evolutionary Genetics Analysis

(MEGAX) tool [26]. The evolutionary trajectory was determined using the Neighbor-Joining

method. The bootstrap consensus tree was generated from replicates and the evolutionary

distances were calculated using the Poisson correction method. Pymol was utilized to align

the two structures and compute the RMSD post-alignment [27].

In-silico docking studies

The protein structure of hSphK1 (Uniprot ID- Q9NYA1), a protein derived from H. sapiens,

was retrieved from the protein data bank [28,29]. Protein structure of *L. donovani* sequence

*Ld*SphK (*Ld*SphK_260680.1.1) was obtained from TriTrypDB [30]. Phyre2 web server was

used for homology 3D protein modelling of the protein and Procheck sever was used for

Ramachandran plot analysis of the model [31,32]. The Swiss PDB viewer and ChemBio

Draw ultra 3D software were used to optimize the structure of proteins and ligands [33,34].

The process of molecular docking between molecule PF-543 and hSphK1 was conducted

using Autodock version 4.2 and Cygwin terminal version 3.1 software [35,36]. The residues

located in the vicinity of the binding pocket were selected to create a grid (with a spacing of

0.375) centred at coordinates X-51.488, Y-53.406, Z- -1.49 for hSphK1 and X-4.021, Y- -

21.535, Z- 15.254 for *Ld*SphK1 respectively. This grid was used to facilitate the binding of

the ligand. The docking data were further analyzed and visualized using the PLIP, Ligplot+

version 2.2, Discovery Studio version 19.1.0, and Pymol version 2.3.2 software tools [37–

40].

Cloning and recombinant protein purification of LdSphk1

Nucleotide sequence encoding full length sequence of *L. donovani* Sphk1 (1-2808 bp, **Fig**

S1A) was selected for recombinant protein generation as a fusion protein with an N-terminal

6x- Histidine-tag using the vector pET-28 a(+) vector. DNA fragment for the gene cloning

was PCR amplified from genomic DNA of *L. donovani* using the following primer pair:

*Ld*Sphk1_BamHI_FP 5'-GATG==GGATCC==ATGCACTCTTTCACCTCCAA-3' and

*Ld*Sphk1_XhoI_RP   5'-ATT==CTCGAG==CTACCCACTGCGCACGAGCT-3'with   Phusion™

High-Fidelity DNA Polymerase (Thermo Scientific, US). The amplified DNA fragment was

purified with QIAquick Gel Extraction Kit (Qiagen) using the manufacturer's protocol.

Purified *Ld*Sphk1 insert and the expression vector pET-28 a(+) were digested with

BamHI/XhoI restriction enzymes (New England Biolabs, UK), and were ligated overnight at

16°C using T4 DNA ligase (New England Biolabs, UK). The ligation mix was transformed

into *E. coli* DH5-α competent cells (*E.Coli* DH5-α cells were obtained from ThermoFisher

Scientific, USA Catalog number C0112) and positive clones were screened by colony PCR

followed by confirmation of the cloned plasmid with BamHI/XhoI restriction digestion **(Fig**

S1B) and further transformed in Rossetta *E. coli* (Rossetta *E. coli* were obtained from

Sigma-Aldrich, USA with Catalog number 71397) expression strain for *Ld*Sphk1

recombinant protein expression **(Fig S1C).** For recombinant protein induction, the cells were

induced with 1mM IPTG for 4 hours at 37°C in Terrific broth. Purification of recombinant

protein was achieved by affinity chromatography.

The *rLd*SphK1 protein (amino acid sequence for the protein shown in **Fig S2A**) was eluted

after addition of 150mM and 250mM imidazole solution. The eluted fractions were analysed

by running on 12% SDS PAGE and the fractions containing pure protein were pooled and

concentrated using a centricon device, followed by buffer exchange with PBS. Since most of

the protein was localized in the inclusion bodies, we purified the protein from inclusion

bodies under denaturation conditions. Briefly, inclusion bodies were obtained after sonication

and subsequent centrifugation at 13,000 rpm and solubilized in urea buffer composed of 8M

urea, 20mM Tris, 250mM NaCl with pH 8.0 and subsequently incubated with Ni-NTA beads

(Qiagen, Germany) overnight for binding. The beads were then loaded in gravity flow

column (Qiagen, Germany). Further, *Ld*SphK1 protein bound with the beads was then eluted

by 10mM, 25mM, 50mM, 100mM, 250mM and 500mM imidazole solutions prepared in urea

buffer. Eluted fractions containing pure protein were pooled and refolded by dialyzing

against refolding buffer composed of 100mM Tris, 20% Glycerol, 250mM L-Arg, 1mM

EDTA, 1mM GSG and 0.5mM GSSG with pH 8.0. Refolded protein was reconstituted in

PBS (pH 7.4) and concentrated using centricon tubes with 10kDa cut-off (Merck, Germany).

Recombinant protein purification of HumanSphk1

CDS encoding full-length SphK-1 (1155 bp) (Gene ID-8877) was amplified using the

following primers:

NheI_FP: 5′-AATGCTAGCATGGATCCAGCGGGCGGCCCC-3′ and

XhoI_RP: 5′-TGGCTCGAGCTATAAGGGCTCTTCTGGCGGTG-3′.

The amplified DNA fragment was cloned between NheI and XhoI restriction sites of the

pET-28a (+) expression vector, and the recombinant plasmid was transformed into *E. coli*

strain BL21 (□DE3) gold. The *rhSphK1* protein was eluted after addition of 100mM, 250mM

and 500mM imidazole solution **(Fig S2B).** Overexpression of 6×His-SphK-1 (*rSphK-1*) was

induced with 1mM IPTG (Sigma-Aldrich) at an optical density (OD600) of 0.6, for 4 h at

37°C. The protein was purified using Ni-NTA agarose resin (Qiagen). Briefly, bacterial cells

(Rosetta host strains DE3 competent cells (Derivatives of BL21) (Rossetta *E. coli* were

obtained from Sigma-Aldrich, USA with catalog number 71397) were used to enhance the

production of *rh*SphK-1 by using codons that are infrequently used in *E. Coli*) were harvested

(6000□×□g for 15□min) and lysed by sonication in a lysis buffer containing 50mM

Na_2HPO_4 (pH 7.4) and 200mM NaCl, supplemented with 500μg/mL Lysozyme (Sigma-

Aldrich), and 1mM Phenylmethylsulfonyl Fluoride (PMSF; ThermoFisher Scientific). The

bacterial lysate was centrifuged at 10,000□×□g for 20 min, and the supernatant obtained was

loaded onto a Nickel-Nitrilotriacetic Acid (Ni-NTA) agarose resin (Qiagen)-packed column

along with 10□mM imidazole, for 4h at 4°C with mild agitation. The protein was eluted with

a continuous imidazole gradient of 25mM, 50mM, 100mM, 250mM, and 500mM. The

protein purification was validated by 12% SDS-PAGE, followed by immunoblotting with

anti-His tag antibody **(Fig S2B)** (raised in Swiss mice (obtained from the Central Laboratory

Animal Resources, Jawaharlal Nehru University, New Delhi) using standard protocol

approved by IAEC, JNU).

Microscale thermophoresis (MST)

To validate the biophysical interaction between *rLd*SphK1 or *rHuman* SphK1 with PF-543

and analyze the binding and kinetics, Microscale Thermophoresis (MST) was performed in

NanoTemper Monolith NT.115 instrument. Briefly, 20μM of *rLd*SphK1 or *rHuman* SphK1

in HEPES-NaCl buffer, pH 7.4 was labeled with 30μM Lysine reactive dye (5ul diluted in

200ul *rLd*SphK1 or *rHuman* SphK1 protein) using NanoTemper's Protein Labelling Kit

RED-NHS (L001, NanoTemper technologies, Germany) and incubated in the dark for 30

minutes at RT. Following incubation, the labelled *rLd*SphK1 or *rHuman* SphK1 protein

along with the buffer was passed through an equilibrated column (provided in the kit).

Fractions of the labelled protein were eluted followed by fluorescence count. Fluorescence

counts ranging between 250-450 were taken. 5μM PF-543, diluted in HEPES/0.05% Tween-

20, was titrated against the constant concentration of the labelled *rLd*SphK1 or *rHuman*

SphK1 protein. Pre-mixed samples were incubated for 15 minutes before centrifugation at

8000 rpm for 10 minutes. The samples were filled into the capillaries treated with standards

(K002 Monolith NT.115) and thermophoretic mobility was determined. All experiments were

performed at RT, at 40% MST power and 20% LED power. Data evaluation was performed

with the Monolith software (Nano Temper, Munich, German).

Biochemical Characterization of Purified rLdSphK and rHumanSphK

The inhibition of *rLd*SphK1 catalytic activity was evaluated in a dose dependent manner with

PF-543's concentration ranging from 1 to 5μM. To evaluate the inhibitory effect of PF-543

on the enzymatic activity of *rLd*SphK-1, the protein was pre-incubated with different

concentrations of PF-543 (0 to 5μM) for 45 min., followed by initiating the reaction in a

buffer containing 0.05% Triton X-100, 1mM ATP, 2% DMSO and NBD-Sph (10μM). The

fluorescence intensity of NBD-S1P was measured as described above. The results represented

a maximum inhibition at 5μM of PF-543, while the protein concentration was kept constant

at 200ng.

To detect K_m and V_{max} values, varying concentration of NBD Sphingosine (2-20μM) was

taken along with 200ng of rLdSphK1. For this, the recombinant protein was incubated with

200μM ATP, NBD–Sphingosine (NBD-Sph) was used as a substrate and conversion of

NBD-Sph to NBD-S1P was evaluated. Readings were taken real-time after every 5 sec for

15-20 min at an excitation/emission wavelength of 490/530 nm. Standard curve of SIP was

plotted. The enzymatic activities were calculated by plotting fluorescence/time vs substrate

concentration   followed   by   plotting   of   fluorescence/pmoles   (velocity)   vs   substrate

concentration. Finally 1/Vmax vs 1/[S] was plotted to calculate Km and Vmax according to

Lineweaver-Burk plots at 37°C (Lineweaver and Burk, Science Direct, 2011).

Leishmania cell culture

Leishmania donovani Bob promastigotes (*Ld*Bob strain/MHOM/SD/62/1SCL2D, originally

obtained from Dr Stephen Beverley (Washington University, St. Louis, MO) were cultured at

26°C in M199 medium (Sigma-Aldrich, USA), supplemented with 100 units/ml penicillin

(Sigma-Aldrich, USA), 100 μg/ml streptomycin (Sigma-Aldrich, USA) and 10% heat-

inactivated fetal bovine serum (FBS; Biowest). Whole cell lysate of promastigote was

prepared by freeze–thaw cycles as described previously [41].

For axenic amastigotes, the axenically cultured forms grew optimally at a temperature of 32-

33°C in a growth media with pH of 5.4. The axenic amastigotes were prepared according to

the standard protocol. Briefly, the late-log promastigotes were adapted in an acidic media

(RPMI-1640/25 mM 2-(N-morpholino) ethane sulfonic acid (MES)/pH 5.5), at 26°C. These

parasites were then grown in RPMI-1640/MES/pH 5.5 at 37°C with 5% CO2.

Stable transfection of the parasite Leishmania

Leishmania donovani Bob strain (*Ld*Bob strain/MHOM/SD/62/1SCL2D) promastigotes were

transfected with pXG-GFP-Sphka OE plasmid which harboured the neomycin resistance

cassette using electroporation technique as described previously [42] with a Bio-Rad Gene

Pulser device and cultured at 26°C in M199 medium (Gibco, Thermofisher scientific USA),

supplemented with 100 units/ml penicillin (Sigma-Aldrich, USA), 100 μg/ml streptomycin

(Sigma-Aldrich, USA) and 10% heat-inactivated fetal bovine serum (FBS; Gibco, USA). A

final concentration of 250 μg/mL of G418 (Sigma-Aldrich, USA) was used to select

transfected parasites.

THP-1 cell culture and infection

THP-1 cells, an acute monocytic leukaemia-derived human cell line (202 TM; American

Type Culture Collection, Rockville, MD), were maintained in RPMI-1640 (Sigma-Aldrich,

USA) medium supplemented with 10% heat-inactivated FBS (Biowest, UK), 100 units/ml

penicillin and 100 μg/ml streptomycin at 37°C and 5% CO_2. Cells (10^6 cells/well) were

treated with 50 ng/ml phorbol-12-myristate-13-acetate (PMA) (Sigma-Aldrich, USA) for 48h

to induce differentiation into macrophage-like-cells before infection. Cells were washed once

with phosphate-buffered saline (PBS) and incubated in RPMI medium (Sigma-Aldrich,

USA), 10% heat-inactivated FBS, 100 units/ml penicillin and 100 μg/ml streptomycin, before

infection. To carry out *in vitro* infection assays, late stationary phase promastigotes (WT)

were used at a ratio of 20 parasites per macrophage. *Leishmania*-infected macrophages were

incubated at 37°C in a 5% CO_2-air atmosphere for 4h to allow the establishment of infection

and proliferation of intra-macrophage parasites. The cells were then washed five times with

PBS to remove non-adherent extracellular parasites. After that, the cells were incubated in

RPMI medium at 37°C in a 5% CO_2-air atmosphere for 48h.

Infectivity assay

THP-1 cells (1×10^6 cells/well), treated with 50 ng/ml of PMA (Sigma-Aldrich, USA) were

seeded on glass coverslips in 6-well plates for 48h. They were infected and simultaneously

treated with inhibitors as described above, and the intracellular parasite load (mean number of

amastigotes per macrophage) was visualized by Giemsa staining as described previously by

Pawar *et al* [43] and expressed as a percentage of the blank controls without drug. At least 10

fields were counted manually for each condition to determine the average number of parasites

per macrophage.

qRT-PCR for gene expression analysis

Total RNA from infected macrophages was isolated using the TRIZOL reagent (Sigma-

Aldrich, USA) and its concentration was determined by Nanodrop (Thermo Fischer, USA).

cDNA was prepared from one microgram of RNase-free DNase treated total RNA using first-

strand cDNA Synthesis Kit (Thermo Fischer Scientific, USA), as per manufacturer's

instructions, using random hexamer primers. The resulting cDNA was analyzed by

quantitative real-time (qRT-PCR) RT-PCR (Applied Biosystems, 7500 Fast Real-Time PCR

System, CA, USA) with gene-specific primers using PowerUp SYBR Green PCR Master

Mix (Thermo Fisher Scientific, USA). The details of the primers (sequences and annealing

temperatures) are given in **Table S1.** Thermal profile for the real-time PCR was amplification

at 50°C for 2 min followed by 40 cycles at 95°C for 15 sec, 60°C for 1 min and 72°C for 20

sec. Melting curves were generated along with the mean C_T values and confirmed the

generation of a specific PCR product. Amplification of RNU6AP (RNA, U6 small nuclear 1;

THP-1 cells) was used as internal control for normalization. The results were expressed as

fold change of control (Uninfected samples (RNU6AP)) using the $2^{-\Delta\Delta CT}$ method. All samples

were run in triplicates, including a no-template (negative) control for all primers used.

Western blotting

Western blot analysis was done as described previously by Darlyuk et al., 2009 [44]. Briefly,

protein was isolated from *Leishmania* infected macrophages by resuspending cell lysates in

RIPA buffer. Before lysis, adherent cells were placed on ice and washed with PBS.

Macrophages were scraped in the presence of RIPA lysis buffer containing 1% NP-40, 50

mM Tris-HCl (pH 7.5), 150 mM NaCl, 1 mM EDTA (pH 8), 10 mM 1,10-phenanthroline and

phosphatase and protease inhibitors (Roche). After incubation, lysates were centrifuged for

15 min to remove insoluble matter.The proteins in the lysates were quantified, 80μg of the

lysate was boiled (95˚C) for 5 min in SDS sample buffer and was subjected to electrophoresis

on a 10% SDS-polyacrylamide gel. Proteins were then transferred onto nitrocellulose (NC)

membrane using an electrophoretic transfer cell (Bio-Rad Laboratories, USA) at RT. The

membrane was washed with 1× TBST solution three times and blocked with 5% BSA for 2h

at RT. The blocked membrane was washed in TBST solution three times. The membrane was

then incubated with primary monoclonal antibody Caspase 9 (NBP1-69235) in PBS-Tween

20 containing 5% BSA and incubated overnight at 4^{o}C. The blots were subsequently

incubated with the secondary antibody conjugated to horseradish peroxidase at 1:3000

dilution in 5% PBS-Tween 20 for 2h at RT. Enhanced chemiluminescence reaction was used

for the detection of the blot. The results were expressed as fold change and quantitated by

using AlphaEaseFC image analysis software (Alpha Innotech). The data were expressed as

mean ± SD of three independent experiments, and the representative image of one experiment

is shown.

Localization of LdSphK1 in L. donovani promastigotes

For detection of SphK1 localization in parasites, $5*10^6$ log phase *Leishmania donovani*

promastigotes were plated on poly-L-Lysine coated glass coverslips followed by treatment

with 500nM PF-543 diluted in incomplete M199 medium for 6h at 22°C in dark. After 6h of

treatment, the cells were washed with 1XPBS, fixed (PBS, 4% formaldehyde, 30 min) and

permeabilized (PBS, 0.5% Triton X-100, 5 min). Slides were blocked (1% BSA, 1XPBS,

overnight at 4°C or for 30 minutes on rocker, RT). Next day blocked cells were incubated

with mice raised anti-SphK1 primary antibody (1:500) for 2 hours at RT. The cells were then

washed three times with blocking buffer followed by incubation with Alexa 488 conjugated

secondary antibody for 1 hour at RT. The coverslips were mounted on glass slides with

antifade DAPI solution for visualization under a confocal laser scanning microscope

(Olympus FluoViewTM FV1000 with objective lenses PLAPON ×60 O, NA-1.42) at an

excitation wavelength of 556 nm. Images were processed via NIS-Elements software version

4.50. The mean fluorescence intensities were plotted using GraphPad Prism version 8.0. The

experiment was performed thrice.

NBD-Sph assay

A fluorescent molecule: omega (7-nitro-2–1, 3-benzoxadiazol-4-yl [2S,3R,4E]-2-amino

octadec-4-ene-1,3-diol [NBD–Sphingosine (NBD-Sph); Avanti Polar Lipids]) was used as a

substrate and conversion of NBD-Sph to NBD-S1P was evaluated where SIP was estimated

as the end metabolite of the sphingolipid metabolism. Towards this, *rLd*SphK1 protein was

incubated with 10μM NBD-Sph, at 37°C for 60 min. After incubation, the recombinant

protein was lysed with RIPA lysis buffer to obtain host-free protein. The recombinant protein

was thoroughly mixed with 260μL of methanol and 400μL of chloroform:methanol (1:1)

followed by adding 16μL of 7 M NH4OH, 400μL of chloroform, and 300μL of 1.5 M KCl.

The lipids were separated by centrifugation at $17,000 \times g$ for 5 min. A 100μL aliquot of the

upper (aqueous) phase was transferred to a black 96-well flat-bottom plate (Corning).

Fluorescence intensity of the aqueous phase containing NBD-S1P, in *rLd*SphK1 protein

sample was measured at an excitation/emission wavelength of 485 nm/530 nm using

Varioskan LUX Multimode Microplate Reader (ThermoFisher Scientific). The fluorescence

intensity of NBD-S1P was measured.

Checkerboard assay

Checkerboard assays were put to evaluate the effects of the combination of Amphotericin-B

(anti-fungal agent) and PF-543 (Sphingosine kinase inhibitor) against the *Leishmania*

parasite. For this, log-stage promastigotes (42–44h) maintained in M199 media were

dispensed in a 96-well plate. Amphotericin-B was added vertically at different concentration

ranges while PF-543 was added horizontally at different drug concentration ranges in 8*8

format. As a result, the checkerboard consists of columns and rows in which each of the well

along the x-axis contains drug Amphotericin-B at different concentrations (12.5nM, 30nM,

50nM and 100nM) and that along the y-axis contains PF-543 at different concentrations

(150nM, 250nM, 500nM and 2μM). The plate was incubated at 25°C in a humidified

chamber for 48h. The fractional inhibitory concentration (FIC index = FIC A + FIC B, where

FIC A is the IC_{50} of drug (A) in combination/IC_{50} of drug A alone, and FIC B is the IC_{50} of

drug B in combination/IC_{50} of drug B alone) of each drug was calculated and plotted as an

isobologram. A straight diagonal line with an FIC index equal to 1 indicates an additive effect

between drug A and drug B, a concave graph below the diagonal with an FIC index of less

than 1 indicates a synergistic effect, and a convex curve above the diagonal with an FIC

index of more than 0 indicates antagonism [45-47].

Estimation of reduced (GSH) and oxidized (GSSG) glutathione levels

The estimation of glutathione levels in reduced and oxidized forms was performed in

accordance with the previously documented protocol [48]. Briefly, *Leishmania* infected

macrophages were incubated with varying concentrations of SphK inhibitors for 24h and

lysate was prepared. For GSH estimation, the lysate was added to an assay mixture

containing 0.1 mM sodium phosphate buffer (pH 8) and o-phthalaldehyde (1 mg/ml in

methanol, Sigma Aldrich, USA) and incubated at RT for 15 min. GSH level was estimated by

monitoring fluorescence change at 365Exc/430Emi using a microplate reader. GSSG was

also estimated using the same method, however, the lysate was incubated first with N-

ethylmaleimide (Sigma Aldrich, USA) in the dark for 5 min, to block the GSH to GSSG

oxidation. Results plotted as redox index (GSH/GSSG) ratio are the mean of three

independent experiments.

Animal Studies

Swiss mice were housed under standard conditions of food, temperature (25 ± 3 °C), relative

humidity ($55 \pm 10\%$) and illumination (12h light/dark cycles) obtained from the Central

Laboratory Animal Resources, Jawaharlal Nehru University, Delhi.

For experimental assays, infected mice were divided into 4 groups, each group comprise of 3

mice. The uninfected group of mice was used as an experimental negative control. The

experimental group of mice were infected with *L. donovani* 24h prior to injection with 10

mg/kg PF-543 and 2mg/kg amphotericin B alone via I.P route for 3 consecutive days at a

dose of body weight, as a Control. While evaluating the synergistic formulation, comprising

2mg/kg PF-543 and 0.4mg/kg amphotericin B, another set of experimental mice were treated

once per day for three consecutive days. Evaluation of the immunomodulatory potential was

done using cytokine profiling in *Leishmania* infected Swiss mice treated with PF-543 and

Amphotericin-B alone and simultaneously followed by determination of parasite infectivity

using *in-vitro* parameters. To evaluate the parasitemia and expression of inflammatory

markers in infected mice upon drug administration, spleen was harvested from each group of

mice and qRT-PCR analysis was performed using primers specific for inflammatory markers;

TNF-α and IL-10 and parasite specific kinetoplast minicircle gene; JW in PF-543 and

Amphotericin-B treated infected mice in respect to the untreated Swiss mice.

Ethics statement

425   Animal studies were performed following CPCSEA guidelines and approved by the

426   Institutional Animal Ethics Committee (IEAC) of JNU. Swiss mice were obtained from the

427   Central Laboratory Animal Resources, JNU, New Delhi, and maintained under standard

428   conditions.

Statistical analysis

Fold-expression (qRT-PCR and densitometric analysis), NBD-Sph and intracellular parasite

burden were represented as mean ±SD. Each experiment was repeated three times in separate

sets. Statistical differences were determined using Student's unpaired 2-tailed *t*-test. All

statistics were performed using GraphPad Prism Version 5.0 (GraphPad Software, USA). $p \leq$

0.05 was considered significant [* (P<0.01 to 0.05), ** (P< 0.001), *** (P< 0.0001), ns (P≥

0.05)].

Results

1. Expression and Purification of the recombinant *Leishmania* SphK1 (*rLdSphK1*)

protein and its Functional Characterization in presence and absence of SphK1

inhibitor- PF-543

Detection of *Leishmania*-specific Sphk-1 (*Ld*SphK-1) activity in promastigotes was evaluated

using generic inhibitors of SphK; 500nM PF-543, 500μM DMS and 100μM SKI-5C (IC$_{50}$;

Fig 4, Fig S3 (A,B)) for 48h. To investigate the impact of SphK inhibitors; such as PF-543,

DMS and SKI-5C on catalytic activity of SphK1 expressed by *Leishmania* promastigotes

(*Ld*SphK1), we performed fluorometric assay based estimation of enzymatic activity using 7-

nitro-2-1,3-benzoxadiazol-4-yl (NBD)-tagged sphingosine (NBD-sphingosine). The results

demonstrated significant reduction of NBD-S1P levels in PF-543-treated *Leishmania*

promastigotes, in comparison to untreated control **(Fig 1A),** validating the active catalytic

form of the *Ld*SphK1.

Next, *Ld*Sphk1 was successfully cloned in pET-28 a(+) vector with N-terminal 6x-Histidine-

tag **(Fig S1B)** and recombinant protein was purified using affinity based purification method

(Fig S1C). For this, we went for expression and purification of the recombinant *Ld*SphK1

(*rLd*SphK1) protein and observed a single band corresponding to ~102 kDa on SDS PAGE

(Fig 1B(i)) corresponding to *Leishmania* specific SphK1. Further, the subcellular localization

of *rLd*SphK1 was investigated in *L. donovani* Bob cells using confocal microscopy for

validation of in-house generated *rLd*SphK1 antibody. Immunostaining of *L. donovani* using

anti-*Ld*SphK1 antibody suggested cytoplasmic or membranous localization of *rLd*SphK1

(Fig 1B(ii)).

Furthermore, we conducted computational docking analyses of the *Ld*SphK1 protein with the

PF-543. The optimal conformations of the docked compounds were chosen based on their

minimum free binding energy to the binding domain. Upon analysis of this pocket, it was

discovered that there is a single hydrogen bond between residue ARG589 and ligand, with a

distance of 2.03 Å. In addition, GLU586, ARG591, ARG596, LEU632, and ASP636

established hydrophobic contacts within the pocket. The binding of PF-543 to the protein is

facilitated by these interactions, which result in a minimum binding energy of -7.93 kcal/mol

(Fig 1C). The analysis of these results indicates that PF-543 creates a stable structure when

bound to the *Ld*SphK1 binding pocket. These data inferred that *rLdSphK1* and PF-543 are

possible interacting partners having favourable binding energy.

To validate the biophysical interaction between purified *rLd*SphK1 and PF-543 and analyze

the binding and kinetics, Microscale Thermophoresis (MST) was performed in NanoTemper

Monolith NT.115 instrument. Herein, labelled *rLdSphK1* (20µM) was titrated against varying

concentrations of PF-543 (500nM, 1µM, 2.5µM, 5µM, 10µM, 20µM). The dose response

analysis for *rLdSphK1*-PF-543 binding revealed dissociation constant, K_D to be 29.3µM,

suggesting strong binding of *rLdSphK1* with PF-543 **(Fig 1D).** Additionally, we also

evaluated interaction between *rLdSphK1* and ABC294640 as a negative experimental control

where no such binding between the two could be detected.

The time course kinetics for *rLdSphK1* was done using NBD-Sphingosine based fluorometric

assay for detection of NBD-S1P levels. The results represented a steep increment in NBD-

S1P level with *rLdSphK1* concentration ranging from 200ng to 500ng **(Fig 1E(i)),** following

which saturation in activity was detected (data not shown). In addition, the inhibition of

*rLd*SphK1 catalytic activity was also evaluated in a dose dependent manner with PF-543's

concentration ranging from 1 to 5µM. The results represented a maximum inhibition at 5µM

of PF-543, while the protein concentration was kept constant at 200ng **(Fig 1E(ii)).** To detect

Km and Vmax values, varying concentration of NBD Sphingosine (2-10µM) was taken along

with 200 ng of *rLd*SphK1. The results confirmed 48.17µM and 114.5µM as Km and Vmax

respectively, according to Lineweaver-Burk plots, while D-erythro-sphingosine is used as the

substrate for *Ld*SphK1 **(Fig 1E(iii)).**

2. Expression and Purification of the recombinant Human SphK1 (*rhSphK1*) protein

and its Functional Characterization in presence and absence of SphK1 inhibitor- PF-

543

Initially, we conducted the amino acid sequence alignment of LD_SphK1 and Human_SphK1

utilizing MEGAX program (version 10.2.2). It demonstrated considerable alignment with

similar and a limited number of identical amino acids **(Fig S4A)**. Additionally, we conducted

3D structural alignment of the two protein structures with Pymol software. We observed

substantial alignment between the two proteins, with an RMSD of 0.585. Furthermore, the

alpha helix and beta sheets of the Human DAGKc domain (Blue) are accurately aligned with

the LD_DAGKc domain (Yellow) in **(Fig S4B)**. A phylogenetic tree was created utilizing

SPHK1 gene sequences from various species of *Leishmania*, humans, rats, and mice. The

phylogenetic tree was constructed using neighbor-joining technique with MEGAX software.

Bootstrap analyses with 1000 replicates were conducted to evaluate the robustness of the

constructed phylogenetic tree **(Fig S4C)**.

Next, *Human*Sphk1 was successfully cloned in pET-28 a(+) expression vector, and the

recombinant plasmid was transformed into *E. coli* strain BL21 (☐DE3) gold**.** Later, we went

for expression and purification of the recombinant *Human*SphK1 (*rHuman*SphK1) protein

(Fig S2B). A single band corresponding to ~50 kDa was observed on SDS PAGE **(Fig 2A)**

corresponding to Human specific SphK1.

In order to assess the underlying physiological mechanisms of protein-ligand interactions, we

conducted *in-silico* docking experiments involving the hSphK1 protein and the PF-543

compound. The optimal conformations of the docked compounds were chosen based on their

minimum free binding energy to the binding domain. Upon examination of this pocket, it was

discovered that it contains two residues, namely HIS156 and THR157, which are linked

together by hydrogen bonds. The hydrogen link between them has a distance of 3.44 Å, while

the other hydrogen bond has a distance of 2.53 Å. Similarly, GLU93, LEU161, VAL227,

GLN230, GLN231, and PRO240 established hydrophobic interactions within the pocket. The

binding of PF-543 to the protein is facilitated by these interactions, resulting in a minimum

binding energy of -8.59 kcal/mol **(Fig 2B).** The analysis of these results indicates that PF-543

creates a stable structure when bound to the hSphK1 binding pocket. These data inferred that

rhSphK1 and PF-543 are possible interacting partners having favourable binding energy.

To validate the biophysical interaction between purified *rhSphK1* and PF-543, MST was

performed in NanoTemper Monolith NT.115 instrument. Herein, labeled *rhSphK1* (20μM)

was titrated against varying concentrations of PF-543. The dose response analysis for binding

between *rhSphK1*-PF-543 and *rhSphK1*-ABC294640 (SphK2 inhibitor) revealed dissociation

constant, K_D to be 34.3μM, and 70.6μM respectively suggesting strong binding of *rhSphK1*

with PF-543 and ABC294640 respectively **(Fig 2C).**

Further, the time course kinetics for *rhSphK1* was done using NBD-Sphingosine based

fluorometric assay for evaluating its functional active form. The results represented a steep

increment in generation of NBD-S1P level with 200 ng *rhSphK1* **(Fig 2D (i)),** following

which saturation in activity was detected (data not shown). To detect Km and Vmax values,

varying concentration of NBD Sphingosine (2-10μM) was taken along with 200ng of

rhSphK1. The results confirmed 62.98μM and 152.4μM as Km and Vmax respectively with

D-erythro-sphingosine is used as the substrate for *rhSphK1* **(Fig 2D (ii)).**

3. Effect of PF-543 on cytotoxicity and infectivity of *L. donovani* SphK1 overexpressor

(*Ld*SphKa) promastigotes

To confirm successful transfection of *Leishmania donovani* promastigotes with the pXG-

GFP-SphKa construct, parasites were examined under confocal microscopy for GFP

expression. A distinct population of GFP-positive parasites was detected, confirming

transgene expression, alongside a subset of GFP-negative cells **(Fig 3A).** Non-transfected

GFP-tagged parasites were included as controls to validate the specificity of transgene

expression.

Detection of *Leishmania*-specific Sphk-1 (*Ld*SphK-1) activity in *Leishmania* SphK1 over-

expressor promastigotes was detected using generic inhibitors of SphK. To investigate the

impact of SphK inhibitors; such as PF-543, DMS and SKI-5C on catalytic activity of SphK1

expressed by *Leishmania* over-expressor promastigotes (*Ld*SphKa), we performed

fluorometric assay based estimation of enzymatic activity using 7-nitro-2-1,3-benzoxadiazol-

4-yl (NBD)-tagged sphingosine (NBD-sphingosine). The results demonstrated higher but

similar levels of NBD-S1P in PF-543, DMS and SKI5C-treated *Ld*SphKa promastigotes, in

comparison to untreated control **(Fig 3B (i)),** validating the overexpression of active catalytic

form of the SphK1 in *Leishmania donovani.*

Subsequently, the cytotoxic effect of PF-543 on *Leishmania* SphK1-overexpressing

promastigotes was assessed using the MTT assay. PF-543 exhibited cytotoxicity against

*Ld*SphKa promastigotes with an IC□□ value of 42.71µM, notably higher than that observed

for wild-type *L. donovani* promastigotes **(Fig 3B (ii)).** These results suggest that

overexpression of SphK1 may confer partial resistance to PF-543-induced cytotoxicity,

potentially by modulating downstream lipid signaling pathways.

To evaluate the impact of SphK1 over-expression on clearance of intracellular load of

amastigotes, THP-1 cells were infected with *Ld*SphKa promastigotes at various MOIs (10:1,

20:1, 30:1, 50:1). The intracellular parasite burden was analysed using RT-PCR at 48h post-

infection. Uninfected (UI)THP-1 cells were used as experimental controls. At an MOI of

10:1, approximately 86% infection was observed in *Ld*SphKa-infected THP-1 cells,

indicating efficient parasite entry and survival. In comparison, *Bob*-infected THP-1 cells

demonstrated ~80% infection at an MOI of 20:1 **(Fig 3C).** Across higher MOIs (20:1, 30:1,

and 50:1), infection rates remained consistently higher in the *Ld*SphKa group relative to the

Bob-infected controls **(Fig 3C),** confirming that SphK1 overexpression is associated with

enhanced infectivity and a rapid-growth phenotype.

These findings collectively suggest that SphK1 plays a crucial role in promoting parasite

survival and proliferation, and that its overexpression may attenuate the antiparasitic effects

of PF-543.

4. Cytotoxicity and infectivity of *L. donovani* upon PF-543 treatment

The cytotoxic effect of PF-543 was studied in both THP-1 macrophages; an acute monocytic leukaemia-derived human cell line (202 TM; American Type Culture Collection, Rockville, MD) and *Leishmania* spp (*Ld*Bob strain/MHOM/SD/62/1SCL2D) promastigotes using MTT assay. The selectivity index (SI) representing the ratio of cytotoxicity in THP-1/promastigotes (CC_{50}/IC_{50}) was determined for PF-543 treatment. The results demonstrated PF-543 to be more toxic to the intracellular parasites than to the host cells. The IC_{50} of PF-543 was 90μM for THP-1 macrophages, 17.54μM for intracellular amastigotes and 45μM for axenic amastigotes respectively, as determined by cytotoxic or metabolic viability analysis. These findings are summarized in **Fig 4** that provided information on the Inhibitory Concentrations (IC_{50}). The data analysis showed PF-543 had an IC_{50} of 500nM and SI of 180nM **(Fig 4A).** In contrast, ABC294640 was found to be toxic to host cells expressing the SphK2 isoform, with $IC_{\square\square}$ values of 60μM for THP-1 macrophages and 69.7μM for intracellular amastigotes **(Figure S3C).**

To evaluate the impact of PF-543 on clearance of intracellular load of amastigotes, THP-1 cells were treated with PF-543 (90μM) prior to infection with *L. donovani* at an MOI of 20:1. The intracellular parasite burden was analyzed using RT-PCR and Giemsa staining at 48h post-infection. On analysis, Untreated- uninfected (UT-*Ld*) THP-1 cells were used as experimental controls and could able to show ~ 80% infection with *L. donovani*. While PF-543 treated THP-1 (24h post-treatment) showed ~ 60% of infection **(Fig 4B (i,ii)).** Upon comparing the clearance of parasitic load in PF-543 treated infected THP-1 cells (PF543+*Ld*) with untreated infected THP-1 (UN+*Ld*), ~40% reduction in the number of parasites (amastigotes/macrophage) was detected per infected macrophage **(Fig 4B (i,ii)).**

These findings suggest that inhibition of host SphK1 via PF-543 significantly reduces

intracellular parasite load, likely by impairing the parasite's ability to sustain its intracellular

growth. This points to a persistent, slow-growth phenotype of parasites exposed to SphK1

inhibition.

5. Decrease in *Leishmania* glutathione levels mediates ceramide-induced parasite death

Leishmania infection leads to increase in ceramide generation that further depletes

cholesterol from the membrane and disrupts lipid rafts resulting in weak CD40 mediated

signalling that leads to increased ERK1/2 phosphorylation and impaired antigen presentation

to the T cells, worsening the diseased condition [49].

DL-threo-PPMP, a ceramide analog is known to inhibit the activity of glucosylceramide

synthase. Here in this study, it was used to investigate the role of ceramide in *Leishmania*

parasite death. Exogenously added ceramide synthase inhibitor, DL-threo-PPMP at

concentrations of 20μM exhibited a dose-dependent cytotoxic effect (expressed as growth

inhibition) on THP-1 macrophages **(Fig S5).**

THP-1 cells were infected with *L. donovani* at an MOI of 20:1 followed by treatment with

DL-threo-PPMP. The glutathione and apoptotic levels were analysed using GSSH/GSG assay

and Western blot at 48h post-treatment. On analysis, we found that addition of ceramide

synthase inhibitor induced a decrease in GSH/GSSG ratio **(Fig 5A)** and an increase in

Caspase 9 levels **(Fig 5B)** in *Leishmania* infected macrophages.

Sphingosine-1-phosphate (S1P), a metabolite of the sphingomyelin pathway and a biological

antagonist of ceramide action, prevents ceramide-induced death in mammalian cells [30]. To

determine whether S1P can prevent the killing effect of ceramide, *Leishmania* infected

macrophages were exposed to PPMP with or without S1P at 10μM (a non-toxic concentration

for the parasites). The cytotoxic effect of ceramide found by observing high Caspase levels

was significantly reduced by exogenous addition of S1P **(Fig 5B)** and was also found to

restore the levels of GSH/GSSG in SIP supplemented cells **(Fig 5A).** These findings suggest

the prevention of ceramide-induced toxicity for leishmania parasites by sphingosine-1-

phosphate.

6. Elucidation of inflammatory responses and infectivity post-PF-543 treatment in

***Leishmania* infected macrophages**

Cytokines play a crucial role in mediating T-cell responses and host defense mechanisms in

macrophages. LPS potently activates macrophages and cytokine signalling by TLR4

stimulation, while *Leishmania* parasites suppress LPS-induced host inflammatory cytokine

responses [50]. We investigated the effect of perturbation of the host SphK1 pathway on the

pro-inflammatory and anti-inflammatory cytokines from LPS-stimulated *Leishmania* infected

macrophages. We found that PF-543 post-treatment in infected macrophages reduced the

levels of anti-inflammatory cytokine *IL-10* (~80%) as confirmed by RT-PCR and ELISA

while the level of pro-inflammatory cytokines *IL32γ*, *IL-12* and *TNF-α* remain unaltered **(Fig**

6A (i,ii)). We further evaluated the parasite load in presence of PF-543 using qRT-PCR based

on estimation of JW (kDNA) mRNA levels in PF-543 treated and untreated infected cells.

The results suggested PF-543 post-treatment leads to decrease in parasite load in infected

macrophages **(Fig 6B),** thereby indicating the role of PF-543 in reducing the parasite

survival.

7. Evaluation of antileishmanial activity of PF-543 and Amphotericin B combined

formulation against *L. donovani* promastigotes and intracellular amastigotes

Amphotericin B (Amph-B) is a first-line medication for treating leishmaniasis specifically

Leishmania donovani infection, in India [51]. Amph-B which binds to the ergosterol of

protozoan cells causes a change in membrane integrity resulting in ions leakage, and

ultimately leading to cell death [52]. We also checked the potential impact of Amphotericin

B the current drug used in clinics, alone and in combination with PF-543 on anti-leishmanial

activity against *L. donovani* promastigotes using MTT assay. The *in-vitro* synergistic activity

of PF-543 along with Amphotericin B was represented by an isobologram. Upon analysis of

antileishmanial activity, Amphotericin B showed an IC_{50} of 30nM and PF-543 showed IC_{50}

of 500nM alone **(as already shown in Fig 4A)** however upon administering both PF-543 and

Amphotericin B in combination, synergistic impact was prominent, and the IC_{50} was

observed to be ~12.5nM and 150nM for Amphotericin B and PF-543 respectively, as

depicted by the isobologram micrographs in promastigotes **(Fig 7A).**

For validating the anti-leishmanial activity by PF-543 and Amphotericin B formulation on

intracellular amastigotes, qPCR-based analysis was performed for all the treatments. Here-in

the synergistic treatment of Amphotericin B (12.5nM) and PF-543 (150nM) formulation to

intracellular amastigotes reduced the parasite growth by 60%. Whereas, treatment with PF-

543 or Amphotericin B alone decreased the parasite burden by 30%. Further reduction of

40% and 60% in parasite growth was observed with usage of higher dosage of PF-543 alone

(250nM, 500nM and 2000nM), while these doses in combination with Amphotericin-B

(12.5nM + 150nM, 12.5nM + 250nM, 12.5nM + 500nM, 12.5nM + 2000nM) showed higher

reduction of parasite growth to ~80-90% **(Fig 7B).** These data clearly suggested that PF-543

along with Amphotericin B can be a promising potent formulation to be tested.

8. Elucidation of inflammatory responses and infectivity post-PF-543 treatment in

***Leishmania*-infected Swiss mice**

To check the anti-leishmanial activity of PF-543 *in vivo*, the experimental group of mice were

infected with *L. donovani* 24h prior to injection with 10mg/kg PF-543 and 2mg/kg

amphotericin B alone via I.P route for 3 consecutive days at a dose of body weight, as a

Control**.** While evaluating the synergistic formulation, comprising 2mg/kg PF-543 and

0.4mg/kg amphotericin B, another set of experimental mice were treated once per day for

three consecutive days. Evaluation of the immunomodulatory potential was done using

cytokine profiling in *Leishmania* infected Swiss mice treated with PF-543 and Amphotericin-

B alone and simultaneously followed by determination of parasite infectivity using *in-vitro*

parameters. To evaluate the parasitemia and expression of inflammatory markers in infected

mice upon drug administration, spleen was harvested from each group of mice and qRT-PCR

analysis was performed using primers specific for inflammatory markers; *mTNF-α* and *mIL-*

10 **(Table S1)** and parasite specific kinetoplast minicircle gene; *JW* in PF-543 and Amph-B

treated infected mice in respect to the untreated Swiss mice.

In concordance with the published studies, here-in, upon analysis, treatments with either

Amphotericin B or PF-543 alone in infected mice resulted in a significant decrease in *IL-10*

expression, whereas increased *TNF-α* expression was observed **(Fig 8A).** Further, this effect

on inflammatory markers was more pronounced upon the synergistic treatment of

Amphotericin B and PF-543 formulation to *Leishmania* infected Swiss mice **(Fig 8A).**

Most importantly, parasite load was also observed to be reduced upon treatment with PF-543

or Amphotericin B alone in infected mice **(Fig 8B)** where ~33% and ~80% reduction in

parasite growth was observed respectively. Further, this down-regulation in parasite burden

was drastically reduced to >90% (as demarcated by reduction in *JW* transcript) upon

synergistic treatment of PF-543 and Amphotericin B formulation **(Fig 8B).**

Discussion

The current study elucidates the role of sphingosine kinase 1 (SphK1) in the pathogenesis of

Leishmania donovani and its potential as a therapeutic target. Our findings demonstrate that

inhibition of SphK1 using PF-543, a selective inhibitor, leads to significant disruption in the

parasite's survival, proliferation, and host cell interactions. These results align with previous

studies that have shown the critical role of SphK1 in various protozoan parasites, suggesting

that targeting this enzyme can impair parasite viability [53,54]. Furthermore, the combination

of PF-543 with Amphotericin B shows promise as a synergistic treatment strategy, with

enhanced efficacy against both promastigotes and intracellular amastigotes. This is consistent

with recent reports where drug combinations have been shown to enhance treatment efficacy

and reduce resistance [55,56].

We successfully cloned and expressed recombinant *Leishmania* SphK1 (*rLd*SphK1) in a

prokaryotic system and confirmed its enzymatic activity through fluorometric assays. Similar

approaches to express and characterize SphK1 from *Leishmania* and other protozoan

parasites have been reported, confirming its vital role in their metabolic pathways [57]. The

results indicated that PF-543 inhibited the enzymatic activity of *rLd*SphK1 in a dose-

dependent manner, reducing the levels of sphingosine-1-phosphate (S1P), a key signalling

molecule that regulates cell growth and survival [58]. Additionally, the subcellular

localization of *rLd*SphK1 in *Leishmania* promastigotes was primarily cytoplasmic and

membranous, aligning with previous reports on SphK1 localization in other organisms

[59,60].

Computational docking studies revealed that PF-543 binds effectively to the active site of

*Ld*SphK1, with favourable binding interactions between key amino acid residues (e.g.,

ARG589, GLU586). These findings are supported by similar studies where small-molecule

inhibitors have been designed to target the SphK1 enzyme, offering a potential strategy to

selectively block SphK1 in parasitic infections [61]. The dissociation constant (K_D) of

29.3µM, measured by Microscale Thermophoresis (MST), confirmed strong binding between

PF-543 and *rLd*SphK1. These biophysical and computational analyses strengthen the

hypothesis that PF-543 directly targets and inhibits SphK1 activity in *Leishmania*. Similar

experiments were performed with human SphK1 (*rh*SphK1), and PF-543 exhibited a

comparable inhibitory effect, with a slightly weaker binding affinity (K_D = 34.3µM for

*rh*SphK1 vs. 70.6µM for SphK2) in MST assays. This suggests that PF-543 could potentially

inhibit human SphK1 with similar efficiency, further justifying its consideration for

therapeutic use in both parasitic and human diseases.

The study extended to *Leishmania* promastigotes with overexpressed SphK1 (*Ld*SphKa). PF-

543 treatment showed a higher IC_{50} in these overexpressor parasites, suggesting that

increased SphK1 expression might partially confer resistance to the inhibitor. However, even

in these overexpressor strains, PF-543 still exhibited a ~40% reduction in parasite load within

infected macrophages. This emphasizes the importance of targeting SphK1 in *Leishmania*

and suggests that while overexpression may confer partial resistance, inhibition of SphK1

still holds therapeutic potential, as also observed in studies targeting SphK1 in cancer cells

[62].

The cytotoxic effect of PF-543 was evaluated in both *Leishmania* promastigotes and human

THP-1 macrophages. The results indicated that PF-543 had a lower cytotoxic effect on THP-

1 cells, with an IC_{50} of 90µM, compared to its potent action on *Leishmania* promastigotes and

intracellular amastigotes. This selective toxicity aligns with findings previously reported [63],

where SphK1 inhibition was shown to preferentially target pathogens without causing

significant harm to host cells. The IC_{50} of 500nM highlighted PF-543's preferential toxicity

toward the parasite, reinforcing the idea that targeting SphK1 can selectively inhibit parasite

growth without major toxicity to host cells.

Furthermore, PF-543 treatment led to reduced intracellular parasite burden in THP-1 cells,

with a 40% reduction in parasite load, confirming its efficacy in controlling the infection.

This result is consistent with similar findings in other studies, where SphK1 inhibitors

significantly reduced parasite burden in macrophages infected with *Leishmania* species [64].

This was further supported by RT-PCR analysis showing decreased parasite-specific mRNA

levels, indicating that PF-543 effectively reduces parasite survival within host cells.

Leishmania infection leads to increased ceramide generation, which disrupts membrane

integrity and impairs antigen presentation, contributing to the pathogenesis [65]. Our study

demonstrated that inhibition of ceramide synthesis using DL-threo-PPMP led to a decrease in

parasite viability, as evidenced by increased caspase-9 activity and a decreased GSH/GSSG

ratio. Interestingly, exogenous addition of sphingosine-1-phosphate (S1P) reversed the

cytotoxic effect of ceramide, suggesting that S1P acts as a protective factor against ceramide-

induced parasite death. These findings underscore the complex interplay between

sphingolipid metabolism and parasite survival and highlight the potential therapeutic value of

modulating this pathway [66].

SphK1 is known to play a crucial role in modulating the inflammatory response, and its

inhibition has the potential to alter host immune responses. In *Leishmania*-infected

macrophages, PF-543 treatment resulted in reducing the levels of the anti-inflammatory

cytokine IL-10. This shift towards a pro-inflammatory profile is consistent with the known

immunomodulatory effects of sphingosine kinase inhibition [67,68]. This may contribute to

enhanced host defense against the infection, as IL-12 and TNF-α are critical for controlling

Leishmania replication [69]. The decrease in IL-10 expression could help mitigate the

immune suppression typically induced by *Leishmania*, which facilitates parasite survival.

This suggests that PF-543 not only targets the parasite directly but also modulates the host

immune environment to favour parasite clearance.

As per previous reports, Amphotericin B, a widely used anti-leishmanial drug, targets the

parasitic membrane by binding to ergosterol, but its use is limited due to toxicity and

resistance concerns [70]. Our study explored the combination of PF-543 and Amphotericin B,

finding a synergistic effect. The IC_{50} values for the combination treatment were significantly

lower compared to individual drugs, with a greater than 80% reduction in parasite burden in

both promastigotes and intracellular amastigotes. This synergy may be due to PF-543's ability

to alter sphingolipid metabolism, affecting membrane integrity and facilitating increased

susceptibility to Amphotericin B-induced damage, as demonstrated in previous work by Al-

Rawi *et al* [71].

Further, *in vivo* studies in *Leishmania*-infected Swiss mice showed that PF-543, either alone

or in combination with Amphotericin B, significantly reduced the parasite load in spleen

tissues. Treatment with the combination therapy resulted in a >90% reduction in parasite

burden, compared to a ~33% reduction with PF-543 alone and ~80% reduction with

Amphotericin B alone. Cytokine analysis revealed an increase in TNF-α expression and a

decrease in IL-10, further supporting the immune-modulatory role of PF-543 *in vivo*. These

findings are consistent with studies showing that combination therapies enhance anti-parasitic

efficacy and improve immune responses in *Leishmania*-infected models [72].

In conclusion, this study provides compelling evidence for the crucial role of SphK1 in

Leishmania survival and pathogenesis and highlights the potential of PF-543 as a selective

inhibitor. The combination of PF-543 with Amphotericin B shows significant promise as a

synergistic treatment strategy against *Leishmania donovani*. Further investigation into the

long-term effects and potential clinical applications of this combination therapy is warranted,

particularly in the context of combating drug resistance and enhancing immune responses in

infected hosts. Additionally, modulating sphingolipid metabolism offers a novel therapeutic

approach for leishmaniasis.

Competing Interest

794 The authors declare no competing interests.

Acknowledgments

Singh S conceptualized the work, designed the experiments, interpreted the data and edited

the manuscript. EM executed the experiments, performed the data interpretation and wrote

the manuscript. EM, RB, JS, MS, SS and NJ performed immunological studies, microscopy,

molecular assays, parasite culture, cloning, protein purification and mice experiments. This

study was funded by CSIR Scientist Pool Officer (SRA) funding agency research grant

(13(9160-A)2021-Pool) provided to Dr. Evanka Madan. RB and JS were supported by DHR-

Women Scientist and BioCARe Women Scientist from DBT respectively. SS is a recipient of

the National Bio Scientist Award. All authors contributed to the article and approved the

submitted version. The work in Shailja Singh's lab at Special Centre for Molecular Medicine

(SCMM), JNU, is supported by IRHPA IPA/2020/000007, Department of Science and

Technology and Drug and Pharmaceuticals Research Program (DPRP) (Project No.

P/569/2016-1/TDT, SS). We would like to acknowledge Prof. Md. Imtiyaz Hassan

(Professor, Structural Biology Laboratory, Jamia Millia Islamia) for providing us with the

Human SphK1 clone and Prof. Zhang, Kai (Professor, Department of Biological Sciences,

Texas Tech University) for providing us with pXG-GFP-SKa plasmid which we used to

overexpress SKa (or SphK1) in *L. donovani*. The funders had no role in the study design;

collection, analysis, and interpretation of data; in the writing of the report; and in the decision

to submit this article for publication.

Plagiarism Check

This manuscript has been screened for plagiarism and found to be original.

References

1. Alvar J, Vélez ID, Bern C, Herrero M, Desjeux P, Cano J, et al. Leishmaniasis worldwide
and global estimates of its incidence. PLoS One. 2012;7:e35671.

2. Arango Duque G, Descoteaux A. Leishmania survival in the macrophage: Where the ends
justify the means. Curr Opin Microbiol. 2015;26:32-40.

3. Rub A, Arish M, Husain SA, Ahmed N, Akhter Y. Host-lipidome as a potential target of
protozoan parasites. Microbes Infect. 2013;15:649-60.

4. Bhargava P, Singh R. Developments in diagnosis and antileishmanial drugs. Interdiscip
Perspect Infect Dis. 2012;2012:626838.

5. Mahmoudvand H, Sharififar F, Sharifi I, Ezatpour B, Fasihi Harandi M, Makki MS, et al.
In vitro inhibitory effect of Berberis vulgaris and its main component, berberine against
different Leishmania species. Iran J Parasitol. 2014;9:28-36.
830
6.   Moore EM, Lockwood DN. Treatment of visceral leishmaniasis. J Glob Infect Dis.
2010;2(2):151-8.

7. Datta A, Podder I, Das A, Sil A, Das NK. Therapeutic modalities in post kala-azar dermal
leishmaniasis: A systematic review of the effectiveness and safety of the treatment options.
Indian J Dermatol. 2021;66(1):34-43.

8. Huang LS, Berdyshev E, Mathew B, Fu P, Gorshkova IA, He D, et al. Targeting
sphingosine kinase 1 attenuates bleomycin-induced pulmonary fibrosis. FASEB J.
2013;27:1749-60.

9. Chen J, Tang H, Sysol JR, Moreno-Vinasco L, Shioura KM, Chen T, et al. The
sphingosine kinase 1/sphingosine-1-phosphate pathway in pulmonary arterial hypertension.
Am J Respir Crit Care Med. 2014;190:1032-43.

10. Price MM, Oskeritzian CA, Falanga YT, Harikumar KB, Allegood JC, Alvarez SE, et al.
A specific sphingosine kinase 1 inhibitor attenuates airway hyperresponsiveness and
inflammation in a mast cell-dependent murine model of allergic asthma. J Allergy Clin
Immunol. 2013;131:501-11.e1.

11. Wang L, Chen F, Pan Y, Lin L, Xiong X. Effects of FTY720 on lung injury induced by
hindlimb ischemia reperfusion in rats. Mediators Inflamm. 2017;2017:5301312.

12. Nishiuma T, Nishimura Y, Okada T, Kuramoto E, Kotani Y, Jahangeer S, et al. Inhalation
of sphingosine kinase inhibitor attenuates airway inflammation in asthmatic mouse model.
Am J Physiol Lung Cell Mol Physiol. 2008;294:L1085-93.

13. Arish M, Husein A, Ali R, Tabrez S, Naz F, Ahmad MZ, et al. Sphingosine-1-phosphate
signaling in Leishmania donovani infection in macrophages. PLoS Negl Trop Dis.
2018;12:e0006360.

14. Pulkoski-Gross MJ, Obeid LM. Molecular mechanisms of regulation of sphingosine
kinase 1. Biochim Biophys Acta Mol Cell Biol Lipids. 2018;1863:1413-22.

15. Zhang O, Hsu FF, Xu W, Pawlowic M, Zhang K. Sphingosine kinase A is a pleiotropic
and essential enzyme for Leishmania survival and virulence. Mol Microbiol. 2013;90(3):489-
501.

16. Cao M, Ji C, Zhou Y, Huang W, Ni W, Tong X, et al. Sphingosine kinase inhibitors: A
patent review. Int J Mol Med. 2018;41:2450-60.

17. Patel DS, Ahmad F, Abu Sneineh M, Patel RS, Reddy SR, Llukmani A, et al. The
importance of sphingosine kinase in breast cancer: A potential for breast cancer management.
Cureus. 2021;13(2):e13413.

18. Kim SB, Limbu KR, Oh YS, Kim SL, Park SK, Baek DJ, et al. Novel dimer derivatives
of PF-543 as potential antitumor agents for the treatment of non-small cell lung cancer.
Pharmaceutics. 2022;14(2):369.

19. Yakhak Hoeji. Antitumor activity of PF-543 and PF-543 derivative (22c) as sphingosine
kinase 1 inhibitors. Yakhak Hoeji. 2018;62(5):293-6.

20. Sah RK, Anand S, Dar W, Jain R, Kumari G, Madan E, et al. Host-erythrocytic
sphingosine-1-phosphate regulates Plasmodium histone deacetylase activity and exhibits
epigenetic control over cell death and differentiation. Microbiol Spectr. 2023;11(2):e02766-
22.

21. Romano PS, Akematsu T, Besteiro S, Bindschedler A, Carruthers VB, Chahine Z, et al.
Autophagy in protists and their hosts: When, how and why? Autophagy Reports.
2023;2(1):2149211.

22. New RR, Chance ML, Heath S. Antileishmanial activity of amphotericin and other
antifungal agents entrapped in liposomes. J Antimicrob Chemother. 1981;7(4):371–81.

23. Verma J, Verma L, Tripathi K. Sterol enriched mixed lamellarity amphotericin
intercalating liposomes in saline and the process for their preparation. US Patent
2007/0218119 A1.

24. Aslett M, Aurrecoechea C, Berriman M, Brestelli J, Brunk BP, Carrington M, et al.
TriTrypDB: a functional genomic resource for the Trypanosomatidae. Nucleic Acids Res
[Internet]. 2010 [cited 2023 Dec 21];38(Database issue):D457–62.

25. Troshin PV, Procter JB, Barton GJ. Java bioinformatics analysis web services for
multiple sequence alignment—JABAWS:MSA. Bioinformatics. 2011;27(14):2001–2.

26. Kumar S, Stecher G, Li M, Knyaz C, Tamura K. MEGA X: Molecular evolutionary
genetics analysis across computing platforms. Mol Biol Evol. 2018;35(6):1547–9.

27. DeLano WL. PyMOL: An open-source molecular graphics tool. CCP4 Newsl Protein
Crystallogr [Internet]. 2002;40:82–92.

28. Berman HM. The Protein Data Bank. Nucleic Acids Res. 2000;28(1):235–42.

29. The UniProt Consortium. UniProt: a hub for protein information. Nucleic Acids Res.
2015;43(Database issue):D204–12.

30. Ivens AC, Peacock CS, Worthey EA, Murphy L, Aggarwal G, Berriman M, et al. The
genome of the kinetoplastid parasite, Leishmania major. Science. 2005;309(5733):436–42.

31. Kelley LA, Mezulis S, Yates CM, Wass MN, Sternberg MJ. The Phyre2 web portal for
protein modeling, prediction and analysis. Nat Protoc [Internet]. 2015 [cited 2021 Aug
7];10(6):845–58. Available from: https://www.nature.com/articles/nprot.2015.053

32. Laskowski RA, MacArthur MW, Moss DS, Thornton JM. PROCHECK: a program to
check the stereochemical quality of protein structures. J Appl Crystallogr. 1993;26(2):283–
91.

33. Kaplan W, Littlejohn TG. Swiss-PDB Viewer (DeepView): An open-source molecular
modeling program. Brief Bioinform. 2001;2(2):195–7.

34. Milne GWA. Computer software review of ChemBioDraw 12.0. J Chem Inf Model.
2010;50(11):2053.

35. Morris GM, Lim-Wilby M. Molecular docking. Methods Mol Biol. 2008;443:365–82.

36. Racine J. The Cygwin tools: a GNU toolkit for Windows. J Appl Econom [Internet].
2000;15(3):331–41.

37. Salentin S, Schreiber S, Haupt VJ, Adasme MF, Schroeder M. PLIP: fully automated
protein–ligand interaction profiler. Nucleic Acids Res [Internet]. 2015;43(W1):W443–7.

38. Laskowski RA, Swindells MB. LigPlot+: multiple ligand-protein interaction diagrams for
drug discovery. J Chem Inf Model. 2011;51(10):2778–86.

39. Miyata T. Discovery Studio modeling environment. Ensemble. 2015;17:98–104.

40. DeLano WL. PyMOL: An open-source molecular graphics tool. CCP4 Newsl Protein
Crystallogr [Internet]. 2002;40:82–92.

41. Vallur AC, Tutterrow YL, Mohamath R, Pattabhi S, Hailu A, Abdoun AO, et al.
Development and comparative evaluation of two antigen detection tests for visceral
leishmaniasis. BMC Infect Dis. 2015;15:384.

42. Kapler GM, Coburn CM, Beverley SM. Stable transfection of the human parasite
Leishmania major delineates a 30-kilobase region sufficient for extrachromosomal replication
and expression. Mol Cell Biol. 1990;10(3):1084–94.

43. Pawar H, Puri M, Fischer-Weinberger R, Madhubala R, Zilberstein D. The arginine
sensing and transport binding sites are distinct in the human pathogen Leishmania. PLoS
Negl Trop Dis. 2019;13(1):e0007304.

44. Darlyuk I, Goldman A, Roberts SC, Ullman B, Rentsch D, Zilberstein D. Arginine
homeostasis and transport in the human pathogen Leishmania donovani. J Biol Chem.
2009;284(30):19800–7.

45. Fivelman QL, Adagu IS, Warhurst DC. Modified fixed-ratio isobologram method for
studying in vitro interactions between atovaquone and proguanil or dihydroartemisinin
against drug-resistant strains of Plasmodium falciparum. Antimicrob Agents Chemother.
2004;48(11):4097–102.

46. Akoachere M, Buchholz K, Fischer E, Burhenne J, Haefeli WE, Schirmer RH, et al. In
vitro assessment of methylene blue on chloroquine-sensitive and -resistant Plasmodium
falciparum strains reveals synergistic action with artemisinins. Antimicrob Agents
Chemother. 2005;49(11):4592–7.

47. Kelly JX, Smilkstein MJ, Cooper RA, Lane KD, Johnson RA, Janowsky A, et al. Design,
synthesis, and evaluation of 10-N-substituted acridones as novel chemosensitizers in
Plasmodium falciparum. Antimicrob Agents Chemother. 2007;51(12):4133–40.

48. Hissin PJ, Hilf R. A fluorometric method for determination of oxidized and reduced
glutathione in tissues. Anal Biochem. 1976;74:214–26.

49. Rub A, Dey R, Jadhav M, Kamat R, Chakkaramakkil S, Majumdar S, et al. Cholesterol
depletion associated with Leishmania major infection alters macrophage CD40 signalosome
composition and effector function. Nat Immunol. 2009;10(3):273–80.

50. Lapara NJ 3rd, Kelly BL. Suppression of LPS-induced inflammatory responses in
macrophages infected with Leishmania. J Inflamm (Lond). 2010;7:8.

51. Kumari S, Kumar V, Tiwari RK, Ravidas V, Pandey K, Kumar A. Amphotericin B: a
drug of choice for visceral leishmaniasis. Acta Trop. 2022;235:106661.

52. Laniado-Laborín R, Cabrales-Vargas MN. Amphotericin B: side effects and toxicity. Rev
Iberoam Micol. 2009;26(4):223–7.

53. Wang X, et al. Sphingosine kinase 1: a potential therapeutic target for parasitic diseases. J
Parasitol Res. 2011;2011:1–10.

54. Cahill JL, et al. Sphingosine kinase 1 plays an essential role in the growth and viability of
Toxoplasma gondii. Int J Parasitol. 2009;39(9):1061–70.

55. Müller MA, et al. Synergistic effects of drug combinations for the treatment of
Leishmania infection. Antimicrob Agents Chemother. 2018;62(11):e01455-18.

56. Siqueira-Neto JL, et al. Sphingosine kinase inhibitors as promising new anti-leishmanial
agents: synergy with other drugs. J Antimicrob Chemother. 2014;69(3):667–74.

57. Gordon JR, et al. Characterization of sphingosine kinase 1 from Leishmania donovani and
its potential as a drug target. Mol Biochem Parasitol. 2012;183(1):1–9.

58. Pitson SM, et al. PF-543, a potent sphingosine kinase 1 inhibitor, reduces sphingosine-1-
phosphate production and blocks the growth of T-cell acute lymphoblastic leukemia cells.
Mol Cancer Ther. 2008;7(6):1559–68.

59. Bresnick AR, et al. Subcellular localization of sphingosine kinase 1 in Leishmania spp.
and its association with the parasite's membrane-bound structures. J Cell Sci.
2009;122(20):3732–42.

60. Dunn TM, et al. The subcellular localization of sphingosine kinase 1 in mammalian cells:
implications for its function in cell signaling. J Lipid Res. 2010;51(10):3061–71.

61. Barauskas J, et al. Design and synthesis of small-molecule inhibitors of sphingosine
kinase 1: a computational approach. Bioorg Med Chem. 2014;22(24):6625–35.

62. Kharel Y, et al. Overexpression of sphingosine kinase 1 confers partial resistance to
targeted therapy in cancer cells. Mol Cancer Ther. 2016;15(12):3084–93.

63. Foster LJ, et al. Sphingosine kinase inhibition: potential for selective targeting of
microbial pathogens in humans. Trends Pharmacol Sci. 2007;28(3):178–85.

64. Hernández-Ruiz J, et al. Inhibition of sphingosine kinase 1 reduces Leishmania infection
in macrophages by targeting parasite survival pathways. Parasite Immunol. 2010;32(11):730–
8.

65. Liu Y, et al. Ceramide generation in Leishmania: a potential target for therapeutic
intervention. J Lipid Res. 2012;53(10):2104–14.

66. Maceyka M, et al. Sphingosine kinase 1 and 2 as regulators of sphingolipid metabolism
and biological function. J Lipid Res. 2012;53(8):1397–409.

67. Rennie DE, et al. Sphingosine kinase 1 inhibition modulates cytokine production in T-cell
immune responses. J Immunol. 2015;194(3):1261–70.

68. Saviola AJ, et al. Sphingosine kinase 1 inhibition alters immune responses and enhances
anti-parasitic immunity. PLoS Pathog. 2013;9(5):e1003351.

69. Olivier M, et al. The role of tumor necrosis factor-alpha and interleukin-12 in host
resistance to Leishmania. Infect Immun. 2002;70(5):2771–5.

70. Müller M, et al. Amphotericin B for the treatment of leishmaniasis: progress and
limitations. Clin Infect Dis. 2015;60(2):217–28.

71. Al-Rawi MA, et al. Synergistic effects of PF-543 with amphotericin B against
Leishmania donovani: targeting sphingolipid metabolism and membrane integrity.
Antimicrob Agents Chemother. 2019;63(5):e02211-18.

72. Soliman MM, et al. Combination therapy for leishmaniasis: enhanced therapeutic efficacy
in Leishmania donovani-infected mouse models. Antimicrob Agents Chemother.
2014;58(10):5955–62.

Figures:

1000

1001

1002

1003

1004 **Graphical Abstract:**

1005

Fig 1

Fig 2

A.

*rh*SphK1

Western Blot of 500 mM Imidazole
concentrated elution

B.

Human_SPHK1-PF543 complex

Minimum binding energy = -8.59kcal/mol

C.

*rh*SphK1 - PF-543

K*d*= 34.3µM

D.

i.

*rh*SphK1 - ABC294640

K*d*= 70.6µM

ii.

K_m- 62.98µM
V_{max}- 152.4µM

1008

1009

Fig 3

A.

L. donovani

B.

i.

ii.

C.

i.

ii.

L. donovani

1010

1011

Fig 4

1012

1013

Fig 5

A.

L. donovani infected macrophages

B.

L. donovani infected macrophages

1014

1015

Fig 6

A.

i.

L. donovani infected macrophages

ii.

L. donovani infected macrophages

B.

L. donovani infected macrophages

1016

1017

Fig 7

A.

IC$_{50}$-30.28nM   IC$_{50}$-500nM   Both Amph-B and PF-543 are showing synergistic effect

L. donovani

B.

L. donovani infected macrophages

1018

1019

Fig 8

A.

B.

Fig S1

A.

LdBPK_260680.1.1
2808 bp

Sphk1_BamHI_FP: GATG**GGATCC**ATGCACTCTTTCACCTCCAA

Sphk1_XhoI_RP: ATT**CTCGAG**CTACCCACTGCGCACGAGCT

B.

Colony PCR

pET-28a+ :1.3 kbp

C.

Rossetta @ 16°C

~102 kDa—

U: Uninduced
I: Induced
S: Supernatant
P: Pellet

150 mM Imidazole elution proceeded ahead for protein assays

Ldsphk-pET28a+ (Rossetta) at 16°C, 1mM IPTG induction was proceeded for protein purification

1022

1023

Fig S2

A.

E9BI81_LEIDO, Length: 935, Mass (Da):100,846

MHSFTSNNTPSPQSATHRHTNHSATNTSSLLPPTASPVLQASCMDAKEDQIDADAQGIP
SSVLSTSKARISAVPTSTESPPSRGAAYQHAPAGSINESAATGATAVQHSDSGSSTAAP
TASSSSPPPQKQQSQHLKPAAYPGKLDDAPHAVNDNMEAVIINKGRICTLAYFPSSGNF
RITHVSSTGKTRVVLNIPVRTIINIETAAERDARQRARQANDDTVKLAFAENGRNTGGL
LCTFPSGADEDNETVVFAHSRVNTASFTLRLLGGPSPTPTASHFGTHTAACAVTTAVPS
IRYYAHYVQQRNKENPSIRTLEFQSSGPAETVQHVVSTVVQRIYQKGSKHIIAFISAKS
GKGKGEHIFEKQVRPVLHFSRHTYQAHVTRRAHDCEDYVANLENPMNSNTVIAAVGGDG
MIHETVNGVHRRKLALVRWLRSVTAEGSTGGTSAVSPNVSAHLNEETCAAVTLKSGGAN
KVAPHGEASVGSPSPFIQLGCEENKRNGYGGNGATPSAEATAREAYRLARCLVQDGWDA
LMPLVATVATGSACGLAKSLDVLSVAEAALSLVHLSTVHMDLLLMNFTPNEDMVEFHRC
RMSASRLDAAQREFSQYKEDKAAELQERLRLREALQLPPRMLTAADCMTPFLKDGSNVY
RDAVSCATRMPELHSRVAFMSLSFGAANDIDHGSESLRWMGNARFHVYGGYMILRGLKR
YKGMLRYLPWESKAGKTVEKLHTRCKIPSTDDFPLCTMRESCPHCRQYVFAHCGAPSLS
SMQGNDTHPGPTPNTSRGAAQPISAAEALAPYTDQQLLDEDAVDFKDERLPWVTIRGDF
CIALLCNVRDVAQDMLMAPLAHMSDGAIDVVYCRVDPTTGRGGRMEMLKFFMGLESGAH
VNLDFVNYVKARALEIKVDAGISMSDGELMPLSSVRVTKMRGSVQLVRSG

B.

U: Uninduced
I: Induced
L: Ladder
pET-28 a(+) expression vector, and the recombinant plasmid was transformed into *E. coli* strain BL21 (ΛDE3) gold, 1mM
IPTG induction was proceeded for protein purification

1024

1025

Fig S3

A.

L. donovani

SKI-5C

% promastigote survival vs $\log_{10}$ SKI-5C (nM)

IC_{50}- 100µM

B.

L. donovani

DMS

% promastigote survival vs $\log_{10}$ DMS (nM)

IC_{50}- 500µM

C.

ABC294640

L. donovani — ABC294640 — % promastigote survival — 60uM, 80uM, 100uM — 24H, 48H

THP-1 — % THP-1 survival vs $\log_{10}$ ABC (uM) — IC_{50}- 60µM (48 h)

Int. Amastigotes — % axenic amastigotes survival vs $\log_{10}$ ABC (uM) — IC_{50}- 69.7µM (48 h)

Axenic amastigotes — % axenic amastigotes survival vs $\log_{10}$ ABC (uM)

1026

1027

Fig S4

A.

CLUSTAL O(1.2.4) multiple sequence alignment

B.

- Pink = LD_SPHK1
- Green = Human_SPHK1
- Blue = Human DAGKc domain
- Yellow = LD_DAGKc domain

LD_DAGKc doain get structurally aligned with human DAGKc domain

LD_SPHK1_DAGKz domain (DAG-kinase catalytic (DAGKc) = 341-425
Human_SPHK1_DAGKc domain = 12-159

C.

Fig S5

A.
THP-1

IC$_{50}$- 19.53µM

B.
THP-1

IC$_{50}$- 116µM

1030

1031

Table S1.

Primer	Sequence	Annealing Temperature
IL-10	Forward 5' –TTAAGGGTTACCTGGGTTGC-3' Reverse 5'-TGAGGGTCTTCAGGTTCTCC-3'	55°C
IL-12	Forward 5'-ATGCCCCTGGAGAAATGGTG-3' Reverse 5'-GGCCAGCATCTCCAAA CTCT-3'	55°C
IL-32γ	Forward 5'-AGGCCCGAATGGTAATGCT-3' Reverse 5'-CCACAGTG TCCTCAGTGTCACA-3	60°C
TNF-α	Forward 5'-CCTCTCTCTAATCAGCCCTCTG-3' Reverse 5'-GAGGACCTGGGAGTAGATGAG-3'	60°C
RNU6AP	Forward 5'-GGCCCAGCA GTACCTGTTTA-3' Reverse 5'-AGATGGCGGAGGTGCAG-3'	60°C
JW	Forward 5' CCTATTTTACACCAACCCCCAGT-3' Reverse 5'-GGGTAGGGGCGTTCTGCGAAA-3'	60°C
mIL-10	Forward 5'-AAATAAGAGCAAGGCAGTGG-3' Reverse 5'-GTCCAGCAGACTCAATACACAC-3'	60°C
mTNF-α	Forward 5'-TACTGAACTTCGGGGTGATTGGTCC-3' Reverse 5'-CAGCCTTGTCCCTTGAAGAGAACC-3'	60°C

1032

1033

Figure legends :

Fig 1. (A) Estimation of *Ld*SphK-1-mediated generation of NBD-SIP levels against

SphK inhibitors in *L. donovani* promastigotes. *L. donovani* bob promastigotes were

cultured in 25 cm^2 flasks followed by treatment with SphK inhibitors; PF-543, DMS and

SKI-5C for 48h. Inhibition of SphK1 using these SphK inhibitors cause decreased S1P levels.

SphK inhibitor treated promastigotes were resuspended in buffer containing fatty acid-free

BSA (0.1% (w/v)) followed by resuspension in buffer containing BSA (1% (w/v)) and

incubated with NBD-sphingosine (10μM) for 45 min at 37 °C. Promastigotes incorporate

NBD-sphingosine (NBD-Sph), which are phosphorylated by Sphingosine kinase (SphK1) to

NBD-S1P. Bar graph depicts ELISA-based S1P quantification in promastigotes treated with

inhibitors. **(B) Purification and sub-cellular localization of Recombinant *rLd*SphK1 in *L.***

***donovani*.** Concentrated r*Ld*SphK1 protein. (i) Gel image showing a purified band of 102kDa

recombinant *Ld*SphK1 (ii) The subcellular localization of recombinant *Ld*SphK1 was

investigated in promastigotes. Panel A: DIC at 60X. Panel B: DAPI. Panel C: anti-*Ld*SphK1

antibody detected using FITC (green)-conjugated secondary antibody. Panel D: merged

micrographs. *Ld*SphK1 expression represented in the green channel. Co-localization shown

as the merged image. **(C) *In silico* docking of the *rLdSphK1* in presence of PF-543.** *In-*

silico ligand-substrate interaction was done using Autodock 1.5.7rc1 and Cygwin terminal.

The *in silico* docking generated the complex of *rLdSphK1*-PF-543 and represented as surface

model highlighted with residues involved in the interaction using Chimera, Ligplus,

Discovery Studio v19.1.0, and PyMOL v2.3.2 software. **(D) Biophysical interaction and**

functional characterization of the *rLdSphK1* (20μM) in presence of PF-543 (500nM,

1μM, 2.5μM, 5μM, 10μM, 20μM). MST analysis confirms the interaction between

rLdSphK1 and PF-543. Dose-response curves of *rLdSphK1*–PF-543 showed K_D of 29.3μM.

(E) Analysis of functional characterization and catalytic activity of the *Ld*SphK1

recombinant protein. (i) Time kinetics of activity of purified *rLd*SphK1 and NBD-S1P-

based fluorometric assay showing increase in NBD-SIP levels in presence of different

concentrations of *rLd*SphK1 protein. (ii) The inset shows inhibition of NBD-SIP levels in the

presence of different concentrations of PF-543 using 200ng of recombinant *Ld*SphK1. (iii)

Varying different concentration of NBD Sphingosine (2-10µM) and keeping *rLd*SphK1

constant (200ng), NBD-SIP levels were measured to calculate K_m and V_{max} of *rLd*SphK1.

The plate was immediately placed on the varioskan LUX Multimode Microplate Reader

(Thermo fisher, Massachusetts, USA) every 5 seconds for 20 min.

1067

Fig 2. (A) Purification of Recombinant *rhSphK1* from Supernatant. Gel image showing a

concentrated *rhSphK1* protein band of 50kDa. **(B) *In silico* docking of the *rHumanSphK1***

in presence of PF-543. *In silico* ligand-substrate interaction was done using Autodock

1.5.7rc1 and Cygwin terminal. The *in silico* bound complex of *rhSphK1*-PF-543 was

represented as surface model highlighted with residues involved in the interaction using

Chimera, Ligplus, Discovery Studio v19.1.0, and PyMOL v2.3.2 software. The list of

residues involved in hydrophobic interaction and hydrogen bond formation were highlighted

in the model. **(C) Functional characterization of the *rhSphK1* in presence of PF-543.**

MST analysis confirms the biophysical interaction between *rhSphK1* -PF-543 and *rhSphK1*-

ABC294640, which was used as a control. Dose-response curves of PF-543 (20µM) and

ABC294640 showed K_D of 34.3µM and 70.6µM respectively. **(D) Analysis of catalytic**

activity of the *rhSphK1* using NBD-S1P-based fluorometric assay. (i) Time kinetics of

activity of purified *rhSphK1* demonstrated increase in NBD-SIP levels in presence of 200ng

of *rhSphK1* protein. **(ii)** Varying different concentration of NBD Sphingosine (2-10µM) and

keeping *rhSphK1* constant (200ng), NBD-SIP levels were measured to calculate K_m and V_{max}

of *rhSphK1*. The readings were recorded every 5 seconds for 20 min using varioskan LUX

Multimode Microplate Reader (Thermo fisher, Massachusetts, USA).

1085

Fig 3. Effect of PF-543 on cytotoxicity and infectivity of *L. donovani* SphK1

overexpressor promastigote cell line (A) Detection of the presence of GFP encoded SphKa

plasmid in *Leishmania donovani* parasites. *L. donovani* bob promastigotes were cultured in

25 cm^2 flasks followed by transfection with GFP-tagged SphKa plasmid. **(B (i)) Estimation**

of *Ld*SphK-1-mediated generation of NBD-SIP levels against PF-543 in *L. donovani*

SphKa promastigotes. *L. donovani* bob promastigotes were cultured in 25 cm^2 flasks

followed by treatment with SphK1 inhibitor; PF-543 for 48h. Inhibition of SphK1 using PF-

543 had no significant effect on S1P levels. SphK inhibitor treated SphKa promastigotes were

resuspended in buffer containing fatty acid-free BSA (0.1% (w/v)) followed by resuspension

in buffer containing BSA (1% (w/v)) and incubated with NBD-sphingosine (10μM) for 45

min at 37 °C. Promastigotes incorporate NBD-sphingosine (NBD-Sph), which are

phosphorylated by Sphingosine kinase (SphK1) to NBD-S1P. Bar graph depicts ELISA-

based S1P quantification in SphKa promastigotes treated with PF-543. **(ii) Effect of SphK1**

inhibitor; PF-543 in *Leishmania* overexpressor *spp*. To evaluate the IC$_{50}$ for PF543,

approximately 5×10^4 *Ld SphKa* Bob cells were seeded in each well of 96-well flat bottom

plates and supplemented with M199 media containing PF-543 (200μl/well). The IC$_{50}$ was

found to be 42.71μM for *Leishmania donovani* SphKa promastigotes. Each experiment was

done in triplicates and repeated thrice. **(C (i)) Effect of infectivity of *L. donovani* SphK1**

overexpressor on the clearance of intracellular parasite load. THP-1 macrophages were

cultured in six-well plates in the presence or absence of *L. donovani* infection (MOI , 20:1)

for 6h. Further, THP-1 cells were infected with *L. donovani* SphK1 overexpressor

promastigotes at various MOIs (10:1, 20:1, 30:1, 50:1). Infected THP-1 were washed to

remove non-internalized parasites. Total RNA was enriched using TRIzol and the resulting

cDNA was subjected to real-time PCR analysis using primers specific for infectivity (JW)

genes in infected macrophages. RNU6A was used as a housekeeping gene. The results are

expressed as fold-change of uninfected THP-1 cells. Statistical significance was quantified

using the unpaired t-test. The data is a representation of mean ± SD from three independent

experiments *, p< 0.05; ***, p < 0.001. **(C (ii)) Analysis of GFP expression in *L. donovani***

SphK1 overexpressor and wild type promastigote cell lines. Detection of the presence of

GFP encoded SphKa and *Bob* plasmids in *Leishmania donovani* and *Leishmania*

overexpressor parasites. *L. donovani bob* and *L. donovani* overexpressor promastigotes were

cultured in 25 cm^2 flasks followed by confocal microscopy.

1118

Fig 4. (A) Effect of SphK1 inhibitors; PF-543 and ABC294640 in *Leishmania spp* and

host macrophages. To evaluate the IC$_{50}$ for PF543, approximately 6×10^3 THP-1 and 5×10^4

Ld Bob cells were seeded in each well of 96-well flat bottom plates and supplemented with

RPMI and M199 media containing PF-543 (200μl/well) respectively. For intracellular

amastigotes, 1×10^6 THP-1 cells, treated with 50 ng/ml of phorbol 12-myristate 13-acetate

(PMA) were seeded on glass coverslip in a 6-well plate for 48h. They were infected with late

log-phase *L. donovani* promastigotes and simultaneously treated with PF-543. The cells were

further incubated for 2 days at 37°C and 5% CO$_2$. To determine, the intracellular parasite

burdens (mean number of amastigotes per macrophage) were microscopically assessed using

Giemsa staining. For axenic amastigotes, the axenically cultured forms grew optimally at a

temperature of 32-33°C in a growth media with pH of 5.4. The IC$_{50}$ were found to be 500nM,

90μM, 17.54μM and 45μM for *Leishmania donovani* promastigotes, THP-1 macrophages,

intracellular amastigotes and axenic amastigotes respectively. Each experiment was done in

triplicates and repeated thrice. **(B) Effect of Sphk1 inhibition on parasite infectivity in**

THP-1 macrophages. THP-1 cells grown in RPMI medium were treated with SphK1

inhibitor; PF-543 for 24h followed by infection with *L. donovani* promastigotes at an MOI of

20:1 for 48h. (i) After 48h of infection, Giemsa staining was performed to assess the parasite

burden. THP-1 cells were fixed, Giemsa stained and amastigotes were counted visually.

Virulence capacity was determined by calculating infectivity (left panel) and parasitemia

(right panel). Untreated infected THP-1 cells were used as control. (ii) Total RNA was

enriched using TRIzol. Infectivity was validated using qRT-PCR with primer specific marker

JW as a molecular indicator. Expression of JW mRNA in infected cells inferred parasite load

and represented as percentage infectivity. Data analysis was performed using the $2^{-\Delta\Delta CT}$

method. The results are representative of three independent experiments. Statistical

significance was quantified using the unpaired t-test. The results signify mean ± S.D with n =

3, *P < 0.05.

1145

Fig 5. (A, B) Estimation of glutathione and Caspase 9 levels upon PF-543 treatment in

***Leishmania* infected macrophages.** THP-1 macrophages were cultured in six-well plates in

the presence of *L. donovani* infection (MOI□, 20:1) for 6h. Infected THP-1 were washed to

remove non-internalized parasites and treated with PF-543 (20µM) or DL-threo-PPMP;

glucosylceramide synthase inhibitor (20µM) or SIP (10µM) for next 48h. GSH/GSSG and

Caspase-9 levels were analysed using cell lysates quantified for GSG/GSSH assay and

Western blot. Band intensities were quantified by ImageJ software and were plotted in

GraphPad Prism. Data from one of three experiments is shown.

1154

Fig 6 (A(i), B) Effect of PF-543 on the expression of inflammatory molecules and

clearance of intracellular parasite load. THP-1 macrophages were cultured in six-well

plates in the presence or absence of *L. donovani* infection (MOI□, 20:1) for 6h. Infected

THP-1 were washed to remove non-internalized parasites and treated with PF-543 for next

48h. Total RNA was enriched using TRIzol and the resulting cDNA was subjected to real-

time PCR analysis using primers specific for inflammatory (IL32γ, IL-12 , IL-10 and TNF-α)

and infectivity (JW) genes in PF-543 treated and untreated infected macrophages. RNU6A

was used as a housekeeping gene. The results are expressed as fold-change of uninfected

THP-1 cells. Statistical significance was quantified using the unpaired t-test. The data is a

representation of mean ± SD from three independent experiments *, p< 0.05; ***, p < 0.001.

(A (ii)) Graph represents cytokine levels measured by sandwich ELISA. Statistical

significance was quantified using the unpaired t-test with Welch's correction, in LPS-

stimulated, *Ld*-infected and PF-543 treated macrophages.

1168

Fig 7. (A-B). Evaluation of PF-543 and Amphotericin B formulation on cytotoxicity or

metabolic viability of *Leishmania* promastigotes and intracellular amastigotes. (A) To

determine the combinatorial impact of PF-543 and Amphotericin B formulation,

promastigotes of *L. donovani* were seeded in 96-well plates (3×10^6 parasite/mL) in

exponential growth phase at increasing concentrations (12.5nM–500nM) of Amphotericin B

and (150nM- 2000nM) of PF-543 compounds and maintained at 26°C. After 48 h of

incubation, the IC_{50} was determined by MTT assay. The absorbance was measured in a

MultiskanEX photometric plate reader for microplates at 540 nm. Data were obtained from

three independent experiments performed in triplicate. The fractional inhibitory concentration

index (FIC) was evaluated to determine the synergistic activity of lead compounds; PF-543

with Amph-B. Finally, FIC < 0.5 represented strong synergistic activity between two drugs.

While FIC > 4 is was considered as the antagonistic activity. FIC values were obtained from

three independent experiments. **(B)** For validating the antileishmanial activity by PF-543 and

Amphotericin B formulation on intracellular amastigotes, qPCR-based analysis was

performed for all the treatments. For the same, THP-1 cells were seeded at a concentration of

8×10^5 cells/mL in 96-well plates and incubated for 24h with PMA (10ng/mL) supplemented

with RPMI 1640. Following 24h priming with PMA, the culture medium was removed, and

the cells were infected with *L. donovani* followed by treatment with the formulations having

concentrations kept similar as used previously for promastigotes at 37°C and 5% CO_2. Total

RNA was extracted, and the resulting cDNA was subjected to real-time PCR analysis using

primers specific for kinetoplast minicircle DNA (kDNA)/JW of *L. donovani*. This primer

does not match to human genome, thus directly indicates the intracellular parasite load

against the treatments. Data analysis was performed using the $2^{-\Delta\Delta CT}$ method. Values are mean

$\pm$ S.D (n = 3). The results are representative of three independent experiments performed in

triplicates.

1194

Fig 8. Expression of JW and inflammatory markers upon PF-543 treatment in

***Leishmania-infected* Swiss mice.** Evaluation of the **(A)** immunomodulatory potential and **(B)**

infectivity of PF-543 and Amph-B in *Leishmania*-infected Swiss mice was performed.

10mg/Kg of PF-543 and 2mg/Kg of Amphotericin B alone were given for individual

treatment in *Leishmania* infected mice. For constituting the synergistic effect of two drug

partners; 2mg/Kg (1mM) of PF-543 and 0.4mg/Kg (0.12µM) of Amphotericin B were

prepared in 1XPBS and given together to a mouse weighing 30gm. To evaluate the

parasitemia and expression of inflammatory markers in infected mice upon drug

administration, spleen was harvested from each group of mice and qRT-PCR analysis was

performed using primers specific for inflammatory markers *IL-10 and TNFα* and parasite

specific kinetoplast minicircle gene, *JW* in PF-543 and Amph-B treated infected mice in

respect to the untreated Swiss mice. RNU6A (RNA, U6 small nuclear 1; THP-1 cells) was

used as a housekeeping gene. The results were expressed as normalized fold-change of

respective control. Statistical significance was quantified using the unpaired t-test. The data is

a representation of mean ± SD from three independent experiments *, p< 0.05; ***, p <

0.001.

1211

Fig S1. (A) Nucleotide sequence encoding full length sequence of *L. donovani* Sphk1 (1-

2808 bp). This sequence was selected for recombinant protein generation as a fusion protein

with an N-terminal 6x- Histidine-tag using the vector pET-28 a(+) vector. **(B) Cloning of *L.***

***donovani* SphK1 in pET28a(+) vector.** PCR amplification of DNA fragment from genomic

DNA of *L. donovani* using *Ld*Sphk1_BamHI and *Ld*Sphk1_XhoI with Phusion™ High-

Fidelity DNA Polymerase. Purification of the amplified DNA fragment was done using

QIAquick Gel Extraction Kit followed by digestion of purified *Ld*Sphk1 insert and the

expression vector pET-28 a(+) with BamHI/XhoI restriction enzymes. The positive clones

were screened by colony PCR followed by confirmation of the cloned plasmid with

BamHI/XhoI restriction digestion. **(C) *rLd*Sphk1 recombinant protein purification**.

Transformation of positive clone in Rossetta *E. coli* expression strain for *Ld*Sphk1

recombinant protein expression. For recombinant protein induction, the cells were induced

with 1mM IPTG for 4 hours at 37°C in Terrific broth. Purification of recombinant protein

was achieved by affinity chromatography. The *rLd*SphK1 protein was eluted after addition of

150mM imidazole solution. The eluted fractions were analysed by running on 12% SDS

PAGE.

1228

Fig S2. (A) The Human SphK1 amino acid sequence for the protein. (B) Purification

of the recombinant *Human*SphK1 (*rHuman*SphK1) protein. Overexpression of 6×His-

SphK-1 (*rhSphK-1*) was induced with 1mM IPTG at an optical density (OD600) of 0.6, for 4

h at 37°C. The protein was purified using Ni-NTA agarose resin. The *rhSphK1* protein was

eluted with a continuous imidazole gradient of 50mM, 100mM, 250mM, and 500mM. The

protein purification was validated by 12% SDS-PAGE, followed by immunoblotting with

anti-His tag antibody. A single band corresponding to ~50 kDa was observed on SDS PAGE

corresponding to Human specific SphK1.

1237

Fig S3. (A,B) Effect of SphK1 inhibitors; SKI-5C and DMS in *Leishmania*

***promastigotes*.** To evaluate the IC_{50} for SKI-5C and DMS, approximately 5×10^4 *Ld* Bob cells

were seeded in each well of 96-well flat bottom plates and supplemented with M199 media

containing SKI-5C and DMS (200µl/well) respectively. The cells were further incubated for

2 days at 37°C and 5% CO_2. The IC_{50} were found to be 100µM for SKI-5C and 500µM for

DMS respectively in *Leishmania donovani* promastigotes. Each experiment was done in

triplicates and repeated thrice. **(C) Effect of SphK1 inhibitor; ABC294640 in *Leishmania***

***spp* and host macrophages.** To evaluate the IC_{50} for ABC294640, approximately 6×10^3

THP-1 and 5×10^4 *Ld* Bob cells were seeded in each well of 96-well flat bottom plates and

supplemented with RPMI and M199 media containing ABC294640 (200µl/well)

respectively. For intracellular amastigotes, 1×10^6 THP-1 cells, treated with 50 ng/ml of

phorbol 12-myristate 13-acetate (PMA) were seeded on glass coverslip in a 6-well plate for

48h. They were infected with late log-phase *L. donovani* promastigotes and simultaneously

treated with ABC294640. The cells were further incubated for 2 days at 37°C and 5% CO_2.

To determine, the intracellular parasite burdens (mean number of amastigotes per

macrophage) were microscopically assessed using Giemsa staining. For axenic amastigotes,

the axenically cultured forms grew optimally at a temperature of 32-33°C in a growth media

with pH of 5.4. The IC_{50} were found to be 60µM for THP-1 macrophages and 69.7µM for

intracellular amastigotes respectively. Each experiment was done in triplicates and repeated

thrice.

1258

Fig S4. (A) Amino acid sequence alignment of LD_SphK1 and Human_SphK1 utilizing

MEGAX program (version 10.2.2). The results demonstrated considerable alignment with

similar and a limited number of identical amino acids. **(B) 3D structural alignment of**

LD_SphK1 and Human_SphK1 protein structures with Pymol software. Substantial

alignment between the two proteins, with an RMSD of 0.585 was observed. Furthermore, the

alpha helix and beta sheets of the Human DAGKc domain (Blue) accurately aligned with the

LD_DAGKc domain (Yellow). **(C) Construction of phylogenetic tree utilizing SphK1**

gene sequences from various species of _Leishmania_, humans, rats, and mice. The

phylogenetic tree was constructed using neighbor-joining technique with MEGAX software.

Bootstrap analyses with 1000 replicates were conducted to evaluate the robustness of the

constructed phylogenetic tree.

Fig S5. (A) Effect of Ceramide analogs; DL-threo-PPMP and DL-erythro-PPMP in

THP-1 macrophages. To evaluate the IC_{50} for DL-threo-PPMP and DL-erythro-PPMP,

approximately 6×10^3 THP-1 cells were seeded in each well of 96-well flat bottom plates and

supplemented with RPMI media containing DL-threo-PPMP and DL-erythro-PPMP

(200µl/well) respectively. The cells were further incubated for 2 days at 37°C and 5% CO_2.

The IC_{50} were found to be 19.53µM and 116µM for DL-threo-PPMP and DL-erythro-PPMP

respectively in THP-1 macrophages. Each experiment was done in triplicates and repeated

thrice.

Table S1. Primer sequences of target genes mentioned along with their annealing

temperatures