AWmeta empowers adaptively-weighted transcriptomic meta-analysis

Yanshi Hu [1,*], **Zixuan Wang**[1], **Yueming Hu** [1], **Cong Feng**[1,2], **Qiuyu Fang**[3], **and Ming Chen** [1,2,*]

[1]Department of Bioinformatics, College of Life Sciences, Zhejiang University, Hangzhou 310058, China
[2]Biomedical Big Data Center, The First Affiliated Hospital, Zhejiang University School of Medicine; Institute of Hematology, Zhejiang University, Hangzhou 310058, China
[3]Department of Thoracic Surgery, The Second Affiliated Hospital Zhejiang University School of Medicine, Zhejiang University, Hangzhou 310058, China
[*]Corresponding author emails: yanshihu@zju.edu.cn; mchen@zju.edu.cn

Abstract

Transcriptomic meta-analysis enhances biological veracity and reproducibility by integrating diverse studies, yet prevailing *P*-value or effect-size integration approaches exhibit limited power to resolve subtle signatures. We present AWmeta, an adaptively-weighted framework that unifies both paradigms. Benchmarking across 35 Parkinson's and Crohn's disease datasets spanning diverse tissues and adaptively down-weighting underpowered studies, AWmeta yields higher-fidelity differentially expressed genes (DEGs) with markedly reduced false positives and establishes superior gene differential quantification convergence at both gene and study levels over state-of-the-art random-effects model (REM) and original studies. AWmeta requires fewer samples and DEGs from original studies to achieve substantial gene differential estimates, lowering experimental costs. We demonstrate AWmeta's remarkable stability and robustness against external and internal perturbations. Crucially, AWmeta prioritizes disease tissue-specific mechanisms with higher functional coherence than those from REM and original studies. By bridging statistical rigor with mechanistic interpretability, AWmeta harmonizes heterogeneous transcriptomic data into actionable insights, serving as a transformative tool for precision transcriptomic integration.

Main

The exponential expansion of publicly available transcriptomic data, propelled by high-throughput sequencing advancements[1], presents unprecedented opportunities and concomitant challenges for uncovering robust biological insights through meta-analysis. By integrating findings across independent studies, this powerful approach transcends the limitations of individual datasets, mitigating issues of statistical power, experimental variability, tissue heterogeneity, and platform-specific biases that often obscure subtle yet pathologically relevant expression signatures[2,3]. As complex diseases increasingly defy dissection by single-study designs, meta-analysis has become indispensable for identifying reproducible biomarkers and elucidating disease pathways with enhanced confidence and precision[4,5].

Contemporary meta-analysis methodologies predominantly fall into three primary categories[6]: *P*-value combination, effect-size integration, and rank aggregation. *P*-value-based methods, such as Fisher's[7], Stouffer's Z-score[8], and the adaptively weighted Fisher's (AW-Fisher) technique[9], efficiently aggregate statistical significance but typically disregard effect magnitude and directionality. Conversely, effect-size approaches quantify expression differences, with the random-effects model (REM)[10] widely adopted due to its capacity to accommodate inter-study heterogeneity and its perceived biological interpretability[11]. REM has underpinned discoveries in diverse areas, including characterizing gut microbiome dysbiosis in Parkinson's disease[12], identifying predictive biomarkers for cancer immunotherapy[13], and assessing pharmacogene expression in nonalcoholic fatty liver disease[14]. Nevertheless, REM and related models can exhibit sensitivity to outlier studies and rely heavily on assumptions about the underlying heterogeneity structure[11,15]. Rank-based techniques (e.g., RankProd and RankSum[16,17]) offer robustness against outliers but often at the cost of statistical resolution and power. Critically, these distinct methodological frameworks typically operate in isolation, failing to synergistically leverage their complementary strengths.

This methodological schism represents a fundamental constraint in transcriptomic meta-analysis, particularly impeding progress in complex disease research where robust biological inference demands both high statistical confidence in identifying dysregulated genes and accurate quantitative estimates of their expression changes[18,19]. Current *P*-value integration schemes, while adept at pinpointing consistently altered genes, offer minimal information on the magnitude or biological relevance of these alterations. Conversely, effect-size methods, though designed to quantify these changes, struggle with the pervasive heterogeneity inherent in pooling diverse experimental designs, tissue sources, or patient cohorts, potentially diminishing the

estimate reliability[20]. Furthermore, existing approaches lack sophisticated mechanisms to dynamically weight studies based on
intrinsic data quality or specific biological context, thereby limiting overall sensitivity and the depth of achievable insights.
To address these critical gaps, we introduce AWmeta, a novel meta-analytic framework that unifies the statistical rigor
of P-value–based method with the quantitative power of effect-size paradigm. The core innovation of AWmeta lies in an
adaptive weighting scheme that (1) identifies and up-weights the most informative studies from a heterogeneous pool, and
(2) explicitly models between-dataset variability to mitigate bias arising from sample size imbalances, data sparsity, and
technical heterogeneity. Validated on 35 diverse transcriptomic datasets spanning Parkinson's and Crohn's disease across
multiple tissues, AWmeta consistently outperforms the state-of-the-art REM. It secures higher-fidelity differentially expressed
gene (DEG) identification with markedly reduced false positives, exhibits greater resilience to limited sample size and DEG
sparsity, and maintains remarkable stability against various perturbations. More importantly, AWmeta delivers gene differential
estimates with enhanced reproducibility and biological mechanism interpretation. By reducing sample-size requirements and
demonstrating robustness to noise, AWmeta empowers researchers to mine the ever-expanding corpus of public transcriptomic
data more effectively, accelerating the discovery of actionable molecular insights across biomedical domains.

Results

Overview of AWmeta
AWmeta is a transcriptomic meta-analytical framework, uniquely synergizing statistical strengths from P-value and effect size
integration methods (Fig. 1a and "Overview of the AWmeta framework" section in Methods). Following preprocessing of
multiple transcriptomic datasets from original studies of the same disease tissue ("Transcriptomic data preprocessing" section
in Methods), AWmeta implements two complementary gene-wise modules: AW-Fisher and AW-REM. The AW-Fisher module
calculates meta P-values by optimizing study-specific weights to minimize the combined probability, effectively filtering out
less informative studies while preserving statistical power. Subsequently, in AW-REM module these optimized weights are
embedded into REM architecture to derive weighted fold change estimates.
In the following sections, we detail a multifaceted rigorous comparison of AWmeta against the state-of-the-art REM, a
representative effect-size integration method[21,22]. Benchmarking on 35 transcriptomic datasets from Parkinson's and Crohn's
disease tissues (Extended Data Fig. 1 and "Transcriptomic datasets" section in Methods), this evaluation assessed key metrics
including DEG detection capability and discrimination, gene- and study-wise gene differential quantification convergence,
stability and robustness, and biological relevance ("Transcriptomic meta-analysis evaluation metrics" section in Methods).
AWmeta consistently demonstrated superior performance, establishing its utility as a powerful tool for precision transcriptomic
integration and providing a solid foundation for downstream biological investigation.

AWmeta secures robust higher-fidelity DEG identification across transcriptomic contexts
A primary goal of transcriptomic meta-analysis is to enhance statistical power for identifying DEGs reliably, i.e., to detect
more subtle yet vital DEGs, typically defined by statistical significance and fold-change thresholds (Fig. 1b and "DEG
detection capability evaluation" section in Methods). Systematically benchmarking across five distinct disease tissue contexts
using nine combinations of statistical significance (0.01, 0.05, and 0.10) and fold-change thresholds ($\log_2 1.2$, $\log_2 1.5$, and
$\log_2 2.0$), AWmeta consistently identified significantly more DEGs than REM ($P < 10^{-4}$, one-tailed Welch's t-test over 100
bootstrap iterations; Fig. 1c and Extended Data Fig. 2). For instance, under a specific threshold combination ($P < 0.01$ and
$|\log_2 \text{FC}| > \log_2 1.2$), AWmeta yielded 69–475% increases in detected DEGs versus REM across all tissues (Fig. 1c), which
demonstrates AWmeta's superior statistical sensitivity in DEG detection.
A key challenge in meta-analysis is to increase statistical power while rigorously controlling false positives. To formally
evaluate this trade-off, we designed a semi-synthetic simulation framework to assess DEG discrimination ("DEG discrimination
evaluation using semi-synthetic simulation strategy" section in Methods). Inspired by Li and colleagues[23], this framework
first creates benchmark datasets with a known ground truth of positives (DEGs) and negatives (non-DEGs) (Fig. 1d). We then
systematically challenged AWmeta's performance by degrading the biological signal in a controlled manner. This was achieved
by permuting sample labels in a progressively increasing number of studies within each tissue context (Fig. 1e). Specifically, we
simulated three distinct noise scenarios by permuting the minimum, median, and maximum allowable number of studies, where
DEG discrimination was quantified by the area under the receiver operating characteristic curve (AUROC) and the area under
the precision-recall curve (AUPRC) over the above defined benchmark genes (Fig. 1f). This perturbation design enabled a
rigorous assessment of AWmeta's resilience across diverse data quality landscapes, a critical feature for real-world applications.
Across all simulated noise levels, AWmeta consistently outperformed or was comparable to REM in DEG discrimination.
Under minimum-permuted low-noise condition, AWmeta demonstrated clear and significant advantage across nearly all tissue
contexts and DEG thresholds ($P < 10^{-4}, 10^{-3}, 10^{-2}$ or 0.05, one-tailed Mann-Whitney test; Fig. 1g and Extended Data Fig. 3).
As expected, performance decayed for both methods with increasing noise from median and maximum study permutations.
However, AWmeta's superiority over REM was not only maintained but often became more pronounced under these more

challenging conditions ($P < 10^{-4}$, one-tailed Mann-Whitney test; Fig. 1h,i and Extended Data Fig. 4 and 5). Notably, AWmeta's
performance remained remarkably robust even under high-noise scenarios, with median AUROC and AUPRC exceeding 0.85
in most cases (Extended Data Fig. 5), highlighting its ability to effectively discount noise from potentially confounding studies.
Taken together, these results demonstrate that AWmeta achieves a superior balance between heightened detection sensitivity
and robust discrimination for higher-fidelity DEG identification from heterogeneous transcriptomic datasets.

AWmeta establishes superior gene- and study-wise convergence in gene differential quantification

To rigorously assess AWmeta's ability to synthesize a consensus biological signal from heterogeneous transcriptomic datasets,
we evaluated its convergence at both gene and study levels. We first quantified gene-wise convergence—the proximity of a
gene's meta effect-size estimates to ones from original studies—using the mean absolute deviation (MAD) between meta and
original fold changes (Fig. 2a and "Gene-wise convergence assessment for gene differential quantification" section in Methods),
where lower MADs signify more accurate biological representations.

AWmeta consistently yielded significantly lower gene-wise convergence scores in all five disease tissues, compared to both
REM and baseline ones (Fig. 2b–f; $P < 10^{-4}$ or 0.05, one-tailed Mann-Whitney test against AWmeta). Notably, while some
original studies occasionally outperformed REM in specific contexts, AWmeta (merely 57–74% of REM) consistently achieved
lower scores than any original study across all tissues, which suggests its capacity to robustly identify and integrate reliable
signals while effectively down-weighting divergent studies, thereby providing a superior consensus representation of the gene
expression landscape. Since disease processes are primarily driven by DEGs, we further confirmed this superior performance
was still evident for these specific genes by the same assessment paradigm using nine distinct thresholds, combining three
significance levels (0.01, 0.05, and 0.10) and three fold-change cutoffs ($\log_2 1.2$, $\log_2 1.5$, and $\log_2 2.0$). Across all threshold and
disease tissue scenarios, AWmeta maintained evidently lower convergence scores, accounting for 56–80% of REM (Extended
Data Fig. 6a–e; $P < 10^{-4}, 10^{-3}, 10^{-2}$ or 0.05, one-tailed Mann-Whitney test over AWmeta), which underscores AWmeta's
effectiveness to derive robust fold-change estimates for disease-relevant genes, independent of specific statistical criteria.

Next, we evaluated study-wise convergence to determine how well the meta-analytic results reflect the collective evidence
across all contributing studies. We employed three complementary approaches: an adjusted rank-sensitive similarity metric
emphasizing top-ranked genes (denoted "adjusted DE list similarity" thereafter), the arithmetic mean of Jaccard and overlap
coefficients (JC/OC) for DEG concordance, and the phi coefficient (PC)[24] to assess classification agreement beyond chance
(Fig. 2g–j, Extended Data Fig. 7 and "Study-wise convergence assessment for gene differential quantification" section in
Methods). Higher scores indicate better study-wise convergence for all metrics.

Across the five disease tissues, AWmeta consistently achieved significantly higher study-wise convergence scores than
baselines representing original inter-study agreement, with dramatic 30–1,166% improvements (Fig. 2k–m; $P < 0.05$ and
$|\log_2 FC| > \log_2 1.2$ where applicable). While overall convergence scores tended to be lower in Parkinson's over Crohn's
disease tissues, potentially reflecting higher inherent variability within these specific disease contexts, AWmeta significantly
outperformed REM in the majority (10 out of 15) of comparisons across different metrics and tissues, performing comparably
otherwise, particularly pronounced in tissues like Parkinson's and Crohn's peripheral blood, where AWmeta's convergence
scores improved by 35–156% compared to REM (Fig. 2k–m). To further validate these findings for the JC/OC and PC metrics,
we confirmed AWmeta's superior performance across nine different DEG cutoffs (Extended Data Fig. 8). These results indicate
that the gene differential quantification results processed by AWmeta are more representative of the faithful consensus signal
across studies than those derived from REM or original studies.

AWmeta attains accelerated meta-analysis convergence with reduced samples and DEGs

While large sample sizes are known to enhance transcriptomic differential quantification efficacy, the feasibility of achieving
reliable meta-analytic effect sizes from studies with limited samples has remained an open question. To fill this gap, we
systematically correlated study-wise convergence with sample size and employed Spearman correlation with a two-tailed
significance test to capture potentially non-linear dependencies. To ensure robustness and minimize sensitivity to arbitrary
cutoffs, this association analysis leveraged results derived from nine distinct DEG thresholds (as previously described) across
both JC/OC and PC metrics.

We observed pronounced positive correlations between sample size and study-wise convergence in Parkinson's substantia
nigra and peripheral blood, as well as in Crohn's ileal mucosa, for both AWmeta and REM (Fig. 3a–c). This reinforces the
principle that larger cohorts generally yield results that more closely approximate the consensus biological signal. Crucially,
at equivalent sample sizes, AWmeta consistently outperformed REM in convergence across all five disease tissues, where
performance gaps widened progressively with increasing sample size. To illustrate, using the adjusted DE list similarity metric
in Crohn's ileal mucosa, while AWmeta and REM exhibited comparable convergence scores (~ 0.38) at the same sample size
of 62, AWmeta's score surged to 0.77 at a sample size of 200—a 54% increase over REM's 0.50 (Fig. 3a), which demonstrates
AWmeta can reach much better convergence level under the same sample-size context. Notably, AWmeta attains comparable
convergence with considerably fewer samples relative to REM. To reach a convergence score of 0.2 in Parkinson's substantia

nigra, for instance, AWmeta necessitated 44% or 43% fewer samples than REM with JC/OC (15 versus 27; Fig. 3b) or PC metric (20 versus 35; Fig. 3c). This marked improvement in sample efficiency translates directly into substantial reductions in experimental cost and resource allocation.

We further hypothesized that DEG abundance would positively contribute to accurate meta-analytic estimates for gene differential quantification and substantiated this premise through the observation of robust positive correlations between DEG number and study-wise convergence across all five disease tissues (Fig. 3d–f), following the analogous procedure to the above sample-size correlation analysis. For studies with similar DEG counts, AWmeta constantly yielded higher convergence scores than REM, with performance disparity more pronounced as DEG abundance ascended, suggesting AWmeta possesses a heightened sensitivity to the underlying biological signal embedded within scattered DEGs. For example, to arrive at a PC metric-based convergence score of 0.4 in Crohn's peripheral blood, AWmeta required only 2,000 DEGs, whereas REM exacted approximately 4,600—a 130% escalation—for equivalent resolution (Fig. 3f). This highlights AWmeta's capacity to achieve robust convergence even from datasets with sparser DEG profiles, a practical advantage for experimental designs where DEG discovery may be limited.

Collectively, these findings imply AWmeta can deliver reliable meta-analytic outcomes in less stringent experimental scenarios, such as studies involving milder treatments or designs with fewer replicates, which substantially mitigates experimental complexity and cost, especially for ambitious large-scale research programs.

AWmeta delivers remarkable stability and robustness in transcriptomic integration

We sought to determine whether AWmeta's adaptively-weighted strategy confers superior stability and robustness to gene differential meta-estimates against REM, and designed quantitative metrics to evaluate consistency over random splits and resilience to systematic perturbations across the five disease tissues.

First, we assessed stability by quantifying the concordance of ranked gene differential lists derived from randomly halved sample sets within each study, a process replicated across 100 iterations (Fig. 4a and "Stability and robustness assessment of transcriptomic integration" in Methods). Across all five disease tissues, AWmeta exhibited markedly higher stability scores relative to REM (Fig. 4b; $P < 10^{-4}$, one-tailed Welch's t-test), underscoring its enhanced consistency under data rationing. The observation that median stability scores for both methods were below 0.7, is likely attributable to the inherently reduced statistical power and study-wise convergence that accompanies halving the sample size (Fig. 3a).

We then challenged the robustness of each method against two distinct forms of perturbation: external interference, simulated by the inclusion of a thematically unrelated study (Fig. 4c and "Stability and robustness assessment of transcriptomic integration" in Methods), and internal fragility, evaluated through a systematic leave-one-study-out procedure (Fig. 4e and "Stability and robustness assessment of transcriptomic integration" in Methods). Against external interference, AWmeta displayed remarkable resilience with median robustness scores above 0.8 and established a significant performance margin over REM across all tissues (Fig. 4d; $P < 10^{-4}, 10^{-2}$ or 0.05, one-tailed Mann-Whitney test). This capacity to resist discordant data is a direct consequence of AWmeta's adaptive weighting scheme, which effectively minimizes the influence of outlier studies. In internal robustness assessment, AWmeta again achieved significantly higher scores than REM in four of the five tissues (Fig. 4f; $P < 0.05$, one-tailed Mann-Whitney test). The sole exception was Crohn's peripheral blood, where the small cohort of only three studies constrained the median robustness scores to below 0.6 for both methods. Notably, in tissues comprising six or more studies, AWmeta achieved exceptional median internal robustness scores around 0.9, demonstrating highly consistent results even upon the exclusion of individual constituent studies.

These rigorous stress tests validate that AWmeta's adaptive weighting architecture endows the meta-analytic process with significantly strengthened stability and robustness. This reinforcement ensures the derivation of more dependable biological insights when integrating diverse and inherently heterogeneous transcriptomic datasets.

AWmeta facilitates discovery of disease-relevant genes

A pivotal determinant of a meta-analysis method's utility is its capacity to prioritize genes of genuine pathological importance. To rigorously assess this, we quantified the biological relevance of gene rankings from AWmeta, REM, and the original studies (as baselines) against authoritative Parkinson's and Crohn's disease-gene benchmarks—compiled from DisGeNET[25], MalaCards[26], and an in-house curated genetic variation corpus[27]—using a custom metric that integrates both statistical significance and effect size magnitude, which provides an objective and threshold-agnostic evaluation of gene prioritization performance (Fig. 5a and "Biological relevance assessment of gene differential quantification" section in Methods).

Prior to assessing performance, we first validated the coherence of our benchmark gene sets, with overlap magnitude quantified using odds ratio (OR) and statistical significance determined by Fisher's exact test. Pairwise comparisons revealed substantial overlaps among the three independent sources for both Parkinson's disease (e.g., DisGeNET versus MalaCards, OR = 138.8, $P = 5.1 \times 10^{-115}$) and Crohn's disease (e.g., DisGeNET versus MalaCards, OR = 242.9, $P = 1.3 \times 10^{-27}$) (Fig. 5b). This strong reciprocal consistency affirmed their utility for a reliable evaluation of biological relevance.

Our primary analysis revealed that AWmeta consistently generates more biologically meaningful gene rankings than REM
and the baseline studies ($P < 10^{-4}, 10^{-3}, 10^{-2}$ or 0.05, Nemenyi post-hoc test; Fig. 5c–e). Specifically, when benchmarked
against our genetic variant corpus, AWmeta achieved significantly higher relevance scores across all interrogated tissues
(Fig. 5c). This superior performance extended to the DisGeNET benchmark in critical disease tissues, including Parkinson's
substantia nigra and Crohn's peripheral blood and ileal mucosa (Fig. 5d). A similar advantage was observed using the MalaCards
benchmark for Parkinson's substantia nigra and Crohn's ileal and colonic mucosa (Fig. 5e). Notably, AWmeta's superiority was
particularly pronounced in the primary disease-affected tissues—Parkinson's substantia nigra and Crohn's ileal mucosa—where
it surpassed baselines across all three independent benchmarks. Cumulatively, in 11 instances of the 15 tissue-benchmark
comparisons (5 tissues $\times$ 3 benchmarks), AWmeta's scores were significantly higher than those of both the baselines and REM.
In stark contrast, REM failed to offer a significant improvement over baseline scores in 11 of 15 comparisons, underscoring its
limited ability to distill new biological insights from existing data.

These results establish that AWmeta's gene prioritization is not merely a statistical refinement but a substantive improvement
in biological fidelity. By more effectively elevating established disease-associated genes to the top of significance rankings,
AWmeta provides a clearer and more accurate representation of the underlying pathology. This enhanced resolution positions
AWmeta as a powerful discovery engine, capable of transforming heterogeneous transcriptomic datasets into a focused and
mechanistically coherent view of disease processes.

AWmeta enables disease tissue-specific mechanism interpretation

225 We further implemented Gene Ontology (GO) enrichment to explore disease mechanism interpretation based on meta-analysis
226 prioritized genes. To avoid arbitrariness, three thresholds (100, 300, and 500) were used to select the number of top integrated
227 rank genes ("Biological relevance assessment of gene differential quantification" section in Methods). The enrichment ratio
228 quantifies the degree to which GO terms are significantly enriched in relevant disease tissues:

$$\text{Enrichment ratio} = \frac{\text{Gene ratio}}{\text{Background ratio}} \tag{1}$$

where gene ratio is the proportion of genes annotated to a specific GO term within the top integrated rank genes, and background
ratio represents the analogous fraction across a reference gene set. GO terms with higher enrichment ratios are more likely to
be involved in a given disease tissue. For comparison, GO enrichments derived from original studies served as baselines.

Compared with REM and baselines, representative GO terms enriched in AWmeta-derived top integrated rank genes
consistently exhibited the highest enrichment ratios across nearly all five disease tissues (Fig. 5f), demonstrating AWmeta's
enhanced capacity for disease-relevant gene prioritization in tissue-specific contexts. In contrast, REM underperformed relative
to some baselines (Fig. 5f), reflecting diminished biological relevance within its gene sets. For instance, biological processes
related to synaptic organization and transmission ("synaptic transmission, dopaminergic", "regulation of synapse organization",
and "distal axon") were significantly enriched in Parkinson's substantia nigra, consistent with their known involvement in
Parkinson's pathogenesis[28, 29]. Likewise, "metal ion transmembrane transporter activity" and "regulation of membrane potential"
were significantly enriched, highlighting their pivotal roles in Parkinson's substantia nigra-involved mechanisms[30–32]. AWmeta
achieved the highest enrichment for these GO terms, a trend robust across all thresholds. The enrichment ratios for AWmeta
and some baselines monotonically decreased with more integrated rank genes included, indicating these term-related genes are
concentrated at the very top of the ranked lists. Strikingly, REM-prioritized genes exhibited minimal or absent enrichment
across all five representative GO terms, further underscoring its impaired capacity to capture contextual biological functions.

Given the well-established inflammatory pathogenesis of Parkinson's and Crohn's disease in non-hematopoietic tissues[33, 34],
we hypothesized that blood-derived gene signatures would reflect systemic immune dysregulation and vascular barrier impair-
ment at disease-relevant interfaces: Parkinson's blood-brain barrier and Crohn's intestinal vasculature. Peripheral blood analyses
revealed significant enrichment of immune-related GO terms in Parkinson's disease, including "MHC protein complex"[35],
"antigen binding"[36], and "immunoglobulin complex". The circulatory specificity was further evidenced by "humoral immune
response mediated by circulating immunoglobulin". Similarly, in Crohn's peripheral blood, neutrophil-related GO terms ("neu-
trophil degranulation", "neutrophil activation involved in immune response", "neutrophil mediated immunity", and "neutrophil
activation") indicated the involvement of immune-inflammatory processes[37]. Furthermore, significant Parkinson's "complement
activation" and Crohn's "blood coagulation" provided disease-specific vascular insights. Aberrant complement system activity
may imply blood-brain barrier disruption in Parkinson's patients[38, 39], whereas increased venous thromboembolism risk in
Crohn's patients due to abnormal coagulation[40] indicates intestinal vascular barrier impairment[41].

The ileal and colonic mucosa constitute primary pathological sites in Crohn's disease[34], where collagen plays a key role
in extracellular matrix remodeling[42], evidenced by the significant "collagen catabolic process". Gut microbiota dysbiosis,
reflected by GO terms "antimicrobial humoral response" and "response to lipopolysaccharide", further aligned with Crohn's
pathogenesis[43, 44]. While Crohn's ileal and colonic mucosa share multifaceted similarities, they displayed two key distinctions:

complement activation and digestion. Although both mucosal tissues showed complement activation, its enrichment ratios were
2-4 times higher in ileal versus colonic mucosa, supported by immunofluorescence staining and single-cell transcriptomics
indicating more active complement activation in ileal mucosa[45,46]. It has been generally acknowledged the ileal mucosa
uniquely mediates hydrolase-driven enzymatic digestion, whereas the colonic mucosa plays no substantive role in chemical
digestion[47]. This functional dichotomy was corroborated by the enrichments of the "digestion" GO term: AWmeta specifically
detected "digestion" within top 100 ileal mucosa genes (highest enrichment), whereas REM and baselines showed delayed
identification (top 300/500 genes; lower enrichment); conversely, AWmeta consistently excluded "digestion" from colonic
mucosa enrichments, in contrast to sporadic false positives by these counterparts. These functional stratifications demonstrate
AWmeta's enhanced biological fidelity in resolving disease tissue-contextual gene functions.

Discussion

Transcriptomic meta-analysis is pivotal for distilling robust biological insights from heterogeneous gene expression studies; yet,
existing frameworks remain confined to either *P*-value combination or effect-size integration, imposing a trade-off between
statistical sensitivity and quantitative fidelity. A unified strategy that seamlessly integrates both paradigms—capitalizing on
their complementary strengths while circumventing their individual limitations—has therefore been a long-standing, unmet
imperative.
AWmeta represents the first successful integration of *P*-value and effect-size aggregation methodologies in transcriptomic
meta-analysis. The core innovation—a cross-module information transfer where optimized weights from *P*-value calculations
directly enhance effect size estimation—effectively addresses between-study heterogeneity while maximizing consistent
biological signal extraction. Indeed, the substantial variability often observed between studies, visually apparent in metrics like
gene-wise convergence (Fig. 2b–f with per-study skewed distributions), highlights the prevalence of such heterogeneity and
strongly supports the usage of random-effects-like frameworks such as AWmeta and REM over simpler fixed-effects models[48].
In our comprehensive evaluation across 35 datasets from Parkinson's and Crohn's disease, AWmeta demonstrated superior
high-fidelity DEG detection that remained robust under substantial experimental noise (Fig. 1c,g–i and Extended Data Fig. 2–5).
This enhanced discrimination capacity enabled identification of subtle yet biologically meaningful expression changes that
conventional methods frequently miss, substantially improving the reliability and reproducibility of transcriptomic discoveries.
Our convergence metrics revealed AWmeta's practical advantages in approximating theoretical true values at both gene and
study levels. It's noteworthy that in our gene-wise convergence assessments, some original studies occasionally outperformed
standard REM, even without larger sample sizes (Fig. 2b–f and Extended Data Fig. 6). While not conclusive, this hints that
inherent study quality or specific experimental contexts might significantly influence reliability, perhaps as much as sample
size itself. We also observed a tendency for these well-performing studies to utilize RNA-seq technology. These observations
underscore the complexity of integrating diverse datasets and highlight the benefit of AWmeta's adaptive capability which
weights studies based on informational content rather than relying solely on metrics like sample size. AWmeta achieved
equivalent study-wise convergence with significantly fewer samples and DEGs than conventional methods—a critical advantage
in resource-constrained research environments that can substantially reduce experimental costs and researcher workload.
The superior biological relevance of AWmeta's findings was rigorously established through two orthogonal and comple-
mentary assessment paradigms: (1) Using authoritative disease-specific gene sets from DisGeNET, MalaCards, and in-house
genetic variant corpus, AWmeta demonstrated significantly enhanced biological meaningfulness. In 11 of 15 tissue-benchmark
combinations, AWmeta outperformed both REM and original studies in biological relevance scoring (Fig. 5c–e). This consistent
advantage provides researchers with more accurate representations of core disease pathways and creates unprecedented oppor-
tunities for discovering novel pathophysiological relationships that remain obscured in conventional analyses. (2) Longitudinal
tracking of GO term enrichment across gene rank thresholds revealed AWmeta's unique capacity to concentrate functionally
critical genes within leading ranks. While terms of secondary importance (e.g., "MHC protein complex", "blood coagulation")
showed delayed enrichment beyond top 300 ranks across all methods (Fig. 5f), pathologically central functions exhibited
exclusive early enrichment in AWmeta. Crucially, terms like "digestion" in Crohn's ileal mucosa reached peak enrichment
exclusively within AWmeta's top 100 genes (Fig. 5f), with no detection at expanded thresholds; REM failed to detect this
pivotal function at both top 100 and 300 thresholds, achieving only marginal detection at top 500 (Fig. 5f)—demonstrating its
fundamental limitations in biological resolution. This enrichment trajectory analysis establishes a dual-purpose paradigm for
quantitatively evaluating gene prioritization performance and objectively stratifying biological mechanisms by pathological
centrality. Together, these orthogonal validation strategies—leveraging curated knowledgebases and temporal enrichment
dynamics—provide compelling evidence that AWmeta uniquely reconciles statistical rigor with biological fidelity, transforming
heterogeneous transcriptomic data into precisely stratified mechanistic insights.
While AWmeta represents a significant advance, several aspects warrant consideration for broader application. The method's
computational complexity, though tractable for typical multi-study analyses, may require optimization for emerging consortia-
level datasets exceeding 100 studies. Performance is also intrinsically linked to input data quality; while adaptive weighting

mitigates variable study quality (Fig. 2b–f and Extended Data Fig. 6), unaddressed technical artifacts (e.g., severe batch effects) could subtly influence results. Furthermore, the current framework focuses on gene-level differential quantification; extension to isoform-resolution or splicing analysis would require adaptation to handle increased dimensionality. Finally, scenarios involving extreme, systematic confounding across studies (e.g., irreconcilable patient stratification) remain challenging—a limitation pervasive among meta-analytic methods.

These considerations highlight clear pathways for AWmeta's evolution, complementary to its core strengths. Algorithmic refinements such as distributed computing could enhance scalability for ultra-large-scale integrations. Furthermore, incorporating tissue-specific molecular networks or multi-omic layers (e.g., epigenomics, proteomics) would refine biological inference beyond expression-centric views. Notably, coupling AWmeta with pharmacological databases holds significant promise for *in silico* drug target prioritization and biomarker discovery. The framework also uniquely empowers researchers to integrate limited-scale local datasets with public repositories, democratizing access to robust meta-analysis and amplifying statistical power for domain investigations.

Crucially, the present AWmeta implementation already delivers immediate, high-impact utility. It establishes a robust new standard for extracting reliable biological signals from complex, heterogeneous transcriptomic data—overcoming a fundamental methodological dichotomy that has long constrained the field. By synergistically combining P-value and effect size paradigms, AWmeta enhances reproducibility and biological interpretability, directly accelerating the translation of transcriptomic discoveries into clinical insights and biotechnological applications. This methodological leap provides an integral tool for navigating the growing complexity of precision transcriptomic integration in biomedical research.

Methods

Overview of the AWmeta framework

The transformative potential of AWmeta stems from its adaptively weighting scheme, which strategically prioritizes the most informative studies while robustly mitigating noise and outliers to yield biologically coherent, high-fidelity meta-analytic estimates. This framework performs gene-by-gene meta-analysis of heterogeneous transcriptomic studies by integrating per-study gene summary statistics (Fig. 1a). For each gene, input from each study in the valid set (S_{gene}) comprises: (1) a P-value (P_i), (2) a $\log_2$-based fold change (FC_i), and (3) its corresponding within-study variance (Var_i), all derived from the original gene differential quantification analyses. AWmeta consists of two sequential modules: *AW-Fisher* for adaptive P-value aggregation and *AW-REM* for adaptive effect-size integration. Studies absent from S_{gene} (e.g., Study_2 with missing P_2, FC_2, or Var_2) are excluded a priori.

AW-Fisher module (adaptive P-value integration)

Within this module, each gene's meta P-value is obtained by selecting an optimal subset of S_{gene} that minimizes a weighted Fisher's statistic-derived combined P-value[9]. Let $N' = |S_{gene}|$ be the number of studies reporting P-values for the gene, with $S'_{gene} = \{1,\ldots,N'\}$ enumerating the study indices, and denote their P-values by $\overrightarrow{\mathbf{P}} = (P_1,\ldots,P_i)_{i \in S'_{gene}} \in (0,1)^{N'}$. The corresponding binary weight vector, $\overrightarrow{\mathbf{w}} = (w_1,\ldots,w_i)_{i \in S'_{gene}} \in \{0,1\}^{N'}$, indicates inclusion ($w_i = 1$) or exclusion ($w_i = 0$) of $\text{Study}_i \in S_{gene}$ in the final subset. The AW-Fisher statistic is defined as:

$$T(\overrightarrow{\mathbf{P}}; \overrightarrow{\mathbf{w}}) = -2 \sum_{i \in S'_{gene}} w_i \ln P_i \tag{2}$$

The significance level of $T(\overrightarrow{\mathbf{P}}; \overrightarrow{\mathbf{w}})$ under the null hypothesis is calculated using the chi-squared distribution:

$$L(T(\overrightarrow{\mathbf{P}}; \overrightarrow{\mathbf{w}})) = 1 - F_{\chi^2_{d(\overrightarrow{\mathbf{w}})}}(T(\overrightarrow{\mathbf{P}}; \overrightarrow{\mathbf{w}})) \tag{3}$$

where the degrees of freedom are $d(\overrightarrow{\mathbf{w}}) = 2\sum_{i \in S'_{gene}} w_i$, and $F_{\chi^2_d}(\cdot)$ is the cumulative distribution function of the chi-squared distribution with d degrees of freedom.

The meta P-value, $s(\overrightarrow{\mathbf{P}})$, is the minimum significance level obtained by optimizing the weight vector over the studies in S_{gene}:

$$s(\overrightarrow{\mathbf{P}}) = \min_{\overrightarrow{\mathbf{w}}} L(T(\overrightarrow{\mathbf{P}}; \overrightarrow{\mathbf{w}})) \tag{4}$$

The optimal weight vector $\hat{w}$ that achieves this minimum is determined by:

$$\hat{\boldsymbol{w}} = w(\overrightarrow{\boldsymbol{P}}) = \operatorname*{argmin}_{\overrightarrow{\mathbf{w}}} L(T(\overrightarrow{\mathbf{P}}; \overrightarrow{\mathbf{w}})) = (\hat{w}_1,\ldots,\hat{w}_i)_{i \in S'_{gene}} \tag{5}$$

This optimal weight vector $\hat{w}$, containing binary weights for each study in S_{gene}, is passed to the following AW-REM module.

AW-REM module (adaptive effect-size integration)

This module calculates the meta effect size ($\log_2 FC$) using an adaptively-weighted REM. It leverages the $\log_2 FC$ (FC_i) and within-study variance (Var_i) from studies in S_{gene}, modulated by the optimal binary weights $\hat{w}$ derived from the AW-Fisher module for those same studies. The contribution weight for $\text{Study}_i \in S_{gene}$ in AW-REM is defined as:

$$W_i = \frac{\hat{w}_i}{Var_i + T^2} \tag{6}$$

where $\hat{w}_i$ is the binary weight (0 or 1) for $\text{Study}_i \in S_{gene}$ from Eq. 5. Var_i is the within-study variance for the gene in Study_i, and T^2 represents the between-study variance, estimated using restricted maximum likelihood (REML) method[49]. W_i is zero whenever $\hat{w}_i = 0$, thus automatically omitting studies not selected by AW-Fisher module.

The final meta fold change, denoted $\mathbf{M}$, is computed as an adaptively calibrated average of the study-wise effect sizes:

$$\mathbf{M} = \frac{\sum_{i \in S'_{gene}} W_i FC_i}{\sum_{i \in S'_{gene}} W_i} \tag{7}$$

This formulation delivers a consensus fold-change estimate both statistically rigorous and quantitatively faithful to the most informative subsets of heterogeneous studies.

Transcriptomic datasets

To proof-of-concept the AWmeta framework, we compiled 35 publicly available human transcriptomic datasets for Parkinson's and Crohn's disease from the Gene Expression Omnibus (GEO)[50], Sequence Read Archive (SRA)[51], and ArrayExpress[52]. These datasets, encompassing both microarray and RNA-sequencing (RNA-seq) platforms, included samples derived from Parkinson's substantia nigra[53–60] and peripheral blood[56,61–68], Crohn's peripheral blood[69–71], ileal mucosa[72–77] and colonic mucosa[76–82]. A complete list of the datasets, detailing data accession IDs, sequencing platform identifiers, dataset and tissue sources, and patient and control sample sizes, is provided in Extended Data Fig. 1.

Transcriptomic data preprocessing

Due to the inclusion of datasets generated on different platforms, specific preprocessing pipelines were applied separately to microarray and RNA-seq data.

Microarray data processing

To ensure accurate and up-to-date probe annotations, microarray probe identifiers were mapped to Entrez Gene IDs using information retrieved from GEO SOFT files, platform-specific Bioconductor annotation packages, and the AnnoProbe R package. For genes represented by multiple probes, we retained the probe with the largest interquartile range (IQR) of intensities across samples to maximize biological informativeness[21,83]. Subsequently, the limma R package[84] was utilized for microarray data preprocessing, normalization, and differential gene identification. Gene differential quantification (case versus control) within each study was determined via empirical Bayes moderated t-statistics, yielding per-gene P_i, FC_i, and Var_i.

RNA-seq data processing

RNA-seq data were processed through an automated snakemake workflow[85]. Raw sequencing reads were processed with Trimmomatic[86] to remove adapter sequences and low-quality bases. Following best practice recommendations[87], the cleaned reads were aligned to the human reference genome (GRCh38 assembly) using HISAT2[88]. Gene-level read counts were quantified from the aligned reads using featureCounts[89]. Finally, we performed gene differential quantification with DESeq2[90], producing per-gene P_i, FC_i, and Var_i for each study.

Transcriptomic meta-analysis evaluation metrics

To impartially evaluate AWmeta's performance advances, we conducted a multi-dimensional comparison against the current gold-standard REM method[6,21,22] across the following critical analytical domains: (i) DEG detection capability, (ii) DEG discrimination, (iii) gene- and study-wise gene differential quantification convergence, (iv) stability and robustness, and (v) biological relevance. Both methods operated on matching inputs and identical gene sets, ensuring an equitable performance assessment.

DEG detection capability evaluation

DEG detection capability is defined as the gene count satisfying pre-defined thresholds for both corrected statistical significance P-value (FDR) and fold change magnitude ($|\log_2 FC|$) (Fig. 1b). To assess the stability and reliability of this capability, we implemented a bootstrap resampling strategy with 100 iterations. In each iteration, we created bootstrapped datasets by randomly sampling with replacement from the original case and control groups while maintaining the original sample sizes, followed by meta-analysis. The resulting DEG counts formed a distribution for statistical comparison with one-tailed Welch's t-test.

DEG discrimination evaluation using semi-synthetic simulation strategy

To evaluate the ability to discriminate between DEGs and non-DEGs, particularly considering potential false positives arising from higher detection sensitivity, we adopted an evaluation metric based on semi-synthetic simulated data, inspired by Li and colleagues[23]. This approach consisted of benchmark dataset generation and evaluation using datasets with simulated noise (Fig. 1d–f) and for each tissue context:

1. Identify the intersection of DEGs and non-DEGs called by both AWmeta and REM under predefined screening thresholds (Fig. 1d).

2. Randomly sample half of the intersected DEGs to form an unbiased positive benchmark; sample an equal-sized negative benchmark from the intersected non-DEGs (Fig. 1d).

3. Construct semi-synthetic datasets by permuting case/control labels within a subset of original studies (e.g., Study$_1$ and Study$_3$; Fig. 1e). Label permutation removes true signal from those studies.

4. Apply AWmeta and REM to the combined set of original and label-permuted studies; compute the AUROC and AUPRC over the previously defined positive and negative benchmark genes (Fig. 1e).

5. Repeat Steps 3 and 4 for 100 times to obtain distributions of AUROC and AUPRC, summarizing performance under minimum-, median-, and maximum-permuted scenarios (Fig. 1f), which ensures the stability and reliability of our assessment. Statistical significance between AWmeta and REM was tested via one-tailed Mann–Whitney test.

Gene-wise convergence assessment for gene differential quantification

To assess per-gene differential quantification agreement between meta-analysis (FC_{meta}) and original constituent studies (FC_i), we computed a MAD-like gene-wise convergence score (Fig. 2a). For each gene $\mathscr{G}$:

$$C_{\mathscr{G}meta} = \frac{1}{|S_{gene}|} \sum_{i \in S'_{gene}} |FC_{meta} - FC_i| \tag{8}$$

where S_{gene} and S'_{gene} denote the valid study set and corresponding indices for the gene (defined in "Overview of the AWmeta framework" section), $|S_{gene}|$ the cardinality of the set S_{gene}, $FC_{meta} = \mathbf{M}$ from Eq. 7 and FC_i is the study-exclusive $\log_2$-based fold change. A lower $C_{\mathscr{G}meta}$ implies better agreement between the meta-analysis and original study estimates within S_{gene}. For baseline comparison, we also computed, for each original Study$_j \in S_{gene}$, a gene-wise convergence score:

$$C_{\mathscr{G}j} = \frac{1}{|S_{gene}|} \sum_{i \in S'_{gene}} |FC_j - FC_i| \tag{9}$$

which represents the MAD of Study$_j$'s fold change from all other contributing studies. This internal consistency benchmark enables direct contrast of AWmeta's and REM's convergence performance against the inherent agreement among the original datasets. All comparisons used one-tailed Mann–Whitney test against AWmeta.

Study-wise convergence assessment for gene differential quantification

To rigorously evaluate the consistency between gene lists derived from meta-analysis methods and those from the original studies, we employed three complementary approaches. For all metrics, higher scores indicate superior study-level convergence.

1. **Adjusted rankeD genE (DE) list similarity**: Our first approach quantifies concordance using a rank-sensitive similarity metric that is critically weighted towards top-ranked genes (Fig. 2g and Extended Data Fig. 7). To construct robustly ordered gene lists (G_{meta} for the meta-analysis; G_i for Study$_i$), we first devised a composite rank for each gene by multiplying its P-value rank (ascending) with its $|\log_2 FC|$ rank (descending), thereby integrating statistical significance and effect size.

   The weighted similarity $S(G_{meta}, G_i)$ between the meta-analysis and each original study gene lists (containing N genes) was computed using a non-linear weighting scheme[91], which emphasizes the top-ranked gene concordance:

$$S(G_{meta}, G_i) = \sum_{n=1}^{N} e^{-\alpha n} O_n(G_{meta}, G_i) \tag{10}$$

   where $O_n(G_{meta}, G_i)$ is the number of common genes in the top n positions and α is a weighting exponent (0.001). This score was then normalized to the interval $[-1, 1]$[22] yielding the adjusted similarity:

$$S_{adj}(G_{meta}, G_i) = \frac{S(G_{meta}, G_i) - E_{\text{null}}(S(G_{meta}, G_i))}{\max(S(G_{meta}, G_i)) - E_{\text{null}}(S(G_{meta}, G_i))} \tag{11}$$

   where $E_{\text{null}}(S(G_{meta}, G_i)) = \sum_{n=1}^{N} \frac{n^2}{N} e^{-\alpha n}$ and $\max(S(G_{meta}, G_i)) = \sum_{n=1}^{N} n e^{-\alpha n}$ are the expected and maximum scores under a null hypothesis of random gene lists.

2. **Set-based overlap similarity**: To circumvent the limitations of the above rank-dependent approach, which is sensitive to gene ranking variations while potentially overlooking consistent differential expression patterns, we assessed study-wise convergence using a set-based overlap metric that exclusively evaluates binary DEG classification concordance (Fig. 2h,i). Here, DEG sets were determined for both the meta-analysis (Set_{meta}) and individual studies (Set_i) using predefined statistical thresholds, thereby focusing analytical power on reproducible differential expression status irrespective of positional gene rankings.

   We calculated two complementary metrics: JC ($JC_i = |Set_{meta} \cap Set_i|/|Set_{meta} \cup Set_i|$) and OC ($OC_i = |Set_{meta} \cap Set_i|/\min(|Set_{meta}|, |Set_i|)$). The convergence metric for the meta-analysis relative to Study$_i$ was the arithmetic mean $(JC_i + OC_i)/2$.

3. **Phi coefficient similarity**: Finally, we measured the association between DEG classifications using phi coefficient[24] (ϕ) (Fig. 2h,j). This approach considers the extreme case where shared DEGs or non-DEGs between two gene sets might be randomly generated.

For each comparison between the meta-analysis and an original Study$_i$, we constructed a 2×2 contingency table categorizing all genes as DEG or non-DEG in both datasets and the phi coefficient (ϕ_i) was then calculated as:

$$\phi_i = \frac{n_{11}n_{22} - n_{12}n_{21}}{\sqrt{n_{1\cdot}n_{2\cdot}n_{\cdot1}n_{\cdot2}}} \tag{12}$$

where n_{11} represents DEGs, n_{22} non-DEGs in both datasets, and n_{12} and n_{21} represent exclusively-classified DEGs for the binary datasets. The row and column sums are denoted by $n_{1\cdot}$, $n_{2\cdot}$, $n_{\cdot1}$, and $n_{\cdot2}$.

To establish a performance baseline, we computed all three convergence metrics for every pairwise combination of the original studies. Overall study-wise convergence differences among AWmeta, REM and baselines were tested by Kruskal–Wallis test, followed by Nemenyi post-hoc test for pairwise comparisons.

For the set-based and ϕ metrics, which rely on binary DEG and non-DEG classification, we note that these outcomes are mutually exclusive and complementary, and therefore report the results derived from the DEG sets for clarity and conciseness.

Stability and robustness assessment of transcriptomic integration

To demonstrate AWmeta's resilience, we evaluated both stability—against stochastic sampling—and robustness—against dataset perturbations—using the adjusted DE list similarity ("Study-wise convergence assessment for gene differential quantification" section and Extended Data Fig. 7).

1. **Within-study subsampling stability**: For each original cohort, we randomly partitioned case and control samples of every study into two equal subcohorts, yielding paired "half-study" datasets. Each half-study set underwent independent DEG analysis and subsequent meta-analysis. The similarity between the resulting ordered gene lists was computed over 100 bootstrap replicates, quantifying stability under within-study sampling (Fig. 4a). AWmeta and REM stability distributions were compared via one-tailed Welch's t-test.

2. **External robustness**: We assessed resilience to new data by sequentially incorporating one independent external study (from a held-out pool) into the original meta-analysis (Fig. 4c). For each addition, we performed meta-analysis pre- and post-inclusion, then computed adjusted DE list similarity between resulting ordered gene lists, measuring the impact of disparate external data (AWmeta versus REM, one-tailed Mann–Whitney test).

3. **Internal robustness**: We evaluated sensitivity to study omission by performing leave-one-study-out analyses (Fig. 4e): each original study was removed in turn, and meta-analyses were rerun on the reduced datasets. The similarity between each leave-one-study-out and the full-cohort ranked gene lists, across all iterations, quantified internal robustness (AWmeta versus REM, one-tailed Mann–Whitney test).

Biological relevance assessment of gene differential quantification

To quantify disease-context relevance of gene differential quantification, we assembled benchmark gene sets for Parkinson's and Crohn's disease from three sources: (1) DisGeNET with gene-disease association (GDA) score > 0.2[92, 93], (2) MalaCards, and (3) our in-house curated disease-related genetic variation corpus. For reference comparisons, original study-derived biological relevance results serve as baselines.

For each method (AWmeta, REM or baselines), all analyzed genes were ranked twice—(i) by descending $|\log_2 FC|$, (ii) by ascending P-value—then each benchmark gene's ranks were multiplied:

$$\text{Integrated Rank} = \text{Rank}_{|\log_2 FC|} \times \text{Rank}_{P\text{-value}} \tag{13}$$

Benchmark genes were then re-ranked according to this Integrated Rank (ascending) to obtain $\text{Rank}_{\text{Integrated Rank}}$. The Biological Relevance score was calculated for each benchmark gene as:

$$\text{Biological Relevance} = 1 - \frac{\text{Rank}_{\text{Integrated Rank}}}{N} \tag{14}$$

where N is the size of gene list from AWmeta, REM or baselines. Higher scores reflect greater biological relevance, signifying that benchmark genes attain superior rankings through combining statistical significance and fold change (Fig. 5a). This rank-based score accounts for gene list size heterogeneity and avoids arbitrary DEG thresholds. We compared biological relevance distributions from AWmeta, REM and baselines using Kruskal-Wallis and Nemenyi post-hoc test.

Data availability

All transcriptomic datasets used in this study are publicly available via GEO, SRA, and ArrayExpress. Parkinson's disease data include substantia nigra (GEO accessions: `GSE114517`, `GSE8397`, `GSE20163`, `GSE20164`, `GSE20292`, `GSE7621`, `GSE49036`, `GSE42966`, `GSE43490`, `GSE26927`, `GSE54282`) and peripheral blood (GEO accessions: `GSE57475`, `GSE54536`, `GSE34287`, `GSE99039`, `GSE72267`, `GSE6613`, `GSE18838`, `GSE165082`). Crohn's disease datasets include peripheral blood (GEO accessions: `GSE119600`, `GSE112057`, `GSE94648`), ileal mucosa (GEO accessions: `GSE102133`, `GSE75214`, `GSE16879`, `GSE68570`, `GSE101794`, `GSE57945`), and colonic mucosa (GEO accessions: `GSE75214`, `GSE16879`, `GSE36807`, `GSE4183`, `GSE9686`, `GSE66207`; ArrayExpress accession: `E-MTAB-184`). Benchmark gene sets for both diseases were retrieved from DisGeNET (`https://www.disgenet.com/`) and MalaCards (`https://www.malacards.org/`).

Code availability

AWmeta is available on GitHub at `https://github.com/YanshiHu/AWmeta`.

References

1. Stark, R., Grzelak, M. & Hadfield, J. Rna sequencing: the teenage years. *Nat. Rev. Genet.* **20**, 631–656 (2019).

2. Shorten, A. & Shorten, B. What is meta-analysis? *Evid. Based Nurs.* **16**, 3–4 (2013).

3. Piras, I. S. *et al.* Peripheral biomarkers in schizophrenia: a meta-analysis of microarray gene expression datasets. *Int. J. Neuropsychopharmacol.* **22**, 186–193 (2019).

4. Zhong, G., Bolitho, S., Grunstein, R., Naismith, S. L. & Lewis, S. J. G. The relationship between thermoregulation and rem sleep behaviour disorder in parkinson's disease. *PLoS One* **8**, e72661 (2013).

5. Cho, H. *et al.* Meta-analysis method for discovering reliable biomarkers by integrating statistical and biological approaches: An application to liver toxicity. *Biochem. Biophys. Res. Commun.* **471**, 274–281 (2016).

6. Makinde, F. L. *et al.* Reviewing and assessing existing meta-analysis models and tools. *Briefings Bioinforma.* **22**, bbab324 (2021).

7. Fisher, R. A. Statistical methods for research workers. In *Breakthroughs in statistics: Methodology and distribution*, 66–70 (Springer, 1970).

8. Stouffer, S. A. A study of attitudes. *Sci. Am.* **180**, 11–15 (1949).

9. Li, J. & Tseng, G. C. An adaptively weighted statistic for detecting differential gene expression when combining multiple transcriptomic studies. *Annals Appl. Stat.* **5**, 994–1019 (2011).

10. Wang, X. *et al.* Comparing methods for performing trans-ethnic meta-analysis of genome-wide association studies. *Hum. molecular genetics* **22**, 2303–2311 (2013).

11. Toro-Domínguez, D. *et al.* A survey of gene expression meta-analysis: methods and applications. *Briefings Bioinforma.* **22**, 1694–1705 (2021).

12. Romano, S. *et al.* Machine learning-based meta-analysis reveals gut microbiome alterations associated with parkinson's disease. *Nat. Commun.* **16**, 1–17 (2025).

13. Yang, J., Liu, Q. & Shyr, Y. A large-scale meta-analysis reveals positive feedback between macrophages and t cells that sensitizes tumors to immunotherapy. *Cancer research* **84**, 626–638 (2024).

14. Powell, N. R. *et al.* Clinically important alterations in pharmacogene expression in histologically severe nonalcoholic fatty liver disease. *Nat. Commun.* **14**, 1474 (2023).

15. Langan, D. Assessing heterogeneity in random-effects meta-analysis. In *Meta-research: methods and protocols*, 67–89 (Springer, 2021).

16. Breitling, R., Armengaud, P., Amtmann, A. & Herzyk, P. Rank products: a simple, yet powerful, new method to detect differentially regulated genes in replicated microarray experiments. *FEBS Lett.* **573**, 83–92 (2004).

17. Del Carratore, F. *et al.* Rankprod 2.0: a refactored bioconductor package for detecting differentially expressed features in molecular profiling datasets. *Bioinformatics* **33**, 2774–2775 (2017).

18. Tseng, G. C., Ghosh, D. & Feingold, E. Comprehensive literature review and statistical considerations for microarray meta-analysis. *Nucleic acids research* **40**, 3785–3799 (2012).

19. Aguzzoli Heberle, B. *et al.* Systematic review and meta-analysis of bulk rnaseq studies in human alzheimer's disease brain
tissue. *Alzheimer's & Dementia* **21**, e70025 (2025).

20. Stogiannis, D., Siannis, F. & Androulakis, E. Heterogeneity in meta-analysis: a comprehensive overview. *The Int. J.*
Biostat. **20**, 169–199 (2024).

21. Ma, T. *et al.* Metaomics: analysis pipeline and browser-based software suite for transcriptomic meta-analysis. *Bioinformat-*
ics **35**, 1597–1599 (2019).

22. Chang, L.-C., Lin, H.-M., Sibille, E. & Tseng, G. C. Meta-analysis methods for combining multiple expression profiles:
comparisons, statistical characterization and an application guideline. *BMC Bioinforma.* **14**, 368 (2013).

23. Li, Y., Ge, X., Peng, F., Li, W. & Li, J. J. Exaggerated false positives by popular differential expression methods when
analyzing human population samples. *Genome Biol.* **23**, 79 (2022).

24. Akoglu, H. User's guide to correlation coefficients. *Turkish J. Emerg. Medicine* **18**, 91–93 (2018).

25. Piñero, J. *et al.* The disgenet knowledge platform for disease genomics: 2019 update. *Nucleic acids research* **48**,
D845–D855 (2020).

26. Rappaport, N. *et al.* Malacards: an amalgamated human disease compendium with diverse clinical and genetic annotation
and structured search. *Nucleic acids research* **45**, D877–D887 (2017).

27. Hu, Y. *et al.* Mapping the comorbid landscape of parkinson's disease and crohn's disease along the gut-blood-brain axis.
medRxiv 2025–10 (2025).

28. Hu, Y., Pan, Z., Hu, Y., Zhang, L. & Wang, J. Network and pathway-based analyses of genes associated with parkinson's
disease. *Mol. neurobiology* **54**, 4452–4465 (2017).

29. Burke, R. E. & O'Malley, K. Axon degeneration in parkinson's disease. *Exp. neurology* **246**, 72–83 (2013).

30. Chen, L. *et al.* Homeostasis and metabolism of iron and other metal ions in neurodegenerative diseases. *Signal Transduct.*
Target. Ther. **10**, 31 (2025).

31. Bjorklund, G. *et al.* Metals and parkinson's disease: mechanisms and biochemical processes. *Curr. Medicinal Chem.* **25**,
2198–2214 (2018).

32. Ma, K. Y., Fokkens, M. R., Reggiori, F., Mari, M. & Verbeek, D. S. Parkinson's disease–associated vps35 mutant reduces
mitochondrial membrane potential and impairs pink1/parkin-mediated mitophagy. *Transl. neurodegeneration* **10**, 1–17
(2021).

33. Poewe, W. *et al.* Parkinson disease. *Nat. reviews Dis. primers* **3**, 1–21 (2017).

34. Roda, G. *et al.* Crohn's disease. *Nat. Rev. Dis. Primers* **6**, 22 (2020).

35. Gu, R. *et al.* The major histocompatibility complex participates in parkinson's disease. *Pharmacol. Res.* 107168 (2024).

36. Tan, E.-K. *et al.* Parkinson disease and the immune system—associations, mechanisms and therapeutics. *Nat. Rev. Neurol.*
16, 303–318 (2020).

37. Danne, C., Skerniskyte, J., Marteyn, B. & Sokol, H. Neutrophils: from ibd to the gut microbiota. *Nat. Rev. Gastroenterol.*
& Hepatol. **21**, 184–197 (2024).

38. Nimmo, J. *et al.* The complement system in neurodegenerative diseases. *Clin. Sci.* **138**, 387–412 (2024).

39. Khosousi, S. *et al.* Complement system changes in blood in parkinson's disease and progressive supranuclear
palsy/corticobasal syndrome. *Park. & Relat. Disord.* **108**, 105313 (2023).

40. Rogler, G., Singh, A., Kavanaugh, A. & Rubin, D. T. Extraintestinal manifestations of inflammatory bowel disease: current
concepts, treatment, and implications for disease management. *Gastroenterology* **161**, 1118–1132 (2021).

41. Stadnicki, A. & Stadnicka, I. Venous and arterial thromboembolism in patients with inflammatory bowel diseases. *World*
journal gastroenterology **27**, 6757 (2021).

42. Mortensen, J. *et al.* The intestinal tissue homeostasis–the role of extracellular matrix remodeling in inflammatory bowel
disease. *Expert. review gastroenterology & hepatology* **13**, 977–993 (2019).

43. Caruso, R., Lo, B. C. & Núñez, G. Host–microbiota interactions in inflammatory bowel disease. *Nat. Rev. Immunol.* **20**,
411–426 (2020).

44. Candelli, M. *et al.* Interaction between lipopolysaccharide and gut microbiota in inflammatory bowel diseases. *Int. journal*
molecular sciences **22**, 6242 (2021).

45. Halstensen, T., Mollnes, T., Garred, P., Fausa, O. & Brandtzaeg, P. Surface epithelium related activation of complement differs in crohn's disease and ulcerative colitis. *Gut* **33**, 902–908 (1992).

46. Kong, L. *et al.* The landscape of immune dysregulation in crohn's disease revealed through single-cell transcriptomic profiling in the ileum and colon. *Immunity* **56**, 444–458 (2023).

47. Feldman, M., Friedman, L. S. & Brandt, L. J. *Sleisenger and Fordtran's gastrointestinal and liver disease E-book: pathophysiology, diagnosis, management* (Elsevier health sciences, 2020).

48. Borenstein, M., Hedges, L. V., Higgins, J. P. & Rothstein, H. R. A basic introduction to fixed-effect and random-effects models for meta-analysis. *Res. synthesis methods* **1**, 97–111 (2010).

49. Kenward, M. G. & Roger, J. H. Small sample inference for fixed effects from restricted maximum likelihood. *Biometrics* **53**, 983–997 (1997).

50. Barrett, T. *et al.* Ncbi geo: archive for functional genomics data sets—update. *Nucleic Acids Res.* **41**, D991–D995 (2012).

51. Sayers, E. W. *et al.* Database resources of the national center for biotechnology information. *Nucleic Acids Res.* **47**, D23–D28 (2019).

52. Kolesnikov, N. *et al.* Arrayexpress update—simplifying data submissions. *Nucleic Acids Res.* **43**, D1113–D1116 (2015).

53. Lewis, P. *et al.* Evidence for immune response, axonal dysfunction and reduced endocytosis in the substantia nigra in early stage parkinson's disease. *PLoS One* **10**, e0128651 (2015).

54. Cookson, M. R. *et al.* Systems-based analyses of brain regions functionally impacted in parkinson's disease reveals underlying causal mechanisms. *PLoS ONE* **9**, e102909 (2014).

55. Simchovitz, A. *et al.* A lncrna survey finds increases in neuroprotective linc-pint in parkinson's disease substantia nigra. *Aging Cell* **19**, e13115 (2020).

56. Duke, D. C., Moran, L. B., Pearce, R. K. B. & Graeber, M. B. The medial and lateral substantia nigra in parkinson's disease: mrna profiles associated with higher brain tissue vulnerability. *Neurogenetics* **8**, 83–94 (2007).

57. Leal, S. M. *et al.* A genomic pathway approach to a complex disease: axon guidance and parkinson disease. *PLoS Genet.* **3**, e98 (2007).

58. Corradini, B. R. *et al.* Complex network-driven view of genomic mechanisms underlying parkinson's disease: analyses in dorsal motor vagal nucleus, locus coeruleus, and substantia nigra. *BioMed Res. Int.* **2014**, 1–16 (2014).

59. Zheng, B. *et al.* Pgc-1alpha, a potential therapeutic target for early intervention in parkinson's disease. *Sci Transl Med* **2**, 52ra73 (2010).

60. Durrenberger, P. F. *et al.* Selection of novel reference genes for use in the human central nervous system: a brainnet europe study. *Acta Neuropathol.* **124**, 893–903 (2012).

61. Cookson, M. R. *et al.* Srrm2, a potential blood biomarker revealing high alternative splicing in parkinson's disease. *PLoS One* **5**, e9104 (2010).

62. Scherzer, C. R. *et al.* Gata transcription factors directly regulate the parkinson's disease-linked gene -synuclein. *Proc. Natl. Acad. Sci.* **105**, 10907–10912 (2008).

63. Calligaris, R. *et al.* Blood transcriptomics of drug-naïve sporadic parkinson's disease patients. *BMC Genomics* **16**, 876 (2015).

64. Mosley, R. L. *et al.* Biosignatures for parkinson's disease and atypical parkinsonian disorders patients. *PLoS One* **7**, e43595 (2012).

65. Locascio, J. J. *et al.* Association between -synuclein blood transcripts and early, neuroimaging-supported parkinson's disease. *Brain* **138**, 2659–2671 (2015).

66. Alieva, A. K. *et al.* Involvement of endocytosis and alternative splicing in the formation of the pathological process in the early stages of parkinson's disease. *BioMed Res. Int.* **2014**, 1–6 (2014).

67. Shamir, R. *et al.* Analysis of blood-based gene expression in idiopathic parkinson disease. *Neurology* **89**, 1676–1683 (2017).

68. Henderson, A. R. *et al.* Dna methylation and expression profiles of whole blood in parkinson's disease. *Front. Genet.* **12**, 640266 (2021).

69. Planell, N. *et al.* Usefulness of transcriptional blood biomarkers as a non-invasive surrogate marker of mucosal healing and endoscopic response in ulcerative colitis. *J. Crohn's Colitis* **11**, 1335–1346 (2017).

70. Mo, A. *et al.* Disease-specific regulation of gene expression in a comparative analysis of juvenile idiopathic arthritis and inflammatory bowel disease. *Genome Medicine* **10**, 48 (2018).

71. Ostrowski, J. *et al.* Common functional alterations identified in blood transcriptome of autoimmune cholestatic liver and inflammatory bowel diseases. *Sci. Reports* **9**, 7190 (2019).

72. Haberman, Y. *et al.* Pediatric crohn disease patients exhibit specific ileal transcriptome and microbiome signature. *J. Clin. Investig.* **124**, 3617–3633 (2014).

73. Haberman, Y. *et al.* Age-of-diagnosis dependent ileal immune intensification and reduced alpha-defensin in older versus younger pediatric crohn disease patients despite already established dysbiosis. *Mucosal Immunol.* **12**, 491–502 (2019).

74. Pérez-Torras, S. *et al.* Transportome profiling identifies profound alterations in crohn's disease partially restored by commensal bacteria. *J. Crohn's Colitis* **10**, 850–859 (2016).

75. Verstockt, S. *et al.* Gene and mirna regulatory networks during different stages of crohn's disease. *J. Crohn's Colitis* **13**, 916–930 (2019).

76. Timmer, A. *et al.* Mucosal gene expression of antimicrobial peptides in inflammatory bowel disease before and after first infliximab treatment. *PLoS One* **4**, e7984 (2009).

77. Vancamelbeke, M. *et al.* Genetic and transcriptomic bases of intestinal epithelial barrier dysfunction in inflammatory bowel disease. *Inflamm. Bowel Dis.* **23**, 1718–1729 (2017).

78. van Beelen Granlund, A. *et al.* Reg gene expression in inflamed and healthy colon mucosa explored by in situ hybridisation. *Cell Tissue Res.* **352**, 639–646 (2013).

79. Carey, R. *et al.* Activation of an il-6:stat3-dependent transcriptome in pediatric-onset inflammatory bowel disease. *Inflamm. Bowel Dis.* **14**, 446–457 (2008).

80. Galamb, O. *et al.* Inflammation, adenoma and cancer: objective classification of colon biopsy specimens with gene expression signature. *Dis. Markers* **25**, 1–16 (2008).

81. Calogero, R. A. *et al.* Identification of novel predictor classifiers for inflammatory bowel disease by gene expression profiling. *PLoS One* **8**, e76235 (2013).

82. Peck, B. C. E. *et al.* Micrornas classify different disease behavior phenotypes of crohns disease and may have prognostic utility. *Inflamm. Bowel Dis.* **21**, 2178–2187 (2015).

83. Gentleman, R., Carey, V., Huber, W., Irizarry, R. & Dudoit, S. *Bioinformatics and computational biology solutions using R and Bioconductor* (Springer Science & Business Media, 2005).

84. Ritchie, M. E. *et al.* limma powers differential expression analyses for rna-sequencing and microarray studies. *Nucleic Acids Res.* **43**, e47–e47 (2015).

85. Köster, J. & Rahmann, S. Snakemake—a scalable bioinformatics workflow engine. *Bioinformatics* **28**, 2520–2522 (2012).

86. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for illumina sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

87. Sahraeian, S. M. E. *et al.* Gaining comprehensive biological insight into the transcriptome by performing a broad-spectrum rna-seq analysis. *Nat. Commun.* **8**, 59 (2017).

88. Kim, D., Paggi, J. M., Park, C., Bennett, C. & Salzberg, S. L. Graph-based genome alignment and genotyping with hisat2 and hisat-genotype. *Nat. Biotechnol.* **37**, 907–915 (2019).

89. Liao, Y., Smyth, G. K. & Shi, W. featurecounts: an efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics* **30**, 923–930 (2014).

90. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for rna-seq data with deseq2. *Genome biology* **15**, 1–21 (2014).

91. Yang, X., Bentink, S., Scheid, S. & Spang, R. Similarities of ordered gene lists. *J. Bioinforma. Comput. Biol.* **4**, 693–708 (2011).

92. Yuan, X. *et al.* The common genes involved in the pathogenesis of alzheimer's disease and type 2 diabetes and their implication for drug repositioning. *Neuropharmacology* **223**, 109327 (2023).

93. Ferreira, J. C., Alshamali, F., Pereira, L. & Fernandes, V. Characterization of arabian peninsula whole exomes: contributing to the catalogue of human diversity. *iScience* **25**, 105336 (2022).

Acknowledgements

The authors thank all members from Ming Chen's Group of Bioinformatics for insightful discussions. This work was supported by the National Key Research and Development Program of China (2023YFE0112300); National Natural Sciences Foundation of China (32261133526; 32270709; 32070677); the 151 talent project of Zhejiang Province (first level), the Science and Technology Innovation Leading Scientist (2022R52035).

Author contributions

Y.S.H. conceptualized the study. M.C. supervised the study. Y.S.H., Z.W., and Y.H. performed data analysis. Y.S.H., Z.W., C.F., and Q.F. contributed to writing the manuscript. All authors reviewed and approved the final manuscript.

Competing interests

The authors declare no competing interests.

Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Fig. 1 | Overview of AWmeta and DEG identification evaluation. a, Schematic of the AWmeta framework, comprising the AW-Fisher module for *P*-value calculation and the AW-REM module for effect size (log_2-transformed fold change) estimation. DEG identification evaluation includes DEG detection capability (**b, c**) and discrimination (**d–i**). **b,** Schematic defining DEG detection capability as the number of DEGs identified by predefined statistical significance (*P*-value) and gene expression fold change (log_2-based) thresholds. **c,** DEG detection capability performance comparisons between AWmeta and REM with corrected *P*-value (FDR) < 0.01 and fold change ($|log_2FC|$) > $log_2 1.2$ across five disease tissues. Statistical significance was determined with one-tailed Welch's *t*-test. **d,** Strategy for generating the semi-synthetic benchmark dataset, sampling equivalent DEGs and non-DEGs from common genes identified by both AWmeta and REM. **e,** Workflow for evaluating DEG discrimination performance using sample label permutation within the semi-synthetic benchmark dataset, followed by AWmeta/REM procedure and AUROC/AUPRC calculation. **f,** Study permutation statistics (number of permuted studies) in the DEG discrimination evaluation procedure across five disease tissues. **g–i**, DEG discrimination performance comparisons between AWmeta and REM using minimum- (**g**), median- (**h**), and maximum- (**i**) permuted semi-synthetic simulation strategy with FDR < 0.01 and $|log_2FC|$ > $log_2 1.2$ across five disease tissues. Statistical significance was determined using one-tailed Mann-Whitney test (**g–i**). Textual details of the AWmeta framework, DEG detection capability, and minimum-, median- and maximum-permuted semi-synthetic simulation strategies for DEG discrimination, reside in "DEG detection capability evaluation" and "DEG discrimination evaluation using semi-synthetic simulation strategy" sections in Methods. Boxplot bounds indicate interquartile ranges (IQR), centers denote median values, and whiskers extend to 1.5×IQR. The following icons represent different tissue sources: 👩: substantia nigra; 🩸: peripheral blood; 🦠: ileal mucosa; 🫃: colonic mucosa.

Fig. 2 | AWmeta establishes superior gene- and study-wise convergence in gene differential quantification. To benchmark meta-analysis efficacy across biological scales, we assessed gene differential quantification convergence in gene- (**a–f**) and study- (**g–m**) wise manners. **a–f**, For gene-wise convergence, mean absolute deviation (MAD) -like similarity measure was utilized to quantify the per-gene fold change ($|\log_2 FC|$) similarity among AWmeta, REM and original studies (**a**), with smaller value indicating better convergence, which demonstrates AWmeta's superior gene-wise convergence over REM and original studies across five disease tissues: Parkinson's substantia nigra (**b**) and peripheral blood (**c**), and Crohn's peripheral blood (**d**), ileal mucosa (**e**) and colonic mucosa (**f**). Statistical significance against AWmeta for gene-wise convergence comparisons was determined by one-tailed Mann-Whitney test. **g–m**, Three complementary similarities, i.e., adjusted DE list similarity (**g**, conceptual schematic; **k**, assessment result), the arithmetic average of Jaccard (JC) and overlap coefficient (OC) (**h, i**, conceptual schematic; **l**, assessment result) and phi coefficient (PC) (**h, j**, conceptual schematic; **m**, assessment result), were used to derive study-wise convergence score, indicating AWmeta exerts better convergence than REM and baselines in five disease tissues with FDR < 0.05 and $|\log_2 FC| > \log_2 1.2$. For comparison purpose, results from original studies serve as reference baselines. Overall study-wise convergence differences among AWmeta, REM and baselines were tested with Kruskal–Wallis test, followed by Nemenyi post-hoc test for pairwise comparisons. Detailed description for MAD-like gene- and three study-wise convergence similarity measures can be referred to in "Gene-wise convergence assessment for gene differential quantification" and "Study-wise convergence assessment for gene differential quantification" sections in Methods and Extended Data Fig. 7. Boxplot bounds show interquartile ranges (IQR), centers indicate median values, and whiskers extend to 1.5×IQR. The following icons represent different tissue sources: 🧠: substantia nigra; 🩸: peripheral blood; 🦠: ileal mucosa; 🫁: colonic mucosa.

Fig. 3 | AWmeta attains accelerated meta-analysis convergence with reduced samples and DEGs. a–f, We dissected the correlations of per-study sample size and DEG number with study-wise convergence measured by three complementary similarities, i.e., adjusted DE list similarity (**a, d**), the arithmetic average of Jaccard (JC) and overlap coefficient (OC) (**b, e**) and phi coefficient (PC) (**c, f**) across five disease tissues, showcasing AWmeta's accelerated study-wise meta-analysis convergence against REM with fewer samples and DEGs, in which non-linear correlations were quantified by spearman's rho with two-tailed significance test. To enhance statistical power and mitigate threshold sensitivity, this correlation analysis summarized results from nine different DEG threshold combinations, spanning varying significance levels (0.01, 0.05 and 0.10) and fold change cutoffs ($\log_2 1.2$, $\log_2 1.5$ and $\log_2 2.0$), for both the average of JC and OC and PC similarity measures. Detailed description for these three study-wise convergence similarity measures can be referred to in "Study-wise convergence assessment for gene differential quantification" section in Methods, Fig. 2g–j and Extended Data Fig. 7. The following icons represent different tissue sources: 🧠: substantia nigra; 🩸: peripheral blood; 🦠: ileal mucosa; 🫃: colonic mucosa.

Fig. 4 | AWmeta delivers remarkable stability and robustness in transcriptomic integration. a, Workflow for evaluating the stability of transcriptomic integration. **b,** Stability assessment against AWmeta and REM with one-tailed Welch's *t*-test. **c–f**, We assessed robustness against external interference (**c,** conceptual schematic; **d,** assessment result) and internal defects (**e,** conceptual schematic; **f,** assessment result) in transcriptomic integration across five disease tissues, with one-tailed Mann-Whitney test, indicating AWmeta's superior performance over REM. Detailed description of adjusted DE list similarity measure can be referred to in "Study-wise convergence assessment for gene differential quantification" section in Methods and Extended Data Fig. 7. Boxplot bounds show interquartile ranges (IQR), centers indicate median values, and whiskers extend to 1.5×IQR. The following icons represent different tissue sources: 👐: substantia nigra; 🩸: peripheral blood; 🧠: ileal mucosa; 🫀: colonic mucosa.

Fig. 5 | AWmeta enhances identification and mechanism interpretation of disease tissue-specific genes. a, Workflow for quantifying gene-wise biological relevance against three benchmark gene sets of Parkinson's and Crohn's disease from genetic association (GA) variation corpus, DisGeNET and MalaCards, with higher scores indicating stronger tissue-contextual disease associations. **b,** Pairwise coherence analysis of the three benchmark gene sets for Parkinson's and Crohn's disease. The degree of overlap between benchmarks was quantified using odds ratios (OR), with statistical significance determined by Fisher's exact test. **c–e**, Biological relevance evaluations on AWmeta (red), REM (blue) and baselines (green) against GA (**c**), DisGeNET (**d**) and MalaCards (**e**) benchmarks, where higher scores (y-axis) denote enhanced biological relevance. Original study-derived biological relevance results serve as reference baselines. We assessed overall differences in biological relevance scores using Kruskal-Wallis test, with Nemenyi post-hoc test for pairwise comparisons (AWmeta versus REM, AWmeta versus baseline, and REM versus baseline). Boxplot bounds indicate interquartile ranges (IQR), centers denote median values, and whiskers extend to 1.5×IQR. **f,** Dynamic GO enrichment trajectories across top-ranked (100, 300 and 500) genes identified by AWmeta (red), REM (blue) and baselines (green), with higher enrichment ratio (y-axis) indicating stronger disease-tissue involvement. Original study-derived enrichments serve as baselines. Connected lines visualize trajectory patterns across gene rank thresholds (x-axis). Textual details of the biological relevance assessment framework and detailed definition of enrichment ratio, reside in "Biological relevance assessment of gene differential quantification" section in Methods and "AWmeta enables disease tissue-specific mechanism interpretation" section in Results. The following icons represent different tissue sources: 👩: substantia nigra; 🩸: Peripheral blood; 🌰: ileal mucosa; 🫙: colonic mucosa.

Dataset	Sequencing platform	Dataset source	Tissue source	Samples	
Data IDs	Microarray or next-generation sequencing platforms	GEO, SRA and ArrayExpress	Blood, SN, IM and CM	No. of patients versus normals	

PD datasets

Data IDs	Sequencing platform	Patients	Normals
GSE57475	Illumina HumanHT-12 V3.0 expression beadchip	93	49
GSE54536	Illumina HumanHT-12 V4.0 expression beadchip	4	4
GSE34287	ExonHit Human Genome Wide SpliceArray 1.0	19	12
GSE99039	Affymetrix Human Genome U133 Plus 2.0 Array	205	233
GSE72267	Affymetrix Human Genome U133A 2.0 Array	40	19
GSE6613	Affymetrix Human Genome U133A Array	50	22
GSE18838	Affymetrix Human Exon 1.0 ST Array	17	11
GSE165082	Illumina HiSeq 2000	12	14
GSE114517	Illumina NextSeq 500	17	12
GSE8397	Affymetrix Human Genome U133A Array	24	13
GSE20163	Affymetrix Human Genome U133A Array	8	9
GSE20164	Affymetrix Human Genome U133A Array	6	5
GSE20292	Affymetrix Human Genome U133A Array	11	15
GSE7621	Affymetrix Human Genome U133 Plus 2.0 Array	16	9
GSE49036	Affymetrix Human Genome U133 Plus 2.0 Array	8	8
GSE42966	Agilent-014850 Whole Human Genome Microarray 4x44K G4112F	9	6
GSE43490	Agilent-014850 Whole Human Genome Microarray 4x44K G4112F	8	5
GSE26927	Illumina humanRef-8 v2.0 expression beadchip	12	8
GSE54282	Affymetrix Human Gene 1.0 ST Array	3	3

CD datasets

Data IDs	Sequencing platform	Patients	Normals
GSE119600	Illumina HumanHT-12 V4.0 expression beadchip	95	47
GSE112057	Illumina HiSeq 2000	60	12
GSE94648	Affymetrix Human Genome U133 Plus 2.0 Array	50	22
GSE102133	Affymetrix Human Gene 1.0 ST Array	65	12
GSE75214	Affymetrix Human Gene 1.0 ST Array	51	11
GSE16879	Affymetrix Human Genome U133 Plus 2.0 Array	18	6
GSE68570	Illumina HumanHT-12 V4.0 expression beadchip	6	5
GSE101794	Illumina HiSeq 2000	198	50
GSE57945	Illumina HiSeq 2000	163	42
GSE75214	Affymetrix Human Gene 1.0 ST Array	8	11
GSE16879	Affymetrix Human Genome U133 Plus 2.0 Array	19	6
GSE36807	Affymetrix Human Genome U133 Plus 2.0 Array	13	7
GSE4183	Affymetrix Human Genome U133 Plus 2.0 Array	5	8
GSE9686	Affymetrix Human Genome U133 Plus 2.0 Array	11	8
E-MTAB-184	Illumina HumanHT-12 v3.0 expression beadChip	15	32
GSE66207	Illumina HiSeq 2500	20	13

No. of Parkinson's (PD) / Crohn's disease (CD) patients: 4 8 16 32 64 128 256

No. of normals:

Legend icons: GEO, SRA, ArrayExpress, Peripheral blood, Substantia nigra (SN), Ileal mucosa (IM), Colonic mucosa (CM), No. of PD / CD patients, No. of normals

Extended Data Fig. 1 | Overview of transcriptomic datasets for Parkinson's and Crohn's disease. Datasets for Parkinson's disease (PD) include substantia nigra[53–60] and peripheral blood[56, 61–68]. Crohn's disease (CD) datasets comprise ileal mucosa[72–77], colonic mucosa[76–82], and peripheral blood[69–71]. Detailed metadata include data accession IDs, sequencing platform identifiers, dataset and tissue sources, and patient and control sample sizes. The following icons represent different tissue sources: 🌼: substantia nigra (SN); 🩸: peripheral blood; 🟤: ileal mucosa (IM); 🟩: colonic mucosa (CM).

Extended Data Fig. 2 | DEG detection capability evaluation across five disease tissues and diverse thresholds. a–e, DEG detection capability (the number of identified DEGs) was assessed against AWmeta and REM using nine distinct thresholds, combining three corrected P-values (FDR) (0.01, 0.05, and 0.10) and three $\log_2$-based fold change ($|\log_2FC|$) cutoffs ($\log_2 1.2$, $\log_2 1.5$, and $\log_2 2.0$), spanning Parkinson's substantia nigra (**a**), Parkinson's (**b**) and Crohn's (**c**) peripheral blood, and Crohn's ileal (**d**) and colonic (**e**) mucosa. Detailed description of DEG detection capability can be referred to in "DEG detection capability evaluation" section in Methods and Fig. 1b. Statistical significance was determined using one-tailed Welch's t-test. Boxplot bounds show interquartile ranges (IQR), centers indicate median values, and whiskers extend to $1.5 \times$IQR. The following icons represent different tissue sources: 🧠: substantia nigra; 🩸: peripheral blood; 🥮: ileal mucosa; 🫃: colonic mucosa.

Extended Data Fig. 3 | DEG discrimination evaluation across five disease tissues using minimum-permuted semi-synthetic simulation strategy. a–e, DEG discrimination (AUROC and AUPRC) was assessed against AWmeta and REM using nine distinct thresholds, combining three corrected *P*-values (FDR) (0.01, 0.05, and 0.10) and three $\log_2$-based fold change ($|\log_2 FC|$) cutoffs ($\log_2 1.2$, $\log_2 1.5$, and $\log_2 2.0$), spanning Parkinson's substantia nigra (**a**), Parkinson's (**b**) and Crohn's (**c**) peripheral blood, and Crohn's ileal (**d**) and colonic (**e**) mucosa. Detailed description of minimum-permuted semi-synthetic simulation strategy can be referred to in "DEG discrimination evaluation using semi-synthetic simulation strategy" section in Methods, "AWmeta secures robust higher-fidelity DEG identification across transcriptomic contexts" section in Results and Fig. 1d–f. Statistical significance was determined using one-tailed Mann-Whitney test. Boxplot bounds show interquartile ranges (IQR), centers indicate median values, and whiskers extend to $1.5\times$IQR. The following icons represent different tissue sources: 🧠: substantia nigra; 🩸: peripheral blood; 🟤: ileal mucosa; 🟢: colonic mucosa.

Extended Data Fig. 4 | DEG discrimination evaluation across five disease tissues using median-permuted semi-synthetic simulation strategy. a–e, DEG discrimination (AUROC and AUPRC) was assessed against AWmeta and REM using nine distinct thresholds, combining three corrected P-values (FDR) (0.01, 0.05, and 0.10) and three $\log_2$-based fold change ($|\log_2 FC|$) cutoffs ($\log_2 1.2$, $\log_2 1.5$, and $\log_2 2.0$), spanning Parkinson's substantia nigra (**a**), Parkinson's (**b**) and Crohn's (**c**) peripheral blood, and Crohn's ileal (**d**) and colonic (**e**) mucosa. Detailed description of median-permuted semi-synthetic simulation strategy can be referred to in "DEG discrimination evaluation using semi-synthetic simulation strategy" section in Methods, "AWmeta secures robust higher-fidelity DEG identification across transcriptomic contexts" section in Results and Fig. 1d–f. Statistical significance was determined using one-tailed Mann-Whitney test. Boxplot bounds show interquartile ranges (IQR), centers indicate median values, and whiskers extend to $1.5 \times$ IQR. The following icons represent different tissue sources: 🧠: substantia nigra; 🩸: peripheral blood; 🌰: ileal mucosa; 🫃: colonic mucosa.

Extended Data Fig. 5 | DEG discrimination evaluation across five disease tissues using maximum-permuted semi-synthetic simulation strategy. a–e, DEG discrimination (AUROC and AUPRC) was assessed against AWmeta and REM using nine distinct thresholds, combining three corrected *P*-values (FDR) (0.01, 0.05, and 0.10) and three log$_2$-based fold change (|log$_2$FC|) cutoffs (log$_2$1.2, log$_2$1.5, and log$_2$2.0), spanning Parkinson's substantia nigra (**a**), Parkinson's (**b**) and Crohn's (**c**) peripheral blood, and Crohn's ileal (**d**) and colonic (**e**) mucosa. Detailed description of maximum-permuted semi-synthetic simulation strategy can be referred to in "DEG discrimination evaluation using semi-synthetic simulation strategy" section in Methods, "AWmeta secures robust higher-fidelity DEG identification across transcriptomic contexts" section in Results and Fig. 1d–f. Statistical significance was determined using one-tailed Mann-Whitney test. Boxplot bounds show interquartile ranges (IQR), centers indicate median values, and whiskers extend to 1.5×IQR. The following icons represent different tissue sources: 🐵: substantia nigra; 🩸: peripheral blood; 🍩: ileal mucosa; 🫙: colonic mucosa.

Extended Data Fig. 6 | AWmeta establishes superior DEG-wise convergence in gene differential quantification. a–e, Considering that DEGs instead of non-DEGs are primarily involved in disease etiology, to explore whether gene- (unfiltered) and DEG-wise convergence assessment results are different, mean absolute deviation (MAD) -like similarity measure was utilized to quantify the per-DEG fold change ($|\log_2 FC|$) similarity among AWmeta, REM and original studies, with smaller values indicating better convergence, which demonstrates AWmeta's consistent superior DEG- with gene-wise convergence over REM and orignal studies using nine distinct thresholds, combining three corrected P-values (FDR) (0.01, 0.05, and 0.10) and three $\log_2$-based fold change ($|\log_2 FC|$) cutoffs ($\log_2 1.2$, $\log_2 1.5$, and $\log_2 2.0$), across five disease tissues: Parkinson's substantia nigra (**a**), Parkinson's (**b**) and Crohn's (**c**) peripheral blood, and Crohn's ileal (**d**) and colonic (**e**) mucosa. For comparison purpose, results from original studies serve as reference baselines. Statistical significance of REM and baselines against AWmeta for DEG-wise convergence comparisons was tested with one-tailed Mann-Whitney test. Detailed description for MAD-like DEG-wise convergence similarity measure can be referred to in "Gene-wise convergence assessment for gene differential quantification" section in Methods, "AWmeta establishes superior gene- and study-wise convergence in gene differential quantification" section in Results and Fig. 2a. Boxplot bounds show interquartile ranges (IQR), centers indicate median values, and whiskers extend to $1.5 \times$ IQR. The following icons represent different tissue sources: 👤: substantia nigra; 🩸: peripheral blood; 🐛: ileal mucosa; 🟢: colonic mucosa.

Extended Data Fig. 7 | Conceptual schematic of adjusted DE list similarity measure. The methodological details reside in "Study-wise convergence assessment for gene differential quantification" section in Methods.

Extended Data Fig. 8 | AWmeta maintains robust superior study-wise convergence in gene differential quantification across diverse thresholds. a–j, To dissect whether set-theory-based similarity-derived study-wise convergence assessment results vary with diverse DEG thresholds, we used nine distinct thresholds, combining three corrected P-values (FDR) (0.01, 0.05, and 0.10) and three $\log_2$-based fold change ($|\log_2 FC|$) cutoffs ($\log_2 1.2$, $\log_2 1.5$, and $\log_2 2.0$), to benchmark study-wise convergence, which showcases AWmeta maintains robust superior study-wise convergence in gene differential quantification across diverse thresholds over REM and baselines in five disease tissues, both by means of the average of Jaccard (JC) and overlap coefficient (OC) (**a–e**) and phi coefficient (PC) (**f–j**). For comparison purpose, results from original studies serve as reference baselines. Overall study-wise convergence differences among AWmeta, REM and baselines were tested by Kruskal–Wallis test, followed by Nemenyi post-hoc test for pairwise comparisons. Details for these two study-wise convergence similarity measures appear in "Study-wise convergence assessment for gene differential quantification" section in Methods and Fig. 2h–j. Boxplot bounds show interquartile ranges (IQR), centers indicate median values, and whiskers extend to $1.5 \times$IQR. The following icons represent different tissue sources: 👩: substantia nigra; 🩸: Peripheral blood; 🌰: ileal mucosa; 🫃: colonic mucosa.