# Disruption of Microhomology-mediated End-joining in Ewing Sarcoma

Shuhei Asada[1#, *], Guangli Zhu[1,2#], Jithma Prasad Abeykoon[1,3], Yutaro Tanaka[4,5], Huy Nguyen[1,6], Yuna Hirohashi[1], Divya R. Iyer[1], Nicholas William Ashton[1], Sirisha Mukkavalli[1], Lige Jiang[1], Miles Del Busso[3], Judith Jebastin Thangaiah[7], Steven I. Robinson[8], Kalindi Parmar[1,6], Eliezer M. Van Allen[5,9,10,11], Riaz Gillani[4,5,9,12,] Geoffrey I. Shapiro[10], Alan D. D'Andrea[1, 6*]

1. Division of Radiation and Genome Stability, Department of Radiation Oncology, Dana-Farber Cancer Institute, Harvard Medical School, Boston, MA, USA.
2. Department of Radiation Oncology, First Affiliated Hospital of Sun Yat-sen University, Guangzhou, China
3. Division of Hematology, Department of Internal Medicine, Mayo Clinic, MN, USA
4. Department of Pediatric Oncology, Dana-Farber Cancer Institute, Boston, MA, USA.
5. Cancer Program, Broad Institute of MIT and Harvard, Cambridge, MA, USA.
6. Center for DNA Damage and Repair, Dana-Farber Cancer Institute, Harvard Medical School, Boston, MA, USA.
7. Department of Pathology, Mayo Clinic, MN, USA
8. Department of Oncology, Mayo Clinic, MN, USA
9. Harvard Medical School, Boston, MA, USA.
10. Department of Medical Oncology, Dana-Farber Cancer Institute, Harvard Medical School, Boston, MA, USA.
11. Center for Cancer Genomics, Dana-Farber Cancer Institute, Boston, MA, USA.
12. Boston Children's Hospital, Boston, MA, USA.
# These authors contributed equally to this work.

*Corresponding authors: Shuhei Asada (shuhei_asada@dfci.harvard.edu) and Alan D. D'Andrea (alan_dandrea@dfci.harvard.edu)

Keywords: Ewing's Sarcoma, DNA double-strand break, Alternative end-joining, TMEJ, DNA-PKc, CDK12

**SUMMARY**

Ewing sarcoma (EwS) is a group of bone and soft tissue cancers in children and young adults. The

EWS-FLI1 protein, the oncogenic driver in EwS, is a fusion of the FET protein, EWSR1, at its

amino terminus, with the carboxy terminus of the transcription factor, FLI1. Since EwS cells have

pronounced sensitivity to radiation and chemotherapy-induced DNA damage, the role of EWS-

FLI1 in DNA repair is likely. Here, we demonstrate that EWS-FLI1 causes a defect in

microhomology-mediated end-joining (MMEJ) repair. EWSR1 is a splicing factor that promotes

the faithful splicing of the *POLQ* pre-mRNA, required for the expression of POLΘ, a critical

protein in the MMEJ pathway. Expression of EWS-FLI1, or loss of EWSR1, causes exon 25

skipping of the *POLQ* transcript, decreased POLΘ protein expression, impaired MMEJ, and

cellular sensitivity to inhibitors of NHEJ repair and HR repair, through the mechanism of synthetic

lethality. Knockdown of EWS-FLI1 expression restores POLΘ mitotic foci and increases MMEJ

activity. Based on this mechanism, inhibitors of NHEJ or HR may provide a targeted therapy for

patients with Ewing sarcoma.

**Highlights:**
52
53 -Ewing sarcoma tumors have a deficiency in POLθ expression and a corresponding loss of MMEJ
54 activity
55
56 -EWSR1 is a splicing factor that interacts with other splicing factors such as FUBP1 and
57 KHSRP/FUBP2 to accurately splice the *POLQ* mRNA.
58
59 -The EWS-FLI1 fusion oncoprotein, or loss of EWSR1, causes a splicing defect, leading to exon25
60 skipping of the *POLQ* pre-mRNA and loss of POLθ expression
61
62 -The MMEJ deficiency of EwS cells results in cellular sensitivity to inhibitors of Homologous
63 Recombination or Non-Homologous End-Joining
64
65 -Exon 25 skipping of *POLQ* mRNA is a predictive biomarker for HR inhibitors in human cancers
66

## INTRODUCTION

Ewing sarcoma (EwS) is a group of bone and soft tissue cancers in children and young adults[1,2]. EwS tumor cells have a chromosome 11;22 translocation, resulting in expression of the EWS-FLI1 fusion oncoprotein. Their elevated sensitivity to ionizing radiation and DNA-damaging chemotherapy, as well as their relatively low level of somatic mutations, suggest that EwS cells have an underlying defect in an error-prone DNA repair (DDR) pathway. Despite this high drug sensitivity, novel treatments for EwS are required, especially for patients with relapsed and/or refractory disease, who have a low 5-year overall survival rate of only 10%.

Recent studies have identified some abnormalities in DDR in Ewing sarcoma cells. Expression of EWS-FLI1 fusion protein promotes R-loop formation and impaired BRCA1 recruitment to chromatin, suggesting a defect in the homologous-recombination repair pathway (HRD) [3]. Despite this preclinical observation, PARP inhibition, an approved treatment for HR-deficient tumors, was ineffective in the treatment of EwS patients[4]. Other cellular defects resulting from EWS-FLI1 expression may indirectly affect DDR, such as defects in BAF complex-mediated transcription [5,6], condensate generation [7-9], and mRNA splicing [10]. However, the mechanism by which EWS-FLI1 specifically influences DDR pathways is unknown, and other more profound DNA repair defects, beyond the HRD phenotype, may exist.

DNA double-strand breaks (DSBs) are the most critical forms of DNA damage in cells. There are three main pathways for repairing DSBs - non-homologous end-joining (NHEJ), homologous recombination (HR), and microhomology-mediated end-joining (MMEJ). For a cancer cell to survive and proliferate, DSBs must be repaired, and the cell must rely on at least one of these pathways. DNA polymerase theta (POLΘ), an error-prone polymerase encoded by the *POLQ* gene, is required for the MMEJ pathway. Overexpression of *POLQ* confers

chemoresistance, whereas depletion of POLΘ results in cancer predisposition and

hypersensitivity to irradiation and chemotherapy [11,12]. *To date, there are no reported human*

*cancers with defects in the MMEJ pathway.*

Here, we demonstrate that EwS cells have an underlying defect in MMEJ. Expression of

EWS-FLI1, or loss of EWSR1, in these cells promotes exon skipping of the *POLQ* pre-mRNA,

generating a premature termination codon and a profound reduction of POLΘ protein expression.

EWS-FLI1 expressing cells exhibit a phenotype like *POLQ* deficient cells, including

hypersensitivity to topoisomerase inhibitors, irradiation, and synthetic lethality with loss of the

NHEJ or HR pathways.  Knockout of the EWS-FLI1 oncoprotein in EwS cells results in reduced

exon skipping, re-expression of the POLΘ protein, and a rescue of MMEJ activity. Accordingly,

the DDR defect in EwS tumors could be exploited for therapeutic benefit by inhibiting another

DSB repair pathway.

104
105

**RESULTS**

***EWSR1* knock-out cells phenocopy MMEJ-deficient cells.**

Previous CRISPR screens and Depmap analyses have demonstrated that genes in the

MMEJ pathway are synthetic lethal in cells with an underlying defect in an HR gene [13,14],

Interestingly, in these screens, knockout of the *EWSR1* gene was highly similar to knockout of

known MMEJ genes,  including *POLQ* [15], *XRCC1* [16], *APEX2* [17], and *RHNO1* [18], in the killing of

HR-deficient tumor cells (**Extended Data Fig. 1A**).  To investigate the function of EWSR1 in

DNA repair, we initially generated an *EWSR1* knock-out in RPE *TP53-/-* cells (**Fig. 1A**).  *EWSR1*

knockout resulted in hypersensitivity to ionizing irradiation, topoisomerase inhibitors, and ATR

inhibitors, but not to the PARP inhibitor, suggesting a loss of MMEJ activity (**Extended Data Fig.**

**1 B-F**). The DDR defect observed in *EWSR1* knockout cells was similar to the defect observed in

*POLQ* knockout cells [11]. Consistently, *EWSR1* is mapped closely to *POLQ* in the previous repair-

seq, indicating a functional relationship between these genes in DNA damage repair [19].

Next, the impact of *EWSR1* loss on the activity of DNA double-strand break repair

pathways in U2OS cells was evaluated, using established Isce-I-based reporter systems [20,21].

*EWSR1* knockout mildly decreased HR activity (**Extended Data Fig. 1G**); however, the *EWSR1-*

deficient cells were not sensitive to the PARP inhibitor, Olaparib.  Interestingly, *EWSR1* knockout

significantly decreased MMEJ activity (**Fig. 1B**). To further confirm a MMEJ reduction upon

*EWSR1* depletion, a novel Cas9-based MMEJ reporter system was generated (**Extended Data Fig.**

**1H**). Again, *EWSR1* knockout decreased MMEJ activity in this Cas9-based reporter cells (**Fig.**

**1C**). Consistent with this result, *EWSR1*-deficient cells were resistant to the *POLQ* inhibitor,

Novobiocin [22](**Extended Data Fig. 1I**).

Recent studies suggest that *EWSR1* knockdown is synthetic lethal with BRCA2 loss [13,14].

To validate these results, depletion of EWSR1 in BRCA2-proficient and BRCA2-deficient DLD1

cells was evaluated.  Like *POLQ* loss, EWSR1 loss profoundly decreased colony-forming activity

in the *BRCA2*-deficient DLD1 cells (**Fig. 1D**).  EWSR1 loss also efficiently decreased clonogenic

activity of the *BRCA1*-mutated breast cancer cell lines, MDA-MB-468 and HCC1937 cells

(**Extended Data Fig1. J, K**).  To further confirm the synthetic lethality of EWSR1 and BRCA1/2,

we generated RPE *TP53-/-* cells stably expressing Cas9 (**Extended Data Fig. 1L**).  These cells

were subsequently lentivirally-transduced with a sgRNA targeting control, *POLQ,* or *EWSR1* (co-

expressing GFP) together with another sgRNA targeting control, *BRCA1* or *BRCA2* (co-expressing

tRFP657), and changes in the frequency of GFP/tRFP657 cells were monitored.  Again, like *POLQ*

loss, *EWSR1* loss resulted in synthetic lethality with *BRCA1/2* loss in this system. Taken together,

*EWSR1* functions as a MMEJ gene.

**EWSR1 promotes MMEJ by maintaining POLΘ expression.**

Since EWSR1 loss caused a reduction of MMEJ activity, we examined the impact of

EWSR1 loss on protein expression of known MMEJ-related proteins. To monitor the POLΘ

protein expression, RPE *TP53-/-* cells expressing 3xFLAG-tagged *POLQ* were generated.

Interestingly, *EWSR1* knockout resulted in decreased POLΘ expression (**Fig. 1E, Extended Data**

**Fig. 2A**). Increased radiosensitivity of the *EWSR1* knockout cells was observed (**Extended Data**

**Fig. 2B**).  To further evaluate the direct and acute effect of EWSR1 loss on POLΘ expression, we

tagged endogenous EWSR1 protein with the mutant cytosolic prolyl isomerase FKBP12$^{F36V}$ in

endogenously 3xFLAG-tagged *POLQ* expressing HEK293T cells [23,24].  In this targeted protein

degradation system, the small bifunctional molecule dTAG$^{v1}$ bridges FKBP12$^{F36V}$ with the E3

ubiquitin ligase VHL, resulting in substrate ubiquitination and proteasomal degradation. Treatment

of cells harboring the FKBP12$^{F36V}$-tagged EWSR1 with dTAG$^{v1}$ resulted in rapid degradation of

EWSR1 protein (**Extended Data Fig. 2C-F**). Interestingly, this acute depletion of EWSR1 rapidly

reduced the expression of POLΘ but did not affect the expression of other known MMEJ-related

proteins, including APEX2, HMCES, FEN1, LIG3, PARP1, and XRCC1 nor HR-related proteins

such as BRCA1 and BRCA2 (**Fig 1F, Extended Data Fig. 2G**).  Since EWSR1, as well as FUS

and TAF15, belongs to the FET family of proteins [25], we speculated that POLΘ loss and MMEJ

deficiency might be a common phenotype of the loss of FET family genes. However, FUS or

TAF15 loss did not affect the protein level of POLΘ or MMEJ activity (**Extended Data 2H, I**).

Consistently, *EWSR1* knockout reduced the POLΘ foci assembly upon irradiation-induced DNA

damage (**Fig. 1G**).

The functional significance of the loss of POLΘ on MMEJ activity in *EWSR1*-depleted

cells was next examined.  Transduction with the WT *POLQ* cDNA rescued MMEJ activity in

*EWSR1* knockout cells (**Fig 1H**). Moreover, *POLQ* cDNA transduction corrected the

hypersensitivity to the topoisomerase inhibitors, indicating that loss of POLΘ protein is

responsible for these phenotypes in *EWSR1* knockout cells (**Fig 1I, Extended Data Fig 2J**).  Like

POLΘ loss, EWSR1 loss increased unrepaired DNA damage and micronuclei in BRCA-deficient

cells (**Extended Data Fig 2K, L**).  Collectively, EWSR1 promotes MMEJ via the maintenance of

POLΘ protein expression.

170
171  **EWSR1 directly binds to the *POLQ* transcript and prevents *POLQ* exon skipping.**
172
To investigate the cause of the reduced POLΘ protein expression following loss of

EWSR1, we evaluated the stability of POLΘ protein in the presence or absence of EWSR1. U2OS

MMEJ-reporter cells were transduced with the *POLQ* cDNA, followed by *EWSR1* knockout by

the CRISPR-Cas9 system.  However, the protein level of POLΘ expressed by *POLQ* cDNA did

not change after *EWSR1* depletion (**Extended Data Fig. 3A**). Furthermore, a proteasome inhibitor

MG132, or the autophagy inhibitor chloroquine, did not rescue endogenous POLΘ expression in

*EWSR1*-depleted cell (**Extended Data Fig. 3B**), suggesting a pre-translational mechanism.

To assess the overall influence of EWSR1 loss on transcript levels, including the POLQ

transcripts, we next performed an RNA-seq analysis on RPE1 *TP53-/-* cells and RPE1 *TP53-/-*

*EWSR1* knockout cells. Alternative splicing events were analyzed by the MISO framework [26] as

well as the rMATs framework[27].  Based on this analysis, various alternative splicing events,

including exon skipping events, were observed in *EWSR1* knock-out cells (**Extended Data Fig.**

**3C-G)**. Among DNA repair genes with exon skipping events, we identified several exon skipping

events of the *POLQ* transcript, including exon 7, exon 21 and exon 25, as top hits (**Fig. 2A**).

Among these exon skipping events within the *POLQ* transcript, the skipping of exon 25 was the

most prominent (**Fig. 2B**). Of note, *POLQ* exon 25 is located between long introns, intron 24 and

intron 25. Furthermore, we re-analyzed previously published RNA-seq datasets using si-control

and si-EWSR1 cells, and again the *POLQ* exon 25 skipping is one of the most significant events

(**Extended Data Fig. 3H, I**). We validated the results in multiple *EWSR1* knockout clones by RT-

PCR (**Extended Data Fig. 3J**).

To evaluate the impact of exon-skipping events on POLΘ protein expression, we

transduced sgRNAs targeting individual exons. Targeting *POLQ* exon 7, exon 21, or exon 25 by

CRISPR-Cas9-mediated gene knockout induced profound protein loss of POLΘ as well as a

reduction of MMEJ activity (**Fig.2C, D**). Moreover, a cycloheximide chase assay revealed that a

shorter isoform of the POLΘ protein, expressed from a *POLQ* cDNA lacking exon 25, is more

unstable than the full-length POLΘ protein (**Fig. 2E**). Exon 25 skipping causes premature

termination of POLΘ protein synthesis, preventing the complete expression of the "Fingers" and

"Palm" of the encoded polymerase. Structural modeling by AlphaFold 3 demonstrated that this

truncation disrupts the structural integrity of this domain, likely explaining the instability

compared to the full-length protein (**Extended Data Fig. 3K**). The *POLQ* cDNA lacking exon 25

or exon7 failed to rescue the MMEJ-deficiencies in *POLQ* knockout cells (**Fig. 2F**).

To further investigate the biological significance of *POLQ* exon 25 skipping in the MMEJ

activity, we designed antisense oligonucleotides (ASOs) that target exon 25 and induce exon 25

skipping (**Extended Data Fig. 4A**). *POLQ* exon 25 skipping decreased POLΘ protein expression

and MMEJ activity (**Extended Data Fig. 4B, C**). Moreover, a mNeongreen-based minigene

reporter, as previously described [28], was generated in which the mNeongreen is split into two exons

with synthetic genes derived from the partial intron 24/25 and exon 25 of *POLQ* inserted in

between (**Extended Data Fig. 4D**). The minigene reporter recapitulated the *POLQ* exon 25

skipping driven by depletion of EWSR1, indicating a direct involvement of EWSR1 in the

reduction of *POLQ* exon 25 skipping by binding the *POLQ* pre-mRNA near the exon 25 (**Fig. 2G,**

**Extended Data Fig. 4E**). Using an RNA immunoprecipitation protocol [29], we determined whether

EWSR1 interacts with the *POLQ* pre-mRNA, using HEK293T cells harboring the FKBP12$^{F36V}$-

tagged EWSR1 (**Extended Data Fig. 4F**). Indeed, EWSR1 efficiently co-immunoprecipitated

with the *POLQ* pre-mRNA near exon 7 and exon 25, as compared to the *GAPDH* control

(**Extended Data Fig. 4G, H**). Taken together, EWSR1 binds directly to *POLQ* pre-mRNA and

prevents exon skipping of the *POLQ* transcript, leading to maintenance of POLΘ protein levels

and MMEJ activity.

Given that one allele of *EWSR1* is lost in EwS cells, we speculated that these phenomena

would also be observed in EwS cells, including patient samples.  A higher level of exon 7 and

exon 25 skipping and decreased expression of POLΘ protein were evident in EwS cell lines,

compared to the osteosarcoma cell line U2OS (**Fig. 2H, I**). Importantly, patient-derived EwS

tumor samples exhibited a higher frequency of *POLQ* exon 25 skipping than primary tumors from

other pediatric solid tumors, including Neuroblastoma, Osteosarcoma, and Wilms tumor (**Fig. 2J**).

Moreover, *POLQ* mRNA levels in Ewing sarcoma were comparable to or lower than other

pediatric solid tumors (**Fig. 2K**).

We also evaluated single nucleotide variants (SNVs), the mutational signature, and small

deletions in patient-derived EwS tumors. Consistent with previous studies [30], the EwS tumors have

a low SNV frequency compared to Neuroblastoma, consistent with a MMEJ deficiency in the

tumors (**Fig. 2L**).  Although the level of HRD hallmarks, ID-6 and signature 3, were overall low,

which is consistent with a previous study [31], Ewing sarcoma tumors showed an even lower level

of signature 3 or ID-6, compared to Neuroblastomas, again confirming that EwS is an HR-

proficient tumor **(Extended Data Fig. 5A-F)**.  In contrast, the indels observed in the EwS samples,

compared with neuroblastoma samples, have a lower frequency of deletions with >3bp micro-

homologies and templated insertions, consistent with a low level of *POLQ*-mediated MMEJ in

EwS [32] **(Fig. 2M, Extended Data Fig. 5G-I)**.  To validate our findings in the primary patient

samples, we measured the degree of microhomology at breakpoints of the *EWSR1* fusion in EwS

tumor samples from two different EwS patient cohorts. Interestingly, the degree of microhomology

at canonical translocation breakpoints was very low in the EwS genomes, consistent with a cellular

defect in the MMEJ pathway and a compensatory increase in NHEJ (**Extended Data Fig. 5J, K**).

Collectively, EwS tumor cells have a bona fide defect in MMEJ, which is probably due to increased

*POLQ* exon 25 skipping.

**The Prion-like domain of EWSR1 is required for the faithful splicing of the *POLQ***

**transcript.**

To identify the functional domain of EWSR1 required for accurate *POLQ* splicing, we next

generated EWSR1 mutants lacking individual domains (**Fig. 3A**). Deletion of the prion-like

domain (M1) or the C-terminus RRG2 domain (M6) resulted in failure to rescue the *POLQ* exon

skipping and POLΘ protein expression level (**Fig. 3B, C, Extended Data Fig. 6A**). Consistent

with these results, MMEJ activity was not restored in *EWSR1*-depleted cells by these two mutant

forms of EWSR1 (**Extended Data Fig. 6B**). In addition, these mutant proteins were unable to

rescue the cellular hypersensitivity to topoisomerase inhibitors or an ATR inhibitor. The mutants

also failed to rescue the synthetic lethality resulting from BRCA2 loss in *EWSR1*-deficient cells

(**Extended Data Fig. 6C-G**).

A nuclear localization signal (NLS) is located at the RRG2 domain, and its loss is likely to

destabilize the EWSR1 protein. The NLS mutant protein failed to rescue *POLQ* exon skipping and

failed to restore POLΘ protein expression. To further evaluate the prion-like domain, the domain

was subdivided into five parts, and mutant proteins lacking the individual regions were generated

(**Extended Data Fig. 6H**). Interestingly, all mutant forms were able to rescue the POLΘ protein

level, albeit with some minor differences, suggesting that the domains have functional

redundancies (**Extended Data Fig. 6I**). Since the prion-like domain of EWSR1 is critical for the

formation of condensates of EWSR1 [33], we infer that the formation of EWSR1 condensates may

play a role in maintaining the faithful splicing of POLΘ. To test this hypothesis, we generated

multiple mutant forms of EWSR1 in which tyrosine residues of the prion-like domain were

substituted with serine residues, as previously described[33]. Strikingly, an EWSR1-7YS mutant, in

which tyrosine residues were substituted by serine residues, failed to rescue the *POLQ* exon

skipping and POLΘ protein expression level in *EWSR1*-depleted cells (**Fig. 3D, E, Extended Data**

**Fig. 6J, K**).  Furthermore, an EWSR1-37YS mutant, in which all thirty-seven tyrosine residues in

the prion-like domain were replaced with serine residues, and known to be defective in condensate

formation, abolished the ability of EWSR1 to prevent exon skipping of the *POLQ* transcript,

further suggesting that condensate formation is critical for faithful splicing.  Consistent with these

findings, the EWSR1-7YS and EWSR1-37YS mutants failed to rescue MMEJ activity in EWSR1-

depleted cells (**Fig. 3F**).  Taken together, the prion-like domain of EWSR1 is required for the

faithful splicing of the *POLQ* transcript and for the expression of the POLΘ protein, suggesting

that condensate formation of EWSR1 is also required for these activities as previously described

[5,33].

279
**The prion-like domain of EWSR1 is critical for interaction with spliceosomes including**

**FUBP1 and KHSRP.**

The condensate-defective EWSR1 mutant may have lost the ability to bind the *POLQ*-pre

mRNA. However, RNA-immunoprecipitation revealed that the EWSR1-7YS and EWSR1-37YS

mutants can still bind to the *POLQ* pre-mRNA (**Extended Data Fig 4. F-I**).   Alternatively, the

condensate-defective EWSR1 mutant may be competent for *POLQ* mRNA binding but defective

in its interaction with RNA-binding proteins in spliceosomes. To test this hypothesis, we

performed an IP/mass spectrometry analysis using HEK293T cells with a biallelic FKBP12-

V36F knock-in at the N-terminus of *EWSR1* (**Fig. 3G**).  The cDNAs encoding the control V5-

tagged *EWSR1*-WT, or the *EWSR1* 37YS mutant, were transduced into the cells and, following

dTAG[v1] treatment, cellular proteins were immunoprecipitated with the anti-V5 antibody. A

previous study demonstrated that the prion-like domain of EWS-FLI1 interacts with BAF

complex subunit components in condensates [5]. Accordingly, wildtype EWSR1, but not the

EWSR1 37YS mutant protein, preferentially bound to BAF subunits, including ARID1A,

SMARCA2, SMARCA4, and SMARCC1 (**Fig. 3G, I, Extended Data Fig. 7A**). In addition to

these BAF subunits, additional RNA binding proteins were identified as binding partners of

wild-type EWSR1. Intriguingly, the interactors contain prion-like domains, further indicating a

functional interaction of EWSR1 with these proteins in spliceosomes (**Fig. 3H**).

Since exon skipping in *EWSR1* KO cells is typically observed for exons that are flanked

by longer introns (**Extended Data Fig. 7B**), we focused on FUBP1, among the prion-like

domain-dependent EWSR1 binding partners. FUBP1 is a spliceosome known to mediate the

splicing of long introns [34] (**Fig. 3H, I**). In addition, KHSRP/FUBP2, known to have a similar

function in splicing, was further evaluated. Depletion of FUBP1 or KHSRP alone cause only a

minor reduction in POLΘ protein expression or MMEJ activity. However, dual depletion of

FUBP1 and KHSRP increased *POLQ* exon25 skipping, reduced the POLΘ protein level, and

reduced the cellular MMEJ activity, indicating a functional redundancy between FUBP1 and

KHSRP (**Fig. 3 J, K, Extended Data Fig. 7C**). A FUBP1-16YS mutant, with tyrosine residues

of its C-terminus region substituted with serine residues, failed to bind to wildtype EWSR1 and

failed to rescue the phenotype of FUBP1/KHSRP DKO cells, suggesting that tyrosine residues of

EWSR1 and FUBP1 are critical for the protein-protein interaction and for *POLQ* exon25

inclusion (**Fig. 3 L-Q, Extended Data Fig. 7D**). Similarly, a KHSRP-17YS mutant failed to

rescue the phenotype of FUBP1/KHSRP DKO cells (**Extended Data Fig. 7E-H**). Taken

together, the protein-protein interaction between EWSR1 and FUBP1 or KHSRP via the prion-

like domain is critical for the faithful splicing of the *POLQ* pre-mRNA.

**The EWS-FLI1 oncoprotein inhibits formation of POLQ mitotic foci at DSB loci and**

**disrupts MMEJ.**

The chromosomal translocation of 11;22 leads not only to the loss of one wildtype EWSR1

allele but also to the generation of an oncogenic fusion protein EWS-FLI1 in EwS cells (**Extended**

**Data Fig. 8A**).  Moreover, EWS-FLI1 is known to heterodimerize with EWSR1 and to disrupt its

activity [5,35,36]. We therefore hypothesized that this oncoprotein may directly or indirectly interfere

with MMEJ activity. To test this hypothesis, the EWS-FLI1 oncoprotein was lentivirally expressed

in U2OS-based MMEJ reporter cells and evaluated for MMEJ activity. Interestingly, EWS-FLI1

diminished the MMEJ activity in U2OS MMEJ reporter cells (**Fig.4A, Extended Data Fig. 8B**).

Consistent with these results, EwS cell lines exhibited resistance to POLΘ inhibitors, Novobiocin

and ART558 [37], compared with osteosarcoma cells U2OS (**Extended Data Fig. 8C, D**).  The

influence of various mutant forms of the EWS-FLI1 fusion oncoprotein on MMEJ activity was

tested (**Fig. 4B**). Expression of the SYGQ1-FLI1, SYGQ2-FLI1, ΔSYGQ1, or ΔSYGQ2 mutant

by cDNA-mediated transfection decreased the MMEJ activity in the U2OS reporter cells,

indicating functional redundancies in the prion-like domain of EWS-FLI1.  Interestingly, the

EWS-FLI1 37YS mutant, which failed to disrupt MMEJ activity (**Fig. 4B, Extended Data Fig.**

**8B**), is defective in condensate formation and in binding to wild-type EWSR1 protein [5,36].

Moreover, expression of EWS-FLI1, but not the EWS-FLI1 37YS mutant, conferred

hypersensitivity to topoisomerase inhibitors (**Extended Data Fig. 8E, F**).

To determine the effect of the EWS-FLI1 oncoprotein on MMEJ activity in Ewing sarcoma

cells, we transduced an Isce-I-based MMEJ reporter into the A673, RH1, and RD-ES EwS lines.

As expected, depletion of EWS-FLI1 from these cells by sgFLI1, as confirmed by western blot,

restored MMEJ activity (**Fig. 4C, Extended Data Fig. 8G**).  These EwS lines, with EWS-FLI1

depletion, exhibited resistance to the POLΘ inhibitor, ART558 (**Extended Data Fig. 8H),**

consistent with their restoration of MMEJ following EWS-FLI1 depletion. The minigene reporter

assay showed that expression of EWS-FLI1 but not the EWS-FLI1 37YS mutant increased the

*POLQ* exon 25 skipping, whereas depletion of EWS-FLI1 in EwS cells ameliorated the *POLQ*

exon 25 skipping (**Fig. 4D, E**). Consistent with a previous study [5], EWS-FLI1 preferentially bound

to EWSR1 compared with the EWS-FLI1 37YS mutant, suggesting an inhibitory effect of EWS-

FLI1 on EWSR1-mediated splicing (**Fig. 4F**). Given that R-loop accumulation driven by the EWS-

FLI1 expression inhibits the recruitment of BRCA1 to the DSB loci, we asked whether R-loop

accumulation may be involved in the MMEJ reduction in Ewing sarcoma. Expression of wildtype

RNaseH1 or mutant RNaseH1 did not affect EWS-FLI1-mediated MMEJ reduction (**Extended**

**Data Fig. 8I, J**).

Recent studies indicate that DSB repair by POLΘ and MMEJ occurs during mitosis of the

cell cycle [38].  POLΘ is phosphorylated by PLK1 and assembles with TOPBP1, along with Rhino

and the 9-1-1 complex, into POLΘ foci [18,39]. To assess the influence of EWS-FLI1 expression on

the assembly of DSB loci, we engineered endogenously V5-tagged POLΘ expressing MMEJ

reporter U2OS cells by CRISPR/Cas9-mediated knock-in. After generation of adenoviral I-SceI-

induced DSBs, we performed chromatin-immunoprecipitation (ChIP) with an anti-V5 antibody,

followed by qPCR analysis.  Overexpression of EWS-FLI1 in these cells disrupted the assembly

of POLΘ with sites of DNA double-strand breaks **(Fig. 4G).**  Consistent with this result, MMEJ

activity and the assembly of POLΘ foci assembly, as measured by immunofluorescence, was

decreased by the overexpression of EWS-FLI1 (**Fig. 4H**).  Moreover, EWS-FLI1 overexpression

in these U2OS cells also increased γH2AX foci, consistent with the loss of MMEJ activity in

mitosis. EWS-FLI1 overexpression can therefore reduce POLΘ-mediated DSB repair in mitosis,

at a time when POLΘ-mediated MMEJ is highly active [18,39].   EWS-FLI1 was also depleted in

endogenously V5-tagged POLΘ expressing A673 cells. As predicted, the loss of EWS-FLI1

restored POLΘ foci assembly (**Fig. 4I).** Collectively, the EWS-FLI1 oncoprotein can reduce the

level of POLΘ mitotic foci, leading to a functional impairment of MMEJ in EwS cells.

366
367
**Inhibition of NHEJ or HR efficiently kills the MMEJ-deficient Ewing sarcoma cells via**

**synthetic lethality.**

Since inhibiting two of three DSB repair pathways is synthetically lethal in cells [40,41], we

asked whether inhibition of NHEJ or HR would provide a viable therapeutic modality in MMEJ-

deficient Ewing sarcoma cells.  Previous studies have shown that inhibitors of NHEJ are

synthetic lethal with MMEJ loss [40,41]. Consistent with our study, EwS cells are sensitive to DNA-

PK inhibitors, known to inhibit NHEJ and enhance the activity of cytotoxic agents in these cells

[42]. We confirmed that EwS cell lines are sensitive to the DNA-PKci (**Fig. 5A**) and have an

elevated baseline level of activated pDNA-PKc (**Fig. 5B**).  EwS cells exhibited high phospho-

RPA, a replication stress marker, and were hypersensitive to the topoisomerase inhibitors,

topotecan, and etoposide (**Extended Data Fig. 8A, 9A, B**). Since etoposide is used in dose-

intensive chemotherapy of Ewing sarcoma, we asked whether inhibition of NHEJ could increase

the therapeutic efficacy of etoposide in EWS cells. Strikingly, the DNA-PK inhibitor, Peposertib,

enhanced the activity of etoposide in two EWS cell lines (**Fig. 5C, D**).  Moreover, the EwS cell

line A673, grown as a xenograft, was also sensitive to peposertib, *in vivo*, and the drug enhanced

the activity of etoposide (**Fig. 5E, F, Extended Data Fig. 9C, D**). Consistent with the enhanced

sensitivity to peposertib, immunohistochemistry analysis of the EwS primary patient samples

revealed an elevated baseline level of pDNA-PKc compared with Osteosarcoma patient samples

(**Fig. 5G**).

On the other hand, EWS-FLI1- but not EWS-FLI1-YS mutant-expressing cells are more

dependent on BRCA2, confirming that the HR pathway is also a rational target in MMEJ-

deficient EwS cells (**Fig. 5H**). Previous studies have also suggested that EwS cells are sensitive

to HR inhibitors, perhaps through a mechanism of synthetic lethality. CDK12 is a known

activator of HR repair, based on its ability to promote the transcription of long HR genes [43]. A

targeted inhibitor of CDK12 is known to enhance the killing of EwS cells [44]. Consistent with this

report, EwS cell lines are sensitive to the CDK12 inhibitors THZ531 and MFH290 (**Fig 5 I, J**).

Depletion of the EWS-FLI1 oncoprotein in the EwS cells resulted in a restoration of MMEJ

activity and acquisition of CDK12i resistance (**Fig 5K, L, Extended Data Fig. 9E, F**). Taken

together, the killing of the EwS cells by this HR inhibitor results from synthetic lethality between

the HR pathway and the MMEJ pathway.

**DISCUSSION**

Despite the high sensitivity to DNA-damaging agents, EwS cells, expressing the EWS-

FLI1 fusion oncoprotein, have a remarkably low somatic mutation burden [30,45,46]. This

observation suggested that EwS cells are deficient in an error-prone DNA repair mechanism,

such as MMEJ. Indeed, CRISPR screens confirmed that the knockout of the EWSR1 gene

functionally resembles the knockout of known MMEJ genes, such as *POLQ*, *XRCC1*, *APEX2*,

and *RHNO1*, suggesting that EWSR1 protein cooperates with these other encoded proteins in the

MMEJ pathway. Moreover, EWSR1-deficient cancer cell lines and EwS tumors have a low ID6

mutational signature, indicating a low level of microhomology-mediated repair of double-strand

breaks. Since EWSR1 has a known function as a mRNA splicing factor [35,47-49], we hypothesized

that EWSR1 may regulate MMEJ indirectly through its splicing of the mRNA encoding a MMEJ

protein. EWSR1 is indeed required for the accurate splicing of the long mRNA encoding the

critical MMEJ polymerase, POLΘ, although it is not required for splicing of mRNAs encoding

other MMEJ proteins. Disruption of EWSR1 activity in EwS cells results in skipping of multiple

exons in the *POLQ* pre-mRNA, especially exon 25, leading to decreased POLΘ expression in

EwS cells and to a strong MMEJ defect.

In contrast to previous reports, our study indicates that EwS tumors are competent for

high-fidelity DNA repair pathways, such as the homologous recombination pathway, and the

tumors are PARP inhibitor-resistant. Other recent studies indicate that EwS cells are dependent

on the HR genes, BRCA1, BARD1, and PALB2 [50,51] for survival. Previous studies have also

demonstrated that HR deficiency is synthetic lethal in cells with a MMEJ deficiency [22,52,53].

Accordingly, the strong MMEJ defect in EwS cells could result in an increased reliance on HR

repair.

The chromosome 11/22 translocation in EwS cells and the generation of the EWS-FLI1

fusion oncoprotein appears to account, at least in part, for the cellular defect in EWSR1 splicing

activity.  First, since one allele of EWSR1 is disrupted by the chromosome translocation, the

EwS cells already have a 50% reduction in EWSR1 protein expression.  Second, the EWS-FLI1

oncoprotein heterodimerizes with EWSR1 and exerts a dominant negative effect on EWSR1

function [5].  Our studies indicate that loss of EWSR1 function can account for the decreased

expression of POLΘ protein, the elevated exon 25 skipping of the *POLQ* pre-mRNA, and the

defective MMEJ activity in EwS cells.  Enhanced expression of the EWSR1 protein, or acute

knockdown of the EWS-FLI1 oncoprotein, can reverse these defects and restore MMEJ activity,

thereby reducing the cellular reliance on other DSB repair pathways, such as HR and NHEJ.

Splicing of specific mRNAs is regulated, at least in part, by RNA Binding Proteins

(RBPs) which function in membrane-less condensates.   Condensates known as nuclear speckles

are required for splicing activity[54].  Many RNPs, including the EWSR1 protein, contain a prion-

like domain (PrLD), allowing the proteins to assemble with other RBPs contained in

condensates. We reasoned that EWSR1 might interact with other RBPs and function

cooperatively as a spliceosome for the *POLQ* pre-mRNA.  Moreover, disruption of the PrLD of

EWSR1, through the generation of multiple Tyrosine to Serine (YS) mutations, disrupts the

assembly of functional spliceosomes. Indeed, the EWSR 37YS mutant, known to be defective in

condensate assembly [5], also fails to correct exon 25 skipping of the *POLQ* mRNA, and fails to

restore POLΘ protein expression.  IP/Mass Spectrometry identified specific RBPs that interact

with EWSR1. Interestingly, multiple RBPs containing prion-like domains were identified in the

immune complexes, and these proteins did not coimmunoprecipitate with the EWSR1 37YS

mutant protein.  Components of SWI/SNF complexes, including subunits of the cBAF complex

with prion-like domains, were also found to interact with EWSR1 and EWS-FLI1, consistent

with a previous report [5]. Unlike EWSR1, other FET proteins containing PrLD domains, such as

FUS and TAF15, were not required for accurate *POLQ* splicing, suggesting that EWSR1 forms

distinct RBP complexes, perhaps functioning in distinct condensates.  Taken together, EWSR1

may act as a "hub" for spliceosomes, using its specific prion-like domain for binding to a distinct

set of RBPs and for splicing of the *POLQ* mRNA and probably other mRNAs.  Expression of

EWS-FLI1 disrupts this splicing mechanism.

Some RNA binding proteins, including FUBP1 and KHSRP/FUBP2, specifically interact

with wildtype EWSR1, but not with the EWSR1 37YS mutant.  FUBP1 and KHSRP have high

similarity in their protein sequence and exhibit functional redundancy in their splicing activity [55].

FUBP1 and KHSRP are multifunctional RNA binding proteins and are known to redundantly

regulate the splicing of the SMN2 mRNA [55].  A carboxy-terminal, tyrosine-rich region is present

in both FUBP1 and KHSRP/FUBP2 [56].  However, the biological functions of the tyrosine-rich

regions of FUBP1 and KHSRP have remained unclear.  Since the tyrosine residues of the prion-

like domain are critical for liquid-liquid phase separation[57,58] and condensate formation,  a YS

mutant of FUBP1 or KHSRP, may fail to form condensates with EWSR1 and may lead to

aberrant splicing events, such as *POLQ* exon skipping.   FUBP1 is crucial for splicing long

introns [34] such as the long intron 24/25 in the *POLQ* pre-mRNA.  Therefore, a condensate

formed by EWSR1-FUBP1/KHSRP may be required for the accurate splicing of exons trapped

between long introns.  Interestingly, the dual knockout of FUBP1 and KHSRP, like the knockout

of EWSR1, results in exon 25 skipping of the *POLQ* mRNA and a reduction in POLΘ protein

expression and MMEJ.  Re-expression of the wildtype FUBP1, but not of the FUBP1-YS

mutant, rescued these defects in the dual knockout cells.  The RBPs, EWSR1, FUBP1, and

KHSRP therefore appear to functionally interact via their PrLD domains and to cooperate in a

EWSR1-initiated splicing pathway for the *POLQ* mRNA (**Extended Data Fig. 9G**).

Given that overexpression of EWS-FLI1 oncoprotein induces extensive DNA damage[59],

we reasoned that the EWS-FLI1 fusion oncoprotein could impair MMEJ activity in EwS cells.

Indeed, the acute degradation of the EWS-FLI1 oncoprotein rapidly rescued MMEJ, suggesting

an acute upregulation of POLΘ expression and POLΘ mitotic foci.  While the EWS-FLI1

protein forms condensate with EWSR1, it disrupts the function of EWSR1, reduces POLΘ

expression, and inhibits MMEJ.  The EWS-FLI1-37YS mutant fails to form condensates with

wild-type EWSR and fails to reduce MMEJ.  While knockout of EWS-FLI1 restores MMEJ, loss

of the oncoprotein is also detrimental to EwS cell growth and results in a reduction in the

transcription of EWS-FLI1 target genes which contributes to EwS cell growth.

Previous studies on the pathogenesis of Ewing Sarcoma have focused primarily on the

transcriptional activity of the EWS-FLI1 oncoprotein, driven by the carboxy-terminal FLI1

transcriptional domain.  Indeed, this oncoprotein upregulates the transcription of genes which in

turn promote R-loops and the neoplastic growth of the EwS tumor.  Our study also provides new

insights into the pathogenesis of the cancer.  While the chromosome 11/22 translocation results

in the formation of the EWS-FLI1 oncoprotein, it also simultaneously inactivates the EWSR1

*tumor suppressor* function.  One allele of this EWSR1 tumor suppressor gene is inactivated by

the translocation itself, and the remaining EWSR1 suppressor function is inactivated by the

binding of the fusion oncoprotein to the remaining EWSR1 protein.  Due to the loss of the

splicing activity of EWSR1, the tumor cells exhibit the characteristic MMEJ defect- namely,

radiation sensitivity, a defect in microhomology-mediated DSB repair, and a hypo-mutable

phenotype.  The splicing defect may also result in the inactivation of other tumor suppressor

genes and a more direct role in the process of oncogenesis.  For instance, the EWS-FLI1 fusion

protein may have additional direct oncogenic functions beyond its ability to promote

transcription, R-loops, and replication stress.  Specifically, the EWS-FLI1 fusion can disrupt

PLK1-activated POLΘ mitotic foci [39] which are known sites of TOPBP1-mediated DSB repair.

Accordingly, the EWS-FLI1 oncoprotein may further block the repair of DSBs in mitosis,

promote the formation of micronuclei, and enhance genomic instability.

Taken together, *Ewing sarcoma is the first known cancer with loss of MMEJ activity.* The

impairment of the MMEJ pathway in EwS prompted us to target NHEJ and HR, the two DNA

repair pathways known to be synthetic lethal with MMEJ loss.  Indeed, EwS cells are sensitive to

the NHEJ inhibitor, peposertib, as a monotherapy.   Peposertib is also synergistic with the

genotoxic drug, Etoposide, in killing EwS cells. Accordingly, the use of this drug combination

could reduce the need for high doses of Etoposide for EWS patients and thereby reduce its long-

term toxicity, such as therapy-related leukemia[60].   EWS cells, due to their MMEJ defect, are also

sensitive to the HR inhibitor, THZ531, a CDK12 inhibitor.  This inhibitor is one of the first

known inhibitors of the HR pathway.  Exon 25 skipping of the *POLQ* mRNA in fresh tumor

samples may provide a biomarker for identifying tumors with an underlying MMEJ defect which

may in turn respond to targeted NHEJ inhibitors or HR inhibitors through the mechanism of

synthetic lethality.

**ACKNOWLEDGMENTS**

We thank members of the D'Andrea laboratory for their helpful suggestions and comments. We thank Harvard Medical School and the Dana-Farber/Harvard Cancer Center in Boston, MA, for the use of the Animal, Flowcytometry, and Taplin Mass Spectrometry core facilities. This work was supported by U.S. National Institutes of Health grants R01HL052725 and R01CA296618 (A.D.D), the Breast Cancer Research Foundation (A.D.D.), the Ludwig Center at Harvard (A.D.D.), the David Liposarcoma Research Initiative at Harvard (A.D.D.), and the Smith Family Foundation (A.D.D.), and the Alex's Lemonade Stand Foundation (S.A.). This work was also supported by U.S. National Institutes of Health grants P50 CA168504 (A.D.D, G.I.S.), P50 CA240243 (A.D.D., G.I.S.), the SENSHIN Medical Research Foundation (S.A.), the Japan Society for the Promotion of Science Overseas Research Fellowships (S.A.), NCI K08 Career Development Award (R.G.), the DoD PRCRP Fellow Career Development Award (R.G.), the ASCO Career Development Award (R.G.), the Rally Foundation Career Development Award (R.G.) and the Boston Children's Hospital TRP Mentored Career Development Award (R.G.). Graphical abstract (Extended Fig, 9G) was created using BioRender.com.

**AUTHOR CONTRIBUTIONS**

S.A and A.D.D. conceived the study. S.A., G.Z., J.P.A., Y.H., D.R.I., S.M. and L.J. designed and performed biological experiments and analyzed the data. S.A. and Y.H designed and performed animal experiments. Y.T., H.N. and R.G. performed computational analysis. N.W.A. performed AlphaFold analysis. M.D.B., J.J.T and S.I.R. performed IHC analysis using patient samples.

K.P., E.M.V.A. and G.I.S. provided scientific inputs. S.A. and A.D.D. wrote the manuscript.

**FIGURE LEGENDS**

**Figure 1: *EWSR1* knock-out cells display MMEJ-deficient phenotypes through POLΘ**

**protein reduction.**

(A) Immunoblotting analysis for the detection of EWSR1 in RPE p53-/- clones edited with

CRISPR-Cas9.

(B, C) U2OS I-SceI-based (B, n=9) or Cas9-based (C, n=4) or were transduced lentivirally

transduced with control sgRNA or sgRNA targeting *EWSR1* or *POLQ* (co-expressing puromycin

resistance gene). Cells were treated with puromycin for 4 days, and then subjected to MMEJ

reporter analysis.

(D) Colony formation assay data of DLD1 *BRCA2* wild-type and *BRCA2* knockout cells

transduced with control, sg*POLQ* or sg*EWSR1* (n=3, left). Shown are representative images of the

colonies (right).

(E) RPE *TP53-/-* 3xFLAG-*POLQ* knock-in cells were lentivirally transduced with Cas9 and a

sgRNA targeting control (NT), *EWSR1* or *POLQ* (co-expressing puromycin resistance gene). After

puromycin selection, cells were subjected to immunoblotting analysis.

(F) HEK293T 3xFLAG-*POLQ* single knock-in cells and HEK293T 3xFLAG-*POLQ*/2xHA-

FKBP12 V36F *EWSR1* double knock-in cells are treated with DMSO or dTAG[V1] for indicated

time. Cell lysates were extracted and followed by immunoblotting analysis.

(G) Immunofluorescence analysis showing DNA damage-induced the foci formation of V5-tagged

endogenous POLΘ was decreased by EWSR1 depletion (scale bars = 2μm).

(H) U2OS MMEJ reporter cells (I-SceI based) were transduced with a control vector or POLQ

cDNA (co-expressing blasticidin resistance gene). After blasticidin selection, cells were

subsequently transduced with Cas9 and a sgRNA targeting control or EWSR1 (co-expressing

puromycin resistance gene). After puromycin selection, cells were subjected to MMEJ reporter

analysis (n=4).

(I) RPE *p53*-/- cells and RPE *p53*-/- *EWSR1* knockout cells were transduced with control, wildtype

*EWSR1* or *POLQ* cDNA. Colony formation assays were performed in the presence of Topotecan

(n=3).

563
564
**Figure 2: Disruption of MMEJ activity in Ewing sarcoma is driven by *POLQ* exon skipping.**

(A) Volcano plots showing alternatively spliced skipped exon (SE) in RPE *p53*-/- *EWSR1* knock-

out cells compared with RPE *p53*-/- cells defined by deep RNA-seq analysis.

(B) Sashimi plots of the alternative splicing events in control and *EWSR1* knock-out RPE *p53*-/-

cells: the region between exon24 and exon26 of the *POLQ* transcript.

(C) U2OS endogenous V5 tagged- POLΘ expressing cells were transduced with Cas9 and sgRNA

targeting the indicated exon of the *POLQ* gene. Cell lysates were extracted and followed by

immunoblotting analysis.

(D) U2OS MMEJ ver.2 (Cas9-based) cells were lentivirally transduced with control sgRNA or

sgRNA targeting different *POLQ* exons (co-expressing puromycin resistance gene). Cells were

treated with puromycin for 4 days, and then subjected to MMEJ reporter analysis (n=4).

(E) *POLQ* mRNA without exon 25 produces shorter POLΘ which is unstable. Blot showing

HEK293T cells transfected with FLAG-*POLQ* cDNA containing *POLQ* full-length CDS or CDS

lacking exon25. Cells were treated with CHX for 3–9 hr.

(F) HEK293T *POLQ* knockout cells were transfected with MMEJ reporter (ver.2) and sgRNA

targeting the reporter together with control vector or indicated *POLQ* cDNA. 48h after

transfection, cells were subjected to FACS analysis to measure the MMEJ activity (n=3).

(G) Schematic representation of the minigene reporter construct (left). K562 cells carrying the

minigene reporter were lentivirally transduced with Cas9 and sgRNA targeting the indicated genes.

Shown are relative frequencies of mNeonGreen positive fraction within mScarlet positive cells

(right, n=3).

(H) RT-PCR showing the exon 7 and exon 25 skippings of *POLQ* transcript in Ewing sarcoma cell

lines.

(I) Immunoblotting analysis showing the lower levels of POLΘ expression in Ewing sarcoma cell

lines compared to U2OS cells.

(J) rMATS analysis showing *POLQ* exon skipping events in primary samples from pediatric cancer

patients. n = 498 (EWS = 44, NBL = 131, OS = 88, WT = 110, RMS = 125).

(K) RNA-seq data from the patient sample shows the *POLQ* mRNA expression is significantly

lower in Ewing sarcoma patients compared with RMS/WT. n = 832 (EWS = 83, NBL = 195, OS

= 182, WT = 162, RMS = 210).

(L, M) Low somatic SNV and MMEJ deletion mutation burden in Ewing sarcoma tumors (ES; n

= 23) as compared to neuroblastoma tumors (NB; n = 47). Short somatic mutations were

determined from the whole genome sequencing data of tumors and matched normals.  Each dot in

K represents the total number of somatic SNVs for each tumor. Each dot in L represents the

proportion of 3+-bp deletions with 2+-bp microhomology among all indels of each tumor.

**Figure 3: Formation of EWSR1 condensates through the prion-like domain is critical for**

**faithful splicing of *POLQ* transcript via interacting with FUBP1 and KHSRP/FUBP2.**

**(A)** Schematic representation of the individual domains of EWSR1 and the deletion mutants used

in Fig.3B, C.

(B, D) RPE *p53-/-* 3xFLAG-*POLQ*/2xHA-FKBP12 V36F *EWSR1* double knock-in cells were

retrovirally transduced with empty vector, wildtype EWSR1, or indicated EWSR1 mutants. After

transduction, cells were treated with blasticidin for four days. Then, cells were treated with DMSO

or 1μM dTAG$^{v1}$ for 24hr. After dTAG$^{v1}$ treatment, cell lysates were subjected to immunoblot

analysis.

(C, E) RT-PCR showing the *POLQ* exon 25 skipping in the indicated cells.

(F) U2OS MMEJ ver.2 (Cas9-based) cells were lentivirally transduced with control sgRNA or

sgRNA targeting *EWSR1* together with empty vector wild-type EWSR1 or the indicated EWSR1

mutants by retroviral transduction. Cells were treated with puromycin and blasticidin for 4days,

and then subjected to MMEJ reporter analysis (n=4).

(G) Volcano plots comparing detected protein levels by IP-MS between EWSR1 wildtype and the

EWSR1 37YS mutant.

(H) A ven diagram showing the frequency of all the genes and positive hits with a prion-like

domain identified by PLAAC.

(I) Immunoprecipitation followed by immunoblotting analysis showing wildtype EWSR1

preferentially bound to FUBP1, KHSRP, SMARCA4 and SMARCC1 compared to the EWSR1

mutants.

(J) FUBP1/KHSRP DKO cells exhibited POLΘ protein reduction (upper) and increased *POLQ*

exon 25 skipping (lower).

(K) MMEJ activity was profoundly decreased in FUBP1/KHSRP DKO cells measured by U2OS

MMEJ ver.2 cells (n=4).

(L) Schematic representation of wildtype and YS-mutated EWSR1 or FUBP1.

(M) Immunoblotting showing wildtype FUBP1 preferentially bound to EWSR1 compared to the

FUBP1 16YS mutant.

(N) U2OS MMEJ ver.2 cells were lentivirally and retrovirally transduced with indicated sgRNA(s)

(co-expressing puromycin resistance gene) and control or FUBP1 cDNA (co-expressing

blastidcidin resistance gene). Cells were treated with puromycin and blasticidin for 4days, and then

subjected to MMEJ reporter analysis (n=4).

(O) RPE *p53*-/- 3xFLAG-*POLQ*/2xHA-FKBP12 V36F *EWSR1* double knock-in cells were

lentivirally transduced with Cas9 and a sgRNA targeting control (NT) or sgRNAs targeting FUBP1

and KHSRP together with retrovirally transduced with a control vector, wildtype FUBP1 or the

FUBP1 16YS mutant. After puromycin and blasticidin selection, cells were subjected to

immunoblotting analysis.

(P) RT-PCR showing the *POLQ* exon25 skipping in the indicated cells.

**Figure 4: Expression of EWS-FLI1 perturbs POLΘ foci assembly, resulting in impairment**

**of MMEJ.**

(A) I-SceI-based MMEJ reporter cells were lentivirally transduced with an empty vector, wild-

type *EWS-FLI1* or the FKBP12 V36F-tagged *EWS-FLI1*. Cells were treated with 1uM dTAG[V1]

for 18hr, and then adenovirally transduced with I-sceI. 48hr after I-sceI transduction, the

frequencies of GFP positive cells were assessed by FACS (n=6).

(B) Schematic representation of the EWS-FLI and mutant forms of EWS-FLI1 used in MMEJ

reporter assay (left). I-SceI-based MMEJ reporter cells were lentivirally transduced with an empty

vector, wild-type *EWS-FLI1* or indicated mutant forms of *EWS-FLI1* (co-expressing puromycin

resistance gene). After puromycin selection, cells were subjected to MMEJ reporter assay (right,

n=6).

(C) I-SceI based MMEJ reporter was lentivirally transduced with U2OS, A673, RH1 and RD-ES

cell lines. Cells were subsequently transduced with Cas9 together with control sgRNA or sgRNA

targeting FLI1. Cells were treated with puromycin for 4days, and then subjected to MMEJ reporter

analysis (n=3).

(D) EWS-FLI1 but not the EWS-FLI1 37YS mutant increased *POLQ* exon 25 skipping measured

by K562 minigene reporter cells (n=6).

(E) A673 and TC32 cells carrying the minigene reporter showing depletion of EWS-FLI1

decreased *POLQ* exon 25 skipping (n=3).

(F) Immunoprecipitation followed by immunoblotting analysis showing wildtype EWS-FLI11

preferentially bound to EWSR1 compared to the EWS-FLI1 37YS mutant.

(G) ChIP-qPCR assay showing expression of EWS-FLI1 perturbed POLΘ assembly to the DSB

loci induced by Isce-I (n=2).

(H) Immunofluorescence analysis showing endogenous V5-tagged POLΘ foci formation was

perturbed by EWS-FLI1 expression both asynchronous and mitotic arrested cells.  Cells were

treated with 2Gy irradiation and then incubated for 4hr (scale bars = 2μm).

(I) Immunofluorescence analysis showing endogenous V5-tagged POLΘ foci formation was

restored by EWS-FLI1 depletion in V5-tagged-*POLQ* knock-in A673 cells. Cells were treated with

2Gy irradiation and then incubated for 4hr (scale bars = 2μm).

**Figure 5: Inhibition of NHEJ or HR efficiently eliminates Ewing sarcoma cells.**

(A) The CTG assay showing that compared with U2OS, Ewing sarcoma cells showed

hypersensitivity to Peposertib.

(B) Immunoblot analysis showing phospho-DNAPK, DNAPK and GAPDH protein level in U2OS

and Ewing sarcoma cell lines.

(C, D) The CTG assay showing that a combination of Peposertib and Etoposide showed a strong

synergistic effect on Ewing sarcoma cell lines A673 (C) and TC32 (D).

(E, F) Xenograft model using TC32 showing therapeutic efficacy of a Peposertib and Etoposide.

NSG mice (n = 10 mice/group) bearing luciferized TC32 cells were treated with 100 mg/kg of

Peposertib (orally), 8 mg/kg etoposide (via IP), or both for 2 weeks. Tumor sizes were measured

every 2-3 days (E) and also monitored by bioluminescence images (F).

(G) Immunohistochemistry analysis from the patient samples of EwS (n=8) showing higher

pDNA-PKc activities compared with that of OS (n=9).

(H) Colony formation assay data of RPE *p53*-/- cells transduced with a control vector, EWS-FLI1

or EWS-FLI1 37YS together with or without BRCA2 depletion (n=3, top) and the representative

colony images (bottom).

(I, J) The CTG assay showing that compared with U2OS, Ewing sarcoma cells showed

hypersensitivity to CDK12 inhibitors, THZ531 (I) and MFH290 (J).

(K, L) The CTG assay showing that depletion of EWS-FLI1 by sg*FLI*1 in TC32 cells led to

acquisition of resistance to CDK12 inhibitors, THZ531 (K) and MFH290 (L).

**Extended Data Figure 1: *EWSR1* loss is synthetic lethal with BRCA1/2 deficiency.**

(A) A Venn diagram showing overlap between genes identified in the previously reported CRISPR

screens to identify synthetic *BRCA2* lethality and genes whose dependence is positively correlated

with *POLQ* in the Depmap analysis.

(B-F, I) Colony formation assay data of RPE *p53*-/- cells and RPE *p53*-/- *EWSR1* knock-out clones

treated with X-ray (B), VE822 (C), Topotecan (D), Etoposide (E), Olaparib (F) and Novobiocin (I

4) (n=4).

(G) U2OS DR-GFP reporter cells (I-SceI based) were transduced lentivirally transduced with

control sgRNA or sgRNA targeting EWSR1, BRCA1 or BRCA2 (co-expressing puromycin

resistance gene). Cells were treated with puromycin for 4days, and then subjected to reporter

analysis (n=3).

(H) Schematic representation of the Cas9-based MMEJ reporter assay (ver.2).

(J, K) Colony formation assay data of MDA-MB-436 (J, n=3) and HCC1937 (K, n=3) cells and

the cells complemented with BRCA1 cDNA transduced with control, sg*POLQ* or sg*EWSR1* (left).

Shown are representative images of the colonies (right).

(L) RPE *p53*-/- cells were lentivirally transduced Cas9 (expressing balasticidin resistance gene).

After balasticicin selection, cells were lentivirally transduced with a sgRNA targeting control,

*POLQ* or *EWSR1* (co-expressing GFP) together with with a sgRNA targeting control, *BRCA1* or

*BRCA2* (co-expressing tRFP657). 72hr after the sgRNA transduction, the frequency of the

GFP/tRFP657 cells was traced by FACS (n=3).

**Extended Data Figure 2: Subacute depletion of EWSR1 results in POLΘ protein loss.**

**(A)** RPE *TP53-/-* 3xFLAG-POLQ knock-in cells were transduced with Cas9 and a sgRNA

targeting *EWSR1* by electroporation. Single clones are isolated 7days after electroporation. Shown

are immunoblotting analyses for FLAF- POLΘ, EWSR1 and GAPDH in control cells and *EWSR1*

knockout clones.

(B) Colony formation assay data of RPE *p53-/-* 3xFLAG-*POLQ* knock-in cells and RPE *p53-/-*

*EWSR1* knock-out 3xFLAG-*POLQ* knock-in clones treated with X-ray (n=3).

(C) RPE *p53-/-* 2xHA-FKBP12 V36F *EWSR1* cells were treated with DMSO, 1μM dTAG[13] or

1μM dTAG[v1] for 24hr. The cell lysates were extracted and then analyzed by immunoblotting.

dTAG[v1] but not dTAG[13] efficiently degraded endogenous EWSR1 protein.

(D) RPE *p53-/-* 3xFLAG-*POLQ*/2xHA-FKBP12 V36F *EWSR1* double knock-in cells are treated

with 1μM dTAG[V1] for the indicated time. The cell lysates were extracted and then analyzed by

immunoblotting.

(E) RPE *p53-/-* 3xFLAG-*POLQ*/2xHA-FKBP12 V36F *EWSR1* double knock-in cells are treated

with dTAG[V1] at the indicated concentration for 24hr. The cell lysates were extracted and then

analyzed by immunoblotting.

(F) RPE *p53-/-* 3xFLAG-*POLQ* single knock-in cells and RPE *p53-/-* 3xFLAG-*POLQ*/2xHA-

FKBP12 V36F *EWSR1* double knock-in cells are treated with dTAG[V1] for indicated time. The cell

lysates were extracted and then analyzed by immunoblotting.

(G) Immunoblotting analyses showed that BRCA1, BRCA2 and other MMEJ proteins, except for

POLΘ, were not affected by *EWSR1* knockout in RPE *p53-/-* cells.

(H) RPE *p53-/-* 3xFLAG-*POLQ* knock-in cells were lentivirally transduced with control sgRNA

or sgRNA targeting *EWSR1, FUS* or *TAF15* (co-expressing puromycin resistance gene). Cells

were treated with puromycin for 4days, and then the cell lysates were extracted and then analyzed

by immunoblotting.

(I) K562 MMEJ ver.2 (Cas9-based) cells were lentivirally transduced with control sgRNA or

sgRNA targeting *EWSR1, FUS* or *TAF15* (co-expressing puromycin resistance gene). Cells were

treated with puromycin for 4days, and then cells were subjected to MMEJ reporter analysis (n=3).

(J) RPE *p53*-/- cells and RPE *p53*-/- *EWSR1* knockout cells were transduced with control, wildtype

*EWSR1* or *POLQ* cDNA. Colony formation assays were performed in the presence of Etoposide

(n=3).

(K) MDA-MB-436 cells and the cells complemented with BRCA1 were transduced with Cas9 and

a sgRNA targeting control or *EWSR1*. Cells were subjected to comet assay to measure unrepaired

DNA damage.

(L) RPE *p53*-/- cells and RPE *p53*-/- *EWSR1* knockout cells were transduced with Cas9 and a

sgRNA targeting control or *BRCA2*. Cells were treated with or without synchronization by 8μM

RO3306. 1 hour after removal of RO3306, cells were treated with 2 Gy irradiation and then

incubated for 24 hours. The frequency of micronuclei positive cells is shown.

**Extended Data Figure 3: EWSR1 prevents multiple exon skipping events of the *POLQ***

**transcript.**

(A) U2OS MMEJ reporter cells (I-SceI based) were transduced with *POLQ* cDNA (co-expressing

blasticidin resistance gene). After blasticidin selection, cells were subsequently transduced with

Cas9 and a sgRNA targeting control or *EWSR1* (co-expressing puromycin resistance gene). After

puromycin selection, cell lysates were extracted and then analyzed by immunoblotting. See also

Fig.2C.

757 (B) RPE *p53-/-* 3xFLAG-*POLQ* knock-in cells were transduced with Cas9 and a control sgRNA

758 or sgRNA targeting *EWSR1* by electroporation. 72hr after electroporation, cells were treated with

759 MG132 or Chloroquine for 4hr. After the treatment, cell lysates were extracted and then analyzed

760 by immunoblotting.

761 (C) A pie chart summarizing alternative splicing event types in RPE *p53-/- EWSR1* knock-out cells

762 obtained by MISO analysis.

763 (D-G) Volcano plots showing retained introns (RI) (D), alternatively spliced 5' splice cites (A5SS)

764 (E), alternatively spliced 3' splice cites (A3SS) (F), mutually exclusive exons (MXE) (G) in RPE

765 *p53-/- EWSR1* knock-out cells compared with RPE *p53-/-* cells.

766 (H) Volcano plots showing alternatively spliced skipped exon (SE) in HEK293T si*EWSR1*

767 transduced cells compared with control cells defined by RNA-seq analysis (GSE154944).

768 (I) Sashimi plots of the alternative splicing events in control and *EWSR1* knock-out RPE *p53-/-*

769 cells: the region between exon 23 and exon 26 of *POLQ* transcript.

770 (J) RT-PCR showing the exon 7, 21, 22 and 25 skipping of *POLQ* transcript in *EWSR1* knock-out

771 cells derived from RPE *p53-/-* (top) and from RPE *p53-/-*3xFLAG-*POLQ* knock-in cells (bottom).

772 (K) The schematics illustrates the location of protein domains with the primary sequence POLΘ.

773 Numbers refer to amino acid residues. Exo = exonuclease, T = thumb, P = palm, F = fingers. The

774 ribbon illustrations are AlphaFold 3-generated predictions of the WT and truncated form of the

775 POLΘ polymerase domain. The two views represent 180º rotation around the y-axis. Colors

776 illustrate the sub-domain architecture.

778 **Extended Data Figure 4: Exon 25 skipping of the *POLQ* transcript decreased POLΘ protein,**

779 **resulting in profound MMEJ deficiencies.**

(A)  Schematics of the *POLQ* exon 25, intron 24/25 and the design of ASO and sgRNAs targeting

the splice sites.

(B) Immunoblotting (upper) and RT-PCR (lower) analysis showing that transduction of two ASOs

efficiently induced exon 25 skipping and POLΘ protein loss in RPE *p53-/-* 3xFLAG-*POLQ* knock-

in cells.

(C) U2OS MMEJ ver.2 (Cas9-based) were transduced control or two ASOs targeting *POLQ* exon

25 splice sites. 48h after ASO transduction, cells were subjected to MMEJ reporter analysis (n=4).

(D) Schematic representation of the endogenous *POLQ* exon 25 and intron 24/25 and the minigene

constructs.

(E) HEK293T 2xHA-FKBP12 V36F *EWSR1* dknock-in cells were transfected with the indicated

minigene construct and then treated with DMSO or dTAG$^{V1}$. 48h after transfection, cells were

subjected to assess the endogenous- and minigene- derived *POLQ* exon 25 skipping (left) and

FACS analysis to measure the reporter activity (right).

(F-H) Binding of EWSR1 to *POLQ* pre-mRNA evaluated using the RNA immuno-precipitation

(RIP) assay, performed using an anti-EWS antibody and an anti-HA antibody (N=2). The assay

was performed in HEK293T 3xFLAG-POLQ/2xHA-FKBP12 V36F *EWSR1* double knock-in cells

in the presence of DMSO or dTAG$^{v1}$. qPCR was performed with primers targeting *POLQ* pre-

mRNA around exon 7 and 25, *POLQ* mature mRNA and *GAPDH* (bottom). Shown were

immunoblotting for cell lysates with or without immunoprecipitation (F), RIP-qPCR assay for the

samples with EWS-IP (G) and HA-IP (H).

(I) Binding of multiple EWSR1 mutants to *POLQ* pre-mRNA evaluated using the RNA immuno-

precipitation (RIP) assay, performed using anti-EWS antibody. The assay was performed in

HEK293T 3xFLAG-*POLQ*/2xHA-FKBP12 V36F *EWSR1* double knock-in cells transduced with

a control vector, EWSR1 WT, EWSR1 7YS mutant or EWSR1 37YS mutant in the presence of

DMSO or dTAG[v1]. qPCR was performed with primers targeting *POLQ* pre-mRNA around exon

7 and 25 and *GAPDH* (bottom).

**Extended Data Figure 5: Samples from Ewing sarcoma patients show no HRD genomic**

**signatures.**

(A-C) Contribution of COSMIC indel (ID)-signature (A) and stacked bar plots of the numbers of

indels attributed to COSMIC ID signatures for Ewing sarcoma tumors (ES) (in B), and

neuroblastoma tumors (NB) (in C). Short somatic mutations were determined from whole genome

sequencing data of tumors and matched normals using GATK4, and SigProfilerAssignment was

used to estimate the COSMIC ID signatures from the mutations. Each bar contains relative

numbers of indels attributed to specific ID signatures, which are shown in different colors. There

are no ES tumors with MMEJ-related ID6 indels (B). In contrast, there are several NB tumors with

ID6 indels (C).

(D-F) Contribution of COSMIC SNV-signature 3 (D) and stacked bar plots of the number of SNVs

attributed to COSMIC SNV signatures for Ewing sarcoma tumors (ES) (in E), and neuroblastoma

tumors (NB) (in F). Short somatic mutations were determined from whole genome sequencing

data of tumors and matched normals using GATK4, and SigProfilerAssignment was used to

estimate the COSMIC SNV signatures from the mutations. Each bar represents estimated numbers

of SNVs attributed to specific SNV signatures, which are shown in different colors. There are no

ES tumors with MMEJ-related Signature 3 SNVs (E). In contrast, there are several NB tumors

with Signature 3 SNVs (F).

(G-H) Distribution of insertions and deletions by size in Ewing sarcoma tumors (ES; n = 23), and

neuroblastoma tumors (NB; n = 47). Short somatic mutations were determined from the whole

genome sequencing data of tumors and matched normals. Each bar represents the proportion of

deletions (G) or insertions (H) of a particular size among all deletions or all insertions, respectively,

of each cohort.

(I) Low templated insertions in Ewing sarcoma tumors (ES; n = 23) as compared to neuroblastoma

tumors (NB; n = 47).

(J, K) The bar graphs show the length of microhomology at canonical translocation breakpoints in

Ewing sarcoma samples from GMKF (J, n=46) and St. Jude (K, N=43) cohorts. All translocations

were called using the Manta structural variant caller. Canonical translocations are defined as those

impacting one of the following genes: EWSR1, FLI1, ETV1, FEV, FUS, ERG.

**Extended Data Figure 6: The prion-like domain of EWSR1 is critical for the faithful splicing**

**of *POLQ* transcript.**

(A) RPE *p53*-/- 3xFLAG-*POLQ* knock-in cells were retrovirally transduced with empty vector,

wildtype *EWSR1*, or indicated *EWSR1* mutants (co-expressing blasticidin resistance gene)

together with lentivirally Cas9 and a control sgRNA or sgRNA targeting *EWSR1* (co-expressing

puromycin resistance gene). Cells were treated with puromycin and blasticidin for 4days, and

then cell lysates were extracted and then analyzed by immunoblotting.

(B) U2OS MMEJ ver.2 (Cas9-based) cells were lentivirally transduced with control sgRNA or

sgRNA targeting *EWSR1* together with empty vector wild-type EWSR1 or the indicated EWSR1

mutants by retroviral transduction. Cells were treated with puromycin and blasticidin for 4days

and then subjected to MMEJ reporter analysis (n=4).

(C-E) Colony formation assay data of RPE *p53-/-* cells and RPE *p53-/- EWSR1* knock-out clones

transduced with an empty vector, wild-type *EWSR1* or the indicated *EWSR1* mutants by

retroviral transduction in the presence of 10nM Topotecan (C), 62.5nM Etoposide (D), or 100nM

VE822 (E), respectively. The data is normalized to the DMSO control (n=3).

(F, G) K562 cells were lentivirally transduced with sgRNAs targeting control or *EWSR1* (co-

expressing puromycin resistance gene) and control or *BRCA2* (co-expressing blasticidin resistance

gene). Cells were treated with puromycin and blasticidin for 4days, and then retrovirally

transduced with indicated constructs (co-expressing GFP). 72hr after retroviral transduction, the

change of frequency of GFP positive fraction was traced by FACS (n=3).

(H) Schematic representation of the deletion mutants within the prion-like domain used in Fig. S6I.

(I) RPE *p53-/-* 3xFLAG-*POLQ*/2xHA-FKBP12 V36F *EWSR1* double knock-in cells were

retrovirally transduced with empty vector, wildtype EWSR1, or indicated EWSR1 mutants. After

transduction, cells were treated with blasticidin for four days. Then, cells were treated with DMSO

or 1μM dTAG$^{v1}$ for 24hr. After dTAG$^{v1}$ treatment, cell lysates were subjected to immunoblot

analysis.

(J) Wildtype EWSR1 but not the EWSR1 7YS or 37YS mutant rescued POLΘ protein expression

in RPE *p53-/-* 3xFLAG-*POLQ* knock-in cells.

(K) Wildtype EWSR1 but not the EWSR1 7YS or 37YS mutant prevented *POLQ* exon 25 skipping

measured by K562 minigene reporter cells (n=3).

**Extended Data Figure 7: Exon skipping in *EWSR1* KO cells was typically observed for exons**

**that are flanked by longer introns.**

(A) Volcano plots comparing detected protein levels by IP-MS between EWSR1 wildtype and

control.

(B) The length of introns surrounding included, excluded or background exons in *EWSR1*

knockout cells identified by RNA-seq analysis.

(C) Dual depletion of FUBP1 and KHSRP increased *POLQ* exon 25 skipping measured by K562

minigene reporter cells (n=3).

(D) The FUBP1 16YS mutant failed to rescue *POLQ* exon 25 skipping induced by FUBP1/KHSRP

dual depletion measured by K562 minigene reporter cells (n=3).

(E) RPE *p53*-/- 3xFLAG-*POLQ*/2xHA-FKBP12 V36F *EWSR1* double knock-in cells were

lentivirally transduced with Cas9 and a sgRNA targeting control (NT) or sgRNAs targeting

FUBP1 and KHSRP together with retrovirally transduced with a control vector, wildtype

KHSRP or the KHSRP 16YS mutant. After puromycin and blasticidin selection, cells were

subjected to immunoblotting analysis.

(F) RT-PCR showing the *POLQ* exon25 skipping in the indicated cells.

(G) U2OS MMEJ ver.2 (Cas9-based) cells were lentivirally transduced with indicated sgRNA(s)

(co-expressing puromycin resistance gene). Cells were treated with puromycin for 4days, and

then subjected to MMEJ reporter analysis (n=4).

(H) The KHSRP 17YS mutant failed to rescue *POLQ* exon 25 skipping induced by

FUBP1/KHSRP dual depletion measured by K562 minigene reporter cells (n=3).

**Extended Data Figure 8: Depletion of EWS-FLI1 ameliorates MMEJ activity in Ewing**

**sarcoma cells.**

(A) The cell lysates were extracted from an Osteosarcoma cell line U2OS and Ewing Sarcoma cell

lines (A673, TC32, TC71, RH1, and RD-ES, and then analyzed by immunoblotting.

(B) Cas9-based MMEJ reporter cells were lentivirally transduced with an empty vector, wild-type

*EWS-FLI1* or indicated mutant forms of *EWS-FLI1* (co-expressing puromycin resistance gene).

After puromycin selection, cells were subjected to MMEJ reporter assay (n=3).

(C, D) the CellTiter-Glo® (CTG) assay showing Ewing sarcoma cells showed resistance to POLΘ

inhibitors, Novobiocin (C) and ART558 (D).

(E, F) Colony formation assay data of RPE *p53*-/- cells lentivirally transduced with control vector,

EWS-FLI WT or the EWS-FLI1 37YS mutant treated with Etoposide (E) and Topotecan (F) (n=2).

(G) The cell lysates used in Fig.4C were extracted and then analyzed by immunoblotting. The

black and red arrows showing EWS-FLI1 and FLI1, respectively.

(H) The CTG assay shows that depletion of EWS-FLI1 in A673 cells sensitized to ART558.

(I) U2OS I-SceI-based MMEJ reporter cells were transduced with control or EWS-FLI1 together

with an empty vector, RNaseH1 WT, the RNaseH1 DN, or the RNaseH1 WKKD mutant. Cells

were treated with puromycin and blasticidin for 4days and then subjected to MMEJ reporter

analysis (n=6).

(J) A673 carrying the Isce-I-based MMEJ reporter were transduced with an empty vector,

RNaseH1 WT, the RNaseH1 DN, or the RNaseH1 WKKD mutant. Cells were treated with

blasticidin for 4days and then subjected to MMEJ reporter analysis (n=3).

**Extended Data Figure 9:  Combination of Peposertib and Etoposide efficiently killed Ewing**

**sarcoma cells in vivo.**

(A, B) The CTG assay showing that compared with U2OS, Ewing sarcoma cells showed

hypersensitivity to Topotecan (A) and Etoposide (B).

(C) Overall survivals showing a therapeutic efficacy of a combination of Etoposide and Peposertib

in a TC32 xenograft model (n=10, each group).

(D) The changes of the body weight of the treated mice.

(E, F) The CTG assay showing that depletion of EWS-FLI1 in A673 cells led to acquisition of

resistance to CDK12 inhibitors, THZ531 (E) and MFH290 (F).

(G) The schematic representation of the findings of this study.

## EXPERIMENTAL MODEL AND SUBJECT DETAILS

### Cell cultures

RPE *p53*-/- cells generated previously[61] were grown in DMEM/F12 containing GlutaMAX

(Gibco) and supplemented with 10% fetal bovine serum (FBS) (Sigma-Aldrich). HEK293T

(ATCC, CRL-3216) cells were grown in DMEM (Gibco) supplemented with 10%FBS. TC32 and

TC71 cell lines were obtained from children cancer repository and grown in RPMI (Gibco)

supplemented with 10%FBS. RH-1 (DSMZ, ACC493) cell grown in RPMI 1640 supplemented

with 10%FBS, 1% P/S. RD-ES (ATCC, HTB-166) cells were grown in RPMI 1640 supplemented

with 20%FBS, 1% P/S. SK-ES1 (ATCC, HTB-86) and SK-PN-DW (ATCC, CRL-2139) cells

were cultured in DMEM (Gibco) supplemented with 10%FBS. U2OS parental (ATCC, HTB-96)

and carrying a DNA repair template cells were grown in DMEM (Gibco) supplemented with

10%FBS. The BRCA1-deficient cell lines HCC1937 (ATCC, CRL-2336) and HCC1954 (ATCC,

CRL-2338) and BRCA1-complemented cells were grown in RPMI (Gibco) supplemented with

10%FBS. RH-1 (DSMZ, ACC493). K562 (ATCC, CCL-243) cells were grown in RPMI 1640

supplemented with 20%FBS, 1% P/S. MDA-MB-436 and MDA-MB-436+BRCA1 cells were a

kind gift from Dr. Geoffrey Shapiro (Dana Farber Cancer Institute) and grown in RPMI (GIBCO)

supplemented with 10% fetal bovine serum. DLD1 BRCA2 -/- and DLD1 Parental were obtained

from Horizon discovery, Waterbeach, UK. U2OS cells with HR reporter (DR-GFP) was a kind

gift from Dr. Jeremy Stark (City of Hope).

To establish the U2OS-MMEJ reporter cells, U2OS cells were lentivirally transduced with the

Isce-I-based (ver.1) or Cas9-based (ver.2) MMEJ reporter and then incubated with puromycin.

After selection with puromycin, a single clone was isolated and subsequently the clones were

validated.

All cells were cultured in an incubator maintained at 37°C, 5% $CO_2$ and a relative humidity of

95%.

**Compounds and antibodies**

Chemical compounds, including Olaparib (#S1060), Novobiocin (#S2492), Etoposide (#S1225),

VE822 (#S7102), Ro-3306 (#S7747), M3814 (#S8586) and THZ0531 (#$6595) were purchased

from Selleckchem. ART558 (#HY-141520) and MFH-290 (#HY-153244) were purchased from

MedChemExpress. Nocodazole (#1404) was purchased from Sigma Aldrich. Chemicals were

dissolved in DMSO and kept in small aliquots at −20°C or 80°C. The key sources of antibodies

are: POLθ (Cell Signaling Technology, #64708), γH2AX (Millipore #05–636), LIG3 (BD

Biosciences, #611876), HMCES (Santa Cruz Biotechnology, SC-514238), HMCES (Atlas

Antibodies #HPA044968), APEX2 (Cell Signaling Technology, #74728), PARP1(Cell Signaling

Technology, #9542), CIP2A (Abcam, ab128179), DNA-PK (Cell Signaling Technology, #12311),

Phospho-DNA-PKcs (Ser2056) (Cell Signaling Technology, #68716), TOPBP1 (Abcam, ab2402),

XRCC1 (Cell Signaling Technology, #76998), FEN1 (Cell Signaling Technology, #82354),

EWSR1 (Cell Signaling Technology, #11910 for WB), EWSR1 (Santa Cruz Biotechnology, sc-

28327 for WB), EWSR1 (Life Technologies, A300417A for IP and RIP), BRCA1 (Merck

Millipore, OP92), BRCA2 (Cell Signaling Technology, #10741), FLI1 (Abcam, ab133485), FUS

(Cell Signaling Technology, #67840), TAF15(Cell Signaling Technology, #28409), Vinculin (Cell

Signaling Technology, #4650), FLAG (Sigma, F1804), HA (Cell Signaling Technology, #3724

for WB), HA (Roche, #11867431001 for IP and RIP), V5 (Cell Signaling Technology, #13202 for

WB and IF), V5(Abcam, ab15828 for ChIP), V5(Abcam, ab27671 for IP and WB), GAPDH (Cell

Signaling Technology, #5174), Histone H3 (Cell Signaling Technology, #9715), FUBP1 (Abcam

ab181111), KHSRP (Cell Signaling Technology, #13398), SMARCA4/BRG1 (Cell Signaling

Technology, #3508), SMARCC1 (Cell Signaling Technology, #11956), MYC (Santa Cruz

Biotechnology, SC-40).

**Generation of CRISPR engineered cell lines**

For generating *EWSR1* knock-out clones in HEK293T and RPE *p53*-/- cells, a ribonucleoprotein

(RNP) complex was formed by Alt-R™ S.p. HiFi Cas9 and CRISPR-Cas9 sgRNA targeting

*EWSR1* (Integrated DNA Technologies). For generating endogenously 3xFLAG-tagged *POLQ*

knock-in HEK293T and RPE *p53*-/- cells, 3xFLAG-tagged *POLQ* HEK293T *EWSR1*-FKBP12

V36F double knock-in cells or V5-tagged *POLQ* knock-in U2OS and A673 cells, the Cas9/sgRNA

RNP complex together with a template was delivered. Cells were resuspended in SE (U2OS cells),

SF (HEK293T and A673 cells) or P3 (RPE *p53*-/- cells) Nucleofector Solution with supplement

(Lonza) and then mixed with the Cas9/sgRNA RNP complex and Alt-R™ Cas9 Electroporation

Enhancer (Integrated DNA Technologies), respectively. The Cas9/sgRNA RNP complex was

delivered to the cells by 4D-Nucleofector™ (Lonza). Media was changed the following day, and

then 2 days after cells were seeded in a 96-well plate to isolate single clones. Individual clones

were tested for genomic editing analyses using immunoblotting and genomic PCR with subsequent

Sanger sequencing.

**Plasmids**

psPAX2 and pMD2.G (Addgene #12260 and #12259) were gifts from Dr. Didier Trono. pLV-

EF1a-IRES-Blast (pLV-Blast) (Addgene #85133) was a gift from Dr. Tobias Meyer.[62]

990 pCDH-puro-EWS-FLI1 was a gift from Jialiang Wang (Addgene plasmid # 102813). pLenti

991 CMV Puro LUC (w168-1) was a gift from Eric Campeau & Paul Kaufman (Addgene plasmid #

992 17477). pCVL.SFFV.d14GFP.EF1a.HA.NLS.Sce(opt).T2A.TagBFP was a gift from Andrew

993 Scharenberg (Addgene plasmid # 32627). EWS-FLI1 37YS mutant was cloned into pCDH-puro

994 vector. EWSR1, FUBP1, KHSRP, RNaseH1 and its mutant forms were cloned into pMYs-IRES-

995 Blastcidin vector, respectively. sgRNAs were cloned into a lentiCRISPR-v2 construct as

996 previously described[63]. Of note, sgEWSR1 #2, sgFUBP1 #2 or sgKHSRP #2 target genomic

997 DNA but not cDNA, so that endogenous knockout and ectopic cDNA expression are possible

998 simultaneously. All the sgRNA sequences used in this study were provided in Supplemental

999 Table 1.

1000

1001 **Virus generation**

1002 Lentivirus was generated by co-transfecting lentiviral constructs with packaging vectors, psPAX2

1003 and pMD2.G, into HEK293T cells using CalPhos Mammalian Transfection Kit (Takara Bio). Two

1004 days after transfection, supernatant containing virus was harvested and filtered through 0.45 μm

1005 filter. Viral transductions were performed in the presence of 8 μg/ml (for RPE *p53*-/-, MDA-MB

1006 436, DLD1, HCC1937, and HCC1954 cells) or 4 μg/ml (for K562 cells) polybrene (Sigma-

1007 Aldrich). For all virus generations and transductions media supplemented with heat inactivated

1008 FBS was used. Retrovirus was generated by co-transfecting retroviral constructs with packaging

1009 vectors, and, into HEK293T cells using CalPhos Mammalian Transfection Kit (Takara Bio). Two

1010 days after transfection, supernatant containing virus was harvested and filtered through 0.45 μm

1011 filter. Viral transductions were performed in the presence of 8 μg/mL polybrene (Sigma-Aldrich).

1012 Puromycin dihydrochloride (Sigma-Aldrich) or Blasticidin S HCl (Thermo Fisher Scientific) was

1013 used after infection of retrovirus or lentivirus.

1014

**Colony formation assay**

Cells were seeded in 6-well plates at 500-10,000 cells/well. The day after seeding, cells were

exposed to each drug or x-ray. X-ray irradiation was performed using RS-2000 Irradiator (Rad

Source Technologies). After incubation, cells were washed with PBS and then fixed in a mixture

of 50% methanol, 10% acetic acid, and 40% water for 20 minutes. The fixed cells were stained

with 0.5% crystal violet (Sigma-Aldrich) in 20% methanol for 2–6 h and then washed with water

twice. The plates were imaged on GE Amersham Imager 600 using colorimetric transillumination

setting. The area fraction of the colonies in each well was estimated using ImageJ.

1023

**CellTiter-Glo assay**

To test drug sensitivity, U2OS or Ewing sarcoma cells were seeded in 96-well plates at 1000–3000

cells/well. The day after seeding, drug was added to the wells, and then the plates were incubated

for 3-5 days. After incubation, ATP in each well was quantified using CellTiter-Glo Luminescent

Cell Viability Assay (Promega). CellTiter-Glo assays were performed in technical duplicate if not

otherwise specified in a figure legend.

1030

**Immunofluorescence**

Mitotic synchronization was achieved using Ro-3306 (8 μM, 16 hours) followed by 30min release

in fresh medium. Then cells were incubated with 100ng/ml nocodazole and subsequently

irradiated. 4h after irradiation or in the absence of treatment, cells were fixed with 4%

paraformaldehyde for 15 minutes at room temperature. Cells were then permeabilized with 0.3%

Triton X-100 for 10 minutes on ice, followed by blocking with 3% non-fat milk for 1 hour at room

temperature. The slides were stained with primary antibodies at 4°C overnight. Afterward, they

were stained with secondary antibodies for 1 hour at room temperature. The slides were scanned

using fluorescence microscope. At least 100 cells were counted for each sample. Foci

quantification was performed using software CellProfiler version 4.2.6.

1041

**Immunoblotting**

Whole cell lysate was prepared by lysing cells in ice-cold RIPA buffer (Cell Signaling

Technology) supplemented with Protease/Phosphatase inhibitor Cocktail (Cell Signaling

Technology). Cell lysate was cleared by centrifuging the samples at 12000 rpm, 4°C for 20

minutes. Supernatant was recovered and protein concentration was estimated using BCA assay kit

(Thermo Fisher Scientific). Protein samples were boiled in 2X Laemmli sample buffer (Bio-rad)

supplemented with 5% 2-mercaptoethanol or NuPAGE LDS sample buffer (Thermo Fisher

Scientific) supplemented with NuPAGE sample reducing agent (Thermo Fisher Scientific).

Samples were run in NuPAGE 4-12% Bis-Tris gels (Thermo Fisher Scientific) using NuPAGE

MES SDS Running Buffer (Thermo Fisher Scientific) or NuPAGE 3-8% Tris-Acetate Gel

(Thermo Fisher Scientific) using NuPAGE Tris-Acetate SDS Running Buffer (Thermo Fisher

Scientific). After electrophoresis, samples were transferred onto a nitrocellulose membrane

(Amersham). Membranes were blocked for 1 h with Blocking Buffer, which was prepared by

diluting Fish Serum Blocking Buffer (Thermo Fisher Scientific) with Tris-buffered saline

containing 0.1% Tween20 (TBST). Membranes were incubated with primary antibodies diluted in

Blocking Buffer overnight at 4°C. Membranes were washed with TBST three times, 5 minutes

each, incubated with secondary antibodies diluted in blocking buffer for 1h, which either

horseradish peroxidase (HRP)-conjugated for chemiluminescent detection or fluorophore-

conjugated for visualization using the LI-COR imaging system.  After secondary antibody

incubation, membranes were washed with TBST and imaged using chemiluminescence setting on

GE Amersham Imager 600 or the LI-COR imaging system.

1063

**DNA repair template assay**

DR-GFP and EJ5-GFP cells were kind gifts from Dr. Jeremy Stark. MMEJ-GFP cells (ver.1 and

ver.2) were generated by lentivirally transducing a cassette into U2OS or Ewing sarcoma cells

using a pLV-EF1a-IRES-puro vector (addgene #85132) as previously described [20]. U2OS cells

carrying a DNA repair template reporter (DR-GFP, EJ5-GFP, and MMEJ-GFP) or Ewing sarcoma

cells carrying a MMEJ reporter were lentivirally transduced with Cas9 and sgRNA (co-expressing

puromycin resistance gene). After puromycin or blasticidin selection, 40,000 DNA repair template

reporter cells were seeded in 12-well plates and adenovirally (for U2OS) or lentivirally (co-

expressing BFP, for Ewing sarcoma cells) transduced with Isce-I the following day. 48h after Isce-

I transduction, GFP signals were analyzed by CytoFLEX (Beckman).  The signals were normalized

to control cells. For U2OS Cas9-based MMEJ reporter assay, the cells were subsequently

transduced with Cas9 and sgRNA targeting the reporter (co expressing BFP). 72h after

Cas9/sgRNA transduction, GFP signals within BFP positive fraction were analyzed by CytoFLEX

(Beckman).  For Cas9-based MMEJ reporter assay using HEK293T *POLQ* WT or knockout cells,

the MMEJ reporter, Cas9 and sgRNA targeting the reporter (co expressing BFP) together with a

control vector, *POLQ* WT cDNA or *POLQ* mutant cDNA were transduced using Lipofectamine

LTX with Plus reagent (Thermo Fisher Scientific). After 72 h incubation, GFP signals within BFP

positive fraction were analyzed by CytoFLEX (Beckman).

1082

**Comet assay**

The alkaline comet assays were performed to detect both single and double-stranded DNA breaks. 1000 cells, suspended in PBS, were mixed with 50µl melted low melting agarose and pipetted onto the Cometslides (R&D Systems). Once the agarose solidified, the slides were immersed in a lysis solution for 18 hours to facilitate cell lysis. Following lysis, the slides were incubated in an alkaline unwinding solution for 1 hour to denature the DNA. Electrophoresis was performed at 21 V for 45 minutes in an alkaline electrophoresis solution. The slides were stained with SYBR green solution, scanned by fluorescence microscope, and analyzed using CometScore 2.0 software.

**Chromatin immunoprecipitation (ChIP) assay**

U2OS biallelic endogenous V5-*POLQ* expressing cells carrying an Isce-I based MMEJ reporter were adenovirally transuduced with Isce-I in the presence or absence of 100ng/ml nocodazole. 12h after Isce-I trunscduction, chromatin immunoprecipitation was performed using SimpleChIP Enzymatic Chromatin IP Kit (Magnetic Beads) (Cell Signaling Technology). Protein-DNA cross-linking was performed with 1% formaldehyde and glycine solution. Chromatin was fragmented with Micrococcal Nuclease. The digested chromatin was incubated at 4°C with each antibody for 4 h. After the antibody reaction, ChIP-Grade Protein G Magnetic Beads were added to the samples, and the samples were incubated at 4°C for 1–2 h. For crosslinking reversal, each sample was incubated at 65°C with Proteinase-K overnight. After DNA purification, immunoprecipitated DNA was analyzed by qPCR using a primer set targeting the integrated MMEJ reporter locus. ChIP-qPCR was performed in technical triplicate. All the primer sequences for ChIP-qPCR used in this study were provided in Supplemental Table 1.

**Reverse transcription PCR (RT-PCR)**

Total RNA was extracted using RNeasy Mini Kit (QIAGEN). Complementary DNA synthesis was performed using SuperScript IV First-Strand cDNA Synthesis Reaction (Thermo Fisher Scientific). The PCR products were separated in 2% agarose gel stained with SYBR Safe (Thermo Fisher Scientific) and imaged using ChemiDoc MP Imaging System (Bio-rad). All the primer sequences for RT-PCR used in this study were provided in Supplemental Table 1.

**ASO transfection**

ASO was designed to target near the splice sites of the *POLQ* exon 25. The ASO was synthesized using 2′-O-methoxyethyl-phosphorothioated bases by Integrated DNA Technologies (IDT). Prior to ASO transfection, cells were seeded in a 6-well plate and cultured overnight. ASO was transfected at a final concentration of 0.1 µM using Lipofectamine LTX with Plus reagent (Thermo Fisher Scientific). After 48 h incubation, cells were harvested and analyzed by RT-PCR and western blotting. All the sequences for ASO used in this study were provided in Supplemental Table 1.

**Splicing minigene reporter assay**

A mini-gene reporter carrying mNeongreen split by synthetic intronic sequence derived from *POLQ* intron 24/25 and exon 25 was generated as previously described [28]. The synthetic intron was made up of 250 bp of the start of the *POLQ* intron 24, 250 bp or 1000bp of the end of the *POLQ* intron 24, the *POLQ* exon 25, 250 bp or 1000bp of the start of the *POLQ* intron 25m and 250 bp of the end of the *POLQ* intron 25. The synthetic introns and exon so generated were placed between the sequences encoding the N-terminus and C-terminus of split mNeongreen.

The mScarlet-P2A-split mNeongreen (N-terminus)-synthetic introns/exon-split mNeongreen (C

terminus) construct was integrated into a pLV-EF1a-IRES-Neo vector (Addgene #85139) to

generate the mini-gene reporter. K562 cells or Ewing sarcoma cell lines were lentivirally

transduced with the mini-gene reporter. 48hr after lentiviral transduction, the cells were enriched

by using G418 (0.8mg/ul) for seven days. After antibiotic selection, mScarlet and mNeongreen

signals in the reporter cells were analyzed by CytoFLEX (Beckman).  For the flow cytometry

analyses, mScarlet positive fractions within living cells were gated. Thereafter, the frequency of

the mNeongreen positive cells within mScarlet expressing cells was analyzed by using Flowjo

ver.10.9.0.

1138

**Mass spectrometry analysis**

HEK293T EWSR1-FKBP12 V36F knock-in cells were seeded into 15cm dishes and cultured

overnight. The cells were transfected with pMYS-IB empty vector (EV) or pMYs-IB-EWSR1

wildtype or pMYs-IB-EWSR1 37YS plasmid. Cells were transfected with 30 μg of vector /dish

using Lipofectamine LTX with Plus reagent. 16h after transfection, media was changed with 1uM

dTAG[v1]. The cells were harvested 40 h after transfection and then were lysed with 40 mM Tris-

HCl pH 8 buffer supplemented with 0.5% NP-40, 200 mM NaCl, 2 mM EDTA,

Protease/Phosphatase inhibitor Cocktail (Cell Signaling Technology), and MG-132. The lysate

was centrifuged, and the pellet was resuspended with 20 mM Tris-HCl pH 7.5 supplemented with

100 mM KCl, 2 mM MgCl2, 1 mM CaCl2, and Protease/Phosphatase inhibitor Cocktail and treated

with Micrococcal Nuclease (Thermo Fisher Scientific) for approximately 30 minutes, followed by

addition of 5 mM EGTA (final concentration) to stop digestion. After three pulses of sonication

(30 s each), samples were centrifuged to collect the supernatants as chromatin fractions.

Immunoprecipitation was performed using Anti-FLAG M2 antibody (Sigma-Aldrich) conjugated

with Dynabeads Protein G for Immunoprecipitation (Thermo Fisher Scientific). Beads were

washed with TGN-150 wash buffer containing 20 mM Tris-HCl pH 7.6, 150 mM NaCl, 3 mM

MgCl2, 10% glycerol, and 0.01% NP-40. Beads were then boiled at 70 C for 20 minutes in 1:1

diluted NuPAGE LDS Sample buffer (4X) (Thermo Fisher Scientific) to elute the proteins. The

eluted proteins were applied to SDS-PAGE, and Coomassie stained gel band samples were used

for the analyses. Two independent biological replicate samples were analyzed by mass

spectrometry.

The gel bands were cut into approximately 1 mm3 pieces. Gel pieces were then subjected to a

modified in-gel trypsin digestion procedure. Gel pieces were washed and dehydrated with

acetonitrile for 10 minutes and completely dried in a SpeedVac (Thermo Fisher Scientific). The

gel pieces were rehydrated with 50 mM ammonium bicarbonate containing 12.5 µg/mL modified

sequencing-grade trypsin (Promega) at 4°C. After rehydration, the excess trypsin solution was

removed and replaced with 50 mM ammonium bicarbonate solution. Samples were then incubated

at 37°C overnight. Peptides were later extracted with a solution containing 50% acetonitrile and

1% formic acid and dried in a SpeedVac. The samples were reconstituted in 5–10 µL of HPLC

solvent A (2.5% acetonitrile, 0.1% formic acid). A nano-scale reverse-phase HPLC capillary

column was created by packing 2.6 µm C18 spherical silica beads into a fused silica capillary (100

µm inner diameter x ~30 cm length) with a flame-drawn tip. After equilibrating the column each

sample was loaded via an EASY-nLC (Thermo Fisher Scientific). A gradient was formed, and

peptides were eluted with increasing concentrations of solvent B (90% acetonitrile, 0.1% formic

acid). As peptides eluted, they were subjected to electrospray ionization and then entered into a

Orbitrap Exploris480 mass spectrometer (Thermo Fisher Scientific). Peptides were detected,

isolated, and fragmented to produce a tandem mass spectrum of specific fragment ions for each

peptide. Peptide sequences were determined by matching protein databases with the acquired

fragmentation pattern by the software program, Sequest (Thermo Fisher Scientific). All databases

include a reversed version of all the sequences and the data was filtered to between a one and two

percent peptide false discovery rate. Perseus (version 2.0.11.0) [64] was used to preprocess the

protein data. The protein intensities were normalized by the median and log2-transformed. Proteins

with only one intensity value or with a sum of all total peptide counts of less than 5 were filtered

out. For each sample, missing log2-intensity values were imputed using the normal distribution

with width of 0.3 and down shift of 1.8. Histograms of log2-instensity values were inspected to

confirm that the imputed values were at the lower end. Finally, limma analysis (version 3.54.2) [65]

was performed on the log2-instensity values to determine enriched and depleted proteins. Volcano

plots were generated using ggplot2 and ggrepel R packages.

1187

**RNA immuno-precipitation (RIP) assay**

Protocol for RIP assay was adapted from Nicholson-Shaw et al., 2022[29]. HEK293T EWSR1-

FKBP12 V36F knock-in cells were cultured in 15 cm dishes and 2-3 plates were used per cell line

for each assay. 16hr after transfection, media was changed with 1uM dTAG[v1]. Cells were harvested

40hr after transfection, washed with ice-cold PBS 2X times, cross-linked with 0.2% PFA for 15

min. and then quenched by adding glycine to a final concentration of 125mM. Cells were lysed in

iCLIP buffer (50mM Tris-HCl pH 7.4, 100mM NaCl, 1% NP-40, 0.1% SDS, 0.5% sodium

deoxycholate) supplemented with protease, phosphatase and RNase inhibitor cocktail. The

samples were rotated at 4°C for 10 min. and then sonicated on 'light' setting twice. The lysate was

incubated with Protein A beads on a rotor for 10 min. at 4°C to pre-clear the extract. The lysate

was centrifuged at max speed for 10 min. and the supernatant was recovered and incubated for 2

hrs on a rotor at 4°C with Protein A beads coupled to EWSR1 Ab (Life Technologies). A part of

the lysate (0.4-0.6%) was kept aside as input. After 2 hrs the beads were washed twice with iCLIP

buffer. The input and the IP samples were all subjected to TURBO DNase treatment (Thermo

Fisher Scientific) for 30 min. at 37°C to get rid of the genomic DNA and the cross-links were

reversed by digesting with Proteinase K (NEB) for 30 min. at 37°C and further digestion and

denaturation of proteins was achieved by adding urea to a final concentration of 2.5M and

incubating the samples at 37°C for additional 20 min. After digestion of the DNA and protein from

the samples, the RNA was purified by adding Trizol and processing the samples through Directzol

RNA mini-prep kit (Zymo Research). The eluted RNA from the input and IP samples were

converted to cDNA using the High-capacity RNA-to-cDNA kit (Thermo Fisher Scientific). The

reverse transcription was performed with and without adding the reverse transcriptase enzyme.

qPCR was set up with primers targeting *POLQ* pre-mRNA, *POLQ* mature mRNA, and GAPDH

in technical replicates for each assay. Ct values were normalized to that from the control samples. .

All the sequences for RIP-qPCR used in this study were provided in Supplemental Table 1.

1213

**Next generation sequencing**

Library preparation and sequencing reactions were conducted at GENEWIZ, Lin/Azenta US, Inc.

Deep RNA-seq (300 million reads/sample, N = 3 for each cell line) was performed for RPE *p53-*

*/-* (parental control) and *p53-/-EWSR1* knock-out cells. RNA samples were prepared using RNeasy

Mini Kit and RNase-Free DNase Set (QIAGEN). RNA samples were quantified using Qubit 2.0

Fluorometer, and the RNA integrity was checked with 4200 TapeStation. NEBNext Ultra II RNA

Library Prep Kit for Illumina was used following the manufacturer's recommendations (New

England BioLabs). Briefly, mRNAs were initially enriched with Oligo d(T) beads. Enriched

mRNAs were fragmented for 15 minutes at 94°C. First strand and second strand cDNA were

subsequently synthesized. cDNA fragments were end repaired and adenylated at 3′ends, and

universal adapters were ligated to cDNA fragments, followed by index addition and library

enrichment by PCR with limited cycles. The sequencing libraries were validated using Agilent

TapeStation and quantified using Qubit 2.0 Fluorometer as well as by quantitative PCR (KAPA

Biosystems). The sequencing libraries were clustered on flowcell lanes. After clustering, the

flowcell was loaded on the Illumina NovaSeq instrument according to manufacturer's instructions.

The samples were sequenced using a 2x150 Paired End configuration. Image analysis and base

calling were conducted by the Control Software (NCS). Raw sequence data (.bcl files) generated

from the Illumina instrument was converted into fastq files and de-multiplexed using Illumina's

bcl2fastq (version 2.17) software. One mismatch was allowed for index sequence identification.

Several bioinformatics analyses were performed on the RNA-seq data to identify differentially

expressed genes, differentially used exons, and differentially spliced events between RPE *p53*-/-

control cells and *p53-/-EWSR1* knock-out cells.

For differential gene expression analysis, Kallisto (version 0.48.0) [66] was used to map the raw

RNA-seq reads against the GENCODE human reference gene annotation (Release 44) [67] and

generate raw transcript abundance estimates, and the *tximport* Bioconductor R package [68] was used

to summarize the transcript abundance estimates to produce gene-level estimated counts.

Differentially expressed genes between control and *EWSR1* knock-out cell samples (N = 3) were

determined using both edgeR (version 3.40.2) [69] and DESeq2 (version 1.38.3) [70] Bioconductor

packages. The high concordance between the estimated $\log_2$ foldchange of the two algorithms was

verified.

For both differential exon usage and differential splicing analyses, STAR (version 2.7.10b) [71] was

used to map RNA-seq reads and generate BAM files. Specifically, the 2-pass STAR was used to

map the raw RNA-seq reads against the GRCh38/hg38 human reference genome, with the

GENCODE human reference gene annotation (Release 44) [67]. The first round of the 2-pass STAR

was run to identify splice junctions, and more confidence junctions then were filtered by requiring

them to have at least 5 total mapped reads. The second round of the 2-pass STAR was run using

the filtered junctions to generate BAM files.

For differential exon usage analysis, DEXSeq (version 1.46.0) [72] was used. In brief, a list of unique

exonic regions was initially generated using the Python script *dexseq_prepare_annotation.py*. This

script "collapsed" exonic regions from different transcripts to be used as exon counting bins. Next,

the *dexseq_count.py* Python script was used to count reads for each sample. Finally, differentially

used exons between control and *EWSR1* knock-out cells (N = 3) were determined using DEXSeq.

For the differential splicing analysis, MISO (version 0.5.4) [26] and other tools were used as

discussed below. The exon-centric splicing event annotations were generated using the *rnaseqlib*

Python package (http://github.com/yarden/rnaseqlib). These annotations were derived

considering all transcripts from Ensembl, UCSC RefSeq and NCBI RefSeq gene annotations,

which were obtained from https://hgdownload.soe.ucsc.edu/goldenPath/hg38/database, in files

*knownGene.txt*, *refGene.txt*, and *ncbiRefSeq.txt*. Each annotation event belongs to one of these 5

types: skipped exons (SE), alternative 3′ splice sites (A3SS), alternative 5′ splice sites (A5SS),

mutually exclusive exons (MXE), and retained introns (RI). These annotated events were linked

to representative overlapping exons or introns of the GENCODE human reference gene

annotation (Release 44) using the intersect function of the *bedtools* toolset (version 2.31.0). The

representative exons and introns were prioritized with a high overlapping percentage (95%),

followed by good transcript quality information including whether they were annotated as

canonical and basic, and with better transcript support level. MISO was performed to quantify

the isoforms and compute percent-spliced-in values (PSIs) of each alternative splicing event

from the sorted BAM of each sample. Lower quality events were filtered out as follows. For each

replicate sample pair, the *compare_miso* program was used to compare the results of the control

and the *EWSR1* knock-out samples. The *filter_events* program was then used to filter the

comparison results, by requiring each sample to have at least 35 total isoform-identifying reads.

Finally, *limma* analysis (version 3.54.2) was performed on the PSI values of the filtered events to

determine differential splicing events between control and *EWSR1* knock-out cells (N = 3).

Significant differential alternative splicing events were called if they had *FDR* smaller than 0.05

and absolute delta PSI of at least 0.05. Events that had significant FDR (<0.05) but small

absolute delta PSI (<0.05) were likely to be artifacts and thus were excluded from *limma* results.

Volcano plots were generated using ggplot2.

To have more confidence in the MISO results, we also independently analyzed differential

splicing using rMATS (version v4.2.0) [27]. Since SE events were the most frequent event type,

we focused on analyzing the concordance of SE events between MISO and rMATS. First,

rMATS was run using all sorted BAMs as inputs and the GENCODE human reference gene

annotation (Release 44).  Lower quality events were filtered out by requiring each sample to

have at least 35 total isoform-identifying reads. Then, the shared events that were highly similar

between MISO and rMATS were identified if their target regions overlapped by at least 95%.

Since one MISO event could be linked to multiple slightly different rMATS events, for each

MISO event, a representative rMATS event was chosen that had the largest absolute difference

in inclusion level. Finally, we assessed the concordance between MISO and rMATS by

comparing their inclusion differences (delta PSI values and differences in inclusion levels,

respectively) from these shared events. We observed that a great majority of significant MISO

events had a shared rMATS event. Furthermore, we observed that the inclusion differences of

MISO and rMATS were positively correlated with $R^2$ of 0.32. The scatterplot comparing the

inclusion differences of the two algorithms was generated using ggplot2. Sashimi plots were

generated using the *rmats2sashimiplot* program (https://github.com/Xinglab/rmats2sashimiplot/).

1296

**AlphaFold analysis**

AlphaFold3 and the AlphaFold Server [73] were used to model full-length (POLQ aa 1824-2579)

and truncated (POLQ aa 1824-2390) POLQ polymerases domains in complex with two

complementary 9 bp DNA molecules (5'-CCAATGACA-3' and 5'-TGTCATTGG-3'). Models

were visualized in Pymol V3.0.4 (Schrödinger) using the top-ranking predictions per seed. The

full-length model was compared with a published crystal structure of the POLQ polymerase

domain (PDB 6XBU) [74] to ensure its accuracy.

1304

**Mouse xenograft study**

All animal experiments were conducted in accordance with Institutional Animal Care and Use

Committee-approved protocol (#08-036) at Dana-Farber Cancer Institute. TC32 cells were

lentivirally transduced with luciferase by using a pLenti CMV Puro LUC (w168-1) vector. Two

million TC32 cells expressing luciferase were subcutaneously injected into 7-week-old male and

female NOD.Cg-*Prkdc*scid Il2rgtm1Wjl/SzJ mice. After confirming tumor engraftment, mice were

treated with Peposertib (100 mg/kg) or vehicle control once a day via oral administration together

with Etoposide (8mg/kg) or vehicle control via intraperitoneal injection (IP injection) twice a week

for 2 weeks. Tumors were measured every 2 to 3 days using an electronic caliper, and tumor

volumes were calculated by using the formula L×W×W/2. For immunohistochemistry analysis,

tumors were excised and fixed with formaldehyde. Mice with a tumor of more than 20 mm in

length or width were euthanized.

1317

**Data acquiring and processing of transcriptomic sequencing from patient tumor samples**

Patient tumor sample bulk transcriptomic sequencing data were obtained from St. Jude's Cloud [75].

For expression analysis, we used the mapped expression counts data; for the exon splicing data,

we used the BAM files provided. For the expression counts data, all of the samples were uniformly

processed and mapped to the same reference genome as part of the St. Jude's Cloud resource, and

as such, no correction between samples or cancer types was performed. To evaluate the *POLQ*

expression, we first computed the TPM (transcript-per-million) counts for all samples. In this

computation, we omitted genes annotated as pseudogenes in GENCODE v.31, and used the

median gene length as the gene length for each gene symbol.

1327

**Computation of exon skipping rates in primary patient samples**

To compute the proportions of reads with exons skipped in the sequencing reads data obtained

from St. Jude's Cloud [76], rMATS-turbo, a computational tool to quantify alternative splicing

patterns from RNA sequencing data, was applied. To compute the proportion of reads that each

exon was skipped in, the rMATS-turbo results inclusive of supporting reads from counted from

both junction and exon reads were used. Of note, for each individual exon, any sample that had no

supporting reads for both the inclusive and skipped cases was omitted. All comparisons between

distributions of skipped read proportions were performed using Mann-Whitney U tests.

1336

**Mutational Analysis of Tumors**

1337

Mutational analysis was performed on the whole genome sequencing (WGS) data of Ewing

sarcoma and neuroblastoma tumors obtained from St. Jude Cloud (https://www.stjude.cloud/).

The samples include 23 Ewing sarcoma (ES) and 48 neuroblastoma (NB) tumor and matched-

normal pairs. Somatic variants were determined as follows. First, for each cohort, short variants

were called from the HG38 Genomic Variant Call Format (gVCF) data using the GATK joint

genotyping workflow (version 4.2.4). Second, bcftools (version 1.14) was used to filter out low-

quality variants with these hard filters: QD < 7.5, MQ < 59.9, -g3, and -G10. Only biallelic SNPs

and INDELs were selected. Third, cohort-level singleton variants were selected if there was only

one sample with read depth (DP) between 17 and 300, and variant allele fraction (VAF) of at

least 0.1. And finally, a variant was considered somatic if both the tumor and matched-normal

had SOR <= 3, MQ > 59.9, and DP > 17, and either, tumor VAF >= 0.175 and matched normal

VAF = 0, or tumor VAF <= 0.825 and matched normal VAF = 1.

Profiles of SNV mutations and INDELs were generated using SigProfilerMatrixGenerator

(version 1.2.13; https://github.com/AlexandrovLab/SigProfilerMatrixGenerator). Mutational

signature analysis was performed using SigProfilerAssignment (version 0.0.21;

https://github.com/AlexandrovLab/SigProfilerAssignment). This involved fitting the mutational

profiles to the COSMIC (https://cancer.sanger.ac.uk/signatures/) SNV signatures (SBS version 2)

and INDEL signatures (ID version 3.3).

The MMEJ signature was characterized by the presence of microhomologies at the

boundaries of deletions and the presence of nearby template sequences in insertions, or templated

insertions [77,78]. The presence of microhomologies in deletions was also independently determined

using the method described by Taheri-Ghahfarokhi *et al* [77,78]. This method essentially finds the
longest possible complementary subsequence between one end of the deleted sequence and the
flanking sequence at the other end. In brief, for each deletion, 50-bp flanking sequences around
the deleted sequence were first retrieved from the reference human genome (HG38) using
bedtools (version 2.27.1; https://bedtools.readthedocs.io/). The deletion was then repeatedly
shifted leftward by one base whenever its rightmost base matched the reference base
immediately to its left. Finally, the microhomology sequence was the longest matching
subsequence between the deleted sequence and the right flanking sequence starting from their
leftmost bases. Deletions with length of at least 3bp with GATK ReadPosRankSum of greater
than -5 and with microhomology of at least 2 bp were designated as MMEJ events [77]. For
comparison between ES and NB, tumors with at least 15 deletions and the proportions of MMEJ
deletions among all indels were used. Local templated insertions were characterized by the
presence of subsequences in the inserted sequence that were identical to the flanking sequences
(templated inserts) [15,79,80]. The templated inserts were detected by aligning the inserted sequence
with the 50bp-flanking sequences around the insertion site using the Smith–Waterman local
alignment algorithm [81]. The analysis was restricted to insertions with GATK ReadPosRankSum
of greater than -5 and with length of at least 5bp. The insertions with templates adjacent to the
insertion position were excluded. The remaining insertions with templates covering at least 60%
of the inserted sequence were designated as MMEJ events. For comparison between ES and NB,
tumors with at least 2 insertions, and the proportions of MMEJ templated insertions among all
insertions were used.
1380
**Immunohistochemistry (IHC)**

Primary patient samples for IHC analysis were obtained from the Mayo Clinic (IRB approval #24-

007702). IHC was performed using a Parhelia autostainer. Slides were deparaffinized using Slide

Brite xylene substitute (23401-01) followed by rehydration in ethyl alcohol. Endogenous blocking

was performed using methanol hydrogen peroxide solution. Slides were heated for 30 minutes in

citrate solution for epitope retrieval. Slides were incubated for 30minutes in anti-DNAPKcs

phosphor S2056 antibody at 1:150 followed by washing in wash buffer (Dako S3006) and

inbubation for 30minutes in Mach 3 rabbit HRP (Buocare Medical RP531L). Primary staining was

performed by incubation in DAB (Dako K3468) for 10minutes. Counterstaing was performed

using hematoxylin (Sigma-Aldrich MHS16) and bluing buffer (Dako C5702). Slides were then

dehydrated in ethyl alcohol before coverslipping using Permount mounting mediun (Fisher

Scientific SP15-100). Slides were digitally imaged using a Motic Easyscan Pro 6.

Histological image analysis was performed using QuPath (version 0.5.1). Whole-slide images

(WSIs) were imported into QuPath, and automated color deconvolution was applied to separate

the hematoxylin and DAB (3,3′-diaminobenzidine) staining channels. The mean optica l density

(OD) of the DAB-positive areas was measured to assess staining intensity. For patients with

multiple tissue sections, and the mean OD across all sections for the same patient was used to

represent the patient's staining intensity.

1399

**Statistical analysis**

Data were analyzed and visualized using GraphPad Prism (Version 10.2.2, GraphPad Software,

LLC). For comparisons between two groups, either a two-tailed Student's t-test or a Mann-

Whitney U test was used. For comparisons involving more than two groups, one-way analysis of

variance (ANOVA) was performed. For *in vivo* survival analysis study, a log-rank test was

performed. A p-value of less than 0.05 ($p < 0.05$) was considered statistically significant.

1406

1407

1408

1409

1410

1411
1412
1413
1414

1415

**REFERENCES**

1416

1    Riggi, N., Suva, M. L. & Stamenkovic, I. Ewing's Sarcoma. *N Engl J Med* **384**, 154-164
(2021). https://doi.org/10.1056/NEJMra2028910
2    Grunewald, T. G. P. *et al.* Ewing sarcoma. *Nat Rev Dis Primers* **4**, 5 (2018).
https://doi.org/10.1038/s41572-018-0003-x
3    Gorthi, A. *et al.* EWS-FLI1 increases transcription to cause R-loops and block BRCA1
repair in Ewing sarcoma. *Nature* **555**, 387-391 (2018).
https://doi.org/10.1038/nature25748
4    Choy, E. *et al.* Phase II study of olaparib in patients with refractory Ewing sarcoma
following failure of standard chemotherapy. *BMC Cancer* **14**, 813 (2014).
https://doi.org/10.1186/1471-2407-14-813
5    Boulay, G. *et al.* Cancer-Specific Retargeting of BAF Complexes by a Prion-like
Domain. *Cell* **171**, 163-178 e119 (2017). https://doi.org/10.1016/j.cell.2017.07.036
6    Davis, R. B., Supakar, A., Ranganath, A. K., Moosa, M. M. & Banerjee, P. R.
Heterotypic interactions can drive selective co-condensation of prion-like low-complexity
domains of FET proteins and mammalian SWI/SNF complex. *Nat Commun* **15**, 1168
(2024). https://doi.org/10.1038/s41467-024-44945-5
7    Chong, S. *et al.* Tuning levels of low-complexity domain interactions to modulate
endogenous oncogenic transcription. *Mol Cell* **82**, 2084-2097 e2085 (2022).
https://doi.org/10.1016/j.molcel.2022.04.007
8    Zuo, L. *et al.* Loci-specific phase separation of FET fusion oncoproteins promotes gene
transcription. *Nat Commun* **12**, 1491 (2021). https://doi.org/10.1038/s41467-021-21690-7
9    Chong, S. *et al.* Imaging dynamic and selective low-complexity domain interactions that
control gene transcription. *Science* **361** (2018). https://doi.org/10.1126/science.aar2555
10    Selvanathan, S. P. *et al.* EWS-FLI1 modulated alternative splicing of ARID1A reveals
novel oncogenic function through the BAF complex. *Nucleic Acids Res* **47**, 9619-9636
(2019). https://doi.org/10.1093/nar/gkz699
11    Wang, Z. *et al.* DNA polymerase theta (POLQ) is important for repair of DNA double-
strand breaks caused by fork collapse. *J Biol Chem* **294**, 3909-3919 (2019).
https://doi.org/10.1074/jbc.RA118.005188
12    Goff, J. P. *et al.* Lack of DNA polymerase theta (POLQ) radiosensitizes bone marrow
stromal cells in vitro and increases reticulocyte micronuclei after total-body irradiation.
*Radiat Res* **172**, 165-174 (2009). https://doi.org/10.1667/RR1598.1
13    Mengwasser, K. E. *et al.* Genetic Screens Reveal FEN1 and APEX2 as BRCA2 Synthetic
Lethal Targets. *Mol Cell* **73**, 885-899 e886 (2019).
https://doi.org/10.1016/j.molcel.2018.12.008
14    Adam, S. *et al.* The CIP2A-TOPBP1 axis safeguards chromosome stability and is a
synthetic lethal target for BRCA-mutated cancer. *Nat Cancer* **2**, 1357-1371 (2021).
https://doi.org/10.1038/s43018-021-00266-w
15    Chan, S. H., Yu, A. M. & McVey, M. Dual roles for DNA polymerase theta in alternative
end-joining repair of double-strand breaks in Drosophila. *PLoS Genet* **6**, e1001005
(2010). https://doi.org/10.1371/journal.pgen.1001005
16    Della-Maria, J. *et al.* Human Mre11/human Rad50/Nbs1 and DNA ligase
IIIalpha/XRCC1 protein complexes act together in an alternative nonhomologous end

joining pathway. *J Biol Chem* **286**, 33845-33853 (2011).
https://doi.org/10.1074/jbc.M111.274159

Fleury, H. *et al.* The APE2 nuclease is essential for DNA double-strand break repair by microhomology-mediated end joining. *Mol Cell* **83**, 1429-1445 e1428 (2023). https://doi.org/10.1016/j.molcel.2023.03.017

Brambati, A. *et al.* RHINO directs MMEJ to repair DNA breaks in mitosis. *Science* **381**, 653-660 (2023). https://doi.org/10.1126/science.adh3694

Hussmann, J. A. *et al.* Mapping the genetic landscape of DNA double-strand break repair. *Cell* **184**, 5653-5669 e5625 (2021). https://doi.org/10.1016/j.cell.2021.10.002

Truong, L. N. *et al.* Microhomology-mediated End Joining and Homologous Recombination share the initial end resection step to repair DNA double-strand breaks in mammalian cells. *Proc Natl Acad Sci U S A* **110**, 7720-7725 (2013). https://doi.org/10.1073/pnas.1213431110

Pierce, A. J., Johnson, R. D., Thompson, L. H. & Jasin, M. XRCC3 promotes homology-directed repair of DNA damage in mammalian cells. *Genes Dev* **13**, 2633-2638 (1999). https://doi.org/10.1101/gad.13.20.2633

Zhou, J. *et al.* A first-in-class Polymerase Theta Inhibitor selectively targets Homologous-Recombination-Deficient Tumors. *Nat Cancer* **2**, 598-610 (2021). https://doi.org/10.1038/s43018-021-00203-x

Nabet, B. *et al.* The dTAG system for immediate and target-specific protein degradation. *Nat Chem Biol* **14**, 431-441 (2018). https://doi.org/10.1038/s41589-018-0021-8

Nabet, B. *et al.* Rapid and direct control of target protein levels with VHL-recruiting dTAG molecules. *Nat Commun* **11**, 4687 (2020). https://doi.org/10.1038/s41467-020-18377-w

Johnson, C. N. *et al.* Insights into Molecular Diversity within the FET Family: Unraveling Phase Separation of the N-Terminal Low Complexity Domain from RNA-Binding Protein EWS. *bioRxiv* (2023). https://doi.org/10.1101/2023.10.27.564484

Katz, Y., Wang, E. T., Airoldi, E. M. & Burge, C. B. Analysis and design of RNA sequencing experiments for identifying isoform regulation. *Nat Methods* **7**, 1009-1015 (2010). https://doi.org/10.1038/nmeth.1528

Shen, S. *et al.* rMATS: robust and flexible detection of differential alternative splicing from replicate RNA-Seq data. *Proc Natl Acad Sci U S A* **111**, E5593-5601 (2014). https://doi.org/10.1073/pnas.1419161111

Harada, N. *et al.* The splicing factor CCAR1 regulates the Fanconi anemia/BRCA pathway. *Mol Cell* **84**, 2618-2633 e2610 (2024). https://doi.org/10.1016/j.molcel.2024.06.031

Nicholson-Shaw, A. L., Kofman, E. R., Yeo, G. W. & Pasquinelli, A. E. Nuclear and cytoplasmic poly(A) binding proteins (PABPs) favor distinct transcripts and isoforms. *Nucleic Acids Res* **50**, 4685-4702 (2022). https://doi.org/10.1093/nar/gkac263

Lawrence, M. S. *et al.* Mutational heterogeneity in cancer and the search for new cancer-associated genes. *Nature* **499**, 214-218 (2013). https://doi.org/10.1038/nature12213

Thatikonda, V. *et al.* Comprehensive analysis of mutational signatures reveals distinct patterns and molecular processes across 27 pediatric cancers. *Nat Cancer* **4**, 276-289 (2023). https://doi.org/10.1038/s43018-022-00509-4

| 1504 | 32 | Schimmel, J., Kool, H., van Schendel, R. & Tijsterman, M. Mutational signatures of non-homologous and polymerase theta-mediated end-joining in embryonic stem cells. *EMBO J* **36**, 3634-3649 (2017). https://doi.org/10.15252/embj.201796948 |
| 1507 | 33 | Johnson, C. N. *et al.* Insights into Molecular Diversity within the FUS/EWS/TAF15 Protein Family: Unraveling Phase Separation of the N-Terminal Low-Complexity Domain from RNA-Binding Protein EWS. *J Am Chem Soc* **146**, 8071-8085 (2024). https://doi.org/10.1021/jacs.3c12034 |
| 1511 | 34 | Ebersberger, S. *et al.* FUBP1 is a general splicing factor facilitating 3' splice site recognition and splicing of long introns. *Mol Cell* **83**, 2653-2672 e2615 (2023). https://doi.org/10.1016/j.molcel.2023.07.002 |
| 1514 | 35 | Graham, G. T. *et al.* Comprehensive profiling of mRNA splicing indicates that GC content signals altered cassette exon inclusion in Ewing sarcoma. *NAR Cancer* **4**, zcab052 (2022). https://doi.org/10.1093/narcan/zcab052 |
| 1517 | 36 | Selvanathan, S. P. *et al.* Oncogenic fusion protein EWS-FLI1 is a network hub that regulates alternative splicing. *Proc Natl Acad Sci U S A* **112**, E1307-1316 (2015). https://doi.org/10.1073/pnas.1500536112 |
| 1520 | 37 | Zatreanu, D. *et al.* Poltheta inhibitors elicit BRCA-gene synthetic lethality and target PARP inhibitor resistance. *Nat Commun* **12**, 3636 (2021). https://doi.org/10.1038/s41467-021-23463-8 |
| 1523 | 38 | Llorens-Agost, M. *et al.* POLtheta-mediated end joining is restricted by RAD52 and BRCA2 until the onset of mitosis. *Nat Cell Biol* **23**, 1095-1104 (2021). https://doi.org/10.1038/s41556-021-00764-0 |
| 1526 | 39 | Gelot, C. *et al.* Poltheta is phosphorylated by PLK1 to repair double-strand breaks in mitosis. *Nature* **621**, 415-422 (2023). https://doi.org/10.1038/s41586-023-06506-6 |
| 1528 | 40 | Kumar, R. J. *et al.* Dual inhibition of DNA-PK and DNA polymerase theta overcomes radiation resistance induced by p53 deficiency. *NAR Cancer* **2**, zcaa038 (2020). https://doi.org/10.1093/narcan/zcaa038 |
| 1531 | 41 | Patterson-Fortin, J. *et al.* Targeting DNA Repair with Combined Inhibition of NHEJ and MMEJ Induces Synthetic Lethality in TP53-Mutant Cancers. *Cancer Res* **82**, 3815-3829 (2022). https://doi.org/10.1158/0008-5472.CAN-22-1124 |
| 1534 | 42 | Collins, V. J. *et al.* Enhancing standard of care chemotherapy efficacy using DNA-dependent protein kinase (DNA-PK) inhibition in preclinical models of Ewing sarcoma. *Mol Cancer Ther* (2024). https://doi.org/10.1158/1535-7163.MCT-23-0641 |
| 1537 | 43 | Dubbury, S. J., Boutz, P. L. & Sharp, P. A. CDK12 regulates DNA repair genes by suppressing intronic polyadenylation. *Nature* **564**, 141-145 (2018). https://doi.org/10.1038/s41586-018-0758-y |
| 1540 | 44 | Iniguez, A. B. *et al.* EWS/FLI Confers Tumor Cell Synthetic Lethality to CDK12 Inhibition in Ewing Sarcoma. *Cancer Cell* **33**, 202-216 e206 (2018). https://doi.org/10.1016/j.ccell.2017.12.009 |
| 1543 | 45 | Gounder, M. M. *et al.* Clinical genomic profiling in the management of patients with soft tissue and bone sarcoma. *Nat Commun* **13**, 3406 (2022). https://doi.org/10.1038/s41467-022-30496-0 |
| 1546 | 46 | Tirode, F. *et al.* Genomic landscape of Ewing sarcoma defines an aggressive subtype with co-association of STAG2 and TP53 mutations. *Cancer Discov* **4**, 1342-1353 (2014). https://doi.org/10.1158/2159-8290.CD-14-0622 |

47    Dutertre, M. *et al.* Cotranscriptional exon skipping in the genotoxic stress response. *Nat Struct Mol Biol* **17**, 1358-1366 (2010). https://doi.org/10.1038/nsmb.1912

48    Paronetto, M. P., Minana, B. & Valcarcel, J. The Ewing sarcoma protein regulates DNA damage-induced alternative splicing. *Mol Cell* **43**, 353-368 (2011). https://doi.org/10.1016/j.molcel.2011.05.035

49    Verdile, V. *et al.* EWS splicing regulation contributes to balancing Foxp1 isoforms required for neuronal differentiation. *Nucleic Acids Res* **50**, 3362-3378 (2022). https://doi.org/10.1093/nar/gkac154

50    Maurer, L. M. *et al.* BRCA1-associated RING domain-1 (BARD1) loss and GBP1 expression enhance sensitivity to DNA damage in Ewing sarcoma. *Cancer Res Commun* **2**, 220-232 (2022). https://doi.org/10.1158/2767-9764.crc-21-0047

51    Menon, S. *et al.* FET fusion oncoproteins disrupt physiologic DNA repair networks and induce ATR synthetic lethality in cancer. *Res Sq* (2023). https://doi.org/10.21203/rs.3.rs-2869150/v1

52    Ceccaldi, R. *et al.* Homologous-recombination-deficient tumours are dependent on Poltheta-mediated repair. *Nature* **518**, 258-262 (2015). https://doi.org/10.1038/nature14184

53    Mateos-Gomez, P. A. *et al.* Mammalian polymerase theta promotes alternative NHEJ and suppresses recombination. *Nature* **518**, 254-257 (2015). https://doi.org/10.1038/nature14157

54    Bhat, P. *et al.* Genome organization around nuclear speckles drives mRNA splicing efficiency. *Nature* **629**, 1165-1173 (2024). https://doi.org/10.1038/s41586-024-07429-6

55    Wang, J., Schultz, P. G. & Johnson, K. A. Mechanistic studies of a small-molecule modulator of SMN2 splicing. *Proc Natl Acad Sci U S A* **115**, E4604-E4612 (2018). https://doi.org/10.1073/pnas.1800260115

56    Duncan, R., Collins, I., Tomonaga, T., Zhang, T. & Levens, D. A unique transactivation sequence motif is found in the carboxyl-terminal domain of the single-strand-binding protein FBP. *Mol Cell Biol* **16**, 2274-2282 (1996). https://doi.org/10.1128/MCB.16.5.2274

57    Farag, M. *et al.* Condensates formed by prion-like low-complexity domains have small-world network structures and interfaces defined by expanded conformations. *Nat Commun* **13**, 7722 (2022). https://doi.org/10.1038/s41467-022-35370-7

58    Wang, J. *et al.* A Molecular Grammar Governing the Driving Forces for Phase Separation of Prion-like RNA Binding Proteins. *Cell* **174**, 688-699 e616 (2018). https://doi.org/10.1016/j.cell.2018.06.006

59    Seong, B. K. A. *et al.* TRIM8 modulates the EWS/FLI oncoprotein to promote survival in Ewing sarcoma. *Cancer Cell* **39**, 1262-1278 e1267 (2021). https://doi.org/10.1016/j.ccell.2021.07.003

60    Chaudhary, U. B. & Haldas, J. R. Long-term complications of chemotherapy for germ cell tumours. *Drugs* **63**, 1565-1577 (2003). https://doi.org/10.2165/00003495-200363150-00004

61    Lim, K. S. *et al.* USP1 Is Required for Replication Fork Protection in BRCA1-Deficient Tumors. *Mol Cell* **72**, 925-941 e924 (2018). https://doi.org/10.1016/j.molcel.2018.10.045

62    Hayer, A. *et al.* Engulfed cadherin fingers are polarized junctional structures between collectively migrating endothelial cells. *Nat Cell Biol* **18**, 1311-1323 (2016). https://doi.org/10.1038/ncb3438

1595 63 Sanjana, N. E., Shalem, O. & Zhang, F. Improved vectors and genome-wide libraries for
1596   CRISPR screening. *Nat Methods* **11**, 783-784 (2014). https://doi.org/10.1038/nmeth.3047
1597 64 Tyanova, S. *et al.* The Perseus computational platform for comprehensive analysis of
1598   (prote)omics data. *Nat Methods* **13**, 731-740 (2016). https://doi.org/10.1038/nmeth.3901
1599 65 Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing
1600   and microarray studies. *Nucleic Acids Res* **43**, e47 (2015).
1601   https://doi.org/10.1093/nar/gkv007
1602 66 Bray, N. L., Pimentel, H., Melsted, P. & Pachter, L. Near-optimal probabilistic RNA-seq
1603   quantification. *Nat Biotechnol* **34**, 525-527 (2016). https://doi.org/10.1038/nbt.3519
1604 67 Frankish, A. *et al.* Gencode 2021. *Nucleic Acids Res* **49**, D916-D923 (2021).
1605   https://doi.org/10.1093/nar/gkaa1087
1606 68 Soneson, C., Love, M. I. & Robinson, M. D. Differential analyses for RNA-seq:
1607   transcript-level estimates improve gene-level inferences. *F1000Res* **4**, 1521 (2015).
1608   https://doi.org/10.12688/f1000research.7563.2
1609 69 Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for
1610   differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139-
1611   140 (2010). https://doi.org/10.1093/bioinformatics/btp616
1612 70 Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion
1613   for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).
1614   https://doi.org/10.1186/s13059-014-0550-8
1615 71 Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21
1616   (2013). https://doi.org/10.1093/bioinformatics/bts635
1617 72 Anders, S., Reyes, A. & Huber, W. Detecting differential usage of exons from RNA-seq
1618   data. *Genome Res* **22**, 2008-2017 (2012). https://doi.org/10.1101/gr.133744.111
1619 73 Abramson, J. *et al.* Accurate structure prediction of biomolecular interactions with
1620   AlphaFold 3. *Nature* **630**, 493-500 (2024). https://doi.org/10.1038/s41586-024-07487-w
1621 74 Chandramouly, G. *et al.* Poltheta reverse transcribes RNA and promotes RNA-templated
1622   DNA repair. *Sci Adv* **7** (2021). https://doi.org/10.1126/sciadv.abf1771
1623 75 McLeod, C. *et al.* St. Jude Cloud: A Pediatric Cancer Genomic Data-Sharing Ecosystem.
1624   *Cancer Discov* **11**, 1082-1099 (2021). https://doi.org/10.1158/2159-8290.CD-20-1230
1625 76 Wang, Y. *et al.* rMATS-turbo: an efficient and flexible computational tool for alternative
1626   splicing analysis of large-scale RNA-seq data. *Nat Protoc* **19**, 1083-1104 (2024).
1627   https://doi.org/10.1038/s41596-023-00944-2
1628 77 Tobalina, L., Armenia, J., Irving, E., O'Connor, M. J. & Forment, J. V. A meta-analysis
1629   of reversion mutations in BRCA genes identifies signatures of DNA end-joining repair
1630   mechanisms driving therapy resistance. *Ann Oncol* **32**, 103-112 (2021).
1631   https://doi.org/10.1016/j.annonc.2020.10.470
1632 78 Taheri-Ghahfarokhi, A. *et al.* Decoding non-random mutational signatures at Cas9
1633   targeted sites. *Nucleic Acids Res* **46**, 8417-8434 (2018).
1634   https://doi.org/10.1093/nar/gky653
1635 79 van Schendel, R., Roerink, S. F., Portegijs, V., van den Heuvel, S. & Tijsterman, M.
1636   Polymerase Theta is a key driver of genome evolution and of CRISPR/Cas9-mediated
1637   mutagenesis. *Nat Commun* **6**, 7394 (2015). https://doi.org/10.1038/ncomms8394
1638 80 Yu, A. M. & McVey, M. Synthesis-dependent microhomology-mediated end joining
1639   accounts for multiple types of repair junctions. *Nucleic Acids Res* **38**, 5706-5717 (2010).
1640   https://doi.org/10.1093/nar/gkq379

1641   81    Smith, T. F. & Waterman, M. S. Identification of common molecular subsequences. *J*
1642         *Mol Biol* **147**, 195-197 (1981). https://doi.org/10.1016/0022-2836(81)90087-5
1643

# Figure 1

# Figure 2

# Figure 3

# Figure 4

# Figure 5