Virus-Positive Merkel Cell Carcinoma Model Supports Unified Cancer Origin

Wendy Yang*, Sara Contente, Sarah Rahman
Department of Pathology, Uniformed Services University of Health Sciences, 4301 Jones Bridge Road, Bethesda, MD 20814
* Corresponding author, email: wendy.yang@usuhs.edu

Summary:

The Germ Cell Theory, rooted in developmental biology[1], has significantly advanced both the understanding and curative treatment of rare germ cell cancers (GCCs)[2]. In contrast, somatic cancer research - long dominated by the Somatic Mutation Theory - has stagnated due to fragmented, mutation-centric approaches that lack a unifying conceptual framework and have yielded only limited clinical progress[3]. GCC research identifies non-somatic human primordial germ cells (hPGCs) as the cells of origin for GCCs[1,4]. In somatic cancers, although the cancer stem cell theory is gaining acceptance, cancer stem cells are assumed to be of somatic origin, with their exact identity undefined[5]. Accumulating experimental evidence and clinical observations challenge the traditional separation between GCCs and somatic cancers, including malignant somatic transformation (MST) - the emergence of somatic cancer phenotypes from GCCs, often without new mutations[2, 6-8] - suggesting that hPGCs may also initiate somatic cancers. However, no experimental model of MST has been established[9]. Merkel cell polyomavirus-positive Merkel cell carcinoma (MCC), a highly aggressive somatic cancer with paradoxically stable genomes resembling those of GCCs[10-12], offers a unique model to test germ cell theory-based somatic oncogenesis. Here we present an MST mouse model linking hPGC origin to somatic cancer by inducing virus-positive MCC-like tumors in vivo from virus-transfected hPGC-like cells or human iPSCs competent for hPGC specification with an obligatory late-hPGC state preceding MST. This genetically simple, molecularly tractable model provides a novel platform to dissect VP-MCC pathogenesis and broadly advocates a developmental biology framework for somatic cancer research beyond mutation-centric and soma-centric paradigms.

Main

Cancer has traditionally been classified into two primary groups based on presumed cellular origin: germ cell cancers (GCCs), arising from reproductive germline cells - also referred to as Type II germ cell tumors (GCTs) - which constitute less than 1% of malignancies[1]; and somatic cancers, originating from non-germline somatic cells, representing the vast majority of malignancies. Despite this traditional separation, all cancers share phenotypic hallmarks, many of which recapitulate early embryonic/germ cell development, suggesting developmental arrest or "blocked ontogeny"[13,14]. This principle, encapsulated as "oncology recapitulates ontogeny," has significantly influenced GCC research and clinical practice yet remains largely overlooked in somatic cancer studies[2,3], despite its clinical relevance to grading and prognosis[15].

The Germ Cell Theory, grounded in developmental biology, has yielded transformative insights into GCT biology and curative treatment strategies[1,2]. GCCs are genetically simple and epigenetically distinct[1,2], originating from human primordial germ cells (hPGCs)[1,4] - founder cells of the germline specified during weeks 2-3 of development[16]. hPGCs exhibit latent pluripotency and intrinsic cancer stem cell-like properties, predisposing them to malignant transformation with minimal genetic alteration[1]. In GCCs, transformed hPGCs and their totipotent derivatives - embryonal carcinoma (EC) cells thought to arise via parthenogenesis in vivo - constitute the cancer stem cell compartments of seminomas and non-seminomatous germ cell cancers (NSGCCs), respectively[1]. Seminomas are composed entirely of transformed hPGCs, while NSGCCs such as ECs or malignant teratomas consist solely of EC cells or arise through multilineage somatic differentiation from ECs respectively[1].

Mounting biological and clinical evidence increasingly challenges the traditional separation between germ cell cancers (GCCs) and somatic cancers, suggesting a shared developmental origin. Notably, approximately 3–6% of germ cell tumors (GCTs) undergo malignant somatic transformation (MST) into somatic cancer phenotypes, often without additional mutations[2,6-8]. GCTs can also manifest as extragonadal GCTs (EGGCTs), demonstrating - like somatic cancers - the capacity to arise at virtually any anatomical site, albeit with a predilection for midline locations[17]. Many somatic cancers aberrantly express cancer-testis antigens normally restricted to germ cells[18,19], and acquisition of a primordial germ cell (PGC)-like state has been shown to be essential for metastasis[20]. Furthermore, global DNA

hypomethylation is a ubiquitous epigenetic hallmark of all cancers, observed in both GCCs and somatic cancers alike[1,21]. Despite these converging features, somatic cancer research remains dominated by the fragmented, mutation-centric Somatic Mutation Theory (SMT), which lacks a unifying biological framework and has yielded limited improvements in patient outcomes[3]. Although the cancer stem cell theory has more recently gained recognition in somatic cancer research, it fundamentally differs from the germ cell theory in its soma-centric view by positing somatic adult stem cells - whose precise identity remains undefined - as the origin of somatic cancers[5]. Given the substantial biological and clinical parallels between GCCs and somatic cancers, it is reasonable to propose that hPGCs - or a PGC-like state - may represent a shared stem cell origin for both. However, no experimental model has yet directly linked hPGCs to somatic cancer via MST[9], despite the longstanding use of teratoma assays to study GCT development and tumorigenesis[22]. These assays, which generate teratomas (a form of non-seminomatous GCT) in immunocompromised mice following injection of human pluripotent stem cells (hPSCs), remain the gold standard for confirming pluripotency, due to the intrinsic coupling of pluripotency and teratoma formation[23]. While MST most commonly arises in clinical teratomas, it has not been recapitulated in these models - likely due to the limited duration of in vivo assays relative to the prolonged latency observed in human disease[9].

Virus-positive Merkel cell carcinoma (VP-MCC), an aggressive high grade neuroendocrine carcinoma (HGNEC) of the skin, comprises ~80% of MCC cases and is defined by the presence of randomly but clonally integrated, C-terminus truncated Merkel cell polyomavirus (MCPyV) genomes (~5 kb)[10-12]. It presents two key features that seem to make it ideally suited for developing an MST model. First, VP-MCC exhibits extreme genomic stability, in stark contrast to virus-negative MCC (VN-MCC), which comprises the remaining ~20% and is among the most highly mutated human cancers. Similarly, other virus-associated cancers, such as HPV- and EBV-positive tumors, typically show higher mutational burdens than their VN counterparts[24]. This exceptional genomic stability mirrors that of germ cell tumors (GCTs) and would enable the development of a genetically simple, experimentally tractable MST model. Although MCPyV is ubiquitous in normal skin flora, VP-MCC is very rare, and MCPyV transduction of cutaneous lineages has failed to reproduce its molecular and histologic features - underscoring the paramount importance of cellular context, a principle also similar to GCT biology[11,12,25]. Second, the small cell neuroendocrine carcinoma (SCNC) phenotype of MCC represents a poorly differentiated, aggressive end-stage state to which diverse somatic cancers converge upon progression[26]. SCNCs, including MCC, also exhibit stem-like properties and multilineage differentiation potential[27], suggesting that VP-MCC may be composed predominantly of somatic cancer stem cells, functionally analogous to seminomas or embryonal carcinomas (ECs) in GCTs. Together, these features support the potential of developing a VP-MCC-based MST model capable of capturing transformation events rapidly with immediate somatic differentiation arrests - within the temporal limits of mouse xenograft assays - where MST has otherwise not been observed[9]. To directly test the Germ Cell Theory in somatic cancer, we proceeded to establish a mouse model based on the hypothesis that VP-MCC arises through MCPyV-driven MST of extragonadal hPGCs residing in the adult cutaneous niche. Multiple lines of evidence support our hypothesis. MCPyV tropism toward human primordial germ cells (hPGCs) is indicated by the relatively high level of MCPyV DNA detected in seminoma[28], a GCC composed of transformed hPGCs[1]. The skin microenvironment is enriched in stem cell factor (SCF) and stromal cell-derived factor 1 (SDF-1), key regulators of hPGC survival and chemotaxis[29,30]. Furthermore, the male predominance and head and neck predilection of MCC reflect Y chromosome-mediated hPGC viability and their characteristic midline migratory trajectory[1,17].

To investigate the proposed MCPyV-driven VP-MCC tumorigenic pathway, we designed a teratoma assay-like experiment aiming to prove MST from MCPyV transfected hPGCs to VP-MCC like tumors (VMLTs) in the adult mouse cutaneous niche. We transduced a truncated MCPyV genome (L82) via lentiviral vector into in vitro surrogates of hPGCs and additional two relevant hPSC types - human embryonic germ cells (hEGCs) and human embryonic stem cells (hESCs). Based on prior studies on hPSC-induced mouse teratoma xenografts, we estimated that three mice per condition would provide >99% success rate of forming mouse xenograft tumors[31]. Eighteen adult NSG mice were divided into three groups (n = 6 per group). The **1st group** used **hPGC-like cells** (**hPGCLCs**) that are derived from hPSCs and recapitulate early-hPGC[32,33]. Both normal hPGCs and hPGCLCs exhibit latent pluripotency - retaining germline unipotency in vivo without chimera[34] or teratoma-forming capacity[23,35] - but can undergo parthenogenetic reversion to pluripotent hEGCs or hEGC-like cells (hEGCLCs) in vitro[36,37]. Due

to limited culture yield, low-number (~3×10^5) cells of either hPGCLC_A4 (fig. S1A, provided by the Shioda lab[36]) or MCPyV-transduced hPGCLC_A4_L82 (fig. S1B) were injected into the right flanks of three mice per condition. The **2nd group** used **Human induced pluripotent stem cells** (**hiPSCs**) that are derived from reprogramming of somatic cells and resemble hESCs[38]. Based on our hypothesis, an indirect route from hPSCs to VP-MCC-like tumors (VMLTs) can theoretically be established through hPGCLC specification from competent hPSCs[32]. Totipotent naïve-hESCs of the inner cell mass (ICM) at the early pre-implantation blastocyst stage and pluripotent formative-hESCs at the peri-implantation blastocyst stage are competent for hPGC induction, while more developmentally advanced pluripotent primed-hESCs of the late post-implantation blastocyst stage are not[16,32]. Only a small subset of conventionally cultured hESCs/hiPSCs maintain a naïve hESC state[39] while the vast majority remain at the primed hESC state[40], reflecting developmental plasticity between totipotent and primed hESC states[1,40]. Given that only a small subset of hiPSCs is capable of hPGCLC differentiation[32,39,40], we tested dose-dependent tumorigenicity by injecting hiPSC_A4_L82 (fig. S1C) at conventional-number (1×10^6) and high-number (2×10^7) cell doses into bilateral flanks of three mice per condition. The **3rd group** used **hEGC-like cells** (**hEGCLCs**) as the surrogate for hEGCs[36,37], which are the in vitro parthenogenic derivatives of unipotent hPGCLCs and hPGCs respectively. Based on normal and teratoma developmental trajectories[1] - and given that VP-MCC is a somatic cancer - we considered the possibility that MST may require parthenogenetic reversion of a MCPyV-transformed hPGC to a pluripotent or totipotent intermediate[1] before somatic neuroendocrine lineage differentiation. This would position hEGCs as a more direct candidate cell of origin than hPGCs, supporting the inclusion of hEGCLCs to test for direct conversion to VMLTs in vivo. It also remained unclear whether MCPyV transfected hEGCLC possess equivalent developmental potency to non-parthenogenetic hPSC counterparts such as MCPyV transfected hiPSCs, particularly in achieving a naïve hESC state to support hPGCLC induction and thus an indirect MST route to VMLTs. To evaluate both possibilities, EGCLC_A4_L82 was injected at both cell doses (1×10^6 and 2×10^7) into bilateral flanks of three mice per condition. For both the 2nd group and the 3rd group, MCPyV-negative controls, hiPSC_A4 (provided by the Shioda lab[36]) or hEGCLC_A4, were not included as it has been well established that pluripotency alone is coupled to teratoma formation without MST[9] (Table S1).

Results

Tumors comprised of both VMLT and teratoma components emerged exclusively from low-number (3×10^5) hPGCLCs_L82 and from high-number (2×10^7) hiPSC_L82 injections. Only teratoma component developed in conventional-number (1×10^6) hiPSC_L82 mice, as well as hECLC_L82 mice regardless of cell numbers (both 1×10^6 and 2×10^7). No tumor formation was observed in MCPyV-negative hPGCLC control mice, consistent with their known "unipotency" in vivo from prior studies[23, 35]. One of the three hPGCLC_A4_L82 mice (#86.1) failed to form any tumor, probably due to a technical failure of cell injection as the mouse was noted to be particularly challenging to apply physical restraint during injection. Bilateral injections of hiPSC_A4_L82 line or hEGCLC_A4_L82 line in the same mouse resulted bilateral mouse xenograft tumors of essentially the same gross and histologic characteristics (Table 1 and Table S1).

Group	1st Group		2nd Group		3rd Group	
Condition	hPGCLC_A4	**hPGCLC_A4_L82**	**hiPSC_A4_L82**		**hEGCLC_A4_L82**	
Cell Number Injected	*3×10^5*	*3×10^5*	*1×10^6*	*2×10^7*	*1×10^6*	*2×10^7*
Number of Mice with VMLT(+) Tumor(s) [VMLT + Teratoma]	0	**2**	0	**3**	0	0
Number of Mice with VMLT(-) Tumor(s) [Teratoma Only]	0	0	**3**	0	**3**	**3**
Number of Mice with NO Tumor	3	1	0	0	0	0
Total Number of Mice per Condition	3	3	3	3	3	3

Table 1. Summary of mouse tumor formation from subcutaneously injected human primeval stem cells.

There was no gross or histological evidence of distant metastasis from any xenograft tumors with VMLTs. This finding is consistent with the known much less aggressive phenotype with only rare distant metastasis from VP-MCC tumor cell line-derived mouse tumors in prior studies[41]. The reason for the

marked difference of VP-MCC tumors' aggressiveness in human versus mouse is unknown and may reflect species differences.

MST to VMLTs confirmed in VMLT(+) tumors
Morphology confirmation of MST to VMLTs
Grossly, VMLT(+) tumors, which were composed of both VMLT and teratoma components, were markedly different from VMLT(-) tumors, which were composed only of a teratoma component. All VMLT(+) tumors were solid and homogeneous with pale and friable texture grossly (Figure 1), while teratoma-only tumors were cystic and mucinous with heterogeneous discoloration (Figure S2). The finding mimicked the typical gross difference between clinical teratomas with MST and pure teratomas[42]. The WHO criteria for MST in human GCTs require a minimum 5 mm expansion of pure somatic type of malignancy component[43]. Our most salient tumor examples of MST to VMLT were T#85.4, consisting of two discrete tumor nodules with the larger nodule (T#85.4.N1 ~20 mm) composed of pure VMLT tissue (Figure S5A), and T#84.5R ~20 mm, composed almost entirely of VMLT (Figure S6A-B). The rest of the VMLTs also occupied substantial portions of tumors in sizable continuous sheets (~20% - ~80% of tumor masses) (Figure S7-8, Figure S6C, Figure S9A, Figure S20A).

Histologically, VMLT(+) tumors exhibited a distinct overall spatial pattern: primitive VMLT components localized superficially and peripherally, surrounding more differentiated teratoma components located deeper and centrally (Figure S6C, S7, S9A, S10A, S18). This organization contrasts with the typical "differentiation gradient" of hPSC-derived mouse xenograft pure teratomas like VMLT(-) tumors, which display a less differentiated core and more mature peripheral regions[23]. Notably, the teratoma components within VMLT(+) tumors did retain the expected maturation gradient of xenograft pure teratomas (Figure S18A–B). The opposing spatial arrangements of the overall VMLT(+) tumors versus their teratoma components implicate a distinct VMLT tumorigenic pathway requiring proximity to the cutaneous microenvironment.

VP-MCC like histopathology of VMLTs
Histologic sections and cytology smears of all VMLTs showed sheets of immature small blue cells with either the typical MCC small round cell cytomorphology in the two hPGCLC_A4_L82 derived tumors T#85.3 or T#85.4 (Figure S5E, Figure S8E, Figure S11), or the atypical MCC large cell variant with admixed larger cells exhibiting prominent nucleoli and more abundant cytoplasm in all six high-number hiPSC_A4_L82 derived tumors, T#84.5L&R, T#85.1L&R, T#85.2L&R (Figure S6F, Figure S9D, Figure S10E, Figure S11). All VMLTs exhibited prominent histological characteristics of malignancy (Figure S12), including extracapsular invasion into dermis, with destruction of dermal striated muscle layer and lymphovascular invasion, with VMLT cells within lymphatic spaces (Figure S7, Figure S8A-D, Figure S6E, Figure S9B-C, Figure S10B-D).

VP-MCC like molecular profile of VMLTs
Molecular characterization by RT-qPCR (Figure 2, Table S2) and/or immunohistochemical (IHC) studies (Figure 3, Figure S13, Figure S14-16, Table S3) of VMLTs showed highly similar molecular profiles to each other and to the six VP-MCC cell lines (CVG-1, MKL-1, MKL-2, MS-1, PeTa and WaGa)[44-50]. They were less similar to that of the two virus-negative variant MCC (VN-vMCC) cell lines (MCC-13 and UISO)[49,51,52] that do not express the key neuroendocrine lineage gene INSM1. The VMLTs showed mRNA and/or protein expression in neuroendocrine lineage genes, viral LT antigen, core pluripotency gene and PGC similar to that of VP-MCCs with few exceptions, and a marked departure from that of the corresponding MCPyV transfected primeval stem cells, hPGCLC_A4_L82 or hiPSC_A4_L82 (Figure 2-3, Figure S13, Figure S14-16, Table S2-S3). The high Oct4 expression of all high-number hiPSC_A4_L82-derived VMLTs, in contrast to very low or absent expression in VP-MCC lines (Figure S17), raised concerns about a possible EC component in these VMLTs. However, it was ruled out by PLAP(-) and CD30(-) IHC stains on these VMLTs (Figure S14-16, Table S3). Significantly lower ATOH1 mRNA levels were noted in all VMLTs than VP-MCC cell lines, which might be due to the incipient tumor status of VMLTs versus the advanced cancer stage of these VP-MCC cell lines[53].

Global DNA methylation reset before MST
Global DNA methylation (GDM) is dynamic during development, but may serve as a relatively stable epigenetic marker that retains lineage memory. Cancer cells may maintain DNA methylation signatures

reflective of their cell of origin despite reprogramming. In normal human development, only two waves of global DNA demethylation occur, each producing a transient hypomethylation nadir during early embryogenesis. The first nadir occurs in totipotent naïve hESC/ICM stage (~40% median GDM); the second nadir - and lowest - in late hPGC stage (~6% median GDM)[54,55] (Figure S19). The second nadir's extremely low GDM with near complete GDM erasure is unique to late hPGCs while the first nadir's moderate DNA hypomethylation characterizes the totipotent naïve hESC/ICM state. In vitro models mirror these developmental states with well-characterized GDM levels: hiPSCs and hEGCLCs (primed hESC-like) show high GDM (~70%), hPGCLCs (early hPGC-like) show intermediate GDM (~50%)[36] (Figure S19). GCCs typically reflect this epigenetic hierarchy with few exceptions: seminomas usually match the extremely low GDM of late-hPGCs, ECs usually show moderate global DNA hypomethylation of naïve hESCs, and malignant teratomas usually exhibit higher GDM due to somatic differentiation (Figure 5). All GCCs thus retain variable degrees of global DNA hypomethylation consistent with their late-hPGC origin[1]. Somatic cancers are also known to exhibit global DNA hypomethylation, positively correlating with degree of malignancy and cancer progression[21]. SCNCs, including MCC, share marked global DNA hypomethylation[56,57] with seminomas and ECs - the stem-like compartments of GCCs - as well as stemness and developmental plasticity. To investigate further, we compared GDM in VMLTs, MCC lines, and the only validated seminoma cell line TCam-2[58]. One important caveat is that DNA methylation is highly sensitive to the microenvironment, and tumor cells maintained in vitro often display artificially elevated GDM[59]. For instance, the only validated seminoma cell line TCam-2 - despite its composition of transformed late-hPGCs known to exhibit extremely low GDM (~6% median)[54,55] - shows a moderate GDM level[60] comparable to the cell lines of EC mirroring naïve hESC/ICM state[1] (GDM ~40%)[54,55], rather than the extreme global hypomethylation observed in primary seminoma tissues[60], likely reflecting culture-induced epigenetic drift[59].

GDM by Flow cytometry and IHC validated
Semi-quantitative flow cytometry and immunohistochemical (IHC) study to measure global 5-methylcytosine (5mC) levels for assessing GDM were validated using reference cell lines with well-characterized GDM profiles. The hiPSC_A4_L82 line, modeling the primed hESC state known to exhibit high somatic-type GDM (~70%)[36], showed a very high 5mC signal (MFI ~113K) by flow cytometry. The seminoma cell line TCam-2, which reflects a moderately hypomethylated state similar to ECs or naïve hESCs (GDM ~40%)[54,55], displayed a much lower 5mC signal (MFI ~7K) by flow cytometry with its isotype negative control yielding a minimal signal (MFI ~3K) and serving as the background reference for absence of GDM (Figure 4A; Table S4). IHC stains of hiPSC_A4_L82 line, TCam2 line and its isotype negative control showed similar trend (Figure 4B1-2,4B6). Positive IHC stains of mouse somatic skin tissue and testis tissue served as positive controls as well (Figure 4B3-4).

Requisite Late-hPGC state before MST to VMLT
VMLTs exhibited unique late-hPGC GDM signature
Flow cytometry analysis of VMLT tumor cells revealed extremely low global 5mC levels (MFI: ~2K–~3.5K) (Figure 4A, Table S4), approaching the isotype negative controls (MFI: ~2K-~3K). Global 5mC IHC studies of VMLT(+) tumors T#85.4 and T#85.2R showed near negative staining for the VMLT components (Figure 4B7, 4B9), and positive staining for the teratoma components (Figure 4B8, 4B10) that is similar to pure teratoma from VMLT(-) tumor T#83.5L (Figure 4B5). These results thus indicated VMLTs exhibited unique late-hPGC GDM signature with near-complete GDM erasure, which suggests a model in which MST occurs directly from the late-hPGC state across the Weismann Barrier (WB), bypassing any intermediary stage marked by moderate to high GDM (Figure S19; Figure 5). The VMLTs were derived either from hPGCLC_A4_L82 (modeling early hPGCs with ~50% GDM)[36], or from hiPSC_A4_L82 (modeling primed hESCs with ~70% GDM)[36], confirming a profound GDM reset preceding MST to VMLTs.

Direct and indirect tumorigenic pathways to VMLTs
VMLTs' late-hPGC GDM signature enables delineation of the tumorigenic pathways for VMLTs derived from hPGCLC_A4_L82 versus those derived from high-number hiPSC_A4_L82 cells. A direct pathway, for hPGCLC_A4_L82-derived VMLTs, involved further germline progression from an early-hPGC like state (modeled by hPGCLC_A4_L82)[36] to the requisite "late-hPGC" state in vivo, followed by MST to VMLTs (Figure 5). In contrast, an indirect pathway, for hiPSC_A4_L82-derived VMLTs, necessitated an extra initial hPGCLC specification step, in which a small subset of naïve hiPSC_A4_L82 cells first

differentiated into hPGCLC_A4_L82 cells. These then proceeded to the "late-hPGC" state before undergoing MST to VMLTs (Figure 5). The observed dosage effect - VMLT formation in high-number (2×10^7) but not conventional-number (1×10^6) hiPSC_A4_L82 injections, with no such effect in low-number (3×10^5) hPGCLC_A4_L82-derived tumors - can be attributed to this additional hPGCLC specification bottleneck. Moreover, clinical observations of rare MST from pure seminomas to somatic malignancies[61] including HGNEC intermixed with seminoma cells[62], offer further clinical support for the plausibility of direct MST from a "late-hPGC"-like state to somatic cancer, reinforcing the pathway proposed here.

MCC and seminoma lines show similar GDM

VP-MCC and TCam2 lines showed similar GDM

To further validate the findings supporting VP-MCC tumorigenesis via direct transformation from an obligatory "late-hPGC" state, we analyzed global 5mC levels in six VP-MCC cell lines and compared them to the seminoma cell line TCam-2, which consists of transformed late-hPGCs[1]. Flow cytometry showed that VP-MCC lines exhibited moderate global DNA hypomethylation (MFI: ~5K-16K), similar to TCam-2 (~7K), though slightly higher (p<0.027) than VPMLs (~2K-3.5K) (Figure 4A; Table S4). As both VP-MCC lines and TCam-2 were cultured under identical conditions (RPMI + 10% FBS), their mildly elevated 5mC levels relative to that of VMLTs likely reflect epigenetic drift due to in vitro culture conditions as well[59].

VN-vMCC and TCam2 lines also showed similar GDM

Interestingly, the two VN-vMCC cell lines, MCC-13 and UISO, grown under the same culture condition also displayed comparable levels of moderate hypomethylation (MFI: ~7K-~9K; Figure 4A, Table S4) to that of VP-MCC cell lines, aligning with previous findings[56,57]. The similarity in GDM between VN-vMCC lines and TCam-2 under matched culture conditions suggests that primary VN-vMCC tumors may also exhibit extreme hypomethylation - an epigenetic hallmark of the "late-hPGC" state observed in primary seminomas. In the case of VN-vMCCs, this "late-hPGC"-like state may arise via reprogramming from somatic cells driven by extensive somatic mutations, including pRB and p53 mutations, characteristic of these tumors[12,20,57] (Figure 5).

Discussion

Functional evidence against parthenogenic intermediates for MST to VMLT

Failure of hEGCLC_A4_L82 to derive VMLT supports no pluripotent intermediate for MST

Failure of hEGCLC_A4_L82 to form VMLTs in vivo indicating a lack of direct conversion to VMLTs from hEGCLC_A4_L82 - the in vitro pluripotent derivative of hPGCLC_A4_L82. This finding therefore provides the functional evidence against the involvement of a parthenogenetic pluripotent hESC-like intermediate in MST to VMLTs, and supports a VMLT tumorigenic pathway that is independent of teratoma formation and uncoupled from parthenogenetic pluripotency.

Decreased potency of hEGCLC_A4_L82 suggests no totipotent intermediate for MST

Previous studies showed dichotomous parthenogenic paradigms for normal hPGCs vs transformed hPGCs in NSGCCs[1,23,36]. The developmental potency of parthenogenetic derivatives of normal hPGCs, such as hEGCLCs or hEGCs, has been shown to decrease along the hPGC developmental trajectory: hEGCs derived from post-migratory late-hPGCs lose bona fide pluripotency, whereas hEGCLCs originating from hPGCLCs (modeling early hPGCs) regain pluripotency, as demonstrated by their inability or ability to form teratomas in vivo, respectively[23, 36]. Thus, prior entry into the germline and a corresponding decline in developmental potency define the normal hPGC parthenogenesis paradigm. In contrast, transformed hPGCs in NSGCCs exhibit a reversed trajectory, with parthenogenetic derivatives showing increased developmental potency along germline progression: transformed late-stage hPGCs give rise to totipotent EC cells, while transformed early-stage hPGCs yield less potent, primed-state hESC-like derivatives. In this transformed paradigm, prior germline entry is associated with an upregulation of developmental potency - a distinct hallmark of NSGCC transformation[1].

The inability of high-number hEGCLC_A4_L82 cells to form VMLTs in mice, compared to the successful VMLT formation by high-number hiPSC_A4_L82 cells, provides functional evidence that hEGCLC_A4_L82 had failed to transition to the totipotent naïve hESC-like state required for VMLT induction via the indirect pathway through hPGCLC_A4_L82 specification. This reduced developmental potency relative to its non-parthenogenetic counterpart aligns with the normal, rather than NSCGC-transformed, hPGC parthenogenesis paradigm. This observation reflects MCPyV's inability to functionally

enhance parthenogenetic potency in hPGCLCs_A4_L82, in contrast to the developmental potency upregulation characteristic of NSGCC transformation. MCPyV's apparent incapacity to induce NSGCC-like transformation indicates that hPGCLC_A4_L82 cells, upon viral transduction, cannot parthenogenetically acquire a totipotent EC-like state. These findings are consistent with the GDM evidence supporting the requirement of a "late hPGC" state before direct MST to somatic VMLTs, without the involvement of a totipotent parthenogenetic intermediate. Furthermore, MCPyV's inability to induce NSGCT-like transformation also suggests that the teratoma components observed in hPGCLC_A4_L82-derived VMLT(+) tumors have arisen directly from an in vivo parthenogenic derivative competent for teratoma formation, from hPGCLC_A4_L82 modeling early-hPGCs[36] as spontaneous germline developmental progression in vivo to a "late-hPGC" state would have resulted in a parthenogenic derivative incapable of teratoma formation[23].

Independent tumorigenic pathways for teratoma vs VMLT

Based on the VMLT tumorigenic pathway supported by the obligatory extremely low GDM unique for "late-hPGC" state and the known intrinsic pluripotency-teratoma coupling with teratoma components exhibiting much higher GDM in VMLT(+) tumors (Figure 4B), an independent tumorigenic pathway is favored for the teratoma components of VMLT(+) tumors, involving multi-lineage somatic differentiation from a pluripotent precursor[23]. Failure of hEGCLC_A4_L82 to form VMLTs also provided the functional evidence to oppose the involvement of parthenogenic intermediates in VMLT tumorigenesis as discussed above. The independent tumorigenic pathways for the VMLT component versus the teratoma component can provide the rationale for the co-presence of VMLT and teratoma components in this VP-MCC mouse xenograft model, while no mixed VP-MCC and teratoma tumor has been reported in clinical settings. The answers lie in the fact that the experimental mouse teratomas derived from hPSCs are known to be polyclonal[23], while clinical cancer is a clonal disease originated from a single transformed cell including VP-MCC that is characterized by a clonal viral insertion site[10].

Early developmental vs carcinogenic pathways

Notably, the "late-hPGC" state is also a requisite intermediate in GCCs, indicating overlapping cancer biology between GCCs and VP-MCC, a somatic cancer. This convergence is exemplified by the globally hypomethylated DNA epigenetic profile - a recognized molecular hallmark of cancer observed across GCCs and somatic cancer alike[1,21]. On the other hand, our model supports a fundamental divergence between normal human development and VP-MCC carcinogenesis at the level of the Weismann Barrier (WB). In canonical early human development, pluripotent hESCs generate somatic tissues through orderly, multi-lineage differentiation. In contrast, VP-MCC appears to originate from a "unipotent" late-hPGC with known intrinsic germline suppression of somatic fate, yet bearing latent or poised pluripotency[34], that aberrantly crosses the WB into the somatic compartment, differentiating exclusively along a neuroendocrine lineage and undergoing developmental arrest. These findings suggest that VP-MCC may arise via an aberrant ontogenic process in which the germline directly gives rise to a somatic malignancy - akin to the emergence of a highly abnormal new life form (Figure 5). This MST model also supports key distinctions between somatic cancers such as VP-MCC and GCCs exhibiting multilineage somatic differentiation, namely malignant teratomas. The latter more closely parallels normal human embryogenesis, requiring a totipotent embryonal carcinoma (EC) cell intermediate to traverse the WB and undergo multilineage somatic differentiation with aberrant developmental arrest. These EC cells are thought to arise via parthenogenetic transformation of late-stage hPGCs in vivo, rather than from fertilization-derived zygotes. Future investigations are needed to determine whether this unifying germ cell theory-based paradigm applies more broadly to other somatic malignances. For instance, in VN-vMCCs, somatic mutations may induce a functionally equivalent "late-hPGC" state, substituting for the native "late-hPGC" state observed in VP-MCCs.

In summary, we developed an MST mouse model demonstrating MCPyV-driven transformation of hPGCLCs into somatic VMLTs. GDM analyses identified a "late-hPGC" state as a necessary developmental precursor for direct MST to VMLTs. Spatial, epigenetic, and functional data collectively support the existence of independent tumorigenic trajectories for the VMLT and teratoma components within the same VMLT(+) tumor, and oppose a parthenogenetic pluripotent or totipotent intermediate in VMLT tumorigenesis. This model thereby establishes a direct developmental link between late-hPGC and a somatic cancer, supporting the same cell (or cell state) of origin for both GCCs and somatic cancer. Finally, the genetic simplicity and experimental tractability of our VP-MCC–based MST model provide a

robust platform for elucidating the molecular mechanisms driving VP-MCC pathogenesis. This system may also yield critical insights into the molecular underpinnings of VN-MCC and other treatment-refractory high-grade neuroendocrine carcinomas (HGNECs) characterized by high mutational burdens and genetic complexity. By directly establishing a developmental origin for VP-MCC, this model supports a shift toward a developmental biology-driven framework for advancing somatic cancer research beyond the mutation-centric and soma-centric paradigm.

Method
Cell Culture
The human iPSC A4 line (hiPSC_A4) generated from commercially obtained Caucasian male (46XY) neonatal foreskin fibroblasts and the derived GFP+ long-term-culture human primordial germ cell like cell (LTC-hPGCLC) A4 line (hPGCLC_A4) were gifted by the Shioda lab[36]. The human embryonic germ cell like cell A4 line (hEGCLC_A4) were derived from hPGCLC_A4 based a 10-day conversion protocol facilitated by SCF and FGF2 from the Shioda lab[36]. Both hiPSC and hEGCLC cultures were maintained in the Essential 8 medium (Gibco, Cat#A1517001) or occasionally the Essential 8 Flex medium (Gibco, Cat#A2858501) on human ESC-qualified Matrigel (Corning, Cat#354277) and passaged at every 3-7 days using ReLeSR (StemCell Technologies, Cat#100-0483) in the presence of the ROCK inhibitor Y-27632 (StemCell Technologies, Cat#72304). Karyotyping analysis were performed every 10-15 passages to assure the genetic integrity of hiPSCs, hEGCLCs and hPGCLCs and their L82 transfected counterparts (Figure S20). The six VP-MCC cell lines, CVG-1, MKL-1, MKL-2, MS-1, PeTa and WaGa were cultured in RPMI 1640 medium (Gibco, Cat#11875093) with 10% FBS (Sigma, Cat#F2442-500ML) as suspension cultures, and passaged every 3-4 days by 1:2 to 1:10 split. The two VN-vMCC cell lines MCC-13 and UISO as well as the TCam2 cell line were also cultured in RPMI 1640 medium (Gibco, Cat#11875093) with 10% FBS (Sigma, Cat#F2442-500ML) as adherent cultures, and passage every 3-4 days by incubation with 0.25% trypsin from Trypsin (Gibco, Cat#15090046) for up to 8 minutes. Mycoplasma contamination testing was performed every six months or when there was a suspicion for contamination was ruled out by performing a PCR based mycoplasma detection kit based on manufacturer's protocol (PCR mycoplasma detection kit (Thermo Scientific, Cat#J66117.AMJ).

Derivation of McPyv+ primeval stem cell lines hPGCLC_A4_L82, hiPSC_A4_L82 hEGCLC_A4_L82
The lentivirus virus vector plenti.puro.MCV.ER.RAZ.2 (a gift from Reety Arora & Sudhir Krishna Addgene plasmid #114382; http://n2t.net/addgene:114382 ; RRID:Addgene_114382) abbreviated as "L82" containing a C-terminus truncated McPyv genome and puromycin resistance gene, a 2nd generation packaging vector psPAX2 (AddGene#12260) and an envelope vector pMD2.G (AddGene#12259) were purchased from AddGene.Org. Lentivirus productions of McPyv containing lentivirus were done with FuGene 6 transfection reagent (Promega, Cat#E2691) and OptiMem serum free medium (Invitrogen Cat#31985). Stable lentivirus transfections followed by puromycin selection to produce McPyv integrated derivatives hPGCLC_A4_L82 and hiPSC_A4_L82 followed standard AddGene protocol with polybrene (Santa Cruz Biotechnology, Cat#sc-134220) and minor modification due to the requirement of daily Essential 8 medium change for pluripotent stem cell line hiPSC_A4. McPyv integrated hEGCLC_A4_L82 was derived from hPGCLC_A4_L82 following the same 10-day in vitro conversion protocol[36] facilitated by SCF and FGF2 used for converting hPGCLCs to hEGCLCs.

McPyv+ human primeval stem cell line-derived mouse xenograft study
All animals used in this study were handled in accordance with an animal study protocol approved by the Uniformed Services University of Health Sciences Institutional Animal Care and Use Committee. As MCC occur in adults with a male dominance, 4–6-week-old adult and male immunocompromised NSG mice [NOD.Cg PrkdcscidIl2rgtm1Wjl/SzJ, Jackson Laboratory Stock# 005557] were used for the study. Simple physical restraint with double-hand method was used for injecting cells subcutaneously to mouse flank. Mice were monitored regularly for health status and were euthanized by CO2 asphyxiation upon reaching humane endpoint criteria. Three mice were used for testing tumorigenicity for each group of total six groups. For the first two groups, only 3×10^5 cells/injection of the hPGCLC_A4 line or the hPGCLC_A4_L82 line was injected to the left flank of three mice for each cell line. For the hiPSC_A4_L82 line, 1×10^6 cells/injection and 2×10^7 cells/injection were injected to bilateral flanks of three mice in two groups. The same injections were also performed for hEGCLC_A4_L82 in two groups as for hiPSC_A4_L82. All human primeval stem cells were suspended in 50 ul with 10 µM ROCK inhibitor Y-27632 and mixed with an equal volume of Matrigel (Corning, Cat#354277) before injected

subcutaneously into the flanks of the mice[31,36]. Mice were monitored regularly for tumor development and sacrificed upon reaching humane endpoint criteria. Tumors were collected by dissection; friable portions of tumors were collected in sterile RPMI to prepare cell suspension by mechanical disruption using scissors and forceps. Spun down cell pellets by centrifuge at 400xg were then used for both single cells suspension for flow cytometry and to extract total RNA using the PureLink RNA extraction mini kit (Thermo Fisher, Cat#2592159), as well as cryopreserved in FBS (Sigma, Cat#F2442-500ML) with 8% DMSO (Thermo Fisher, Cat#M3148-250ml). Cryopreserved tumor cells were later thawed to extract DNA by PureLink™ Genomic DNA Mini Kit (Termo Fisher, Cat#182001). The rest of the tumors were fixed 4% paraformaldehyde, and subjected to the standard H&E as well as IHC staining. Cell line authenticity for all VMLT (+) tumors was confirmed by no variation at any loci of five mononucleotide repeats and two pentanucleotide repeats (NR-21, BAT-26, BAT-25, NR-24, MONO-27; and Penta C, Penta D) among themselves and all three human primeval stem cell lines (hPGCLC_A4_L82 and hiPSC_A4_L82), using Promega OncoMate MSI Analysis System (Cat. # MD2140; Madison, WI) and DNA extracted from the tumor cells and cell lines (Figure S21).

RT-qPCR
Total RNA of cell lines was also extracted using the PureLink RNA extraction mini kit (Thermo Fisher, 2592159). Reverse transcription was performed using Luna one-step RNA-to-Ct RT-qPCR kit (New England Biolab, E3005E), 5 uM gene-specific primer pairs and 5ng RNA per final 20-ul RT-qPCR reaction and Azure Ciel 3 qPCR detection system (Azure Biosystems, CA) with the cycling/melting curve RT-qPCR program recommended by the manufacturers. All primer sequences are listed in table S4. mRNA expression of each gene was normalized relative to the mRNA expression of housekeeping gene RnaseP as ΔCq. All RT-qPCR experiments were repeated at least three times.

Immunohistochemistry (IHC) Study: Sections of CDX mouse tumors or pellets of cell lines were first deparaffinized by incubating in xylene, followed by rehydration with ethanol solutions (100% and 95%) and distilled water. Antigen retrieval was performed using EDTA Antigen Retrieval Buffer (pH 8.0, 10X) (Vitro Vivo Biotech, Cat#VB-6007) in a steamer, with sections boiled for 40 minutes-1 hour in 1X EDTA Buffer (Table S6). After washing, endogenous peroxidase activity was blocked with 3% hydrogen peroxide prepared from Hydrogen Peroxide 30% Solution (Lab Alley, Cat#HPL30) and sections were then blocked with 5% goat serum (Gibco, Cat#16210064) in TBST (QuickSilver Powdered Buffer Packs, Cat#EB-1203) for 1 hour. Primary antibodies were diluted based on manufacturer's recommendation and empirical testing (table S5) and the Rabbit IgG isotype control (Invitrogen, Cat# 31235) or mouse IgG isotype control (Invitrogen, Cat#31903) was diluted to the same final concentrations. If the primary antibodies are of rabbit host, they were incubated overnight at 4°C with the only exception of 5mC antibody (Invitrogen, Cat#MA5-24694) for one hour at room temperature. Following antibody incubation, SignalStain® Boost IHC Detection Reagent (HRP, Rabbit) (Cell Signaling Technology, Cat#8114S) was applied per the manufacturer's instruction, and sections were incubated for 30 minutes. However, if the primary antibodies were of mouse host, to exclude false positive signal from mouse tissue, the sections were stained for primary and secondary antibodies by M.O.M.® (Mouse on Mouse) ImmPRESS® HRP (Peroxidase) Polymer Kit (Vector Laboratories, Cat#MP-2400) per the manufacturer's protocol. SignalStain® DAB Substrate Kit (Cell Signaling Technology, Cat#8059S) was applied to visualize the antigen-antibody complex, and counterstaining was performed with Hematoxylin (Cell Signaling Technology, Cat#14166S). The sections were subsequently dehydrated through 95% and 100% ethanol followed by xylene before being mounted with SignalStain® Mounting Medium II (Cell Signaling Technology, Cat#84583S) under a coverslip.

Flow cytometry 5mC Analysis
Single cell suspensions were prepared for VMLT+ mouse tumors, VP-MCC cell lines and hiPSC_A4_L82 line, as well as VN-MCC cell lines and TCam2 cell line via different methods. Single cell suspensions were made for friable mouse VMLTs by mechanical disruption with scissors and forceps in sterile RPMI. Single cell suspensions of floating VP-MCC cell lines and attached hiPSC_A4_L82 line required cell separation by incubation with ReLeSR™ (Stem Cell Technologies, Cat#100-0483) for 20+ minutes and up to 10 minutes at 37°C respectively. Single cell suspensions of attached VN-MCC cell lines and TCam2 cell line required cell separation by incubation with 1.25% trypsin (Gibco, Cat#15090046) for up to 8 minutes. All separated cells were then filtered through 100 um cell strainers (Ibis Scientific, Cat#229486). Prepared single cell suspensions were then stained with eBioscience™ Fixable Viability Dye eFluor™

780 (Invitrogen, Cat#65-0865-14) for 30 minutes in a blocking buffer of 2% FBS (MilliporeSigma, Cat#F2442-500ML) in PBS (Cytiva, Cat#SH30256.02). eBioscience™ Foxp3/Transcription Factor Staining Buffer Set (Invitrogen, Cat#00-5523-00) was used for cell fixation and antibody staining per the manufacturer's protocol. Briefly, the cell suspensions were fixed and stained with 5-Methylcytosine (5mC) Recombinant Rabbit Monoclonal Antibody (RM231) (Invitrogen, Cat#MA5-24694) at 1:1000 final dilution and/or the Rabbit IgG Isotype Control (Invitrogen, Cat#31235) at the same final concentration for 30 minutes on ice, followed by staining with Goat anti-Rabbit IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 790 (Invitrogen, Cat#A11369) at 1:500 final dilution for 30 minutes. The cells were then stained with Alexa Fluor™ 594 conjugated Rabbit Sox2 antibody (Santa Cruz Biotechnology, Cat# sc-365823) for 30 minutes on ice. Cell suspensions were then placed in the eBioscience™ Flow Cytometry Staining Buffer (Invitrogen, Cat#00-4222-26) and analyzed by flow cytometer Cytek Aurora. FlowJo software was used to analysis flow data by gating the Sox2(+) population for VMLT cells and the Median Fluorescence Intensity (MFI) at Alexa Fluor™ 790 as well as the Standard Error of the Mean (SEM) were calculated by the FlowJo.

Statistical Analysis
All RT-qPCR experiments were repeated at least three times. p values were calculated by either paired or unpaired one tail Student T-test. A p-value with <0.05 was considered statistically significant. Based on a binomial distribution of single injection success rate of 80-100% for teratoma formation from subcutaneous injection of hPSCs mixed with Matrigel[31], we estimated three-mice per group being the minimum number to ensure the achievement of a high success rate (99.2%) of tumor formation.

Data availability the datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

Acknowledgement
We would like to thank Toshi Shioda for providing LTC-hPGCLC_A4 and hiPSC_A4 cell lines, technical guidance and protocols; Yuan Chang, Patrick Moore, Masahiro Shuda, Isaac Brownell and James DeCaprio for providing virus-positive and virus-negative Merkel cell carcinoma cell lines, technical guidance, and/or protocols; Alan Epstein for providing TCam2 cell line and protocols; Reety Arora and Sudhir Krishna for AddGene plasmids; Joel Moncur for mouse xenograft tumor histopathological evaluation and manuscript review; Guanghua Wang for cell line and mouse xenograft tumor authentication, mouse tumor histopathological and manuscript review; Geeta Upadhyay for technical advices and manuscript review; Steve Dumler and Dennis Grab for manuscript review; as well as the USUHS Genomics Core and Flow Cytometry Core Lab in Biomedical Instrumentation Center for DNA primer synthesis and flow cytometry services respectively. This study was funded by Uniformed Services University of the Health Sciences StartUp fund.

Ethics Declaration The authors declare no competing interests.

Disclaimer The opinions and assertions expressed herein are those of the author(s) and do not reflect the official policy or position of the Uniformed Services University of the Health Sciences or the Department of Defense.

Author Contributions: Conceptualization: WY; Methodology: WY; Investigation: WY, SR; Visualization: SC, WY; Funding acquisition: WY; Project administration: WY; Supervision: WY; Writing - original draft: WY; Writing - review & editing: WY, SC

Materials & Correspondence Any correspondence and material requests should be addressed to corresponding author Wendy Yang.

References

1. J. W. Oosterhuis, L. H. J. Looijenga, Human germ cell tumours from a developmental perspective. *Nature Reviews Cancer* **19**, 522-537 (2019).
2. S. M. Tu, L. L. Pisters, Curing Cancer: Lessons from a Prototype. *Cancers (Basel)* **13**, (2021).

3. S. Huang, A. M. Soto, C. Sonnenschein, The end of the genetic paradigm of cancer. *PLoS Biol* **23**, e3003052 (2025).

4. J. Taguchi *et al.*, DMRT1-mediated reprogramming drives development of cancer resembling human germ cell tumors with features of totipotency. *Nat Commun* **12**, 5041 (2021).

5. J. E. Visvader, G. J. Lindeman, Cancer stem cells in solid tumours: accumulating evidence and unresolved questions. *Nat Rev Cancer* **8**, 755-768 (2008).

6. J. B. Kum *et al.*, Molecular genetic evidence supporting the origin of somatic-type malignancy and teratoma from the same progenitor cell. *Am J Surg Pathol* **36**, 1849-1856 (2012).

7. J. Taylor *et al.*, Germ cell tumors and associated hematologic malignancies evolve from a common shared precursor. *J Clin Invest* **130**, 6668-6676 (2020).

8. N. Wyvekens *et al.*, Molecular correlates of male germ cell tumors with overgrowth of components resembling somatic malignancies. *Mod Pathol* **35**, 1966-1973 (2022).

9. I. Damjanov, P. W. Andrews, Teratomas produced from human pluripotent stem cells xenografted into immunodeficient mice - a histopathology atlas. *Int J Dev Biol* **60**, 337-419 (2016).

10. H. Feng, M. Shuda, Y. Chang, P. S. Moore, Clonal integration of a polyomavirus in human Merkel cell carcinoma. *Science* **319**, 1096-1100 (2008).

11. P. W. Harms *et al.*, The biology and treatment of Merkel cell carcinoma: current understanding and research priorities. *Nat Rev Clin Oncol* **15**, 763-776 (2018).

12. E. A. Pedersen, M. E. Verhaegen, M. K. Joseph, K. L. Harms, P. W. Harms, Merkel cell carcinoma: updates in tumor biology, emerging therapies, and preclinical models. *Front Oncol* **14**, 1413793 (2024)

13. D. Hanahan, R. A. Weinberg, Hallmarks of cancer: the next generation. *Cell* **144**, 646-674 (2011).

14. S. Sell, G. B. Pierce, Maturation arrest of stem cell differentiation is a common pathway for the cellular origin of teratocarcinomas and epithelial cancers. *Lab Invest* **70**, 6-22 (1994).

15. A. Jögi, M. Vaapil, M. Johansson, S. Påhlman, Cancer cell differentiation heterogeneity and aggressive behavior in solid tumors. *Ups J Med Sci* **117**, 217-224 (2012).

16. D. Chen *et al.*, Human Primordial Germ Cells Are Specified from Lineage-Primed Progenitors. *Cell Rep* **29**, 4568-4582 e4565 (2019).

17. M. De Felici *et al.*, To Be or Not to Be a Germ Cell: The Extragonadal Germ Cell Tumor Paradigm. *Int J Mol Sci* **22**, (2021).

18. D. S. Nin, L. W. Deng, Biology of Cancer-Testis Antigens and Their Therapeutic Implications in Cancer. *Cells* **12**, (2023).

19. A. J. Simpson, O. L. Caballero, A. Jungbluth, Y. T. Chen, L. J. Old, Cancer/testis antigens, gametogenesis and cancer. *Nat Rev Cancer* **5**, 615-625 (2005).

20. Z. Ma, F. Zhang, A. Liu, H. Lin, C. Liu, Somatic-to-primordial germ cell-like transformation is critical in tumor initiation of mouse breast tumor 4T1 cells. *Am J Cancer Res* **13**, 1471-1485 (2023).

21. M. Ehrlich, DNA hypomethylation in cancer cells. *Epigenomics* **1**, 239-259 (2009)

22. J. Montilla-Rojo *et al.*, Teratoma Assay for Testing Pluripotency and Malignancy of Stem Cells: Insufficient Reporting and Uptake of Animal-Free Methods-A Systematic Review. *Int J Mol Sci* **24**, (2023).

23. M. Aleckovic, C. Simon, Is teratoma formation in stem cell research a characterization tool or a window to developmental biology? *Reprod Biomed Online* **17**, 270-280 (2008).

24. Lu Z, Fan P, Huo W, Feng Y, Wang R. Genomic profiles and their relationships with clinical characteristics and immune features in cervical cancer. Transl Oncol. 2024 Jun;44:101923. doi: 10.1016/j.tranon.2024.101923. Epub 2024 Mar 2. Erratum in: Transl Oncol. 2024 Sep;47:102037. doi: 10.1016/j.tranon.2024.102037. PMID: 38432114; PMCID: PMC10920960.

25. T. Kervarrec *et al.*, Histogenesis of Merkel Cell Carcinoma: A Comprehensive Review. *Front Oncol* **9**, 451 (2019).

26. N. G. Balanis *et al.*, Pan-cancer Convergence to a Small-Cell Neuroendocrine Phenotype that Shares Susceptibilities with Hematological Malignancies. *Cancer Cell* **36**, 17-34.e17 (2019).

27. Z. Zhang *et al.*, Stemness and inducing differentiation of small cell lung cancer NCI-H446 cells. *Cell Death Dis* **4**, e633 (2013).

28. Csoboz B, Rasheed K, Sveinbjørnsson B, Moens U. Merkel cell polyomavirus and non-Merkel cell carcinomas: guilty or circumstantial evidence? APMIS. 2020 Feb;128(2):104-120. doi: 10.1111/apm.13019. Epub 2020 Jan 28. PMID: 31990105.

29. B. M. Foster, K. L. Langsten, A. Mansour, L. Shi, B. A. Kerr, Tissue distribution of stem cell factor in adults. *Exp Mol Pathol* **122**, 104678 (2021).
30. C. Quan *et al.*, Dermal fibroblast expression of stromal cell-derived factor-1 (SDF-1) promotes epidermal keratinocyte proliferation in normal and diseased skin. *Protein Cell* **6**, 890-903 (2015).
31. T. A. Prokhorova *et al.*, Teratoma formation by human embryonic stem cells is site dependent and enhanced by the presence of Matrigel. *Stem Cells Dev* **18**, 47-54 (2009).
32. N. Irie *et al.*, SOX17 is a critical specifier of human primordial germ cell fate. *Cell* **160**, 253-268 (2015).
33. K. Sasaki *et al.*, Robust In Vitro Induction of Human Germ Cell Fate from Pluripotent Stem Cells. *Cell Stem Cell* **17**, 178-194 (2015).
34. H. G. Leitch, A. Smith, The mammalian germline as a pluripotency cycle. *Development* **140**, 2495-2501 (2013).
35. F. Sugawa *et al.*, Human primordial germ cell commitment in vitro associates with a unique PRDM14 expression profile. *EMBO J* **34**, 1009-1024 (2015).
36. M. Kobayashi *et al.*, Expanding homogeneous culture of human primordial germ cell-like cells maintaining germline features without serum or feeder layers. *Stem Cell Reports* **17**, 507-521 (2022).
37. M. J. Shamblott *et al.*, Derivation of pluripotent stem cells from cultured human primordial germ cells. *Proc Natl Acad Sci U S A* **95**, 13726-13731 (1998).
38. K. Takahashi, S. Yamanaka, Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors. *Cell* **126**, 663-676 (2006).
39. J. K. Cinkornpumin *et al.*, Naive Human Embryonic Stem Cells Can Give Rise to Cells with a Trophoblast-like Transcriptome and Methylome. *Stem Cell Reports* **15**, 198-213 (2020).
40. T. Messmer *et al.*, Transcriptional Heterogeneity in Naive and Primed Human Pluripotent Stem Cells at Single-Cell Resolution. *Cell Rep* **26**, 815-824 e814 (2019).
41. L. R. Dresang *et al.*, Response of Merkel cell polyomavirus-positive merkel cell carcinoma xenografts to a survivin inhibitor. *PLoS One* **8**, e80543 (2013).
42. T. M. Ulbright, Germ cell tumors of the gonads: a selective review emphasizing problems in differential diagnosis, newly appreciated, and controversial issues. *Mod Pathol* **18 Suppl 2**, S61-79 (2005).
43. D. M. Berney *et al.*, An introduction to the WHO 5th edition 2022 classification of testicular tumours. *Histopathology* **81**, 459-466 (2022).
44. A. Guastafierro *et al.*, Characterization of an early passage Merkel cell polyomavirus-positive Merkel cell carcinoma cell line, MS-1, and its growth in NOD scid gamma mice. *J Virol Methods* **187**, 6-14 (2013).
45. R. Houben *et al.*, Mechanisms of p53 restriction in Merkel cell carcinoma cells are independent of the Merkel cell polyoma virus T antigens. *J Invest Dermatol* **133**, 2453-2460 (2013).
46. R. Houben *et al.*, Merkel cell polyomavirus-infected Merkel cell carcinoma cells require expression of viral T antigens. *J Virol* **84**, 7064-7072 (2010).
47. E. M. Martin *et al.*, Parathyroid hormone-related protein, chromogranin A, and calcitonin gene products in the neuroendocrine skin carcinoma cell lines MKL1 and MKL2. *Bone Miner* **14**, 113-120 (1991).
48. S. T. Rosen *et al.*, Establishment and characterization of a neuroendocrine skin carcinoma cell line. *Lab Invest* **56**, 302-312 (1987).
49. M. Shuda *et al.*, T antigen mutations are a human tumor-specific signature for Merkel cell polyomavirus. *Proc Natl Acad Sci U S A* **105**, 16272-16277 (2008).
50. C. Velásquez *et al.*, Characterization of a Merkel Cell Polyomavirus-Positive Merkel Cell Carcinoma Cell Line CVG-1. *Front Microbiol* **9**, 713 (2018).
51. S. G. Ronan, A. D. Green, A. Shilkaitis, T. S. Huang, T. K. Das Gupta, Merkel cell carcinoma: in vitro and in vivo characteristics of a new cell line. *J Am Acad Dermatol* **29**, 715-722 (1993).
52. J. A. Thomson, J. H. Leonard, K. McGregor, R. A. Sturm, P. G. Parsons, A nonconsensus octamer-recognition sequence (TAATGARAT-motif) identifies a novel DNA binding protein in human Merkel cell carcinoma cell lines. *Int J Cancer* **58**, 285-290 (1994).
53. T. Gambichler *et al.*, Prognostic relevance of high atonal homolog-1 expression in Merkel cell carcinoma. *J Cancer Res Clin Oncol* **143**, 43-49 (2017).
54. H. Guo *et al.*, The DNA methylation landscape of human early embryos. *Nature* **511**, 606-610 (2014).

55. L. Wen, F. Tang, Human Germline Cell Development: from the Perspective of Single-Cell Sequencing. *Mol Cell* **76**, 320-328 (2019).
56. J. Gravemeyer *et al.*, DNA-methylation patterns imply a common cellular origin of virus- and UV-associated Merkel cell carcinoma. *Oncogene* **41**, 37-45 (2022).
57. P. W. Harms *et al.*, Viral Status Predicts the Patterns of Genome Methylation and Decitabine Response in Merkel Cell Carcinoma. *J Invest Dermatol* **142**, 641-652 (2022).
58. Y. Mizuno, A. Gotoh, S. Kamidono, S. Kitazawa, [Establishment and characterization of a new human testicular germ cell tumor cell line (TCam-2)]. *Nihon Hinyokika Gakkai Zasshi* **84**, 1211-1218 (1993).
59. D. J. Smiraglia *et al.*, Excessive CpG island hypermethylation in cancer cell lines versus primary human malignancies. *Hum Mol Genet* **10**, 1413-1419 (2001).
60. D. Nettersheim *et al.*, Analysis of TET expression/activity and 5mC oxidation during normal and malignant germ cell development. *PLoS One* **8**, e82881 (2013).
61. K. P. Dieckmann *et al.*, Pure Testicular Seminoma Relapsing Late with Somatic Type Malignancy. *Case Rep Oncol Med* 2017, 2457023 (2017).
62. Kumaki, N. *et al.* Immunohistochemical analysis of neuroendocrine (NE) differentiation in testicular germ cell tumors (GCTs): use of confocal laser scanning microscopy (CLSM) to demonstrate direct NE differentiation from GCTs. *Acta Histochem Cytochem* **40**, 143-151, doi:10.1267/ahc.07025 (2007).

Author contributions: Conceptualization: WY; Methodology: WY; Investigation: WY, SR; Visualization: SC, WY; Funding acquisition: WY; Project administration: WY; Supervision: WY; Writing - original draft: WY; Writing - review & editing: WY, SC

Competing interests: Authors declare that they have no competing interests.

Data and materials availability: hPGCLC_A4 and hiPSC_A4 cell lines and their derivative cell lines are subject to the restrictions stated in a materials transfer agreement (MTA) with Dr. Toshi Shioda lab at Massachusetts General Hospital. All data are available in the main text or the supplementary materials.

Disclaimer
The opinions and assertions expressed herein are those of the author(s) and do not reflect the official policy or position of the Uniformed Services University of the Health Sciences or the Department of Defense.

Supplementary Materials
Figures S1 to S21
Tables S1 to S6

Fig. 1. Gross appearance and representative histology of mouse VP-MCC-like tumors (VMLTs) derived from human primeval stem cells: Injected with hPGCLC_A4_L82 (~ 3 × 10^5 cells): **(A)** Mouse #85.3. **(B)** Mouse #85.4. Injected with hiPSC_A4_L82 (~ 2 × 10^7 cells): **(C)** Mouse # 85.2. **(D)** Mouse #85.1 **(E)** Mouse #84.5.

Fig. 2. Heatmap of mRNA expression. Comparison of mRNA expression of key neuroendocrine lineage genes, core pluripotent genes, PGC marker genes and McPyv LT viral oncogene among transfected primeval stem cell lines (hiPSC_A4_L82 and hPGCLC_A4_L82), their derivative mouse CDX tumors, VP-MCC and VN-MCC cell lines.

Fig. 3. Immunohistochemical studies of mouse VMLTs. (A) Protein expression of a panel of genes in VMLT of T#85.4.N1. **(B)** High Ki67 proliferative indexes of VMLTs from five mice. **(C)** Positive expression of McPyv LT oncoprotein in VMLT from five mice. MKL-2 cell pellet was used as a positive control.

Fig. 4. Global 5mC levels to estimate GDM by (A) flow cytometry and (B) immunohistochemical (IHC) studies.
(A) Flow cytometry estimated global 5mC levels semi-quantitively of hiPSC_A4_L82 cell line, seminoma cell line TCam2 and its isotype negative control, VMLTs from five mice, six VP-MCC cell lines and two VN-MCC cell lines. Global 5mC levels were measured as Median Fluorescence Intensity (MFI) with error bars indicating Standard Error of the Mean (SEM). hiPSC_A4_L82 line and seminoma cell line Tcam2 exhibited very high (~113K) and low (~7K) global 5mC levels indicating GDM levels consistent with previous studies[36,60]. The isotype negative control of TCam2 exhibited extremely low (~3K) global 5mC level as the background signal for absence of GDM, similar to that of VMLTs indicating nearly complete GDM erasure. VP-MCC and VN-MCC cell lines showed low global 5mC levels similar to that of the TCam-2 cell line, which is composed of transformed late-hPGCs. **(B)** Global 5mC IHC studies of hiPSC_A4_L82 cell line (B1), TCam2 line (B2) and its isotype control (B6), as well as the VMLT components from VMLT(+) tumors T#85.4 (B7) and T#85.2R (B9) showed similar findings as that of flow cytometry. Positive IHC stains of mouse#85.4 somatic skin (B3) and testis (B4) served as positive controls. Positive IHC stains are observed in the teratoma components of VMLT(+) tumors T#85.4 (B8) and T#85.2R (B10), similar to that of pure teratoma of VMLT(-) tumor T#83.5L (B5). Scattered positive cells (black arrows), next to the VMLT tumor cells exhibiting near negative IHC staining, in VMLT(+) T#85.4 and T85.2R tumors are mouse somatic white blood cells (B7) or dermal stromal cells (B9) respectively.

Fig. 5. Schematic of normal human embryonic and germline development in comparison to the carcinogenesis of somatic VP-MCC and germ cell tumors (GCTs). Light green arrows show normal early embryogenesis from fertilization to primed pluripotent hESCs before crossing the Weismann Barrier (WB) into somatic differentiation to form the soma. The intrinsic plasticity between the 2-cell stage and the primed hESC stage is reflected in the bidirectionality of the light green arrows[1]. Normal human germline development (dark green arrows) begins at the specification of hPGCs from formative hESCs and ends with the formation of sperm or eggs, guided by an intrinsic program suppressing somatic fate, yet exhibiting latent/poised pluripotency[34] with the germline being the only human cell lineage capable of physiological conversion to pluripotency (parthenogenesis) without genetic manipulation. Thus, the germline cycle may be considered a "pluripotency cycle"[34], consisting of both early embryogenesis (light green arrows) and germline development (dark green arrows), which are separated from the soma compartment by the Weismann Barrier (WB). Based on our mouse model, carcinogenesis of somatic VP-MCC (red straight arrow) also crosses the WB to the somatic compartment, similar to teratomas - a type of non-seminomatous GCT (NSGCT) (dark purple and light purple straight arrows) - and normal human embryogenesis[1]. However, unlike teratomas or embryogenesis, VP-MCC tumorigenesis crosses the WB directly from the germline (late hPGCs) to the somatic side without a pluripotent intermediary, resulting in mono-lineage (neuroendocrine) somatic differentiation accompanied by developmental arrest. In contrast, tumorigenesis of malignant teratomas more closely mirrors normal embryogenesis, requiring a totipotent EC cell (ECC) intermediate to cross the WB and enable multilineage somatic differentiation. These totipotent stem cells of malignant teratomas are believed to arise via parthenogenetic transformation of transformed hPGCs in vivo (dark purple and light purple curved arrows), rather than from fertilization-derived zygotes. Seminomas, another malignant GCT subtype, consist of transformed late hPGCs and therefore retain the epigenetic signatures characteristic of this stage[1,55]. Two waves of global DNA demethylation occur during development, each followed by de novo methylation. The first wave begins at the 2-cell stage and reaches its nadir at the naïve hESC/ICM stage, with partial erasure of GDM (~40%)[54,55]. The second, germline-specific wave occurs only during hPGC development, resulting in near-complete GDM erasure and reaching the lowest nadir at the late hPGC stage (GDM ~6%)[54,55]. The moderate global hypomethylation observed in two VN-vMCC cell lines similar to that of seminoma cell line TCam2 and VP-MCC lines suggests a possible somatic-to-hPGC transformation (slanted dashed blue arrow with a question mark), potentially triggered by somatic mutations or other mechanisms. The somatic-to-hPGC transformation may also undergo indirect pathway through reprogramming of somatic cells into hiPSCs followed by specification into a hPGC-like state (vertical dashed blue arrow). The three human primeval stem cell types used in our mouse model - hPGCLCs (modeling early hPGCs), hiPSCs (modeling primed hESCs), and hEGCLCs (modeling hEGCs) - are highlighted in yellow.