Discover the Maze-like Pathway for Glabridin Biosynthesis

Zhen Zhang[1], Wenqiang Li[1], Fanze Meng[1], Yanming Jin[1], Wentao Sun[1], Qina Kuang[1], Shichao Ren[2], Wei Liu[1], Liang Zhang[2], Lei Qin[1,3,4,5,*], Bo Lv[2,*], Haiyang Jia[2,*] and Chun Li[1,2,3,4,5,*]

[1]Department of Chemical Engineering/ Key Laboratory for Industrial Biocatalysis of Ministry of Education, Tsinghua University, Beijing, China

[2] Institute of Biochemical Engineering, School of Chemistry and Chemical Engineering/ Key Laboratory of Medical Molecule Science and Pharmaceutical Engineering, Ministry of Industry and Information Technology, Beijing Institute of Technology, Beijing, 100081, P. R. China

[3]Center for Synthetic and Systems Biology, Tsinghua University, Beijing, China

[4]Beijing Key Laboratory of Recombinant Protein Synthetic Biomanufacturing, Beijing, China

[5]State Key Laboratory of Green Biomanufacturing, Beijing, China

*Correspondence to: lichun@tsinghua.edu.cn; haiyangjia@bit.edu.cn; qinleithu@tsinghua.edu.cn and lv-b@bit.edu.cn

Abstract

Tailoring enzymes diversify plant metabolite scaffolds through complex, context-dependent modifications, generating maze-like biosynthetic networks that complicate metabolic pathway reconstruction. Here, we systematically deconstructed and reconstructed the biosynthetic network of glabridin, a valuable skin-whitening isoflavone from *Glycyrrhiza*. By integrating metabolic pathway mining with genome and 183 transcriptomes, we identified four functional routes among sixteen theoretical possibilities and uncovered a previously uncharacterized, ladder-like multi-route tailoring network. Reconstruction of such architecture in yeast revealed metabolic redundancy and interconnectivity confer unexpected robustness, enabling higher production efficiencies compared to a single-route design. Further modular engineering enabled the *de novo* biosynthesis of glabridin in yeast. Our work establishes a generalizable framework for reconstructing metabolic mazes and demonstrates that multi-route architectures can be harnessed to enhance the robustness and productivity of cell factories.

Key words：glabridin, tailoring enzymes, synthetic biology, plant natural protucts, yeast

Introduction

Tailoring enzymes(*1-3*) are essential in plant secondary metabolism, where they steer the synthesis of complex bioactive compounds by modifying a foundational metabolite scaffold. Unlike the initial assembly of a basic molecular framework, tailoring introduces a series of chemical modifications that determine activity, solubility, stability, and subcellular localization(*4, 5*). This post-assembly process is not only critical for generating the diverse array of natural products, but it also adds an additional layer of regulation that ultimately shapes the biological activity of these compounds. However, this tailoring complexity presents a challenge when attempting to reconstruct or engineer the biosynthesis of specific target compounds. The intricate sequence of reactions due to the promiscuity often gives rise to a metabolic network that resembles a maze—each step interwoven and highly context-dependent(*6, 7*). Such complexity makes it exceptionally difficult to fully map the biosynthetic pathways of certain specialized metabolites, especially those with multiple modifiable sites and obscure tailoring logic.

In the realm of flavonoid biosynthesis, more than 9,000 distinct compounds have been tailored from common precursors, representing one of the most structurally diverse classes of natural products(*8*). Among them, glabridin(*9*), a naturally occurring prenylated isoflavone, is renowned for its skin-whitening, anti-inflammatory, and anti-aging properties(*9, 10*). It commands a multimillion-dollar global market in high-end whitening cosmetics. However, glabridin is difficult to chemically synthesize due to the challenge of stereoselective construction of the C-3 chiral center and the chemoselective reduction of conjugated systems(*11*). Currently, the world's supply chain for glabridin relies on low-yielding extraction and purification from the plant root of rare plant *Glycyrrhiza glabra L*, which only accounting for 0.08-0.35% of the roots' dry weight(*10*). Microbial or synthetic biosynthesis of glabridin remains challenging due to a complex maze of post-tailoring modifications, and even its native biosynthetic pathway has not yet been fully elucidated.

In this study, we applied a metabolic pathway AI search algorithm integrated with experimental metabolic data to comprehensively elucidate and map a maze-like biosynthetic network of glabridin. By systematically mining the transcriptome and genome data of *Glycyrrhiza* species, we identified all key tailoring enzymes involved in glabridin biosynthesis. These enzymes were then used to reconstruct a multi-route synthetic pathway, in which a single set of promiscuous enzymes supports multiple parallel biosynthetic routes. Through this reconstruction, we uncovered distinct biosynthetic modes underlying glabridin formation and ultimately achieved *de novo* biosynthesis of glabridin in *Saccharomyces cerevisiae*.

Glabridin pathway design and core nodes identification

To discover the biosynthesis pathway of glabridin, we employed a metabolic pathway search algorithm based on known enzymatic reactions in combination with a retrosynthetic algorithm derived from reaction rules(*12-14*), which enables pathway design by deconstructing target molecule and medicarpin (**8**) as key intermediates into simpler precursors in yeast step-by-step. Using this approach, we designed 16 potential biosynthetic pathways leading to the production of glabridin, involving L-phenylalanine (**1**) as precursor and 31 intermediate metabolites (Fig.1a). Among these, 12 compounds are located at pathway branch nodes, and over 64% of the predicted pathways include more than six branch nodes. The presence of such diverse and numerous branch points significantly increases the complexity of glabridin biosynthesis.

To evaluate the most feasible biosynthetic routes among those predicted, we conducted metabolomic profiling of three *Glycyrrhiza* species: *Glycyrrhiza glabra*, *Glycyrrhiza uralensis*, and *Glycyrrhiza inflata* (Fig.1a-b). Metabolomic analysis revealed the major metabolites, such as compounds **5**, **11**, **13**, **15** and **25**, are predominantly accumulated in the roots compared to leaves and stems. Glabridin was detected only in *G. glabra* and *G. inflata*, with the highest accumulation observed in *G. glabra*. Furthermore, several predicted intermediates (**4**, **8**, **9**, **10**, **12** and **14**) were also identified in *G. glabra*, providing support for their roles as key metabolic network nodes(*15*). Thus, through node-centric analysis of the metabolomic data, the predicted routs major incorporated with key intermediates, such as medicarpin (**8**), were further considered. Even though, 12 possible alternative biosynthetic pathways for glabridin still persist due to the diverse nature of tailoring reactions. This complexity complicates pathway design and hinders efficient metabolic engineering of glabridin.

To address this complexity, we simplified the pathway by categorizing glabridin biosynthesis into four classes of tailoring reactions: pterocarpin reduction, prenylation, prenyl cyclization, and demethylation (Fig. 1c). These transformations are likely mediated by four classes of tailoring enzymes: pterocarpin reductases (PTRs), prenyltransferases (PTs), oxidative cyclases (OCs), and demethylases (DMTs). Among these, PTRs, along with VRs and IFRs, belong to the class of NADPH-dependent reductases(*16-18*), sharing similar catalytic mechanisms and often exhibiting high sequence similarity(*18*), which poses a challenge for accurate PTRs identification. In addition, DMTs catalyze critical branch-forming reactions, with the high branching complexity, which direct the metabolic flux toward different outcomes. PTs and OCs enzymes were found to broaden the range of possible intermediate metabolites. The four tailoring reactions can proceed in different sequential orders through various branching nodes, all ultimately leading to produce glabridin.

Figure 1 The maze-like glabridin biosynthesis pathway. a) Summary of predicted glabridin biosynthesis. b) Metabolomic profiling of different organs and species of *Glycyrrhiza, including Glycyrrhiza glabra* (*Gg*), *Glycyrrhiza uralensis* (*Gu*), and *Glycyrrhiza inflata* (*Gi*). c) The key tailoring steps in glabridin biosynthesis. Dash arrows in a. and c. represent uncharacterized step, gray arrows represent characterized steps, and the overlapping arrows represent a multi-step reaction. The color bar in b. indicates the log$_2$-transformed peak area. Metabolite abundance is categorized into four levels based on accumulation in *G. glabra* roots: high (>24), medium (>18), low (>12) and undected. The bar chart in the upper right corner of each compound in a. indicates its dection in *G. glabra* roots, where the number of solid brown bars corresponds to the abundance level, and hollow bars indicate that the compound was not detected. Abbreviations not already defined in the text are as follows: isoflavanone 2'-O-methyltransferase (2'OMT), aldehyde ketone reductase (AKR), sativan synthase (SR),

isoflavone synthase (IFS), 2,7,4'-trihydroxyisoflavanone 4'-O-methyltransferase (4'OMT), 2-hydroxyisoflavanone dehydratase (HID), isoflavone 2'-hydroxylase (I2'H), isoflavone reductase (IFR), vestitone reductase (VR), and pterocarpan synthase (PTS), 2'-hydroxydihydrodaidzein synthase (THIS), daidzein reductase (DZNR), dihydrodaidzein reductase (DHDR) and tetrahydrodaidzein reductase (THDR).

Systematic omics analysis for enzyme discovery

To identify suites of candidate genes from *G. glabra* that can be used to reconstitute glabridin biosynthesis, especially for the four classes of tailor enzymes, we used systematic seasonal and organ specific transcriptome, genome mining and co-expression analysis (Fig. 2). We first collected wild perennial samples of *G. glabra.* from natural habitats in Xinjiang of China and performed genome sequencing and assembly (Fig. 2a). We assembled a chromosome-level *G. glabra* genome with a total size of 415 Mb. Genome annotation predicted 34941 protein-coding genes with high-confidence. Using orthologous gene groupings, we constructed a phylogenetic tree comparing *G. glabra* with *G. inflata*, *G. uralensis*, and 10 other legume-related species. The analysis revealed that *G. glabra* shares the closest evolutionary relationship with *G. inflata*. Building on this, we then conducted a systematic genome-wide annotation, with a particular focus on candidate tailoring genes.

To investigate the transcriptional patterns of genes involved in glabridin biosynthesis, we conducted a large-scale transcriptomic study across major *Glycyrrhiza* resource in China. We sampled five *Glycyrrhiza* species, across seven locations, at four key seasonal growth stages, and from different plant organs, resulting in a total of 183 distinct samples. The clustering analysis revealed that the organ-specific transcriptome profiles displayed clear differences, grouping mainly into three clusters: 1) tap roots and horizontal roots, 2) stems and sprouts, 3) leaves. In contrast, the overall transcriptomic differences among species were relatively minor, with variations observed only in certain specific genes. Based on genome-wide alignments using *G. glabra* as the reference, genes that failed to match in *G. inflata* and *G. uralensis* were filtered out. Further clustering of expression patterns across organs and months highlighted key biosynthetic enzymes in *G. glabra*, including 7 PTRs, 18 PTs, 39 OCs, and 6 DMTs. Notably, these genes exhibited chromosomal clustering, primarily located on chromosomes 5 and 6 (Fig. 2b), and are primarily distributed in open and active chromatin regions—areas enriched in genes and transcriptionally active(*19*).

To further identify genes involved in glabridin synthesis, we performed hierarchical clustering to generate co-expressed patterns of transcripts across five organs (Fig. 2c). The clustering revealed a root-enriched gene expression cluster, including 1 PTR, 9 PTs, 8 OCs,

and 2 DMTs. To refine candidate genes, we used previous reported key gene *Gg*I2'H1(*20*) as a bait for co-expression analysis. This analysis prioritized additional candidate enzymes, including 1 PTR, 7 PTs, 2 OCs, and 2 DMTs.

Figure 2 Genomic and transcriptomic analysis of *G. glabra* resources. a) *G. glabra* pseudochromosome genome assembly and gene annotation statistics. b) Genomic distribution of biosynthetic enzyme candidates for glabridin in *G. glabra*. The outermost layer indicates the density of candidate genes across the genome (measured as number of genes per 1Mb). c) The transcriptional profiles of these genes across five tissues of *G. glabra*: tap root, horizontal root, stem, leaf and sprout (121 genes compared across 53 tissue samples). The expression levels are represented as Z-scores derived from $\log_2$-normalized TPM. A specific cluster enriched in root-expressed genes is highlighted in red. d) Co-expression analysis of the *G. glabra* transcriptome using *Gg*I2'H1 as a bait gene. The co-expressed genes were ranked based on their Pearson's correlation coefficient (PCC) with *Gg*I2'H1. Genes shown in black are newly identified glabridin biosynthetic genes (this study). Homologs of the blue-labeled genes (I2'H, IFR, VR, and PTS) were previously identified (*20*). e) Identification of candidate genes for

165 four tailoring enzymes involved in glabridin biosynthesis. The dots represent individual
166 candidate genes. Blue circles indicate all annotated candidates identified through annotation;
167 red circles represent candidates with high expression in root tissues; yellow circles denote
168 genes that co-expressed with GgI2'H1.

Stepwise discovery of Glabridin biosynthetic key enzymes

To characterize the initial tailoring enzyme involved in the glabridin pathway, we first
focused on pterocarpin reductase that catalyzes the reductive cleavage of the dihydrofuran ring
within potential core scaffolds, medicarpin (**8**) and demethylmedicarpin (**12**) (Fig. 3). We
tested the one PTR candidate (GgPTR1) identified through co-expression analysis. This
enzyme catalyzed the conversion of medicarpin (**8**) and demethylmedicarpin (**12**) into vestitol
(**9**) and demethylvestitol (**13**), with conversion rates of 90.6±3.6% and 52.7±0.9%, respectively
(Fig. 3a). This confirms co-expression analysis as an effective strategy for prioritizing
candidate enzymes. To identify more efficient PTRs, we further evaluated the left 6 PTR genes.
Among these, the homolog GgPTR5 exhibited detectable PTR activity but was less active than
GgPTR1, whereas GgPTR2 with a different transcription pattern—highly expressed in the
stems and leaves—also exhibited activity, with conversion rates of 58.8 ± 7.6% for compound
(**8**) and 4.7 ± 2.5% for compound (**12**), respectively (Fig. 3b). In addition, although sharing 71%
sequence similarity, GgPTR2 can convert **25** to 4'-O-Methylpreglabridin (**10**), whereas
GgPTR1 showed no activity, indicating selective recognition of prenylated substrates by
GgPTR2 (Fig. 3a).

To further explore the functional divergence, we modeled their structures in complex with
medicarpin (**8**) and licoagrocarpin (**25**) (Fig. 3c). The electrostatic potential maps(*21*) indicated
that both enzymes possess a positively charged catalytic pocket, attracting the electronegative
phenolic hydroxyl groups of the substrates. Additionally, GgPTR2 exhibited a similar binding
orientation for both **8** and **25**, consistent with the binding of **8** in GgPTR1. In contrast,
compound **25** adopted a 180-degree flipped orientation in GgPTR1, which may explain its lack
of catalytic activity toward this substrate (Fig. 3c).

Next, to identify prenyltransferases of *G. glabra*, we tested the 7 PT candidates and found
GgPT1 exhibited detectable activity, catalyzing the conversion of compounds **9** and **13** into 4'-
O-methylpreglabridin (**10**) and preglabridin (**14**) (Fig. 3d). Other PT genes, despite sharing a
certain degree of phylogenetic similarity with GgPT1, showed no detectable catalytic activity
toward the tested substrates (Fig. 3d). In addition, a reported PcM4DT that catalyzes the
production of **10** from **25**(*22*).

After specific prenylation, intramolecular cyclization is required to form the characteristic cyclic ether structure(*23-26*). We then identified the two genes, *Gg*OC1 and *Gg*OC2. Interestingly, the *GgOC1* exhibited an asymmetric co-expression pattern: the expression of *OC* was highly dependent on the transcriptional activity of the upstream genes(i.e. *GgI2'H1*), whereas the upstream genes showed relatively weak association with *OC* expression. This strongly suggests that the *Gg*OC1 is most likely to participate in glabridin biosynthesis. Indeed, functional assays of the two OCs demonstrated that only *Gg*OC1 catalyzes the oxidative cyclization of 4'-O-methylpreglabridin (**10**) and preglabridin (**14**), yielding 4'-O-methylglabridin (**11**) and glabridin (**15**) (Fig. 3e). Further analysis revealed that *Gg*OC1 belongs to the BBE-type enzyme family and clusters with previously reported *Ma*OC1 in phylogenetic tree, which is responsible for the oxidative cyclization of prenylated chalcones(*27*) .

Next, to find the DMTs in licorice species to realize the demethylation tailoring, we identified a gene ranked second, *Gg*DMT1 from the 2 candidates. It encodes a 2-oxoglutarate-dependent dioxygenase and reveals the ability to completely realize the demethylation of compound **11** (Fig. 3f). Interestingly, when checking the substrate profiling, we found *Gg*DMT1 shows high promiscuity and efficiently demethylates both vestitol (**9**) and 4'-O-methylpreglabridin (**10**). It also shows moderate activity toward formononetin (**5**), but displays very weak on medicarpin (**8**) (Fig. 3f). To further expand the enzyme toolbox for specific 4'-O-demethylation, we screened a panel of previously reported demethylases, including cytochrome P450s enzymes(*28-32*) from plant, fungal and human. Among these, only the fungal enzyme *Nh*PDA1 displayed demethylation activity, displaying promiscuity as well across toward multiple 4'-O-methylated substrates (**5**, **8**, **9**, **10**, **15**) (Fig. 3f). These observations are consistent with prior reports that demethylases often exhibit broad substrate specificity(*27, 32, 33*).

Given that *Gg*DMT1, originating from *G. glabra*, exhibits strong substrate promiscuity and exceptionally high catalytic activity, it raises the question of why methylated intermediates still accumulate in this specie. Analysis of seasonal transcriptional patterns revealed an opposite trend between O-methyltransferase and demethylase expression, suggesting that precursor synthesis and final product accumulation may occur in temporally distinct stages. Furthermore, subcellular localization studies showed that *Gg*DMT1 is cytoplasmic, whereas *Gg*OC1 and *Gg*PT1 are more like a membrane-associated proteins (Fig. 3g). This spatial separation implies that the four tailoring enzymes may be compartmentalized—potentially within specific

organelles—thereby facilitating intermediate segregation and stepwise regulation of the biosynthetic pathway(*34-36*).

Additionally, we found that throughout the pathway, methylation occurs first on the same hydroxyl group, followed by demethylation. This raised the question of whether such complexity is essential for pathway. When we characterized alternative routes that bypass methylation reactions, we found that the I2'H1–IFR1–VR1 module is capable of catalyzing the conversion from intermediate **4** to **12**. However, its catalytic efficiency was extremely low. These results suggest that methylation tailoring effectively blocks the non-methylated route and instead directs metabolic flux toward the more efficient methylated pathway.

Interestingly, when we examined the presence of these key enzymes in the other two *Glycyrrhiza* species, we found that both harbor homologous genes with >93% protein sequence identity, and their genomic loci are relatively conserved (Fig. 3h). We also analyzed the transcriptional differences of the four key enzymes across the three species and found that the transcription level of *PT* in *G. glabra* is significantly higher than in the other two species. *PTR*, *OC and DMT* also show significantly higher transcript levels in *G. glabra* compared to *G. inflata* (Fig. 3i). These findings help explain the significantly higher accumulation of glabridin in *G. glabra*, as observed in our metabolomic analyses (Fig. 1b). This may result from evolutionary changes in unknown regulatory networks that have activated the robust expression of glabridin biosynthetic genes in *G. glabra*, enabling the efficient production and accumulation of this bioactive compound.

Figure 3 Characterization of the four tailoring enzymes. a) Catalytic activity characterizations of PTRs. The tissue-specific expression heatmap demonstrates the transcriptome data from tap root, horizontal root, stem, and leaf. Expression levels are log2-transformed and normalized as Z-scores. b）Functional characterization of PTRs in catalyzing the conversion of prenylated pterocarpan substrate (**25**). c) Structural modeling of *Gg*PTR1 and *Gg*PTR2 in complex with substrates (**5 and 25**) using AlphaFold3(*37*). Boxed regions highlight major differences in the catalytic pocket. Electrostatic potential calculated using

PyMOL APBS tools. Connolly surface shown within ±5 eV. Functional characterization and phylogenetic analysis of d) PTs and e) OCs. f) Substrate profiling of DMTs in catalyzing the demethylation of intermediate compounds. g) Yeast confocal microscopy showing subcellular location of mCherry-*Gg*PT1, *Gg*OC1-mCherry, *Gg*PTR1-mCherry, *Nh*PDA1-mCherry and *Gg*DMT1-mCherry. Scale bar, 5 μm. Images are representative of three independent experiments. Comparison of h) protein sequence homology and i) transcriptional expression levels of *Gg*PT1, *Gg*OC1, *Gg*PTR1, and *Gg*DMT1 across three *Glycyrrhiza* species (*G. glabra,* n=26; *G. inflata*, n=12; *G. uralensis,* n=39) in roots. Box plots in i.: lines are median, box limits are quartiles 1 and 3, whiskers are 1.5×interquartile range and points are outliers. The bar chart shows the molar conversion rates in a., d., e., and f.. Functional assays were performed by feeding 20 mg/L substrate in a. and 50 mg/L substrate in d., e., and f., and conversion rates were determined based on the relative peak areas of the corresponding products. Data represent the mean of at least three biological replicates.

De novo biosynthesis of Glabridin in engineered yeast

To realize the biosynthesis of glabridin in yeast from the simple sugar glucose for the future industrilaztion, we designed the pathway with two modules, core scaffold module (glucose to **8**) and tailoring module (**8** to **15**) (Fig. 4a). We began by assembling the tailoring module using key enzymes we had characterized. Given that all involved enzymes-including PTR, PT, OC, and DMT-exhibit broad substrate promiscuity, we reconstructed the metabolic landscape from compound **8** to **15** and discovered a complex, interconnected tailoring network. This network supports at least two major tailoring routes: PT-PTR-OC and PTR-PT-OC. We then explored different enzyme combinations to further unravel this "tailoring maze", aiming to realize glabridin production.

We first tested the *Pc*M4DT-*Gg*PTR2-*Gg*OC1, which proceed as expected catalytic sequence: PT-PTR-OC, and continuous convert **8** to **11**, or **12** to **15** (Fig. 4b). Due to the promiscuity of *Gg*PTR2, side products such as **9** and **13** were also detected. In this design, we observed significant accumulation of intermediate **8**, indicating that the prenyltransferase reaction is inefficient—likely due to low the prenyl donor DMAPP. To address this, we dynamically regulated *ERG20* gene and overexpressed key rate-limiting enzymes (*tHMG1*, *IDI1*) for regulation of DMAPP supply.

In an alternative configuration PTR-PT-OC, we combined *Gg*PTR1-*Gg*PT1-*Gg*OC1, and when paired with a demethylation step, successfully reconstructed a ladder-like metabolic network. In this architecture, the shared enzymes-PTR, PT, and OC-act on different intermediates along two parallel metabolic routes, which are further interconnected by DMT,

resulting in a flexible and interwoven biosynthetic network. Notably, the choice of demethylase
significantly influenced the network's behavior. Using *Nh*PDA1, a moderately active
demethylase, the metabolic flux was distributed across multiple routes, forming a multi-route
architecture in which various intermediates converge toward glabridin as the final product (Fig.
4d). This metabolism pattern is very close to that we detected in *G. glabra* (Fig. 1a-b),
indicating *G. glabra* may contain a multi-route architecture as well. Moreover, we observed
that flux preferentially flowed toward compound **10**, while conversion to **13** remained low.
After 48 hours, the demethylation of **10** became dominant, initiating glabridin production. This
setup yielded 0.9 mg/L of glabridin (Fig. 4d). However, compound **14** accumulated over time,
indicating a bottleneck at oxidative cyclase (OC) step. In contrast, *Gg*DMT1, with its
exceptionally high demethylation activity, drives the flux predominantly from compound **9** to
13, effectively creating a streamlined, single-route pathway. In this mode, **10, 11** and **12** were
not observed. When checking yield of glabridin, surprisingly, the simplified single-route design
achieved much lower productivity than the complex, multi-route "ladder" network (Fig. 4),
challenging the conventional view that flux distribution in multi-route systems reduces overall
efficiency.
To investigate this phenomenon, we deeply analyzed metabolism dynamics of this network
and observed intermediates secretion during fermentation (Fig. 4f). Specifically, prenylation
reduces extracellular secretion, potentially due to increased hydrophobicity, while
demethylation promotes secretion into the medium. These observations suggest that both
methylation-demethylation and prenylation act as self-retention strategies, helping to retain
intermediates within the cell and minimizing loss through secretion. Among intermediates, **13**
as a key metabolic node, demonstrates highest extracellular ratio, reaching 88.4±4.2%
compared with the others. This extensive leakage explains the lower overall yield observed in
the simplified single-route design. In contract, despite with the strong efflux of **9** and **13**, multi-
route network maintains higher level of robustness that the single one, suggesting that the
inherent redundancy and interconnectedness of the multi-route network provide robustness. To
validate these observations, we built a metabolic flux model, which confirms the experimental
trends and further support the network's behavior (Fig. 4g). Interestingly, this model also
indicated that shifting the demethylation step to the end of the pathway could substantially
improve glabridin yield, offering a promising strategy for optimizing *de novo* biosynthesis (Fig.
4h, S9b-c).
To achieve *de novo* biosynthesis of glabridin in *S. cerevisiae*, 14 enzymes were introduced
to establish the core scaffold module (glucose to compound **8**), enabling the stepwise

biosynthesis of liquiritigenin, formononetin and ultimately medicarpin. Specifically, seven genes-*Gg*PAL1, *Gg*C4H1, *Gg*4CL1, *Gg*CHS1, *Gg*CHR1, *Gg*CHI1 and *Sts*TAL-were incorporated to drive liquiritigenin biosynthesis from glucose. Liquiritigenin was subsequently converted to formononetin via three additional enzymes: *Gg*IFS1, *Gg*HID1, and *Gg*OMT1. To complete the formation of the core scaffold medicarpin, four more enzymes-*Gg*I2'H1, *Gg*IFR1, *Gg*VR1, and *Gg*PTS1-were expressed in yeast. Following establishment of the core pathway, the tailoring module was introduced, with *Gg*DMT1 expression placed under the control of the cyanamide-inducible DDI2 promoter. In addition, to relieve limitations in the de novo biosynthesis of glabridin, the rate-limiting enzymes were overexpressed and NADPH supply were enhanced. Finally, this fully engineered yeast strain enabled *de novo* production of glabridin in shake-flask fermentation, reaching a final titer of 0.5 mg/L (Fig. 4i).

Figure **4 Glabridin biosynthesis module assembly and de novo production in yeast.** a）

Schematic of the glabridin biosynthetic pathway reconstructed in *S. cerevisiae*. b）Stepwise

reconstitution of the glabridin biosynthetic pathway (PT–PTR–OC) in *S. cerevisiae*. 20 mg/L

(**8**) and 10 mg/L (**12**) were added separately to engineered yeast strains with *Pc*M4DT-

*Gg*PTR2-*Gg*OC1 combination. The squares on the bottom indicate different enzyme

combinations: grey indicates enzymes present in the chassis, blue indicates newly introduced

enzymes, and white indicates enzymes not introduced. c-e) Ladder-like biosynthetic network of glabridin (PTR–PT–OC–DMT). d-e) Metabolic dynamics of the ladder-like glabridin biosynthetic network incorporating with either d) *Nh*PDA1 or e) *Gg*DMT1. f) Extracellular distribution ratio of key intermediates in the ladder-like glabridin biosynthetic network. Compounds detected in the supernatant were considered extracellular, while those in the pellet were considered intracellular. Box plots in f.: lines are median, box limits are quartiles 1 and 3, whiskers are 1.5×interquartile range and points are outliers. g) Metabolic simulation of multi-route biosynthetic network and single-route design. h) Metabolic simulation of induced pathway. i) *De novo* biosynthesis of glabridin in engineered *S. cerevisiae*.

Conclusion

In this study, we systematically mapped the maze-like biosynthetic network of glabridin and successfully reconstructed it in *S. cerevisiae*, achieving *de novo* production from glucose. By identifying and assembling key tailoring enzymes, we revealed a ladder-like network architecture mediated by a set of promiscuous enzymes that enhances pathway robustness through multiple interconnected routes. Unlike classical linear metabolic pathways, the multi-route design tolerates metabolic flux fluctuations(*38, 39*), contributing to improved overall productivity. Our study also provides new insights into the evolutionary adaptation of biosynthetic pathways in *Glycyrrhiza* species. The observed metabolic promiscuity and redundancy may represent an evolutionary strategy to maximize ecological adaptability while minimizing genetic burden, by utilizing a limited number of gene copies to ensure metabolic resilience and flexibility across diverse environmental conditions(*40, 41*).

While we have successfully reconstructed the major biosynthetic routes, several challenges remain for further pathway optimization. Notably, the oxidative cyclization step catalyzed by *Gg*OC1 exhibited low catalytic efficiency, emerging as the rate-limiting step for glabridin biosynthesis. Moreover, although we established two major biosynthetic routes, certain branches of the network (Fig. 1) remain incomplete due to missing or unidentified enzymes. Future efforts to discover and integrate these enzymes could fully realize the potential of the glabridin biosynthetic network, enabling more efficient and versatile production of glabridin and structurally related derivatives.

Overall, our work not only provides a blueprint for glabridin biosynthesis but also highlights a generalizable principle: maze-like, multi-route architectures can endow microbial cell factories with enhanced metabolic resilience, offering a new strategy for the synthesis of complex plant natural products.

Materials and Methods

Chemical standards

All chemical standards used in this study were obtained from commercial sources or custom synthesis. Preglabridin was obtained from TargetMol (Boston, MA, USA), while 4'-O-methylglabridin and 4'-methylpreglabridin were custom-synthesized by Acme Biopharma Co., Ltd. (Shanghai, China). All other chemicals were purchased from YuanYe Biotechnology Co., Ltd. (Shanghai, China).

Retrobiosynthesis analysis

The bio-retrosynthesis algorithm was employed to predict the biosynthetic pathway of glabridin and design its synthesis route in yeast cell factories. First, the molecular fingerprint similarity method (FingerprintSimilarity) from the RDKit package in Python was utilized to identify structural analogs of glabridin among metabolites in the KEGG database (https://www.genome.jp/kegg/pathway.html). Second, the acyclic path search method (shortest_simple_paths function) from NetworkX package in Python was applied to explore potential biosynthetic routes for these structural analogs. The pathway origins were set as both the identified glabridin analogs and 2,808 endogenous metabolites from the *S. cerevisiae* genome-scale metabolic model YEAST8(*42*). Subsequently, RetroRules(*43*) (containing 65,688 reaction rules with diameters = 4 and 6) was implemented for retrosynthetic pathway prediction from glabridin to its structural analogs identified in the previous step. To address combinatorial explosion challenges, Monte Carlo Tree Search (MCTS) was employed to establish a retrosynthetic route from glabridin to its structural analog medicarpin. This foundation was further expanded through human-computer interaction single-step retrosynthetic analysis to develop a comprehensive retrosynthetic network.

Genome sequencing, assembly, and annotation

Samples of *Glycyrrhiza glabra* were collected from the Alar region of Xinjiang, China. A PacBio sequencing library was constructed and sequenced using six SMRT cells on the PacBio Sequel platform (Pacific Biosciences) by BGI Genomics Co., Ltd. The third-generation sequencing data were first corrected using the Canu(*44*) (v2.2), followed by preliminary assembly with the Smartdenovo(*45*) software (v1.0.0,) to obtain initial contig sequences. To further improve assembly quality, the initial contigs were sequentially polished using Racon(*46*) (v1.4.3) for third-generation data correction and Pilon(*47*) (v1.24) for error correction with second-generation sequencing data, resulting in the final preliminary assembly. To remove redundant regions, the genome assembly was processed using the Purge haplotigs(*48*) (v1.1.2)

tool. Genome completeness was assessed using BUSCO(*49-51*) (v5.8.3). A Hi-C sample was sequenced for high-quality data. By integrating Hi-C linkage signals, the contig sequences were oriented, clustered, and anchored to the chromosomal level using Juicer(*52*) (v1.6) and 3d-dna(*53*).(v201008) Illumina and PacBio sequencing reads were mapped on chromosome level genome assembly with Bowtie2 (v2.5.4) and minimap2 (v2.28), respectively, where default parameters were adopted (*54, 55*). Then sequencing coverage was calculated by 'depth' function of Samtools (v1.18) in '-aa' mode and displayed in a Circos plot with TBtools (v2.007) (*56, 57*). Phylogeny relationship of *G. glabra*, *G. inflata*, *G. uralensis* and thirteen neighbor species in Leguminosae were constructed according to profiles of shared and unique orthology genes with OrthoFinder (v2.5.4) by default parameters(*58*).

To annotate the SDR family enzymes identified from licorice, enzymes from the SDR406A family were retrieved using the Uniprot database (https://www.uniprot.org). The BLAST tool (https://blast.ncbi.nlm.nih.gov/Blast.cgi) was used to search for homologous sequences of the target enzymes. From the identified candidates, enzymes with known functions were selected, and a multiple sequence alignment was performed using MUSCLE(*59*) (version 5.1). Phylogenetic tree construction was carried out using IQ-TREE(*60*)(version 2.1.4-beta), applying the Maximum Likelihood (ML) method. The inference of the phylogenetic tree was based on the JTT (Jones-Taylor-Thornton) amino acid substitution model, with 1000 ultrafast bootstrap(*61*) resampling to assess branch support and an additional 1000 approximate likelihood ratio tests to further evaluate branch reliability. The final tree was visualized using online iTOL (https://itol.embl.de), with clusters containing only branches that include the enzymes to be annotated retained. The branch lengths of the tree represent evolutionary distances.

Total RNA isolation, RNA-Seq, and gene expression quantification

Based on the geographical distribution of wild licorice resources, sampling sites were selected in Turpan, Urumqi, Emin, Tumxuk, Aksu in Xinjiang, and Longdong in Gansu. From the same individual plant at each location, distinct samples of tap roots, horizontal roots, stems, leaves and sprouts were collected. To facilitate subsequent metabolite analysis and RNA extraction for transcriptome profiling, the collected tissues were immediately flash-frozen in liquid nitrogen. Aliquots intended for metabolomic analysis were stored on dry ice, while aliquots for transcriptomic analysis were preserved in an RNA stabilization solution. Licorice was chopped and ground by freezing with liquid nitrogen. Total RNA was extracted by using the TranZol™ kit (TransGen Biotech, China) according to the manufacturer's instructions, and was subsequently used to synthesize first-strand complementary DNA (cDNA) with the

TransScript® One-Step gDNA Removal and cDNA Synthesis SuperMix (TransGen Biotech, China). Transcriptome sequencing of various *G. glabra* tissues was performed by BGI Genomics Co., Ltd. (Shenzhen, China). The quality of RNA was assessed by the Agilent 2100 Bioanalyzer, and mRNA was purified using oligomagnetic beads. The reverse-transcribed cDNA library was sequenced using BGI's BGISEQ500RS platform (BGI Genomics Co., Ltd). The raw data and measured parameters of the transcriptome have been stored in Jingweikaiwu database (kw.ibc.tsinghua.edu.cn). For transcriptome analysis, RNA-seq reads were aligned to the genome sequence using Hisat2 (v2.2.1)(*62, 63*). Subsequent gene prediction was performed using GeMoMa(*64, 65*), leveraging genomic and transcriptomic data from a closely related species. The gene set was evaluated for completeness using BUSCO. For transcriptome quantification, transcript abundance was quantified using Stringtie(*66*).

Co-expression analysis of candidate genes

A heatmap was generated to visualize the transcriptional profiles of candidate genes across five tissues of *Glycyrrhiza glabra*—tap roots, horizontal roots, stems, leaves and sprouts. The heatmap was constructed using the R package pheatmap (v1.0.12), with row-wise scaling. Expression levels were shown as Z-scores calculated from $\log_2$-normalized TPM values. Pearson correlation analysis was performed using *Gg*I2'H1 as the bait gene, based on TPM values across 53 *G. glabra* transcriptomes covering the five tissues.

Plasmid construction

Codon-optimized synthetic genes were synthesized by Tsingke. Two classes of plasmids were constructed: functional assay plasmids and pathway integration plasmids. For functional assays, candidate genes were cloned between a galactose-inducible GAL2 promoter and the ECM10 terminator, flanked by 500–800 bp genomic homology arms to enable targeted integration into the yeast genome. All vector components except for the gene insert were synthesized in advance. A pair of inward-facing BsaI restriction sites was placed between the promoter and terminator to allow seamless gene insertion via Golden Gate assembly. Gene fragments (either PCR-amplified or synthesized) were digested with BsaI and ligated using T4 DNA ligase (both from New England Biolabs). Pathway integration plasmids were constructed using the same BsaI-based cloning strategy. In these plasmids, expression cassettes consisted of GAL2 or GAL1 promoters from various species, and native yeast terminators. Each expression cassette was flanked by 200 bp synthetic orthogonal sequences that were manually designed to serve as overlap regions, enabling one-step assembly of multi-gene constructs.

Yeast Strain construction

Yeast genomic modifications were performed using the CRISPR-based method(*67*). DNA fragments intended for genomic integration were PCR-amplified using Q5 DNA polymerase (NEB) and co-transformed with a CRISPR-Cas9 plasmid targeting the desired locus. For yeast transformations, an overnight culture of the parental strain was inoculated into 4 mL of 2×YPD medium in a 24-deep well plate at an initial OD_{600} of 0.2 and incubated at 30 °C with shaking at 400 rpm until OD_{600} reached 1.0. Cells corresponding to 1 OD unit were harvested by centrifugation at 3,000 × g for 2 min and washed with an equal volume of sterile water. The cell pellet was then resuspended in a transformation mixture containing 1,000 ng of DNA fragment, 500 ng of CRISPR-Cas9 plasmid, and 260 µL of 50% PEG3350, 36 µL of 1 M LiOAc, and 10 µL of denatured salmon sperm ssDNA(*68*). The mixture was incubated at 42 °C for 30 min. Following centrifugation at 3,000 × g for 2 min, the cell pellet was resuspended in 600 µL of YPD and recovered at 30 °C for 90 min. Cells were then pelleted again, resuspended in 100 µL of sterile water, and plated onto selective agar plates. Correct integration events were confirmed by colony PCR. Verified clones were used for subsequent engineering steps after curing the CRISPR-Cas9 plasmid. Yeast culture media were obtained from Coolaber (Beijing, China).

Strains and culture conditions

Yeast colonies were initially inoculated in triplicate into 2 mL YPD and grown overnight to saturation at 30 °C and 400 rpm, and then 100 µL transferred into 4 mL deep-well 24-well plates sealed with Breathable Sealing Film (Biotss). Cultures were incubated for 120 h at 30 °C, 400 rpm in a high-speed orbital shaker (ZSZY-88BH; Shanghai Zhichu Instrument Co., Ltd.). After 24 h of incubation, galactose was added to a final concentration of 20 g/L to induce gene expression, along with substrate addition when required.

Metabolomic profiling

Dried roots, stems, and leaves of *Glycyrrhiza glabra* were ground into coarse powder under liquid nitrogen and further disrupted using 0.5 mm and 3 mm zirconia beads in a cryogenic bead mill (Guangzhou Luka Sequencing Instrument Co., Ltd.) at 75 Hz for five cycles (each cycle: 60 s grinding, 30 s pause) to obtain fine plant powder. Accurately weighed samples were extracted three times with 25 mL of methanol and 25 mL of ethyl acetate, respectively, in an ultrasonic bath at 25 °C for 30 min. The combined extracts were concentrated to dryness, re-dissolved in methanol, and filtered through a 0.22 µm membrane. Metabolite profiling was performed using a Thermo Scientific Ultimate 3000 UHPLC system coupled to a Q Exactive Plus Orbitrap high-resolution mass spectrometer (Thermo Fisher Scientific, USA). Chromatographic separation was carried out on a Waters ACQUITY UPLC HSS T3 C18

column (2.1 × 100 mm, 1.8 μm; Waters Corporation, USA), with 0.1% (v/v) formic acid in water as mobile phase A and 0.1% (v/v) formic acid in acetonitrile as mobile phase B. The column temperature was maintained at 35 °C, the flow rate was set to 0.2 mL/min, and the injection volume was 5 μL. The elution was performed using the following gradient: 0.00–10.00 min, 100% B; 10.00–20.00 min, 100–70% B; 20.00–25.00 min, 70–60% B; 25.00–30.00 min, 60–50% B; 30.00–40.00 min, 50–30% B; 40.00–45.00 min, 30–0% B; 45.00–60.00 min, 0% B; 60.00–60.10 min, 0–100% B; 60.10–70.00 min, 100% B. Mass spectrometry was conducted in both positive and negative ionization modes using a heated electrospray ionization (HESI) source. The sheath gas flow rate was set to 40 arbitrary units, auxiliary gas to 15 units, capillary temperature to 320 °C, and auxiliary gas heater temperature to 350 °C. The spray voltages were +3.2 kV (positive mode) and –3.0 kV (negative mode). The resolution was set to 70,000 for MS and 17,500 for MS/MS. Data were acquired in full scan mode over an m/z range of 100–1500. Compound identification was performed using Compound Discoverer 3.3 software, with reference to mzCloud and mzVault databases. The peak value of each detected compound was normalized to the corresponding sample weight and used for metabolomic heatmap visualization.

Analysis of metabolite production

Yeast cultures were centrifuged at 3,000 g for 5 min to separate the cell pellets and supernatants. Cell pellets were resuspended in saturated saline and mixed with 0.5 mm zirconia beads. Cell disruption was performed using a cryogenic bead mill (Guangzhou Luka Sequencing Instrument Co., Ltd.). Ethyl acetate was added to both the disrupted cell suspension and the culture supernatant for metabolite extraction. The organic phase was collected and evaporated to dryness under vacuum using a concentrator (Concentrator plus; Eppendorf). Residues were re-dissolved in methanol and filtered through a Nylon 66 membrane (Bioexploration, Guangdong, China) to obtain samples for subsequent HPLC analysis. Metabolites were analyzed using an Agilent 1260 Infinity Binary HPLC system. Chromatographic separation was performed on an InfinityLab Poroshell 120 EC-C18 column (3.0 × 100 mm, 2.7 μm; Agilent Technologies) with 0.1% (v/v) formic acid in water as mobile phase A and acetonitrile as mobile phase B. The flow rate was set to 0.5 mL/min, with a column temperature of 40 °C and an injection volume of 5 μL. The elution was carried out using the following gradient: 0.00–5.00 min, 10–20% B; 5.00–20.00 min, 20–60% B; 20.00–32.00 min, 60–85% B; 32.00–34.00 min, 85–100% B; 34.00–36.00 min, 100% B; 36.00–38.00 min, 100–10% B; 38.00–40.00 min, 10% B. LC–MS analysis was performed following the same procedure described in the "Metabolomic profiling of licorice" section. The production of

metabolite was purified and characterized by liquid Chromatography-Mass Spectrometry (LC-MS) and structural determination by NMR.

Protein structure modeling and docking

Protein structures and ligand docking were predicted using the local version of AlphaFold 3(*69*). The SMILES notation of the ligands was obtained from the PubChem database (https://pubchem.ncbi.nlm.nih.gov). PyMOL (version 2.5.7, http://www.pymol.org) molecular visualization software was used for further analysis of the ligand-receptor complexes. The APBS Electrostatics feature in PyMOL was applied to calculate the electrostatic potential maps of the proteins, utilizing the pdb2pqr method with a grid spacing of 0.50 Å. The predicted isoprenoid-binding pockets were determined by comparing protein structures and sequence similarities.

Fluorescence image and analysis

Individual colonies of yeast strains transformed with plasmids encoding biosynthetic enzymes fused to fluorescent protein reporters were inoculated into 4 mL YPD medium and grown overnight (~12 h) at 30 °C with shaking at 400 rpm. Overnight cultures were then diluted 1:4 into fresh YPD medium and incubated for an additional 6–8 h at 30 °C, 400 rpm to allow proper folding of slow-maturing fluorescent proteins. Cells were harvested by centrifugation and resuspended in 100 μL of sterile PBS for imaging. For microscopy, 5–10 μL of cell suspension was spotted onto a glass microscope slide and covered with a glass coverslip (Shanghai Titan Scientific Co., Ltd.). Imaging was performed using a Nikon AXR multi-SIM multimodal super-resolution confocal microscope (Nikon Instruments Inc., Japan) equipped with a 100×oil immersion objective. Fluorescence excitation was carried out using 488 nm for the green channel (GFP) and 561 nm for the red channel (mCherry). Images were acquired using NIS-Elements software and processed using ImageJ (v1.54f).

Ladder network dynamics measurement

Yeast colonies were initially inoculated in triplicate into 4 mL YPD medium and grown overnight at 30 °C with shaking at 400 rpm until saturation. Subsequently, 1 mL of each culture was transferred into 20 mL fresh YPD medium in 100-mL shake flasks. Cultures were incubated for 120 h at 30 °C and 250 rpm in a shaking incubator. After 24 h of incubation, galactose was added to a final concentration of 20 g/L to induce gene expression, along with substrate addition when applicable. Samples were collected every 24 h and processed using the same method described above.

Model Framework and Simplifications

A computational modeling approach was employed to describe metabolic networks formed

by multiple promiscuous enzymes and their substrate compounds. To address the inherent

complexity of cellular metabolic environments, essential simplifications were implemented

throughout the simulations. Cellular compartmentalization and compound diffusion rates were

neglected in the model formulation.

The Michaelis-Menten equation was adapted to simulate catalytic processes, accounting for

competitive interactions between: (1) different substrates for shared enzyme active sites, and

(2) multiple enzymes competing for identical substrates. This framework yielded:

589
$$v_{1i} = \frac{V_{max1,i}[S_1]}{K_{m1,i}(1 + \frac{[S_2]}{K_{m2}}) + [S_1] + \Sigma_{j \neq i}[S_1] \cdot \alpha_{ij}}$$

590   Km values for Michaelis-Menten equation were predicted through reported model (*70*).

591   Computational analysis revealed Km $\gg$ [S], where substrate concentrations ([S]) were

592   significantly lower than corresponding Km values. This permitted simplification of the

593   Michaelis-Menten equation to:

594
$$v_{1i} = \frac{V_{max1,i}[S_1]}{K_{m1,i}}$$

To align with laboratory conditions for microbial cell factories, the model assumed constant

cellular biomass and enzyme concentrations. Enzyme concentrations were fixed based on

biomass measurements at 24 hr post-inoculation, coinciding with substrate supplementation

timing in experimental protocols. Additionally, concerning the critical parameter of compound

transport across cellular membranes (influx/efflux) that significantly impacts this reaction

network, we postulate that all compounds achieve rapid equilibrium through passive transport

mechanisms, represented by:

602
$$\frac{d[S_{out}]}{dt} = -k_{trans}([S_{out}] - [S_{in}])$$

where transport rate constants (k_{trans}) were experimentally determined through separate

permeability assays. The resulting system of ordinary differential equations was numerically

solved using LSODA (Low-order SObolov differential equation solver), solve_ivp function in

Python (v3.8.5, scipy 1.10.1).

Quantification and statistical analysis

Data were processed by Excel, GraphPad Prism and R. The significant comparisons of two

different groups are indicated in the graphs statistical analysis was performed using two-tailed

unpaired Student's t-test. P values were given in the figures. The graphs represented means ±
SD unless otherwise indicated, as described in the figure legends.

Acknowledgements

613 This work was supported by the Natural Science Foundation of China (22138006, 22278240,
614 22478032, 22478031).

Author contributions

C.L., H.J., L.Q., and B.L. conceived the project. Z.Z. contributed to experimental design,
planning, and execution. WQ.L. and W.L. performed genome assembly and analysis. B.L.,
W.S., S.R., and L.Z. collected Glycyrrhiza samples. L.Q. and Z.Z. analyzed the transcriptome
data. FZ. M. developed the theoretical model. Q.K. and Z.Z. performed metabolomic analysis.
Y.J. carried out enzyme structural modeling. WQ.L. and Y.J. performed enzyme kinetic
simulations. All authors contributed to data analysis and interpretation of results. H.J. and Z.Z.
drafted the manuscript. H.J., L.Q., B.L., and C.L. revised the manuscript and supervised the
study.

Competing interests

625 The authors declare no competing interests.

References

1.   C. T. Walsh, Tailoring enzyme strategies and functional groups in biosynthetic
pathways. *Natural product reports* **40**, 326-386 (2023).
2.   R. S. Nett *et al.*, Plant carbonic anhydrase-like enzymes in neuroactive alkaloid
biosynthesis. *Nature* **624**, 182-191 (2023).
3.   R. De La Peña *et al.*, Complex scaffold remodeling in plant triterpene biosynthesis.
Science **379**, 361-368 (2023).
4.   O. Robles, D. Romo, Chemo-and site-selective derivatizations of natural products
enabling biological studies. *Natural product reports* **31**, 318-334 (2014).
5.   S. Wang, S. Alseekh, A. R. Fernie, J. Luo, The Structure and Function of Major Plant
Metabolite Modifications. *Molecular Plant* **12**, 899-919 (2019).
6.   W. Kreis, J. Munkert, Exploiting enzyme promiscuity to shape plant specialized
metabolism. *Journal of experimental botany* **70**, 1435-1445 (2019).

7.   R. A. Notebaart, B. Kintses, A. M. Feist, B. Papp, Underground metabolism: network-
level perspective and biotechnological potential. *Current Opinion in Biotechnology* **49**,
108-114 (2018).

8.   H. Dvora, M. Koffas, in *Microbial production of food ingredients, enzymes and*
nutraceuticals. (Elsevier, 2013), pp. 234-261.

9.   J. Zhang *et al.*, Review on the diverse biological effects of glabridin. *Drug design,*
development and therapy, 15-37 (2023).

10. C. Simmler, G. F. Pauli, S.-N. Chen, Phytochemistry and biological properties of
glabridin. *Fitoterapia* **90**, 160-184 (2013).

11. W.-H. Ji *et al.*, Total Synthesis of (±)-Glabridin. *Synthetic Communications* **44**, 540-
546 (2014).

12. T. Duigou, M. Du Lac, P. Carbonell, J.-L. Faulon, RetroRules: a database of reaction
rules for engineering biology. *Nucleic acids research* **47**, D1229-D1235 (2019).

13. A. Sveshnikova, H. MohammadiPeyhani, V. Hatzimanikatis, ARBRE: Computational
resource to predict pathways towards industrially important aromatic compounds.
Metabolic Engineering **72**, 259-274 (2022).

14. M. Koch, T. Duigou, J.-L. Faulon, Reinforcement Learning for Bioretrosynthesis. *ACS*
Synthetic Biology **9**, 157-168 (2020).

15. W. Song *et al.*, Biosynthesis-Based Quantitative Analysis of 151 Secondary
Metabolites of Licorice To Differentiate Medicinal Glycyrrhiza Species and Their
Hybrids. *Analytical Chemistry* **89**, 3146-3153 (2017).

16. X. Wang *et al.*, Crystal Structure of Isoflavone Reductase from Alfalfa (Medicago
sativa L.). *Journal of Molecular Biology* **358**, 1341-1352 (2006).

17. H. Shao, R. A. Dixon, X. Wang, Crystal Structure of Vestitone Reductase from Alfalfa
(Medicago sativa L.). *Journal of Molecular Biology* **369**, 265-276 (2007).

18. T. Akashi, S. Koshimizu, T. Aoki, S.-i. Ayabe, Identification of cDNAs encoding
pterocarpan reductase involved in isoflavan phytoalexin biosynthesis in Lotus
japonicus by EST mining. *FEBS Letters* **580**, 5666-5670 (2006).

19. E. Lieberman-Aiden *et al.*, Comprehensive mapping of long-range interactions reveals
folding principles of the human genome. *Science* **326**, 289-293 (2009).

20. C. Lu *et al.*, Heterologous biosynthesis of medicarpin using engineered *Saccharomyces*
cerevisiae. *Synthetic and Systems Biotechnology* **8**, 749-756 (2023).

21. E. Jurrus *et al.*, Improvements to the APBS biomolecular solvation software suite.
Protein Science **27**, 112-128 (2018).

22. J. He *et al.*, Regio-specific prenylation of pterocarpans by a membrane-bound
prenyltransferase from Psoralea corylifolia. *Organic & Biomolecular Chemistry* **16**,
6760-6766 (2018).

23. B.-B. He *et al.*, Enzymatic Pyran Formation Involved in Xiamenmycin Biosynthesis.
ACS Catalysis **9**, 5391-5399 (2019).

24. X.-C. Huang *et al.*, The gradual establishment of complex coumarin biosynthetic
pathway in Apiaceae. *Nature Communications* **15**, 6864 (2024).

25. J.-l. Zou *et al.*, Complete biosynthetic pathway of furochromones in *Saposhnikovia*
divaricata and its evolutionary mechanism in Apiaceae plants. *Nature Communications*
16, 3109 (2025).

26. T. Akashi, K. Uchida, T. Aoki, CYP71D8 and CYP82A2 catalyze the last committed
step in biosynthesis of glyceollin isomers in soybean. *Plant Biotechnology* **42**, 51-56
(2025).

27. Q. Ding *et al.*, The evolutionary origin of naturally occurring intermolecular Diels-
Alderases from Morus alba. *Nature Communications* **15**, 2492 (2024).

28. A. Nag *et al.*, Spatial transcriptional dynamics of geographically separated genotypes
revealed key regulators of podophyllotoxin biosynthesis in Podophyllum hexandrum.
Industrial Crops and Products **147**, 112247 (2020).

29. N. Ikezawa, K. Iwasa, F. Sato, Molecular Cloning and Characterization of CYP80G2,
a Cytochrome P450 That Catalyzes an Intramolecular C–C Phenol Coupling of (S)-
Reticuline in Magnoflorine Biosynthesis, from Cultured Coptis japonica Cells. *Journal*
of Biological Chemistry **283**, 8810-8821 (2008).

30. S. Qiu *et al.*, Functional evolution and diversification of CYP82D subfamily members
have shaped flavonoid diversification in the genus *Scutellaria*. *Plant Communications*
6, (2025).

31. A. P. Maloney, H. D. VanEtten, A gene from the fungal plant pathogen *Nectria*
haematococca that encodes the phytoalexin-detoxifying enzyme pisatin demethylase
defines a new cytochrome P450 family. *Molecular and General Genetics MGG* **243**,
506-514 (1994).

32. W. H. Tolleson, D. R. Doerge, M. I. Churchwell, M. M. Marques, D. W. Roberts,
Metabolism of biochanin A and formononetin by human liver microsomes in vitro.
Journal of agricultural and food chemistry **50**, 4783-4790 (2002).

33.   H. Nagayoshi *et al.*, Preference for O-demethylation reactions in the oxidation of 2′-,
3′-, and 4′-methoxyflavones by human cytochrome P450 enzymes. *Xenobiotica* **50**,
1158-1169 (2020).

34.   L. Bar-Peled, N. Kory, Principles and functions of metabolic compartmentalization.
Nature Metabolism **4**, 1232-1244 (2022).

35.   H. C. Kistler, K. Broz, Cellular compartmentalization of secondary metabolism. *Front*
Microbiol **6**, 68 (2015).

36.   L. V. Roze, A. Chanda, J. E. Linz, Compartmentalization and molecular traffic in
secondary metabolism: a new understanding of established cellular processes. *Fungal*
Genet Biol **48**, 35-48 (2011).

37.   J. Abramson *et al.*, Accurate structure prediction of biomolecular interactions with
AlphaFold 3. *Nature* **630**, 493-500 (2024).

38.   F. Liang *et al.*, Elucidation of the final steps in Taxol biosynthesis and its
biotechnological production. *Nature Synthesis*, (2025).

39.   S. M. Narasimha, T. Malpani, O. S. Mohite, J. S. Nath, K. Raman, Understanding flux
switching in metabolic networks through an analysis of synthetic lethals. *npj Systems*
Biology and Applications **10**, 104 (2024).

40.   M. E. Glasner, D. P. Truong, B. C. Morse, How enzyme promiscuity and horizontal
gene transfer contribute to metabolic innovation. *The FEBS journal* **287**, 1323-1342
(2020).

41.   G. I. Guzmán *et al.*, Enzyme promiscuity shapes adaptation to novel growth substrates.
Molecular Systems Biology **15**, e8462 (2019).

42.   H. Z. Lu *et al.*, A consensus *S. cerevisiae* metabolic model Yeast8 and its ecosystem
for comprehensively probing cellular metabolism. *Nature Communications* **10**, (2019).

43.   T. Duigou, M. du Lac, P. Carbonell, J. L. Faulon, RetroRules: a database of reaction
rules for engineering biology. *Nucleic Acids Research* **47**, D1229-D1235 (2019).

44.   S. Koren *et al.*, Canu: scalable and accurate long-read assembly via adaptive k-mer
weighting and repeat separation. *Genome Research* **27**, 722-736 (2017).

45.   H. Liu, S. Wu, A. Li, J. Ruan, SMARTdenovo: a de novo assembler using long noisy
reads. *GigaByte (Hong Kong, China)* **2021**, gigabyte15 (2021).

46.   R. Vaser, I. Sović, N. Nagarajan, M. Šikić, Fast and accurate de novo genome assembly
from long uncorrected reads. *Genome Research* **27**, 737-746 (2017).

47.   B. J. Walker *et al.*, Pilon: An Integrated Tool for Comprehensive Microbial Variant
Detection and Genome Assembly Improvement. *PLOS ONE* **9**, e112963 (2014).

48.   M. J. Roach, S. A. Schmidt, A. R. Borneman, Purge Haplotigs: allelic contig
740   reassignment for third-gen diploid genome assemblies. *BMC Bioinformatics* **19**, 460
741   (2018).

742   49.   M. Manni, M. R. Berkeley, M. Seppey, F. A. Simão, E. M. Zdobnov, BUSCO Update:
743   Novel and Streamlined Workflows along with Broader and Deeper Phylogenetic
744   Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes. *Molecular*
745   *Biology and Evolution* **38**, 4647-4654 (2021).

746   50.   M. Manni, M. R. Berkeley, M. Seppey, E. M. Zdobnov, BUSCO: Assessing Genomic
747   Data Quality and Beyond. *Current Protocols* **1**, e323 (2021).

748   51.   F. A. Simão, R. M. Waterhouse, P. Ioannidis, E. V. Kriventseva, E. M. Zdobnov,
749   BUSCO: assessing genome assembly and annotation completeness with single-copy
750   orthologs. *Bioinformatics* **31**, 3210-3212 (2015).

751   52.   N. C. Durand *et al.*, Juicer Provides a One-Click System for Analyzing Loop-
752   Resolution Hi-C Experiments. *Cell Systems* **3**, 95-98 (2016).

753   53.   O. Dudchenko *et al.*, De novo assembly of the Aedes aegypti genome using Hi-C yields
754   chromosome-length scaffolds. *Science (New York, N.Y.)* **356**, 92-95 (2017).

755   54.   B. Langmead, S. L. Salzberg, Fast gapped-read alignment with Bowtie 2. *Nature*
756   *Methods* **9**, 357-U354 (2012).

757   55.   H. Li, New strategies to improve minimap2 alignment accuracy. *Bioinformatics* **37**,
758   4572-4574 (2021).

759   56.   C. J. Chen *et al.*, TBtools-II: A "one for all, all for one"bioinformatics platform for
760   biological big-data mining. *Molecular Plant* **16**, 1733-1742 (2023).

761   57.   P. Danecek *et al.*, Twelve years of SAMtools and BCFtools. *Gigascience* **10**, (2021).

762   58.   D. M. Emms, S. Kelly, OrthoFinder: phylogenetic orthology inference for comparative
763   genomics. *Genome Biology* **20**, (2019).

764   59.   R. C. Edgar, High-accuracy alignment ensembles enable unbiased assessments of
765   sequence homology and phylogeny. *bioRxiv*, 2021.2006.2020.449169 (2022).

766   60.   B. Q. Minh *et al.*, IQ-TREE 2: New Models and Efficient Methods for Phylogenetic
767   Inference in the Genomic Era. *Molecular Biology and Evolution* **37**, 1530-1534 (2020).

768   61.   D. T. Hoang, O. Chernomor, A. von Haeseler, B. Q. Minh, L. S. Vinh, UFBoot2:
769   Improving the Ultrafast Bootstrap Approximation. *Molecular Biology and Evolution*
770   **35**, 518-522 (2017).

62. D. Kim, J. M. Paggi, C. Park, C. Bennett, S. L. Salzberg, Graph-Based Genome
Alignment and Genotyping with HISAT2 and HISAT-genotype. *Nature biotechnology*
37, 907-915 (2019).

63. D. Kim, B. Langmead, S. L. Salzberg, HISAT: a fast spliced aligner with low memory
requirements. *Nature methods* **12**, 357-360 (2015).

64. J. Keilwagen, F. Hartung, J. Grau, GeMoMa: Homology-Based Gene Prediction
Utilizing Intron Position Conservation and RNA-seq Data. *Methods in Molecular*
Biology (Clifton, N.J.) **1962**, 161-177 (2019).

65. J. Keilwagen *et al.*, Using intron position conservation for homology-based gene
prediction. *Nucleic Acids Research* **44**, e89 (2016).

66. M. Pertea *et al.*, StringTie enables improved reconstruction of a transcriptome from
RNA-seq reads. *Nature Biotechnology* **33**, 290-295 (2015).

67. Y. Zhang *et al.*, A gRNA-tRNA array for CRISPR-Cas9 based rapid multiplexed
genome editing in *Saccharomyces cerevisiae. Nature Communications* **10**, 1053 (2019).

68. R. Gietz, R. Schiestl, High-efficiency yeast transformation using the LiAc/SS carrier
DNA/PEG method. *Nature protocols* **2**, 31-34 (2007).

69. J. Abramson *et al.*, Accurate structure prediction of biomolecular interactions with
AlphaFold 3. *Nature* **630**, 493-500 (2024).

70. A. Kroll, M. K. M. Engqvist, D. Heckmann, M. J. Lercher, Deep learning allows
genome-scale prediction of Michaelis constants from structural features. *Plos Biology*
19, (2021).

795
796
797