Title: Targeting Siglec-10/α3β1 Integrin Interactions Enhances Macrophage-Mediated Phagocytosis of Pancreatic Cancer

Running Title: Role of Siglec-10 in Immune Evasion in Pancreatic Cancer

Author list: Pratima Saini[1, 2, 3], Gauri Mirji[4], S M Shamsul Islam[1, 2, 3], Lacy M. Simons[1], Sajad Ahmed Bhat[4], Amanda P. Bonfanti[1,2,3], Kar Muthumani[4], Priyesh Agrawal[5], Joel Cassel[4], Hsin-Yao Tang[4], Hiroaki Tateno[6], Rugang Zhang[7], Judd F. Hultquist[1], Rahul S. Shinde[4], Mohamed Abdel-Mohsen[1,2,3†]

Affiliations: [1]Division of Infectious Diseases, Department of Medicine, Feinberg School of Medicine, Northwestern University, Chicago, Illinois, USA; [2] Center for Human Immunobiology, Department of Medicine, Feinberg School of Medicine, Northwestern University, Chicago, Illinois, USA; [3]Potocsnak Longevity Institute, Department of Medicine, Feinberg School of Medicine, Northwestern University, Chicago, Illinois, USA; [4]The Wistar Institute, Philadelphia, PA, USA; [5] Data Science and Artificial Intelligence Institute, Johns Hopkins University, Baltimore MD, USA; [6]Cellular and Molecular Biotechnology Research Institute, National Institute of Advanced Industrial Science and Technology (AIST), Tsukuba, Ibaraki, Japan; [7]University of Texas MD Anderson Cancer Center, Houston, TX, USA.

[†]Corresponding author: Mohamed Abdel-Mohsen, Ph.D. Associate Professor, Northwestern University; Chicago, IL 60611. Phone: 312-503-4946. Email: mmohsen@northwestern.edu

ABSTRACT

Tumor-associated macrophages (TAMs) in the pancreatic ductal adenocarcinoma (PDAC) tumor microenvironment (TME) exhibit immunosuppressive phenotypes and impaired phagocytic activity, facilitating tumor progression and immune evasion. Here, we identify integrin α3β1, composed of ITGA3 and ITGB1 subunits, as a sialylated glycoprotein ligand for Siglec-10, an inhibitory glyco-immune checkpoint receptor highly expressed on TAMs in PDAC. The interaction between Siglec-10 on TAMs and α3β1 on PDAC cells suppresses macrophage-mediated phagocytosis, thereby promoting immune evasion. Consistently, disrupting Siglec-10 interactions with monoclonal antibodies significantly enhances macrophage phagocytosis of PDAC cells and alleviates myeloid cell-mediated inhibition of T cell proliferation and activation *in vitro*. In both a PDAC xenograft mouse model engrafted with human macrophages and a human Siglec-10 transgenic mouse model, targeting Siglec-10 with monoclonal antibodies reduces PDAC tumor growth. These findings suggest that Siglec-10 interactions are key mediators of TAM-driven immune evasion in PDAC and highlight the therapeutic potential of targeting these interactions to restore anti-tumor immunity.

SIGNIFICANCE

Pancreatic tumor cells exploit integrin α3β1 to engage the immunosuppressive checkpoint receptor Siglec-10 on myeloid cells, driving immune evasion, and antibody-mediated blockade of Siglec-10 restores myeloid cell–mediated anti-PDAC immunity.

KEYWORDS

Pancreatic ductal adenocarcinoma (PDAC); Siglec-10; Integrin α3β1; Macrophage phagocytosis; immunotherapy; Siglecs; Sialic acid; ITGA3; ITGB1.

INTRODUCTION

There is an urgent need to identify novel immunological targets to improve survival rates for patients with pancreatic ductal adenocarcinoma (PDAC). This cancer remains exceptionally challenging, with a five-year survival rate of only 13% (1,2), and it shows substantial resistance to current immunotherapies (3). PDAC's resistance to immunotherapy is thought to arise from multiple mechanisms within the tumor microenvironment (TME). Among these, tumor-associated myeloid cells, including tumor-associated macrophages (TAMs), which represent a major immune infiltrate in the TME, play a critical role in promoting treatment resistance. These myeloid cells are often reprogrammed into dysfunctional, immunosuppressive phenotypes that impair T cell effector functions and fail to phagocytose tumor cells (4-6). However, the specific factors driving this myeloid reprogramming remain poorly defined, hindering the development of effective immunotherapies for PDAC.

One emerging mechanism of myeloid immunosuppression involves alterations in tumor cell glycosylation. Advances in the field of glycobiology have revealed that many tumor cells exploit aberrant glycosylation patterns to evade immune responses (7-10). A particularly common glycomic alteration is hyper-sialylation, the process by which cancer cells upregulate sialic acid-containing glycans (sialoglycans) on their surface (11). This enables these cancer cells to bind members of the Siglec family, a group of mostly inhibitory receptors predominantly expressed on innate immune cells, including myeloid cells. Many Siglecs contain immunoreceptor tyrosine-based inhibitory motifs (ITIMs); their engagement triggers downstream inhibitory signaling cascades via SHP-1 or SHP-2 phosphorylation. These signals promote immunosuppressive phenotypes of myeloid cells (12-18). Thus, Siglecs on myeloid cells function as "glyco-immune checkpoints," analogous to PD-1 on CD8⁺ T cells. Targeting the interactions between Siglecs on myeloid cells and sialic acid on cancer cells therefore holds significant potential for reprograming myeloid cells toward an immune-activated, anti-tumor state.

Multiple studies (19-24) highlight the potential of targeting Siglec-10 interactions to reprogram macrophages and restore anti-tumor immunity. For instance, sialic acid on the glycoprotein CD24, expressed by breast and ovarian cancer cells, binds to Siglec-10 on myeloid cells (20).

This interaction transmits an inhibitory signal that induces myeloid cell immunosuppressive

features and impairs macrophage-mediated phagocytosis of cancer cells (20). In an ovarian

cancer mouse model, blockade of CD24 significantly reduced tumor growth and enhanced

macrophages' phagocytic activity (20). While the roles of other Siglecs (Siglec-7, Siglec-9, and

Siglec-15) (25-31) have been explored in the context of PDAC, Siglec-10's contribution to the

immunosuppressive PDAC TME is understudied, despite its important role in other cancers (20).

In this study, we investigated the role of Siglec-10 interactions in enabling PDAC cells to evade

myeloid-mediated anti-tumor activities. Unlike in ovarian and breast cancers, where CD24 is a

major ligand for Siglec-10, we found that most PDAC cells do not predominantly express CD24.

They do, however, bind Siglec-10, suggesting the presence of an alternative sialic acid-

containing glycoprotein ligand. Using affinity pulldown assays followed by mass spectrometry,

we identified integrin α3β1 (composed of ITGA3 and ITGB1 subunits) as a novel glycoprotein

ligand for Siglec-10 on PDAC cells. These integrins had not previously been implicated in

PDAC immune evasion. We further confirmed that ITGA3 and ITGB1 contribute to PDAC

resistance to macrophage-mediated phagocytosis using *in vitro* immune-killing assays. Given

that Siglec-10 on macrophages interacts with multiple glycoproteins on PDAC cells, we

hypothesized that directly targeting Siglec-10 would be a more effective strategy than blocking

individual ligands. To test this, we developed novel monoclonal antibodies against Siglec-10 and

evaluated their ability to enhance macrophage phagocytosis of PDAC cells and to reverse

myeloid cell-mediated suppression of T cell proliferation and activation. Disrupting Siglec-10

interactions with these antibodies significantly increased macrophage-mediated phagocytosis of

PDAC cells and restored T cell proliferation and activation *in vitro*. Furthermore, these

antibodies reduced tumor burden in both a PDAC xenograft mouse model engrafted with human

macrophages   and   a   transgenic   mouse   model   expressing   human   Siglec-10.

MATERIALS AND METHODS

Single cell analysis of PDAC tissues and TCGA datasets. We analyzed a publicly available single-cell RNA-Seq dataset from human PDAC tissues (32). Data processing and analysis were performed using Bioturing's Talk2Data V4 platform. To explore gene expression in specific cell populations, we utilized the sub-clustering function.

Siglec-10 staining on immune cells. Monocytes isolated from healthy donors were differentiated into either immune-activating or immunosuppressive macrophages. For immune-activating conditions, monocytes were cultured with M-CSF (50 ng/mL) for 6 days, followed by stimulation with LPS (10 ng/mL) and IFN-γ (20 ng/mL) along with M-CSF (50 ng/mL) for 24 hours on day 6. For immunosuppressive differentiation, monocytes were treated with M-CSF (50 ng/mL) for 3 days, then stimulated with M-CSF (50 ng/mL) and IL-4 (20 ng/mL) from day 3 to day 8. Siglec-10 expression was assessed on monocytes at day 0 and on macrophages at their respective time-points. Cells were detached using TrypLE, stained with APC-conjugated anti-Siglec-10 antibody for 30 minutes, and washed twice with PBS. Samples were analyzed on a BD FACS Symphony A3 flow cytometer. Data analysis was performed using FlowJo (RRID:SCR_008520) version 9.

Siglec-10 ligand and CD24 staining on cancer cells. Siglec-10 ligand expression was detected using either a recombinant chimeric protein consisting of the Siglec-10 binding region fused to human IgG Fc domains (Siglec-10-Fc, R&D Systems) or a biotin-tagged Siglec-10 mutant (R119A) (Kactus, catalog# SIG-HM411). To form the Siglec-10-Fc/anti-human IgG-BV421 complex, Siglec-10-Fc (2.5 μg/mL) and BV421-conjugated anti-human IgG were incubated on ice for 1 hour. Cancer cells were detached using TrypLE Express, washed, pelleted, and adjusted to a density of $1 \times 10\square$ cells/mL. The cells were then resuspended in the Siglec-10-Fc/anti-human IgG-BV421 precomplex solution together with CD24-Alexa 488 antibody. To form the Siglec-10 mutant (R119A)-streptavidin conjugate, biotin-tagged Siglec-10 mutant (R119A) (2.5 μg/test) was incubated with APC-streptavidin on ice for 1 hour. Cancer cells were detached using TrypLE Express, washed, pelleted, and adjusted to a density of $5 \times 10\square$ cells/test. The cells were then resuspended in the Siglec-10 mutant (R119A)/streptavidin-APC precomplex solution together with CD24-Alexa 488 antibody. After a 30-minute incubation at 4 □°C, the cells were pelleted by centrifugation at $300 \times g$ for 5 minutes and washed twice. Samples were

analyzed on a BD FACS Symphony A3 cytometer, acquiring at least 30,000 events per sample.

Data analysis was performed using FlowJo v9.

Pull-Down and mass spectrometry analysis of Siglec-10 ligands on PDAC cells. To identify

Siglec-10 ligands, we employed a proximity labeling approach based on the tyramide

radicalization principle (33). Briefly, recombinant Siglec-10-Fc fusion protein (10 μg) was

incubated with an anti-human Fc HRP conjugate (5 μg), a monoclonal antibody against human

Fc tag directly conjugated to horseradish peroxidase (HRP), on ice for 60 minutes to facilitate

complex formation. PDAC cell lines (AsPC-1 (RRID:CVCL_0152), BxPC-3

(RRID:CVCL_0186), MiaPaCa-2 (RRID:CVCL_0428), PANC-1 (RRID:CVCL_0480)) were

incubated with the Siglec-10-HRP complex on ice for 1 hour ($20 \times 10\square$ cells per reaction). After

incubation, cells were washed twice with 10 mL of 140 mM NaCl in 20 mM Tris-HCl buffer (pH

8.0, TBS). The cells were then treated with 10 μM biotin tyramide and 10 mM $H\square O\square$ in TBS at

room temperature for 10 minutes. Following the reaction, cells were washed three times with

TBS and lysed in 200 μL of lysis buffer (50 mM Tris-HCl, pH 8.0, 150 mM NaCl, 1% Nonidet

P-40, 0.5% sodium deoxycholate, 0.1% SDS) supplemented with a protease inhibitor cocktail.

Lysis was performed on ice for 30 minutes. A Siglec-5 control and a secondary antibody-only

control were included for each cell line. The lysates were cleared by centrifugation at $15,000 \times g$

for 30 minutes and used for the purification of biotinylated proteins. Biotinylated proteins were

purified using streptavidin-functionalized paramagnetic beads. One milligram of Dynabeads

Streptavidin-MyOne C1 (Thermo Fisher Scientific) was mixed with 100 μL of cleared lysate and

incubated at room temperature for 1 hour with constant shaking. The beads were collected using

a magnet (DynaMag-2, Thermo Fisher Scientific) and washed extensively with D-PBS

containing 0.1% SDS. The samples were reduced with tris(2-carboxyethyl)phosphine, alkylated

with iodoacetamide, and digested on-bead with trypsin (Promega). Tryptic digests were cleaned

using BioPureSPN C18 spin columns (Nest Group) and analyzed by liquid chromatography-

tandem mass spectrometry (LC-MS/MS) using a 1.5-hour LC gradient on a Thermo Q Exactive

Plus mass spectrometer (34). MS data were searched with full tryptic specificity against the

UniProt human proteome (downloaded on 8/21/2023) and a contaminant database using

MaxQuant 2.4.7.0 (35) (RRID:SCR_014485). Protein and peptide false discovery rates were set

at 1%. A total of 4,044 proteins were identified, which were further filtered based on a two-fold

enrichment of Siglec-10 compared to the Siglec-5 control.

Surface Plasmon Resonance (SPR): SPR experiments were performed to evaluate the binding

of various recombinant proteins to Siglec-10. Recombinant Siglec-10 Fc protein was

immobilized on a protein A/G sensor chip at three different surface densities (~200 response

units [RU], ~600 RU, and ~1000 RU). One flow cell was left blank to serve as a reference

control for non-specific binding to the chip surface. The following recombinant proteins were

tested: CD47 (His tag, Acrobiosystems, Catalog# CD7-H5227), CD59 (His, Avitag,

Acrobiosystems, Catalog# CD9-H82E3), NT5E/CD73 (His, Avitag, Acrobiosystems, Catalog#

CD3-H82E3), ITGB6 (C-Myc/DDK, Origene, Catalog# TP317387), ITGA3 (C-Myc/DDK,

Origene, Catalog# TP320975), and ITGB1 (C-Myc/DDK, Origene, Catalog# TP303818). All test

proteins were desalted into running buffer before analysis. Proteins were tested at two

concentrations (100 nM and 1000 nM), except for ITGA3, which was tested at 30 nM and 300

nM. For each injection, the association phase was set to 120 seconds and the dissociation phase

to 600 seconds, with a flow rate of 25 µL/min. After each binding cycle, the Siglec-10/protein

complex was dissociated from the chip surface using 20 mM glycine (pH 2.0), and fresh Siglec-

10 protein was re-immobilized to maintain consistent binding capacity across injections.

Sialidase treatment and lectin staining. ITGA3 and ITGB1 proteins were treated with Sialidase

A (AdvanceBio Sialidase A; Agilent; catalog# GK80040) following the manufacturer's

instructions. Briefly, 5 µg of ITGA3 and ITGB1 glycoproteins were combined with 14 µL of

deionized water, 4 µL of 5× Reaction Buffer, and 2 µL of Sialidase A. The reaction mixture was

incubated at 37°C for 1 hour. To verify the removal of sialic acid, a lectin microarray platform

was used to profile sialic acid-dependent glycan structures. This array utilizes a panel of

immobilized lectins, each with known glycan structure binding specificity. Both sialidase-

digested and undigested proteins were labeled with Cy3 dye (Sigma-Aldrich) and hybridized to

the lectin microarray. The lectin chips were scanned for fluorescence intensity at each lectin-

coated spot using an evanescent-field fluorescence scanner (Rexxam Co., Ltd.). All samples

were run in triplicate, and the average fluorescence intensity of the triplicates was used for

analysis. Data were normalized using the global normalization method.

CRISPR/Cas9-mediated knockout of ITGA3 in PDAC Cells. Knockout of ITGA3 was

performed using a synthetic guide RNA (gRNA) targeting ITGA3 (Horizon, Catalog# CM-

004571-01; Target Sequence: CCAGGGTACACGATGCAGGT). A non-targeting gRNA served

as a control (catalog# U-009502-01-02). CRISPR ribonucleoprotein (crRNP) complexes were

assembled; 1 µL gRNA was combined with 1 µL tracrRNA and incubated at 37°C for 30

minutes to allow duplex formation. 1.6 µL of Poly-L-glutamic acid sodium salt (100mg/mL) was

added to the duplex solution followed by the slow addition of recombinant Cas9 protein (2 µL).

The final crRNP complex was incubated at 37°C for an additional 15 minutes to complete

assembly. Assembled crRNPs (5.6 µL per reaction) were aliquoted into PCR strip tubes and

stored on ice prior to transfection. PANC-1 cells were prepared at a density of $2 \times 10^{\square}$ cells per

reaction. Electroporation was performed using the Lonza 4D-Nucleofector system with the SE

Cell Line 4D-Nucleofector X Kit (Lonza), following the manufacturer's protocol. A 20 µL

nucleofection buffer was prepared by combining 3.6 µL of SE Supplement with 16.4 µL of SE

Solution. Cells were suspended in nucleofection buffer and mixed with 3.5 µL of the crRNP

complex prior to nucleofection using program DN-100. Immediately following electroporation,

100 µL of pre-warmed complete growth medium was added to each reaction for recovery. Cells

were then transferred to 6-well plates containing 2 mL of medium per well and incubated under

standard conditions for 48 hours. Knockout efficiency was assessed 72 hours post-transfection

by flow cytometry. Cells were stained with an APC-conjugated anti-ITGA3 antibody

(Biolegend; catalog# 343808) and analyzed for surface expression. A reduction in ITGA3 signal

compared to non-targeting control indicated successful gene knockout. Edited cell populations

were maintained in culture and routinely monitored by flow cytometry at regular intervals to

confirm stable loss of ITGA3 expression.

Macrophage differentiation and in vitro phagocytosis assay. Macrophages were differentiated

as described previously (20,36). Briefly, monocytes from healthy donors were differentiated into

macrophages by culturing them for 7–9 days in Iscove's Modified Dulbecco's Medium (IMDM)

supplemented with 10% AB human serum (Life Technologies). During the initial 3-4 days,

macrophages were stimulated with 50 ng/mL M-CSF. Subsequently, M-CSF (50 ng/ml) and IL-4

(20 ng/mL) were added to support differentiation, and this stimulation was maintained until the

macrophages were used on Days 7-9. PDAC cells were treated with TrypLE Express and labeled

with pHrodo Red, SE (Thermo Fisher Scientific) according to the manufacturer's protocol.

Labeling was performed at a concentration of 250 ng/mL in PBS for 10 minutes at 37°C,

followed by two washes with 1X PBS. Healthy donor-derived macrophages were harvested

using TrypLE Express, and 25,000 macrophages were seeded into black 96-well flat-bottom

plates, allowing them to adhere for 30 minutes at 37°C. After adhesion, 25,000 pHrodo Red-

labeled PDAC cells were added in serum-free IMDM. The plates were incubated at 37°C, and

phagocytosis was monitored by imaging every hour using an Incucyte (Sartorius). The initial

image (t = 0) was captured within 30 minutes of co-culture. Phagocytosis was quantified by

measuring the total red area (µm² per well) across technical replicates for each donor. Thresholds

for identifying pHrodo Red-positive events were determined based on intensity measurements

from labeled cells in the absence of macrophages.

ITGA3^{low} and ITGB1^{low} PDAC cell enrichment. PDAC cells were incubated with anti-human

ITGA3 and ITGB1 antibodies (Clone ASC-1 and TS2/16, respectively, from Thermo Fisher

Scientific) at a concentration of 1 µg/mL in 1× PBS at 4°C for 15 minutes. The cells were then

washed twice and incubated with 20 µL of anti-mouse MicroBeads (Miltenyi Biotec, catalog#

130-048-402) for 15 minutes at 4°C. Following another wash, the cells were loaded onto pre-

equilibrated LS columns (Miltenyi Biotec, catalog# 130-042-401) in accordance with the

manufacturer's protocol. After washing, the cells in the eluate fraction were pelleted,

resuspended in serum free IMDM, and subsequently used for in vitro cell-killing assays.

Siglec-10 monoclonal antibody production. Monoclonal antibodies targeting the extracellular

domain (ECD) of Siglec-10 were developed and characterized. Ten BALB/c

(RRID:MGI:2683685) and C57BL/6 (RRID:IMSR_JAX:000664) mice were initially immunized

with 50µg of recombinant Siglec-10 protein (GenScript), followed by four booster

immunizations of 25µg each at two-week intervals. Blood samples were collected after each

immunization, and serum antibody titers were assessed using indirect ELISA. For ELISA, 96-

well plates were coated with Siglec-10 antigen or Siglec-5 antigen (1µg/mL in PBS) and

incubated overnight at 4°C. Plates were blocked with 1% BSA in PBS to prevent nonspecific

binding. Diluted serum samples were incubated for 1 hour at 37°C, followed by washes and

incubation with horseradish peroxidase (HRP)-conjugated goat anti-mouse IgG secondary

antibody. The reaction was developed using tetramethylbenzidine (TMB) substrate, stopped with

acid, and the absorbance was measured at 450 nm. Mice with serum OD450 values exceeding
1.0 at dilutions ≥1:8,000 were selected for hybridoma generation. Splenocytes from selected
mice were fused with SP2/0 myeloma cells via electrofusion. The hybridomas were cultured in
HAT medium, and supernatants were screened for Siglec-10-specific antibodies using same
ELISA protocol described above. Hybridoma clones were also tested for specificity by flow
cytometry (FACS) using CHO-K1 cells (RRID:CVCL_0214) expressing Siglec-10 or Siglec-5
(as a control). Positive hybridoma supernatants demonstrated significant fluorescence with
Siglec-10-expressing cells compared to controls. For FACS, 50 μL of a CHO-K1 cell suspension
($1 \times 10\square$ cells per well) was incubated with hybridoma supernatants at 4°C for 30 minutes. After
washing, Alexa Fluor 488-conjugated goat anti-mouse IgG secondary antibody (Jackson
ImmunoResearch) was added, and samples were analyzed using a flow cytometer. Clones with
high median fluorescence intensity (MFI) against Siglec-10 cells were selected for subcloning.

Selected hybridoma clones underwent limiting dilution for subcloning. Subclones were cultured,
and supernatants were collected for antibody purification. Antibodies were purified using Protein
A chromatography and assessed for purity by SDS-PAGE. Concentrations were determined
using UV spectrophotometry. The specificity of purified antibodies was validated using the same
ELISA and FACS protocols described above against both Siglec-10 and Siglec-5 (as a control).
Antibodies from positive clones were sequenced. RNA was extracted from hybridoma cells
using the RNeasy Isolation Kit, and antibody sequences were determined via RT-PCR using
universal primers. Sequenced antibodies were expressed recombinantly in CHO-S cells and
purified. Functional characterization of these recombinant antibodies was performed using the
same ELISA and FACS assays described above.

Co-culture assays with monocyte-derived macrophages and CD8⁺ T cells. Human CD8⁺ T cells
and monocytes were obtained from the Human Immunology Core at the University of
Pennsylvania. CD8⁺ T cells were labeled with 5$\square$μM CellTrace Violet (Thermo Fisher
Scientific) and co-cultured with autologous monocytes at a 1:2 ratio (T cell:monocyte) for 5 days
in the presence of anti-CD3/CD28 Dynabeads (Thermo Fisher Scientific) for polyclonal T cell
activation. At the end of the culture period, cells were harvested and analyzed using a BD
FACSymphony flow cytometer (BD Biosciences). Proliferation was assessed based on CellTrace
dye dilution. For GranzymeB staining, cells were fixed and permeabilized using True nuclear

transcription factor buffer set (Biolegend, Catalog# 424401), according to the manufacturer's

instructions. Flow cytometry data were analyzed using FlowJo software (v10.6.1, TreeStar), and

statistical analyses and graphical representations were performed using GraphPad Prism

(v10.0.0).

Cancer associated fibroblasts (CAFs), macropahges, and cancer cell triple co-culture assays.

CAFs were cultured in DMEM supplemented with 10% FBS. Macrophages were differentiated

from healthy donor monocytes as previously described. After differentiation, macropahges were

detached using TrypLE Express, and 25,000 cells per well were seeded into a 96-well plate in the

presence of either a Siglec-10 blocking antibody or an isotype control (10 µg/well). TAMs were

allowed to adhere for 1 hour at 37°C. CAFs and PANC-1 cells were detached using TrypLE

Express. PDAC cells were labeled with pHrodo Red as previously mentioned. Once TAMs

adhered, 25,000 pHrodo Red-labeled PDAC cells and 10,000 CAFs were added per well in

serum-free IMDM. Plates were incubated at 37˚C, and phagocytosis was monitored hourly using

an Incucyte live-cell imaging system (Sartorius).

Experiments using xenograft mouse model. NSG (NOD.Cg-Prkdc^{scid} Il2rg^{tm1Wjl}/SzJ) mice

(RRID:BCBC_4611) were anesthetized with isoflurane, and the hair on the right flank was

shaved. A 100 µL suspension of AsPC-1 cells ($2 \times 10\square$ cells per mouse) was prepared by

mixing the cells with Matrigel and $1\times$ PBS at a 1:1 ratio, which was then injected subcutaneously

into each mouse. After one-week, subcutaneous tumors of approximately 40 mm³ had developed.

The mice were randomly assigned to two groups: isotype control or Siglec-10 antibody

treatment. Monocyte-derived macrophages were incubated with 200 µg of either isotype or

Siglec-10 antibody for 30 minutes. Each mouse received intravenous injections of antibody-

incubated macrophages three times per week for three weeks. Tumor size was measured using a

vernier caliper, and tumor volume was calculated using the formula: tumor volume = ½ (length $\times$

width²). All mice were maintained inappropriate environmental conditions. All animal protocols

were approved by Wistar Institutional Animal Care and Use Committee (protocol # 201363).

Euthanasia was performed using carbon dioxide (CO_2) delivered as bottled gas, which is listed

among the acceptable agents for euthanasia of laboratory rodents in the American Veterinary

Medical Association (AVMA) Guidelines for the Euthanasia of Animals: 2013 Edition.

RNA-Seq analysis of macrophages. Subcutaneous tumors were excised from each of the
xenograft mice, and human cells were sorted by FACS. Total RNA was extracted from the sorted
human macrophages using the Single Cell RNA Purification Kit (Norgen). RNA quality was
assessed using the TapeStation High Sensitivity RNA ScreenTape (Agilent). Libraries were
prepared using the Low Input RNA Library Prep Kit (Takara Biosciences) with 500 ng of DNase
I-treated total RNA. Final library quality control was performed using the Bioanalyzer High
Sensitivity DNA Kit (Agilent). Next-generation sequencing (NGS) was conducted with 100 bp
paired-end reads on a NovaSeq 6000 system (Illumina) using the SP v1.5 200-cycle kit. RNA-
seq analysis was performed by Bencos Research Solutions Pvt. Ltd. (Mumbai) using their
proprietary TWINE BI platform. For the analysis, the RNA-seq data were aligned to the hg19
human genome using STAR-aligner (v 2.7.10a), and salmon (v 1.10.1) was used to estimate read
counts and RPKM values. Raw counts were analyzed for differential expression using DESeq2
(RRID:SCR_000154).

Experiments using human Siglec-10 transgenic mouse model. B-hSIGLEC10 mice
[C57BL/6N-(Siglecgtm2(SIGLEC10) Bcgen/Bcgen)] were obtained from Biocytogen (Catalog #
110843). In this genetically engineered strain, exons 3–10 of the murine *Siglec-G* gene, encoding
the extracellular domain, were replaced with the corresponding human *SIGLEC10* exons,
resulting in a humanized extracellular domain. Knockout of mouse Siglec-G and knock-in
expression of human SIGLEC10 were confirmed by flow cytometry analysis using the following
fluorophore-conjugated antibodies: F4/80-BUV805 (BM8, ThermoFisher, 368-4801-80), CD3-
BV510 (OKT3, BioLegend, 317331), MHC-II-BV711 (M5/114.15.2, BioLegend, 107643), Ly-
6C-Pacific Blue (HK1.4, BioLegend, 128013), CD11c-PE-Cy7 (N418, BioLegend, 117317),
CD19-PE-Cy5 (6D5, BioLegend, 115509), mouse Siglec-G-PE (SH1, BioLegend, 163304),
CD45-APC-Cy7 (30-F11, BioLegend, 103115), CD11b-Alexa Fluor 700 (M1/70, BioLegend,
101222), human Siglec-10-APC (5G6, BioLegend, 347606), and the LIVE/DEAD Fixable Blue
Dead Cell Stain (ThermoFisher, L23105). B-hSIGLEC10 mice were anesthetized using
isoflurane, and the hair on the right flank was removed. A 100 µL cell suspension containing 3 ×
10□ KPCY (2838c3) tumor cells was prepared in a 1:1 mixture of Matrigel and 1× PBS. The
suspension was injected subcutaneously into the right flank of each mouse. The cell line was
tested for contamination by Center for Comparative Medicine at Northwestern University. After

5 days, tumors reached an approximate volume of 40 mm³, at which point mice were randomized
into two treatment groups: isotype control or anti-Siglec-10 antibody. Mice received daily
intraperitoneal injections of either Siglec-10 blocking antibody or respective isotype until the
study endpoint. Tumor growth was monitored using vernier calipers, and tumor volume was
calculated using the formula: tumor volume = ½ (length × width²).

Data availability: All data are available within the manuscript and its figures and/or available
upon   request.

RESULTS

Single-cell analysis of the PDAC TME showed that Siglec-10 is predominantly expressed in multiple myeloid cell subtypes. We began our investigation by asking which PDAC TME cells expressed Siglec-10. We analyzed a publicly available single-cell RNA sequencing (scRNAseq) dataset from human PDAC tumors (**Fig. 1a**) (32) and found that Siglec-10 is expressed in several myeloid cell subsets within the PDAC TME. Specifically, 60% of Siglec-10$^+$ cells were macrophages, followed by myeloid-derived suppressor cells (MDSCs) and dendritic cells (DCs) (**Fig. 1b**). Turning to which Siglecs were most highly expressed on these myeloid cells, we found that Siglec-10 had higher expression than the other Siglecs previously implicated in PDAC immune evasion, in particular Siglec-7, -9, and -15 (**Fig. 1c**).

These findings suggest that Siglec-10 expression on TAMs, MDSCs, and DCs may contribute to their immunosuppressive phenotypes. Supporting this hypothesis, Siglec-10☐ cells exhibited elevated expression of genes linked to immunosuppressive pathways, particularly TGF-β signaling (22,37-42), and lower expression of pro-inflammatory and activation-associated genes. Siglec-10☐ macrophages also showed increased expression of SHP-1 and SHP-2, key mediators of Siglec-inhibitory signaling. In DCs, Siglec-10☐ cells exhibited upregulation of genes impairing antigen presentation, such as HLA-DOB and HLA-DOA (**Fig. 1d**). Additionally, Siglec-10 expression within the PDAC TME was associated with more advanced and metastatic disease (**Fig. 1e**).

To assess the clinical relevance of Siglec-10 expression in pancreatic cancer, we analyzed The Cancer Genome Atlas (TCGA; version 1, www.gepia.cancer-pku.cn) data for pancreatic adenocarcinoma (PAAD). PAAD tissues exhibited higher Siglec-10 expression than normal pancreatic tissues (**Fig. 1f**), and elevated Siglec-10 levels correlated with poorer survival outcomes in the Human Protein Atlas dataset (www.proteinatlas.org; **Fig. 1g**). In addition to examining Siglec-10 expression on macrophages, we investigated the potential role of its ligands, sialic acids, in PDAC progression. Sialic acids are added to glycoproteins through the activity of various glycan-related enzymes, including sialyltransferases (**Supplementary Fig. 1a**). We found that multiple genes involved in the synthesis of sialic acid ligands for Siglec-10 were significantly upregulated in PAAD tissues compared to normal tissues (**Supplementary**

Fig. 1b–c), suggesting that Siglec-10–sialic acid interactions may contribute to PDAC
progression.

Together, these data show that Siglec-10, a glyco-immune checkpoint, is highly expressed in the
PDAC TME and is associated with disease progression and poorer prognosis. To validate these
findings at the protein level, we differentiated primary human monocytes into macrophages
using M-CSF, followed by IL-4 or LPS plus IFN-γ to generate immunosuppressive or
immunoactivating phenotypes, respectively. Siglec-10 protein levels, measured by flow
cytometry, were higher on immunosuppressive macrophages than on immunoactivating
macrophages (**Supplementary Fig. 2**). Overall, these results suggest that Siglec-10 expression
on myeloid cells within the PDAC TME is associated with an immunosuppressive phenotype.

PDAC cells bind to Siglec-10 through sialic acid expressed on proteins other than CD24. Our
next goal was to identify the glycoprotein ligand(s) of Siglec-10 on PDAC cells. CD24 was
recently identified as a major glycoprotein ligand of Siglec-10 on breast and ovarian cancer cells
(20). To test whether PDAC cells express CD24 and whether this expression coincides with
Siglec-10 binding, we used a Siglec-10–Fc chimera protein followed by a secondary antibody,
along with anti-CD24 antibodies, to detect Siglec-10 ligands and CD24 expression by flow
cytometry on multiple PDAC cell lines as well as ovarian and breast cancer cells. Multiple
human PDAC cell lines bound Siglec-10, indicating the presence of compatible sialylated
glycoproteins on their surface (**Fig. 2a–f**). However, only a subset of these cells expressed CD24,
with a large population of CD24-negative cells capable of binding Siglec-10 (**Fig. 2a–f**). This
was in contrast to ovarian and breast cancer cells, which expressed high levels of CD24 that
coincided with the presence of Siglec-10 ligands (**Fig. 2i–l**). Importantly, using a Siglec-10
protein containing the inhibitory R119A mutation, which abrogates sialic acid binding, we
confirmed that Siglec-10 binding to PDAC cells is sialic acid–dependent (**Fig. 2**). These data
suggest that PDAC cells engage Siglec-10 on myeloid cells through glycoproteins other than
CD24.

Siglec-10 binds to ITGA3 and ITGB1, the subunits of α3β1 Integrin, on PDAC cells. To

identify glycoproteins on PDAC cells that bind to Siglec-10, we performed a pull-down assay

using recombinant Siglec-10 protein and cell-membrane proteins isolated from several human

PDAC cell lines. Proteins bound to Siglec-10 were identified via mass spectrometry analysis

(**Fig. 3a**). From a total of 4,044 proteins whose binding was enriched compared to a no-protein

control (a control with anti-Fc HRP secondary antibody only), 110 proteins showed enrichment

compared to Siglec-5 protein (a closely related Siglec used as a control). Of these, six proteins,

CD47, CD59, CD73, ITGB6, ITGA3, and ITGB1, were significantly overexpressed in PAAD

tissues compared to normal tissues based on TCGA data (**Fig. 3b**).

To determine whether these six proteins bind directly to Siglec-10, we performed surface

plasmon resonance (SPR) analysis with immobilized recombinant Siglec-10 protein using

recombinant versions of these proteins (produced in human cell lines to maintain their

glycosylation). Among the six candidates, only ITGA3 and ITGB1, the subunits of α3β1

integrin, exhibited strong binding affinity to Siglec-10 (**Fig. 3c**). To assess whether this binding

is dependent on the sialic acid content of these glycoproteins, we treated ITGA3 and ITGB1 with

sialidase to enzymatically remove sialic acid. The removal of sialic acid was confirmed using

lectin staining with a sialic acid–specific lectin (SNA), which showed a significant reduction in

sialic acid levels on ITGA3 and ITGB1 after sialidase treatment (**Fig. 3d**). Sialidase treatment

abolished the ability of ITGA3 and ITGB1 to bind to Siglec-10 (**Fig. 3e-f**). These findings show

that Siglec-10 binds to the sialic acid on ITGA3 and ITGB1.

The expression of ITGA3 and ITGB1 on PDAC cells correlates with disease progression and

prognosis. Having identified ITGA3 and ITGB1 as glycoprotein ligands that engage Siglec-10 to

potentially suppress immune surveillance, we next investigated whether their expression is

associated with PDAC progression and patient outcomes. First, we measured the surface

expression of ITGA3 and ITGB1 on several human PDAC cell lines and observed consistently

high levels across multiple lines (**Supplementary Fig. 3**). To evaluate their *in vivo* relevance, we

analyzed TCGA data comparing gene expression in PAAD tissues to normal pancreatic tissues.

While CD24 levels were similar between PAAD and normal tissues, ITGA3 and ITGB1

expression was significantly higher in PAAD samples (**Fig. 4a–c**). Furthermore, ITGA3 and

ITGB1 expression positively correlated with elevated Siglec-10 levels (**Supplementary Fig.**
4a,c) and with higher expression of glycosylation-related genes such as ST3GAL1 and B3GNT3,
which are responsible for adding sialic acids to glycoproteins (**Supplementary Fig. 4b,d**).

Next, we examined the association of CD24, ITGA3, and ITGB1 expression with PDAC patient
survival. While CD24 expression showed no correlation with survival outcomes (**Fig. 4d**), higher
expression of ITGA3 (**Fig. 4e**) and ITGB1 (**Fig. 4f**) was significantly linked to worse prognosis.
Finally, single-cell expression analyses within the PDAC TME revealed that elevated ITGA3
and ITGB1 expression correlated with more aggressive disease features, whereas CD24
expression did not (**Fig. 4g–i**). Together, these findings support the notion that ITGA3 and
ITGB1 contribute to PDAC progression, potentially by engaging Siglec-10 and facilitating
immune evasion.

ITGA3 and ITGB1 facilitate PDAC cell evasion from macrophage-mediated phagocytosis.
Given that (1) sialic acid on ITGA3 and ITGB1 in PDAC cells binds strongly to Siglec-10, (2)
ITGA3 and ITGB1 expression is associated with PDAC progression and poorer prognosis, and
(3) Siglec-10 interactions on macrophages inhibit their phagocytic activity, we next investigated
whether PDAC cells with high ITGA3 or ITGB1 expression exhibit greater evasion of
macrophage-mediated phagocytosis compared to cells with low expression of these
glycoproteins.

We first employed the PDAC cell line PANC-1 and knocked out ITGA3 using CRISPR-Cas9.
As shown in **Fig. 5a**, this effectively reduced cell-surface ITGA3 expression by over 95%, as
measured by flow cytometry. We then examined whether this reduction enhanced the
susceptibility of PDAC cells to macrophage-mediated phagocytosis. ITGA3-KO cells and non-
targeting gRNA control cells were labeled with pHrodo dye and co-cultured with macrophages
differentiated from monocytes of multiple healthy donors using M-CSF and IL-4 (to generate
immunosuppressive monocyte-derived macrophages). We then assessed phagocytosis using live-
imaging staining over time (**Fig. 5b**). As shown in **Fig. 5c**, ITGA3-KO PDAC cells were
significantly more susceptible to macrophage-mediated phagocytosis than control PDAC cells
with high ITGA3 expression.

To validate these results, we used the PDAC cell lines MIA PaCa-2 and PANC-1 and employed

antibody-coated columns targeting ITGA3 to enrich for ITGA3^{low} cells (cells not bound by the

columns) (**Fig. 5d**). Flow cytometry confirmed a significant enrichment of ITGA3^{low} cells

compared to controls, which consisted predominantly of ITGA3^{high} cells that passed through the

columns. **Fig. 5e** illustrates an example where depletion of ITGA3^{high} cells in MIA PaCa-2

resulted in over a 70% reduction in ITGA3 expression. Consistently and in subsequent

phagocytosis assays (**Fig. 5f**), PDAC cells with high ITGA3 expression evaded macrophage-

mediated phagocytosis more effectively than ITGA3^{low} cells (**Fig. 5g-h**). Similar results were

observed when depleting ITGB1^{high} cells (**Supplementary Fig. 5**). Together, these findings

indicate that ITGA3 and ITGB1 significantly contribute to the immune evasion of PDAC cells

by inhibiting macrophage-mediated phagocytosis.

Disrupting Siglec-10 interactions, via monoclonal antibodies, enhances macrophage-mediated

phagocytosis of PDAC cell and prevent myeloid cell–mediated inhibition of T cell proliferation

and activation in vitro. Our findings thus far suggest that Siglec-10 on macrophages binds to

multiple glycoprotein ligands on PDAC cells, including ITGA3, ITGB1, and potentially others.

This binding inhibits the phagocytic capacity of macrophages against PDAC cells (**Fig. 6a**).

Therefore, directly targeting Siglec-10 itself, rather than its individual ligands, may be necessary

to enhance macrophage-mediated phagocytosis of PDAC cells. Such an approach would disrupt

interactions between Siglec-10 and sialic acids across its various glycoprotein ligands. However,

the lack of potent, commercially available anti-Siglec-10 antibodies has necessitated the

development of these antibodies in-house.

To generate mAbs targeting Siglec-10, mice were immunized with recombinant human Siglec-10

protein, and serum antibody titers were measured using indirect ELISA. High reactivity to

Siglec-10 was observed in several mice, which were subsequently selected for hybridoma

generation. Over 200 hybridoma clones were initially screened for antibody production using

binding specificities by ELISA against Siglec-10 and Siglec-5 (a control). Sixteen clones

(sequences in **supplementary Table 1**) were selected for subcloning. Antibodies from these

clones were further characterized by ELISA and flow cytometry-based immune assays. Several

antibodies exhibited high affinity for Siglec-10 in binding ELISA (**Fig. 6b**). To confirm the

binding specificity of these antibodies to Siglec-10, CHO-K1 cells expressing Siglec-10 or Siglec-5 (as a control) were incubated with hybridoma supernatants and stained with an Alexa Fluor 488-conjugated secondary antibody. The antibodies showed significant binding to Siglec-10-expressing cells, but not to Siglec-5 controls (**Fig. 6c**).

The clones were then evaluated for their ability to enhance macrophage-mediated phagocytosis of PDAC cells using the same assay described in Fig. 5, compared against an isotype control, an anti-CD24 antibody, and a commercially available anti-Siglec-10 antibody (**Fig. 6d**). Several clones significantly improved the capacity of macrophages to phagocytose PDAC cells (**Fig. 6d**). The top five clones were further tested using macrophages derived from monocytes of multiple donors to assess reproducibility. Among these, one clone, 68A11A1, demonstrated superior performance and was selected for recombinant expression (**Fig. 6e–f**). The recombinant 68A11A1 antibody (IgG1 isotype) showed high specificity for Siglec-10 without cross-reactivity to Siglec-5 or other irrelevant His-tagged proteins (**Fig. 6g**). Notably, this recombinant antibody significantly enhanced macrophage-mediated phagocytosis across multiple human PDAC cell lines compared to the anti-CD24 antibody, using macrophages derived from several donors (**Fig. 6h**). It also enhanced phagocytosis by monocyte-derived macrophages against PDAC PANC-1 cells even in the presence of cancer-associated fibroblasts (CAFs) in tri-culture assays with monocyte-derived macrophages, PDAC cells, and CAFs (**Fig. 6i**), and did so without altering the levels of ITGA3, ITGB1, or Siglec-10 ligands on PDAC cells (**Supplementary Fig. 6**).

In addition, we assessed the ability of this antibody to overcome myeloid cell-mediated suppression of T cell responses. In a monocyte–T cell suppression assay, Siglec-10 blockade increased CD8+ T cell proliferation (**Fig. 6j**) and granzyme B expression (**Fig. 6k**) compared to the isotype control. Together, these results demonstrate the successful generation and characterization of monoclonal antibodies with high affinity and specificity for Siglec-10. These antibodies not only enhanced macrophage-mediated phagocytosis of PDAC cells but also prevented myeloid cell-mediated inhibition of T cell proliferation and activation *in vitro*, underscoring the critical role of Siglec-10 in PDAC immune evasion and its potential as a therapeutic target.

Disrupting Siglec-10 interactions enhances macrophage-mediated phagocytosis of PDAC cells
and induces human macrophage activation in vivo. Since mice lack Siglec-10 and instead
express its homolog Siglec-G, testing a Siglec-10 antibody in wild-type mice is not feasible. To
overcome this limitation, we employed two complementary animal models. First, we used a
xenograft model: immunodeficient NSG mice engrafted with AsPC-1 human PDAC cells and
human monocyte-derived macrophages. Mice were treated with either an isotype control
antibody or the Siglec-10-specific antibody clone 68A11A1 (**Fig. 7a**). Targeting Siglec-10
significantly reduced PDAC tumor growth *in vivo* (**Fig. 7b**).

At the end of the experiment, tumors were harvested, and human macrophages were isolated by
FACS (**Supplementary Fig. 7**). Transcriptomic analysis of these isolated macrophages by RNA
sequencing revealed significant gene expression changes consistent with enhanced activation and
phagocytic capacity upon Siglec-10 blockade (**Fig. 7c–e**). Specifically, macrophages from
Siglec-10 antibody–treated mice exhibited enrichment of PI3K/AKT, MAPK, AP-1, and NF-κB
signaling pathways, consistent with known Siglec-mediated signaling mechanisms described in
previous studies (20,43-46). Upregulated genes included PIK3C2A, PTEN, and PRKAA1,
important for PI3K/AKT signaling and macrophage metabolic reprogramming. Genes such as
MAP3K1, MAP3K2, JUN, and FOS supported activation of MAPK and AP-1 pathways,
promoting a pro-inflammatory and phagocytic phenotype. Additionally, upregulation of genes
related to cytoskeletal remodeling (e.g., RAB5A, DOCK1, WASL) and downregulation of
inhibitory genes (e.g., FCGR2B, CFL1) further underscored enhanced phagocytic function.
Furthermore, macrophages displayed evidence of metabolic reprogramming, with upregulation
of ACADM and PDK4 (supporting fatty acid oxidation) and downregulation of ALDOA
(indicating reduced glycolysis), consistent with a profile favoring sustained immune activation.
Together, these data show that Siglec-10 blockade *in vivo* reprograms macrophages toward an
activated, phagocytic, and metabolically adaptable state.

To complement these findings, we next employed a genetically modified human Siglec-10
transgenic immunocompetent mouse model. Human Siglec-10 knock-in (B-hSIGLEC10)
C57BL/6 mice are engineered to knock out Siglec-G and express human Siglec-10 instead (**Fig.**
7f, Supplementary Fig. 8). We subcutaneously engrafted these mice with 2838c3 cells, a highly
aggressive murine PDAC cell line derived from the well-established KPC (LSL-Kras$^{G12D/+}$; LSL-

Trp53$^{R172H/+}$; Pdx1-Cre) model. The KPC model is a spontaneous, autochthonous mouse model
that faithfully recapitulates the genetic landscape, desmoplastic stroma, immune evasion, and
rapid progression characteristic of human PDAC. We then treated these mice with either the
Siglec-10 antibody or a control (**Fig. 7g**). Siglec-10 targeting significantly reduced tumor volume
in this model without signs of toxicity (**Fig. 7h–i**). Overall, these *in vivo* results validate that
blocking Siglec-10 reprograms macrophages by activating well-established immune signaling
pathways (PI3K/AKT, MAPK, AP-1, NF-κB), consistent with mechanisms previously defined in
the literature (20,43-46). Together, these findings support Siglec-10 blockade as a promising
immunotherapeutic approach for PDAC.

DISCUSSION

In this study, we identified a novel immunological interaction between α3β1 integrin on PDAC cell surfaces and Siglec-10, a glyco-immune checkpoint receptor on macrophages, which facilitates immune evasion by PDAC cells. This previously unrecognized interaction can suppress macrophage-mediated immune surveillance, enabling tumor cells to escape immune clearance. Notably, we show that disrupting Siglec-10 interactions with monoclonal antibodies restores macrophage phagocytic capacity against PDAC cells. By tapping into glycan-based interventions, a largely underexplored avenue in cancer immunotherapy, our findings lay the foundation for innovative strategies that could enhance therapeutic outcomes for PDAC patients. With PDAC projected to account for approximately 67,440 new cases and 51,980 deaths in the United States in 2025 alone (47), this work addresses an urgent and critical clinical need.

In 2019, a pivotal study identified interactions between Siglec-10 and CD24 as a key "don't eat me" signal in breast and ovarian cancers, mediating immune evasion through macrophage suppression (20). Our findings uncover a distinct mechanism in PDAC, where α3β1 integrin, rather than CD24, emerges as one of the primary ligands for Siglec-10. Indeed, our comparative analyses of CD24 expression across PDAC, breast, and ovarian cancer cells showed that PDAC cells exhibit low CD24 expression, and many PDAC cells that bind Siglec-10 do not express CD24. This supports the presence of alternative Siglec-10 ligands in PDAC and underscores the importance of identifying context-dependent ligands mediating Siglec-10–driven immune evasion.

Integrin α3β1, composed of ITGA3 and ITGB1 subunits, is a well-characterized cell surface receptor known for roles in adhesion, migration, and metastasis (48-53). In PDAC, ITGA3 and ITGB1 are often overexpressed, contributing to epithelial-mesenchymal transition, invasion, and poor prognosis (54-60). Our study, for the first time, suggests a previously unrecognized immune regulatory function for α3β1 integrin, showing that its sialylated form acts as a critical ligand for Siglec-10, likely transmitting immunosuppressive signals that facilitate immune evasion. Our CRISPR-Cas9 knockout experiments of ITGA3 and depletion experiments of both ITGA3 and ITGB1 confirmed that loss of these integrins significantly increases susceptibility to macrophage-mediated phagocytosis. However, additional studies are needed to understand the

exact glycomic modulation of these integrin interactions, as they are heavily glycosylated with
multiple *N*-linked glycans and, in some contexts, may also carry *O*-linked glycans (61-66). This
could be addressed through comprehensive glycoproteomic studies and genetic modulation of
specific sialyltransferases and other glycosyltransferases that control the addition of Siglec-10
ligands to glycoproteins. These future investigations can broaden our understanding of $\alpha3\beta1$
integrin, expanding its role to include immune modulatory properties within the tumor
microenvironment.

Current immunotherapy efforts focusing on targeting proteins and nucleic acids have
revolutionized cancer treatment. However, these approaches remain insufficient to fully realize
the potential of immunotherapy for many patients, including those with PDAC (67,68).
Targeting glycomic interactions between glycans and glyco-immune checkpoints such as Siglec-
10 represents a novel and complementary approach to enhance the efficacy of cancer
immunotherapy. Our findings underscore the therapeutic promise of disrupting Siglec-10
interactions as a strategy to restore macrophage function and enhance anti-PDAC immunity.
Unlike traditional immune checkpoint inhibitors that primarily act on T cells (e.g., PD-1/PD-L1
blockers), Siglec-10 targeting directly reactivates myeloid cells like macrophages, which play a
pivotal role within the tumor microenvironment. This creates opportunities for combining Siglec-
10 targeting with other therapies. For example, combining it with therapies that target T cell
checkpoints could further amplify anti-tumor immunity. Additionally, Siglec-10 targeting can be
integrated with therapies designed to enhance macrophage-mediated phagocytosis, such as
CD47-blocking antibodies, which are already in clinical development. CD47 inhibits
phagocytosis by interacting with macrophage SIRPα (69-71). Since Siglec-10 and CD47 regulate
phagocytosis through distinct pathways, targeting both simultaneously could produce additive or
synergistic effects. Together, Siglec-targeting therapies, when combined with immune
checkpoint inhibitors acting on T cells or macrophages, have the potential to significantly expand
the arsenal of immunotherapeutic tools available for cancer treatment.

In our study, monoclonal antibodies against Siglec-10 restored macrophage phagocytic capacity
in vitro and reduced tumor growth *in vivo*, providing proof of concept for targeting this glyco-
immune checkpoint. However, targeting Siglecs on myeloid cells could have broader
implications for immunity by activating antigen-presenting cells and potentially enhancing

downstream T cell responses indirectly (72-74). While we focused on the role of Siglec-10
interactions in modulating phagocytosis, the broader impact of disrupting these interactions on
overall anti-tumor immunity, including effects on T cells via modulation of MDSCs and DCs,
warrants further investigation.

While our study provides significant insights into the role of Siglec-10 interactions in PDAC
immune evasion, it is not without limitations. First, a deeper mechanistic investigation into the
exact glycomic structures governing the binding between integrin α3β1 and Siglec-10 is needed,
which could be accomplished by genetic manipulation of key sialyltransferases and other
glycosyltransferases involved in adding Siglec-10 ligands to glycoproteins. Similarly, future
studies could benefit from direct genetic manipulation of Siglec-10 in primary myeloid cells.
Second, while we identified α3β1 integrin as a critical ligand for Siglec-10 in PDAC, it is likely
that additional ligands exist in other cancer types, and these remain to be explored. Third,
potential off-target effects and long-term consequences of Siglec-10 blockade, such as non-
specific inflammation or autoimmunity, require further investigation in more complex and
clinically relevant models. If toxicity is observed when blocking Siglecs using monospecific
antibodies, alternative approaches may be required. Bispecific or trispecific antibodies targeting
both Siglec-10 and its ligands, such as the α3β1 integrin identified in this study, could represent a
promising strategy. Such approaches could enhance immune responses by disrupting Siglec-10-
mediated immune suppression while minimizing the risk of non-specific inflammatory effects.
Fourth, our study primarily focused on establishing the functional role of the Siglec-10 axis as an
immune checkpoint that inhibits phagocytosis of PDAC cells using peripheral monocyte-derived
macrophages to provide a controlled, reproducible system. However, the impact of macrophage
polarization within the TME is highly relevant. Our single-cell analyses show that Siglec-10 is
highly expressed on TAMs from human PDAC tumors and is associated with
immunosuppressive transcriptional programs. This suggests that the Siglec-10 axis may be active
even within immunosuppressive TAMs. Further studies directly examining reprogramming of
TAMs isolated from PDAC patients would be highly valuable.

Despite these limitations, our findings represent a substantial advancement in understanding
glyco-immune interactions in PDAC biology. By identifying α3β1 integrin as a novel ligand for

Siglec-10 and showing the therapeutic potential of disrupting Siglec-10 interactions, we provide

a new framework for targeting immune evasion in PDAC.

REFERENCES

1.   Siegel RL, Miller KD, Jemal A. Cancer statistics, 2020. CA Cancer J Clin **2020**;70(1):7-30 doi 10.3322/caac.21590.

2.   Siegel RL, Miller KD, Wagle NS, Jemal A. Cancer statistics, 2023. CA Cancer J Clin **2023**;73(1):17-48 doi 10.3322/caac.21763.

3.   Timmer FEF, Geboers B, Nieuwenhuizen S, Dijkstra M, Schouten EAC, Puijk RS, *et al.* Pancreatic Cancer and Immunotherapy: A Clinical Overview. Cancers (Basel) **2021**;13(16) doi 10.3390/cancers13164138.

4.   Beatty GL, Eghbali S, Kim R. Deploying Immunotherapy in Pancreatic Cancer: Defining Mechanisms of Response and Resistance. Am Soc Clin Oncol Educ Book **2017**;37:267-78 doi 10.1200/EDBK_175232.

5.   Clark CE, Hingorani SR, Mick R, Combs C, Tuveson DA, Vonderheide RH. Dynamics of the immune reaction to pancreatic cancer from inception to invasion. Cancer Res **2007**;67(19):9518-27 doi 10.1158/0008-5472.CAN-07-0175.

6.   Helm O, Held-Feindt J, Grage-Griebenow E, Reiling N, Ungefroren H, Vogel I, *et al.* Tumor-associated macrophages exhibit pro- and anti-inflammatory properties by which they impact on pancreatic tumorigenesis. Int J Cancer **2014**;135(4):843-61 doi 10.1002/ijc.28736.

7.   RodrIguez E, Schetters STT, van Kooyk Y. The tumour glyco-code as a novel immune checkpoint for immunotherapy. Nat Rev Immunol **2018**;18(3):204-11 doi 10.1038/nri.2018.3.

8.   Fernandes A, Azevedo CM, Silva MC, Faria G, Dantas CS, Vicente MM, *et al.* Glycans as shapers of tumour microenvironment: A sweet driver of T-cell-mediated anti-tumour immune response. Immunology **2023**;168(2):217-32 doi 10.1111/imm.13494.

9.   Pinho SS, Reis CA. Glycosylation in cancer: mechanisms and clinical implications. Nat Rev Cancer **2015**;15(9):540-55 doi 10.1038/nrc3982.

10.   Silva MC, Fernandes A, Oliveira M, Resende C, Correia A, de-Freitas-Junior JC, *et al.* Glycans as Immune Checkpoints: Removal of Branched N-glycans Enhances Immune Recognition Preventing Cancer Progression. Cancer Immunol Res **2020**;8(11):1407-25 doi 10.1158/2326-6066.CIR-20-0264.

11. Rodrigues E, Macauley MS. Hypersialylation in Cancer: Modulation of Inflammation and Therapeutic Opportunities. Cancers (Basel) **2018**;10(6) doi 10.3390/cancers10060207.

12. Gu Y, Zhou G, Tang X, Shen F, Ding J, Hua K. The biological roles of CD24 in ovarian cancer: old story, but new tales. Front Immunol **2023**;14:1183285 doi 10.3389/fimmu.2023.1183285.

13. Laubli H, Nalle SC, Maslyar D. Targeting the Siglec-Sialic Acid Immune Axis in Cancer: Current and Future Approaches. Cancer Immunol Res **2022**;10(12):1423-32 doi 10.1158/2326-6066.CIR-22-0366.

14. Marciel MP, Haldar B, Hwang J, Bhalerao N, Bellis SL. Role of tumor cell sialylation in pancreatic cancer progression. Adv Cancer Res **2023**;157:123-55 doi 10.1016/bs.acr.2022.07.003.

15. Saini P, Adeniji OS, Abdel-Mohsen M. Inhibitory Siglec-sialic acid interactions in balancing immunological activation and tolerance during viral infections. EBioMedicine **2022**;86:104354 doi 10.1016/j.ebiom.2022.104354.

16. Stanczak MA, Laubli H. Siglec receptors as new immune checkpoints in cancer. Mol Aspects Med **2023**;90:101112 doi 10.1016/j.mam.2022.101112.

17. Yu Y, Peng W. Recent progress in targeting the sialylated glycan-SIGLEC axis in cancer immunotherapy. Cancer Biol Med **2023**;20(5):369-84 doi 10.20892/j.issn.2095-3941.2023.0046.

18. Zhou X, Chi K, Zhang C, Liu Q, Yang G. Sialylation: A Cloak for Tumors to Trick the Immune System in the Microenvironment. Biology (Basel) **2023**;12(6) doi 10.3390/biology12060832.

19. Bandala-Sanchez E, Zhang Y, Reinwald S, Dromey JA, Lee BH, Qian J, *et al.* T cell regulation mediated by interaction of soluble CD52 with the inhibitory receptor Siglec-10. Nat Immunol **2013**;14(7):741-8 doi 10.1038/ni.2610.

20. Barkal AA, Brewer RE, Markovic M, Kowarsky M, Barkal SA, Zaro BW, *et al.* CD24 signalling through macrophage Siglec-10 is a target for cancer immunotherapy. Nature **2019**;572(7769):392-6 doi 10.1038/s41586-019-1456-0.

21. Hazra R, Chattopadhyay S, Mallick A, Gayen S, Roy S. Unravelling CD24-Siglec-10 pathway: Cancer immunotherapy from basic science to clinical studies. Immunology **2024**;173(3):442-69 doi 10.1111/imm.13847.

22.   Li X, Tian W, Jiang Z, Song Y, Leng X, Yu J. Targeting CD24/Siglec-10 signal pathway for cancer immunotherapy: recent advances and future directions. Cancer Immunol Immunother **2024**;73(2):31 doi 10.1007/s00262-023-03606-0.

23.   van Bruggen JAC, Peters FS, Mes M, Rietveld JM, Cerretani E, Cretenet G*, et al.* T-cell dysfunction in CLL is mediated through expression of Siglec-10 ligands CD24 and CD52 on CLL cells. Blood Adv **2024**;8(17):4633-46 doi 10.1182/bloodadvances.2023011934.

24.   Xiao N, Zhu X, Li K, Chen Y, Liu X, Xu B*, et al.* Blocking siglec-10(hi) tumor-associated macrophages improves anti-tumor immunity and enhances immunotherapy for hepatocellular carcinoma. Exp Hematol Oncol **2021**;10(1):36 doi 10.1186/s40164-021-00230-5.

25.   Boelaars K, Rodriguez E, Huinen ZR, Liu C, Wang D, Springer BO*, et al.* Pancreatic cancer-associated fibroblasts modulate macrophage differentiation via sialic acid-Siglec interactions. Commun Biol **2024**;7(1):430 doi 10.1038/s42003-024-06087-8.

26.   Chen X, Mo S, Zhang Y, Ma H, Lu Z, Yu S*, et al.* Analysis of a novel immune checkpoint, Siglec-15, in pancreatic ductal adenocarcinoma. J Pathol Clin Res **2022**;8(3):268-78 doi 10.1002/cjp2.260.

27.   Dharmadhikari G, Stolz K, Hauke M, Morgan NG, Varki A, de Koning E*, et al.* Siglec-7 restores beta-cell function and survival and reduces inflammation in pancreatic islets from patients with diabetes. Sci Rep **2017**;7:45319 doi 10.1038/srep45319.

28.   Li H, Zhu R, Liu X, Zhao K, Hong D. Siglec-15 Regulates the Inflammatory Response and Polarization of Tumor-Associated Macrophages in Pancreatic Cancer by Inhibiting the cGAS-STING Signaling Pathway. Oxid Med Cell Longev **2022**;2022:3341038 doi 10.1155/2022/3341038.

29.   Longevity O. Retracted: Siglec-15 Regulates the Inflammatory Response and Polarization of Tumor-Associated Macrophages in Pancreatic Cancer by Inhibiting the cGAS-STING Signaling Pathway. Oxid Med Cell Longev **2023**;2023:9793031 doi 10.1155/2023/9793031.

30.   Rodriguez E, Boelaars K, Brown K, Eveline Li RJ, Kruijssen L, Bruijns SCM*, et al.* Sialic acids in pancreatic cancer cells drive tumour-associated macrophage differentiation via the Siglec receptors Siglec-7 and Siglec-9. Nat Commun **2021**;12(1):1270 doi 10.1038/s41467-021-21550-4.

31.   Swanson BJ, McDermott KM, Singh PK, Eggers JP, Crocker PR, Hollingsworth MA. MUC1 is a counter-receptor for myelin-associated glycoprotein (Siglec-4a) and their

interaction contributes to adhesion in pancreatic cancer perineural invasion. Cancer Res **2007**;67(21):10222-9 doi 10.1158/0008-5472.CAN-06-2483.

32. Werba G, Weissinger D, Kawaler EA, Zhao E, Kalfakakou D, Dhara S*, et al.* Single-cell RNA sequencing reveals the effects of chemotherapy on human pancreatic adenocarcinoma and its tumor microenvironment. Nat Commun **2023**;14(1):797 doi 10.1038/s41467-023-36296-4.

33. Chang L, Chen YJ, Fan CY, Tang CJ, Chen YH, Low PY*, et al.* Identification of Siglec Ligands Using a Proximity Labeling Method. J Proteome Res **2017**;16(10):3929-41 doi 10.1021/acs.jproteome.7b00625.

34. Crissey MAS, Versace A, Bhardwaj M, Jain V, Liu S, Singh A*, et al.* Divergent effects of acute and chronic PPT1 inhibition in melanoma. Autophagy **2024**:1-13 doi 10.1080/15548627.2024.2403152.

35. Cox J, Mann M. MaxQuant enables high peptide identification rates, individualized p.p.b.-range mass accuracies and proteome-wide protein quantification. Nat Biotechnol **2008**;26(12):1367-72 doi 10.1038/nbt.1511.

36. Martinez FO. Analysis of gene expression and gene silencing in human macrophages. Curr Protoc Immunol **2012**;Chapter 14:14 28 1-14 28 3 doi 10.1002/0471142735.im1428s96.

37. Luo Q, Hu Z, Zhao H, Fan Y, Tu X, Wang Y*, et al.* The role of TGF-beta in the tumor microenvironment of pancreatic cancer. Genes Dis **2023**;10(4):1513-24 doi 10.1016/j.gendis.2022.10.019.

38. Principe DR, Timbers KE, Atia LG, Koch RM, Rana A. TGFbeta Signaling in the Pancreatic Tumor Microenvironment. Cancers (Basel) **2021**;13(20) doi 10.3390/cancers13205086.

39. Crocker PR, Varki A. Siglecs in the immune system. Immunology **2001**;103(2):137-45 doi 10.1046/j.0019-2805.2001.01241.x.

40. Liu Q, Sasaki T, Kozieradzki I, Wakeham A, Itie A, Dumont DJ*, et al.* SHIP is a negative regulator of growth factor receptor-mediated PKB/Akt activation and myeloid cell survival. Genes Dev **1999**;13(7):786-91 doi 10.1101/gad.13.7.786.

41. Yeoh WJ, Krebs P. SHIP1 and its role for innate immune regulation-Novel targets for immunotherapy. Eur J Immunol **2023**;53(12):e2350446 doi 10.1002/eji.202350446.

42. Yin SS, Gao FH. Molecular Mechanism of Tumor Cell Immune Escape Mediated by CD24/Siglec-10. Front Immunol **2020**;11:1324 doi 10.3389/fimmu.2020.01324.

43. Chen GY, Tang J, Zheng P, Liu Y. CD24 and Siglec-10 selectively repress tissue damage-induced immune responses. Science **2009**;323(5922):1722-5 doi 10.1126/science.1168988.

44. Jandus C, Boligan KF, Chijioke O, Liu H, Dahlhaus M, Demoulins T, *et al.* Interactions between Siglec-7/9 receptors and ligands influence NK cell-dependent tumor immunosurveillance. J Clin Invest **2014**;124(4):1810-20 doi 10.1172/JCI65899.

45. Kang EA, Soh H, Park S, Lee HJ, Im JP, Kim JS. Soluble Siglec-9 alleviates intestinal inflammation through inhibition of the NF-kappaB pathway. Int Immunopharmacol **2020**;86:106695 doi 10.1016/j.intimp.2020.106695.

46. Laubli H, Varki A. Sialic acid-binding immunoglobulin-like lectins (Siglecs) detect self-associated molecular patterns to regulate immune responses. Cell Mol Life Sci **2020**;77(4):593-605 doi 10.1007/s00018-019-03288-x.

47. Siegel RL, Kratzer TB, Giaquinto AN, Sung H, Jemal A. Cancer statistics, 2025. CA Cancer J Clin **2025**;75(1):10-45 doi 10.3322/caac.21871.

48. Ganguly K, Cox JL, Ghersi D, Grandgenett PM, Hollingsworth MA, Jain M, *et al.* Mucin 5AC-Mediated CD44/ITGB1 Clustering Mobilizes Adipose-Derived Mesenchymal Stem Cells to Modulate Pancreatic Cancer Stromal Heterogeneity. Gastroenterology **2022**;162(7):2032-46 e12 doi 10.1053/j.gastro.2022.02.032.

49. Li R, Ji Q, Fu S, Gu J, Liu D, Wang L, *et al.* ITGA3 promotes pancreatic cancer progression through HIF1alpha- and c-Myc-driven glycolysis in a collagen I-dependent autocrine manner. Cancer Gene Ther **2024** doi 10.1038/s41417-024-00864-7.

50. Miskin RP, Warren JSA, Ndoye A, Wu L, Lamar JM, DiPersio CM. Integrin alpha3beta1 Promotes Invasive and Metastatic Properties of Breast Cancer Cells through Induction of the Brn-2 Transcription Factor. Cancers (Basel) **2021**;13(3) doi 10.3390/cancers13030480.

51. Subbaram S, Dipersio CM. Integrin alpha3beta1 as a breast cancer target. Expert Opin Ther Targets **2011**;15(10):1197-210 doi 10.1517/14728222.2011.609557.

52. Subbaram S, Lyons SP, Svenson KB, Hammond SL, McCabe LG, Chittur SV, *et al.* Integrin alpha3beta1 controls mRNA splicing that determines Cox-2 mRNA stability in breast cancer cells. J Cell Sci **2014**;127(Pt 6):1179-89 doi 10.1242/jcs.131227.

53.   Varzavand A, Drake JM, Svensson RU, Herndon ME, Zhou B, Henry MD, *et al.* Integrin alpha3beta1 regulates tumor cell responses to stromal cells and can function to suppress prostate cancer metastatic colonization. Clin Exp Metastasis **2013**;30(4):541-52 doi 10.1007/s10585-012-9558-1.

54.   Hu F, Zhao L, Wang Y, Ye H, Tang H, Zhou J. Integrated bioinformatics analysis shows integrin alpha 3 is a prognostic biomarker for pancreatic cancer. Open Med (Wars) **2022**;17(1):1935-43 doi 10.1515/med-2022-0606.

55.   Idichi T, Seki N, Kurahara H, Fukuhisa H, Toda H, Shimonosono M, *et al.* Involvement of anti-tumor miR-124-3p and its targets in the pathogenesis of pancreatic ductal adenocarcinoma: direct regulation of ITGA3 and ITGB1 by miR-124-3p. Oncotarget **2018**;9(48):28849-65 doi 10.18632/oncotarget.25599.

56.   Iwatate Y, Yokota H, Hoshino I, Ishige F, Kuwayama N, Itami M, *et al.* Transcriptomic analysis reveals high ITGB1 expression as a predictor for poor prognosis of pancreatic cancer. PLoS One **2022**;17(6):e0268630 doi 10.1371/journal.pone.0268630.

57.   Jiao Y, Li Y, Liu S, Chen Q, Liu Y. ITGA3 serves as a diagnostic and prognostic biomarker for pancreatic cancer. Onco Targets Ther **2019**;12:4141-52 doi 10.2147/OTT.S201675.

58.   Li J, Peng L, Chen Q, Ye Z, Zhao T, Hou S, *et al.* Integrin beta1 in Pancreatic Cancer: Expressions, Functions, and Clinical Implications. Cancers (Basel) **2022**;14(14) doi 10.3390/cancers14143377.

59.   Lu Y, Hu J, Sun W, Li S, Deng S, Li M. MiR-29c inhibits cell growth, invasion, and migration of pancreatic cancer by targeting ITGB1. Onco Targets Ther **2016**;9:99-109 doi 10.2147/OTT.S92758.

60.   Zhang W, Han S, Yuan Y, Xu M, Ding A, Li M. FTO Knockdown-Mediated Maturation of miR-383-5p Inhibits Malignant Advancement of Pancreatic Cancer by Targeting ITGA3. Biochem Genet **2024**;62(4):2652-66 doi 10.1007/s10528-023-10560-0.

61.   Marsico G, Russo L, Quondamatteo F, Pandit A. Glycosylation and Integrin Regulation in Cancer. Trends Cancer **2018**;4(8):537-52 doi 10.1016/j.trecan.2018.05.009.

62.   Pochec E, Litynska A, Amoresano A, Casbarra A. Glycosylation profile of integrin alpha 3 beta 1 changes with melanoma progression. Biochim Biophys Acta **2003**;1643(1-3):113-23 doi 10.1016/j.bbamcr.2003.10.004.

63.   Ranjan A, Bane SM, Kalraiya RD. Glycosylation of the laminin receptor (alpha3beta1) regulates its association with tetraspanin CD151: Impact on cell spreading, motility,

degradation and invasion of basement membrane by tumor cells. Exp Cell Res **2014**;322(2):249-64 doi 10.1016/j.yexcr.2014.02.004.

64.   Ren Y, Hao P, Law SK, Sze SK. Hypoxia-induced changes to integrin alpha 3 glycosylation facilitate invasion in epidermoid carcinoma cell line A431. Mol Cell Proteomics **2014**;13(11):3126-37 doi 10.1074/mcp.M114.038505.

65.   Zhao Y, Nakagawa T, Itoh S, Inamori K, Isaji T, Kariya Y*, et al.* N-acetylglucosaminyltransferase III antagonizes the effect of N-acetylglucosaminyltransferase V on alpha3beta1 integrin-mediated cell migration. J Biol Chem **2006**;281(43):32122-30 doi 10.1074/jbc.M607274200.

66.   Zhou Y, Fukuda T, Hang Q, Hou S, Isaji T, Kameyama A*, et al.* Inhibition of fucosylation by 2-fluorofucose suppresses human liver cancer HepG2 cell proliferation and migration as well as tumor formation. Sci Rep **2017**;7(1):11563 doi 10.1038/s41598-017-11911-9.

67.   Johnson PC, Gainor JF, Sullivan RJ, Longo DL, Chabner B. Immune Checkpoint Inhibitors - The Need for Innovation. N Engl J Med **2023**;388(16):1529-32 doi 10.1056/NEJMsb2300232.

68.   Ju Y, Xu D, Liao MM, Sun Y, Bao WD, Yao F*, et al.* Barriers and opportunities in pancreatic cancer immunotherapy. NPJ Precis Oncol **2024**;8(1):199 doi 10.1038/s41698-024-00681-z.

69.   Matlung HL, Szilagyi K, Barclay NA, van den Berg TK. The CD47-SIRPalpha signaling axis as an innate immune checkpoint in cancer. Immunol Rev **2017**;276(1):145-64 doi 10.1111/imr.12527.

70.   Weiskopf K, Jahchan NS, Schnorr PJ, Cristea S, Ring AM, Maute RL*, et al.* CD47-blocking immunotherapies stimulate macrophage-mediated destruction of small-cell lung cancer. J Clin Invest **2016**;126(7):2610-20 doi 10.1172/JCI81603.

71.   Willingham SB, Volkmer JP, Gentles AJ, Sahoo D, Dalerba P, Mitra SS*, et al.* The CD47-signal regulatory protein alpha (SIRPa) interaction is a therapeutic target for human solid tumors. Proc Natl Acad Sci U S A **2012**;109(17):6662-7 doi 10.1073/pnas.1121623109.

72.   Egan H, Treacy O, Lynch K, Leonard NA, O'Malley G, Reidy E*, et al.* Targeting stromal cell sialylation reverses T cell-mediated immunosuppression in the tumor microenvironment. Cell Rep **2023**;42(5):112475 doi 10.1016/j.celrep.2023.112475.

73. Mei Y, Wang X, Zhang J, Liu D, He J, Huang C*, et al.* Siglec-9 acts as an immune-checkpoint molecule on macrophages in glioblastoma, restricting T-cell priming and immunotherapy response. Nat Cancer **2023**;4(9):1273-91 doi 10.1038/s43018-023-00598-9.

74. Wang J, Manni M, Barenwaldt A, Wieboldt R, Kirchhammer N, Ivanek R*, et al.* Siglec Receptors Modulate Dendritic Cell Activation and Antigen Presentation to T Cells in Cancer. Front Cell Dev Biol **2022**;10:828916 doi 10.3389/fcell.2022.828916.

FIGURE LEGENDS

Figure 1. Siglec-10 is predominantly expressed on myeloid cells within the PDAC TME and is associated with worse disease progression. (**a**) UMAP visualization of immune cell populations within the PDAC TME. (**b**) UMAP of Siglec-10 expression and a pie chart illustrating the percentage of Siglec-10□ cells within each myeloid cell subset. (**c**) A comparison of the expression levels of various Siglecs in myeloid cells within the PDAC TME. Kruskal–Wallis test with Dunn's multiple comparisons correction. Means with standard deviations (SD) are shown. Percentages shown represent the proportion of myeloid cells expressing each of these Siglecs. (**d**) A comparison of the gene expression of the indicated genes between Siglec-10□ and Siglec-10□ macrophages, MDSCs, and DCs. Analysis was performed using the Bioturing's Talk2Data V4 platform. (**e**) Siglec-10 expression in the PDAC TME across different disease states. Kruskal–Wallis test with Dunn's multiple comparisons correction. Means with standard errors of the mean (SEM) are shown. (**f**) Comparison of Siglec-10 expression between normal tissues and pancreatic adenocarcinoma (PAAD) tissues from the TCGA dataset. Unpaired t test. (**g**) Survival analysis of pancreatic tumor patients in the Human Protein Atlas dataset showing the correlation between Siglec-10 expression and overall survival.

Figure 2. PDAC cell lines express Siglec-10 ligands on glycoproteins other than CD24. Flow cytometric analysis of Siglec-10 ligand and CD24 expression on (**a–f**) PDAC cell lines, (**g–j**) ovarian cancer cell lines, and (**k–l**) a breast cancer cell line. Siglec-10 ligands were detected using either (1) a recombinant Siglec-10-Fc chimeric protein and a secondary antibody (x-axis), with CD24 detected using a specific antibody (y-axis), or (2) a recombinant Siglec-10-Fc chimeric protein containing the inhibitory R119A mutation, which abrogates sialic acid binding (x-axis), with CD24 detected using a specific antibody (y-axis). Panels on the left (**a, c, e, g, i, k**) show representative flow plots. Panels on the right (**b, d, f, h, j, l**) show the percentage of positive cells and the median fluorescence intensity (MFI) for Siglec-10 ligands (using the non-mutated recombinant Siglec-10 protein), CD24, or cells expressing both. Means with SD are shown.

Figure 3. ITGA3 and ITGB1 are sialylated glycoprotein ligands of Siglec-10 glycoproteins on PDAC cells. (a) Experimental design for the pull-down of Siglec-10 ligands on PDAC cells. Recombinant Siglec-10 Fc (as well as a no-protein control or a Siglec-5 control) was allowed to bind its physiological ligands on the surface of PDAC cells, followed by an HRP-conjugated anti-Fc secondary antibody. In the presence of H_2O_2, HRP generated short-lived radicals that facilitated the transfer of biotin to proximal Siglec-10 ligands. Biotinylated Siglec-10 ligands were pulled down using streptavidin beads and identified by mass spectrometry. **(b)** A total of 4,044 proteins were identified, with enriched binding compared to a control using the anti-Fc antibody only without Siglec-10 protein. Of these, 110 proteins showed enrichment relative to a Siglec-5 control. Six proteins, CD47, CD59, CD73, ITGB6, ITGA3, and ITGB1, were significantly overexpressed in PAAD tissues compared to normal tissues in the TCGA dataset. **(c)** Response curves showing interactions between Siglec-10 and the six glycoproteins measured by surface plasmon resonance (SPR). Two concentrations (1000 nM, green; 100 nM, red) were tested for all glycoproteins, while ITGA3 was also tested at 300 nM (green) and 30 nM (red). **(d)** Binding of the SNA lectin (specific for sialic acid) to ITGA3 and ITGB1 recombinant glycoproteins was measured by a lectin array. Sialidase-treated glycoproteins (red bars) showed significantly reduced binding compared to untreated glycoproteins (blue bars). Unpaired t-tests. Means with SEM are shown. **(e)** SPR response curves comparing the binding of intact (sialylated) ITGA3 and desialylated ITGA3 to immobilized Siglec-10. **(f)** SPR response curves comparing the binding of intact (sialylated) ITGB1 and desialylated ITGB1 to immobilized Siglec-10.

Figure 4. ITGA3 and ITGB1 are overexpressed in pancreatic cancer tissues and associated with poor survival. (a–c) Comparison of CD24 (a), ITGA3 (b), and ITGB1 (c) expression between normal tissues and PAAD tissues in the TCGA dataset. Unpaired t tests. **(d–f)** Survival analyses of pancreatic tumor patients in the TCGA dataset showing the correlation between CD24 (d), ITGA3 (e), and ITGB1 (f) expression and overall survival. **(g–i)** Expression of CD24 (g), ITGA3 (h), and ITGB1 (i) in the PDAC TME across different disease states. Kruskal–Wallis test with Dunn's multiple comparisons correction. Means with SEM are shown.

Figure 5. ITGA3-expressing PDAC cells evade macrophage-mediated phagocytosis. (a) ITGA3 knock-down in PANC-1 PDAC cells was achieved using CRISPR-Cas9. Flow cytometry confirmed the reduction of ITGA3 expression in knockout cells (ITGA3-KO) compared to non-targeting gRNA-treated cells and untreated controls. Data from three independent replicates show the percentage of ITGA3-expressing cells and the mean fluorescence intensity (MFI). Means and SEM are shown. Unpaired t-tests. (**b**) Experimental schematic illustrating co-culture of red-labeled PDAC cells (ITGA3-KO or non-targeting gRNA-treated) with monocyte-derived macrophages. Phagocytosis was monitored over time using live-cell imaging on the Incucyte Live-Cell Analysis System. (**c**) (left) Phagocytosis over time and area under the curve (AUC). Data show macrophage-mediated phagocytosis of PDAC cells, normalized to the 0-hour time point. Paired t-tests. N = 3 donors, with experiments performed in triplicate for each donor. (right) Representative images showing increased red fluorescence as an indicator of enhanced phagocytosis. (**d**) Schematic of ITGA3^{low} PDAC cell enrichment. ITGA3^{high} PDAC cells were separated using anti-ITGA3 antibody-coated columns, and the flow-through, containing ITGA3^{low} cells, was collected. (**e**) Flow cytometry analysis of ITGA3 expression in MIA PaCa-2 cells after enrichment for ITGA3^{low} cells and comparison with non-enriched controls (Ctrl cells). (**f**) Experimental schematic illustrating the phagocytic assay setup, where monocyte-derived macrophages were co-cultured with either ITGA3^{high} (control) or ITGA3^{low} PDAC cells. (**g-h**) Phagocytosis analysis of ITGA3^{high} (control) versus ITGA3^{low} MIA PaCa-2 (g) or PANC1 (h) cells by macrophages derived from different donors. N = 5–6, with experiments performed in triplicate for each donor. Top: Representative images showing increased red fluorescence as an indicator of enhanced phagocytosis. Bottom left: Live imaging data from individual donors (each symbol represents one donor's data done in triplicate per donor). Bottom right: AUC data from multiple donors, showing significantly higher phagocytic activity with ITGA3^{low} cells. Paired t-tests.

Figure 6. Development of Siglec-10 blocking antibodies to enhance macrophage-mediated phagocytosis of PDAC cells and prevent myeloid cell–mediated inhibition of T cell proliferation and activation *in vitro*. (a) Schematic model illustrating Siglec-10-mediated suppression of macrophage phagocytosis. In the left panel, Siglec-10 on macrophages binds to

glycan ligands on PDAC cells, including ITGA3, ITGB1, and CD24, triggering inhibitory signaling and suppressing phagocytosis. In the right panel, blocking Siglec-10 with an antibody prevents inhibitory signaling and enhances macrophage phagocytic capacity. (**b**) ELISA screening of recombinant antibodies from the top hybridoma clones for Siglec-10 binding. Binding to immobilized Siglec-10 (blue) and Siglec-5 (gray) proteins is shown. (**c**) Flow cytometric analysis of antibody selectivity, showing binding to CHO-K1 cells expressing Siglec-10 (blue) but not Siglec-5 (gray). (**d**) AUC analysis of *in vitro* phagocytosis assays screening various Siglec-10 antibody clones, along with commercially available anti-CD24 and anti-Siglec-10 antibodies, for their ability to enhance macrophage-mediated phagocytosis of AsPC-1 PDAC cells. Means with SEM are shown. (**e**) AUC analysis of the *in vitro* phagocytosis assay using the top-performing Siglec-10 antibody clone with macrophages differentiated from monocytes of four healthy donors. Statistical significance was determined using Friedman's ANOVA test. Means with SEM are shown. (**f**) Time-course analysis of the *in vitro* phagocytosis assay comparing the top Siglec-10 blocking antibody clone (68A11A1, blue) with the isotype control (gray). Data represent n = 4 independent experiments. (**g**) ELISA-based binding analysis of the recombinant 68A11A1 antibody to immobilized recombinant Siglec-10 and Siglec-5 proteins across different dilutions. (**h**) Evaluation of the recombinant Siglec-10 antibody (clone 68A11A1) and anti-CD24 antibody in enhancing macrophage-mediated phagocytosis of multiple PDAC cell lines (AsPC-1, MIA PaCa-2, and PANC-1). Phagocytosis was normalized to the isotype control for each cell line and conducted using macrophages derived from monocytes of 5–8 healthy donors. Each symbol represents an individual donor; statistical significance was assessed using ratio paired t-tests compared to isotype control. Means with SEM are shown. (**i**) Triple co-culture assay involving cancer-associated fibroblasts (CAFs), PANC-1 PDAC cells, and monocyte-derived macrophages, showing phagocytosis kinetics, AUC quantification, and representative images. Statistical significance assessed using paired t-tests. (**j-k**) Flow cytometry analysis of CellTrace Violet (CTV)-labeled human CD8$^+$ T cells co-cultured with human monocytes ± anti-Siglec-10 antibody in the presence of anti-CD3/CD28 beads for 5 days. (**j**) T cell proliferation; (**k**) granzyme B expression. ANOVA with post hoc comparisons.

Figure 7. Blocking Siglec-10 interactions enhances macrophage-mediated phagocytosis and reduces PDAC tumor growth *in vivo*. (**a**) Experimental design for subcutaneous tumor development and Siglec-10 antibody treatment. AsPC-1 PDAC cells were subcutaneously injected into immunodeficient NSG mice. After 7 days, when tumors reached ~40 mm³, mice were randomized into treatment groups. On the indicated days, ~$3 \times 10^{\square}$ monocyte-derived macrophages were intravenously injected along with 200 µg of either isotype control or anti-Siglec-10 antibody. Tumor size was measured on specified days using a vernier caliper. (b) Tumor volume over time (left) and at the last time point (right) in mice treated with isotype control or anti-Siglec-10 antibody. n = 10 mice per group. Means with SEM are shown for the longitudinal analysis. For the last time point analysis, box and whiskers represent the minimum to maximum range, with all individual data points shown. Mann–Whitney U test. (**c–e**) Heatmaps showing differential gene expression in FACS-sorted human macrophages from tumors of NSG mice treated with anti-Siglec-10 antibody, highlighting: (c) immune activation pathways, (d) phagocytosis-related genes, and (e) metabolism-related genes. Red indicates higher expression; blue indicates lower expression. (**f**) Flow cytometry analysis of peritoneal cells from wild-type B6 mice and transgenic B-hSIGLEC10 mice. B6 mice express Siglec-G but not Siglec-10, whereas B-hSIGLEC10 mice express Siglec-10 but not Siglec-G on myeloid cells. Means with SEM are shown. (**g**) Experimental design for tumor challenge in B-hSIGLEC10 transgenic mice. KPC-derived murine PDAC cells (2838c3) were subcutaneously injected. After 5 days, mice were randomized and treated intravenously with either isotype control or anti–Siglec-10 antibody. (**h**) Tumor volume in B-hSIGLEC10 mice treated with isotype control or anti–Siglec-10 antibody. Tumor volume over time (left) and at the last time point (right) in mice treated with isotype control or anti-Siglec-10 antibody. Means with SEM are shown for the longitudinal analysis. N=4-5 per group. For the last time point analysis, box and whiskers represent the minimum to maximum range, with all individual data points shown. Unpaired t test. (**i**) Mouse body weight during treatment.

SUPPLEMENTARY FIGURE LEGENDS

Supplementary Figure 1. Pancreatic cancer is associated with high expression of sialyltransferases involved in producing Siglec-10 ligands. (a) A schematic representation of the glycan ligands of Siglec-10 on *N*-linked and *O*-linked glycans, along with the enzymes involved in their formation. **(b–c)** Expression levels of sialyltransferase enzymes responsible for Siglec-10 ligand formation compared between normal tissues and PAAD tissues in the TCGA dataset. Unpaired t tests. *$P < 0.05$.

Supplementary Figure 2. Monocyte-derived macrophages express Siglec-10. A flow cytometry-based analysis of monocytes and macrophages derived from monocytes for Siglec-10 expression.

Supplementary Figure 3. ITGA3 and ITGAB1 highly expresses in PDAC cells. Several PDAC cells were examined for their expression of (a) ITGA3 and (b) ITGB1 (x-axis) by flow cytometry.

Supplementary Figure 4. The expression of Siglec-10 and Siglec-10 ligand forming enzymes associated with the expression of ITGA3 and ITGB1. Spearman r correlation analysis in TCGA of (a) Siglec-10 with ITGA3, (b) ITGA3 with Siglec-10 ligand forming enzymes (ST3GAL1 and B3GNT3), (c) Siglec-10 with ITGB1, and (d) ITGB1 with Siglec-10 ligand forming enzymes (ST3GAL1 and B3GNT3).

Supplementary Figure 5. ITGB1-expressing PDAC cells evade macrophage-mediated phagocytosis. Columns coated with anti-ITGB1 antibodies were used to isolate ITGB1^{low} cells (cells passing through the columns without binding). Cells passing through uncoated columns (expressing high levels of ITGB1) were used as controls. **(a)** Schematic representation of the experimental setup to test the phagocytic capacity of monocyte-derived macrophages targeting control cells (enriched with ITGB1^{hi} cells) or ITGB1^{low} cells (n = 5–6). **(b-c)** Phagocytosis of

control and ITGB1^{low} MIA PaCa-2 (f) or PANC1 (g) PDAC cells by macrophages from different donors. Top: Representative images showing phagocytosis (increased red indicates higher phagocytosis). Bottom left: Live imaging results (each symbol represents data from one donor). Bottom right: AUC data compiled from multiple donors. Statistical significance was assessed using ratio paired t-tests.

Supplementary Figure 6. Targeting Siglec-10 using anti–Siglec-10 antibodies does not impact the levels of ITGA3, ITGB1, or Siglec-10 ligands on PDAC cells. Percentage and MFI of (**a–b**) ITGA3, (**c–d**) ITGB1, and (**e–f**) Siglec-10 ligands measured by flow cytometry. ANOVA tests corrected with Benjamini, Krieger, and Yekutieli method. Means and SEM are shown.

Supplementary Figure 7. Gating strategy for sorting the human tumor associated macrophages from mouse subcutaneous tumor. Single cell suspension from mouse subcutaneous tumor was used to obtain human monocytes derived macrophages injected 12hr intravenously for RNA seq analysis. Mouse BV711-CD45⁻ and Human APC-CD45⁺ cells were sorted for RNA seq analysis.

Supplementary Figure 8. B-hSIGLEC10 mice express human Siglec-10 on their myeloid cells. (a–b) Gating strategy for cells from (**a**) B-hSIGLEC10 mice and (**b**) C57BL/6 mice. (**c**) Representative example of the expression of human Siglec-10 and mouse Siglec-G on B cells, monocytes, macrophages, and DCs from both mouse models.

SUPPLEMENTARY TABLES

Supplementary Table 1. Sequences of the monoclonal antibodies against Siglec-10

ACKNOWLEDGMENTS

M.A-M is supported by NIH grants (NIH R01AG092241, R01AI165079, R01AA029859, R01DK123733, and R01NS117458). M.A-M is also funded by the NIH-funded BEAT-HIV Martin Delaney Collaboratory to cure HIV-1 infection (1UM1Al126620). H.Y.T is supported by NCI R50CA221838, and the Wistar Shared Resources are supported by P30 CA010815.

AUTHOR CONTRIBUTIONS

M.A.-M conceived and designed the study. P.S carried out the majority of the experiments. G.M, S.M.S.I, S.A.B, and A.P.B performed the animal experiments. R.S. supervised and designed the animal experiments. L.M.S. and J.F.H. supervised and helped perform the CRISPR-Cas9 experiments. J.C and H.-Y.T conducted the SPR and mass spectrometry analyses. K.M. consulted on the generation of the antibodies. H.T and R.Z interpreted the data. P.S and M.A.-M wrote the manuscript, and all authors edited it.

COMPETING INTERESTS STATEMENT

J.F.H. received research support, paid to Northwestern University, from Gilead Sciences and is a paid consultant for Merck and Ridgeback Biotherapeutics. All other authors declare no competing interests.

Figure 1

a Single cells analysis of PDAC tissues

Macrophages (21.8%)
MDSCs (6.4%)
DCs (3%)
B cells (5.8%)
CD4⁺ T cells (27.8%)
NK cells (8.5%)
CD8⁺ T cells (19.5%)
UMAP2
UMAP1

Lymphocytes (2.3%)
Myeloid leukocytes (1.3%)
Monocytes (1.2%)
Mast cells (1%)
T cells (0.8%)
pDCs(0.5%)

b % of Siglec-10-expressing cells within immune cells

Macrophages
MDSCs
DCs
Monocytes
60%
11%
8%
8%
UMAP2
UMAP1

c Siglec expression on myeloid cells in PDAC TME

P<0.0001
P<0.0001
P<0.0001
20%
17%
14%
2%
Siglec-10 Siglec-7 Siglec-9 Siglec-15

d

Siglec-10⁺ MΦ vs. Siglec-10⁻ MΦ

Gene	FDR
EBI3	3.0E-22
PDCD1LG2	2.3E-16
CMKLR1	1.9E-48
PROS1	4.0E-11
GAS6	1.8E-26
LILRB2	2.8E-48
LILRB1	3.2E-45
SMURF2	5.9E-20
MERTK	1.0E-37
CD274	1.1E-05
LILRB4	4.8E-54
LAIR1	4.0E-71
VSIG4	1.7E-48
TGFBR2	4.5E-31
CD47	1.9E-31
IL10RA	6.9E-18
TGFBRAP1	1.0E-03
SIRPB1	1.2E-04
CCL18	2.9E-07
SHP-1	7.3E-17
LILRA2	7.4E-03
HAVCR2	8.4E-14
SIRPA	1.6E-12
TGFBI	3.5E-19
SHP-2	4.3E-04
TGFBR1	1.1E-05
TGFB1	1.3E-07
PLAUR	6.2E-09

Log₂ fold difference

Siglec-10⁺ MDSCs vs. Siglec-10⁻ MDSCs

Gene	FDR
CMKLR1	1.5E-04
MERTK	4.1E-09
VSIG4	5.8E-08
FCGR3A	2.1E-15
SERPING1	2.6E-03
FCGR2BMSR1	2.3E-05
LILRB4	4.0E-03
LAIR1	1.4E-05
TGFBR1	4.6E-08
HMOX1	8.1E-03
CD47	3.8E-04
TGFBI	4.6E-04
CD163	1.6E-05
IL10RA	2.8E-03
MNDA	2.8E-02
FCN1	1.2E-03
VCAN	1.1E-04
CD36	6.5E-08
CSF3R	1.1E-06
FCAR	3.0E-05
SERPINB2	2.6E-02
	5.2E-03

Log₂ fold difference

Siglec-10⁺ DCs vs. Siglec-10⁻ DCs

Gene	FDR
ERLIN1	1.2E-02
HLA-DOB	3.4E-03
IL27RA	8.3E-03
FPR3	1.7E-04
AXL	2.0E-04
PDLIM2	2.0E-02
SLAMF8	9.4E-03
LILRB3	4.6E-02
TGFBI	1.0E-02
TGFBR2	2.1E-02
TGFB1	5.7E-03
HLA-DOA	9.8E-03
VSIR	2.0E-02
VCAN	4.2E-03
IL32	2.4E-04
CCL22	1.2E-07
CCR7	1.6E-12

Up-regulated
Down-regulated

Log₂ fold difference

e Siglec-10 expression in PDAC TME

P<0.0001
P<0.0001
P<0.0001
P=0.02
P=0.01
Borderline Resectable Locally-advanced Metastatic

f Siglec-10

P<0.05
Normal tissues PAAD tissues

g Siglec-10 — Pancreatic cancer

P = 0.0016
Low Siglec-10
High Siglec-10
Time (years)

Figure 2

Figure 2

a — AsPC-1 (PDAC); b — AsPC-1 (PDAC); c — MIA PaCa-2 (PDAC); d — MIA PaCa-2 (PDAC); e — PANC-1 (PDAC); f — PANC-1 (PDAC); g — SKOV3 (Ovarian cancer); h — SKOV3 (Ovarian cancer); i — OVCAR8 (Ovarian cancer); j — OVCAR8 (Ovarian cancer); k — MCF-7 (Breast cancer); l — MCF-7 (Breast cancer)

Columns: Control, Stained, Control, Stained

Y-axis: CD24

X-axis (left panels): Binding to recombinant Siglec-10 protein

X-axis (right panels): Binding to Siglec-10 protein with inactivating R119A mutation

Legend:
- Siglec-10 ligands⁺ cells
- CD24⁺ cells
- Cells expressing both Siglec-10 ligands and CD24

Figure 3

a

Anti-Fc antibody / HRP / Biotin-Tyramide / H2O2 / Siglec-10 recombinant protein Fc (or controls) / Biotinylation of Siglec-10 ligand by short-lived radical / Pull-down by streptavidin beads / Identification of glycoproteins by mass spec / PDAC cells

b

4,044 proteins were enriched compared to a no-protein control (FDR < 1%).

110 proteins were enriched compared to the Siglec-5 protein control (≥ 2 fold)

6 proteins are expressed at higher levels in PAAD tissues compared to normal tissues and are associated with poor survival

c

CD47 — No binding

CD59 — No binding

CD73 — No binding

ITGB6 — No binding

ITGA3
$k_{on} = 5.3 \times 10^5$ M^{-1} s^{-1}
$k_{off} = 0.0054$ s^{-1}
$K_D = 10$ nM
$R_{max} = 24$ RU

ITGB1
$k_{on} = 4.7 \times 10^4$ M^{-1} s^{-1}
$k_{off} = 2.7 \times 10^{-4}$ s^{-1}
$K_D = 5.8$ nM
$R_{max} = 35$ RU

d

Sialic acid levels (relative binding to SNA lectin)

$P = 0.02$ $P = 0.05$

ITGA3 protein: –Sialidase / +Sialidase
ITGB1 protein: –Sialidase / +Sialidase

e

Intact ITGA3
$k_{on} = 1.9 \times 10^5$ M^{-1} s^{-1}
$k_{off} = 6.2 \times 10^{-4}$ s^{-1}
$K_D = 3.3$ nM
$R_{max} = 194$ RU

Desialylated ITGA3 — No binding

f

Intact ITGB1
$k_{on} = 3.6 \times 10^4$ M^{-1} s^{-1}
$k_{off} = 4.6 \times 10^{-4}$ s^{-1}
$K_D = 13$ nM
$R_{max} = 120$ RU

Desialylated ITGB1 — No binding

Figure 4

Figure 5

Figure 6

Figure 7