Mechanistic Insights into Human Antimicrobial Peptide-Induced

Activation of a Broadly Conserved Bacterial Signaling System

Samuel A. Adeleye[1], Abraham F. Mechesso[2], Arpita Mukherjee[1], Guangshun Wang*[,2], Srujana

S. Yadavalli[1]*

[1]Waksman Institute of Microbiology and Department of Genetics, Rutgers University,

Piscataway, New Jersey 08854 USA

[2]Department of Pathology and Microbiology, College of Medicine, University of Nebraska

Medical Center, Omaha, Nebraska 68198 USA

*Corresponding authors: sam.yadavalli@rutgers.edu, gwang@unmc.edu

Short title: Peptide-induced activation of a bacterial signaling system

Classification: Biological Sciences, Microbiology

Keywords: Bacteria, antimicrobial peptides, two-component signaling systems, stress

response, cell elongation, reporter gene assays

Abstract

Antimicrobial peptides (AMPs) represent a promising class of therapeutics against bacterial pathogens. While their direct bactericidal mechanisms are well-characterized, how bacteria sense and respond to these peptides at sublethal concentrations remains poorly understood. Here, we investigate the activation of the *Escherichia coli* PhoQ-PhoP signaling system by the human antimicrobial peptide LL-37 and its derivatives (KR-12 and RI-10). We demonstrate that these peptides exhibit variable antimicrobial potency but surprisingly similar abilities to activate the PhoQ-PhoP pathway, indicating that signaling function is separable from bactericidal activity. Notably, sublethal concentrations of these peptides induce significant cell elongation, a phenotype dependent on PhoQ and mediated by the upregulation of QueE, which interferes with bacterial cell division. Contrary to the previous model suggesting peptides activate PhoQ passively by displacing its inhibitor MgrB, we observed enhanced cell elongation in *ΔmgrB* strains across all tested peptides, including RI-10, lacking antibacterial activity. Our findings suggest peptides actively stimulate PhoQ through a mechanism independent of MgrB dissociation, providing a more refined understanding of the peptide signaling through the PhoQ-PhoP system. These insights into bacterial adaptation mechanisms against host-derived peptides may guide the development of peptide therapeutics with enhanced efficacy against drug-resistant pathogens.

Significance statement

Antimicrobial peptides (AMPs) are promising alternatives to conventional antibiotics, yet how bacteria detect and respond to these host defense molecules remains poorly understood. This study investigates the interaction between bacterial sensing systems and AMPs at the genetic and molecular levels. Unlike other eukaryotes with multiple cathelicidin peptides, humans have only one cathelicidin that produces various active fragments through processing. Rather than creating multiple detectors, *E. coli* deploys an elegant solution, the PhoQ-PhoP signaling system that recognizes the conserved antimicrobial region shared by all LL-37 fragments. We demonstrate that peptides directly stimulate PhoQ independent of their bactericidal activity, even in the absence of inhibitor MgrB, inducing cell elongation. These insights may inform the development of effective peptide-based therapeutics against drug-resistant pathogens.

Introduction

When exposed to specific cationic antimicrobial peptides (AMPs) at sublethal concentrations, *Escherichia coli* activates the PhoQ/PhoP signal transduction pathway—a conserved two-component system crucial for stress adaptation in Gram-negative bacteria (1, 2). The PhoQ/PhoP signaling system is also an important pathway for adaptation and survival in conditions such as low Mg^{2+}, mildly acidic pH, and hyperosmolarity in *E. coli* and closely related enterobacteria (3, 4). Activated PhoQ-PhoP regulate numerous genes, including those responsible for remodeling the outer membrane, activation of efflux pumps, and proteolytic degradation of the peptides that help bacteria resist cationic AMPs (5–8). Strong activation of the PhoQ-PhoP system is observed when cells are treated with a human platelet-derived AMP (C18G), causing the upregulation of QueE, an enzyme primarily known for its role in the queuosine biosynthetic pathway that modifies specific tRNAs (2). Remarkably, elevated QueE levels lead to the accumulation of this protein at the bacterial division septum, where it interacts with the cell division machinery. This interaction disrupts normal septation processes downstream of Z-ring formation, resulting in inhibited cell division and the development of filamentous bacterial morphology. This mechanism represents a sophisticated bacterial stress response that temporarily halts proliferation when facing host antimicrobial challenges.

Human cathelicidin, LL-37, is one of the best-studied AMPs (9–11) with a broad spectrum of activity against bacteria, fungi, parasites, viruses, and cancer cells (11–15). LL-37 is known to activate the PhoQ-PhoP system and exert its antimicrobial action primarily by interacting with bacterial membranes, inserting itself into the lipid bilayer, and leading to membrane destabilization and the formation of pores (1, 16–18). This disruption results in the loss of membrane integrity, causing leakage of cellular contents and ultimately leading to bacterial cell death. LL-37's efficacy against pathogens is influenced by the composition of the growth

medium, particularly the presence of high salt concentrations and divalent cations (19). In our

current understanding, cationic antimicrobial peptides interact with the acidic patch in the

periplasmic sensing domain of PhoQ, displacing divalent cations (particularly Mg^{2+}) that

normally repress PhoQ activity, leading to PhoQ autophosphorylation and subsequent

phosphorylation of PhoP (1). Besides peptides and other signals, a 47 amino acid long small

membrane protein called MgrB interacts with and inhibits PhoQ by affecting its kinase activity

(20, 21). Specific residues in the transmembrane and periplasmic regions of MgrB are important

for the interaction with PhoQ (22). It is proposed that the sensing of AMPs by PhoQ at

physiological concentrations of magnesium is dependent on MgrB (23). A conformational

change in PhoQ upon AMP binding is thought to disrupt the PhoQ-MgrB complex, resulting in

PhoQ activation via derepression. This model suggests that AMPs play an inert role in

stimulating PhoQ.

Human cathelicidin LL-37 contains 37 amino acids and starts with a pair of leucine residues.

Triple-resonance NMR structural studies established that the C-terminal tail of LL-37 (residues

33-37) is disordered and not required for membrane-targeting mediated antimicrobial activity

(24). This led to the use of several derivatives of the LL-37 peptide from the central region for

therapeutic development (11, 24, 25). KR-12 is a 12-amino acid-long, truncated peptide derived

from human LL-37, which is toxic to bacteria but not human cells (24). It has been widely utilized

in peptide engineering (26). To effectively develop AMPs as novel peptide therapeutics, it is

imperative to study the bacterial response to these peptides, including the biochemical and

regulatory pathways that underlie this process. While there has been tremendous progress in

establishing the direct antimicrobial action of AMPs such as LL-37 against bacteria (27), much

less is known about the phenotypic changes elicited by these peptides at sub-lethal doses and

how their length and sequence contribute to bacterial response. In addition, it is unclear if AMPs

have a direct and active role in stimulating PhoQ.

To gain deeper insights into the stress response mechanism to AMPs in *E. coli*, we utilized LL-

37, its shortest anti-*E. coli* peptide KR-12, and an even shorter non-antibacterial peptide RI-10,

resulting from the truncation of the N-terminal lysine and C-terminal arginine of KR-12. In

particular, we first examined the antibacterial activity of LL-37 and its smaller fragments, KR-12

and RI-10, using different *E. coli* strains under physiologically relevant conditions. We

determined the killing kinetics and characterized the peptides using membrane permeabilization

and depolarization assays. Next, we investigated whether LL-37 and its fragments, at

concentrations below MIC, are effective in activating the PhoQ/PhoP signaling system by

analyzing both the changes in cell morphology and PhoP-dependent transcriptional reporter

activity as readouts. Finally, based on our analysis of cells lacking the PhoQ inhibitor, MgrB, we

propose a model of how peptides like LL-37 and its derivatives may interact with bacterial

histidine kinase and affect bacterial physiology. Our study sheds light on the molecular

mechanism of bacterial detection of biologically relevant peptide LL-37, which is known to play a

key role in human health via both antimicrobial and immune regulatory roles (9-12,27). Our

results reveal that the bacterial PhoQ-PhoP two-component system detects a variety of LL-37

antimicrobial fragments in a specific manner by recognizing the consensus core antimicrobial

region.

Results

Antibacterial activity of LL-37 and its derivatives against *E. coli*.

To better delineate the antibacterial activity of LL-37, KR-12, and RI-10 against *E. coli*, we

compared both their minimum inhibitory concentration (MIC) and minimum bactericidal

concentration (MBC) against four *E. coli* strains (MG1655, E423-17, E416-17, and 25922,

Tables 1, S1, S2) under three different media conditions: Mueller Hinton Broth (MHB), tryptic

soy broth (TSB), or supplemented minimal A medium (MinA). Gentamicin was included as a

non-peptide antibiotic control. LL-37 exhibited the highest potency with MIC values between 2

and 8 μM, depending on the media used (Table 1). Interestingly, the KR-12 peptide, which is

shorter than LL-37 by 25 amino acids, is also active against the *E. coli* isolates tested. The MIC

and MBC values for KR-12 are similar or 2-4 fold higher than those obtained for LL-37. The

values obtained in minimal medium were in a similar range as those obtained in other media,

especially for *E. coli* K-12 MG1655 strain (Tables 1, S1). The shortest LL-37 variant, RI-10, did

not show antibacterial activity against any of the isolates tested here across different media

conditions, or after dilution of MHB to 10%. From here on, for characterization of the

antibacterial and physiological effects of these peptides, we used the *E. coli* MG1655 strain

grown in the supplemented minimal medium. Next, we investigated the time-dependent killing

kinetics of LL-37, KR-12, and RI-10 to further substantiate their bactericidal properties using a

concentration corresponding to 2x MIC values (Figure S1). LL-37 (2× MIC) completely

eradicated the exponential phase of *E. coli* K-12 MG1655 in 1 hour, while KR-12 caused about a

4-log fold reduction in 2 hours. As expected, RI-10 displayed no activity at the concentration

tested. Together, these results show that full-length LL-37 and KR-12 exhibit antibacterial

properties against *E. coli* but not RI-10.

LL-37 and KR-12, but not RI-10, disrupt bacterial membranes.

Given that LL-37 targets bacterial membranes, we wondered if the shorter peptide variants, KR-

12 and RI-10, may retain and exhibit activity against *E. coli* membranes. To assess membrane

disruption, we conducted membrane permeabilization and depolarization experiments. We

analyzed membrane permeability by the uptake of a membrane-impermeable red dye,

propidium iodide (PI), either in the presence of a peptide, 0.1% Triton-X (positive control) or

PBS (untreated). Treatment with either LL-37 or KR-12 led to an increase in membrane

permeability in a dose-dependent manner, although LL-37 was more effective (Figure 1A-C). To

assay membrane depolarization, we used DiBAC$_4$(3), a probe for membrane potential, whose

fluorescence increases upon entering depolarized cells. As expected, LL-37 depolarized

membranes well with increasing concentration of the peptide (Figure 1D-F). Similarly, KR-12

also caused membrane depolarization but to a lower extent relative to LL-37. Consistent with

the data from MIC assay and killing kinetics, RI-10 did not have a discernible effect on

membrane permeability and depolarization. Together, these results suggest that KR-12, like LL-

37, targets the bacterial membrane as its mode of action.

Sub-MIC levels of LL-37 and KR-12 activate the PhoQ/PhoP signaling system and cause

cell elongation.

Cationic antimicrobial peptides, LL-37 and C18G activate the PhoQP pathway in *E. coli* and

related bacteria (1, 2). In addition, *E. coli* cells elongate upon exposure to a sub-lethal dose of

C18G, leading to filamentous growth (2). This filamentation is associated with conditions that

strongly activate PhoQ/PhoP. We asked if the truncated LL-37 derivatives retain the ability to

activate the PhoQP system. We used a promoter-reporter strain containing P$_{mgrB}$-*yfp* to measure

the transcriptional activity of the *mgrB* gene, whose expression is PhoQP-dependent. In parallel,

we observed the cells by phase contrast microscopy to test if the peptides induce any changes

in *E. coli* cell morphology. Our analysis revealed that there is a 3 to 3.5-fold increase in PhoQP-

regulated transcriptional activity when cells are treated with a sub-MIC dose of LL-37 at 1.25 µM

and KR-12 at 5 µM (Figure 2A). RI-10 showed weak activation of ~1.7-fold higher than the

untreated control. As the amino acid sequence of RI-10 (i.e., residues 19-28 of LL-37) is

included in both KR-12 (residues 18-29) and LL-37, we also asked how a different LL-37

segment, KE-10 (residues 10-19), with a minimal sequence overlap with KR-12 and RI-10

(Table S2), behaves. Like RI-10, KE-10 showed no antibacterial activity against *E. coli* (MIC >

64 µM in Table 1). In contrast to RI-10, the KE-10 peptide showed negligible activation of the

PhoQP system based on the transcription activity measurement of the *mgrB* gene, implying that

there is a requirement for a particular peptide sequence for this activation. Although LL-37 was

long known to activate the PhoQP system, its impact on *E. coli* cell morphology, especially at

concentrations below the MIC, has not been studied previously. Upon observation by phase

contrast microscopy, both LL-37 and KR-12-treated cells displayed elongation with

heterogeneous cell lengths (Figure 2B). The average cell lengths increased by ~1.5-fold for cells

exposed to LL-37 and KR-12 when compared to the control cells (Figure 2C). However, both RI-

10 and KE-10 treated cells did not show a significant change in cell length (Figure 2B-C).

Together, these data elucidate that LL-37, KR-12, and RI-10 are functional in stimulating the

PhoQP pathway and eliciting a physiological response, but not KE-10. However, only full-length

LL-37 and its minimal antibacterial peptide KR-12 are strong enough to cause morphological

changes similar to what was observed for the C18G peptide previously (2) , whereas two

shorter LL-37 segments (RI-10 and KE-10), which lost antibacterial activity, were unable to.

Cell elongation caused by sub-MIC levels of peptides is dependent on PhoQ and QueE.

To gain additional insight into the cell elongation caused by peptides, we examined the

morphology of the cells upon deletion of *phoQ*. As expected, Δ*phoQ* cells did not elongate when

treated with LL-37 and KR-12 (Figure 3A). There was no increase in PhoQP-dependent

transcriptional activity of P$_{mgrB}$, even in the presence of LL-37 and KR-12 in Δ*phoQ* cells (Figure

3B), and the average cell lengths were similar relative to the no-peptide control (Figure 3C),

confirming that the cell elongation phenotype associated with these AMPs is dependent on

PhoQ activation. In previous work, sub-MIC levels of C18G are shown to cause filamentation in

wild-type *E. coli* cells, which is mediated by increased expression of a PhoP-dependent gene

called *queE*, encoding a protein with dual activities in the inhibition of cell division and the

synthesis of queuosine tRNA modification (2, 28). We asked if the morphological phenotype

observed here for cells exposed to LL-37 and KR-12 is also mediated by QueE or through an

independent mechanism. Using a *queE* deletion strain, we carried out microscopy and

measured cell lengths and transcriptional reporter activity. Δ*queE* suppressed the elongation

phenotype observed when cells were treated with either LL-37 or KR-12 (Figures 3A, 2B). It is

worth noting that Δ*queE* cells grown in the presence of these peptides still display an increase in

PhoQP-regulated transcriptional reporter activity similar to the wild-type cells (Figures 3D, 2A).

Consistently, Δ*queE* cells exposed to peptides did not show a difference in the average cell

length relative to the controls (no peptide, KE-10, Figure 3E). Overall, filamentation and

heterogeneity in cell size observed upon treatment with LL-37 and KR-12 peptides are both

PhoQ- and QueE-dependent.

Deletion of PhoQ inhibitor, MgrB, increases cell responsiveness to LL-37 and its

derivatives.

E. coli cells, carrying an *mgrB* deletion, show an overall, robust increase in PhoQP-controlled

gene expression (20). These cells experience higher stimulation of PhoQ under magnesium-

limiting conditions, similar to the strong activation of this system observed in wild-type cells

grown in the presence of C18G at intermediate magnesium concentrations (2). Consequently,

Δ*mgrB* cells grow as long filaments, and this phenotype is dependent on upregulation of QueE

expression by PhoQ/PhoP (2). It is proposed that MgrB mediates the sensing of cationic AMPs

by PhoQ at intermediate concentrations of magnesium, therefore, LL-37, KR-12, and RI-10 may

activate PhoQ only when MgrB is present. In this case, we expect Δ*mgrB* cells to show no

significant change in either the PhoP-dependent transcriptional reporter activity or cell

morphology when treated with peptides vs. no treatment. We examined how LL-37 and its

derivatives influence the activity of the PhoQP system and cell lengths in a Δ*mgrB* strain. As

expected, in the absence of peptides, Δ*mgrB* cells showed an increase in P$_{mgrB}$ transcription

relative to the wild-type cells under the same growth conditions (Figures 4A-B). When the AMPs

were added to Δ*mgrB* cells, there was a modest increase of ~1.4-fold in the PhoQP-regulated

transcriptional reporter activity for LL-37, as observed previously using the same media

conditions carrying 0.1 mM Mg^{2+} (Figure 4A) (20). The addition of peptides KR-12, RI-10, and

KE-10 did not lead to a substantial increase in reporter activity, suggesting that PhoP-mediated

transcription is not sensitive in response to the two peptides in cells lacking MgrB. This result is

consistent with a previous study where ∆*mgrB* cells did not show an increase in PhoP-regulated

transcription in the presence of another AMP, C18G (23).

To investigate this further, we analyzed the morphology of ∆*mgrB* cells in the presence of

peptides, wherein cell elongation is PhoQ-dependent and a direct function of strong PhoQP

activation. As expected, we noticed filamentation in ∆*mgrB* cells treated with LL-37 and KR-12,

but only short, typical cells in the KE-10-treated or no-peptide control cells, under this growth

condition with 0.1 mM Mg^{2+} (Figure 4C). It was intriguing, however, to see that RI-10-treated

∆*mgrB* cells also displayed elongation, suggesting that RI-10 may interact more effectively with

PhoQ to activate it, in the absence of MgrB. In other words, loss of the inhibitor MgrB appears to

unmask the effect of RI-10 peptide on PhoQ stimulation, resulting in an evident cell elongation

phenotype. Interestingly, cell size quantification revealed greater heterogeneity and higher

average cell lengths across all three peptides for ∆*mgrB* of 6-8 µm (Figure 4D). The mean cell

lengths for LL-37, KR-12, and RI-10-treated ∆*mgrB* cells were higher than the corresponding

values obtained for the wild type by 2.5-, 2.3-, and 1.9-fold, respectively; however, there is no

difference in cell length in the KE-10-treated cells (Figures 4D-E), indicating a central role of the

RI-10 containing LL-37 segments in activation in the absence of *mgrB*.

It is known that high stimulation of PhoQ from exposure to LL-37 results in differential regulation

of specific PhoP-regulated promoters, including that of *mgrB* (29). Also, upon activation by LL-

37, a mutated *mgrB-t7a* promoter (with a T7A substitution in the PhoP-binding site) shows a

higher fold change in transcriptional activity relative to the wild-type *mgrB* promoter. To

substantiate whether there may be an increased transcriptional response to peptide treatment

via this promoter, we analyzed wild-type and ∆*mgrB* strains carrying the *mgrB-t7a* promoter. We

observed a significant increase in cell size heterogeneity in wild-type cells treated with LL-37

and KR-12, and ∆*mgrB* cells treated with LL-37, KR-12, and RI-10 (Figure S2). The cells did not

display an increase in cell length when treated with control peptide KE-10 or in the absence of a

peptide. If the exacerbated cell elongation observed in peptide-treated Δ*mgrB* cells is indeed

due to a PhoQP-dependent mechanism that is mediated by QueE, we should see this

phenotype suppressed upon deletion of *queE*. Consistent with our hypothesis, Δ*mgrB* Δ*queE*

cells treated with LL-37, KR-12, and RI-10 did not show an increase in PhoQ-dependent

transcriptional reporter activity and displayed a loss of filamentation and a decrease in cell size

heterogeneity (Figure S3). Together, these data suggest that LL-37 and its derivatives bind and

actively stimulate PhoQ. In the absence of MgrB, these peptides further enhance PhoQ

activation. Importantly, cell elongation serves as a sensitive phenotype for peptide-mediated

PhoQ stimulation in *E. coli*. Wild-type cells exposed to sub-MIC peptide concentrations exhibit

moderate elongation, reflecting intermediate PhoQ activity, while Δ*mgrB* cells display

significantly increased cell lengths, indicating robust PhoQ activation (Figure 4E).

281 **Discussion**

Despite extensive investigation, fundamental questions remain about the molecular

mechanisms of peptide-mediated signal transduction by the PhoQ-PhoP system. Moreover, the

effects of PhoPQ activation by cationic AMPs in cells and the physiological importance of these

interactions have not been elucidated. We address two critical questions in this work: (i) Can

peptides lacking potent antimicrobial activity still interact with PhoQ and act as a signal for this

pathway? Using LL-37 and derivatives with variable MICs and antibacterial activities against *E.*

coli cells, we show that the signaling function of peptides through the PhoQ-PhoP system is

separable from their antimicrobial activity (Figure 5A), suggesting that bacteria may use this

pathway primarily as a detection mechanism for host environments rather than simply as a

response to membrane damage. (ii) Do the peptides actively stimulate PhoQ, or do they exert

their effect on PhoQ passively by displacing the inhibitor protein, MgrB? Many studies on

peptide-induced signaling of PhoQ measure the activity of peptides using PhoP-dependent

transcriptional reporters as a proxy. In agreement with the previous studies, upon exposure to

peptides, we do not observe a significant increase in PhoQP-driven transcriptional reporter

activity in *ΔmgrB* cells relative to the wild type. It is possible that the derepression of PhoP-

regulated gene expression in the absence of MgrB could lead to lower sensitivity of the reporter

signal, masking small increments in the amplitude. While such reporters are incredibly useful in

determining the activation status of this pathway and measuring the transcription levels, we

have a second, sensitive readout – a cell elongation or filamentation phenotype in response to

peptide stress in *E. coli*. Based on our microscopy and cell length measurements, we find that

cells elongate significantly upon exposure to sublethal concentrations of peptides, LL-37 and

KR-12, even though the promoter reporters do not show a significant change in their activities.

The extent of cell elongation may differ in response to specific peptides, as we note that

filamentation induced by an AMP, C18G, in wild-type *E. coli* cells is quite robust, with an

average cell length of > 20 μm (2). Upon deletion of MgrB, the average cell lengths increased

further, and even RI-10-treated cells displayed cell elongation; this observation suggests that

AMPs may activate PhoQ directly in addition to causing MgrB dissociation as proposed

previously (23). Our findings elucidate PhoQ-dependent effects of these peptides and support a

model where peptides play an active and direct role in PhoQ activation (Figure 5B).

Our understanding of what characteristics of AMPs are required to activate the PhoP-PhoQ

system is still evolving. A previous report found a correlation between the extent of positive

charge, hydrophobicity, and amphipathicity of an AMP and its ability to activate PhoQ-PhoP (8).

Intriguingly, the authors discovered that even a peptide with minimal positive charge (+1) but

containing histidines was highly potent at activating the system, suggesting the existence of an

activation mechanism more complex than thought. A later study with 117 AMPs from the

Antimicrobial Peptide Database (30) showed that PhoQ is activated by a diverse set of AMPs,

including the human cathelicin LL-37 and non-cathelicidin peptides that carry either an α-helical

or a combination of α-helical/β-sheet structural regions (31). In our work, we see that LL-37

derivatives induce PhoQ-dependent morphological changes, and KR-12, the smallest

antibacterial peptide, is also the smallest peptide sequence for the PhoPQ induction. Further,

RI-10, but not KE-10, became responsive upon the deletion of *mgrB*. Interestingly, LL-25, the N-

terminal segment of LL-37 which does not kill bacteria under normal conditions (31), does not

activate PhoPQ based on the recent human AMPs screen study although it contains partial

sequence of KR-12 (30). LL-25 can be obtained by extending the sequence of KE-10 from both

ends (Tables S2, S3). Together, the data from the current study suggest that part of the LL-37

amino acid sequence that overlaps with both KR-12 and RI-10 is likely mediating the interaction

with PhoQ. More broadly, our findings imply that the bacterial molecular detection machinery is

elegantly tuned to specifically target the core antimicrobial region of human antimicrobial

peptide LL-37 but not fragments outside this region. When broadly viewed from the host-

pathogen interaction at the molecular level, bacterial detection of the core antimicrobial region of

human cathelicidin LL-37 (24) is remarkably efficient since this region is commonly shared by all

the active fragments of LL-37 naturally produced in human skin (Table S3), including KS-27,

KR-20, KS-30, RK-31, and LL-29 (32). Indeed, all these human cathelicidin fragments

responded to the PhoPQ system based on a recent screening of human AMPs registered in the

APD (30). Collectively, these findings underscore that the bacterial PhoQ-PhoP system likely

senses host defense peptides based on evolutionarily conserved structural and chemical

properties independent of their antimicrobial activity, allowing bacteria to mount preemptive

responses rather than simply react to membrane damage.

Finally, the dramatic rise of antimicrobial resistance is a major public health concern. AMPs and

other peptide derivatives are relatively small, typically comprised of around 10 to 50 amino acids

(30), and constitute a key component of host innate immunity for defense against microbial

pathogens (32, 33). These cationic peptides are relatively cheap to produce, have good thermal

stability and water solubility, and exhibit a broad spectrum of activity against bacteria and fungi,

making them promising candidates for drug development (34–38). Studying the bacterial

responses to antimicrobial peptides in general provides us with a fundamental understanding of

how pathogens can sense, adapt, and develop tolerance to these host defense molecules and

guide the strategic design of novel therapeutics with enhanced efficacy to overcome the growing

antibiotic resistance problem.

Materials and Methods

Strains

Details of the strains used in this study are provided in Table S4. AML20, JNC21, NRS5, NRS6,

SAM31, TIM92, TIM100, and TIM166 are *E. coli* K-12 MG1655 derivatives. Additionally, clinical

strains of *E. coli*, E423-17, E416-17, and ATCC strain 25922 were used for specific assays as

indicated. NRS6 was made by transducing the KEIO strain JW1815 into the TIM148 and

TIM166 strains, respectively, using $P1_{vir}$.

Media, reagents, and growth conditions

Routine bacterial growth on solid agar was performed using LB Miller medium (IBI Scientific)

containing 1.5% bacteriological grade agar (VWR Lifesciences) at 37°C. Liquid cultures were

grown at 37°C with aeration in either LB miller medium or minimal A medium (MinA) (39)

(K_2HPO_4 (10.5 g l^{-1}), KH_2PO_4 (4.5 g l^{-1}), $(NH_4)_2SO_4$ (1.0 g l^{-1}), Na_3 citrate.$2H_2O$ (0.5 g l^{-1}),

supplemented with 0.1% casamino acids, 0.2% glucose and 1 mM $MgSO_4$ unless otherwise

indicated. Throughout this study, supplemented MinA, as indicated above, will be referred to as

a minimal medium. In addition, Mueller-Hinton broth (MHB, catalog info) and tryptic soy broth

(TSB, catalog info) were used for minimum inhibitory/bactericidal concentration (MIC/MBC)

assays. Peptides and/or AMPs used in the study are indicated in Table S2.

For the measurement of cell length and transcriptional reporter activity, overnight cultures of

SAM31, TIM92, TIM100, AML20, SAA58, and JNC21 were grown in supplemented MinA

containing 1 mM $MgSO_4$. Saturated cultures were back diluted 1:500 into 5 ml supplemented

MinA with 0.1 mM Mg^{2+} and grown for 2 hours in a shaker at 300 rpm. Each culture was split

into two tubes, with one tube designated as a no-peptide control and the second tube where a

test peptide was added. LL-37, KR-12, RI-10, and KE-10 were added to a final concentration of

1.25 µM, 5 µM, 5 µM, and 5 µM, respectively. These cultures were further grown for 4 hours at

37°C in a roller drum at 60 rpm.

Microscopy and image analysis

Cell morphology was observed and single-cell fluorescence to measure transcriptional reporter

activity was performed using phase contrast and fluorescence microscopy as previously

described (2). 3 µl of the cells were immobilized on a 1% agarose pad and observed using a

Nikon TiE fluorescent microscope with a TI2-S-HU attachable mechanical stage and a perfect

focus system (PFS). The images were captured using a Teledyne Photometrics Prime 95B

sCMOS camera with 1x1 binning. YFP and CFP fluorescence was imaged using a 100 ms

exposure time at 20% intensity. Image acquisition was done using Metamorph software version

7.10.3.279 (Molecular Devices, Sunnyvale, CA). The background fluorescence was determined

using MG1655 cells grown under the same conditions. The average cell fluorescence was

quantified using ImageJ (40) and the MicrobeJ plugin (41). Data from independent replicates

were plotted using R.

Minimum inhibitory/bactericidal concentration (MIC/MBC) testing

The antibacterial activity of the peptides against *E. coli* MG1655, E423-17, E416-17, and 25922

strains was evaluated using a previously established protocol (42). Briefly, a peptide

concentration gradient with twofold dilution was prepared in the 96-well polystyrene microplate

at 10 µl per well. Bacteria were grown to the exponential phase in MHB, TSB, or supplemented

MinA. It was then diluted to ~10^5 CFU/ml in MHB, TSB, or supplemented MinA with 0.1 mM

$MgSO_4$, and partitioned into the 96-well microplate at 90 µl per well. After overnight incubation at

37°C, readings were taken at 600 nm using a ChroMate 4300 Microplate Reader at (GMI,

Ramsey, MN). The minimum inhibitory concentration (MIC), represented as ranges, was

determined for wells without bacterial growth. For minimum bactericidal concentration (MBC)

determination, aliquots of 30 µl from growth-absent wells were plated on MH agar plates, and

the lowest peptide concentration with no isolated colony was considered the MBC. Results were

expressed in µM.

Minimum inhibitory/bactericidal concentration (MIC/MBC) testing

The antibacterial activity of the peptides against *E. coli* MG1655, E423-17, E416-17, and 25922

strains was evaluated using a previously established protocol (42). Briefly, a peptide

concentration gradient with twofold dilution was prepared in the 96-well polystyrene microplate

at 10 µl per well. Bacteria were grown to the exponential phase in MHB, TSB, or supplemented

MinA. It was then diluted to ~10^5 CFU/ml in MHB, TSB, or supplemented MinA with 0.1 mM

$MgSO_4$, and partitioned into the 96-well microplate at 90 µl per well. After overnight incubation at

37°C, readings were taken at 600 nm using a ChroMate 4300 Microplate Reader at (GMI,

Ramsey, MN). The minimum inhibitory concentration (MIC), represented as ranges, was

determined for wells without bacterial growth. For minimum bactericidal concentration (MBC)

determination, aliquots of 30 µl from growth-absent wells were plated on MH agar plates, and

the lowest peptide concentration with no isolated colony was considered the MBC. Results were

expressed in µM.

Membrane permeabilization

The peptide ladder was prepared as described for the antibacterial assay in a black COSTAR

96-well plate. Propidium iodide (PI) (MP Biomedicals, Solon, OH) was prepared in the dark and

dissolved in DMSO (Thermo Fisher Scientific, NY) to 20 mM. This PI stock solution was further

diluted to 1 mM with water and 2 µl of 1 mM PI was added to each well. An overnight culture of

E. coli K12 MG1655 was inoculated into 1xMinA supplemented with 0.2% glucose, 0.1%

casamino acids, 0.1mM $MgSO_4$, and grown to exponential phase. It was then diluted to OD_{600}

0.11 in the same media, and 88 µl was added to each well. The plate was incubated at 37℃ with

continuous shaking in a FLUOstar Omega (BMG LABTECH, NC) microplate reader. The sample

was read every 5 minutes for 24 cycles with excitation and emission wavelengths of 584 nm and

620 nm, respectively.

Killing kinetics

E. coli K12 MG1655 was grown to the exponential phase (OD_{600} ~ 0.3) in 1xMinA supplemented

with 0.2% glucose, 0.1% casamino acids, 1 mM $MgSO_4$. The culture was diluted to OD_{600} 0.001

(~10^5 CFU/ml) in 1xMinA supplemented with 0.2% glucose, 0.1% casamino acids, 0.1mM

$MgSO_4$. One milliliter of bacterial suspension was then mixed with LL-37 (4 µM, 2× MIC), KR-12

(8 µM, 2× MIC) or RI-10 (32 µM) and incubated at 37°C. After 15, 30, 60, 90, and 120 minutes

of incubation, 50 µl was taken and serially diluted with 1×PBS. Then, 50 µl of the diluted

suspension was plated on MH agar Agar plates. The plates were incubated overnight at 37°C

for bacterial CFU determination.

Membrane depolarization

An overnight culture of *E. coli* K12 MG1655 was inoculated into 1xMinA supplemented with

0.2% glucose, 0.1% casamino acids, 1 mM $MgSO_4$, and grown to exponential phase. Bacteria

were washed with 1×PBS, re-suspended in twice the volume of 1×PBS containing 25 mM

glucose, and incubated at 37°C for 15 minutes. Then, 500 nM (final concentration) of the

DiBAC4 (3) bis-(1,3-dibutylbarbituric acid) trimethine oxonol (ANASPEC, CA) was added and

vortexed gently. Aliquots of 90 µl of the energized bacteria solution were loaded to the 96 well

plates (Corning COSTAR, AZ) and placed in a FLUOstar Omega microplate reader (BMG

LABTECH, NC). Fluorescence was read for 15 min at excitation and emission wavelengths of

485 nm and 520 nm, respectively. Then, 10 µl of peptide solutions were added and fluorescence

readings were recorded for 45 min. Triton X-100 (0.1%) was used as a positive control.

Acknowledgments

The authors thank Dr. Mark Goulian for sharing transcriptional reporter strains for *mgrB*, Dr.

Nitish Sharma for preparing *ΔmgrB* strains carrying the reporters, Dr. Premal Shah for critical

feedback and assistance with data visualization and figure creation, Dr. Manuela Roggiani for

critical feedback on the manuscript, and members of the Yadavalli and Wang labs for helpful

discussions.

Funding

S.S.Y. is supported by the National Institutes of Health - National Institute of General Medical

Sciences (NIH-NIGMS) ESI-MIRA R35 GM147566 and institutional start-up funds from Rutgers.

A.M. is supported by the Waksman Institute Busch Predoctoral fellowship (2024-25) at Rutgers.

G.W. is supported by the NIH/NIAID R56 AI175209. The funders did not play any role in the

study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Author contributions

458   S.S.Y. and G.W. conceived the experiments. All authors planned the experiments, designed the

459   methodology, and performed data analysis. S.A.A., A.F.M., and A.M. performed the

460   experiments. All authors contributed to the writing and editing of the manuscript.

Competing interests

463   The authors declare no competing financial interests.

Materials and resource availability

Requests for strains and plasmids generated in this study should be directed to and will be

fulfilled by the lead contact, Srujana S. Yadavalli (sam.yadavalli@rutgers.edu). Further

information and requests for resources and reagents should be directed to and will be fulfilled by

Srujana S. Yadavalli (sam.yadavalli@rutgers.edu) and Guangshun Wang (gwang@unmc.edu).

Data availability

The main manuscript and its supporting files include all data generated or analyzed during this

study.

References

1. M. W. Bader, *et al.*, Recognition of antimicrobial peptides by a bacterial sensor kinase. *Cell* **122**, 461–
472 (2005).

2. S. S. Yadavalli, *et al.*, Antimicrobial peptides trigger a division block in Escherichia coli through
stimulation of a signalling system. *Nat. Commun.* **7**, 12340 (2016).

3. E. A. Groisman, The pleiotropic two-component regulatory system PhoP-PhoQ. *J. Bacteriol.* **183**,
1835–1842 (2001).

4. J. Yuan, F. Jin, T. Glatter, V. Sourjik, Osmosensing by the bacterial PhoQ/PhoP two-component
system. *Proc. Natl. Acad. Sci. U. S. A.* **114**, E10792–E10798 (2017).

5. S. Matamouros, S. I. Miller, S. Typhimurium strategies to resist killing by cationic antimicrobial
peptides. *Biochim. Biophys. Acta* **1848**, 3021–3025 (2015).

6. Z. D. Dalebroux, S. I. Miller, Salmonellae PhoPQ regulation of the outer membrane to resist innate
immunity. *Curr. Opin. Microbiol.* **17**, 106–113 (2014).

7. R. K. Ernst, T. Guina, S. I. Miller, Salmonella typhimurium outer membrane remodeling: role in
resistance to host innate immunity. *Microbes Infect.* **3**, 1327–1334 (2001).

8. T. Shprung, A. Peleg, Y. Rosenfeld, P. Trieu-Cuot, Y. Shai, Effect of PhoP-PhoQ activation by broad
repertoire of antimicrobial peptides on bacterial resistance. *J. Biol. Chem.* **287**, 4544–4551 (2012).

9. M. F. Burton, P. G. Steel, The chemistry and biology of LL-37. *Nat. Prod. Rep.* **26**, 1572–1584 (2009).

10. D. Vandamme, B. Landuyt, W. Luyten, L. Schoofs, A comprehensive summary of LL-37, the factotum
human cathelicidin peptide. *Cell. Immunol.* **280**, 22–35 (2012).

11. G. Wang, *et al.*, Design of antimicrobial peptides: Progress made with human cathelicidin LL-37. *Adv.*
Exp. Med. Biol. **1117**, 215–240 (2019).

12. F. Lu, Y. Zhu, G. Zhang, Z. Liu, Renovation as innovation: Repurposing human antibacterial peptide
LL-37 for cancer therapy. *Front. Pharmacol.* **13**, 944147 (2022).

13. K. Niemirowicz, *et al.*, Magnetic nanoparticles enhance the anticancer activity of cathelicidin LL-37
peptide against colon cancer cells. *Int. J. Nanomedicine* **10**, 3843–3853 (2015).

14. K. Kuroda, K. Okumura, H. Isogai, E. Isogai, The human cathelicidin antimicrobial peptide LL-37 and
mimics are potential anticancer drugs. *Front. Oncol.* **5**, 144 (2015).

15. S. Tripathi, *et al.*, Antiviral activity of the human cathelicidin, LL-37, and derived peptides on seasonal
and pandemic influenza A viruses. *PLoS One* **10**, e0124706 (2015).

16. I. Ginsburg, Bactericidal cationic peptides can also function as bacteriolysis-inducing agents
mimicking beta-lactam antibiotics?; it is enigmatic why this concept is consistently disregarded. *Med.*
Hypotheses **62**, 367–374 (2004).

17. Y. Zhang, J. Lakshmaiah Narayana, Q. Wu, X. Dang, G. Wang, Structure and activity of a selective
antibiofilm peptide SK-24 derived from the NMR structure of human cathelicidin LL-37.
Pharmaceuticals (Basel) **14**, 1245 (2021).

18. K. A. Henzler Wildman, D.-K. Lee, A. Ramamoorthy, Mechanism of lipid bilayer disruption by the
human antimicrobial peptide, LL-37. *Biochemistry* **42**, 6545–6558 (2003).

19. J. Turner, Y. Cho, N. N. Dinh, A. J. Waring, R. I. Lehrer, Activities of LL-37, a cathelin-associated
antimicrobial peptide of human neutrophils. *Antimicrob. Agents Chemother.* **42**, 2206–2214 (1998).

20. A. M. Lippa, M. Goulian, Feedback inhibition in the PhoQ/PhoP signaling system by a membrane
peptide. *PLoS Genet.* **5**, e1000788 (2009).

21. M. E. Salazar, A. I. Podgornaia, M. T. Laub, The small membrane protein MgrB regulates PhoQ
bifunctionality to control PhoP target gene expression dynamics. *Mol. Microbiol.* **102**, 430–445
(2016).

22. S. S. Yadavalli, *et al.*, Functional determinants of a small protein controlling a broadly conserved
bacterial sensor kinase. *J. Bacteriol.* **202** (2020).

23. S. Jiang, *et al.*, The inhibitory mechanism of a small protein reveals its role in antimicrobial peptide
sensing. *Proc. Natl. Acad. Sci. U. S. A.* **120**, e2309607120 (2023).

24. G. Wang, Structures of human host defense cathelicidin LL-37 and its smallest antimicrobial peptide
KR-12 in lipid micelles. *J. Biol. Chem.* **283**, 32637–32643 (2008).

25. K. Chen, W. Gong, J. Huang, T. Yoshimura, J. M. Wang, The potentials of short fragments of human
anti-microbial peptide LL-37 as a novel therapeutic modality for diseases. *Front. Biosci. (Landmark
Ed.)* **26**, 1362–1372 (2021).

26. J. Lakshmaiah Narayana, *et al.*, Origami of KR-12 designed antimicrobial peptides and their potential
applications. *Antibiotics (Basel)* **13**, 816 (2024).

27. N. Mookherjee, M. A. Anderson, H. P. Haagsman, D. J. Davidson, Antimicrobial host defence
peptides: functions and clinical potential. *Nat. Rev. Drug Discov.* **19**, 311–332 (2020).

28. S. A. Adeleye, S. S. Yadavalli, Queuosine biosynthetic enzyme, QueE moonlights as a cell division
regulator. *PLoS Genet.* **20**, e1011287 (2024).

29. T. Miyashiro, M. Goulian, Stimulus-dependent differential regulation in the Escherichia coli PhoQ
PhoP system. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 16305–16310 (2007).

30. G. Wang, X. Li, Z. Wang, APD6: the antimicrobial peptide database is expanded to promote research
and development by deploying an unprecedented information pipeline. *Nucleic Acids Res.* **in Press**
(2025).

31. K. R. Brink, *et al.*, An E. coli display method for characterization of peptide-sensor kinase
interactions. *Nat. Chem. Biol.* **19**, 451–459 (2023).

32. G. Wang, *et al.*, Structure, dynamics, and antimicrobial and immune modulatory activities of human
LL-23 and its single-residue variants mutated on the basis of homologous primate cathelicidins.
Biochemistry **51**, 653–664 (2012).

33. M. Murakami, B. Lopez-Garcia, M. Braff, R. A. Dorschner, R. L. Gallo, Postsecretory processing
generates multiple cathelicidins for enhanced topical antimicrobial defense. *J. Immunol.* **172**, 3070–
3077 (2004).

34. G. Laverty, S. P. Gorman, B. F. Gilmore, The potential of antimicrobial peptides as biocides. *Int. J.*
Mol. Sci. **12**, 6566–6596 (2011).

35. L. Segev-Zarko, M. L. Mangoni, Y. Shai, "Antimicrobial peptides: multiple mechanisms against a
variety of targets" in *Antimicrobial Peptides: Discovery, Design and Novel Therapeutic Strategies*,
(CABI, 2017), pp. 119–134.

36. R. E. W. Hancock, K. L. Brown, N. Mookherjee, Host defence peptides from invertebrates--emerging
antimicrobial strategies. *Immunobiology* **211**, 315–322 (2006).

37. T. L. Raguse, E. A. Porter, B. Weisblum, S. H. Gellman, Structure-activity studies of 14-helical
antimicrobial beta-peptides: probing the relationship between conformational stability and
antimicrobial potency. *J. Am. Chem. Soc.* **124**, 12774–12785 (2002).

38. D. L. Lee, R. S. Hodges, Structure-activity relationships of de novo designed cyclic antimicrobial
peptides based on gramicidin S. *Biopolymers* **71**, 28–48 (2003).

39. J. H. Miller, *A Short Course in Bacterial Genetics*, 1987th Ed. (Cold Spring Harbor Laboratory Press,
1992).

40. C. A. Schneider, W. S. Rasband, K. W. Eliceiri, NIH Image to ImageJ: 25 years of image analysis.
Nat. Methods **9**, 671–675 (2012).

41. A. Ducret, E. M. Quardokus, Y. V. Brun, MicrobeJ, a tool for high throughput bacterial cell detection
and quantitative analysis. *Nat. Microbiol.* **1**, 16077 (2016).

42. B. Mishra, T. Lushnikova, R. M. Golla, X. Wang, G. Wang, Design and surface immobilization of short
anti-biofilm peptides. *Acta Biomater.* **49**, 316–328 (2017).

Figure legends

Figure 1. Membrane permeabilization (A-C) and depolarization (D-F) activity of a series of

LL-37, KR-12, and RI-10 in *E. coli* K-12 MG1655. The assay was performed in supplemented

MinA with 0.1 mM $MgSO_4$. RFU: Relative fluorescence intensity. Data are represented as mean

± range for two independent replicates.

Figure 2. PhoQP-regulated transcription and cell morphology of *E. coli* cells treated with

sub-MIC levels of antimicrobial peptides. (A) PhoQP-dependent promoter P*mgrB*-*yfp* reporter

activity, **(B)** phase contrast micrographs, and **(C)** quantification of cell lengths of *E. coli* MG1655

(TIM92) cells grown in the presence of sub-MIC levels of antimicrobial peptides LL-37 (1.25

µM), KR-12 (5 µM), RI-10 (5 µM), and KE-10 (5 µM) in supplemented MinA with 0.1 mM $MgSO_4$.

Data are obtained from at least three biological replicates; mean and median values are shown

in red text and black bars, respectively. The number of cells analyzed is indicated for each

sample. P-values indicate the results of a Student's t-test: ****$P \leq 0.0001$, *$P \leq 0.05$, and "ns" =

$P > 0.05$. Scale bar = 5 µm.

Figure 3. PhoQP-regulated transcription and cell morphology of $\Delta queE$ and $\Delta phoQ$ cells

treated with sub-MIC levels of antimicrobial peptides. (A) Phase contrast micrographs, **(B,**

D) PhoQP-dependent promoter P_{mgrB}-*yfp* reporter activity, and **(C, E)** quantification of cell

lengths of $\Delta phoQ$ (TIM100) cells and $\Delta queE$ (SAM31) cells grown in the presence of sub-MIC

levels of the antimicrobial peptides LL-37 (1.25 µM), KR-12 (5 µM), RI-10 (5 µM), and KE-10 (5

µM) in supplemented MinA with 0.1 mM $MgSO_4$. Data are obtained from at least three biological

replicates; mean and median values are shown in red text and black bars, respectively. The

number of cells analyzed is indicated for each sample. P-values indicate the results of a

Student's t-test: ****$P \leq 0.0001$, ***$P \leq 0.001$, *$P \leq 0.05$, and "ns" = $P > 0.05$. Scale bar = 5 µm.

Figure 4. PhoQP-regulated transcription and cell morphology of $\Delta mgrB$ cells treated with

sub-MIC levels of antimicrobial peptides. (A) PhoQP-dependent promoter P_{mgrB}-*yfp* reporter

activity in $\Delta mgrB$ (AML20) cells, **(B)** comparison of PhoQP-dependent promoter P_{mgrB}-*yfp*

reporter activity in $\Delta mgrB$ (AML20) cells with that of WT *E. coli* (TIM92) from Fig. 2A, and **(C)**

phase contrast micrographs of WT *E. coli* (TIM92) and $\Delta mgrB$ (AML20) cells, **(D)** quantification

of cell lengths of $\Delta mgrB$ (AML20) cells, and **(E)** comparison of cell lengths in $\Delta mgrB$ (AML20)

cells with that of WT *E. coli* (TIM92) from Fig. 2C. Cells were grown in the presence of sub-MIC

levels of the antimicrobial peptides LL-37 (1.25 µM), KR-12 (5 µM), RI-10 (5 µM), and KE-10 (5

µM) in supplemented MinA with 0.1 mM $MgSO_4$. Arrows in panel (E) indicate the PhoQ

activation status. Data are obtained from at least three biological replicates; mean and median

values are shown in red text and black bars, respectively. The number of cells analyzed is

indicated for each sample. P-values indicate the results of a Student's t-test: ****$P \leq 0.0001$, **P

≤ 0.01, *$P \leq 0.05$, and "ns" = $P > 0.05$. Scale bar = 5 µm.

Figure 5. Schematic overview of peptide stress response through the *E. coli* PhoQ-PhoP

signaling system. (A) At high concentrations, antimicrobial peptides affect membrane integrity

and cause damage, leading to cell death. At sub-MIC levels, peptides are sensed by the

signaling systems, such as PhoQ/PhoP, which control the gene expression under peptide

stress, leading to physiological changes, including cell elongation and modification of the outer

cell membrane. **(B)** At physiological magnesium concentration, PhoQ is strongly stimulated by

specific cationic peptides, and some PhoQ molecules are inhibited by the small protein

regulator, MgrB, leading to an intermediate level of PhoQ activation. In the absence of MgrB,

PhoQ is not inhibited, and when cells are exposed to peptides, PhoQ is strongly activated,

resulting in maximal activation of this signaling system.

Tables and Figures

Table 1. Minimum inhibitory concentration (MIC) and minimum bactericidal concentration

(MBC) values for LL-37 and its derivatives against *E. coli* K-12 MG1655. MIC values (µM) of

antimicrobial peptide LL-37 and its derivatives KR-12, RI-10, and KE-10 against *E. coli* K-12

MG1655 in supplemented minimal medium (Min A), Mueller Hinton broth (MHB), and tryptic soy

broth (TSB). The corresponding MBC values (µM) are shown within parentheses. Gentamicin is

included as a bactericidal antibiotic control. Data represent values obtained from three

independent biological replicates.

Strain	Media	LL-37	KR-12	RI-10	KE-10	Gentamicin
E. coli K-12 MG1655	Min A	2 (2)	8-16 (16)	>64 (>64)	>64 (>64)	0.5-1 (1)
	MHB	8 (8)	16-32 (32)	>64 (>64)	>64 (>64)	0.5-1 (1)
	10% MHB	2 (4)	2 (2)	>64 (>64)	>64 (>64)	ND
	TSB	8 (16)	16 (32)	>64 (>64)	>64 (>64)	1-2 (2)
E. coli K-12 MG1655 Δ*mgrB*	Min A	2 (2)	8 (16)	>64 (>64)	>64 (>64)	-
	MHB	8 (8)	8-16 (16)	>64 (>64)	>64 (>64)	-
	10% MHB	2 (2)	2 (2)	>64 (>64)	>64 (>64)	-
	TSB	8-16 (32)	16 (32)	>64 (>64)	>64 (>64)	-

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Supporting Information

Tables

Table S1. Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal Concentration (MBC) values for LL-37 and its derivatives against *E. coli* strains. MIC values (µM) of antimicrobial peptide LL-37 and its derivatives KR-12 and RI-10 against different *E. coli* strains in supplemented minimal medium (Min A), Mueller Hinton broth (MHB), and tryptic soy broth (TSB). The corresponding MBC values (µM) are shown within parentheses. Data represent values obtained from three independent biological replicates.

E. coli strains	Media	LL-37	KR-12	RI-10	Gentamicin
E.coli E423-17	Min A	2-4 (4)	8 (8)	>64 (>64)	2 (2)
	MHB	4-8 (8)	32 (32)	>64 (>64)	1 (1)
	TSB	4-8 (8)	32 (>32)	>64 (>64)	2 (4)
E.coli E416-17	Min A	4 (4)	4-8 (8)	>64 (>64)	2 (2)
	MHB	8 (8)	16 (16)	>64 (>64)	1 (2)
	TSB	8-16 (32)	16-32 (32)	>64 (>64)	4 (4)
E.coli 25922	Min A	2-4 (8)	1-2 (16)	>64 (>64)	2 (4)
	MHB	8 (8)	16 (32)	>64 (>64)	1 (2)
	TSB	8 (32)	32 (>32)	>64 (>64)	2-4 (4)

Table S2. List of peptides used in the study.

Peptide	Length	Amino acid sequence	Reference
LL-37	37	LLGDFFRKSKEKIGKEFKRIVQRIKDFLRNLVPRTES	(19)
KR-12	12	KRIVQRIKDFLR	(24)
RI-10	10	RIVQRIKDFL	(3)
KE-10	10	KEKIGKEFKR	(4)

Table S3. List of human LL-37 fragments that share the key recognition sequence for bacterial sensing (bolded).

Name	Peptide amino acid sequence
LL-37	LLGDFFRKSKEKIGKEF**KRIVQRIKDFLR**NLVPRTES
LL-23	LLGDFFRKSKEKIGKEFKRIVQR
LL-29	LLGDFFRKSKEKIGKEF**KRIVQRIKDFLR**
KR-20	**KRIVQRIKDFLR**NLVPRTES
KS-27	KSKEKIGKEF**KRIVQRIKDFLR**NLVPR
KS-30	KSKEKIGKEF**KRIVQRIKDFLR**NLVPRTES
RK-31	RKSKEKIGKEF**KRIVQRIKDFLR**NLVPRTES

Table S4. List of strains used in the study.

Strain	Relevant genotype	Reference/source
E. coli K12 MG1655	λ^- *rph-1*	*E. coli* Genetic Stock Center #7740
E. coli E423-17	Clinical antibiotic-resistant isolate	Wang lab, University of Nebraska Medical Center
E. coli E416-17	Clinical antibiotic-resistant isolate	Wang lab, University of Nebraska Medical Center
E. coli 25922	O6 isolate, reference strain for antibiotic susceptibility testing	ATCC
AML20	MG1655 $\Delta mgrB$ λ_{att} (P$_{mgrB}$-yfp) HK$_{att}$ (P$_{tetA}$-*cfp*)	(5)
JNC21	MG1655 $\Delta mgrB$ $\Delta queE$::kan λ_{att} (P$_{mgrB}$-*yfp*) HK$_{att}$ (P$_{tetA}$-*cfp*). kan^r	(6)
NRS6	MG1655 $\Delta mgrB::kan$ $\Delta phoQ$ λ_{att}::(P$_{mgrBt7a}$-*yfp*) HK$_{att}$::(P$_{tetA}$-*cfp*)	This work
SAM31	MG1655 $\Delta queE$ λ_{att} (P$_{mgrB}$-*yfp*) HK$_{att}$ (P$_{tetA}$-*cfp*)	(6)
TIM92	MG1655 λ_{att} (P$_{mgrB}$-yfp) HK$_{att}$ (P$_{tetA}$-*cfp*)	(7)
TIM100	MG1655 $\Delta phoQ$ λ_{att}::(P$_{mgrB}$-*yfp*) HK$_{att}$::(P$_{tetA}$-*cfp*)	(7)
TIM166	MG1655 $\Delta phoQ$ λ_{att}::(P$_{mgrBt7a}$-*yfp*) HK$_{att}$::(P$_{tetA}$-*cfp*)	(7)

Figures

Figure S1. Killing kinetics LL-37, KR-12, and RI-10 against *E. coli* K-12 MG1655. Cells were

treated with peptides at indicated concentrations in supplemented MinA with 0.1 mM MgSO$_4$.

Data are represented as mean ± range for two biological replicates.

Figure S2. PhoQP-regulated transcription and cell lengths in *mgrB-t7a* reporter cells treated with sub-MIC levels of antimicrobial peptides. (A, C) PhoQP-dependent promoter reporter activity in wild-type (TIM166) or Δ*mgrB* (NRS6) cells, and **(B, D)** quantification of cell lengths of wild-type or Δ*mgrB* cells, grown in the presence of sub-MIC levels of the antimicrobial peptides LL-37 (1.25 μM), KR-12 (5 μM), RI-10 (5 μM), and KE-10 (5 μM) in supplemented MinA with 0.1 mM MgSO₄. Data are obtained from three biological replicates; mean and median values are shown in red text and black bars, respectively. The number of cells analyzed is indicated for each sample. P-values indicate the results of a Student's t-test: ****P ≤ 0.0001, ***P ≤ 0.001, **P ≤ 0.01, *P ≤ 0.05, and "ns" = P > 0.05.

Figure S3. PhoQP-regulated transcription and cell lengths in Δ*mgrB* Δ*queE* cells treated with sub-MIC levels of antimicrobial peptides. (A) PhoQP-dependent promoter reporter activity, **(B)** quantification of cell lengths, and **(C)** phase contrast micrographs of Δ*mgrB* Δ*queE* (JNC21) cells grown in the presence of sub-MIC levels of the antimicrobial peptides LL-37 (1.25 μM), KR-12 (5 μM), RI-10 (5 μM), and KE-10 (5 μM) in supplemented MinA with 0.1 mM $MgSO_4$. Data are obtained from three biological replicates; mean and median values are shown in red text and black bars, respectively. The number of cells analyzed is indicated for each sample. P-values indicate the results of a Student's t-test: ****P ≤ 0.0001, ***P ≤ 0.001, **P ≤ 0.01, *P ≤ 0.05, and "ns" = P > 0.05.

References

1. J. Turner, Y. Cho, N. N. Dinh, A. J. Waring, R. I. Lehrer, Activities of LL-37, a cathelin-associated antimicrobial peptide of human neutrophils. *Antimicrob. Agents Chemother.* **42**, 2206–2214 (1998).

2. G. Wang, Structures of human host defense cathelicidin LL-37 and its smallest antimicrobial peptide KR-12 in lipid micelles. *J. Biol. Chem.* **283**, 32637–32643 (2008).

3. R. F. Epand, G. Wang, B. Berno, R. M. Epand, Lipid segregation explains selective toxicity of a series of fragments derived from the human cathelicidin LL-37. *Antimicrob. Agents Chemother.* **53**, 3705–3714 (2009).

4. A. F. Mechesso, Y. Su, J. Xie, G. Wang, Enhanced antimicrobial screening sensitivity enabled the identification of an ultrashort peptide KR-8 for engineering of LL-37mini to combat drug-resistant pathogens. *ACS Infect. Dis.* **9**, 2215–2225 (2023).

5. A. M. Lippa, M. Goulian, Feedback inhibition in the PhoQ/PhoP signaling system by a membrane peptide. *PLoS Genet.* **5**, e1000788 (2009).

6. S. S. Yadavalli, *et al.*, Antimicrobial peptides trigger a division block in Escherichia coli through stimulation of a signalling system. *Nat. Commun.* **7**, 12340 (2016).

7. T. Miyashiro, M. Goulian, Stimulus-dependent differential regulation in the Escherichia coli PhoQ PhoP system. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 16305–16310 (2007).