Carbon and microbes from thawing palsas and peats do not reach a wider fluvial system

Tuomela, Nea[1*]; Elovaara, Samu[2]; Hultman, Jenni[3]; Kaartokallio, Hermanni[2]; Thomas, David N.[1]

[1] Faculty of Biological and Environmental Sciences, University of Helsinki, Helsinki, Finland

[2] Finnish Environment Institute (SYKE), Helsinki, Finland

[3] Natural Resources Institute Finland (LUKE), Helsinki, Finland

*Corresponding author, nea.tuomela@helsinki.fi, Viikinkaari 1, 00790 Helsinki, Finland

Abstract

Climate change is altering the biogeochemical cycling of carbon and nutrients in the northern peatland and permafrost regions, which provide one of the largest terrestrial carbon storages. Lateral transfer of carbon needs to be more widely studied, especially in smaller streams and catchments, as they receive high loading of organic matter and are hotspots of carbon degradation. In this study we combined measurements of dissolved organic matter (DOM) quality and quantity with microbial community data from a small Arctic catchment. Our aim was to understand how the catchment is affected by two sub-catchments, degrading palsa permafrost mire and peatland thawing in spring. The small thaw ponds in the palsa mire were clearly distinct from the rest of the catchment and ponds in the peatland. Palsa ponds had higher DOM quantity, more aromatic DOM and distinct microbial communities. Thus, our results point towards analogous and low export rates of organic matter from the permafrost mire and peatland to the receiving small streams. Instead, the DOM is being retained and metabolised within the small ponds in both palsa mire and peatland. In our study we were also able to detect high abundances of ultra-small Patescibacteria. Patescibacteria dominated the microbial community composition in all the sampled waters.

Introduction

Northern peatlands are two of the largest reservoirs of organic carbon on Earth, storing a total of 415 Pg C, of which 185 Pg of carbon is in permafrost peatlands [1]. The Arctic is undergoing rapid warming [2,3], which is leading to the mobilization and release of soil organic carbon from permafrost [4]. Palsas, peat mounds with a permafrost core, are especially vulnerable to increasing temperatures and changing rainfall patterns [5] and drastic degradation of palsas has been detected pan-Arctic [6–8]. Following permafrost thaw, organic matter and nutrients can be released into streams, to be transported and transformed [1,9–11]. Fluxes of dissolved organic matter (DOM) are highly seasonal, with highest export rates during spring freshet [12]. However, with permafrost thaw changing the hydrology, DOM export patterns and quality can change [11,13].

The properties of DOM along with temperature and nutrient availability determine whether it is degraded or transferred further downstream [14]. In general DOM with lower molecular weight and less aromatic structure is more available for microbial degradation and consumed rapidly in lower-order streams [15]. Ancient permafrost DOM has been characterised as being highly available to microbial degradation, due to containing material comprised of DOM with lower molecular weight and lower aromaticity [10,16,17]. Other studies have found permafrost derived DOM to be resistant to degradation as it reaches freshwaters, due to rapid degradation within soil porewaters prior to reaching freshwaters [13,18]. Regardless of the quality of organic matter released from soils to freshwaters, the increasing runoff of organic matter affects the function of fluvial networks in the Arctic region [4,19]. The role of headwaters and thaw ponds in processing permafrost DOM is pertinent because they receive high loading of DOM from soils [14,20,21].

Microbial communities and their functionality are one of the key factors in DOM processing [22,23] and vice versa, DOM quantity and quality is an important driver for microbial community assembly [24,25]. Microbial assemblages in headwaters are also highly affected by the input of microbial taxa from soil [26–29]. Of the soil microbial taxa, some will increase in abundance downstream from the soils, acting as seed taxa for the downstream communities [30]. The community formation processes change from stochastic in the headwaters towards deterministic further downstream [28,30].

Headwater streams and ponds have been studied to an extent, but there is a knowledge gap on microbial community composition and function [14,31]. For example, less known phylum Candidatus Patescibacteria can represent most of the free-living microbial communities in surface and groundwaters [32–35]. Patescibacteria have been understudied due to their ultrasmall size, leading to underestimation of their abundance with standard water sampling protocols [33,36,37]. Patescibacteria might possess genes involved in the biogeochemical cycling of carbon and

nitrogen [36] and degrading peatland and permafrost derived organic matter to usable forms for methanogens [38].

In this study, we aimed to examine the impact of permafrost degradation on DOM export rate, DOM properties and microbial communities in an Arctic catchment. Our hypotheses were 1) DOM export rate increases due to degrading permafrost, 2) DOM quality differs between permafrost and peatland affected sub-catchments, 3) Microbial communities are affected by the differences in DOM quantity and quality along the catchment. To test these hypotheses, we conducted a sampling campaign in a small river catchment with both peatland and permafrost characteristics. We were able to capture the state of the catchment towards end of snowmelt period, when presumably the highest thawing of permafrost and peat occurred simultaneously while lateral transport into the fluvial system still occurred.

Materials and Methods

Sampling site and sampling strategy

Samples were collected in July 2023 from Kidisjoki River catchment, a sub-catchment of Utsjoki River close to Kevo Subarctic Research Institute (69º 48' N, 27º 06.5' E) in Finnish Lapland (Figure 1). Kidisjoki has a pristine catchment, with only direct human impact being caused by a reindeer herding facility (operational during October-November). The location has subarctic climate with cumulative rainfall in of 7.5 mm and average temperature of 11.9 °C in July 2023 [39]. Kidisjoki catchment area covers 22 km² of which approximately 1 km² is palsa permafrost mire [40]. The vegetation consists mainly of mountain birch, with treeless alpine heaths, peatland and Scots pine present in the lower parts of the basin. More detailed description of the Kidisjoki catchment vegetation can be found in [41].

Figure 1. Map of the sampling scheme in Kidisjoki River catchment, Kevo, Finland. Sampling sites are indicated with black dots. Water samples were collected from all sites and accompanied with soil samples from sampled pond sites, 13-17 and 21-26. Kidisjoki River was sampled from three locations (Kidisjoki 1-3). LTM stands for long term monitoring site. Green colour is indicative of peatland, and base map was acquired from National Land Survey of Finland [42].

Water samples were collected as grab samples from Utsjoki River, along Kidisjoki River and from ponds and outlets in peatland and palsa mire (Figure 1). Water was collected into clean acid-washed plastic bottles and transported in coolers to the field station for further processing. When present, large particles were removed by pre-filtering the water with plankton net (sieve size 40 μm). Temperature and conductivity of were measured with YSI 6600 V2 CTD sonde (YSI Corporation, Yellow Springs, Ohio, USA) and stream flow rates were measured with AVD FlowTracker (SonTek, San Diego, CA, USA).

All water samples for microbial communities were processed within the sampling day. Autoclaved filtration units (Millipore, 250ml) were used to sequentially filter the sample with 0.45 μm and 0.1 μm pore size polyether sulfone membrane filters (Sartorius, Göttingen, Germany). Samples were

filtered until maximum of 30 minutes had passed to keep cells intact. Filtered volumes are listed in Supplementary Table 1. Filters were folded with sterile tweezer and stored in cryovials at -80 °C until DNA extraction.

Soil samples were collected next to the ponds from peatland and palsa sites (Figure 1). Six replicates were collected from palsa site and five from peatland. Soil was collected from three depts. From peatland sampled depths were 2, 4 and 10 cm below soil surface. Soil corer was used to sample palsa soil, from active, transitional and permafrost layers. Samples were sliced with ethanol-disinfected knife, put into sterile plastic bags and stored in cooler with -80 °C icepacks until transfer to -80 °C freezer within hours.

DOM degradation experiment

Microbial degradation of DOM was investigated by incubating water samples in the dark at 10 °C and collecting subsamples for analyses at the start and after approximately 15, 100, 150 and 220 days. The first sampling interval aimed to capture the short-term processes that could be related to the microbial community observed in situ. The subsequent samplings assessed the long-term resistance of DOM to general microbial degradation, as the microbial community was expected to change radically, and we thus did not attempt to connect this change to the observed components of the microbial community. 500 mL of sample was filtered through 0.8 µm to remove flagellate grazers, stored in a glass bottle and stirred periodically. In addition to filtered samples, unfiltered samples were collected at sites Palsa outlet 2, Peat outlet 2 and Main 3 to investigate how the presence of full planktonic community would affect degradation of DOM. Samples for the measurement of dissolved O_2 were collected separately into glass stoppered vials with no head space and measured using Winkler titration to calculate microbial respiration.

Respiration was calculated only at the first subsampling interval. Optical properties of DOM were measured at each subsampling, bacterial abundance was measured at the first, second and the last subsampling and nutrient concentrations were measured at the first and the last subsampling.

Microbial analysis and cell enumeration

DNA extractions for soil and water samples were done with DNeasy PowerSoil Pro Kit (Qiagen, Hilden, Germany). For soil samples, 0.25-0.30 g of soil was used for DNA extraction avoiding the exposed surface of the sample. For water samples, DNA was extracted from the filters and separately for the two size fractions. Filters were transferred to PowerBead tubes, Solution CD1 was added, and filter was broken with sterile pipette tip. Cell disruption step was done with FastPrep homogenizer (MP Biomedicals, Heidelberg, Germany; 30s, 5.5m/s). The rest of the extraction was completed following manufacturer's instructions. DNA concentration was measured with Qubit Fluorometer, using Qubit High sensitivity dsDNA kit (Invitrogen by Thermo Fisher

Scientific, Life Technologies Corporation, USA). DNA from the two size fractions was combined prior to sequencing. Libraries were prepared with Nextera™ DNA Flex Library Preparation Kit (Illumina, San Diego, CA, USA) and paired-end sequencing was done with AVITI (Element Biosciences, 2x150bp) at the Institute of Biotechnology, University of Helsinki. Bacterial cell abundance was determined using flow cytometry with LSRFortessa (BD Biosciences) at University of Helsinki Flow Cytometry Unit (Supplementary Methods 1).

Bioinformatics

Raw sequences were trimmed using cutadapt v4.9 [43] to remove Nextera adapters. Quality was checked with FastQC v0.11.9 [44] and MultiQC v1.19 [45]. After trimming, the sequences from three individual lanes were combined for each sample. For taxonomic annotation, trimmed and combined reads were mapped against SILVA 138.1 database with PhyloFlash v3.4.2 [46]. Mitochondrial and chloroplast reads were removed prior to analysis with phyloseq package v.1.42.0 [47] in R v4.2.3 [48] and RStudio v2024.12.0 [49]. Bacterial and archaeal communities are presented as mean relative abundance for each sample type. Community compositions for individual samples can be found in Supplementary Figures 1-3.

DOC, DOM and nutrient analysis

Samples for dissolved organic carbon (DOC) and coloured dissolved organic matter (CDOM) were prepared by filtering 20mL of water through pre-combusted GF/F filters (450 ∘C, 4 h) into acid washed and pre-combusted glass vials. DOC samples were acidified to pH 2 with concentrated orthophosphoric acid, sealed with septum cap and stored at −20 ∘C. DOC concentrations were analysed with TOC-L total organic carbon analyser (Shimadzu). Long term data on DOC concentration and export rates in Utsjoki River were obtained from Finnish Environment Institute [50].

CDOM samples were stored in +4∘C until analysis after field campaign, within 10 days of sampling. Optical properties of DOM were conducted following [51].

CDOM absorption was measured with a Shimadzu 2401PC spectrophotometer using 1cm quartz cuvette over spectral length range 200-800 nm with 1 nm resolution, using ultrapure water (MQ) as blank. Ultraviolet specific absorbance at 254nm ($SUVA_{254}$) was calculated by normalizing DOC absorbance at $254nm^{-1}$ to DOC concentration in mg L^{-1} [52]. CDOM slopes ($S_{275-295}$, $S_{350-400}$, $S_{300-650}$) were calculated with R package cdom v0.1.0 [53].

Excitation–emission matrices (EEMs) of FDOM were measured with a Varian Cary Eclipse fluorometer (Agilent). The characteristics and the quality of the DOM were assessed with fluorescence peaks [54] extracted from the EEMs utilizing the R package eemR [55]. A blank

sample was subtracted from the EEMs, and the Rayleigh and Raman scattering bands were removed from the spectra after calibration. EEMs were calibrated by normalizing to the area under the Raman water scatter peak 11 (excitation wavelength of 350 nm) of an MQ water sample run on the same session as the samples and were corrected for inner filter effects with absorbance spectra [56].

Nutrient concentrations (nitrate, nitrite, phosphorus, silicate and total dissolved nitrogen) were measured with SEAL AA500 AutoAnalyzer (Seal Analytical, US) using standard methods. Ammonium concentrations were determined fluorometrically [57] using Varian Cary Eclipse fluorometer (Agilent).

Soil properties

Soil physicochemical properties were measured from lyophilized and ground soils. Soil pH was measured following international standard ISO10390. Loss on ignition (LOI) was used to analyse soil organic matter (SOM) content according to Finnish standard SFS3008. Soil total carbon and total nitrogen analysis was performed with a high temperature induction furnace (LECO, 828 series).

Statistical analysis

All data analysis were conducted in R v4.2.3 [48] and RStudio v2024.12.0 [49] with package tidyverse v2.0.0 [58]. Variation of biogeochemical properties was analysed with Principal Components Analysis (PCA, prcomp-command) utilizing R package ggbilot v0.55 [59]. Mean values of technical replicates were used and variables which were not determined for all samples were excluded. Screeplot was used to justify the selection of principal components (Supplementary Figure 5). For microbial community analysis phyloseq package v1.42.0 was used [47] and variation between samples was visualized with Principal Coordinate Analysis (PCoA) on genus level, utilizing Bray-Curtis distance. The microbial community taxonomic diversity was measured with Shannon Index (Supplementary Figure 6).

Results

Biogeochemical properties

DOC concentrations were highest in palsa ponds, ranging from 2 992 to 9 330 µmol L^{-1}, followed by peat ponds with values from 210 to 5 750 µmol L^{-1} (Table 1). Peat and palsa outlets had similar DOC concentrations of 407 and 401 µmol L^{-1} respectively. Kidisjoki River and Utsjoki River had similar range of DOC with each other. In peat ponds, a254 was relatively low compared to the DOC concentration, indicating possibly higher ratio of non-coloured DOM of the total DOM pool

than in e.g. palsa ponds. The humification index was highest in palsa ponds, followed by peat outlet and Kidisjoki River. Strikingly, the humification index decreased from palsa ponds to the palsa outlet, while the opposite was measured between the peat ponds and peat outlet. The lowest $SUVA_{254}$ values were measured in peat ponds and the highest in palsa ponds. Palsa outlets had slightly higher $SUVA_{254}$ than rest of the catchment. $SUVA_{254}$ is a proxy for aromaticity [52], and the values would indicate higher aromaticity in palsa ponds than rest of the samples and the lowest aromaticity in peat ponds. Aromaticity increased from the peat ponds to the peat outlet and decreased from palsa ponds to the palsa outlet site.

Table 1. Physicochemical properties of freshwater sampling sites in the studied Kidisjoki River catchment in Utsjoki region, Finland. Number of sampling sites are expressed in column header.

Parameter	Unit	Palsa pond (6)	Palsa outlet (2)	Peat pond (5)	Peat outlet (2)	Kidisjoki River (3)	Utsjoki River (1)
DOC	$\mu mol\ L^{-1}$	5 561 (2217)	407 (71.00)	3706 (2882)	401 (56)	223 (5.9)	237
Organic N	$\mu mol\ L^{-1}$	110.47	7.81 (2.04)	4.30 (0.37)	6.66 (0.65)	3.80 (0.29)	6.84
DOC:DON		46.13	53.37 (4.84)	870.64 (709.65)	59.91 (2.64)	58.92 (4.93)	34.69
$SUVA_{254}$	$L\ mg^{-1}\ m^{-1}$	16.04 (3.99)	8.41 (0.97)	2.11 (2.69)	7.98 (0.29)	7.96 (0.06)	7.19
NO_2^-	$\mu mol\ L^{-1}$	1.02 (0.34)	0.08 (0.01)	0.04 (0.02)	0.07 (0.01)	0.08 (0.01)	0.04
NO_3^-	$\mu mol\ L^{-1}$	0.02 (0.04)	0.02 (0.02)	0.09 (0.03)	0.08 (0.06)	0.34 (0.34)	0.13
PO_4^{3-}	$\mu mol\ L^{-1}$	5.66 (4.37)	0.03 (0.01)	0.03 (0.01)	0.04 (0.00)	0.04 (0.00)	0.03
Si	$\mu mol\ L^{-1}$	1.44	26.88 (16.28)	60.21 (14.45)	18.45 (5.24)	91.85 (36.50)	82.39
TDN	$\mu mol\ L^{-1}$	119.13 (38.02)	7.95 (2.05)	4.51 (0.34)	6.88 (0.74)	4.34 (0.60)	7.11
NH_4^+	$\mu mol\ L^{-1}$	2.13	0.04 (0.00)	0.08 (0.00)	0.06 (0.03)	0.12 (0.03)	0.11
pH		4.84 (0.99)	6.25 (0.91)	6.91 (0.16)	7.19 (0.30)	7.22 (0.18)	5.57
Conductivity	$\mu S\ cm^{-1}$	95.15 (12.33)	79.23 (1.77)	79.83 (4.48)	72.00 (2.00)	77.00 (0.82)	82.08
Temperature	°C	14.89 (1.89)	11.00 (2.78)	13.25 (1.42)	13.29 (1.43)	6.80 (0.59)	15.55
ODO	%	67.45 (15.52)	75.25 (3.93)	73.64 (10.80)	89.50 (0.26)	91.80 (0.25)	99.46
ODO	$mg\ L^{-1}$	6.81 (1.51)	8.35 (0.97)	7.72 (1.16)	9.38 (0.27)	11.20 (0.15)	9.91
fi		1.03 (0.10)	1.10 (0.02)	1.18 (0.08)	1.16 (0.10)	1.17 (0.03)	1.04
hix		33.48 (6.80)	23.64 (3.82)	18.96 (7.70)	37.97 (4.92)	35.14 (5.69)	17.43
bix		0.34 (0.03)	0.51 (0.07)	0.49 (0.07)	0.41 (0.04)	0.39 (0.02)	0.42
S_R		0.95 (0.07)	0.75 (0.03)	0.74 (0.02)	0.72 (0.01)	0.77 (0.01)	0.86
$S_{275-295}$	$10^{-3}\ nm^{-1}$	8.88 (0.50)	11.93 (0.13)	12.92 (0.31)	12.58 (0.29)	13.00 (0.23)	14.45
$S_{350-400}$	$10^{-3}\ nm^{-1}$	9.38 (1.04)	16.03 (0.41)	17.42 (0.14)	17.50 (0.14)	16.83 (0.41)	16.73

$S_{300-650}$	10^{-3} nm^{-1}	18.23 (1.02)	14.67 (0.14)	15.36 (0.32)	15.32 (0.25)	15.48 (0.09)	15.71
Peak T	R.U.	0.03 (0.00)	0.08 (0.03)	0.05 (0.03)	0.04 (0.01)	0.02 (0.00)	0.04
Peak A	R.U.	0.45 (0.12)	0.50 (0.01)	0.34 (0.02)	0.62 (0.06)	0.34 (0.01)	0.28
Peak M	R.U.	0.27 (0.08)	0.31 (0.01)	0.23 (0.01)	0.39 (0.04)	0.20 (0.00)	0.19
Peak C	R.U.	0.31 (0.08)	0.35 (0.02)	0.24 (0.01)	0.43 (0.04)	0.24 (0.01)	0.19
a_{254}	m^{-1}	981 (252.13)	40.28 (2.41)	19.83 (1.37)	38.20 (3.99)	21.28 (0.61)	20.47
HNA bacteria	10^6 cells ml^{-1}	5.74 (2.43)	2.58 (0.57)	2.39 (0.52)	2.39 (0.76)	1.22 (0.30)	5.15
LNA bacteria	10^6 cells ml^{-1}	4.73 (3.84)	2.24 (0.78)	1.75 (0.31)	1.66 (0.59)	0.95 (0.41)	1.59

Total dissolved nitrogen (TDN) and phosphate (PO_4^{3-}) had highest concentrations in the palsa ponds. Concentration of TDN was 10-fold lower in other sampling sites. Palsa ponds had highest conductivity, followed by Utsjoki River. Peat ponds, palsa outlet and Kidisjoki River had similar conductivity, whereas peat outlet had slightly lower value. Oxygen concentration and saturation lowest in palsa ponds, with slight increase to palsa outlets. Similar pattern was also observed in peat ponds and peat outlet. Bacterial numbers were highest in palsa ponds for both high and low nucleic acid bacteria, whereas palsa outlet and peatland had similar range of bacterial cells. Oxygen measurements and bacterial numbers indicated higher bacterial activity in palsa ponds than rest of the catchment. Palsa ponds had the highest temperature of all sites, they were dark coloured and warmed in the sunlight during the sampling day.

Principal component analysis of the nutrients, and DOM quality and quantity revealed distinction between the palsa pond sites and rest of the water samples (Figure 2). PC1 explained 44.7% and PC2 15.3% of the variation. PC1 variation was contributed by oxygen, $S_{275-295}$, pH, SUVA$_{254}$, conductivity, DOC, phosphate and nitrogen. The largest contributions to PC2 were Peak A, Peak C and Peak M. Palsa ponds were apparently clustering due to DOC concentration, $S_{350-600}$, phosphate, conductivity and SUVA$_{254}$. Whereas peat ponds, Kidisjoki River and Utsjoki River were clustering by spectral slope $S_{275-295}$, oxygen, nitrate concentration and pH values. This would point towards lower molecular weight and more labile organic matter. Palsa and peat outlets are forming slightly into their own group, driven by Peak T values, which is for fluorescence from protein-like compounds [54].

Figure 2. Principal component analysis of biogeochemical properties of the studied arctic catchment water samples from small ponds and streams. Measured variables include DOC concentration (DOC), nutrient concentrations and physicochemical properties. PCA includes also optical property measurements of DOM (Peaks A, C, M and T, Fluorescence index (FI), Biological index (BIX), Humification index (HIX), Spectral slopes ($S_{275-295}$, $S_{350-400}$ and $S_{300-650}$) and SUVA (SUVA$_{254}$). Sample type is indicated by shape and site by colour. Labels refer to the PCA loading arrows.

Soil properties

Palsa soils were acidic with pH between 3.75 and 4.10. Palsa soils had C:N (total C : total N) ratios of between 35.2 and 38.7 and high SOM contents of 97% in all depths (Table 2). In peatlands the pH values were between 5.16 and 5.31, C:N ratio ranged from 17.81 to 25.85, and SOM was between 19 and 67%. SOM content in peatland was highest in the soil surface.

Table 2. Soil properties in palsa and peatland soils, grouped by depth. Values are means of 6 replicates for each depth in palsa site and 5 for peatland samples. C:N ratio is the ratio of total carbon to total nitrogen. Standard deviation is given in parentheses. Palsa samples are divided to active, transitional and permafrost layers. Peat sample depths are expressed as the depth below soil surface.

Site	Depth	pH	C:N ratio	Soil organic matter %
Palsa	Active layer	3.75 (0.08)	37.35 (7.71)	96.98 (2.95)
n = 6	Transitional	3.78 (0.10)	35.20 (4.09)	97.25 (1.27)
	Permafrost	4.10 (0.39)	38.67 (9.50)	97.93 (0.70)
Peat	2 cm	5.31 (0.22)	25.85 (7.68)	67.14 (28.37)
n = 5	4 cm	5.19 (0.26)	23.51 (4.50)	59.68 (35.04)
	10 cm	5.16 (0.15)	17.81 (1.40)	18.88 (19.88)

Microbial communities

Bacterial community composition

Based on the 16S rRNA gene annotations from metagenomes, the bacterial communities in the water samples were dominated by phyla Patescibacteria, Proteobacteria, Bacteroidetes, Acidobacteriota and Verrucomicrobiota (Figure 3a). Acidobacteriota, Actinobacteriota and Proteobacteria were abundant in palsa soil communities. In peat soils more relative abundance was more evenly distributed among phylum, including e.g. Chloroflexi, and Verrucomicrobiota. Patescibacteria was highly abundant phylum in palsa outlet, peat ponds, peat outlet and Kidisjoki River. In Utsjoki River, the community is dominated by Actinobacteriota, Bacteroidetes and Gammaproteobacteria. In order level the Utsjoki River bacterial community consisted of Burkholderiales, Frankiales, Flavobacteriales, Cytophagales and Chitinophagales (Figure 3b). Palsa pond community consisted of Burkholderiales, Acidobacteriales and Acetobacteriales. Palsa outlets, peat outlets, peat ponds and the Kidisjoki River were dominated by the orders Omnitrophales, Burkholderiales and from the Patescibacteria phylum, orders Candidatus Nomurabacteria and Candidatus Kaiserbacteria.

Figure 3. Bacterial community composition of the study site in (a) phylum level (relative abundance ≥ 0.01% shown) and (b) order level (relative abundance ≥ 0.38% shown). Number of replicates from each site is indicated below the site label in x-axis and reported relative abundances are mean values of the replicates.

Archaeal community composition

Archaeal communities consisted of phyla Nanoarchaeota, Micrarchaeota, Halobacterota and Crenarchaeota (Figure 4). Peat soils had highest relative abundance of archaeal 16S rRNA genes with approximately 4% of total relative abundance. The archaeal communities in palsa soils consisted of orders Group 1.1c and Methanomicrobiales, while in palsa ponds of Methanobacteriales and Micrarchaeales were most abundant. Peat soil archaeal community was composed of Methanomicrobiales, Methanobacteriales and Methanosarciniales. Woesearchaeales were present in all sites, and were abundant in palsa outlet, peat soil, peat ponds, peat outlets and Kidisjoki River. Micrarchaeles were also abundant in all water samples except Utsjoki River.

Figure 4. Archaeal community composition in Order level of the Kidisjoki River catchment sampling sites. Number of replicate samples indicated below the site label in x-axis and relative abundances are reported as mean values of the replicates.

Microbial community variability

PCoA shows clear division of the microbial communities among sampled sites, with x-axis explaining 35.9 % of variation and y-axis 20.7% (Figure 5a). Palsa ponds, palsa soils and peat soils formed three distinct groups. Permafrost soil from site 26 and Utsjoki River are grouping together with palsa ponds. Kidisjoki River, outlet sites and peat ponds cluster together, forming a fourth distinct group.

Palsa ponds cluster separately from other sites also in PCoA with only water samples (Figure 5b). Utsjoki River sample seems to group closer to palsa ponds than to the rest of the catchment, likely due to Patescibacteria and Proteobacteria relative abundance being similar range in both sites. Peat and palsa outlets were grouped together with the rest of the water samples, unlike in the PCA of biogeochemical properties, where they formed a slightly distinct group (Figure 2).

Figure 5. Principal Coordinate Analysis to detect variation between microbial communities of relative abundance in genus level in (a) all sampling sites and (b) sampled freshwater sites. Palsa and peat soils formed separate groups from water samples. Notably palsa ponds formed a distinct group from rest of the water sites. Sample type is indicated by shape and sample site by colour.

Degradation experiment

The local microbial community in the palsa pond was more active compared to any other site, as seen by higher bacterial respiration rate (Supplementary Figure 7a) and growth (Supplementary Figure 7b, Supplementary Table 2). Microbial transformation of DOM resulted in an initial increase in CDOM absorption, especially in palsa ponds (Supplementary Figure 8, Supplementary Table 2). At other sites, a254 generally decreased while absorption coefficients at higher wavelengths increased. Spectral slopes changed negligibly (Supplementary Figure 9, Supplementary Table 2). Fluorescence peaks generally increased throughout the experiment, except at palsa ponds, where they abruptly dropped between the 2nd and 3rd subsamplings (Supplementary Figure 10, Supplementary Table 2). At the end of the degradation experiment most CDOM and FDOM parameters had either decreased from initial values or increased very little, except for HIX which increased at each site. With a few exceptions (ammonium increased in Utsjoki River and decreased in the palsa pond) nutrient concentrations did not change considerably during the experiment (Supplementary Figure 11, Supplementary Table 2). The exception is silicate, which increased in every site except the palsa pond.

The changes in the CDOM absorption coefficients during the degradation experiment in the palsa pond were modest compared to other sites, when accounting for the high differences in starting conditions. This suggests that the high bacterial respiration and growth could more strongly be sustained by the non-coloured DOM in the ponds. This could in turn lead to the export of predominantly refractory coloured DOM.

DOC consumption in the degradation experiment was estimated using three methods for predicting DOC concentration from CDOM absorption coefficients [60–62]. Predicted DOC concentrations are highly variable, and especially the results for the palsa pond should be assessed critically, because the conversion methods were not created using such high absorption coefficient values. Yet the predicted DOC degradation of other sites is much lower than in the palsa pond, generally less than 20 µmol C L^{-1} consumed during the total duration of the experiment (Supplementary Table 2).

Non-filtered samples showed that grazer limit bacterial growth at the selected control sites (Supplementary Figure 7c). In these samples CDOM absorption coefficients decrease already during the first measurement interval (Supplementary Table 3), showing the role of bacteria in producing CDOM.

DOC export rates

Estimated export rates of DOC were similar in both palsa and peat outlets (Table 3). Higher rates of export were estimated in Kidisjoki River sites, where DOC export rate increased downstream. DOC export rates of Utsjoki River long term monitoring site in May show variation between 1990-

2024, but no clear trend of increased DOC export with time is discernible (Supplementary Figure 12).

Table 3. Concentration and export rates of dissolved organic carbon (DOC) of the catchment sampling locations estimated based on DOC and flow rate measurements in early June. Utsjoki River export rate is calculated from DOC concentration measured in this study and day specific flow rate from Finnish Environment Institute [50].

Site name	DOC g m^{-3}	Flow rate m^3 s^{-1}	DOC export kg C day^{-1}
Palsa outlet 2	4.04	0.02	5.48
Peat outlet 2	4.14	0.02	5.47
Kidisjoki River 1	2.74	0.09	20.47
Kidisjoki River 2	2.71	0.28	64.42
Kidisjoki River 3	2.58	0.30	67.03
Utsjoki River	2.85	17.74	4 366

Discussion

The main aim of the study was to identify how DOM quantity and quality was affected by peatland and degrading permafrost in a small subarctic catchment and how DOM properties affect the microbial community composition. We were able to capture a unique time towards the end of the spring freshet, with the catchment being hydrologically connected and thaw depth of the permafrost soil highest. During our sampling campaign palsa thaw ponds were evaporating rapidly, and likely had disappeared within days following our sampling.

Dissolved organic matter is processed within site and not transferred

Contradictory to general view of increased DOC flux from permafrost and peat soils into streams and rivers [4,9,19,63], our results do not show clear impact of enhanced DOC export to the stream network. Instead of transfer to streams, our results point towards majority of organic matter being retained and processed before it could reach the first headwater streams. This conclusion is supported by high bacterial cell numbers, higher bacterial growth and respiration and distinct bacterial and archaeal communities in palsa ponds compared to the rest of the catchment. While our sampling consisted of one sampling, long-term monitoring data of spring freshet from Utsjoki River does not show a drastic increase in DOC export rates either (Supplementary Figure 12).

Physicochemical properties of the palsa pond were also distinct from the other sites, supporting the immediate processing and relatively low export rate of DOM from palsa mire to the catchment. Carbon export rate from peatland was analogous to the palsa export rate. However, the peat ponds had bacterial community reflective to the rest of the catchment, unlike the permafrost affected site.

DOC export rates have been detected to remain stable or increase following palsa degradation in Stordalen palsa mire [13]. They detected that as palsa permafrost degrades, palsa mound will be transforming into bog or fen, determining the DOC quality and export rate. In bogs DOC export did not increase, had higher aromaticity and higher molecular size, reflecting a terrestrial origin. Conversely in a more hydrologically connected fen, increased DOC export is detected, and DOC characterized predominantly of microbial origin and labile [13]. In Siberian Arctic, peat bog depressions of similar size as our sampled ponds had comparable DOC concentrations [18]. The study site had less dramatic decrease in DOC concentration from the depressions to small streams compared to our results. Higher DOC concentrations in thaw ponds than non-permafrost affected ponds have also been detected in Canadian Arctic [64], supporting our findings. However, our thaw ponds had overall slightly higher DOC values. This could be due to sampling time, as our sampling was done towards the end of the spring snowmelt and theirs in August during the base flow period.

Our results are limited to a one-time sampling of a small catchment, and thus might not be applicable for large-scale generalisation of DOC export rates and processing following permafrost thaw in the Arctic. Yet similar trends in lower DOC export and higher processing have been detected in other studies [13,65,66]. Streams and ponds have been measured to be sources of CO_2 and CH_4 [21,65–68]. While greenhouse gas measurements were not in the scope of our study, in a similar palsa site with comparable DOC concentrations dissolved CH_4 concentrations were 2 000-5 000 times of the atmospheric equation in palsa ponds [65]. Based on their values, similar range could be expected in our study site. Relative to their areal coverage, small water bodies are an unproportionally large carbon emitter and can affect carbon balance even at landscape scales [67]. The range of carbon lateral transfer and greenhouse gas emissions remains uncertain, due to limited wider sampling campaigns on small streams [67].

Dissolved organic matter properties are distinct between palsa ponds and rest of the catchment

The properties of DOM divided the catchment to two distinct groups: palsa ponds and the rest of the catchment. Palsa ponds CDOM was characterized as more refractory, with higher aromaticity ($SUVA_{254}$) and larger molecular size ($S_{275-295}$). The palsa ponds were also characterized by high total nitrogen and lower pH relative to the other sites. Lower pH, high DOC concentrations and

refractory carbon quality are typical for emerging thaw ponds, observed in Canadian discontinuous permafrost zone [38,63] and palsa mires [13,65]. Different soil properties in peat and palsa sites could lead to variability in DOC properties. Palsa soils had high SOM, whilst peat soils were shallow with lower SOM content. The quality of organic matter varies, with palsa soils containing more degraded and recalcitrant organic matter in aerobic active layer and more labile organic matter in permafrost [69]. In peatland the anoxic state of the soil prevents rapid degradation [69].

Our results on CDOM properties agree with earlier work, where DOC from palsa sites was characterized as poorer substrate for microbial consumption [13]. Higher SUVA values, higher C:N ratio and lower FI were detected in palsa mire compared to fens in Swedish Arctic [13]. The changes in hydrology from palsa degrading into fen resulted into higher groundwater connectivity, which could result into more labile DOC. Similar dynamics were also detected in our study sites. Similar trend in permafrost degradation leading into higher terrestrial and refractory carbon in freshwaters is detected also in the circumpolar permafrost zone [11].

Refractory DOC properties could be explained by the sampling time being towards the end of spring freshet and start of baseflow period. During spring freshet more organic matter is transferred, and microbial degradation is limited by shorter residence times and lower temperatures [70]. In subarctic peatland stream, lower quantity and quality of organic matter for microbial conditions was detected during baseflow conditions [66]. During baseflow organic matter is likely degraded closer to source soil [66].

Elevated concentrations of DOC and more labile characteristics downstream from the thaw slumps were detected in Canadian peat plateau [63]. In our study site, DOC concentration downstream from both permafrost and palsa sites decreased drastically. Notably our thaw pond sites were evaporating rapidly, likely leading to elevated DOC concentrations and reduced hydrological connectivity. These elevated DOC concentrations were still not detected in the outlet sites, which were minimum of meters from the ponds, strongly pointing towards DOC residing in the ponds and not transferred further.

Microbial communities comprised mainly of ultra-small bacterial and archaeal taxa

Microbial communities were distinct between palsa ponds and rest of the catchment, reflecting the biogeochemistry of the sites. This supports our hypothesis that DOC quantity and quality, along with the environmental properties like pH, are affecting the microbial community composition. Microbial community assembly in palsa ponds was likely affected the thawing palsa soils, which caused low pH and high input of DOC. In addition to the distinct community composition, we detected high relative abundance of ultra-small taxa in both bacterial and archaeal communities.

Stream and peat ponds bacterial communities were dominated by reads assigned to Patescibacteria 16S rRNA genes, consisting of over half of the relative abundance in the stream sites and peat ponds. While this bacterial phylum is relatively unknown, it has been detected in high abundances in e.g. groundwater [71,72], wastewaters [73] and thermokarst ponds [33,38]. Of Patescibacteria, the orders Candidatus Kaiserbacteria and Candidatus Nomurabacteria were the most abundant in the stream sites, which have been also detected in Arctic thermokarst lakes [33] and groundwater [74]. Patescibacteria relative abundance was lower in Utsjoki River than in the smaller streams and ponds. While our data had only one sampling point from Utsjoki River, similar trend of decrease in Patescibacteria with increasing river size has been detected in a large-scale riverine study [31].

Patescibacteria lifestyle is considered host dependent, either symbiotic or parasitic, due to the reduced genome size and lack of metabolic pathways [72,75]. Patescibacteria have been detected to ferment carbon compounds to acetate, and benefitting e.g. methanogens, methanotrophs, Actinobacteria, Betaproteobacteria, Planctomycetes, and Chloroflexi [33]. Patescibacteria are facultative anaerobes but possess genetic ability to resist oxygen [33].

Ultra-small archaea Nanoarchaeota and Micrarchaeota were also detected along the catchment. They have similar lifestyle as ultra-small bacteria, with host attached lifestyle and ability to degrade complex organic matter by fermentation [36,37]. Archaeal communities were distinct between palsa ponds and rest of the catchment, with Nanoarchaeota and order Woesearchaeales dominating in peat ponds and stream sites, and Methanobacteriales in palsa ponds.

Besides Patescibacteria, palsa pond bacterial community consisted of Acidobacteriota, Proteobacteria, Bacteroidota and Verrucomicrobiota. Similar bacterial communities have been detected in thaw ponds of Canadian Arctic [38]. Palsa soil was dominated by Acidobacteriota, Actinobacteriota and Proteobacteria, similarly as earlier findings in Stordalen palsa mire in Sweden [76]. Stordalen fen soils were comprised of a higher number of bacterial phyla, and higher archaeal relative abundance, than the palsa soils [76], supporting our findings.

Soils are the main source of bacteria into the freshwaters, where the bacterial communities are shaped by stochastic and deterministic processes [26,27,30]. Our sampled freshwaters were highly connected to soils, yet the community composition between soil and adjacent pond was distinct. Since our sampling was conducted towards the end of spring freshet, microbial communities in the ponds had likely been developed over few weeks, creating distinct community structure from the soils. Stochastic processes might dominate community assembly in the stream sites and ponds immediately after spring freshet [27], whereas towards the end of freshet, selection might dominate microbial community assembly, especially in palsa ponds, where DOC and dissolved nutrient concentrations were high.

It is likely that peatland and Kidisjoki River were also dominated by bacterial taxa derived from snowmelt or groundwater, rather than soil input. During high flow events, selection hotspots are focused on areas with longer residence time [27]. Our sampling was towards the end of high flow snowmelt, and apart from the less connected palsa ponds, the residency time could have been too short for selection to occur, and rather the mass effect of microbial input from soils and groundwater to dominate the assembly processes. While this effect might not be detected here, assembly-based methods might give more insight into the formation and occurrence of similar taxa between soil and water samples. Our study is limited to the free-living and particle associated microbes. Benthic microbial communities might display the community assembly processes along the stream network [77].

Conclusions

The impact of thawing palsas and peatland on the export and properties of DOM and microbial community composition was studied on a small arctic river catchment. As hypothesised, terrestrial DOM loading was high from both peat and permafrost soils to ponds, indicated by high DOC concentration and optical properties. However, our results indicate that DOM was not transferred further to the catchment, but rather retained in the small ponds. Similar range of DOC export detected from both permafrost and peatland sites. Microbial community composition reflected the DOM quality and quantity, alongside with other environmental values, with their taxonomic composition in palsa ponds being distinct from the rest of the sites. We were able to detect high relative abundance of ultra-small Patescibacteria along the whole catchment. Our results highlight the need for more research on the biogeochemistry and role of microbial communities in the carbon cycling, especially in the diverse Arctic and sub-Arctic small-scale freshwaters.

Data availability

DNA sequences have been deposited in ENA under project number PRJEB85624. Biogeochemical data is available in Pangaea (https://doi.pangaea.de/10.1594/PANGAEA.981156).

CRediT authorship contribution statement

NT: Conceptualization; funding acquisition; data curation; formal analysis; investigation; visualization; writing—original draft; writing—review and editing. **SE:** Conceptualization; data curation; formal analysis; investigation; visualization; writing—review and editing. **HK:** Conceptualization; Supervision; Investigation; Resources; Writing – review & editing. **JH:** Conceptualization; Supervision; Investigation; Resources; Writing – review & editing. **DT:**

Conceptualization; Project administration; Investigation; Resources; Supervision; Writing – review & editing

Acknowledgements

We want to thank the funders of this study, Kone Foundation (202204330) and Emil Aaltonen Foundation (220226). JH acknowledges the Research Council of Finland (354462). We thank Kevo Subarctic Research Institute of University of Turku and staff for their help. Metsähallitus is acknowledged for sampling permit. We thank Janina Kariluoto for her assistance during the field campaign and with the laboratory work. We acknowledge DNA Sequencing and Genomics Laboratory (supported by HiLIFE and Biocenter Finland funding), Institute of Biotechnology, University of Helsinki for sequencing. We acknowledge CSC – IT Center for Science, Finland for the provision of computational resources. The study utilized marine research infrastructure at the Finnish Environment Institute as a part of the national FINMARI RI consortium. The flow cytometry analysis was performed at the HiLife Flow Cytometry Unit, University of Helsinki.

Declaration of competing interest

The authors declare no conflict of interest.

References

1. Hugelius G, Loisel J, Chadburn S et al. Large stocks of peatland carbon and nitrogen are vulnerable to permafrost thaw. *Proc. Natl. Acad. Sci. U.S.A*. 2020;117:20438–46. https://doi.org/10.1073/pnas.1916387117.

2. Rantanen M, Karpechko AY, Lipponen A et al. The Arctic has warmed nearly four times faster than the globe since 1979. Commun. *Earth Environ*. 2022;3:168, https://doi.org/10.1038/s43247-022-00498-3.

3. IPCC. In: Team C.W., Lee H., Romero J.(eds.), Climate Change 2023: Synthesis Report. IPCC, 2023.

4. Vonk JE, Tank SE, Bowden WB et al. Reviews and syntheses: Effects of permafrost thaw on Arctic aquatic ecosystems. *Biogeosciences* 2015;12:7129–67. https://doi.org/10.5194/bg-12-7129-2015.

5. Seppälä M. Synthesis of studies of palsa formation underlining the importance of local environmental and physical characteristics. *Quat Res* 2011;75:366–70. https://doi.org/10.1016/j.yqres.2010.09.007.

6. Verdonen M, Störmer A, Lotsari E et al. Permafrost degradation at two monitored palsa mires in north-west Finland. *Cryosphere* 2023;17:1803–19. https://doi.org/10.5194/tc-17-1803-2023.

7. Mamet SD, Chun KP, Kershaw GGL et al. Recent Increases in Permafrost Thaw Rates and Areal Loss of Palsas in the Western Northwest Territories, Canada. *Permafr Periglac Process* 2017;28:619–33. https://doi.org/10.1002/ppp.1951.

8. Borge AF, Westermann S, Solheim I et al. Strong degradation of palsas and peat plateaus in northern Norway during the last 60 years. *Cryosphere* 2017;11:1–16. https://doi.org/10.5194/tc-11-1-2017.

9. Tank SE, Striegl RG, McClelland JW et al. Multi-decadal increases in dissolved organic carbon and alkalinity flux from the Mackenzie drainage basin to the Arctic Ocean. *Environ. Res. Lett*. 2016;11:054015. http://dx.doi.org/10.1088/1748-9326/11/5/054015.

10. Abbott BW, Larouche JR, Jones JB et al. Elevated dissolved organic carbon biodegradability from thawing and collapsing permafrost. *J Geophys Res Biogeosci* 2014;119:2049–63. https://doi.org/10.1002/2014JG002678.

11. Wauthy M, Rautio M, Christoffersen KS et al. Increasing dominance of terrigenous organic matter in circumpolar freshwaters due to permafrost thaw. *Limnol Oceanogr Lett* 2018;3:186–98. https://doi.org/10.1002/lol2.10063.

12. Gandois L, Tananaev NI, Prokushkin A et al. Seasonality of DOC Export From a Russian Subarctic Catchment Underlain by Discontinuous Permafrost, Highlighted by High-Frequency Monitoring. *J Geophys Res Biogeosci* 2021;126(10):e2020JG006152, https://doi.org/10.1029/2020JG006152.

13. Olefeldt D, Roulet NT. Effects of permafrost and hydrology on the composition and transport of dissolved organic carbon in a subarctic peatland complex. *J Geophys Res Biogeosci* 2012;117:1005. https://doi.org/10.1029/2011JG001819.

14. Iannucci FM, Jones JB, Olson KL et al. Temperature and Flow Control Organic Carbon Metabolism in Boreal Headwater Streams. *J Geophys Res Biogeosci* 2024;129:e2024JG008281. https://doi.org/10.1029/2024JG008281.

15. Sipler RE, Kellogg CTE, Connelly TL et al. Microbial community response to terrestrially derived dissolved organic matter in the coastal Arctic. *Front Microbiol* 2017;8:255-275. https://doi.org/10.3389/fmicb.2017.01018.

16. Drake TW, Wickland KP, Spencer RGM et al. Ancient low-molecular-weight organic acids in permafrost fuel rapid carbon dioxide production upon thaw. *Proc. Natl. Acad. Sci. U.S.A*. 2015;112:13946–51. https://doi.org/10.1073/pnas.1511705112.

17. Mann PJ, Eglinton TI, McIntyre CP et al. Utilization of ancient permafrost carbon in headwaters of Arctic fluvial networks. *Nature Comm* 2015;6:1–7. https://doi.org/10.1038/ncomms8856.

18. Shirokova LS, Chupakov A, Zabelina S et al. Humic surface waters of frozen peat bogs (permafrost zone) are highly resistant to bio- and photodegradation. *Biogeosciences* 2019;16:2511–26. https://doi.org/10.5194/bg-16-2511-2019.

19. Vonk JE, Alling V, Rahm L et al. A centennial record of fluvial organic matter input from the discontinuous permafrost catchment of Lake Torneträsk. *J Geophys Res Biogeosci* 2012;117:3018. https://doi.org/10.1029/2011JG001887.

20. Chauhan A, Patzner MS, Bhattacharyya A et al. Interactions between iron and carbon in permafrost thaw ponds. *Sci. Total Environ*. 2024;946:174321. https://doi.org/10.1016/j.scitotenv.2024.174321.

21. Kuhn M, Lundin EJ, Giesler R et al. Emissions from thaw ponds largely offset the carbon sink of northern permafrost wetlands. *Sci. Rep*. 2018;8:1–7. https://doi.org/10.1038/s41598-018-27770-x.

22. Barbato RA, Jones RM, Douglas TA et al. Not all permafrost microbiomes are created equal: Influence of permafrost thaw on the soil microbiome in a laboratory incubation study. *Soil Biol Biochem* 2022;167:108605. https://doi.org/10.1016/j.soilbio.2022.108605.

23. Emsens WJ, van Diggelen R, Aggenbach CJS et al. Recovery of fen peatland microbiomes and predicted functional profiles after rewetting. *ISME J* 2020;14:1701–12. https://doi.org/10.1038/s41396-020-0639-x.

24. Muscarella ME, Boot CM, Broeckling CD et al. Resource heterogeneity structures aquatic bacterial communities. *ISME J* 2019;13:2183–95. https://doi.org/10.1038/s41396-019-0427-7.

25. Zhou L, Wu Y, Zhou Y et al. Terrestrial dissolved organic matter inputs drive the temporal dynamics of riverine bacterial ecological networks and assembly processes. *Water Res* 2024;249:120955. https://doi.org/10.1016/j.watres.2023.120955.

26. Crump BC, Amaral-Zettler LA, Kling GW. Microbial diversity in arctic freshwaters is structured by inoculation of microbes from soils. ISME J 2012;6:1629–39. https://doi.org/10.1038/ismej.2012.9.

27. Stadler M, del Giorgio PA. Terrestrial connectivity, upstream aquatic history and seasonality shape bacterial community assembly within a large boreal aquatic network. *ISME J* 2022;16:937–47. https://doi.org/10.1038/s41396-021-01146-y.

28. Savio D, Sinclair L, Ijaz UZ et al. Bacterial diversity along a 2600km river continuum. *Environ Microbiol* 2015;17:4994–5007. https://doi.org/10.1111/1462-2920.12886.

29. Caillon F, Besemer K, Peduzzi P et al. Soil microbial inoculation during flood events shapes headwater stream microbial communities and diversity. *Microb Ecol* 2021;82:591–601. https://doi.org/10.1007/s00248-021-01700-3.

30. Ruiz-González C, Niño-García JP, del Giorgio PA. Terrestrial origin of bacterial communities in complex boreal freshwater networks. *Ecol Lett* 2015;18:1198–206. https://doi.org/10.1111/ele.12499.

31. Borton MA, McGivern BB, Willi KR et al. A functional microbiome catalogue crowdsourced from North American rivers. *Nature* 2025;637:103–12. https://doi.org/10.1038/s41586-024-08240-z.

32. Anantharaman K, Brown CT, Hug LA et al. Thousands of microbial genomes shed light on interconnected biogeochemical processes in an aquifer system. *Nat Commun* 2016;7: 13219. https://doi.org/10.1038/ncomms13219

33. Vigneron A, Cruaud P, Langlois V et al. Ultra-small and abundant: Candidate phyla radiation bacteria are potential catalysts of carbon transformation in a thermokarst lake ecosystem. *Limnol Oceanogr Lett* 2020;5:212–20. https://doi.org/10.1038/ncomms13219.

34. Purkamo L, Ó Dochartaigh B, MacDonald A et al. Follwing the flow—Microbial ecology in surface- and groundwaters in the glacial forefield of a rapidly retreating glacier in Iceland. *Environ Microbiol* 2022;24:5840–58. https://doi.org/10.1111/1462-2920.16104 .

35. Chaudhari NM, Pérez-Carrascal OM, Overholt WA et al. Genome streamlining in Parcubacteria transitioning from soil to groundwater. *Environ Microbiome* 2024;19:1–12. https://doi.org/10.1186/s40793-024-00581-6.

36. He C, Keren R, Whittaker ML et al. Genome-resolved metagenomics reveals site-specific diversity of episymbiotic CPR bacteria and DPANN archaea in groundwater ecosystems. *Nat. Microbiol*. 2021;6:354–65. https://doi.org/10.1038/s41564-020-00840-5.

37. Gios E, Mosley OE, Weaver L et al. Ultra-small bacteria and archaea exhibit genetic flexibility towards groundwater oxygen content, and adaptations for attached or planktonic lifestyles. *ISME Commun* 2023;3(1):13, https://doi.org/10.1038/s43705-023-00223-x.

38. Wurzbacher C, Nilsson RH, Rautio M et al. Poorly known microbial taxa dominate the microbiome of permafrost thaw ponds. *ISME J* 2017;11(8):1938–41. https://doi.org/10.1038/ismej.2017.54.

39. Finnish Meteorological Institute. Download observations. https://en.ilmatieteenlaitos.fi/download-observations (2nd of June 2024 last access).

40. Huitu E, Arvola L. Water Chemistry and Bacterioplankton in Two Subalpine Rivers in Finnish Lapland. *Nord. Hydrol.* 2003;34:139-146. https://doi.org/10.2166/nh.2003.0033.

41. Dankers R. Sub-arctic hydrology and climate change a case study of the Tana River Basin in Northern Fennoscandia. *PhD. Thesis.* Utrecht University, 2002.

42. National Land Survey of Finland. MapSite. https://www.maanmittauslaitos.fi/en/opendata-licence-cc40 (2nd of July 2024, last accessed).

43. Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet J* 2011;17:10–2. https://doi.org/10.14806/ej.17.1.200.

44. Andrews S. FastQC A Quality Control tool for High Throughput Sequence Data. 2019.

45. Ewels P, Magnusson M, Lundin S et al. MultiQC: summarize analysis results for multiple tools and samples in a single report. *Bioinformatics* 2016;32:3047–8. https://doi.org/10.1093/bioinformatics/btw354.

46. Gruber-Vodicka HR, Brandon KS, Pruesse E. phyloFlash: Rapid SSU rRNA profiling and targeted assembly from metagenomes. *mSystems* 2020;5(5):e00920-20. https://doi.org/10.1128/msystems.00920-20.

47. McMurdie PJ, Holmes S. phyloseq: An R Package for Reproducible Interactive Analysis and Graphics of Microbiome Census Data. *PLOS ONE* 2013;8(4):e61217. https://doi.org/10.1371/journal.pone.0061217.

48. R Core Team. R: a language and environment for statistical computing. Vienna: R Foundation for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria 2023.

49. Posit team. RStudio: Integrated Development Environment for R. Posit Software, PBC, Boston, MA 2024.

50. Finnish Environment Institute. Hertta information system. https://www.syke.fi/en-US/Open_information (21st October 2024, last accessed).

51. Elovaara S, Zhao L, Asmala E et al. Changes in riverine dissolved organic matter caused by gypsum-induced flocculation. *J. Environ. Qual.* 2025:1-13. https://doi.org/10.1002/jeq2.70001.

52. Weishaar JL, Aiken GR, Bergamaschi BA et al. Evaluation of specific ultraviolet absorbance as an indicator of the chemical composition and reactivity of dissolved organic carbon. *Environ. Sci. Technol.* 2003;37:4702–8. https://doi.org/10.1021/es030360x

53. Massicotte P, Markager S. Using a Gaussian decomposition approach to model absorption spectra of chromophoric dissolved organic matter. *Mar. Chem.* 2016;180:24–32. https://doi.org/10.1016/j.marchem.2016.01.008.

54. Coble PG. Characterization of marine and terrestrial DOM in seawater using excitation-emission matrix spectroscopy. *Mar. Chem.* 1996;51:325–46. https://doi.org/10.1016/0304-4203(95)00062-3.

55. Massicotte P. eemR: Tools for Pre-Processing Emission-Excitation-Matrix (EEM) Fluorescence Data, R package version 0.1.5.9000 [code]. 2016. available at: https://github.com/PMassicotte/eemR

56. Murphy KR, Butler KD, Spencer RGM et al. Measurement of dissolved organic matter fluorescence in aquatic environments: An interlaboratory comparison. *Environ. Sci. Technol.* 2010;44:9405–12. https://doi.org/10.1021/es102362t.

57. Holmes RM, Aminot A, Kérouel R et al. A Simple and Precise Method for Measuring Ammonium in Marine and Freshwater Ecosystems. *Can. J. Fish. Aquat. Sci.* 1999: **56**(10): 1801-1808. https://doi.org/10.1139/f99-128.

58. Wickham H, Averick M, Bryan J et al. Welcome to the Tidyverse. J Open Source Softw 2019;4:1686. https://doi.org/10.21105/joss.01686.

59. Vu V. ggbiplot: a ggplot2 based biplot. R package version 0.55, 755. 2011. https://github.com/vqv/ggbiplot.

60. Asmala E, Stedmon CA, Thomas DN. Linking CDOM spectral absorption to dissolved organic carbon concentrations and loadings in boreal estuaries. *Estuar Coast Shelf Sci* 2012;111:107–17. https://doi.org/10.1016/j.ecss.2012.06.015.

61. Hulatt CJ, Kaartokallio H, Asmala E et al. Bioavailability and radiocarbon age of fluvial dissolved organic matter (DOM) from a northern peatland-dominated catchment: Effect of land-use change. *Aquat Sci* 2014;76:393–404. https://doi.org/10.1007/s00027-014-0342-y.

62. Massicotte P, Asmala E, Stedmon C et al. Global distribution of dissolved organic matter along the aquatic continuum: Across rivers, lakes and oceans. *Sci. Total Environ*. 2017;609:180–91. https://doi.org/10.1016/j.scitotenv.2017.07.076.

63. Moore MRN, Tank SE, Kurek MR et al. Ultrahigh resolution dissolved organic matter characterization reveals distinct permafrost characteristics on the Peel Plateau, Canada. *Biogeochemistry* 2024;167:99–117. https://doi.org/10.1007/s10533-023-01101-3.

64. Wauthy M, Rautio M. Permafrost thaw stimulates primary producers but has a moderate effect on primary consumers in subarctic ponds. *Ecosphere* 2020;11(6): e03099. https://doi.org/10.1002/ecs2.3099.

65. Knutson JK, Clayer F, Dörsch P et al. Water chemistry and greenhouse gas concentrations in waterbodies of a thawing permafrost peatland complex in northern Norway. *EGUsphere* [preprint] 2025. https://doi.org/10.5194/egusphere-2025-184.

66. Mustonen KR, Marttila H, Lehosmaa K et al. From thaw till fall: Interacting hydrology, carbon cycle, and greenhouse gas dynamics in a subarctic stream-lake continuum. *Sci. Total Environ*. 2024;957:177434. https://doi.org/10.1016/j.scitotenv.2024.177434.

67. Lundin EJ, Klaminder J, Giesler R et al. Is the subarctic landscape still a carbon sink? Evidence from a detailed catchment balance. *Geophys Res Lett* 2016;43:1988–95. https://doi.org/10.1002/2015GL066970.

68. Yvon-Durocher G, Hulatt CJ, Woodward G et al. Long-term warming amplifies shifts in the carbon cycle of experimental ponds. *Nat Clim Chang* 2017;7:209–13. https://doi.org/10.1038/nclimate3229.

69. Pengerud A, Cécillon L, Johnsen LK et al. Permafrost Distribution Drives Soil Organic Matter Stability in a Subarctic Palsa Peatland. *Ecosystems* 2013;16:934–47. https://doi.org/10.1007/s10021-013-9652-5.

70. Holmes RM, McClelland JW, Raymond PA et al. Lability of DOC transported by Alaskan rivers to the Arctic Ocean. *Geophys Res Lett* 2008;35. https://doi.org/10.1029/2007GL032837.

71. Tian R, Ning D, He Z et al. Small and mighty: Adaptation of superphylum Patescibacteria to groundwater environment drives their genome simplicity. *Microbiome* 2020;8:1–15. https://doi.org/10.1186/s40168-020-00825-w.

72. Brown CT, Hug LA, Thomas BC et al. Unusual biology across a group comprising more than 15% of domain Bacteria. *Nature* 2015 523:7559 2015;523:208–11. https://doi.org/10.1038/nature14486.

73. Dutkiewicz Z, Singleton CM, Sereika M et al. Proposal of Patescibacterium danicum gen. nov., sp. nov. in the ubiquitous ultrasmall bacterial phylum Patescibacteriota phyl. nov. *ISME Commun* 2025;5(1):ycae147. https://doi.org/10.1093/ismeco/ycae147.

74. Abiriga D, Jenkins A, Klempe H. Microbial assembly and co-occurrence network in an aquifer under press perturbation. *Ann Microbiol* 2022;72:1–13. https://doi.org/10.1186/s13213-022-01698-0.

75. Srinivas P, Peterson SB, Gallagher LA et al. Beyond genomics in Patescibacteria: A trove of unexplored biology packed into ultrasmall bacteria. *Proc. Natl. Acad. Sci. U.S.A* 2024;121:e2419369121. https://doi.org/10.1073/pnas.2419369121.

76. Mondav R, McCalley CK, Hodgkins SB et al. Microbial network, phylogenetic diversity and community membership in the active layer across a permafrost thaw gradient. *Environ Microbiol* 2017;19:3201–18. https://doi.org/10.1111/1462-2920.13809.

77. Besemer K, Singer G, Quince C et al. Headwaters are critical reservoirs of microbial diversity for fluvial networks. *Proc. R. Soc. B*. 2013;280:20131760. https://doi.org/10.1098/rspb.2013.1760.