A novel sorting method reveals superior metabolic activity of mononucleated over binucleated tetraploid hepatocytes

Yusuke Watanabe[1], Jonathan H. Sussman[2], Tomoya Anami[3], Moena Hattori[3], Saaya Yamane[3], Masaki Nishikawa[3], Yasuyuki Sakai[1,3], Takeshi Katsuda[3,*]

[1]Department of Bioengineering, Graduate School of Engineering, The University of Tokyo, Tokyo, Japan

[2]Medical Scientist Training Program, Perelman School of Medicine, University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA

[3]Department of Chemical System Engineering, Graduate School of Engineering, The University of Tokyo, Tokyo, Japan

*Correspondence: Takeshi Katsuda, email: tkatsuda@g.ecc.u-tokyo.ac.jp

Author contributions:

Y.W., M.N., and T.K. conceived the study. Y.W., T.A., and T.K. performed the methodology. Y.W. and T.K. designed the experiments. Y.W., M.H., and S.Y. conducted the experiments. Y.W., J.H.S., and T.K. analyzed the bioinformatics and wrote the manuscript. M.N., Y.S., and T.K. acquired the resources and funding. T.K. supervised the study.

ABSTRACT

Hepatocytes display notable ploidy diversity, varying both in the number of genomic copies and in the number of nuclei. Most mouse hepatocytes are tetraploid (4n), which can exist as either mononucleated (1x4n) or binucleated (2x2n) cells. Despite this distinction, these two cell types have traditionally been grouped and studied as a single population. One likely reason for this is that conventional ploidy-sorting methods classify hepatocytes based solely on total DNA content, without distinguishing between mononucleated and binucleated states. In this study, we developed a novel flow cytometry strategy to distinguish and isolate 1x4n and 2x2n hepatocytes. Our approach leverages Hoechst-Area to assess total ploidy and Hoechst-Height to differentiate mononucleated and binucleated hepatocytes. Transcriptome analysis comparing these two populations revealed that 1x4n hepatocytes exhibit a broader and more metabolically active gene expression profile. Importantly, this enhanced metabolic activity was independent of liver zonation, a well-known driver of metabolic heterogeneity in hepatocytes. Our findings uncover a previously underappreciated layer of functional diversity in the liver and provide a new framework for studying the physiological and pathological roles of nuclear configuration in hepatocytes.

MAIN

Unlike most other mammalian cell types, hepatocytes exhibit variable numbers of genomic copies (ploidy). The most common state in mouse hepatocytes is a binucleated tetraploid cell with each nucleus containing two genomic copies (2x2n), while other cells are mononucleated with four genomic copies (1x4n). A smaller percentage of hepatocytes contain two tetraploid nuclei (2x4n) or one nucleus with 8 genome copies (1x8n). While tetraploid hepatocytes have traditionally been studied as a single entity, several recent studies reported that binucleated hepatocytes decrease in pathological conditions such as after a partial hepatectomy[1] and carbon tetrachloride (CCl_4)-induced injury[2], suggesting that there are functional differences between mononucleated and binucleated hepatocytes.

A major limitation in studying the differences between 1x4n and 2x2n hepatocytes has been the lack of a method to isolate them. A flow cytometry study in plants reported that DNA-stained pollen populations could be distinguished based on their fluorescence and intensity parameters when using propidium iodide (PI), allowing for selective sorting of mononucleated and binucleated populations[3]. We hypothesized that a similar approach could be applied to hepatocytes, using Hoechst 33342, a DNA dye compatible with live cells. First, we confirmed that the conventional gating strategy enabled the sorting of freshly isolated mouse hepatocytes with different ploidy states (**Figure 1A, Supplementary Figure S1**), yielding mononucleated or binucleated cells whose nuclear sizes increased in accordance with their expected ploidy states (**Figure 1B**). We then analyzed hepatocytes using Hoechst-Area, representing the total integrated fluorescence

signal, and Hoechst-Height, representing the maximum intensity of the signal pulse. As

expected, 4n hepatocytes separated into two distinct populations, one with a higher

Hoechst-Height signal (termed 4n-top) and one with a lower signal (termed 4n-bottom)

(**Figure 1C**). To examine whether these populations correspond to 1x4n and 2x2n

hepatocytes, we sorted the whole 4n (4n-whole), 4n-top and 4n-bottom hepatocytes. The

proportions of mononucleated and binucleated cells were determined by microscopy after

plating, confirming that 4n-top and 4n-bottom populations were enriched for

mononucleated and binucleated cells, respectively (**Figure 1D-E**). Specifically, the

4n-whole population consisted of 48.0% of mononucleated and $52.0 \pm 3.7\%$ of binucleated

cells (mean $\pm$ SEM). In contrast, the mononucleated fraction in the 4n-top population

increased to $59.6 \pm 1.6\%$ while the binucleated fraction in the 4n-bottom population

reached $87.5 \pm 2.1\%$ of 2x2n cells (n = 9) (**Figure 1E**). To further test our hypothesis that

the 4n-top population is enriched for 1x4n hepatocytes, we performed three rounds of

sequential sorting. This gradually increased the proportion of 1x4n cells to $75.5 \pm 1.2\%$ (n =

3) (**Figure 1F-G**). In conclusion, our new gating strategy enables the selective sorting of

the 1x4n and 2x2n hepatocytes.

Using this sorting method, we next investigated differences of gene expression

profiles among hepatocytes with different ploidy. We sorted 2n-whole, 4n-whole, 4n-top

and 4n-bottom populations and performed RNA sequencing (RNA-seq). For 4n-top

hepatocytes, we used cells after a single round of sorting, rather than repeated sorting, to

ensure enough cells for RNA-seq. We first performed a global principal component analysis

(PCA) across all the samples, which revealed a clear separation between the 2n-whole

population and the three 4n populations along the first principal component (PC1) (**Figure**

2A). A binary comparison of the 4n-whole versus the 2n-whole population identified 1363

differentially expressed genes (DEGs, p-adj<0.05, |log2FC|>0.5) (**Figure 2B**) and 934

differentially enriched pathways (FDR<0.05) from the Gene Ontology Biological Process

(GOBP) category using a Gene Set Enrichment Analysis (GSEA) (top enriched pathways

shown in **Figure 2C**). These results indicated that our RNA-seq dataset reliably captures

the difference between 2n and 4n cells, while representing the overall similarity among the

4n populations.

We next investigated differences between 1x4n and 2x2n hepatocytes. Notably, the

4n-bottom population was positioned closer to the 2n-whole population along PC1 than the

other 4n populations were. This observation is consistent with a previous report suggesting

that the gene expression profile of 2x2n nuclei is more similar to that of 1x2n nuclei than

1x4n nuclei[4]. Despite the overall similarity, a direct comparison between 4n-top and

4n-bottom populations identified 54 DEGs (**Figure 2D**, representative genes shown in

Figure 2E). Moreover, GSEA revealed differential enrichment across multiple pathway

categories, including 99 in the GOBP (top enriched pathways shown in **Figure 2F**), 14 in

KEGG, 122 in Reactome and 10 in Hallmark (**Supplementary Figure S2A**). Notably,

many pathways enriched in 4n-top were involved in metabolism, including *branched chain*

amino acid catabolism and *porphyrin-containing compound metabolism*. Given that the

liver is spatially portioned into three metabolic zones (known as zonation), we considered

whether these differences were driven by zonation rather than ploidy. To assess the

potential influence of zonation, we curated zone-specific gene lists based on published

RNA-seq datasets[5, 6] GSEA using these lists revealed that Zone 1 genes were enriched

in 4n-bottom, while Zone 2 and 3 genes were enriched in 4n-top (**Figure 2G**). This result

suggests that 2x2n hepatocytes are more prevalent in Zone 1, whereas 1x4n hepatocytes are

enriched in Zone 2 and 3, consistent with previous reports[4, 7].

To further explore the relationship between ploidy and zonation, we examined

enrichment of zonation-associated metabolic pathways[8, 9]. Unexpectedly, Zone 1-related

fatty acid β-oxidation and Zone 3-related *xenobiotic metabolism* were both enriched in

4n-top hepatocytes (**Supplementary Figure S2B**). Moreover, 4n-top showed enrichment of

many other metabolism-related pathways that were not previously linked to zonation,

including   *nucleoside   bisphosphate   metabolism*   and   *tetrapyrrole   metabolism*

(**Supplementary   Figure   S2C**).   In   contrast,   we   found   only   two   metabolic   pathways

enriched in the 4n-bottom population: *formate metabolism* and *glutamine family amino acid*

metabolism (**Supplementary   Figure   S2D**).   Taken   together,   these   analyses   suggest   that

zonation   alone   cannot   fully   explain   the   observed   differences   between   1x4n   and   2x2n

hepatocytes.   Therefore,   it   is   likely   that   these   two   populations   have   inherently   distinct

metabolic   programs,   with   1x4n   hepatocytes   exhibiting   broader   and   more   active   metabolic

profiles.

In summary, our novel sorting strategy enabled the selective isolation of 1x4n and

2x2n hepatocytes, allowing us to uncover transcriptomic differences between these two

populations. Although the greatest differences existed between 4n and 2n cells, 1x4n and

2x2n   hepatocytes   exhibited   clear   transcriptional   differences,   including   more   active

metabolic   profiles   in   1x4n   hepatocytes   compared   to   2x2n   hepatocytes.   These   findings

motivate future work on the roles of different nucleation states on cellular function in the

liver. Recent developments in transcriptional profiling, such as spatial transcriptomics will

facilitate a more granular dissection of the phenotypes contained within each ploidy state.

METHODS

Mice

C57BL/6JJcl mice were purchased from CLEA Japan. Mice were fed with a standard diet CE-2 (10 kGy) (CLEA Japan) and kept in a 12-hour light–dark cycle. 7- to 15-week-old male mice were used for experiments.

Hepatocyte isolation

Hepatocyte isolation from the mouse liver was performed as previously described［10］. Briefly, livers were perfused with HBSS (Wako), Liver Perfusion Medium (Gibco), and HBSS with 5 mM CaCl2 (Wako) and 500 μg/ml collagenase (Wako) in order. Following perfusion, livers were mechanically dispersed with tweezers, resuspended in wash medium (DMEM (Wako) with 5% FBS (BioWest)) supplemented with 40 μg/ml DNase (Millipore), and filtrated with a 70 μm cell strainer. The cells were centrifuged at 4 °C at 48 × g for 5 min. Then, the cells were resuspended in complete percoll solution (10.8 ml percoll (Cytiva), 12.5 ml wash medium, and 1.2 ml 10× HBSS per liver) and centrifuged at 4 °C at 48 × g for 10 min. After a single wash with wash medium, hepatocytes and were use d for the downstream experiments.

Fluorescent staining

Hepatocytes were suspended with DMEM supplemented with 10% FBS and 1 x Penicillin-Streptomycin-Amphotericin B Suspension (P.S.A.) (Wako) at a concentration of 2e6 /ml. After 15 μg/ml Hoechst 33342 (Wako), 5 μM reserpine (Wako) and 40 μg/ml DNase were added, hepatocytes were incubated at 37 °C for 30 min, with inverted every 10 minutes. Following the centrifuge at 4 °C at 48 × g for 5 min, hepatocytes were

resuspended with flow buffer (HBSS supplemented with 1 M HEPES (Sigma), 1 M

MgCl$_2$(Wako), 200 mg/ml glucose (Sigma), 1 x P.S.A., 1 x MEM Non-essential Amino

Acids Solution (Wako), 1 X GlutaMAX Supplement (Wako), 1 mM sodium pyruvate

(Wako), 40 μg/ml DNase, 15 μg/ml Hoechst 33342 and 5 % FBS and pH was adjusted to

7.4 with NaOH (Wako)) at a concentration of 1e7 /ml. Then, hepatocytes were filtrated with

a 40 μm cell strainer, and 100 nm/ml TO-PRO-3 Iodide (Invitrogen) was added for staining

dead cells.

Flow cytometry

Flow cytometry was performed using SH800S (Sony) with a 130 μm nozzle.

Events other than hepatocytes and doublets were removed by gating on BSC-A vs. FSC-A

and FSC-H vs. FSC-A, respectively. Dead cells were removed according to TO-PRO-3

fluorescence by gating on FSC-H vs. APC-A. Then, hepatocytes were plotted by FSC-W vs.

Hoechst 33342-A or Hoechst 33342-H vs. Hoechst 33342-A. Sorted cells were collected

into 1:1 mixture of DMEM and FBS. After sorting, cells were immediately centrifuged at

4 °C at 48 × g for 5 min for 4n and 8n cells, or 800 × g for 10 min for 2n cells, respectively.

Cells were then resuspended with DMEM supplemented with 10% FBS and 1 x P.S.A. and

kept on ice before the downstream experiments.

Counting of mononucleated and binucleated hepatocytes

A portion of sorted hepatocytes were plated onto a 96-well plate with DMEM

supplemented with 10% FBS and 1 x P.S.A. Cells were incubated at 37 °C to for a few

hours, stained with 1 μg/ml Hoechst 33342 again and observed by fluorescence microscope

BZ-X810 (Keyence). The numbers of mononucleated and binucleated cells were counted,

respectively, and the proportions of these cells in sorted samples were calculated.

RNA extraction

178   80 ~ 300 K of hepatocytes were sorted by flowcytometry. RNA was extracted with

179   FastGene RNA basic kit including DNase I treatment. The concentration, A260/A280 and

180   A260/A230 of extracted RNA were measured with NanoDrop One (Thermo). RNA with

181   low purity was cleaned up with Monarch Spin RNA Cleanup Kit (10 μg) (New England

182   Biolabs).

RNA-seq

184   Library preparation and sequencing were performed by Azenta (Tokyo, Japan)

185   using NovaSeq (Illumina).

Bioinformatics for RNA-seq

Reads were aligned to the mouse genome (GRCm39) using STAR aligner with

default parameters[11]. The gene-count matrix was produced by featureCounts[12]. Batch

effects due to individual differences among mice were adjusted using ComBat. Then, to

compare gene expression between samples, expression levels were normalized on the

"median of ratios" method using DESeq2[12]. Binary comparisons were performed using

the results() function in DEseq2. Differentially expressed genes were defined following the

criteria: adjusted p value < 0.05 and $|\log_2(\text{fold-change})| > 0.5$. Differentially expressed

pathways were identified by gene sets enrichment analysis with the cut-off values of false

discovery rate < 0.05.

Preparation of lists of zonated genes

For Zone1- and 3-related genes, the counts per million matrix from

PRJNA556572[5] was used. Eight layers along hepatic zones were merged in pairs to

create the expression values for four layers, and Kruskal-Walis test was performed for

every gene. 188 of Zone1- and 259 of Zone3-related genes were identified by two criteria:

FDR < 0.05 and the median values of the layers exhibit a monotonic increase or decrease.

For Zone2-related genes, we used 46 of previously identified genes by Halpern *et al*[6].

ACKNOWLEDGEMENTS

205 This work was supported by grants from the Uehara Memorial Foundation and the

206 Nakatomi Foundation.

REFERENCES

1. Sladky VC et al. *Developmental Cell* 2020; **52**: 335-349.e7.

2. Lin YH et al. *Gastroenterology* 2020; **158**: 1698-1712.e14.

3. Kron P et al. *Cytometry Part A* 2015; **87**: 943-957.

4. Yang L et al. *Developmental Cell* 2023; **58**: 1996-2010.e6.

5. Ben-Moshe S et al. *Nature Metabolism* 2019; **1**: 899-911.

6. Halpern KB et al. *Nature* 2017; **542**.

7. Richter ML et al. *Nature Communications* 2021; **12**.

8. Paris J et al. *Hepatology* 2022; **76**: 1219-1230.

9. Martini T et al. *Annu. Rev. Pathol. Mech. Dis* 2025; **46**: 439-466.

10. Katsuda T et al. *Nature Communications* 2024; **15**.

11. Dobin A et al. *Bioinformatics* 2013; **29**: 15-21.

12. Love MI et al. *Genome Biology* 2014; **15**.

FIGURES

Figure 1

Figure 1. Development of a novel strategy for isolating 1x4n and 2x2n hepatocytes

(A) The conventional gating strategy for sorting whole 2n, 4n and 8n hepatocytes.

(B) Representative images of sorted hepatocytes with distinct ploidy states.

(C) The newly developed gating strategy for presumable 1x4n and 2x2n hepatocyte

populations.

(D) Representative images of plated 4n hepatocytes sorted with the conventional (**A**) or

new (**C**) gating strategy.

(E) Proportions of binucleated cells in each of the sorted 4n populations (n = 9).

Significance was assessed using a one-sided Welch's *t*-test.

(F) A representative image of plated cells after three rounds of sorting of 4n-top

hepatocytes.

(G) Proportions of mononucleated cells after three rounds of sorting of 4n-top hepatocytes

(n = 3). Significance was assessed using a one-sided Welch's t-test.

Figure 2

Figure 2. Transcriptomic analysis of hepatocytes with different ploidy states.

(A) Principal component analysis (PCA) across all samples.

(B) A volcano plot comparing gene expression between 4n-whole and 2n-whole hepatocytes. The log fold change shrunken data was plotted. Significant genes defined as p-adj < 0.05 and |Log2FC| > 0.5.

(C) Representative pathways differentially enriched between 4n-whole and 2n-whole

hepatocytes. The Gene Ontology Biological Process (GOBP) database was used. FDR,

false discovery rate.

(D) A volcano plot comparing 4n-top and 4n-bottom hepatocytes. The log fold change

shrunken data was plotted. Significant genes defined as p-adj < 0.05 and |Log2FC| > 0.5.

(E) A heatmap comparing 4n-top and 4n-bottom hepatocytes. Among the 54 DEGs, the top

10 genes enriched in 4n-top and 4n-bottom are shown, respectively. Genes are colored by

z-score normalized expression.

(F) Representative pathways differentially enriched between 4n-top and 4n-bottom

hepatocytes.

(G) GSEA using manually curated gene lists characteristic to each zone. The analysis was

conducted by comparing 4n-top (left) and 4n-bottom (right) hepatocytes. FDR, false

discovery rate. NES, normalized enrichment score.

SUPPLEMENTARY FIGURES

Supplementary Figure S1. Flow cytometry gating strategy.

Supplementary Figure S2. Additional GSEA results.

(A) Top enriched pathways by GSEA using KEGG, Hallmark, and Reactome categories.

(B-D) GSEA plots comparing 4n-top (left) and 4n-bottom (right). Zonated pathways (B).

pathways enriched in 4n-top (C), and pathways enriched in 4n-bottom (D).