The influence of HIV sense and antisense transcripts on stochastic HIV gene expression and reactivation

Kamil Więcek[1], Janusz Wiśniewski[1], and Heng-Chang Chen[1,2,*]

[1]Quantitative Virology Research Group, Population Diagnostics Center, Łukasiewicz Research Network – PORT Polish Center for Technology Development, Stablowicka 147, 54-066 Wrocław, Poland
[2]The Laboratory of Quantitative Virology, Centre for Advanced Materials and Technologies, Warsaw University of Technology, 19 Poleczki St, 02-822, Warsaw, Poland

*Correspondence: heng-chang.chen@port.lukasiewicz.gov.pl

Summary

In this study, we established dozens of single provirus-infected cellular clones offering various transcriptional phenotypes of HIV. We proposed that stochastic fluctuations in HIV transcription can appear at, at least, two levels: (1) the chromosomal landscape and (2) the in situ HIV integration site. In the former case, proviruses integrating at different genomic locations demonstrated a variety of transcriptional bursting and can be classified in noise space constructed based on the parameters associated with the coefficient of variation and using a mathematical model fitting a curve of exponential decay. In the latter case, stochastic HIV gene expression tended to be a pure epigenetic phenomenon: the identical provirus demonstrated fluctuations in its transcription with an elevated frequency. We observed similar expression patterns between sense and antisense RNA transcripts. Notably, both HIV long terminal repeats reacted to drug stimulation and may reveal distinct behaviors. Overall, our data suggest that HIV antisense transcripts could be involved in the stochastic nature of HIV transcription.

Keywords

HIV antisense transcripts; stochastic HIV gene expression; noise space; exponential decay model; provirus reactivation

Introduction

Noise in gene expression can be generated at multiple levels, ranging from transcription and translation to chromatin remodeling and its associated pathway, as well as gene-specific factors[1–7]. Studying such noise thus aids in a better understanding of how genes are regulated and their biological functions.

Human immunodeficiency virus (HIV) possesses two promoters, the 5' and 3' long terminal repeat (5'LTR and 3'LTR) promoters, and its transcription is an inherently stochastic process, generating strong fluctuations in HIV gene products[8–10]. Such spontaneous viral gene transcription[11–17] and translation[18–20] can persist during the administration of antiretroviral therapy (ART), HIV reservoirs thus are most likely transcriptionally heterogeneous.

At the chromosome level, the heterogeneity of HIV transcription has been observed from proviruses that integrate at different genomic locations associated with various epigenetic features, demonstrating inconsistency in HIV transcription[21–26]. At the virus level, HIV transcription has a long and rich history of investigations and is well known to be tightly regulated by the 5'LTR promoter[27–29], with which HIV hijacks host cellular transcription factors to control its gene expression[30]. One well-studied mechanism is the impact of RNA polymerase II (RNAPII) elongation on HIV transcription[27–29]; another is the effect of chromatin remodeling and nucleosome positioning[31,32]. Either way, evidence explaining how some viruses remain active and others turn into latency, reflecting the stochastic phenotype of HIV transcription, remains lacking.

Although it is currently unknown whether the HIV Tat positive-feedback loop is the only intrinsic molecular mechanism that governs the provirus transcriptional state (i.e., 5'LTR in the ON versus the OFF state), at present, the two-state promoter model (ON versus OFF) remains the most popular model for HIV transcriptional noise, an episodic process of gene expression[10,33] and has been applied to explain the molecular mechanism behind this bursty phenotype of gene expression. This two-state model relies on a specific regulatory structure to determine the thresholds that allow a biological system to either respond to or disregard an input signal. Tat transactivation is sufficient to induce a burst of HIV 5'LTR promoter activity that leads to fluctuations in HIV gene expression[34]; however, this model fails to generate a clear threshold response with deterministic bistability[35], implying that either the two-state model for the HIV 5'LTR promoter is atypical with a transient threshold[35,36], or an additional unknown determinant is involved in this circuit. Indeed, in physiological HIV infection conditions, latent HIV infections can be established at the early stage of infection[37–39], indicating that perhaps an additional molecular determinant that functions as an antagonist to the HIV Tat transactivator is most likely concomitantly present when the Tat positive-feedback circuit is initiated.

A recent study (preprint) suggested that, although the expression of HIV antisense transcripts (ASTs) could be influenced by Tat-dependent sense HIV transcription based on *in vitro* cellular models, the antisense RNA synthesis and the expression remain detectable in cells infected with Tat-deficient viruses, suggesting that most likely the expression of HIV ASTs is Tat-independent[40]. A profound mechanism behind the regulation between HIV ASTs and Tat, still requires further investigation. Antisense RNAs are a type of non-coding RNA and were discovered by Singer et al. in 1963[41]. Different endogenous antisense RNAs have been identified in mammalian cells[42]. In 1988, the existence of HIV ASTs and the encoded protein was first

postulated using computer analyses[43] and confirmed more than ten years later[44,45]. To date, HIV is well established as being able to perform antisense transcription from 3'LTR of their proviral genome[46,47]. HIV ASTs most likely encode at least one major gene product, named HIV antisense protein (ASP), 189 amino acids in length[48]. Like many other antisense RNAs discovered in animals and plants, HIV antisense RNAs repress the expression of their sense RNAs[49–51] and most likely contribute to maintaining proviruses in latency[51].

Although, at present, very little is known about the involvement of HIV ASTs in its pathogenesis, one recent study (preprint) provided evidence that the expression of HIV ASTs is associated with the efficiency of HIV integration[40]: the expression of HIV ASTs showed an increase, while the interaction between HIV integrase and the host-factor lens epithelium-derived growth factor (LEDGF)/p75 was impaired[40], suggesting that the expression of HIV ASTs might be tightly connected to HIV DNA integration, a critical step for HIV persistence infection. Furthermore, the presence of HIV ASTs is observed to be restricted to the group M HIV type 1[52], which is responsible for the human pandemic. These findings imply that HIV ASTs are most likely a functional and pathogenic mechanism left behind after a long period of evolutionary selection and might be critical in fulfilling its pathogenesis. Li et al.[53] (preprint) showed that the ectopic expression of ASTs in CD4+ T cells from people living with HIV on ART blocks latency reversal in response to pharmacologic and T-cell receptor stimulation, suggesting the functional role of HIV ASTs in enforcing transcriptional silencing. Understanding how HIV ASTs interact with their sense RNA transcription, subsequently governing the fluctuation of HIV gene expression may be beneficial to gaining better insights into HIV entry and relief from its latency.

Given that the amount of HIV ASTs in bulk infected cells is much lower than that of the sense RNA transcripts (from 1/100 to 1/2500 compared with sense RNAs)[45,50], quantitatively investigating HIV ASTs remains challenging. For this reason, in this study, we attempted to establish dozens of cellular models harboring a single provirus demonstrating various transcriptional phenotypes (**Figures 1A** and **2A**, n = 33), allowing for better characterization of the interplay between sense and antisense transcripts at the single-virus level. Such cellular models were named "single provirus-integrated cellular clones" (hereinafter, sinpro clones). We first utilized parameters measured via flow cytometry to define noise spaces[10], enabling the classification of sinpro clones harboring proviruses that demonstrated different transcriptional phenotypes. We further performed strand-specific reverse transcription (RT) followed by either quantitative PCR (qPCR) or digital PCR (dPCR) to track the dynamics of sense and antisense RNA transcription independently and provided the potential threshold levels of sense and antisense RNAs that determine stochastic fluctuations in HIV transcription across sinpro clones. We proposed mathematical models to fit curves of exponential decay, enabling better discrimination of distinct intensities of transcription and transcriptional noise in proviruses across sinpro clones. Finally, both HIV LTRs between sinpro clones in different groups reacted to drug (phorbol myristate acetate, PMA/ionomycin or ingenol-3-angelate, PEP005) stimulation.

Results

Establishment of sinpro clones and parameter setting

The selection procedure for sinpro clones is described in detail in the **STAR Methods**. After a screening of ten 96-well plates, 33 sinpro clones were obtained (**Figures 1A** and **2A**). We used a flow cytometry analysis to record and calculate a total of 16 parameters (**Figure 1B**) at 11 subsequent time points (**Figure S1A**). Descriptions of each parameter can be found in **Figure 1B** and the **STAR Methods**. Of note, given that the viability of each clone differs (some clones were robust in culture for a long period of time; some were not), not all sinpro clones could be included in every experiment planned in this work.

Based on the sequential measurement of the percentage of GFP-positive cells [(%) GFP(+)], categorizing these 33 clones into four groups was visually feasible: clones in which proviruses remained transcriptionally active over 11 subsequent time points, namely the active group (the A group, 10 clones); clones in which proviruses demonstrated a decrease in transcriptional activity over time, namely the attenuated active group (the A- group, 17 clones); clones in which the transcription of proviruses showed a sharp decrease and remained transcriptionally silent, namely the silent group (the S group, 5 clones) (**Figure 1A**; **Figures S1A** and **S1B**); and one clone (clone #814) in which the turnover of the state of HIV transcription showed elevated frequency (the F group) (**Figure 2A**). Given such a unique phenotype of provirus transcription in clone #814, more details will be discussed separately in the following content.

With respect to clones in the A, A-, and S groups, a tendency for a positive correlation was observed between (%) GFP(+) and the intensity of GFP fluorescence measured from GFP(+)-gated cells (GFP-fluo-gated; R^2 = 0.193; cor = 0.939) (**Figure S1C**) or from a population of total live cells (GFP-fluo-live; R^2 = 0.724; cor = 0.85) (**Figure S1D**). Of note, in addition to the mean coefficient of variation (CV) measured via flow cytometry at independent time points (namely population-based CV, pb-CV), we calculated CV for a time series across 11 time points (namely time-based CV, tb-CV; detailed in the **STAR Methods**) for comparison. Overall, a tendency for a negative correlation was observed between pb-CV, separated by the measurement from GFP(+)-gated cells (pb-CV-gated) or a population of total live cells (pb-CV-live), and GFP-fluo-gated and GFP-fluo-live, respectively (**Figure S1I**). A similar observation of a modestly negative correlation was observed when tb-CV was applied (**Figure S1J**). This observation indicated that the transcription of proviruses in the A, A-, and S groups aligned with that in previous findings[9,54]: a gene that exhibits a higher level of transcription has a lower degree of CV.

To better characterize sinpro clones across different groups, we constructed noise space (**Figures 1C-1G**) using FACS-relevant parameters, divided into four sets (**Figure 1B**). The definition of noise space was borrowed from the study of Dar et al.[10] with modifications. In principle, noise space consists of three parameters: CV, GFP expression, and the autocorrelation time ($\tau_{1/2}$). An analysis of the $\tau_{1/2}$ axis is critical given that it facilitates a direct comparison of data containing a wide variety of expression levels by removing the reciprocal dependence of noise magnitude on expression level[55]. In this work, $\tau_{1/2}$ was independently calculated based on mean transcription levels (namely $\tau_{1/2}$-fluo-live and $\tau_{1/2}$-fluo-gated) or mean-centered transcriptional noise (namely $\tau_{1/2}$-noise-live and $\tau_{1/2}$-noise-gated). The rationales and calculations are described in the **STAR Methods**. Overall, four sets of parameters were applied to construct

noise space: (1) pb-CV-live, GFP-fluo-live, and $\tau_{1/2}$-noise-live (**Figure 1D**), (2) pb-CV-gated, GFP-fluo-gated, and $\tau_{1/2}$-noise-gated (**Figure 1E**), (3) tb-CV-live, GFP-fluo-live, and $\tau_{1/2}$-fluo-live (**Figure 1F**), and (4) tb-CV-gated, GFP-fluo-gated, and $\tau_{1/2}$-fluo-gated (**Figure 1G**).

Visualization of transcriptional characteristics of sinpro clones in noise space

We first examined whether the mentioned four parameter sets were sufficient to classify sinpro clones across different (A, A-, and S) groups in two-dimensional (2D) projections of the principal component analysis (PCA) (**Figures S1E-S1H**). Overall, only a modest classification of clones across different groups in different parameter sets was observed (**Figures S1E-S1H**). A separation was demonstrated as the parameters related to the population of live cells were implemented (**Figures S1E** and **S1G**) compared with PCA conducted using the parameters related to GFP(+)-gated cells (**Figures S1F** and **S1H**). In each set, a positive correlation between the intensity of GFP fluorescence (GFP–fluo-live and GFP-fluo-gated) and a standard deviation (SD) was shown; a lack of correlation between GFP fluorescence and $\tau_{1/2}$, as well as CV was present (**Figures S1E-S1H**). Given that a linear correlation is expected between CV and the intensity of GFP fluorescence as $CV \propto 1/\sqrt{mean}$, a lack of correlation between these parameters may suggest either an atypical phenotype of stochastic HIV gene expression[35] in sinpro clones or the need for additional parameters to characterize its fluctuations in transcription.

A clear separation of clones across the three groups (A, A-, and S) was shown in noise space when the total number of live cells was applied irrespective of $\tau_{1/2}$ (**Figure 1D**, $\tau_{1/2}$-noise-live; **Figure 1F**, $\tau_{1/2}$-fluo-live). The separation between clones among the A, A-, and S groups was less effective in noise space constructed based on the parameters related to the GFP(+)-gated cells (**Figures 1E** and **1G**). Using a network-based analysis, we verified that the parameters related to the live cells (**Figures 1H** and **1J**) enabled the structural topology of an assortative network (assortativity = 0.045, **Figure 1H**; assortativity = 0.021, **Figure 1J**), whereas networks structured based on the parameters related to the GFP(+)-gated cells (assortativity = -0.088, **Figure 1I**; assortativity = -0.088, **Figure 1K**) failed to separate clones in which the proviruses possessed different transcriptional phenotypes. The same observation was seen in the correlation between every pair of the three parameters used for defining noise space assayed in a 2D projection (**Figures S1I-S1N**). We stress that transcriptional bursts varied across clones within the same group while comparing the noise frequency range (F_N)[33] (**Figure 1B**) calculated for the four parameter sets (**Figure S1O**: F_N-fluo-live, F_N-fluo-gated, F_N-noise-live, and F_N-noise-gated). Altogether, these observations suggest that the transcriptional characteristics in sinpro clones could be better defined based on the parameters related to a population of total live cells, rather than a focus on GFP(+)-sorted cells only. In the following analysis, the focus will be on these parameter sets.

Characteristics of the unique model, clone #814, unveiling stochastic HIV transcription with elevated frequency

We obtained one clone, clone #814, in which the turnover of the state of HIV transcription demonstrated an elevated frequency (**Figure 2B**), as visualized in the measurement of (%) GFP(+) cells over time (**Figure 2B**). Such a stochastic phenotype of HIV transcription reappeared as GFP-fluo-gated and GFP-fluo-live were measured at 11 subsequent time points (**Figure 2B**). We also observed an inconsistency between the

measures of (%) GFP(+) and GFP-fluo-gated: even though a small subset of (%) GFP(+) cells was present in the clonal population of clone #814, the highest level of proviral transcription was still detected (**Figure 2B**, highlighted by a red dotted line), whereas similar curves were observed between the measures of (%) GFP(+) and GFP-fluo-live (**Figure 2B**), implying that, in clone #814, GFP fluorescence measured from total live cells could better represent the heterogeneity of the clonal population shown by (%) GFP(+) cells. Remarkably, given that the site of HIV integration is identical regardless of the state of provirus transcription in clone #814, such a phenotype of stochastic HIV transcription should be a pure epigenetic phenomenon. The integration site of clone #814 will be discussed in the following content. Such stochastic fluctuations measured via (%) GFP(+) remained as additional independent measurements in a time series were conducted in 2025.

We wondered whether such a fluctuation in HIV transcription could be inheritable in both subpopulations of GFP(+) and GFP(-) of clone #814. To test this, we FACS-sorted a clonal population of clone #814 based on its GFP expression into two subpopulations, #814-GFP(+) and #814-GFP(-) (**Figure S2A**), and, once again, recorded the FACS-related parameters (**Figure 1B**) across six time points (**Figure 2C**). Although similar dynamics between GFP-fluo-gated and GFP-fluo-live were observed (**Figure 2C**), GFP-fluo-live showed a higher intensity in the #814-GFP(+) subpopulation compared with that measured in the #814-GFP(-) subpopulation (**Figure S2C**), whereas GFP-fluo-gated was indistinguishable between these two subpopulations (**Figure S2B**). In addition, the #814-GFP(-) subpopulation demonstrated a significant increase in the measures of CV-gated and CV-live compared with those measured in the #814-GFP(+) subpopulation (**Figure 2C**; **Figures S2D** and **S2E**). When examining the correlation between CV and GFP-fluo in both gated (**Figure S2F**) and live cells (**Figure S2G**) at six subsequent time points, we observed that even though clonal cells in the #814-GFP(-) subpopulation were more busty than those in the #814-GFP(+) subpopulation, all measures showed a negative regression, suggesting that the regulation of transcriptional bursting is likely inherited in both subpopulations. An undistinguishable F_N between these two subpopulations further verified this assumption (**Figure S2L**).

When we assayed the spatial correlation between all sinpro clones, including #814-GFP(+) and #814-GFP(-) in noise spaces (**Figures 2D-2G**), we observed that both subpopulations were in the spatial vicinity of each other, indicating a continuation of frequent turnover of transcriptional bursting in both subpopulations. The same observation was observed in the network properties structured based on parameters related to the total live cells: unsorted clone #814, #814-GFP(+), and #814-GFP(-) were in the vicinity of each other (**Figures 2H** and **2J**). Once again, networks structured based on parameters related to the GFP(+)-gated cells failed to clearly discriminate between sinpro clones across different groups (**Figures 2I** and **2K**). We noticed that after the addition of clone #814 and its subpopulations, clones from the A and A- groups were indistinguishable (**Figures 2H** and **2J**) compared with those from our previous observation, shown in **Figures 1H** and **1J**, suggesting the presence of additional subsets of clones between the A and A- groups. Additional mathematical approaches for the discrimination of sinpro clones and clone #814 and its subpopulations will be discussed in **Figure 4**. While applying 2D projections of PCA, the distribution of #814-GFP(+) and #814-GFP(-) was also irrelevant to the transcriptional pattern of sinpro clones (**Figures S2H-S2K**). Altogether, these findings suggest that the mechanism that most likely governs the transcription of the provirus in clone

#814 differed from that of other sinpro clones and that transcriptional bursting in clone #814 could be inheritable at a minimal level.

The provirus in clone #814 integrated in repetitive regions associated with active histone modifications

Given that we hypothesize that the fluctuation in HIV transcription in clone #814 could be a pure epigenetic phenomenon, we wondered whether any (epi)genomic features were involved. To identify the provirus integration site in clone #814, we developed a high-throughput method based on *in vitro* transcription, namely lentiviral integration site sequencing (LIS-seq) (**Figures S2M** and **S2N**; detailed in the **STAR Methods**). The specific PCR product was obtained before Illumina high-throughput sequencing (**Figure S2O**). We obtained 403,596 raw reads after sequencing, and 252,571 reads remained after quality control and read filtering using our tailored pipeline (see the **STAR Methods**), of which a total of 221,470 were mapped to the locus at 110,913,078 base pairs (bp) within the region of the protein-coding gene ankyrin repeat domain 10 (*ankrd10*) at the chromosome 13 (chr13) (**Figure S2P**). The identified integration was confirmed via PCR amplification using a pair of primers: one annealed to the local genomic content surrounding the integration site and another annealed to the start of the HIV 5'LTR (**Figure 2L**). We first assayed genomic features within the 10 kilobase pair (kb) region upstream and downstream from the site (chr13:110,913,078) of provirus integration. The first peak of repetitive sequences classified in the family short interspersed nuclear elements (SINEs) and Alu elements was present ~1.6 kb upstream from the integration site (**Figure 2M**). Furthermore, we observed intense ChIP-seq signals (fold change over control) of the acetylation of lysine 27 on histone H3 (H3K27ac) and trimethylation of lysine 4 on histone H3 (H3K4me3) within this 10 kb region: the first noticeable peak of H3K27ac was present at the position 661 bp upstream from the integration site, and the first peak of H3K4me3 was present at the position 108 bp upstream from the integration site (**Figure 2N**). Other ChIP-seq signals, including monomethylation of lysine 4 on histone H3 (H3K4me1), trimethylation of lysine 36 on histone H3 (H3K36me3), trimethylation of lysine 9 on histone H3 (H3K9me3), and trimethylation of lysine 27 on histone H3 (H3K27me3), appeared at a basal level (**Figure 2N**). The landscape of assayed ChIP-seq signals in the 20 kb genomic region surrounding the integration site is visualized in **Figure S2P**. Altogether, the provirus integration site in clone #814 was in the vicinity of the SINEs' and Alu elements' repetitive regions associated with H3K27ac and H3K4me3 histone modifications.

The involvement of HIV ASTs in characterizing distinct transcriptional phenotypes across sinpro clones

Although, at present, the fundamental mechanism behind the role of HIV ASTs and its consequence is not fully understood, accumulating evidence has shown the potential interference of HIV ASTs with its sense RNA transcription[45,49–51,56]. Here, we further hypothesize that HIV ASTs may also be involved in the regulation of stochastic transcription of the proviruses. In order to independently measure sense and antisense transcription, we applied the method of strand-specific reverse transcription (RT), followed by PCR amplification, which was previously published by Landry et al.[46], with modifications (**Figures S3**; see the **STAR Methods**). We first verified the expression of antisense RNAs in an HIV-based vector[23,57] in the sinpro clones; then, we observed the PCR products resulting from RT in the antisense direction (**Figure S3B**, lanes 10 and 11). The size of the antisense transcripts was no longer detectable starting from the middle of the gene encoding GFP in the antisense direction (**Figure S3B**, lanes 7 and 8). To increase the specificity of

this assay, we redesigned the RT primers, allowing for independent amplification of the sense (HCP#266) and antisense (HCP#264) transcripts (**Figure S3C**), optimized the RT-PCR condition (**Figure S3D**), and confirmed the specificity of the PCR product (**Figure S3E**). In parallel, we implemented a strand-specific RT assay, followed by digital PCR (dPCR), enabling an absolute quantitative measurement of RNA molecules at near-single-molecule resolution (**Figure S3F**). The dose-dependent (10X and 100X) detection of sense and antisense transcripts was achieved across 12 sinpro clones at three time points (**Figure S3G**). Altogether, these observations indicate the feasibility of using our methods to independently measure sense and antisense transcripts.

In a follow-up study, we applied strand-specific RT, followed by qPCR and dPCR, on nine clones, including clone #812 in the A group, three clones (#789, #795, and #802) in the A- group, four clones (#778, #788, #800, and #815) in the S group, and clone #814, to observe the sense and antisense RNA transcription at six subsequent time points (**Figures 3A** and **3B**; **Figure S4**) and to match them to corresponding FACS measurements recorded at the same time (**Figure S5**).

Given that the RNAs were isolated from the total cell population, here, we only took into account the parameters measured from the total live cells. We observed a positive correlation for GFP-fluo-live versus either sense or antisense RNAs measured via qPCR (**Figures S5A** and **S5B**) and dPCR (**Figures S5C** and **S5D**), respectively (dotted lines of linear regression shown in each panel). Moreover, a positive correlation between sense and antisense RNAs was observed over time (**Figures 3A** and **3B**). A similar trend for sense and antisense RNA transcription is presented in **Figure S4**. Although the magnitude of transcriptional bursting varied across individual clones, based on our methods, the abundance of sense transcripts was about 15 to 20 times (qPCR: mean, 28.32 and median, 12.51; dPCR: mean, 20.83 and median, 9.69) higher than that of antisense RNAs in the sinpro clones. Altogether, these findings suggest that fluctuations in HIV transcription in the sinpro clones most likely resulted from the transcriptional competition between sense and antisense transcripts, rather than alternative transcription between each other.

Intriguingly, while assaying the ratio between sense and antisense transcripts for clones in different groups, no difference was observed (**Figure 3C**). However, as we independently assayed the abundance of sense and antisense transcripts, an obvious deterministic threshold level was observed (**Figures 3D**, qPCR and **3E**, dPCR). Using both methods, a superior abundance of sense and antisense transcripts was observed in clones #812 and #814, followed by clones in the A- and S groups.

We further sought the threshold for determining HIV transcription in sinpro clones. We computed the minimum (Min.), lower quartile (Q1), median (Q2), upper quartile (Q3), and maximum (Max.) values, representing the ensemble of provirus transcripts in clones in each group, and observed a clear separation between clone #812 (the A group), clone #814 (the F group), and other clones (the A- and S groups) (**Figures 3F-3I**). As the Min. measured via qPCR was over 0.1 (arbitrary unit, a.u.; **Figure 3F**, sRNAs) and 0.05 (a.u.; **Figure 3G**, asRNAs), the clone tended to be in the A group. Using dPCR, a clear separation between the A group and other clones was observed at Q1, with values over 2 (a.u.; **Figure 3H**, sRNAs) and 0.25 (a.u.; **Figure 3I**, asRNAs). In all circumstances, the separation of clones between the A- and the S groups faltered: a visible separation lay at Q3 and Max. (**Figures 3F-3I**). The abundance of sense and antisense transcripts

measured in clone #814 either revolved around the threshold generated based on clone #812 using qPCR (**Figures 3F and 3G**) or lay between the thresholds between the A and A- groups (**Figures 3H** and **3I**) using dPCR. Altogether, our results suggest that the yield of sense and antisense transcripts, rather than their ratio, determined the stochastic phenotype of provirus transcription in sinpro clones.

Exponential decay model demonstrated a better fit of stochastic HIV transcription in sinpro clones

To strengthen the classification of these clones with an improving systematic quantification of stochastic variation across sinpro clones, we further implemented the mathematical model previously proposed by Singh et al.[9] to fit their transcriptional phenotypes. As demonstrated in their study, in most of the clones infected by minimal lentiviral vector noise aligned with the equation $CV^2 = C/\langle m \rangle$ (Equation 1), where C is the proportionality factor and $\langle m \rangle$ is the average number of GFP molecules per cell ($\langle \ \rangle$ denotes the average)[9,54] with a proportionality factor in the range between 15,000 and 65,000[9]. In this study, we call this equation the "reciprocal curve model" (**Figure 4B**) in order to distinguish it from our proposed model described below. While plotting the correlation between CV^2 and GFP fluorescence across 32 clones in the A, A-, and S groups, a lack of dependence between these two variables was observed (**Figure 4B**, R[2] = 0). A fragile correlation between these two parameters across eight clones in different groups was also observed as measurements were performed in a time series (**Figures S6A-S6F**).

We wondered about the cause of this observation. It might have resulted from the relatively few input data in our experimental setting compared with that in the previous studies that investigated stochastic variation in HIV gene expression[9,33] and in other organisms[54]. In an attempt to seek models that may better fit HIV transcription in sinpro clones, we formulated an equation of exponential decay, namely the "exponential decay model" (**Figure 4C**)

$$CV^2 = A*e^{(-B<m>)} + C \tag{Equation 2}$$

where CV and m are identical to the parameters present in Equation 1; e is Euler's number; and A, B, and C are fitted constants (**Figure 4E**). Intriguingly, we observed a slight redistribution of clones across the A, A-, and S groups, whereas clone #814 remained independent from other clones (**Figure 4C**). Based on the curve fitting using the exponential decay model, five new models were identified: (1) model I—clones in which a relatively high degree of transcriptional bursting (as represented by CV^2) was detected (R[2] = 0.77), (2) model II—clones in which moderate transcriptional bursting was detected (R[2] = 0.61), (3) model III—clones in which relatively low transcriptional bursting was detected (R[2] = 0.33), (4) model IV—the identical clones previously assigned to the S group (R[2] = 0), and (5) model V—clone #814 (R[2] = 0.58) (**Figure 4C**). Here, we note that no convincing model can be constructed for the clones in model IV because of the wide variety of transcription noise and the extremely low level of GFP expression, resulting in the appearance of a vertical line aligning all data points (**Figure 4C**). In addition, as we FACS-sorted the GFP(+) and GFP(-) subpopulations from clone #814, both subpopulations fell into the curve of model V (**Figure 4C**). This observation somehow support our previous conclusion that the mechanism governing transcriptional bursting in clone #814 could be inheritable at a minimal level. Individual clones assigned to the corresponding models are listed in **Figure 4C**.

Noticeably, here, better discrimination across clones in the A and A- groups was observed using the exponential decay model. Clones #758, #764, and #802, in which a high degree of noise was measured (model I), were clustered in the vicinity of those in model IV in the network property due to a high degree of noise appearing in clones in both models (**Figure 4D**), whereas the wide distribution of clones in model III may reflect a relative weakness of curve fitting ($R^2 = 0.33$) given the low degree of measured noise (**Figures 4C** and **4D**). Although the exponential model representing clone #814 remained isolated from others (**Figure 4C**), the subpopulation of #814-GFP(-), in which more noise was measured over time, tended to be close to the clones in models I and IV (**Figures 4D** and **4F**), whereas the unsorted #814 and the #814-GFP(+) subpopulations, in which relatively low levels of noise were measured (**Figure 4F**), were embedded in the clones in model II (**Figure 4D**). The A, B, and C constants are summarized in **Figure 4E**. Nevertheless, these findings suggest that the exponential decay model enabled a better classification of sinpro clones, in which the proviruses demonstrated distinct levels of fluctuations in HIV gene expression. In addition, this result, once again, highlighted that the mechanism governing stochastic HIV transcription in clone #814 may differ from that regulating provirus transcription in the clones in the other models.

To verify whether HIV sense and antisense transcription can also be depicted via the exponential decay model constructed based on GFP expression, we applied measures of sense and antisense RNAs from nine clones, shown in **Figure 3**, to fit the models. The curve of exponential decay recapitulated sense RNA transcription in model I with a better fit when transcription was measured using qPCR (**Figure 4G**, $R^2 = 0.92$) than when measured using dPCR (**Figure 4I**, $R^2 = 0.0$). The robustness of curve fitting faltered in the rest of the models as $\langle m \rangle$ was replaced with sRNAs: models II (**Figure 4G**, qPCR, $R^2 = 0.52$; **Figure 4I**, dPCR, $R^2 = 0.0$), III (**Figure 4G**, qPCR, $R^2 = 0.17$; **Figure 4I**, dPCR, $R^2 = 0.0$), IV (**Figure 4G**, qPCR, $R^2 = 0.009$; **Figure 4I**, dPCR, $R^2 = 0.0$), and V (**Figure 4G**, qPCR, $R^2 = 0.0$; **Figure 4I**, dPCR, $R^2 = 0.0$). Antisense RNA transcription measured via qPCR modestly fit the curves in models II (**Figure 4H**, $R^2 = 0.21$) and III (**Figure 4H**, $R^2 = 0.67$). Similar to the observation in sRNAs, a poor overall fit was observed when antisense transcription was measured using dPCR (**Figure 4J**). These findings imply that either different promoter activities occur between the HIV 5'LTR and 3'LTR or perhaps additional parameters are required to better describe the transcriptional phenotype of asRNAs. Of note, we also verified the effectiveness and robustness of our models using GFP expression measured via flow cytometry in the same nine clones: model I remained robust regarding its fit to the provirus transcription in clone #802 based on the GFP expression at six subsequent time points (**Figure S6G**, $R^2 = 0.99$), whereas semi-robustness of models II ($R^2 = 0.43$) and V ($R^2 = 0.35$) was observed (**Figure S6G**). A poor fit was observed in models III (**Figure S6G**, $R^2 = 0.0$) and IV (**Figure S6G**, $R^2 = 0.04$). We assumed that this phenomenon was due to either the few clones included here or the few time points used for the measurements. Clones in which proviruses demonstrated a high degree of noise and a low level of transcription (clones #778, #788, #800, and #815), once again, failed to formulate a curve of exponential decay (**Figure S6G**). Altogether, the proposed exponential decay model with the parameter measured at the protein level (GFP expression) enabled better recapitulation of the sinpro clones, in which the proviruses demonstrated distinct transcriptional phenotypes. The models were fragile at the transcription level, especially the test fitting asRNAs, which might be due to the more intense transcriptional bursting at the RNA level compared with that at the protein level.

Both HIV LTRs were responsive to drug stimulation

As a final step, we wondered whether the HIV 5' and the 3'LTRs that drive sense and antisense RNA transcription, respectively, behave in the same manner while clones are subjected to different activators and drugs. To test this, we continuously treated the previous clones, as well as #814-GFP(+) and #814-GFP(-) subpopulations, with phorbol 12-myristate 13-acetate (PMA) plus ionomycin (hereinafter, "PMA/ Ionomycin"), ingenol-3-angelate (PEP005), a PKC activator, or dimethyl sulfoxide (DMSO) as a negative control (**Figures S7A-S7I**). The fact that the proviruses in these clones and in two #814 subpopulations remained functional and were responsive to PMA/Ionomycin and PEP005 was evidenced by an increase in the measures of (%) GFP(+), GFP-fluo-gate, and GFP-fluo-live (**Figure S7A-S7I**). The reactivation shown in clone #814 and its two subpopulations suggests that the HIV 5'LTR remained transcriptionally accessible despite the presence of the integration site proximal to the genomic repetitive regions (**Figure 2M**).

Intriguingly, we observed that proviruses in clones assigned to different groups and models showed different responses to different drugs (**Figure S8**). In clones #812 (**Figure S8A**) and #789 (**Figure S8B**), provirus reactivation tended to elevate burst frequency[9,10], which led to increasing expression (GFP-fluo-live), showing a negative but linear relationship with burst size[9,10] [represented by $(pb\text{-}CV\text{-}live)^2$]. In contrast, in clones #795 (**Figure S8C**) and #802 (**Figure S8D**), provirus reactivation tended to synergetically affect burst frequency and burst size, which showed a positive but linear relationship. Of note, although both pairs of clones— #812 and #789, and clones #795 and #802—were assigned to different groups or models, clones in the same pair demonstrated topological proximity to the network property (**Figure 4D**). This may partially explain their similar response toward drugs. The reactivation pattern of the provirus in the S group was more diverse (**Figures S8E-S8H**). We assumed that this phenomenon might be due to the relatively high intrinsic background noise (shown in the Mock 1 and 2 controls and cells treated with DMSO) that lasted until the proviruses were reactivated. At present, it remains challenging for us to tell whether the intense magnitude of noise shown in reactivated proviruses in the S group resulted as an effect of drugs or was due to the viruses themselves. Different reactivation patterns were observed between clone #814 and its subpopulations (**Figure S8I**): unsorted clone #814 and #814-GFP(-) showed similar reactivation patterns to those observed for clones #812 and #789 (**Figure S8I**), whereas all data points from #814-GFP(+) shifted to the lower right quadrant (**Figure S8I**). In the former group, the presence of PMA/Ionomycin or PEP005 enhanced the mean value for GFP expression, and the presence of DMSO enhanced transcription noise (**Figure S8I**). In the latter case, the enrichment of the mean GFP expression was comparable between drug reactivation and the DMSO control (**Figure S8I**).

At the RNA level, the enrichment of both sense and antisense transcripts appeared in the presence of PMA/ Ionomycin or PEP005 across clones in all groups (**Figure 5A-5H**). A more intense augmentation was shown in sRNAs compared with asRNAs, reflecting a general expectation that the HIV 5'LTR activity should be superior to that of 3'LTR. Intriguingly, we observed that in clone #814, the provirus preferentially responded to PEP005, especially the 5'LTR that drives sense RNA transcription (**Figure 5H**); an extremely weak enrichment for both sense and antisense transcripts against the presence of PMA/Ionomycin (**Figure 5H**). Notably, although the overall abundance of asRNAs was less than that of sRNAs, we observed that the comparable enrichment between both sense and antisense RNAs when the fold changes were normalized by measures from clonal cells treated with DMSO in the A, A-, and S groups (**Figure 5I**), suggesting that the

activity of two HIV LTRs from the proviruses in clones assigned to these three groups were inducible irrespective of the presence of PMA/Ionomycin or PEP005. Although a tendency that two HIV LTRs were more susceptible to PMA/Ionomycin than PEP005 was observed in proviruses in clone #812 (the A group) (**Figure 5I**), a bias resulting from few data points here should also be taken into account. Notably, a discrepancy between regression lines formed by the enrichments of sense and antisense transcripts in clone #814 (the F group) indicated that the 5'LTR was more susceptible to PEP005, whereas the 3'LTR was more sensitive to PMA/Ionomycin (**Figure 5I**). Nevertheless, these findings offered a proof-of-concept that both HIV LTRs are responsive to provirus reactivation in the presence of different drugs for provirus reactivation.

Discussion

The stochastic expression of HIV proteins immediately after infection has been shown to be able to critically influence viral fate following active replication or post-integration latency[8,9,36,58]; however, the source of this noise is not fully understood. Mechanisms, including (1) stochastic fluctuations in the expression of the HIV Tat protein (Tat transcriptional pulses)[8,35,36,58], (2) the selection of HIV integration sites associated with epigenetic features[10,21–23], (3) RNAPII pausing[59], and (4) chromatin architecture and nucleosome occupancy[60], were reported to contribute to stochastic HIV gene expression.

In this study, we proposed that stochastic fluctuations in HIV transcription can appear at, at least two levels: (1) the chromosomal landscape and (2) the in situ HIV integration site (**Figure 6**). The former case, represented by sinpro clones in the A, A-, and S groups (**Figure 1A**), resonated with previous findings, indicating that the level of provirus transcription can be influenced by the local genomic context surrounding HIV integration sites coupled with genomic (e.g., chromatin accessibility)[21–23] and histone modifications[23,61]. Our results further set the stage for advancements in the classification of proviruses with varying levels of stochastic transcription in noise space (**Figures 1D** and **1F**) or based on mathematical modeling (exponential decay models) (**Figure 4C**).

In the latter case, represented by clone #814 (**Figure 2A**), the turnover of HIV transcription (active versus silent) showed elevated frequency (**Figure 2B**). In other words, stochastic fluctuations in HIV gene expression in this circumstance should be a pure epigenetic phenomenon, given that the provirus is identical regardless of the state of HIV transcription. We observed relatively intense signals of H3K27ac and H3K4me3 (**Figure 2N**) within the 10 kb genomic region, suggesting that the HIV-1 5'LTR is most likely transcriptionally accessible even though the integration site was identified in the vicinity of SINEs and Alu elements (**Figure 2M**). This conclusion can be supported by the provirus reactivation assay on clone #814 using PMA/Ionomycin or PEP005 (**Figure S7I**). Of note, Alu repeats, which are preferred for HIV integration, have been identified since the mechanism behind the bias of HIV integration site selection started to be addressed[62]. Heterochromatin is presently considered one of the genomic hubs where (deeply) latent proviruses persist via immune-mediated selection[63–67]. Future studies will be required to explore the correlation between clones harboring the same transcriptional phenotype as clone #814 and integration near repetitive genomic regions. Interestingly, unintegrated HIV DNA was proposed to adopt a repressive chromatin structure that competes with the transcription mechanism, leading to its silencing[68,69]; whether HIV antisense transcription can also be manifested around 2-LTR circles of unintegrated HIV DNA is presently unknown and will be studied in the future.

Notably, the phenotype of clone #814 resembles a subset of clonal populations that generated a variegated expression phenotype, namely phenotypic bifurcation (PheB), which can exhibit two subpopulations, one with bright GFP fluorescence and another GFP fluorescence-free[8]. As we separated the GFP(+) subpopulation of clone #814 from its GFP(-) subpopulation, we observed a minimal level of inheritance of such stochastic fluctuations (**Figure 2C**; **Figures S2F** and **S2G**): #814-GFP(+) and #814-GFP(-) were indistinguishable in noise space (**Figures 2D** and **2F**) and network properties (**Figures 2H** and **2J**). The clear separation between these two subpopulations based on the measures of (%) GFP(+) cells may suggest unequal epigenetic memory between these two subpopulations. Nevertheless, cumulative studies have shown

that long-term epigenetic inheritance mainly occurs in broad heterochromatin domains[70,71]; further investigations into these two subpopulations at higher order chromatin architectures will be required to gain more insights into the molecular mechanism underlying provirus transcription in such a clone.

In this study, we further hypothesized that HIV ASTs may also be involved in mechanisms that govern this stochastic phenotype of HIV gene expression, perhaps subsequently promoting the establishment of its latency. Indeed, viruses have developed various strategies to establish a latent state to ensure their persistence in their hosts. For instance, herpes viruses encode latency proteins that are responsible for switching the viral cycle from a lytic to a latent state[72]. Another example is the detection of ASTs, namely CMV latency-associated transcripts in bone marrow aspirating from naturally infected, healthy seropositive donors infected by human cytomegalovirus[73]. Retroviruses, in contrast, are known to persist through transcriptional silencing and do not encode latency proteins. In fact, in other human retroviruses, like human T-cell leukemia virus type 1 (HTLV-1), one of the best characterized ASTs in a viral system, its antisense gene product, HTLV-1 basic leucine zipper factor (HBZ), has been proven to functionally promote the survival and proliferation of HTLV-1-infected cells, thereby influencing progression into adult T-cell leukemia/lymphoma or HTLV-1-associated myelopathy/tropical spastic paraparesis[74]. HIV ASTs are supposed to suppress sense RNA transcription; however, at present, the fundamental mechanism behind HIV ASTs and their biological consequences remain elusive. A comprehensive understanding of the role of HIV ASTs involved in the regulation of HIV transcription may not only unveil an unprecedented mechanism used by HIV to establish a persistent infection but also reinforce concepts related to the fundamental mechanism of antisense RNA transcription.

To date, HIV ASTs have been demonstrated *in vitro* to be capable of promoting the initiation and maintenance of HIV latency by recruiting polycomb repressor complex 2 or chromatin modifying enzymes to the 5'LTR of the provirus, resulting in repression of the expression of their sense RNAs[49–51]. However, the expression of HIV ASTs *in vivo* has not been convincingly demonstrated, and only a few studies have examined its expression. Zapata et al.[50] reported low levels of ASTs in three people living with HIV on ART using real-time PCR in unstimulated CD4+ T cells. Mancarella et al.[56] detected ASTs in CD4+ T cells collected from both untreated and ART-treated patients after *ex vivo* stimulation with anti-CD3/CD28. Recently, a preprint reported by Capoferri et al.[75] provided sequence evidence for the expression of ASTs in unstimulated peripheral blood mononuclear cells collected from people living with HIV.

Presently, the fundamental mechanism behind the mechanistic interplay between HIV sense and antisense transcripts is unclear. In this study, we performed strand-specific RT followed by qPCR and dPCR to independently measure sRNAs and asRNAs in sinpro clones over time (**Figures 3A** and **3B**; **Figure S4**). We observed similar expression patterns (**Figure S4**) and a positive correlation between the abundance of sRNAs and asRNAs (**Figures 3A** and **3B**), indicating that, in sinpro clones, both the HIV 5' and 3'LTR are most likely active, whereas their strengths differ. This phenomenon may partially align with the previous study, indicating a predominant usage of the 5'LTR promoter by HIV (so-called "promoter suppression" in this study)[76]; however, in our study, we observed an indistinguishable ratio between sRNAs and asRNAs across sinpro clones (**Figure 3C**). Nevertheless, in our setting, based on the measurements conducted via qPCR and dPCR, more sense transcripts than antisense transcripts with clear threshold levels were detected

(**Figures 3D-3I**), which aligned with previous findings[45,50,77]. Intriguingly, Capoferri et al.[75] observed that in the subgenomic region of the HIV *env* gene, more HIV ASTs than their sense transcript counterparts can be targeted, highlighting that perhaps diverse transcript abundances between sRNAs and asRNAs are present throughout the HIV genome.

It is worth noting that whether the two-state promoter model (ON versus OFF) can completely capture stochastic fluctuations of HIV gene expression at the transcription level, especially antisense transcription, is presently unclear. It has been evidenced that perhaps the HIV Tat positive-feedback loop tends to be a monostable[8] or atypical two-state model with a transient threshold[35,36]. The robustness of our exponential decay models also faltered when sRNAs and asRNAs were applied (**Figures 4G-4J**), implying the need for additional factors to understand the entire regulatory mechanism. Nevertheless, further investigations into the interaction between sRNAs and asRNAs of individual genes will be needed.

The shock-and-kill strategy attempts to reactivate latent proviruses back into active replication states and to simultaneously purge HIV infections via ART[78,79]. Despite remarkable efforts, expectations for this shock-and-kill strategy have not been met yet[80]. As previously proposed, such limited success may be due to the stochastic nature of latency[8,36,81–83]. To compensate for the limitations of this strategy, alternative approaches are required. In the preprint reported by Li et al.[53], the ectopic expression of HIV ASTs in CD4+ T cells from people living with HIV under ART can block latency reversal in response to pharmacologic and T-cell receptor stimulation, underscoring the potential of HIV ASTs to be used in the block-and-lock functional cure strategy. Our results, as analyzed from another aspect, implied that both HIV LTRs were responsive to drug stimulation (**Figure 5A-5H**) and may react with different strengths from proviruses in sinpro clones assigned to different groups at both the protein (**Figure S8**) and RNA (**Figure 5I**) levels. Indeed, it has been known that two HIV LTRs possess different activities[76]; however, how they respond to drug reactivation remains unclear. Of note, the distinct effectiveness of curve fitting in exponential decay models observed between sense and antisense RNAs also supports the assumption that different promoter activities between two HIV LTRs exist (**Figures 4G-4J**). Distinct behaviors between 5' and 3' LTRs of the provirus in clone #814 highlighted the opportunity that perhaps in unique scenarios, e.g., pure epigenetics-driven stochastic fluctuations, two HIV LTRs may respond to stimuli in different ways (**Figure 5I**). A more precise investigation into the reactivity and the affinity between two HIV LTRs will thus be critical in determining the threshold for either increasing the efficacy of latency reversing agents by minimizing the transcription level of HIV ASTs or permanently locking proviruses in a state of (deep) latency by increasing the transcription of ASTs. We stress, as a previous study indicated, that drugs with different functionalities influence provirus transcriptional bursting differently[83]: transcriptional activators, such as tumor necrosis factor, are prone to increase burst frequency[9,10], whereas chromatin remodelers, such as histone deacetylase inhibitors, methylation inhibitors, and bromodomain inhibitors tend to cause modest increases in burst size[83]. Nevertheless, elucidating the transcriptional behaviors of proviruses in the presence of different drugs may result in better designed antiretroviral drugs and strategies.

The limitations of this study
This study was based on an experimental setting with 33 sinpro clones in total. Although we cannot rule out any bias introduced by these small-scale cellular models, we demonstrated a clear separation between sinpro

clones in noise space and using exponential decay models. Most importantly, a unique provirus transcriptional phenotype, shown in clone #814, may not be simply identified if a pool of infected cells is studied although only one clone harboring such a unique phenotype was obtained. We have confirmed the stability of the stochastic phenotype of provirus transcription in clone #814 based on two independent measurements in a time series: one was performed in 2023 (**Figure 2B**) and another in 2025 (data not shown). The phenotype reappeared, but faltered. At present, we cannot answer whether the flickering of such a phenotype will fade with time. And if so, does it result from multiple rounds of cell manipulation (freeze and thaw cells) or inheritance of unequal epigenetic memory in daughter cells, as previously discussed. More clones, in which proviruses demonstrate the same phenotype of stochastic transcription as that in clone #814 will be requisite to tackle this question. Finally, we cannot offer single-cell measurements for parameters, as previously performed[10,33]. Further investigations in large-scale experimental settings at the single transcripts resolutions in individual cells will be necessary to verify our observations on sinpro clones.

Acknowledgments

HCC acknowledges funding from the Narodowe Centrum Nauki (Sonata Bis Grant UMO-2022/46/E/NZ6/00022). We would like to thank Dr. Z. Debyser (KU Leuven) for his proofreading and critical feedback on the manuscript.

Author contributions

Conceptualization, H.-C.C.; methodology, H.-C.C., K.W. and J.W.; software, H.-C.C., and J.W., formal analysis, H.-C.C., K.W. and J.W., investigation, H.-C.C., K.W., and J.W.; resources, H.-C.C. and J.W.; data curation, H.-C.C., and J.W.; writing of original draft manuscript, H.-C.C., K.W., and J.W.; writing, manuscript review and editing, H.-C.C., K.W., and J.W.; visualization, H.-C.C., and J.W.; supervision, H.-C.C., project administration, H.-C.C.; funding acquisition, H.-C.C.

Declaration of interests

The authors declare no conflict of interest.

Declaration of generative AI and AI-assisted technologies

The authors declare no generative AI and AI-assisted technologies are used in this work.

Supplemental information index

Document S1. Figures S1–S8 and Tables S1-S5

Table S1. Excel file containing additional data too large to fit in a PDF, related to Figures 1 and 4.

Table S4. Excel file containing additional data too large to fit in a PDF, related to Figures 3 and 4.

Figure Legends

Figure 1

Figure 1. Characteristics of sinpro clones in the A, A-, and S groups

(**A**) Schematic representation of the experimental strategy of the establishment of single provirus-integrated cellular clones (sinpro clones). A total of 33 clones, in which proviruses demonstrated different transcriptional phenotypes were assigned to four different groups, including the active group (the A group, 10 clones), the attenuated active group (the A- group, 17 clones), the silent group (the S group, 5 clones) and the fluctuation group (the F group, 1 clone; see **Figure 2A**). (**B**) Table listing parameters used in this work. Detailed descriptions, resources, and calculations of parameters are provided where appropriate in the main text and the **STAR Methods**. (**C**) Schematic representation of the rationale of noise space previously reported by Dar et al. (2012). Three-dimensional (3D) noise space consists of noise magnitude represented by coefficient of variation (CV), the mean GFP expression, and noise autocorrelation time. Two-dimensional projections on the right-hand side of 3D noise space illustrate different behaviors between busty (marked in magenta) and constitutive (marked in grey) gene expression. (**D-G**) Noise space demonstrating a spatial distribution of sinpro clones in the A, A-, and S groups. Space is independently constructed using four

different sets of parameters, including (**D**) population-based (pb)-CV-live, GFP-fluo-live, and $\tau_{1/2}$-noise-live; (**E**) pb-CV-gated, GFP-fluo-gated, and $\tau_{1/2}$-noise-gated; (**F**) time-based (tb)-CV-live, GFP-fluo-live, and $\tau_{1/2}$-fluo-live; and (**G**) tb-CV-gated, GFP-fluo-gated, and $\tau_{1/2}$-fluo-gated. Spots marked in red, orange, and grey represent sinpro clones in the A, A-, and S groups, respectively. (**H-K**) Graphic networks illustrating the topological distribution of sinpro clones in the A, A-, and S groups in the network property. Once again, network properties are structured using four different sets of parameters mentioned above and correspond to respective noise space. Feature parameters used to construct corresponding networks and assortativity coefficients are placed beneath graphic networks. Spots marked in red, orange, and grey represent sinpro clones in the A, A-, and S groups, respectively. See also **Figure S1** and **Tables S1** and **S2**.

Figure 2

Figure 2. Characteristics of sinpro clone #814

(**A**) Schematic representation of the experimental strategy of the establishment of clone #814, in which the turnover of the state of HIV transcription showed elevated frequency. (**B**) Line plots illustrating the dynamics

of measures of parameters, including (%) GFP(+), GFP-fluo-gated, GFP-fluo-live, CV-gated, and CV-live recorded by flow cytometry with unsorted clone #814 at 11 subsequent time points. (**C**) Line plots illustrating the dynamics of measures of parameters, including (%) GFP(+), GFP-fluo-gated, GFP-fluo-live, CV-gated, and CV-live recorded by flow cytometry with clone #814-sorted GFP(+) and GFP(-) subpopulations at six subsequent time points. (**D-G**) Noise space demonstrating a spatial distribution of sinpro clones in the A, A-, and S groups as well as clone #814-sorted GFP(+) and GFP(-) subpopulations. Space is independently constructed using four different sets of parameters, including (**D**) population-based (pb)-CV-live, GFP-fluo-live, and $\tau_{1/2}$-noise-live; (**E**) pb-CV-gated, GFP-fluo-gated, and $\tau_{1/2}$-noise-gated; (**F**) time-based (tb)-CV-live, GFP-fluo-live, and $\tau_{1/2}$-fluo-live; and (**G**) tb-CV-gated, GFP-fluo-gated, and $\tau_{1/2}$-fluo-gated. Spots marked in red, orange, dark grey, light green, and light grey represent sinpro clones in the A, A-, S groups as well as clone #814-sorted GFP(+) and GFP(-) subpopulations, respectively. (**H-K**) Graphic networks illustrating the topological distribution of sinpro clones in the A, A-, and S groups, clone #814 as well as its two subpopulations in the network property. Once again, network properties are structured using four different sets of parameters mentioned above and correspond to the respective noise space. Feature parameters used to construct corresponding networks and assortativity coefficients are placed beneath graphic networks. Spots marked in red, orange, and grey represent sinpro clones in the A, A-, and S groups, respectively. Spots marked in green include clone #814 and its two subpopulations. (**L**) Expected results from PCR amplification confirming the provirus integration site identified using LIS-seq. 2.0% (w/v) agarose gel displaying a PCR product obtained from PCR amplification using a pair of primers flanking the junction between the HIV integration site and the host genome (lane 1, HCP#364: 560 bp; lane 2, HCP#365: 650 bp). (**M**) Line plot illustrating the signal profile of human repetitive regions. Smith Waterman (SW) alignment score representing the peak intensity is plotted in the y-axis in the genomic region of 50 kb upstream and downstream from the provirus integration site (chr13:110,913,078) (x-axis) in clone #814. (**N**) Line plot illustrating ChIP-seq signal profiles in the genomic region of 500 kb upstream and downstream from the provirus integration site (chr13:110,913,078) (x-axis) in clone #814. ChIP-seq signal value (fold change over control) represents the intensity of signal peaks shown on the y-axis. Red lines, including H3K27ac, H3K4me1, H3K4me3, and HeK36me3, represent active histone marks, whereas blue lines, including H3K9me3 and H3K27me3, represent repressive histone marks. See also **Figure S2** and **Tables S1-S3**.

Figure 3. Similar trends of consecutive expression between sRNAs and asRNAs

(**A**, **B**) Scatter plots representing the correlation between sense and antisense transcripts measured via strand-specific RT, followed by qPCR (**A**) or dPCR (**B**) at six subsequent time points. Facets separate the measurements at different time points. Spots marked in red, orange, black and green represent sinpro clones in the A (#812), A- (#789, #795, and #802), S (#778, #788, #800, #815), and F (#814) groups. (**C**) Boxplot representing the ratio between sense and antisense transcripts (sRNAs versus asRNAs) across nine sinpro clones. Facets separate the performance of measurements using qPCR or dPCR. Boxes marked in red, orange, black and green represent sinpro clones in the A (#812), A- (#789, #795, and #802), S (#778, #788, #800, #815), and F (#814) groups. (**D**, **E**) Boxplots representing the abundance of sense and antisense transcripts measured via qPCR (**D**) or dPCR (**E**) across nine sinpro clones. Facets separate the measures of sRNAs versus asRNAs. Boxes marked in red, orange, black and green represent sinpro clones in the A (#812), A- (#789, #795, and #802), S (#778, #788, #800, #815), and F (#814) groups. (**F**, **G**, **H**, **I**) Line plots representing the transcription threshold between sense and antisense transcripts measured via qPCR (**F**, sRNAs; **G**, asRNAs) or dPCR (**H**, sRNAs; **I**, asRNAs) across nine sinpro clones. The abundance of transcripts from clones in each group is compared in the range from the minimum (Min.), lower quartile (Q1), median (Q2), upper quartile (Q3), to the maximum (Max.) values (x-axis). Lines in red, orange, black and green represent sinpro clones in the A (#812), A- (#789, #795, and #802), S (#778, #788, #800, #815), and F (#814) groups, respectively. See also **Figures S3-S5** and **Table S4**.

Figure 4

Figure 4. Curve fitting of exponential decay depicting the transcription phenotypes of sinpro clones

(**A**) Schematic summary of distinct transcriptional phenotypes across 33 sinpro clones assigned to four different groups based on the measurement of (%) GFP(+) cells. (**B**) Plot of mean GFP expression (measured via GFP-fluo-live) versus transcriptional noise (measured via CV-live) across sinpro clones in the A, A-, and

S groups. Solid back line represents the fitting curve ($R^2 = 0$) based on the reciprocal curve model (Equation 1) mentioned in the main text. GFP-fluo-live is plotted with the values divided by 1,000; CV-live is plotted with the values divided by 100. (**C**) Plot of mean GFP expression (measured via GFP-fluo-live) versus transcriptional noise (measured via CV-live) across sinpro clones in all groups. Solid lines represent fitting curves based on the exponential decay model (Equation 2) described in the main text, including (1) model I marked in red (the exponential high noise group, $R^2 = 0.77$), (2) model II marked in orange (the exponential moderate noise group, $R^2 = 0.61$), (3) model III marked in azure (the exponential low noise group, $R^2 = 0.33$), (4) model IV marked in grey (the silent group, $R^2 = 0$), and (5) model V marked in green (the fluctuation group, $R^2 = 0.85$). Facets separate individual curves fitting in each model; sinpro clones assigned to respective groups are labeled on the right-hand side of facets. Of note, measures from #814-GFP(+) and #814-GFP(-) subpopulations are included in the analysis and assigned to the same model (model V) as unsorted clone #814. GFP-fluo-live is plotted with the values divided by 1,000; CV-live is plotted with the values divided by 100. (**D**) Graphic networks (assortativity coefficient = 0.0502) illustrating the topological distribution of sinpro clones across five models. Spots in red, orange, azure, grey and green represent clones in models I, II, III, IV, and V, respectively. (**E**) The formula of Equation 2 used for the simulation of curve fitting. The A, B, and C constants corresponding to individual models are summarized in the table beneath the formula. (**F**) Boxplot representing transcriptional noise (measured via CV-live) across 33 sinpro clones and FACS-sorted #814 GFP(+) and GFP(-) subpopulations. Clones assigned to models I, II, III, IV, and V are written in red, orange, azure, grey, and green, respectively. The y-axis is in logarithmic scale. **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$. (**G**, **H**, **I**, **J**) Plot of the abundance of RNA transcripts, separated between sRNAs and asRNA versus transcriptional noise (measured via CV-live) across nine sinpro clones used in **Figure 3**. The abundance of sense and antisense transcripts are measured using strand-specific RT, followed by either qPCR (**G**, sRNAs; **H**, asRNAs) or dPCR (**I**, sRNAs; **J**, asRNAs). Solid lines represent fitting curves based on the exponential decay model (Equation 2) described in the main text. Solid lines and spots corresponding to models I, II, III, IV, and V are marked in red, orange, azure, grey, and green, respectively. The A, B, and C constants corresponding to individual models are summarized in the table beneath each plot. See also **Figure S6** and **Table S1**.

Figure 5. Both HIV LTRs were responsive to drug stimulation.

(**A-H**) Bar charts representing the fold change (drugs versus DMSO) between sense and antisense RNA transcripts measured via strand-specific RT-qPCR (ratio to RPL27A) in sinpro clones treated with PMA/ Ionomycin or PEP005. Sipro clones used in this assay include clones (**A**) #812 (the A group), (**B**) #789 (the A- group), (**C**) #795 (the A- group), (**D**) #778 (the S group), (**E**) #788 (the S group), (**F**) #800 (the S group), (**G**) #815 (the S group), and (**H**) #814 (the F group). Results are the mean ± standard deviation (SD) of two independent experiments with duplicate samples. Bar in green and magenta represent the measures from sense and antisense transcripts, respectively. (**I**) Scatter plots representing the correlation between the enrichment of sense and antisense transcripts between sinpro clones treated with PMA/Ionomycin (x-axis) and those treated with PEP005 (y-axis). The enrichment is calculated using strand-specific RT-qPCR readouts from cells treated with drugs (PMA/Ionomycin or PEP005) normalized by readouts from cells treated with DMSO. Experiments were carried out from two independent biological replicates with duplicate samples. Both axes are in logarithmic scale. Dots and dotted lines marked in green and magenta represent measures from sense and antisense transcription, respectively.

Figure 6. The involvement of HIV asRNAs in stochastic fluctuations in HIV gene expression.

We propose that fluctuations in HIV transcription can appear at, at least two levels: (1) the chromosomal landscape and (2) the in situ HIV integration site (**A**). In the former case, proviruses integrating in different genomic locations (sinpro clones in the A, A-, and S groups) demonstrate different levels of HIV gene expression coupled with a variety of transcriptional bursting (**B**). In the latter case represented by clone #814 (the F group), stochastic HIV gene expression tends to be a pure epigenetic phenomenon: the identical provirus demonstrates the turnover of HIV infection with elevated frequency (**B**). We observed a minimum level of inheritance of such stochastic fluctuations (the provirus in the F group) and identified the provirus integration site, which is in the vicinity of the genomic repetitive regions associated with intense signals of H3K27ac and H3K4me3. In both cases, similar expression patterns between sense and antisense RNA transcripts in a time series were observed (**C**), suggesting that the independent yields of both transcripts determine the stochastic phenotype of provirus transcription, rather than their ratios.

STAR Methods

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Bacterial and virus strains		
Barcoded HIV-based vector (pHCC1)	Chen et al.[23,57]	N/A
pCMVΔR8.2	Chen et al.[23,57]	N/A
pVSV-G	Chen et al.[23,57]	N/A
Biological samples		
HEK293TN cell line	Cell line was kindly supported by the Bioengineering Group affiliated with Łukasiewicz Research Network - PORT Polish Center for Technology Development	N/A
Jurkat T cell line	Cell line previously obtained from the cell collection of the Center for Genomic Regulation, Barcelona	N/A
Chemicals, peptides, and recombinant proteins		
RPMI 1640 medium	Gibco	Cat. No. 42401-018
Fetal bovine serum (FBS)	Gibco	Cat. No. 10270-106
Penicillin-streptomycin	Gibco	Cat. No. 10378016
GlutaMAX (100x)	Gibco	Cat. No. 35050-061
Dulbecco's modified Eagle's medium (DMEM)	Gibco	Cat. No. 41965-039
7-Aminoactinomycin D (7-AAD)	BioLegend	Cat. No. 420404
RNase-free DNase I	New England Biolabs	M0303S
HpyCH4III	New England Biolabs	R0618S
Glycogen	ThermoFisher Scientific™	R0561
3M Sodium Acetate	Sigma-Aldrich	Cat. No. 71196-100ml
Isopropanol	CHEMPUR	Cat. No. 117515002
99.9% Ethanol	CHEMPUR	Cat. No. 113964810

E. coli Poly(A) Polymerase	NEB	M0276
Phusion™ High-Fidelity DNA Polymerases	Thermo Scientific	F534S
Phorbol myristate acetate (PMA)	Sigma-Aldrich	Cat. No. P8139
Ionomycin	ThermoFisher Scientific™	Cat. No. I24222
Ingenol-3-angelate (PEP005)	Sigma-Aldrich	Cat. No. SML1318
Dimethyl sulfoxide (DMSO)	NEB	cat. no. B0515A
Critical commercial assays		
AllPrep DNA/RNA Mini kit(50)	Qiagen	Cat. No. 80204
QubitTM RNA High Sensitivity (HS) kit	Thermo Fisher	Q32852
NG dART RT-PCR kit	EURx	E0802-02
PowerTrackTM SYBR Green Master Mix	Thermo Fisher	A46109
Probe PCR kit	Qiagen	Cat. No. 250102
HiScribe® T7 High Yield RNA Synthesis kit	NEB	E2040S
MinElute PCR Purification kit	Qiagen	Cat. No. 28004
NovaSeq 6000 SP Reagent Kit v1.5 (100 cycles)	Illumina	Cat. No. 20028401
Deposited data		
FACS measurements (measured in 2023) on 33 sinpro clones at 11 subsequent time points	This study	Table S1
Measures of four sets of parameters used to construct noise space	This study	Table S2
FACS measurements (measured in 2024) on FACS-sorted clone #814 GFP(+)- and GFP(-) subpopulations at 6 subsequent time points	This study	Table S3
FACS measurements (measured in 2024) coupled with sRNAs and asRNAs measured by strand-specific qPCR and dPCR from 9 sinpro clones at 6 subsequent time points	This study	Table S4
Illumina sequencing reads of clone #814 HIV integration sites	This study	PRJNA1244909
RepeatMasker	Smit, AFA, Hubley, R & Green, P. *RepeatMasker Open-4.0*. 2013-2015	http://www.repeatmasker.org
ChIP-seq data from Jurkat clone E61 and CD4+, αβ T cells	the ENCODE Project Consortium	

Human reference genome GRCh38 no-alt loci analysis set		https://ftp.ncbi.nlm.nih.gov/genomes/all/GCA/000/001/405/GCA_000001405.15_GRCh38/seqs_for_alignment_pipelines.ucsc_ids/GCA_000001405.15_GRCh38_no_alt_analysis_set.fna.bowtie_index.tar.gz
Experimental models: Cell lines		
Sinpro clone: clone #758 (the A- group & model I)	This study	Table S1
Sinpro clone: clone #760 (the A- group & model II)	This study	Table S1
Sinpro clone: clone #761 (the A- group & model II)	This study	Table S1
Sinpro clone: clone #762 (the A group & model II)	This study	Table S1
Sinpro clone: clone #764 (the A- group & model I)	This study	Table S1
Sinpro clone: clone #766 (the A- group & model II)	This study	Table S1
Sinpro clone: clone #767 (the A group & model II)	This study	Table S1
Sinpro clone: clone #768 (the A group & model II)	This study	Table S1
Sinpro clone: clone #770 (the A group & model III)	This study	Table S1
Sinpro clone: clone #775 (the S group & model IV)	This study	Table S1
Sinpro clone: clone #776 (the A- group & model II)	This study	Table S1
Sinpro clone: clone #777 (the A- group & model III)	This study	Table S1
Sinpro clone: clone #778 (the S group & model IV)	This study	Table S1
Sinpro clone: clone #779 (the A group & model II)	This study	Table S1
Sinpro clone: clone #780 (the A- group & model II)	This study	Table S1
Sinpro clone: clone #783 (the A- group & model II)	This study	Table S1
Sinpro clone: clone #784 (the A- group & model II)	This study	Table S1
Sinpro clone: clone #788 (the S group & model IV)	This study	Table S1
Sinpro clone: clone #789 (the A- group & model III)	This study	Table S1
Sinpro clone: clone #790 (the A- group & model III)	This study	Table S1
Sinpro clone: clone #791 (the A group & model II)	This study	Table S1
Sinpro clone: clone #795 (the A- group & model II)	This study	Table: S1
Sinpro clone: clone #799 (the A group & model III)	This study	Table S1

Sinpro clone: clone #800 (the S group & model IV)	This study	Table S1
Sinpro clone: clone #802 (the A- group & model I)	This study	Table S1
Sinpro clone: clone #803 (the A- group & model II)	This study	Table S1
Sinpro clone: clone #807 (the A- group & model II)	This study	Table S1
Sinpro clone: clone #809 (the A group & model II)	This study	Table S1
Sinpro clone: clone #811 (the A group & model II)	This study	Table S1
Sinpro clone: clone #812 (the A group & model II)	This study	Table S1
Sinpro clone: clone #813 (the A- group & model II)	This study	Table S1
Sinpro clone: clone #814 (the F group & model V)	This study	Table S1
Sinpro clone: clone #815 (the S group & model IV)	This study	Table S1
Oligonucleotides		
Primer sequences for RT, PCR, and qPCR	This study	Table S5
Probe sequences for dPCR	This study	Table S5
Software and algorithms		
Code and analytical pipelines	This study	https://github.com/ HCAngelC/ HIV_sinpro_clones _project
Python 3.11 statsmodels 0.14.4	Seabold, Skipper, and Josef Perktold. "statsmodels: Econometric and statistical modeling with python." *Proceedings of the 9th Python in Science Conference.* 2010	https:// www.statsmodels.o rg/stable/ index.html
Script for mapping HIV integration sites (LIS-seq) (Version 1.1.1)	This study	https://github.com/ wis-janusz/LISseq
R package "plot3D" (Version 1.4.1)	Soetaert, K. (2024). plot3D: Plotting Multi-Dimensional Data	https://cran.r-project.org/web/ packages/plot3D/ index.html
R package "stats" (Version 4.3.3)	R Core Team (2024). _R: A Language and Environment for Statistical Computing_. R Foundation for Statistical Computing, Vienna, Austria.	https://stat.ethz.ch/ R-manual/R-devel/ library/stats/html/ stats-package.html
R package "FactoMineR" (Version 2.11)	Lê et al. (2008)[85]	10.32614/ CRAN.package.Fa ctoMineR

R package "factoextra" (Version 1.0.7)	Kassambara, A., & Mundt, F. (2020)[93]. factoextra: Extract and Visualize the Results of Multivariate Data Analyses	https://cran.r-project.org/web/packages/factoextra/index.html
R package "Hmisc" (Version 5.2-2)	Harrell Jr., F., & Dupont, Ch. (2019). Hmisc: Harrell Miscellaneous	https://CRAN.R-project.org/package=Hmis
R package "igraph" (Version 2.1.4)	Csárdi et al. (2025)[86]	https://igraph.org
Python 3.11 SciPy 1.15.1	Virtanen et al. (2020)[87]	https://scipy.org/citing-scipy/
Pandas 2.2.3	The pandas development team (2024)[88]	https://pandas.pydata.org/docs/
Biopython 1.84	Cock et al. (2009)[89]	https://biopython.org/docs/1.84/
pysam 0.22.1	Li et al. (2009)[90]	https://pysam.readthedocs.io/en/v0.22.1/release.html
bowtie2 version 2.5.4	Langmead and Salzberg (2012)[91]	https://bowtie-bio.sourceforge.net/bowtie2/index.shtml
Other		
BD FACSAria™	BD Biosciences	https://www.bdbiosciences.com/en-us/products/instruments/flow-cytometers/research-cell-sorters/bd-facsaria-iii
BD FACSCanto™ II	BD Biosciences	https://www.bdbiosciences.com/en-se/products/instruments/flow-cytometers/clinical-cell-analyzers/facscanto
Qubit 4 Fluorometer	Invitrogen™	https://www.thermofisher.com/order/catalog/product/Q33226

CFX ConnectTM instrument	Bio-Rad	https://www.bio-rad.com/en-pl/product/cfx-connect-real-time-pcr-detection-system?ID=LN5TFG15
QIAcuity Digital PCR System (QIAcuity One, 5plex instrument)	Qiagen	Cat. No. 911020
QIAcuity 26-k, 24-well Nanoplates	Qiagen	Cat. No. 250001
NovaSeq 6000 sequencer	Illumina	https://www.illumina.com/systems/sequencing-platforms/novaseq.html

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Heng-Chang Chen (heng-chang.chen@port.lukasiewicz.gov.pl).

Materials availability

This study did not generate new unique reagents.

Data and code availability

Data availability.

The raw sequencing data generated during this study are available from Sequence Read Archive (SRA) (PRJNA1244909).

FACS measurements (measured in 2023) on 33 sinpro clones at 11 subsequent time points (**Figures 1A**, **2A, and 2B**; **Figure S1A**; **Figures 4B** and **4C**) are available at **Supplementary Information**, **Supplementary Table S1**.

Measures of four sets of parameters used to construct noise space (**Figures 1D-1G**; **Figures 2D-2G**) are available at **Supplementary Information**, **Supplementary Table S2**.

FACS measurements (measured in 2024) on FACS-sorted clone #814 GFP(+) and GFP(-) subpopulations at 6 subsequent time points (**Figure 2C**) are available at **Supplementary Information**, **Supplementary Table S3**.

FACS measurements (measured in 2024) coupled with sRNAs and asRNAs measured by strand-specific qPCR and dPCR from 9 sinpro clones at 6 subsequent time points (**Figures 3**; **Figures 4G-4I**; **Figure S4C**; **Figure S5**; **Figure S6**) are available at **Supplementary Information**, **Supplementary Table S4**.

Sequences of primers and probes are available at **Supplementary Information**, **Supplementary Table S5**.

Human, Homo sapiens, (hg38) DNA sequences for interspersed repeats and low complexity DNA sequences data were downloaded from RepeatMasker (https://www.repeatmasker.org/species/hg.html).

ChIP-seq data from Jurkat clone E61 and CD4+, αβ T cells for H3K27ac (ENCFF338LAI, ENCFF701MUZ), H3K4me1 (ENCFF002WRW, ENCFF704DNO, ENCFF068SDD, ENCFF545FKD), H3K4me3 (ENCFF860LFO, ENCFF017AFO, ENCFF923RAA, ENCFF067KWB, ENCFF280EJY, ENCFF855AXA), H3K36me3 (ENCFF536HDR, ENCFF275ZSW, ENCFF215CDE), H3K9me3 (ENCFF010VZL, ENCFF154QXP, ENCFF353AHL, ENCFF201NFN) and H3K27me3 (ENCFF720HUE, ENCFF514ZIY, ENCFF153LTQ, ENCFF669USD) were downloaded from the ENCODE Project Consortium[84] (see **KEY RESOURCES TABLE** for the download link).

Code availability.

Analytical pipelines and scripts generated in this work are available at GitHub (https://github.com/HCAngelC/HIV_sinpro_clones_project). The script for mapping provirus integration sites is available at GitHub (https://github.com/wis-janusz/LISseq). Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request. This paper reports the original code.

METHOD DETAILS

Transfection and viral Infection. For preparation of viral stocks, two million HEK293TN cells in 10-cm dishes were transfected with 3 μg pCMVΔR8.2, 2 μg pVSV-G and 6 μg barcoded pHCC1[23,57]. After 16 hr, the medium was replaced. The supernatant was collected 48 and 72 hr after transfection. Transfection efficiency was validated on the basis of the percentage of GFP(+) cells determined through FACS analysis. To determine the viral stock titer, five-fold serial dilutions of virus stocks were titrated in quadruplicate in 100 μL culture medium in 96-well culture plates. After 48 h, the percentage of GFP(+) cells was measured through FACS analysis. 1 million Jurkat cells in 6-well plates were infected with barcoded viral inocula at an MOI of approximately 0.03. The medium was replaced with 3 mL fresh RPMI 24 hr after infection. 48 hr after infection, the efficiency of infection was monitored by FACS analysis.

Establishment of single provirus-integrated cellular clone (sinpro clones) models. Sinpro clones established from a pool of infected Jurkat cells with HIV-based vector, pHCC1[23,57]. Infected cells were FACS-sorted using BD FACSAria™ (BD Biosciences) four days post-infection. The population of GFP(+) cells was diluted down to one cell per well in 96 well plates to obtain clones (**Figure 1A** and **2A**). After a period of about two weeks of growth, we screened the plate for GFP expression by measuring the percentage of GFP(+) cells using BD FACSCanto™ II (BD Biosciences). Starting from 10 96-well plates, eventually, 33 clones with expression of GFP were obtained. To further characterize the transcriptional phenotype of each clone, parameters, including the percentage of GFP(+) cells [(%) GFP(+)], GFP expression (GFP-fluo-gated and GFP-fluo-live), standard deviation (SD), the coefficient of variation (pb-CV-gated amd pb-CV-live) (**Figure 1B**) were recorded using BD FACSCanto™ II (BD Biosciences) twice every week for five weeks

(**Figure S1A**, a total of 11 subsequent time points). GFP expression and the coefficient of variation were independently measured from either GFP(+)-gated cells or a population of total live cells.

Cell culture. The human Jurkat T cell line (obtained from the cell collection of the Center for Genomic Regulation, Barcelona) was grown at 37 °C under a 95% air and 5% CO_2 atmosphere, in RPMI 1640 medium (Gibco, Cat. No. 42401-018) supplemented with 10% Fetal bovine serum (FBS; Gibco, Cat. No. 10270-106), 1% penicillin-streptomycin (Gibco, Cat. No. 10378016) and 1% GlutaMAX (100x) (Gibco, Cat. No. 35050-061). Jurkat cells were passaged every 2 d with a 1:5 dilution. HEK293TN cells were grown under the same conditions in Dulbecco's modified Eagle's medium (DMEM; Gibco, Cat. No. 41965-039). Sinpro clones were cultured in the same condition as the human Jurkat T cell line. Cells were tested for mycoplasma yearly.

The framework of noise space and computation of its components. The concept of noise space was borrowed from the study of Dar et al.[10]. It consists of three parameters: coefficient of variation (CV), GFP expression and the autocorrelation time ($\tau_{1/2}$). In this work, we divided parameters related to noise space into four sets: (1) population-based (pb)-CV-live, GFP-fluo-live, and $\tau_{1/2}$-noise-live, (2) pb-CV-gated, GFP-fluo-gated, and $\tau_{1/2}$-noise-gated, (3) time-based (tb)-CV-live, GFP-fluo-live, and $\tau_{1/2}$-fluo-live, and (4) tb-CV-gated, GFP-fluo-gated, and $\tau_{1/2}$-fluo-gated (**Figure 1B**). To summarize, noise space was visualized in GFP expression measured in either population of total live cells or GFP(+)-gated cells; each scenario was further coupled with either pb-CV and $\tau_{1/2}$-noise (**Figure 1D** and **1E**; **Figure 2D** and **2E**) or tb-CV and $\tau_{1/2}$-fluo (**Figure 1F** and **1G**; **Figure 2F** and **2G**). Code used for the visualization of noise space is described in the following section. Computation of the parameters CV and $\tau_{1/2}$ is described below.

Coefficient of variation (CV)

CV values for both gated and all live cells for each clone and each measurement time point were recorded directly from the flow cytometry results. The values were converted from percentage to ratio and their respective means were taken as pb-CV-gated and pb-CV-live. tb-CV-gated and tb-CV-live were measured based on the formula $cv_{tb} = \delta_{tb}/m_{tb}$; that is, the standard deviation δ_{tb} of GFP fluorescence normalized by the GFP fluorescence mean m_{tb}.

Noise autocorrelation time ($\tau_{1/2}$)

In this study, the calculation of $\tau_{1/2}$ was computed based on mean transcription levels (namely $\tau_{1/2}$-fluo-live and $\tau_{1/2}$-fluo-gated) or mean-centered transcriptional noise (namely $\tau_{1/2}$-noise-live and $\tau_{1/2}$-noise-gated). First, data for GFP-fluo-gated, GFP-fluo-live, CV-gated and CV-live were linearly interpolated to a regular spacing of one day. To calculate $\tau_{1/2}$-fluo-live and $\tau_{1/2}$-fluo-gated, autocorrelation function available in Python 3.11 statsmodels 0.14.4 (https://www.statsmodels.org/stable/index.html), the function acf(), was immediately applied for all possible time lags in days to the regularized GFP-fluo values. $\tau_{1/2}$-fluo-live and $\tau_{1/2}$-fluo-gated were recorded as the earliest time lags at which the autocorrelation was lower than 0.5.

The calculation of $\tau_{1/2}$-noise-live and $\tau_{1/2}$-noise-gated followed the concept described in Austin et al.[33], where $\tau_{1/2}$ was calculated based on transcriptional noise of individual clones against the noise averaged from clones in the same group[10,33]. Given such a limitation of our experimental instrument with which we cannot record CV for individual cells in a population of cells, using the same concept proposed by Austin et al.[33], we reformulated the calculation: regularized CV values were first mean-centered, followed by the application of autocorrelation functions available in Python 3.11 statsmodels 0.14.4 (https://www.statsmodels.org/stable/index.html), the function acf(). The first time lag, at which the autocorrelation function value dropped below 0.5 was recorded as $\tau_{1/2}$. The analytical pipeline is available at GitHub (https://github.com/HCAngelC/HIV_sinpro_clones_project).

Computing the noise frequency range (F_N)

The calculation of F_N was based on the definition $F_N = 1/\tau_{1/2}$, published by Austin et al.[33]. In this study, we took the values of $\tau_{1/2}$-fluo-live, $\tau_{1/2}$-fluo-gated, $\tau_{1/2}$-noise-live, $\tau_{1/2}$-noise-gated to compute F_N, respectively (**Figure S1O**).

Flow cytometry measurements. Flow cytometry experiments were conducted using the BD FACSCanto™ II (BD Biosciences) machine and standard protocols. FACS-sorting of GFP(+) and GFP(-) subpopulations of clone #814 was conducted using the BD FACSAria™ (BD Biosciences) machine. Single cells were filtered based on FSC-H and FSC-W filters. Alive cells were filtered using a fluorescent intercalator 7-Aminoactinomycin D (7-AAD; BioLegend, Cat. No. 420404) for staining the dead cells.

RNA extraction and DNase I treatment. Total RNA was extracted using the AllPrep DNA/RNA Mini Kit(50) (Qiagen, Cat. No. 80204) following the manufacturer's instructions. Subsequently, total RNA was subjected to DNase I treatment (New England Biolabs, M0303S) according to the producer's recommendations.

Strand-specific reverse transcription (RT). Total RNA concentration after DNaseI treatment was measured using a Qubit 4 Fluorometer with QubitTM RNA High Sensitivity (HS) kit (Thermo Fisher, Q32852). Subsequently, RNA samples were reverse transcribed using an NG dART RT-PCR kit (EURx, e0802-02) according to the producer's instructions. RT reactions were performed using 100 ng of RNA with respective RT primers: sRNAs, primer HCP#266; asRNAs, primer HCP#264; RPL27A (internal control gene), primer HCP#270. Primers HCP#264 and HCP#270 were added to the dPCR mixture. The RT reaction was incubated at 50 °C for 1 hr and inactivated at 85 °C for 5 min. Primer sequences are available at **Table S5**.

Quantitative polymerase chain reaction reaction (qPCR). qPCR was conducted using a CFX ConnectTM (Bio-Rad) instrument and PowerTrackTM SYBR Green Master Mix (Thermo Fisher, A46109) according to the producer's instructions. 2 μl cDNA was added in each reaction containing respective primers in the final concentration of 0.3 μM (sRNAs, HCP#265 and HCP#266; asRNAs, HCP#263 and HCP#264; RPL27A, HCP#269 and HCP#270). The reaction mix was predenaturated at 95°C for 10 min and ran for 30 cycles under following conditions: 95 °C for 10 min, 58 °C for 30 s, and 72 °C for 30 s, finished by melting curve analysis. Primer sequences are available at **Table S5**.

Digital PCR (dPCR). The dPCR procedure was performed using QIAcuity Digital PCR System (Qiagen, Cat. No. 911020 QIAcuity One, 5plex instrument), QIAcuity 26-k, 24-well Nanoplates (Qiagen, Cat. No. 250001) and Probe PCR Kit (Qiagen, Cat. No. 250102) according to the manufacturer's recommendations. In brief, primer-probe mixes targeting both sense [HCP#265, HCP#266 and HCP#272 (5'-end fluorescence: FAM; 3'-end quencher: BHQ1)] and antisense [HCP#263, HCP#264 and HCP#271 (5'-end fluorescence: HEX; 3'-end quencher: BHQ1)] transcripts were added in a total 10 µl mixture containing non-diluted cDNA. The reaction mix was predenatured at 95°C for 10 min and ran for 30 cycles under presented conditions: 95°C for 10 min, 58°C for 30 s, and 72°C for 30 s. Primer and probe sequences are available at **Table S5**.

Integration Site mapping. Schematic representation of the experimental workflow is shown in **Figure S2M**. Genomic DNA extraction was carried out using the AllPrep DNA/RNA Mini Kit(50) (Qiagen, Cat. No. 80204) following the manufacturer's instructions and digested using 2 µL 5,000 U/mL HpyCH4III (New England Biolabs, R0618S) and 2 µL 10 U/µL BplI (Thermo Scientific, ER1011) in Tango buffer complemented with 1× SAM in a 50 µL final volume at 37 °C for 3 hr. The digestion was terminated by incubating samples for 20 min at 65 °C. Digested DNA was used for *in vitro* transcription (IVT) using HiScribe® T7 High Yield RNA Synthesis Kit (NEB, E2040S). The reaction was performed according to the producer's recommendations with the addition of DTT at a final concentration of 5 mM and incubated at 37 °C for 16 hr. IVT products were digested with RNase-free DNase I (NEB, M0303S). DNase I-treated IVT products were diluted in a final volume of 500 µl RNase-free water, containing 1 µl 20 mg/mL Glycogen (Thermo Scientific™, R0561). 50 µl 3M Sodium Acetate (Sigma, Cat. No. 71196-100ml) and 500 µl isopropanol (CHEMPUR, Cat. No. 117515002) at room temperature were subsequently added to the mixture. The reaction was incubated at room temperature for 20 min. RNA was precipitated by centrifugation at 12,000 x g for 15 min, followed by two times washing using 70% ice-cold ethanol.

Purified IVT products were first verified using RT-PCR with primers HCP#237* and HCP#238* (**Table S5**) (**Figure S2N**). Subsequently, poly(A) tailing was subjected to purified IVT products using *E. coli* Poly(A) Polymerase (NEB, M0276) at 37°C for 30 min, followed by reverse transcription (NG dART RT-PCR kit, EURx, E0802-02) using a string of polythymine in the length of 33 bp, followed by two additional random nucleotides with options guanine, adenine, or cytosine at the 3' end of the primer sequence (HCP#273, **Table S5**) at 48°C for 1 hr. The obtained cDNA was used in two independent PCR reactions: one aims to identify the viral integration site (primers HCP#279 and HCP#290, **Table S5**), and another targets the HIV 5'LTR (HCP#237* and HCP#238*, **Table S5**), serves as a PCR positive control. Both reactions were performed using Phusion™ High-Fidelity DNA Polymerases (2 U/µL) (Thermo Scientific, F534S) according to the protocol recommended by the producer. 5 µl non-diluted cDNA was used for the reaction in a total 20 µl reaction mixture. The PCRs ran in the following cycling conditions: 30 cycles, initial denaturation at 98 °C for 10 min, denaturation at 98 °C for 10 s, annealing at 59 °C for 30 s, extension at 72 °C for 30 s, final extension at 72 °C for 5 min. The PCR products from the first round of integration site amplification were purified using MinElute PCR Purification Kit (Qiagen, Cat. No. 28004), followed by the second round of PCR with tailored indexing primers (HCP#304 and HCP#305, **Table S5**). The same reaction condition and

cycling parameters as the first round of PCR amplification were used. The final amplification products were purified, once again, using MinElute PCR Purification Kit (Qiagen, Cat. No. 28004) and visualized on 1.5% agarose gel (**Figure S2O**). The identified integration site of the provirus in clone #814 was verified using PCR amplification with two pairs of the primers, HCP#364 and HCP#238*, as well as HCP#365 and HCP#238*, which yield a specific 560 bp and 650 bp PCR product, respectively (**Figure 2L**). Primer sequences are available at **Table S5**.

Illumina high-throughput sequencing of HIV integration sites (IS). Integration site mapping amplicons were pooled at a final concentration of 2 nM for high-throughput sequencing. Libraries were sequenced as 50 bp single reads on a NovaSeq 6000 sequencer (Illumina) with the NovaSeq 6000 SP Reagent Kit v1.5 (100 cycles) (Illumina, Cat. No. 20028401).

Drug reactivation assay. 1 million sinpro clonal cells were seeded per well in 6-well plates for the performance of drug reactivation. 81 nM phorbol myristate acetate (PMA) (Sigma-Aldrich, Cat. No. P8139) with 1.134 µM ionomycin (ThermoFisher, Cat. No. I24222) or 200 nM ingenol 3-angelate (PEP005) (Sigma-Aldrich, Cat. No. SML1318) were applied to provirus reactivation. DMSO (NEB, cat. no. B0515A) was used as a negative control at a final concentration of 0.1% (v/v). After 24 hr incubation, cells were collected for cytometric measurements. Total RNA was isolated from cells treated with PMA/Ionomycin, PEP005, and DMSO, respectively, and used for RT-qPCR analysis.

Data analysis.

Plotting 2D and 3D projection of the parameters under the framework of noise space

2D projection

The correlation between every pair of three parameters, including CV, GFP expression and $\tau_{1/2}$, required for the construction of noise space was plotted using the R package "ggplot"[92] (**Figure S1I-S1N**; **Figure S2F and S2G**). Four different sets of the parameters, including (1) pb-CV-live, GFP-fluo-live, and $\tau_{1/2}$-noise-live, (2) pb-CV-gated, GFP-fluo-gated, and $\tau_{1/2}$-noise-gated, (3) tb-CV-live, GFP-fluo-live, and $\tau_{1/2}$-fluo-live, and (4) tb-CV-gated, GFP-fluo-gated, and $\tau_{1/2}$-fluo-gated, were plotted independently.

3D projection

In noise space, three coordinate axes were given as follows: (1) x-axis was referred to as GFP expression in a logarithmic scale (log2), (2) y-axis was referred to as $\tau_{1/2}$, and (3) z-axis was referred to as the square of CV in a logarithmic scale (log2) (**Figures 1D-1G** and **Figures 2D-2G**). Noise space was visualized using the function scatter3D() implemented in the R package "plot3D" (version 1.4.1) (see **KEY RESOURCES TABLE** for the download link), separated by the four sets of the parameters (**Figures 1D-1G**; **Figures 2D-2G**).

The performance of principal component assay (PCA) on parameters measured by flow cytometry across sinpro clones

Four parameters, including CV, SD, the mean of GFP expression and $\tau_{1/2}$ were included for the performance of PCA (**Figures S1E-S1H**; **Figures S2H-S2K**) using the function PCA() imbedded in the R package

"FactoMineR" (version 2.11)[85] and visualized using the function fviz_pca_biplot() embedded in the R package "factoextra" (version 1.0.7)[93]. PCA was independently performed, aligning with the criteria assigning four different sets of parameters in this study.

The network-based analysis on parameters measured by flow cytometry across sinpro clones

Five parameters, including CV, SD, the mean of GFP expression, $\tau_{1/2}$, and F_N were utilized for the calculation of correlation coefficients between two adjacent vertices in a network property using the function rcorr() embedded in the R package "Hmisc" (Version 5.2-2) (https://CRAN.R-project.org/package=Hmis). Subsequently, correlation matrices were transformed into data frames containing four columns. The first two columns served as ID columns, designating two adjacent enriched signatures irrespective of the direction. The remaining columns included the correlation coefficient and the associated *p*-value. To filter out weak or spurious connections, correlation coefficients with a *p*-value > 0.05 are excluded, thereby generating a correlation matrix. We further utilized this correlation matrix as the edge list to depict simple graphs (**Figures 1H-1K**; **Figures 2H-2K**; **Figure 4D**). In this list, the correlation coefficients representing the edge were calculated between two adjacent vertices that represent individual sinpro clones. The node list consisted of a series ID for individual sinpro clones, information about groups (**Figures 1H-1K**; **Figures 2H-2K**) or models (**Figure 4D**) assigned to respective clones. Once again, we structured individual network properties for four different sets of parameters, respectively. The network structure was then established using the function graph_from_data_frame() from the R package "igraph" (Version 2.1.4) (https://igraph.org)[86] with the following arguments: d for the edge list, vertices for the node list, and directed = FALSE to account for undirected edges in the network.

Mapping HIV integration site (IS) in clone #814

The code for mapping HIV integration sites was written in Python 3.12 using pandas 2.2.3[88], Biopython 1.84[89] and pysam 0.22.1[90] packages. The source code is available at https://github.com/wis-janusz/LISseq, while the latest release (Version 1.2.1) is available at https://github.com/wis-janusz/LISseq/releases.

50 bp single-read raw reads were filtered as follows: first, the 5'LTR sequence GGAGTGAATTAGCCCTTCCA was removed from the 5' end of reads. Any reads that do not contain the 5'LTR sequence with 1 mismatch allowed were discarded. Next, the poly(A) sequence in length of minimal 5 bp was removed from the 3' ends of the reads. Finally, reads that contain at least 1 ambiguous base or position with the phred quality score (q-score) lower than 20 were discarded.

The filtered reads were mapped to the human reference genome GRCh38 no-alt loci analysis set (see the **KEY RESOURCES TABLE** for the download link) using bowtie2 version 2.5.4[91] with default parameters. Alignments with quality lower than 20, as reported by bowtie2, were discarded. Filtered alignments were grouped by chromosome number and position. Loci with sequencing depth, i.e., the number of reads aligned to the same chromosome and position, of less than 1000 were discarded. The reported integration site indicated the single locus with the highest sequencing depth in this study.

Visualization of the landscape of (epi)genomic features surrounding the provirus integration site in clone #814

The landscape of repeating elements in the genomic region of 50,000 base pairs upstream and downstream of the provirus integration site (chr13:110,913,078) in clone #814 was plotted using R with default functions (**Figure 2M**). Smith Waterman (SW) alignment score was applied to represent the peak intensity. Human, Homo sapiens, (hg38) DNA sequences for interspersed repeats and low complexity DNA sequences data were downloaded from RepeatMasker (https://www.repeatmasker.org/species/hg.html).

ChIP-seq reads, including signals from H3K27ac, H3K4me1, H3K4me3, H3K36me3, H3K9me3 and H3K27me3 were downloaded from the ENCODE Project Consortium[84] (detailed in the **KEY RESOURCES TABLE** and **Data availability**) and plotted in the genomic region of 500,000 base pairs upstream and downstream of the provirus integration site (chr13:110,913,078) in clone #814 using R with default functions (**Figure 2N**). ChIP-seq signal value (fold change over control) was applied to represent the intensity of signal peaks. The landscape of chosen histone modifications is also visualized using the Integrative Genomics Viewer (IGV) (IGV version 2.19.1 for Linux)[94] (**Figure S2P**).

Obtaining the best-fitted curve of exponential decay model

Reciprocal curve model

The formula embedded in the reciprocal curve model was taken from the Equation previously reported by Singh et al.[9] to systematically quantify variability in GFP expression driven by the HIV 5'LTR across different HIV integration sites in the human genome. The Equation is written as follows:

$$CV^2 = C / \langle m \rangle \qquad \text{(Equation 1)}$$

, where C is the proportionality factor and $\langle m \rangle$ is the average number of GFP molecules per cell ($\langle \ \rangle$ denotes the average)[9,54]. The performance of curve fitting on 32 sinpro clones in A, A-, and S groups was conducted using tailored functions implemented in Python 3.11 SciPy 1.15.1[87] (available at GitHub https://github.com/HCAngelC/HIV_sinpro_clones_project). The curve was visualized in **Figure 4B**.

Exponential decay model

The performance of curve-fitting of exponential decay was based on the Equation written as follows:

$$CV^2 = A * e^{(-B<m>)} + C \qquad \text{(Equation 2)}$$

, where CV is the coefficient of variation, m is the mean GFP fluorescence, e is the Euler's number and A, B, and C are fitted constants (**Figure 4F**). Five distinct models of the exponential decay model across 33 sinpro clones were determined based on Equation 2 using tailored functions implemented in Python 3.11 SciPy 1.15.1[87] (available at GitHub https://github.com/HCAngelC/HIV_sinpro_clones_project). The curve was visualized in **Figure 4C**. Values of the constants A, B, C in each model were shown in **Figure 4F**.

To further probe the influence of sense- and antisense RNAs on stochastic variation driven by the HIV 5'LTR and the 3'LTR, independently at the transcription level, the parameter value of $\langle m \rangle$ was replaced by the values measured from either sRNAs (**Figures 4H** and **4J**) or asRNAs (**Figures 4I** and **4K**) using

strand-specific qPCR and dPCR. To determine a set of initial constants A, B, and C to fit the model, the constant C was estimated as its value has to be smaller than the minimum value of CV^2 and be greater than zero. The constants A and B were determined based on a linear model transformed from Equation 2. The mathematical operation was demonstrated as follows:

$$CV^2 = A*e^{(-B<m>)} + C$$

$$CV^2 - C = A*e^{(-B<m>)}$$

$$log(CV^2 - C) = log(A) - B<m>*e \quad \textit{\# A linear model.}$$

Step by step operation code is available at GitHub (https://github.com/HCAngelC/ HIV_sinpro_clones_project). In brief, models were tested using the function nls() implemented in the R package "stats" (version 4.3.3) (see the **KEY RESOURCES TABLE** for the download link). Estimated values of each constant in models were recorded using the function summary() implemented in the R. Of note, in the circumstance that the default Gauss-Newton algorithm in the function nls() cannot support the estimated values for the initiation of curve fitting, a feasible alternative of Golub-Pereyra algorithm for Partial Least Squares model with the argument alg = "plinear" built-in in the function nls() was applied. The plot of fitted curves (**Figure 4**) were visualized using R with default functions.

QUANTIFICATION AND STATISTICAL ANALYSIS

Statistics.

All statistical tests were performed using Python or R (version 4.3.3) with default options and specific details are provided in the main text and figure legends where applicable.

References

1. Thattai, M., and van Oudenaarden, A. (2001). Intrinsic noise in gene regulatory networks. Proc Natl Acad Sci U S A *98*, 8614–8619. https://doi.org/10.1073/pnas.151588598.

2. Ozbudak, E.M., Thattai, M., Kurtser, I., Grossman, A.D., and van Oudenaarden, A. (2002). Regulation of noise in the expression of a single gene. Nat Genet *31*, 69–73. https://doi.org/10.1038/ng869.

3. Paulsson, J. (2004). Summing up the noise in gene networks. Nature *427*, 415–418. https://doi.org/10.1038/nature02257.

4. Raser, J.M., and O'Shea, E.K. (2004). Control of stochasticity in eukaryotic gene expression. Science *304*, 1811–1814. https://doi.org/10.1126/science.1098641.

5. Becskei, A., Kaufmann, B.B., and van Oudenaarden, A. (2005). Contributions of low molecule number and chromosomal positioning to stochastic gene expression. Nat Genet *37*, 937–944. https://doi.org/10.1038/ng1616.

6. Colman-Lerner, A., Gordon, A., Serra, E., Chin, T., Resnekov, O., Endy, D., Pesce, C.G., and Brent, R. (2005). Regulated cell-to-cell variation in a cell-fate decision system. Nature *437*, 699–706. https://doi.org/10.1038/nature03998.

7. Rosenfeld, N., Young, J.W., Alon, U., Swain, P.S., and Elowitz, M.B. (2005). Gene regulation at the single-cell level. Science *307*, 1962–1965. https://doi.org/10.1126/science.1106914.

8. Weinberger, L.S., Burnett, J.C., Toettcher, J.E., Arkin, A.P., and Schaffer, D.V. (2005). Stochastic gene expression in a lentiviral positive-feedback loop: HIV-1 Tat fluctuations drive phenotypic diversity. Cell *122*, 169–182. https://doi.org/10.1016/j.cell.2005.06.006.

9. Singh, A., Razooky, B., Cox, C.D., Simpson, M.L., and Weinberger, L.S. (2010). Transcriptional bursting from the HIV-1 promoter is a significant source of stochastic noise in HIV-1 gene expression. Biophys J *98*, L32–L34. https://doi.org/10.1016/j.bpj.2010.03.001.

10. Dar, R.D., Razooky, B.S., Singh, A., Trimeloni, T.V., McCollum, J.M., Cox, C.D., Simpson, M.L., and Weinberger, L.S. (2012). Transcriptional burst frequency and burst size are equally modulated across the human genome. Proc. Natl. Acad. Sci. U. S. A. *109*, 17454–17459. https://doi.org/10.1073/pnas.1213530109.

11. Dornadula, G., Zhang, H., VanUitert, B., Stern, J., Livornese, L., Jr, Ingerman, M.J., Witek, J., Kedanis, R.J., Natkin, J., DeSimone, J., et al. (1999). Residual HIV-1 RNA in blood plasma of patients taking suppressive highly active antiretroviral therapy. JAMA *282*, 1627–1632. https://doi.org/10.1001/jama.282.17.1627.

12. Ishizaka, A., Sato, H., Nakamura, H., Koga, M., Kikuchi, T., Hosoya, N., Koibuchi, T., Nomoto, A., Kawana-Tachikawa, A., and Mizutani, T. (2016). Short Intracellular HIV-1 Transcripts as Biomarkers of Residual Immune Activation in Patients on Antiretroviral Therapy. J Virol *90*, 5665–5676. https://doi.org/10.1128/JVI.03158-15.

13. Yukl, S.A., Kaiser, P., Kim, P., Telwatte, S., Joshi, S.K., Vu, M., Lampiris, H., and Wong, J.K. (2018). HIV latency in isolated patient CD4 T cells may be due to blocks in HIV transcriptional elongation, completion, and splicing. Sci Transl Med *10*. https://doi.org/10.1126/scitranslmed.aap9927.

14. Fischer, M., Günthard, H.F., Opravil, M., Joos, B., Huber, W., Bisset, L.R., Ott, P., Böni, J., Weber, R., and Cone, R.W. (2000). Residual HIV-RNA levels persist for up to 2.5 years in peripheral blood mononuclear cells of patients on potent antiretroviral therapy. AIDS Res Hum Retroviruses *16*, 1135–1140. https://doi.org/10.1089/088922200414974.

15. Günthard, H.F., Havlir, D.V., Fiscus, S., Zhang, Z.Q., Eron, J., Mellors, J., Gulick, R., Frost, S.D., Brown, A.J., Schleif, W., et al. (2001). Residual human immunodeficiency virus (HIV) Type 1 RNA and DNA in lymph nodes and HIV RNA in genital secretions and in cerebrospinal fluid after suppression of viremia for 2 years. J Infect Dis *183*, 1318–1327. https://doi.org/10.1086/319864.

16. Halvas, E.K., Joseph, K.W., Brandt, L.D., Guo, S., Sobolewski, M.D., Jacobs, J.L., Tumiotto, C., Bui, J.K., Cyktor, J.C., Keele, B.F., et al. (2020). HIV-1 viremia not suppressible by antiretroviral therapy can originate from large T cell clones producing infectious virus. J Clin Invest *130*, 5847–5857. https://doi.org/10.1172/JCI138099.

17. White, J.A., Wu, F., Yasin, S., Moskovljevic, M., Varriale, J., Dragoni, F., Camilo-Contreras, A., Duan, J., Zheng, M.Y., Tadzong, N.F., et al. (2023). Clonally expanded HIV-1 proviruses with 5'-leader defects can give rise to nonsuppressible residual viremia. J Clin Invest *133*. https://doi.org/10.1172/JCI165245.

18. Ferdin, J., Goričar, K., Dolžan, V., Plemenitaš, A., Martin, J.N., Peterlin, B.M., Deeks, S.G., and Lenassi, M. (2018). Viral protein Nef is detected in plasma of half of HIV-infected adults with undetectable plasma HIV RNA. PLoS One *13*, e0191613. https://doi.org/10.1371/journal.pone.0191613.

19. Passaes, C., Delagreverie, H.M., Avettand-Fenoel, V., David, A., Monceaux, V., Essat, A., Müller-Trutwin, M., Duffy, D., De Castro, N., Wittkop, L., et al. (2021). Ultrasensitive Detection of p24 in Plasma Samples from People with Primary and Chronic HIV-1 Infection. J Virol *95*, e0001621. https://doi.org/10.1128/JVI.00016-21.

20. Wu, G., Zuck, P., Goh, S.L., Milush, J.M., Vohra, P., Wong, J.K., Somsouk, M., Yukl, S.A., Shacklett, B.L., Chomont, N., et al. (2021). Gag p24 Is a Marker of Human Immunodeficiency Virus Expression in

Tissues and Correlates With Immune Response. J Infect Dis *224*, 1593–1598. https://doi.org/10.1093/infdis/jiab121.

21. Jordan, A., Defechereux, P., and Verdin, E. (2001). The site of HIV-1 integration in the human genome determines basal transcriptional activity and response to Tat transactivation. EMBO J *20*, 1726–1738. https://doi.org/10.1093/emboj/20.7.1726.

22. Jordan, A., Bisgrove, D., and Verdin, E. (2003). HIV reproducibly establishes a latent infection after acute infection of T cells in vitro. EMBO J *22*, 1868–1877. https://doi.org/10.1093/emboj/cdg188.

23. Chen, H.-C., Martinez, J.P., Zorita, E., Meyerhans, A., and Filion, G.J. (2017). Position effects influence HIV latency reversal. Nat Struct Mol Biol *24*, 47–54. https://doi.org/10.1038/nsmb.3328.

24. Battivelli, E., Dahabieh, M.S., Abdel-Mohsen, M., Svensson, J.P., Tojal Da Silva, I., Cohn, L.B., Gramatica, A., Deeks, S., Greene, W.C., Pillai, S.K., et al. (2018). Distinct chromatin functional states correlate with HIV latency reactivation in infected primary CD4 T cells. Elife *7*. https://doi.org/10.7554/eLife.34655.

25. Miklík, D., Šenigl, F., and Hejnar, J. (2018). Proviruses with Long-Term Stable Expression Accumulate in Transcriptionally Active Chromatin Close to the Gene Regulatory Elements: Comparison of ASLV-, HIV- and MLV-Derived Vectors. Viruses *10*. https://doi.org/10.3390/v10030116.

26. Vansant, G., Chen, H.-C., Zorita, E., Trejbalová, K., Miklík, D., Filion, G., and Debyser, Z. (2020). The chromatin landscape at the HIV-1 provirus integration site determines viral expression. Nucleic Acids Res *48*, 7801–7817. https://doi.org/10.1093/nar/gkaa536.

27. Nabel, G., and Baltimore, D. (1987). An inducible transcription factor activates expression of human immunodeficiency virus in T cells. Nature *326*, 711–713. https://doi.org/10.1038/326711a0.

28. Kinoshita, S., Su, L., Amano, M., Timmerman, L.A., Kaneshima, H., and Nolan, G.P. (1997). The T cell activation factor NF-ATc positively regulates HIV-1 replication and gene expression in T cells. Immunity *6*, 235–244. https://doi.org/10.1016/s1074-7613(00)80326-x.

29. Perkins, N.D., Felzien, L.K., Betts, J.C., Leung, K., Beach, D.H., and Nabel, G.J. (1997). Regulation of NF-kappaB by cyclin-dependent kinases associated with the p300 coactivator. Science *275*, 523–527. https://doi.org/10.1126/science.275.5299.523.

30. Ne, E., Palstra, R.-J., and Mahmoudi, T. (2018). Transcription: Insights From the HIV-1 Promoter. Int. Rev. Cell Mol. Biol. *335*, 191–243. https://doi.org/10.1016/bs.ircmb.2017.07.011.

31. Verdin, E., Paras, P., Jr, and Van Lint, C. (1993). Chromatin disruption in the promoter of human immunodeficiency virus type 1 during transcriptional activation. EMBO J. *12*, 3249–3259. https://doi.org/10.1002/j.1460-2075.1993.tb05994.x.

32. Mahmoudi, T. (2012). The BAF complex and HIV latency. Transcription *3*, 171–176. https://doi.org/10.4161/trns.20541.

33. Austin, D.W., Allen, M.S., McCollum, J.M., Dar, R.D., Wilgus, J.R., Sayler, G.S., Samatova, N.F., Cox, C.D., and Simpson, M.L. (2006). Gene network shaping of inherent noise spectra. Nature *439*, 608–611. https://doi.org/10.1038/nature04194.

34. Razooky, B.S., Pai, A., Aull, K., Rouzine, I.M., and Weinberger, L.S. (2015). A hardwired HIV latency program. Cell *160*, 990–1001. https://doi.org/10.1016/j.cell.2015.02.009.

35. Aull, K.H., Tanner, E.J., Thomson, M., and Weinberger, L.S. (2017). Transient Thresholding: A Mechanism Enabling Noncooperative Transcriptional Circuitry to Form a Switch. Biophys. J. *112*, 2428–2438. https://doi.org/10.1016/j.bpj.2017.05.002.

36. Weinberger, L.S., Dar, R.D., and Simpson, M.L. (2008). Transient-mediated fate determination in a transcriptional circuit of HIV. Nat. Genet. *40*, 466–470. https://doi.org/10.1038/ng.116.

37. Whitney, J.B., Hill, A.L., Sanisetty, S., Penaloza-MacMaster, P., Liu, J., Shetty, M., Parenteau, L., Cabral, C., Shields, J., Blackmore, S., et al. (2014). Rapid seeding of the viral reservoir prior to SIV viraemia in rhesus monkeys. Nature *512*, 74–77. https://doi.org/10.1038/nature13594.

38. Colby, D.J., Trautmann, L., Pinyakorn, S., Leyre, L., Pagliuzza, A., Kroon, E., Rolland, M., Takata, H., Buranapraditkun, S., Intasan, J., et al. (2018). Rapid HIV RNA rebound after antiretroviral treatment interruption in persons durably suppressed in Fiebig I acute HIV infection. Nat Med *24*, 923–926. https://doi.org/10.1038/s41591-018-0026-6.

39. Gantner, P., Buranapraditkun, S., Pagliuzza, A., Dufour, C., Pardons, M., Mitchell, J.L., Kroon, E., Sacdalan, C., Tulmethakaan, N., Pinyakorn, S., et al. (2023). HIV rapidly targets a diverse pool of CD4 T cells to establish productive and latent infections. Immunity *56*, 653–668.e5. https://doi.org/10.1016/j.immuni.2023.01.030.

40. Tedbury, P.R., Mahboubi, D., Puray-Chavez, M., Shah, R., Ukah, O.B., Wahoski, C.C., Fadel, H.J., Poeschla, E.M., Gao, X., McFadden, W.M., et al. (2024). Disruption of LEDGF/p75-directed integration derepresses antisense transcription of the HIV-1 genome. bioRxiv. https://doi.org/10.1101/2024.12.06.627169.

41. Singer, M.F., Jones, O.W., and Nirenberg, M.W. (1963). The effect of secondary structure of the template activity of polyribonucleotides. Proc. Natl. Acad. Sci. U. S. A. *49*, 392–399. https://doi.org/10.1073/pnas.49.3.392.

42. Røsok, Ø., and Sioud, M. (2004). Systematic identification of sense-antisense transcripts in mammalian cells. Nat. Biotechnol. *22*, 104–108. https://doi.org/10.1038/nbt925.

43. Miller, D. (1988). HIV and social psychiatry. Br. Med. Bull. *44*, 130–148. https://doi.org/10.1093/oxfordjournals.bmb.a072236.

44. Tagieva, N.E., and Vaquero, C. (1997). Expression of naturally occurring antisense RNA inhibits human immunodeficiency virus type 1 heterologous strain replication. J. Gen. Virol. *78 (Pt 10)*, 2503–2511. https://doi.org/10.1099/0022-1317-78-10-2503.

45. Kobayashi-Ishihara, M., Yamagishi, M., Hara, T., Matsuda, Y., Takahashi, R., Miyake, A., Nakano, K., Yamochi, T., Ishida, T., and Watanabe, T. (2012). HIV-1-encoded antisense RNA suppresses viral replication for a prolonged period. Retrovirology *9*, 38. https://doi.org/10.1186/1742-4690-9-38.

46. Landry, S., Halin, M., Lefort, S., Audet, B., Vaquero, C., Mesnard, J.-M., and Barbeau, B. (2007). Detection, characterization and regulation of antisense transcripts in HIV-1. Retrovirology *4*, 71. https://doi.org/10.1186/1742-4690-4-71.

47. Laverdure, S., Gross, A., Arpin-André, C., Clerc, I., Beaumelle, B., Barbeau, B., and Mesnard, J.-M. (2012). HIV-1 antisense transcription is preferentially activated in primary monocyte-derived cells. J Virol *86*, 13785–13789. https://doi.org/10.1128/JVI.01723-12.

48. Barbeau, B., and Mesnard, J.-M. (2015). Does chronic infection in retroviruses have a sense? Trends Microbiol. *23*, 367–375. https://doi.org/10.1016/j.tim.2015.01.009.

49. Saayman, S., Ackley, A., Turner, A.-M.W., Famiglietti, M., Bosque, A., Clemson, M., Planelles, V., and Morris, K.V. (2014). An HIV-encoded antisense long noncoding RNA epigenetically regulates viral transcription. Mol Ther *22*, 1164–1175. https://doi.org/10.1038/mt.2014.29.

50. Zapata, J.C., Campilongo, F., Barclay, R.A., DeMarino, C., Iglesias-Ussel, M.D., Kashanchi, F., and Romerio, F. (2017). The Human Immunodeficiency Virus 1 ASP RNA promotes viral latency by recruiting the Polycomb Repressor Complex 2 and promoting nucleosome assembly. Virology *506*, 34–44. https://doi.org/10.1016/j.virol.2017.03.002.

51. Kobayashi-Ishihara, M., Terahara, K., Martinez, J.P., Yamagishi, M., Iwabuchi, R., Brander, C., Ato, M., Watanabe, T., Meyerhans, A., and Tsunetsugu-Yokota, Y. (2018). HIV LTR-Driven Antisense RNA by Itself Has Regulatory Function and May Curtail Virus Reactivation From Latency. Front. Microbiol. *9*, 1066. https://doi.org/10.3389/fmicb.2018.01066.

52. Cassan, E., Arigon-Chifolleau, A.-M., Mesnard, J.-M., Gross, A., and Gascuel, O. (2016). Concomitant emergence of the antisense protein gene of HIV-1 and of the pandemic. Proc Natl Acad Sci U S A *113*, 11537–11542. https://doi.org/10.1073/pnas.1605739113.

53. Li, R., Daneshvar, K., Ji, X., Pleet, M., Igbinosun, G., Iqbal, M.S., Kashanchi, F., Mullen, A., and Romerio, F. (2024). Epigenetic silencing and blockade of latency reversal by an HIV-1 encoded antisense transcript. bioRxiv. https://doi.org/10.1101/2024.12.06.627241.

54. Bar-Even, A., Paulsson, J., Maheshri, N., Carmi, M., O'Shea, E., Pilpel, Y., and Barkai, N. (2006). Noise in protein expression scales with natural protein abundance. Nat Genet *38*, 636–643. https://doi.org/10.1038/ng1807.

55. Cox, C.D., McCollum, J.M., Allen, M.S., Dar, R.D., and Simpson, M.L. (2008). Using noise to probe and characterize gene circuits. Proc Natl Acad Sci U S A *105*, 10809–10814. https://doi.org/10.1073/pnas.0804829105.

56. Mancarella, A., Procopio, F.A., Achsel, T., De Crignis, E., Foley, B.T., Corradin, G., Bagni, C., Pantaleo, G., and Graziosi, C. (2019). Detection of antisense protein (ASP) RNA transcripts in individuals infected with human immunodeficiency virus type 1 (HIV-1). J Gen Virol *100*, 863–876. https://doi.org/10.1099/jgv.0.001244.

57. Chen, H.-C., Zorita, E., and Filion, G.J. (2018). Using Barcoded HIV Ensembles (B-HIVE) for Single Provirus Transcriptomics. Curr Protoc Mol Biol *122*, e56. https://doi.org/10.1002/cpmb.56.

58. Weinberger, L.S., and Shenk, T. (2007). An HIV feedback resistor: auto-regulatory circuit deactivator and noise buffer. PLoS Biol *5*, e9. https://doi.org/10.1371/journal.pbio.0050009.

59. Tantale, K., Garcia-Oliver, E., Robert, M.-C., L'Hostis, A., Yang, Y., Tsanov, N., Topno, R., Gostan, T., Kozulic-Pirher, A., Basu-Shrivastava, M., et al. (2021). Stochastic pausing at latent HIV-1 promoters generates transcriptional bursting. Nat Commun *12*, 4503. https://doi.org/10.1038/s41467-021-24462-5.

60. Dey, S.S., Foley, J.E., Limsirichai, P., Schaffer, D.V., and Arkin, A.P. (2015). Orthogonal control of expression mean and variance by epigenetic features at different genomic loci. Mol Syst Biol *11*, 806. https://doi.org/10.15252/msb.20145704.

61. Wang, G.P., Ciuffi, A., Leipzig, J., Berry, C.C., and Bushman, F.D. (2007). HIV integration site selection: analysis by massively parallel pyrosequencing reveals association with epigenetic modifications. Genome Res *17*, 1186–1194. https://doi.org/10.1101/gr.6286907.

62. Schröder, A.R.W., Shinn, P., Chen, H., Berry, C., Ecker, J.R., and Bushman, F. (2002). HIV-1 integration in the human genome favors active genes and local hotspots. Cell *110*, 521–529. https://doi.org/10.1016/s0092-8674(02)00864-4.

63. Einkauf, K.B., Lee, G.Q., Gao, C., Sharaf, R., Sun, X., Hua, S., Chen, S.M., Jiang, C., Lian, X.,

Chowdhury, F.Z., et al. (2019). Intact HIV-1 proviruses accumulate at distinct chromosomal positions during prolonged antiretroviral therapy. J Clin Invest *129*, 988–998. https://doi.org/10.1172/JCI124291.

64. Jiang, C., Lian, X., Gao, C., Sun, X., Einkauf, K.B., Chevalier, J.M., Chen, S.M.Y., Hua, S., Rhee, B., Chang, K., et al. (2020). Distinct viral reservoirs in individuals with spontaneous control of HIV-1. Nature *585*, 261–267. https://doi.org/10.1038/s41586-020-2651-8.

65. Lian, X., Gao, C., Sun, X., Jiang, C., Einkauf, K.B., Seiger, K.W., Chevalier, J.M., Yuki, Y., Martin, M., Hoh, R., et al. (2021). Signatures of immune selection in intact and defective proviruses distinguish HIV-1 elite controllers. Sci Transl Med *13*, eabl4097. https://doi.org/10.1126/scitranslmed.abl4097.

66. Einkauf, K.B., Osborn, M.R., Gao, C., Sun, W., Sun, X., Lian, X., Parsons, E.M., Gladkov, G.T., Seiger, K.W., Blackmer, J.E., et al. (2022). Parallel analysis of transcription, integration, and sequence of single HIV-1 proviruses. Cell *185*, 266–282.e15. https://doi.org/10.1016/j.cell.2021.12.011.

67. Lian, X., Seiger, K.W., Parsons, E.M., Gao, C., Sun, W., Gladkov, G.T., Roseto, I.C., Einkauf, K.B., Osborn, M.R., Chevalier, J.M., et al. (2023). Progressive transformation of the HIV-1 reservoir cell profile over two decades of antiviral therapy. Cell Host Microbe *31*, 83–96.e5. https://doi.org/10.1016/j.chom.2022.12.002.

68. Machida, S., Depierre, D., Chen, H.-C., Thenin-Houssier, S., Petitjean, G., Doyen, C.M., Takaku, M., Cuvier, O., and Benkirane, M. (2020). Exploring histone loading on HIV DNA reveals a dynamic nucleosome positioning between unintegrated and integrated viral genome. Proc Natl Acad Sci U S A *117*, 6822–6830. https://doi.org/10.1073/pnas.1913754117.

69. Thenin-Houssier, S., Machida, S., Jahan, C., Bonnet-Madin, L., Abbou, S., Chen, H.-C., Tesfaye, R., Cuvier, O., and Benkirane, M. (2023). POLE3 is a repressor of unintegrated HIV-1 DNA required for efficient virus integration and escape from innate immune sensing. Sci Adv *9*, eadh3642. https://doi.org/10.1126/sciadv.adh3642.

70. Rando, O.J., and Chang, H.Y. (2009). Genome-wide views of chromatin structure. Annu Rev Biochem *78*, 245–271. https://doi.org/10.1146/annurev.biochem.78.071107.134639.

71. Kaufman, P.D., and Rando, O.J. (2010). Chromatin as a potential carrier of heritable information. Curr Opin Cell Biol *22*, 284–290. https://doi.org/10.1016/j.ceb.2010.02.002.

72. Lekstrom-Himes, J.A., Wang, K., Pesnicak, L., Krause, P.R., and Straus, S.E. (1998). The comparative biology of latent herpes simplex virus type 1 and type 2 infections: latency-associated transcript promoter activity and expression in vitro and in infected mice. J Neurovirol *4*, 27–37. https://doi.org/10.3109/13550289809113479.

73. Kondo, K., Xu, J., and Mocarski, E.S. (1996). Human cytomegalovirus latent gene expression in granulocyte-macrophage progenitors in culture and in seropositive individuals. Proc Natl Acad Sci U S A *93*, 11137–11142. https://doi.org/10.1073/pnas.93.20.11137.

74. Bangham, C.R.M., Miura, M., Kulkarni, A., and Matsuoka, M. (2019). Regulation of Latency in the Human T Cell Leukemia Virus, HTLV-1. Annu Rev Virol *6*, 365–385. https://doi.org/10.1146/annurev-virology-092818-015501.

75. Capoferri, A.A., Sklutuis, R., Famuyiwa, T.O., Pathak, S., Li, R., Rausch, J.W., Luke, B.T., Hoh, R., Deeks, S.G., Mellors, J.W., et al. (2024). *In vivo*detection of HIV-1 antisense transcripts in untreated and ART-treated individuals. bioRxiv. https://doi.org/10.1101/2024.12.06.627170.

76. Klaver, B., and Berkhout, B. (1994). Comparison of 5' and 3' long terminal repeat promoter function in human immunodeficiency virus. J Virol *68*, 3830–3840. https://doi.org/10.1128/JVI.68.6.3830-3840.1994.

77. Lefebvre, G., Desfarges, S., Uyttebroeck, F., Muñoz, M., Beerenwinkel, N., Rougemont, J., Telenti, A., and Ciuffi, A. (2011). Analysis of HIV-1 expression level and sense of transcription by high-throughput sequencing of the infected cell. J Virol *85*, 6205–6211. https://doi.org/10.1128/JVI.00252-11.

78. Archin, N.M., Liberty, A.L., Kashuba, A.D., Choudhary, S.K., Kuruc, J.D., Crooks, A.M., Parker, D.C., Anderson, E.M., Kearney, M.F., Strain, M.C., et al. (2012). Administration of vorinostat disrupts HIV-1 latency in patients on antiretroviral therapy. Nature *487*, 482–485. https://doi.org/10.1038/nature11286.

79. Deeks, S.G. (2012). HIV: Shock and kill. Nature *487*, 439–440. https://doi.org/10.1038/487439a.

80. Abner, E., and Jordan, A. (2019). HIV "shock and kill" therapy: In need of revision. Antiviral Res *166*, 19–34. https://doi.org/10.1016/j.antiviral.2019.03.008.

81. Ho, Y.-C., Shan, L., Hosmane, N.N., Wang, J., Laskey, S.B., Rosenbloom, D.I.S., Lai, J., Blankson, J.N., Siliciano, J.D., and Siliciano, R.F. (2013). Replication-competent noninduced proviruses in the latent reservoir increase barrier to HIV-1 cure. Cell *155*, 540–551. https://doi.org/10.1016/j.cell.2013.09.020.

82. Weinberger, A.D., and Weinberger, L.S. (2013). Stochastic fate selection in HIV-infected patients. Cell *155*, 497–499. https://doi.org/10.1016/j.cell.2013.09.039.

83. Dar, R.D., Hosmane, N.N., Arkin, M.R., Siliciano, R.F., and Weinberger, L.S. (2014). Screening for noise in gene expression identifies drug synergies. Science *344*, 1392–1396. https://doi.org/10.1126/science.1250220.

84. ENCODE Project Consortium, Moore, J.E., Purcaro, M.J., Pratt, H.E., Epstein, C.B., Shoresh, N., Adrian, J., Kawli, T., Davis, C.A., Dobin, A., et al. (2020). Expanded encyclopaedias of DNA elements in the human and mouse genomes. Nature *583*, 699–710. https://doi.org/10.1038/s41586-020-2493-4.

85. Lê, S., Josse, J., and Husson, F. (2008). FactoMineR: AnRPackage for Multivariate Analysis. J. Stat. Softw. *25*. https://doi.org/10.18637/jss.v025.i01.

86. Csárdi, G., Nepusz, T., Müller, K., Horvát, S., Traag, V., Zanini, F., and Noom, D. (2025). igraph for R: R interface of the igraph library for graph theory and network analysis (Zenodo) https://doi.org/10.5281/ZENODO.14736815.

87. Virtanen, P., Gommers, R., Oliphant, T.E., Haberland, M., Reddy, T., Cournapeau, D., Burovski, E., Peterson, P., Weckesser, W., Bright, J., et al. (2020). SciPy 1.0: fundamental algorithms for scientific computing in Python. Nat Methods *17*, 261–272. https://doi.org/10.1038/s41592-019-0686-2.

88. The pandas development team (2024). pandas-dev/pandas: Pandas (Zenodo) https://doi.org/10.5281/ZENODO.3509134.

89. Cock, P.J.A., Antao, T., Chang, J.T., Chapman, B.A., Cox, C.J., Dalke, A., Friedberg, I., Hamelryck, T., Kauff, F., Wilczynski, B., et al. (2009). Biopython: freely available Python tools for computational molecular biology and bioinformatics. Bioinformatics *25*, 1422–1423. https://doi.org/10.1093/bioinformatics/btp163.

90. Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., Durbin, R., and 1000 Genome Project Data Processing Subgroup (2009). The Sequence Alignment/Map format and SAMtools. Bioinformatics *25*, 2078–2079. https://doi.org/10.1093/bioinformatics/btp352.

91. Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2. Nat Methods *9*, 357–359. https://doi.org/10.1038/nmeth.1923.

92. The R Project for Statistical Computing https://www.R-project.org/.

93. Kassambara, A., and Mundt, F. (2016). Factoextra: Extract and visualize the results of multivariate data analyses. (The R Foundation). https://doi.org/10.32614/cran.package.factoextra.

94. Robinson, J.T., Thorvaldsdottir, H., Turner, D., and Mesirov, J.P. (2023). igv.js: an embeddable JavaScript implementation of the Integrative Genomics Viewer (IGV). Bioinformatics *39*. https://doi.org/10.1093/bioinformatics/btac830.