Population-Level Activity Dissociates Preparatory Overt from Covert Attention

Damian Koevoet[1,*], Vicky Voet[1], Henry M. Jones[2], Edward Awh[2], Christoph Strauch[1,+], and Stefan Van der Stigchel[1,+]

[1]Experimental Psychology, Helmholtz Institute, Utrecht University

[2]Department of Psychology, Institute of Mind and Biology, The University of Chicago

[+]Shared authorship

[*]Correspondence to d.koevoet@uu.nl

Abstract

The neural signatures of preparing overt eye movements and directing covert spatial attention overlap as they recruit the same brain areas. Yet, these neural signatures are dissociable at the single-cell level: Specific cells within these visuo-oculomotor areas are exclusively involved in either motor preparation or covert attention. Nevertheless, it has been proposed that many cells in visuo-oculomotor areas are involved in both motor preparation and covert attention, and consequently their neural signatures should functionally overlap to a large degree. Here, we put this proposal to the test: we combined EEG with sensitive decoding techniques to investigate whether the neural signatures of preparatory overt and covert attention are dissociable across large-scale neuronal populations. We found that neural decoding reliably discerned whether overt or covert attention was shifted well before saccade initiation. Further, inverted encoding modeling revealed sharper spatially tuned activity in preparatory overt than in covert attention. We then asked whether preparatory overt attention achieved sharper spatially-tuned activity by using 'more-of-the-same' covert attention, or by recruiting additional spatially selective neural processing. Cross-decoding results demonstrated that preparatory overt attention recruited at least one additional, frontal process. This additional spatially selective process emerged early and likely reflects motor preparation or predictive remapping. To summarize, we found that the neural signatures of overt and covert attention overlap, yet diverge rapidly, in part because overt attention employs an additional spatially selective neural process. Extending beyond a dissociation on the single-cell level, our findings demonstrate that population-level neural activity dissociates preparatory overt from covert attention.

Keywords: Attention, Eye movements, Neural decoding, Population, Spatial selectivity

Introduction

Spatial attention is shifted overtly with, or covertly without an accompanying eye movement [1–5]. Preparatory overt and covert attention are tightly coupled [6, 7]. For example, saccade preparation leads to an attentional shift to the target prior to eye movement initiation [8–10] and saccade trajectories deviate from attended locations [11]. While such behavioral results sparked tremendous interest in the neural signatures of preparatory overt and covert attention, it currently remains unclear to what degree these signatures overlap [4, 6, 7, 10, 12–14].

On the one hand, preparatory overt and covert attention neurally overlap to a large degree. For example, they recruit the same areas including frontal eye fields (FEF) and superior colliculi [4, 15–21]. Moreover, subthreshold microstimulation of FEF cells involved in saccade generation (which would trigger saccades if stimulated above threshold) leads to covert attentional selection [22–24]. This demonstrates a tight neural link between preparatory overt and covert attention [although it is possible that neighboring cells were also stimulated, see 25]. On the other hand, it has become increasingly clear that preparatory overt and covert attention are dissociable at the single-cell level: Specific cells code exclusively for motor preparation or covert attention [26–30]. Reconciling these data, Awh *et al.* [4] proposed that a large proportion of cells in visuo-oculomotor areas are sensitive to both motor preparation and covert attention, and that therefore preparatory overt and covert attention overlap to a large degree - even despite dissociations at the single-cell level [also see 13, 31].

If the neural signatures of preparatory overt and covert attention indeed overlap to a large degree [4, 13, 31], dissociations should occur at the single-cell, but not the population level. By contrast, extant work implies that population-level neural activity may dissociate preparatory overt from covert attention [32–35]. Recently, transcranial magnetic stimulation (TMS) - which affects a large population of neurons - of FEF had differential effects on perception during preparatory overt and covert attentional selection. Specifically, TMS applied to FEF ameliorated perceptual benefits

at validly cued and costs at invalidly cued locations in voluntary covert attention [34]. However, TMS at FEF reduced costs at invalidly cued locations but did not mediate benefits at validly cued locations in preparatory overt attention [33]. Another example is that pupil size - indicative of activity in attentional networks [36, 37] - is larger during preparatory overt compared with covert attentional shifts, which was attributed to the recruitment of additional processes such as motor preparation or remapping [32]. Despite these tentative findings indicating larger-scale neural dissociations between preparatory overt and covert attention, direct evidence for a neural dissociation on the population-level is currently lacking.

We here combined EEG - which reflects synchronous activity of large-scale neuronal populations [38, 39] - with sensitive decoding techniques [40–47] to investigate whether preparatory overt and covert attention are dissociable at the population level. To preface our results, we could decode whether preparatory overt or covert attention was shifted already ∼700 ms prior to saccade onset, indicating an early population-level dissociation between shift types. Further analyses demonstrated sharper spatially selective population-level activity in preparatory overt than covert attention. Cross-decoding demonstrated that this sharper spatial selectivity was achieved by recruiting an additional spatially selective process in frontal cortex, which likely reflects motor preparation or predictive remapping [19, 28, 31–33, 48–57]. Our results extend the dissociation between preparatory overt and covert attention from the single-cell level to activity in large-scale neuronal populations. This demonstrates that preparatory overt and covert attention are neurally distinct at a much broader scale than previously thought.

Results

Behavioral and Pupillometry Results

Twenty-three participants performed a cued visual selection task (Figure 1A, [45]) wherein they shifted overt or covert attention while scalp EEG was recorded. In each

trial, a cue (100% valid) indicated where the target diamond would later appear. Participants reported whether the gap was on the left or right side of the target diamond.

The task required participants to attend overtly or covertly in alternating blocks. In the overt condition, participants were instructed to saccade toward the cued location upon array onset. In the covert condition, participants were asked to maintain central fixation throughout the trial. In Figure 1B, we present five example gaze traces per condition from a single participant for illustration.

We analyzed accuracy and response time as behavioral outcomes. Accuracy was generally high (Figure 1C). Participants were more accurate in the overt compared with the covert condition ($t(22) = 3.11$, $p = .005$, $d = 0.65$). Response times did not differ significantly between conditions (Figure 1D, $t(22) = 1.48$, $p = .153$, $d = 0.31$). These data indicate that participants were well able to perform the task, and that participants were more accurate in overt compared with covert trials.

We recently demonstrated that preparatory overt attention causes more pupil dilation than shifting covert attention [32]. We here compared pupil size prior to saccade onset (-200 - 100 ms around array onset) [32, 58] between shift types while controlling for known covariates of pupil size (e.g., task difficulty [36, 59–64] and shift direction [32, 58]; see Methods). In line with our previous work, we observed that preparatory overt attention caused a larger pupil size prior to saccade initiation compared with covert attention (Figure 1E; one-sample t-test: $t(22) = 2.98$, $p = .007$, $d = .62$). This is a first sign that population-level activity [36, 37] may differ between shift types.

Preparatory Overt and Covert Attention Diverge Rapidly

Our main goal was to investigate whether activity in large-scale neuronal populations (as reflected in EEG [38, 39]) discerned between preparatory overt and covert attention. Our neural analyses focused on the period preceding array onset because participants did not (yet) initiate a saccade toward the target location in either con-

Figure 1: Task and behavioral results. **A** Participants performed a cued visual selection task while shifting overt or covert attention. Participants indicated whether a gap was missing from the left ('z') or right ('/') part of the target diamond. Cues (100% valid) indicated the target location. **B** Five example gaze traces from a single participant for each condition upon array onset. As is evident from the figure, participants made a saccade toward the target location upon array onset in overt blocks. In covert blocks, participants maintained central fixation throughout the trial. **C** Gap identification accuracy per condition. **D** Response times per condition. **E** Pupil size difference between shift type conditions while controlling for known covariates of pupil size. Higher values indicate a larger pupil size in the overt compared with the covert attention condition. Error bars reflect standard errors of the mean across participants. Black points indicate individual participants. **$p < .01$.

dition (as in the pupillometry analysis). This allowed for a clean comparison of neural activity between shift types without ocular artifacts. In our first neural analysis, we used time-resolved multivariate classification to decode whether preparatory overt or covert spatial attention was shifted.

We could decode whether participants shifted preparatory overt or covert attention early in the trial. Specifically, we observed significant decoding already from 70 ms after cue onset, which remained reliable throughout the delay period (70-700 ms; $p < .0001$; Figure 2A). This dissociation preceded saccade initiation by at least 630 ms. Therefore, neural processes unfolding well before saccade initiation readily dissociated between preparatory overt and covert attention. Decoding was highly robust: We averaged classification accuracy across the delay period (100-700 ms after cue onset; Figure 2B), and observed above empirical chance classification for

Figure 2: Multivariate classification results. **A** Time-resolved classifiers were trained to identify trials as overt or covert. The purple line represents classification accuracy, while the gray line depicts accuracy on shuffled trial labels (i.e. empirical chance). Shaded error bands reflect standard errors of the mean across participants. Horizontal black line indicates a significant cluster where classifier accuracy was higher than empirical chance ($p < .0001$). **B** The averaged classification accuracy (100-700 ms after cue onset) difference between the actual and shuffled data. Positive values indicate better decoding performance in the non-shuffled compared with the shuffled data. Black points represent individual participants. Error bar holds standard error of the mean.

each participant (one-sample t-test: $t(22) = 10.06$, $p < .0001$, $d = 2.10$). Control analyses demonstrated that none of the decoding results throughout the paper can be ascribed to (premature or miniature) eye movements (Supplementary Material). These results demonstrate that activity in large-scale neuronal populations discerned between preparatory overt and covert attention rapidly after cue onset, and well before saccade initiation.

Sharper Spatial Tuning in Preparatory Overt than in Covert Attention

The above analysis established that preparatory overt and covert attention diverge rapidly after cue onset. But *how* do these shift types differ? Likely candidates that contribute to differences between preparatory overt and covert attention include motor preparation and predictive remapping [10, 32, 65, 66], which are inherently spatial processes. We therefore focused on *how* preparatory overt and covert attention may differ in terms of spatially selective neural activity. To this end, we employed inverted encoding modeling [IEM; 67, 68], which allows for the spatiotemporal tracking of attention [40, 44]. Specifically, we employed IEMs to track whether spatial selectivity of large-scale neuronal populations differed between preparatory overt and covert attention (Figure 3).

In line with previous work [e.g. 40, 44], our results revealed reliable spatial selectivity of neural activity when shifting *covert* attention. Spatial selectivity emerged 164 ms after cue onset and remained throughout the delay ($p < .0001$; Figure 3A,B,C). We here extend this by demonstrating that preparatory *overt* attention also caused spatially-tuned neural activity (117-700 ms, $p < .0001$; Figure 3A,B,C) [also see 69]. Thus, IEMs precisely tracked spatial attention across time in both conditions.

We then tested whether the spatial selectivity of neural activity differed between shift types. We observed sharper spatially-tuned neural activity in overt compared with covert trials (Figure 3A,B,C). Specifically, we found two early (cluster 1: 144-273 ms, $p = .002$, cluster 2: 289-323 ms, $p = .015$) and a later significant cluster in the delay period (cluster 3: 516-700 ms, $p < .001$). The preparation of an imminent

Figure 3: Inverted encoding modeling results. **A** CTF slopes across time for the overt (blue) and covert (red) conditions. The reconstructed CTF's slope is a measure of spatial selectivity, where higher values represent stronger spatial selectivity. Shaded error bands reflect standard errors of the mean across participants. Blue and red horizontal lines indicate significant CTF slope clusters for overt and covert data, respectively ($p < .0001$). Horizontal black lines indicate significant clusters between conditions ($p < .05$). **B** Averaged (100-700 ms after cue onset) reconstructed CTFs for the overt (blue) and covert (red) conditions. For both tuning curves, responses are strongest for the cued location (i.e. channel offset of 0) and drop off at further distances. Error bars reflect standard errors of the mean across participants. **C** Heat maps of channel responses across time per shift type. **D** Searchlight-based visualizations of channel contributions to CTF slopes across the scalp. Difference topography was calculated by subtracting the covert from the overt topography; positive values thus correspond to higher slopes in the overt condition (and vice versa).

saccade caused sharper spatially selective neural activity compared with directing covert attention already early after cue onset, and thus well before saccade initiation (>550 ms). These results indicate that preparatory overt and covert attentional shifts diverge early, in part due to sharper spatial tuning of large-scale neural activity to a cued location.

We visualized topographies of spatial selectivity in the preparatory overt and covert conditions using a searchlight approach [Figure 3D; 70]. To this end, we iteratively selected groups of 5 neighboring electrodes and repeated the IEM procedure on each group, and averaged CTF slopes across the delay period (100-700 ms after cue onset). This allowed us to obtain the contribution of each electrode to the observed spatial selectivity (i.e. CTF slope) per condition but collapsed across cued locations. As expected, we observed considerable contributions to spatial selectivity in posterior (occipital and parietal) sites for both shift types [43]. In addition, frontal channels contributed more to spatial selectivity during preparatory overt compared with covert attentional shifts. Together, the topographies reveal overlapping spatial selectivity in posterior areas across shift types, and a possible spatial selectivity in frontal areas that may only be present during preparatory overt but not covert attention.

Preparatory Overt Attention Recruits an Additional Spatially Selective Neural Process

Preparatory overt attention was accompanied by sharper spatially-tuned neural activity tracking the cued location compared with covert attention. Pushing this further, we asked whether this difference was due to overt attention recruiting 'more-of-the-same' covert attention, or because overt attention recruits (at least) one additional spatially selective neural process (Figure 4). We previously hypothesized that the larger pupil size in preparatory overt attention (Figure 1E) may be attributed to the recruitment of additional necessary neural processes such as motor preparation and remapping [32].

Figure 4: Hypothetical and actual inverted encoding modeling (IEM) results when training on only covert or overt trials. For each graph, we visualized CTF slopes across time for the overt (blue) and covert (red) conditions when training only on covert (left) and overt (right) data. **A** Hypothetical IEM results from a 'more-of-the-same' account. **B** Hypothetical IEM results from an 'additional source' account. **C** Actual IEM results when training on only covert or overt trials. Shaded error bands reflect standard errors of the mean across participants. Blue and red horizontal lines indicate significant CTF slope clusters for overt and covert data, respectively ($p < .0001$). Horizontal black lines indicate significant clusters between conditions ($p < .05$).

We reasoned that a 'more-of-the-same' account would predict that preparatory overt attention makes stronger use of the neural signals present when shifting covert attention (Figure 4A). In contrast, if overt attention employs an additional spatially selective neural process, this process should be absent when shifting covert attention (Figure 4B). To arbitrate between these accounts, we now trained IEMs on only one shift type but tested them on both conditions (i.e. cross-decoding).

When exclusively training on covert data, a 'more-of-the-same' account would predict a difference between shift types in terms of neural spatial selectivity (Figure 4A). In contrast, when training exclusively on covert data, the difference in neural spatial selectivity between overt and covert conditions disappeared (all ps > .16; Figure 4C). We did observe reliable neural spatial selectivity for the cued location in both conditions, demonstrating overlap in spatially-tuned neural activity between conditions (overt: 133-700 ms, p < .0001; covert: 172-700 ms, p < .0001). This argues against a more-of-the-same account and instead favors the interpretation that overt attention employs an additional spatially selective neural process.

If so, when training exclusively on overt data - where the IEM would once again have access to the putative additional spatially selective process - the difference in neural spatial selectivity between conditions should reemerge (Figure 4C). We indeed observed sharper spatial tuning in preparatory overt compared with covert attention when training on overt data exclusively (cluster 1: 164-352 ms, p = .001; cluster 2: 383-438 ms, p = .018; cluster 3: 453-700 ms, p < .001). Again, we also found spatially selective neural activity in both conditions separately, indicating overlap in the employed spatially selective neural processes (overt: 125-700 ms, p < .0001; covert: 117-700 ms, p < .0001).

To formally test whether the higher spatial selectivity in overt compared with covert attention depended on the training data, we calculated the difference in averaged CTF slopes (100-700 ms after cue onset) between shift types when training only on overt or covert data. Comparisons confirmed that the difference between spatial selectivity in overt compared with covert trials is higher when training only on overt compared with training only on covert data (t(22) = 4.11, p < .001, d = 0.86).

Together, these analyses demonstrate preparatory overt attention to recruit (at least) one additional spatially selective neural process that was not present when covert attention was shifted. This additional spatial selectivity was most prominent in frontal channels, and explains the sharper spatial selectivity in preparatory overt than in covert attention. We speculate that motor preparation and/or predictive remapping underlie this additional spatially selective process (see Discussion).

Discussion

We here demonstrated that population-level activity as reflected in EEG [38, 39] dissociates preparatory overt from covert attention. This dissociation emerged rapidly after cue onset, and was in part driven by the recruitment of (at least) one additional spatially selective process during preparatory overt attention. Our results dissociate preparatory overt from covert attention at a much broader scale than previously proposed.

By definition the neural signatures of preparatory overt and covert attention must diverge as a saccade is initiated in the former but not the latter case [4]. But when do these neural signatures diverge: Does this already occur prior to saccade initiation? Complementing evidence from single-cell recordings [26], our decoding results demonstrate that population-level neural activity dissociated rapidly after cue onset. These data resonate with psychophysical data demonstrating differential effects of presaccadic and (voluntary) covert attention on perception rapidly after cue onset [10, 56, 71]. Our results reveal a rapid neural dissociation between preparatory overt and covert attention well before saccade onset.

We previously found a larger pupil size during preparatory overt attention compared with covert attentional shifts [32] - which we replicated here - and hypothesized that this larger pupil size could be attributed to the recruitment of additional neural processes in preparatory overt attention. We here found direct neural evidence for this hypothesis (Figure 4), as cross-decoding revealed an additional spatially selective process in frontal cortex in preparatory overt attention. What could

this additional spatially selective process reflect?

Although our data cannot cannot answer this question with certitude, we here discuss two possible candidates that are inherently spatial and recruit FEF: Motor preparation and predictive remapping [4, 6–10, 12–14, 19–22, 26, 55, 72–74]. Jonikaitis *et al.* [28] demonstrated that pharmacological inactivation (using muscinol) of FEF impaired motor preparation but not spatial memory of the target location [or covert spatial attention, see 75–79]. In line with this, single-unit recordings in FEF revealed enhanced spatial selectivity when a saccade can be prepared to the target compared with solely attending/memorizing the target location [27]. As for the timecourse, the independent contribution of motor preparation to spatial selectivity emerges well before saccadic onset, which closely fits with our population-level results [26, 27] [also see 57]. The above puts forth motor preparation as a strong candidate to underlie the additional spatially selective process we observed in frontal cortex during preparatory overt but not covert attention.

Another candidate that could underlie the additional spatially selectivity is predictive remapping. Predictive remapping entails the remapping of receptive fields from the presaccadic stimulus location to the post-saccadic stimulus location. Such remapping accounts for displacements of the visual world on the retina due to saccades, which facilitates visual continuity [52, 53, 80–88]. Importantly, predictive remapping should therefore only occur in preparatory overt but not covert attention [55, 65, 82]. Although the above argues for a potential role of predictive remapping in explaining our results, the timecourse of predictive remapping may not fit our findings. For example, recent M/EEG decoding studies demonstrated that the post-saccadic stimulus position could be decoded only briefly ($\sim$50-100 ms) preceding saccade onset [89, 90], and similar timecourses are observed using single-unit FEF recordings [e.g. 91]. By contrast, in the current data preparatory overt attention already showed sharper spatial selectivity than covert attention well before saccade initiation (>550 ms). One discrepancy is that participants initiated saccades as fast as possible upon cue onset in previous work [89, 90] but delayed their saccade in our task, making it difficult to compare timecourses between them. Future work

should address whether preparatory overt also shows sharper spatial selectivity than covert attention when there is no delay (i.e. saccades are initiated as fast as possible upon cue onset). Note that we are unsure whether participants even *could* predictively remap array stimuli since they were not presented prior to saccade onset. Nevertheless, we cannot rule out that the fixation point was remapped predictively. Thus, predictive remapping may account for the sharper spatial tuning involved in preparatory overt than covert attention but its timecourse may not be compatible with the current results. We speculate that the additional frontal spatially selective process likely reflects motor preparation as previous work provides more traction for this than predictive remapping but more work is necessary to definitively arbitrate between these candidates.

Where do our results leave motor-based theories of visual attention? For example, the popular premotor theory of attention [6, 7] equates covert attention to preparing but not executing a saccade. Our results oppose a 'hard' interpretation of the premotor theory since we demonstrate dissociable neural activity between preparatory overt and covert attention. Nevertheless, we did observe cross-decodability of cued locations across conditions, demonstrating overlap in attentional mechanisms. One way that a weaker interpretation of the premotor theory may still hold, is if the additional spatially selective process observed here reflects predictive remapping [32, 65]. In such a view, preparatory overt attention is tantamount to covert attention but with an added spatially selective non-motor process. However, if the observed additional spatially selective process reflects motor preparation, our data would show that this weaker version of the premotor theory of attention breaks down even at the population level.

Together, we extend previous dissociations between preparatory overt and covert attention from single cells to the population level. We found that preparatory overt and covert attention dissociated rapidly. Specifically, an additional spatially selective process in frontal cortex was recruited in preparatory overt but not in covert attention. We conclude that population-level activity dissociates preparatory overt from covert attention at a much larger scale than previously thought.

Methods

All data and analysis scripts necessary to reproduce the reported results are available via the Open Science Framework: https://osf.io/kd4qp/.

Participants

We set our target sample to 22 participants based on previous EEG decoding work [e.g. 40, 41, 45, 92, 93]. A total of 28 participants took part in the experiment. We excluded three participants due to poor task performance (<50% accuracy in at least one condition) and two other participants because eye-tracking data were unavailable due to technical errors. Our final sample consisted of 23 participants (M_{age} = 23.22, range: [19-27], 15 female, 8 male) with normal or corrected-to-normal vision and without a diagnosis of, or history with, attention-deficit (hyperactivity-)disorder or autism spectrum disorder. We overshot our target sample size by one participant due to performing artifact rejection and task performance checks in batches. Participants were compensated with course credits or received €8 per hour. The experimental procedure was approved by the Utrecht University ethical review board (approval code: 24-0197).

Data acquisition and stimuli

Gaze data were recorded from the right eye at 1000 Hz using an EyeLink 1000+ (SR Research, Mississauga, Ontario, Canada). Participants were positioned in a chinrest 67.5 cm from the monitor (2560 x 1440; 100 Hz). We calibrated the eye-tracker using a 9-point procedure, and recalibration was performed whenever necessary between blocks. Stimuli were presented using PsychToolbox in MATLAB [94, 95].

EEG was recorded at 2048 Hz from 30 active electrodes using a BioSemi ActiveTwo system (BioSemi, Amsterdam, The Netherlands). Specifically, we recorded from International 10/20 sites: Fp1, AF3, F7, F3, FC1, FC5, C3, CP1, CP5, P7, P3, Pz, PO3, O1, Oz, O2, PO4, P4, P8, CP6, CP2, C4, FC6, FC2, F4, F8, AF4, Fp2, Fz, and Cz. We additionally

placed electrodes ~1 cm from the outer canthi of the left and right eyes to record horizontal EOG, and placed electrodes above and below the right eye to monitor vertical EOG. We also placed electrodes on the left and right mastoid bones that were used for later offline re-referencing.

Stimuli were displayed on a gray background. The central fixation stimulus was a slightly darker grey dot (0.15°). Cues consisted of luminance-matched blue and red (counterbalanced) bars (0.12° x 0.25°) placed around the fixation point. Arrays were presented at 4° eccentricity in an invisible circle, and consisted of a single diamond target (1.6° x 1.6°; with a gap on the left or right) and square distractors (1.6° x 1.6°). All stimuli in the array were equidistant (steps of 45°).

Procedure

Participants performed a cued visual selection task [based on 45] and identified whether the target diamond stimulus had a missing gap on its left or right side (Figure 1A). Trials started with a brief fixation period (500-600 ms). Then a cue (100% valid) was presented that indicated the upcoming target location. The post-cue delay period was variable (600-900 ms) to discourage participants from making premature saccades toward the target stimulus [following 9, 32]. A search array was briefly presented (150 ms) and followed by a mask array (400 ms). Participants then indicated whether the missing gap was on the left ('z' key) or right ('/' key) of the target diamond stimulus. Blank inter-trial intervals ranged between 800 and 1100 ms. Participants were instructed to only blink during the blank inter-trial interval.

Whereas in the covert condition, participants were instructed to maintain central fixation throughout the trial, in the overt condition, participants were instructed to make a saccade as fast as possible to the target location upon array onset (see Figure 1B). Participants were given feedback based on their gaze behavior. If gaze deviated more than 2° from central fixation in the covert condition, participants were shown "Keep still!". In the overt condition, participants received feedback when they made premature (i.e. prior to array onset; "Too early!") or slow saccades (>400 ms;

"Too slow!").

Each block consisted of 64 trials and blocks of overt and covert trials alternated. We counterbalanced across participants whether they would start with the overt or covert condition. Participants completed at least 16 blocks but could choose to complete more until a maximum of 20 blocks was reached. Prior to the experimental blocks, participants completed one block of practice trials per condition to get familiar with the experiment. Participants were free to take breaks between blocks. Participants saw their mean accuracy and response time performance in their last completed block during each break.

Data processing and artifact rejection

Data processing and analyses were performed using custom Python scripts [based on 40–42, 45]. Data were re-referenced to the average of the left and right mastoids, epoched around cue onset (-250 - 700 ms), downsampled to 128 Hz and bandpass filtered (0.5-30 Hz). In the selected epoch, no saccades should be present in the data in both the overt and covert conditions. Therefore, neural activity in this epoch can be compared between conditions without ocular artifacts.

We used an automated artifact detection algorithm based on Foster *et al.* [45]. Eye-tracking data were used to detect trials with ocular artifacts (e.g. saccades or blinks). Gaze data were drift-corrected by subtracting the median gaze position 100-0 ms prior to cue onset for each trial. We took two steps to detect trials with saccades. First, we discarded trials wherein drift-corrected gaze deviated more than 1° from central fixation. Second, we used a 60 ms sliding window of mean gaze that advanced in steps of 10 ms. If at any point in the trial the difference in gaze between the first and last halve of the sliding window exceeded a threshold of 0.5°, we excluded the trial. All trials wherein the pupil size data were missing at any point were discarded to omit trials with blinks.

Next, the algorithm checked the EEG voltage data itself for potential artifacts. Trials with voltage drifts (i.e. skin potentials) were identified by calculating the ab-

solute voltage difference between the first and last quarter of the trial. We discarded trials if the difference exceeded 40 μV. High-frequency noise (i.e. muscle artifacts) was identified by calculating peak-to-peak amplitudes in ~15 ms windows (i.e. two samples at 128 Hz) advanced in ~15 ms steps across the entire trial. Trials were determined to contain high-frequency noise if peak-to-peak amplitudes exceeded 120 μV, and were discarded. We checked for sudden steps in voltage changes by calculating peak-to-peak amplitudes in a window of 250 ms that advanced in steps of ~21 ms. We examined peak-to-peak amplitudes obtained with the sliding window above, and if the amplitude exceeded 60 μV at any point, the trial was discarded. Lastly, we discarded trials if absolute voltage values exceeded 80 μV at any time during the trial [42]. An average of 990.09 trials (*SD* = 127.16) were included per participant in the final analyses.

Data were baseline corrected by subtracting the median voltage within 250 ms prior to cue onset in each trial for each electrode separately. These baseline-corrected data served as the input for further neural analyses.

Multivariate classification analysis

We conducted a multivariate classification analysis to discern between preparatory overt and covert attention [based on 42, 96]. To this end, we trained logistic regression models to classify trials as overt or covert based on baseline-corrected voltage data [46, 47]. Although classification was reliable even at the single trial level, we averaged randomly chosen groups of five trials to increase power [46, 96]. Data were split into training (75%) and test sets (25%), while ensuring an equal number of trials for each location and condition (i.e. cross-validation). Following Thyer *et al.* [42] and Jones *et al.* [41], we standardized the training data (using *sklearn.preprocessing.StandardScaler*) and standardized the test data using the mean and standard deviation from the training data [97]. To evaluate model performance, the trained classifier classified trials as overt or covert in the held-out test set, as well as in a shuffled held-out test set (i.e. shuffled condition labels). This

allows to compare actual classifier accuracy with empirical chance performance. Classification was performed at every time point with 1,000 test-training splits of the data.

Inverted encoding modeling

To track the spatiotemporal dynamics of preparatory overt and covert attention, we employed inverted encoding modeling [IEM, Figure 5; 67, 68]. Although IEMs in combination with alpha oscillations sensitively track the allocation of spatial attention [40, 43–45], alpha oscillations in isolation may not be able to pick up on all relevant processes involved in saccade preparation. For example, results from Günseli *et al.*

Train

Test

Figure 5: Visualization of the inverted encoding model. Data were first split into train and test sets. Basis channel responses were created for each location separately. Activity was assumed to be highest for the attended location, and followed a half sinusoid (raised to the 7th power) to model the spatial selectivity. To obtain weights, we used these basis channel responses for each location and fit them to the training data. Inverting the obtained weights matrix allows for the reconstruction of channel-tuning functions. Specifically, test data were fit to the inverted weights matrix to reconstruct channel-tuning functions for each location separately.

[69] indicate that the spatial selectivity of saccade preparation may not be limited to alpha-band activity. Therefore, we here instead focused on analyzing voltage patterns across the scalp [46, 47] as voltage measures have popularly been employed in studying presaccadic processes [57, 98–102]. For completeness, we report results obtained with alpha-band (8-12 Hz) power in the Supplementary Material.

We reconstructed spatially sensitive channel tuning functions (CTFs) from the voltage topography across time. We assumed that voltage patterns at each electrode reflected the weighted average of eight spatial channels, which were each selective to a specific location. We modeled these basis channel tuning functions using a half sinusoid raised to the 7th power that were centered on all possible target locations [as in 40, 43, 78].

For each iteration, data were split into training (2 blocks) and test (1 block) data while an equal number of trials at each cued location as well as for each shift type was ensured per block. Note that for all IEM analyses, unless explicitly mentioned, we trained and tested on data from both shift type conditions. We then fit the basis channel tuning functions to the training data using a general linear model (using *numpy.linalglstsq*), and we obtained weights at each electrode. We then inverted the weights matrix to reconstruct putative spatially sensitive CTFs separately per condition in the test data. The procedure was conducted at each time point (every ~7.81 ms), and to consider all trials in the analyses we ran 100 iterations per participant over time. We recentered CTFs (i.e. the center corresponds to the cued location) and averaged the reconstructed CTFs across the eight possible locations. This allowed us to track spatial selectivity of neural activity with a high temporal precision.

In keeping with previous work [40, 41, 43, 45], we quantified spatial selectivity of the reconstructed CTF by calculating its slope. More specifically, we first averaged channel responses equidistant to the target location, and then estimated the slope of the reconstructed CTF using linear regression. A higher CTF slope indicates greater spatial selectivity.

To visualize CTF slope topographies, we used a searchlight approach [70]. We

iteratively selected groups of 5 neighbouring electrodes around the scalp and conducted the IEM procedure and calculated the average CTF slope for the time window of 100-700 ms after cue onset (using 25 iterations per electrode group) [similar to 103]. We then averaged CTF slopes across all channel-groups that a specific electrode was included in. This allowed us to obtain specific CTF slope contributions for each electrode separately per condition but collapsed across positions.

Pupillometry

For the pupillometry analysis, we first ran online saccade detection around array onset. We applied a Savitzky-Golay filter [104] and detected saccade onsets using a velocity threshold of 20°/s [32, 105]. For this analysis, we discarded trials with premature, very fast or very slow saccade onsets (<100 ms and >600 ms) in the overt condition, and trials with any detected saccade in the covert condition. (Note that using this subset of trials for the neural decoding analyses yields identical results.) We reconstructed blinks [106], downsampled the data to 100 Hz, linearly interpolated remaining missing data and baseline corrected using the median pupil size 250-0 ms prior to cue onset [107]. Following Koevoet *et al.* [58], we then averaged pupil size prior to saccade initiation (-200 - 100 ms around array onset). We z-scored pupil size and conducted multiple linear regression for each participant separately (Wilkinson notation: z-scored pupil size $\sim$ shift type * y target position + x target position + oblique + correct). This allowed us to control for individual differences in pupil size as well as to control for effects of known covariates of pupil size, such as task difficulty and shift direction [32, 58–61]. We used a two-tailed one-sample *t*-test to test whether the shift type betas reliably differed from 0, effectively testing the difference in pupil size between preparatory overt and covert attention (Figure 1E).

Statistical analysis

To analyze behavioral data, we calculated average accuracies and median response times per condition (in all artifact-free trials). We compared conditions using two-

tailed paired-samples *t*-tests and report Cohen's d_z as a measure of effect size [108].

For the classification analysis, we tested whether classification accuracy was above empirical chance. To this end, we first subtracted decoding accuracies in the shuffled from the non-shuffled data. We then tested whether the difference between these accuracies was reliably above 0. To circumvent the issue of multiple comparisons, we used one-tailed one-sample permutation tests (using *mne.stats.permutation_cluster_1samp_test*) to identify putative clusters and test their reliability [109].

As for spatial selectivity, we tested whether CTF slopes were reliably above 0 (i.e. indicative of spatial selectivity) separately for the overt and covert conditions using one-tailed one-sample permutation tests. To compare the spatial selectivity between overt and covert attention, we subtracted CTF slopes of the covert from the overt condition and used a two-tailed one-sample test to assess significance. All cluster-based tests were conducted with $\alpha = .05$ using 10,000 iterations.

Funding: This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement n° 863732).

Data and code availability: All data and analysis scripts necessary to reproduce the reported results are available via the Open Science Framework: `https://osf.io/kd4qp/`.

Competing interests: The authors have no competing interests to declare.

References

1. Carrasco, M. Visual Attention: The Past 25 Years. *Vision Research* **51,** 1484–1525. ISSN: 0042-6989 (2011).

2. Posner, M. I. Orienting of Attention. *Quarterly Journal of Experimental Psychology* **32,** 3–25. ISSN: 0033-555X (1980).

3. Helmholtz, H. *Concerning the Perceptions in General* (1866).

4. Awh, E., Armstrong, K. M. & Moore, T. Visual and Oculomotor Selection: Links, Causes and Implications for Spatial Attention. *Trends in Cognitive Sciences* **10,** 124–130. ISSN: 1364-6613, 1879-307X (2006).

5. Findlay, J. M. & Gilchrist, I. D. *Active Vision: The Psychology of Looking and Seeing* ISBN: 978-0-19-852479-3 (Oxford University Press, 2003).

6. Rizzolatti, G., Riggio, L., Dascola, I. & Umiltá, C. Reorienting Attention across the Horizontal and Vertical Meridians: Evidence in Favor of a Premotor Theory of Attention. *Neuropsychologia* **25,** 31–40. ISSN: 0028-3932 (1987).

7. Rizzolatti, G., Riggio, L. & Sheliga, B. M. in *Attention and Performance 15: Conscious and Nonconscious Information Processing* 232–265 (The MIT Press, Cambridge, MA, US, 1994). ISBN: 978-0-262-21012-6.

8. Kowler, E., Anderson, E., Dosher, B. & Blaser, E. The Role of Attention in the Programming of Saccades. *Vision Research* **35,** 1897–1916. ISSN: 0042-6989 (1995).

9. Deubel, H. & Schneider, W. X. Saccade Target Selection and Object Recognition: Evidence for a Common Attentional Mechanism. *Vision Research* **36,** 1827–1837. ISSN: 0042-6989 (1996).

10. Li, H.-H., Hanning, N. M. & Carrasco, M. To Look or Not to Look: Dissociating Presaccadic and Covert Spatial Attention. *Trends in Neurosciences* **44,** 669–686. ISSN: 0166-2236 (2021).

11. Sheliga, B. M., Riggio, L. & Rizzolatti, G. Orienting of Attention and Eye Movements. *Experimental Brain Research* **98,** 507–522. ISSN: 1432-1106 (1994).

12. Corbetta, M. Frontoparietal Cortical Networks for Directing Attention and the Eye to Visual Locations: Identical, Independent, or Overlapping Neural Systems? *Proceedings of the National Academy of Sciences* **95,** 831–838 (1998).

13. Moore, T., Armstrong, K. M. & Fallah, M. Visuomotor Origins of Covert Spatial Attention. *Neuron* **40,** 671–683. ISSN: 0896-6273 (2003).

14. Jonikaitis, D. & Moore, T. The Interdependence of Attention, Working Memory and Gaze Control: Behavior and Neural Circuitry. *Current Opinion in Psychology. Attention & Perception* **29,** 126–134. ISSN: 2352-250X (2019).

15. de Haan, B., Morgan, P. S. & Rorden, C. Covert Orienting of Attention and Overt Eye Movements Activate Identical Brain Regions. *Brain Research* **1204,** 102–111. ISSN: 0006-8993 (2008).

16. Nobre, A. C., Gitelman, D. R., Dias, E. C. & Mesulam, M. M. Covert Visual Spatial Orienting and Saccades: Overlapping Neural Systems. *NeuroImage* **11,** 210–216. ISSN: 1053-8119 (2000).

17. Corbetta, M. & Shulman, G. L. Control of Goal-Directed and Stimulus-Driven Attention in the Brain. *Nature Reviews Neuroscience* **3,** 201–215. ISSN: 1471-0048 (2002).

18. Fairhall, S., Indovina, I., Driver, J. & Macaluso, E. The Brain Network Underlying Serial Visual Search: Comparing Overt and Covert Spatial Orienting, for Activations and for Effective Connectivity. *Cerebral Cortex* **19,** 2946–2958. ISSN: 1047-3211 (2009).

19. Bruce, C. J., Goldberg, M. E., Bushnell, M. C. & Stanton, G. B. Primate Frontal Eye Fields. II. Physiological and Anatomical Correlates of Electrically Evoked Eye Movements. *Journal of Neurophysiology* **54,** 714–734. ISSN: 0022-3077 (1985).

20. Sparks, D. L. The Brainstem Control of Saccadic Eye Movements. *Nature Reviews Neuroscience* **3,** 952–964. ISSN: 1471-0048 (2002).

21. Perry, R. J. & Zeki, S. The Neurology of Saccades and Covert Shifts in Spatial Attention: An Event-Related fMRI Study. *Brain* **123,** 2273–2288. ISSN: 0006-8950 (2000).

22. Moore, T. & Fallah, M. Control of Eye Movements and Spatial Attention. *Proceedings of the National Academy of Sciences* **98,** 1273–1276 (2001).

23. Moore, T. & Fallah, M. Microstimulation of the Frontal Eye Field and Its Effects on Covert Spatial Attention. *Journal of Neurophysiology* **91,** 152–162. ISSN: 0022-3077 (2004).

24. Moore, T. & Armstrong, K. M. Selective Gating of Visual Signals by Microstimulation of Frontal Cortex. *Nature* **421,** 370–373. ISSN: 1476-4687 (2003).

25. Tehovnik, E. J. Electrical Stimulation of Neural Tissue to Evoke Behavioral Responses. *Journal of Neuroscience Methods* **65,** 1–17. ISSN: 0165-0270 (1996).

26. Messinger, A., Cirillo, R., Wise, S. P. & Genovesio, A. Separable Neuronal Contributions to Covertly Attended Locations and Movement Goals in Macaque Frontal Cortex. *Science Advances* **7,** eabe0716 (2021).

27. Park, J., Holmes, C. D. & Snyder, L. H. *Compositional Architecture: Orthogonal Neural Codes for Task Context and Spatial Memory in Prefrontal Cortex* 2025.

28. Jonikaitis, D., Noudoost, B. & Moore, T. Dissociating the Contributions of Frontal Eye Field Activity to Spatial Working Memory and Motor Preparation. *Journal of Neuroscience* **43,** 8681–8689. ISSN: 0270-6474, 1529-2401 (2023).

29. Juan, C.-H. *et al.* Segregation of Visual Selection and Saccades in Human Frontal Eye Fields. *Cerebral Cortex* **18,** 2410–2415. ISSN: 1047-3211 (2008).

30. Schall, J. D. On the Role of Frontal Eye Field in Guiding Attention and Saccades. *Vision Research. Visual Attention* **44,** 1453–1467. ISSN: 0042-6989 (2004).

31. Thompson, K. G. & Bichot, N. P. in *Progress in Brain Research* 249–262 (Elsevier, 2005).

32. Koevoet, D., Strauch, C., Naber, M. & Van der Stigchel, S. The Costs of Paying Overt and Covert Attention Assessed With Pupillometry. *Psychological Science* **34,** 887–898. ISSN: 0956-7976 (2023).

33. Hanning, N. M., Fernández, A. & Carrasco, M. Dissociable Roles of Human Frontal Eye Fields and Early Visual Cortex in Presaccadic Attention. *Nature Communications* **14,** 5381. ISSN: 2041-1723 (2023).

34. Fernández, A., Hanning, N. M. & Carrasco, M. Transcranial Magnetic Stimulation to Frontal but Not Occipital Cortex Disrupts Endogenous Attention. *Proceedings of the National Academy of Sciences* **120,** e2219635120 (2023).

35. Wu, T., Mackie, M.-A., Chen, C. & Fan, J. Representational Coding of Overt and Covert Orienting of Visuospatial Attention in the Frontoparietal Network. *NeuroImage* **261,** 119499. ISSN: 1053-8119 (2022).

36. Strauch, C., Wang, C.-A., Einhäuser, W., Van der Stigchel, S. & Naber, M. Pupillometry as an Integrated Readout of Distinct Attentional Networks. *Trends in Neurosciences* **45,** 635–647. ISSN: 0166-2236 (2022).

37. Joshi, S. & Gold, J. I. Pupil Size as a Window on Neural Substrates of Cognition. *Trends in Cognitive Sciences* **24,** 466–480. ISSN: 1364-6613, 1879-307X (2020).

38. Nunez, P. L. *Electric Fields of the Brain: The Neurophysics of EEG.* 2nd ed. (Oxford University Press, 2006).

39. Lopes da Silva, F. EEG and MEG: Relevance to Neuroscience. *Neuron* **80,** 1112–1128. ISSN: 1097-4199 (2013).

40. Foster, J. J., Sutterer, D. W., Serences, J. T., Vogel, E. K. & Awh, E. Alpha-Band Oscillations Enable Spatially and Temporally Resolved Tracking of Covert Spatial Attention. *Psychological Science* **28,** 929–941. ISSN: 0956-7976 (2017).

41. Jones, H. M., Diaz, G. K., Ngiam, W. X. Q. & Awh, E. Electroencephalogram Decoding Reveals Distinct Processes for Directing Spatial Attention and Encoding Into Working Memory. *Psychological Science,* 09567976241263002. ISSN: 0956-7976 (2024).

42. Thyer, W. *et al.* Storage in Visual Working Memory Recruits a Content-Independent Pointer System. *Psychological Science* **33,** 1680–1694. ISSN: 0956-7976 (2022).

43. van Moorselaar, D. *et al.* Spatially Selective Alpha Oscillations Reveal Moment-by-Moment Trade-offs between Working Memory and Attention. *Journal of Cognitive Neuroscience* **30,** 256–266. ISSN: 0898-929X (2018).

44. Samaha, J., Sprague, T. C. & Postle, B. R. Decoding and Reconstructing the Focus of Spatial Attention from the Topography of Alpha-band Oscillations. *Journal of Cognitive Neuroscience* **28,** 1090–1097. ISSN: 0898-929X (2016).

45. Foster, J. J., Bsales, E. M. & Awh, E. Covert Spatial Attention Speeds Target Individuation. *Journal of Neuroscience* **40,** 2717–2726. ISSN: 0270-6474, 1529-2401 (2020).

46. Grootswagers, T., Wardle, S. G. & Carlson, T. A. Decoding Dynamic Brain Patterns from Evoked Responses: A Tutorial on Multivariate Pattern Analysis Applied to Time Series Neuroimaging Data. *Journal of Cognitive Neuroscience* **29,** 677–697. ISSN: 0898-929X (2017).

47. Carrasco, C. D., Bahle, B., Simmons, A. M. & Luck, S. J. Using Multivariate Pattern Analysis to Increase Effect Sizes for Event-Related Potential Analyses. *Psychophysiology* **61,** e14570. ISSN: 1469-8986 (2024).

48. Thompson, K. G., Biscoe, K. L. & Sato, T. R. Neuronal Basis of Covert Spatial Attention in the Frontal Eye Field. *Journal of Neuroscience* **25,** 9479–9487. ISSN: 0270-6474, 1529-2401 (2005).

49. Juan, C.-H., Shorter-Jacobi, S. M. & Schall, J. D. Dissociation of Spatial Attention and Saccade Preparation. *Proceedings of the National Academy of Sciences* **101,** 15541–15544 (2004).

50. Jonikaitis, D., Xia, R. & Moore, T. Robust Encoding of Stimulus–Response Mapping by Neurons in Visual Cortex. *Proceedings of the National Academy of Sciences* **122,** e2408079122 (2025).

51. Bruce, C. J. & Goldberg, M. E. Primate Frontal Eye Fields. I. Single Neurons Discharging before Saccades. *Journal of Neurophysiology* **53,** 603–635. ISSN: 0022-3077 (1985).

52. Rolfs, M. Attention in Active Vision: A Perspective on Perceptual Continuity Across Saccades. *Perception* **44,** 900–919. ISSN: 0301-0066 (2015).

53. Cavanagh, P., Hunt, A. R., Afraz, A. & Rolfs, M. Visual Stability Based on Remapping of Attention Pointers. *Trends in Cognitive Sciences* **14,** 147–153. ISSN: 1364-6613, 1879-307X (2010).

54. Melcher, D. Predictive Remapping of Visual Features Precedes Saccadic Eye Movements. *Nature Neuroscience* **10,** 903–907. ISSN: 1546-1726 (2007).

55. Duhamel, J.-R., Colby, C. L. & Goldberg, M. E. The Updating of the Representation of Visual Space in Parietal Cortex by Intended Eye Movements. *Science* **255,** 90–92 (1992).

56. Li, H.-H., Pan, J. & Carrasco, M. Different Computations Underlie Overt Presaccadic and Covert Spatial Attention. *Nature Human Behaviour* **5,** 1418–1431. ISSN: 2397-3374 (2021).

57. Gutteling, T. P., van Ettinger-Veenstra, H. M., Kenemans, J. L. & Neggers, S. F. W. Lateralized Frontal Eye Field Activity Precedes Occipital Activity Shortly before Saccades: Evidence for Cortico-cortical Feedback as a Mechanism Underlying

Covert Attention Shifts. *Journal of Cognitive Neuroscience* **22,** 1931–1943. ISSN: 0898-929X (2010).

58. Koevoet, D. *et al.* Effort Drives Saccade Selection. *eLife* **13,** 1–25. ISSN: 2050-084X (2025).

59. Kahneman, D. *Attention and Effort* ISBN: 978-0-13-050518-7 (Prentice-Hall, 1973).

60. Beatty, J. Task-Evoked Pupillary Responses, Processing Load, and the Structure of Processing Resources. *Psychological Bulletin* **91,** 276–292. ISSN: 1939-1455 (1982).

61. Koevoet, D., Strauch, C., Van der Stigchel, S., Mathôt, S. & Naber, M. Revealing Visual Working Memory Operations with Pupillometry: Encoding, Maintenance, and Prioritization. *WIREs Cognitive Science,* e1668. ISSN: 1939-5086 (2024).

62. Mathôt, S. Pupillometry: Psychology, Physiology, and Function. *Journal of Cognition* **1,** 16. ISSN: 2514-4820 (2018).

63. Sirois, S. & Brisson, J. Pupillometry. *WIREs Cognitive Science* **5,** 679–692. ISSN: 1939-5086 (2014).

64. van der Wel, P. & van Steenbergen, H. Pupil Dilation as an Index of Effort in Cognitive Control Tasks: A Review. *Psychonomic Bulletin & Review* **25,** 2005–2015. ISSN: 1531-5320 (2018).

65. Smith, D. T. & Schenk, T. The Premotor Theory of Attention: Time to Move On? *Neuropsychologia. Special Issue: Spatial Neglect and Attention* **50,** 1104–1114. ISSN: 0028-3932 (2012).

66. Casteau, S. & Smith, D. T. Associations and Dissociations between Oculomotor Readiness and Covert Attention. *Vision* **3,** 17. ISSN: 2411-5150 (2019).

67. Sprague, T. C. *et al.* Inverted Encoding Models Assay Population-Level Stimulus Representations, Not Single-Unit Neural Tuning. *eNeuro* **5.** ISSN: 2373-2822 (2018).

68. Brouwer, G. J. & Heeger, D. J. Decoding and Reconstructing Color from Responses in Human Visual Cortex. *Journal of Neuroscience* **29,** 13992–14003. ISSN: 0270-6474, 1529-2401 (2009).

69. Günseli, E. *et al.* Encoded and Updated Spatial Working Memories Share a Common Representational Format in Alpha Activity. *iScience* **27.** ISSN: 2589-0042 (2024).

70. Kriegeskorte, N., Goebel, R. & Bandettini, P. Information-Based Functional Brain Mapping. *Proceedings of the National Academy of Sciences* **103,** 3863–3868 (2006).

71. Rolfs, M. & Carrasco, M. Rapid Simultaneous Enhancement of Visual Sensitivity and Perceived Contrast during Saccade Preparation. *Journal of Neuroscience* **32,** 13744–13752a. ISSN: 0270-6474, 1529-2401 (2012).

72. Schall, J. D. Visuomotor Functions in the Frontal Lobe. *Annual Review of Vision Science* **1,** 469–498. ISSN: 2374-4642, 2374-4650 (2015).

73. Rao, H. M., Mayo, J. P. & Sommer, M. A. Circuits for Presaccadic Visual Remapping. *Journal of Neurophysiology* **116,** 2624–2636. ISSN: 0022-3077 (2016).

74. Mayo, J. P., DiTomasso, A. R., Sommer, M. A. & Smith, M. A. Dynamics of Visual Receptive Fields in the Macaque Frontal Eye Field. *Journal of Neurophysiology* **114,** 3201–3210. ISSN: 0022-3077 (2015).

75. Awh, E., Jonides, J. & Reuter-Lorenz, P. A. Rehearsal in Spatial Working Memory. *Journal of Experimental Psychology: Human Perception and Performance* **24,** 780–790. ISSN: 1939-1277 (1998).

76. Williams, M., Pouget, P., Boucher, L. & Woodman, G. F. Visual–Spatial Attention Aids the Maintenance of Object Representations in Visual Working Memory. *Memory & Cognition* **41,** 698–715. ISSN: 1532-5946 (2013).

77. Awh, E. & Jonides, J. Overlapping Mechanisms of Attention and Spatial Working Memory. *Trends in Cognitive Sciences* **5,** 119–126. ISSN: 1364-6613 (2001).

78. Foster, J. J., Bsales, E. M., Jaffe, R. J. & Awh, E. Alpha-Band Activity Reveals Spontaneous Representations of Spatial Position in Visual Working Memory. *Current biology* **27,** 3216–3223.e6. ISSN: 1879-0445 (2017).

79. Foster, J. J., Sutterer, D. W., Serences, J. T., Vogel, E. K. & Awh, E. The Topography of Alpha-Band Activity Tracks the Content of Spatial Working Memory. *Journal of Neurophysiology* **115,** 168–177. ISSN: 0022-3077 (2016).

80. Koevoet, D., Naber, M., Strauch, C. & Van der Stigchel, S. Presaccadic Attention Shifts Up- and Downwards: Evidence From the Pupil Light Response. *Psychophysiology* **62,** e70047. ISSN: 1469-8986 (2025).

81. Mathôt, S. & Theeuwes, J. Visual Attention and Stability. *Philosophical Transactions of the Royal Society B: Biological Sciences* **366,** 516–527 (2011).

82. Van der Stigchel, S. & Hollingworth, A. Visuospatial Working Memory as a Fundamental Component of the Eye Movement System. *Current Directions in Psychological Science* **27,** 136–143. ISSN: 0963-7214 (2018).

83. Currie, C. B., McConkie, G. W., Carlson-Radvansky, L. A. & Irwin, D. E. The Role of the Saccade Target Object in the Perception of a Visually Stable World. *Perception & Psychophysics* **62,** 673–683. ISSN: 1532-5962 (2000).

84. Deubel, H., Bridgeman, B. & Schneider, W. X. Immediate Post-Saccadic Information Mediates Space Constancy. *Vision Research* **38,** 3147–3159. ISSN: 0042-6989 (1998).

85. Deubel, H. & Schneider, W. X. Perceptual Stability and Postsaccadic Visual Information: Can Man Bridge a Gap? *Behavioral and Brain Sciences* **17,** 259–260. ISSN: 1469-1825, 0140-525X (1994).

86. McConkie, G. W. & Currie, C. B. Visual Stability across Saccades While Viewing Complex Pictures. *Journal of Experimental Psychology. Human Perception and Performance* **22,** 563–581. ISSN: 0096-1523 (1996).

87. Szinte, M., Jonikaitis, D., Rangelov, D. & Deubel, H. Pre-Saccadic Remapping Relies on Dynamics of Spatial Attention. *eLife* **7** (ed King, A. J.) e37598. ISSN: 2050-084X (2018).

88. Fabius, J. H., Fracasso, A., Nijboer, T. C. W. & Van der Stigchel, S. Time Course of Spatiotopic Updating across Saccades. *Proceedings of the National Academy of Sciences* **116,** 2027–2032 (2019).

89. Moran, C., Johnson, P. A., Landau, A. N. & Hogendoorn, H. Decoding Remapped Spatial Information in the Peri-Saccadic Period. *Journal of Neuroscience* **44.** ISSN: 0270-6474, 1529-2401 (2024).

90. Fabius, J. H., Fracasso, A., Acunzo, D. J., der Stigchel, S. V. & Melcher, D. Low-Level Visual Information Is Maintained across Saccades, Allowing for a Postsaccadic Handoff between Visual Areas. *Journal of Neuroscience* **40,** 9476–9486. ISSN: 0270-6474, 1529-2401 (2020).

91. Wang, X. *et al.* Perisaccadic and Attentional Remapping of Receptive Fields in Lateral Intraparietal Area and Frontal Eye Fields. *Cell Reports* **43.** ISSN: 2211-1247 (2024).

92. Duncan, D. H., van Moorselaar, D. & Theeuwes, J. Pinging the Brain to Reveal the Hidden Attentional Priority Map Using Encephalography. *Nature Communications* **14,** 4749. ISSN: 2041-1723 (2023).

93. Bae, G.-Y. & Luck, S. J. Reactivation of Previous Experiences in a Working Memory Task. *Psychological Science* **30,** 587–595 (2019).

94. Brainard, D. H. The Psychophysics Toolbox. *Spatial Vision* **10,** 433–436. ISSN: 0169-1015 (1997).

95. Pelli, D. G. The VideoToolbox Software for Visual Psychophysics: Transforming Numbers into Movies. *Spatial vision* **10,** 437–442. ISSN: 0169-1015 (1997).

96. Adam, K. C. S., Vogel, E. K. & Awh, E. Multivariate Analysis Reveals a Generalizable Human Electrophysiological Signature of Working Memory Load. *Psychophysiology* **57,** e13691. ISSN: 1469-8986 (2020).

97. Pedregosa, F. *et al.* Scikit-Learn: Machine Learning in Python. *The Journal of Machine Learning Research* **12,** 2825–2830. ISSN: 1532-4435 (2011).

98. van der Lubbe, R. H. J., Neggers, S. F. W., Verleger, R. & Kenemans, J. L. Spatiotemporal Overlap between Brain Activation Related to Saccade Preparation and Attentional Orienting. *Brain Research* **1072,** 133–152. ISSN: 0006-8993 (2006).

99. Talcott, T. N., Kiat, J. E., Luck, S. J. & Gaspelin, N. Is Covert Attention Necessary for Programming Accurate Saccades? Evidence from Saccade-Locked Event-Related Potentials. *Attention, Perception, & Psychophysics* **87,** 172–190. ISSN: 1943-393X (2025).

100. Talcott, T. N. & Gaspelin, N. Eye Movements Are Not Mandatorily Preceded by the N2pc Component. *Psychophysiology* **58,** e13821. ISSN: 1469-8986 (2021).

101. Parks, N. A. & Corballis, P. M. Electrophysiological Correlates of Presaccadic Remapping in Humans. *Psychophysiology* **45,** 776–783. ISSN: 1469-8986 (2008).

102. Huber-Huber, C., Ditye, T., Marchante Fernández, M. & Ansorge, U. Using Temporally Aligned Event-Related Potentials for the Investigation of Attention Shifts Prior to and during Saccades. *Neuropsychologia. Spatial and Non-Spatial Aspects of Visual Attention: Interactive Cognitive Mechanisms and Neural Underpinnings* **92,** 129–141. ISSN: 0028-3932 (2016).

103. Hajonides, J. E., Nobre, A. C., van Ede, F. & Stokes, M. G. Decoding Visual Colour from Scalp Electroencephalography Measurements. *NeuroImage* **237,** 118030. ISSN: 1053-8119 (2021).

104. Savitzky, A. & Golay, M. J. E. Smoothing and Differentiation of Data by Simplified Least Squares Procedures. *Analytical Chemistry* **36,** 1627–1639 (1964).

105. Van der Stigchel, S. & de Vries, J. P. There Is No Attentional Global Effect: Attentional Shifts Are Independent of the Saccade Endpoint. *Journal of Vision* **15,** 17. ISSN: 1534-7362 (2015).

106. Mathôt, S. & Vilotijević, A. Methods in Cognitive Pupillometry: Design, Preprocessing, and Statistical Analysis. *Behavior Research Methods.* ISSN: 1554-3528 (2022).

107. Koevoet, D., Naber, M., Strauch, C. & Van der Stigchel, S. The Intensity of Internal and External Attention Assessed with Pupillometry. *Journal of Cognition* **7,** 8. ISSN: 2514-4820 (2024).

108. Lakens, D. Calculating and Reporting Effect Sizes to Facilitate Cumulative Science: A Practical Primer for t-Tests and ANOVAs. *Frontiers in Psychology* **4.** ISSN: 1664-1078 (2013).

109. Maris, E. & Oostenveld, R. Nonparametric Statistical Testing of EEG- and MEG-data. *Journal of Neuroscience Methods* **164,** 177–190. ISSN: 0165-0270 (2007).