When things add up: environmental structure and microbial interactions drive
antibiotic-resistance plasmid evolution

Laurens E. Zandbergen[1], Mishelle Bustamante[1], Jan-Maarten Luursema[2], Thomas Hackl[1],
Marjon G.J. de Vos[1*]

1 Groningen Institute of Evolutionary Life Sciences (GELIFES), University of Groningen,
Groningen, The Netherlands

2 Radboud University Medical Center, Nijmegen, the Netherlands

*corresponding author: m.g.j.de.vos@rug.nl

Competing interests

The authors declare no conflict of interest.

13
14

Abstract

Antimicrobial resistance is recognized as a major global health threat. Pathogens can rapidly evolve resistance which diminishes the impact of antimicrobial treatments. The presence of other microbes in polymicrobial infections can impact the development of antimicrobial resistance of pathogens, yet it is unclear how the interplay between the environmental structure and the presence of a polymicrobial community affect the rate of plasmid mediated antibiotic resistance evolution. This study investigates the effect of environmental structure and microbial interactions on antibiotic resistance evolution in a uropathogenic *Escherichia coli.* We conducted a serial transfer experiment across well-mixed and spatially structured environments with increasing concentrations of trimethoprim-sulfamethoxazole, in the presence and absence of a polymicrobial community. Our results revealed that *E. coli* in community context displayed parallel evolutionary trajectories, leading to higher final antibiotic tolerance, while spatial structures allowed for prolonged resistance evolution. Copy number variation of the plasmid-borne resistance locus varied significantly across the different evolutionary contexts. *E. coli* lineages evolved in the absence of the community, in the well-mixed isolation environments demonstrated the highest copy number increases, while those in the presence of the community showed minimal changes compared to the ancestor. These findings underscore the intricate relationship between the genetic basis of resistance, the environmental structure and the microbial ecology in plasmid-mediated antimicrobial resistance evolution.

Introduction

Polymicrobial communities can have a profound effect on the severity of infections[1–4] and can diminish treatment efficacy of pathogens.[5–8] Pathogens are increasingly becoming resistant to antimicrobials, complicating treatments.[9] According to the World Health Organization, antimicrobial resistance (AMR) is among the top ten of threats to global health.[10]

Emerging research shows that AMR can be a collective trait that develops within bacterial communities.[7,10,11] For example, tolerance to antibiotics can be increased in a community by decreasing antibiotic pressures in the local environment through antibiotic degradation, leading to increased pathogen density.[12–14] Higher population densities have been shown to reduce the susceptibility to certain antibiotics, which is known as the inoculum effect.[15,16] However, microbial communities are more often driven by competition rather than cooperation, which would lead to a reduction in average population size per member and therefore a change in survival.[17] Yet, cooperation occurs and it has been observed that interspecies interactions can lead to antibiotic protective effects.[10,18]

The environmental structure of bacterial communities can have a large impact on bacterial growth. Virtually all known bacteria are capable of creating biofilms or attach to surfaces, which enhances tolerance to environmental stressors, both biotic and abiotic.[19,20] Spatial growth versus growth in a well-mixed environment has different effects on the selective pressures and growth of bacterial populations, such as alterations in competitive and mutualism dynamics.[21,22] Environmental structure has been implicated as a critical factor for the stability and coexistence of species in a community.[23–25] For example, gene expression related to quorum sensing, and consequently antibiotic susceptibility,[26] is different in a spatially structured system compared to a well-mixed environment.[27] Importantly, theorical and empirical research suggests that social interactions of bacteria in spatially structured environments are significantly altered compared to environments with better mixing, leading to different survival outcomes.[28–31]

Different environmental structures can alter the ecological interactions of microbes. Such different ecological interactions can subsequently affect microbial co-existence and evolution.[23–25] Here we seek to investigate the effect of the environmental structure on antibiotic resistance evolution of a focal pathogenic species in the presence and absence of a four-member microbial community, and in two contrasting environments, a well-mixed liquid and a spatially structured solid environment. Previous studies indicated that populations in spatially structured environments adapt more slowly than those in well-mixed environments and evolution in a spatially structured environments increases genetic drift and founder effects on local populations reducing the rate of adaptation.[21,32] It has also been shown that spatial

structures promote species coexistence and that there is a decrease in metabolite exchange in continuously mixed environments.[33,34] Based on these studies, we hypothesized that the evolution of resistance will be slower in a spatial environment,[21,32,35,36] and that in the well-mixed environment the focal species will be less able to benefit from the presence of the community members.[33,34]

Results

Environmental structure and microbial community impact evolutionary trajectories

To study the effects of structure and community context on the evolution of antibiotic resistance, we evolved a uropathogenic isolate of *E. coli* carrying a plasmid encoding resistance genes that only confer low resistance in a serial transfer evolution experiment. Evolving cultures were transferred bi-daily over 120 days, in artificial urine medium (AUM) containing increasing levels of trimethoprim-sulfamethoxazole, under four different conditions. Specifically, five replicates of *E. coli* per condition were evolved in both a well-mixed and spatially structured AUM environment, both in the presence and absence of a community of three other bacterial species. These other species are often co-isolated with *E. coli*; *Klebsiella pneumoniae, Enterococcus faecium* and *Staphylococcus epidermidis* in polymicrobial urinary tract infections (Materials and Methods).[37] During the serial transfer experiment antibiotic concentrations were increased by 20%, depending on whether the population size of the *E. coli* focal species exceeded a certain threshold. Thresholds were set at colony size exceeding >1 mm (which is an increase of 20% colony size at minimum) for the spatially structured environments. And at $>1*10^6$ colony forming units (CFU)/mL for the liquid environments. Because colonies could only be counted the day after a transfer, the population size at the previous transfer was taken as the threshold. Therefore, the fastest rate of antibiotic increase was every other transfer. Community members, other than *E. coli*, already had high resistances to trimethoprim and sulfamethoxazole (>1 mg/mL), and were always present at a low density in the community at the start of each transfer ($<10^5$). To prevent their loss during the serial transfer experiment, a 100x dilution of wild-type stationary phase monocultures of the community members was added at each transfer together with a 100x dilution of the *E. coli* from the previous step (Materials and Methods). The initial concentration of antibiotics was 50% of the *E. coli* minimum inhibitory concentration (MIC) for the 1:5 ratio of trimethoprim-sulfamethoxazole, 42 µg/mL trimethoprim (TMP) and 210 µg/mL sulfamethoxazole (SMX). This ratio was picked to resemble the ratio used in clinical settings.[38]

Observing the level of antibiotic increase of *E. coli* per replicate lineage in each condition, we
find that the five replicate phenotypic evolutionary trajectories overlapped in three of the four
conditions. Particularly, the replicate lineages overlapped for *E. coli* evolved in the spatially-
structured environments in the presence and absence of the community, and *E. coli* evolved
in the well-mixed environment in the presence of the community (Figure 1AB). The presence
of the community seemingly leads to the canalization of the phenotypic evolutionary
trajectories of *E. coli* in the well-mixed environment. In contrast, *E. coli* evolved in isolation in
this same well-mixed environment had markedly increased divergent phenotypic evolutionary
trajectories. We observed a 3.5x difference in antibiotic concentration between the replicates
with the highest (1.92:9.58 mg/mL TMP:SMX) and lowest (0.54:2.68 mg/mL TMP:SMX)
antibiotic concentration at the final transfer (Figure 1A). Evolution in the spatially structured
environments did not lead to a difference in the divergence of phenotypic evolutionary
trajectories, as all five replicate trajectories overlapped (Figure 1B).

For the three overlapping replicate trajectories, the rate of evolution in the first few weeks were
close to the hypothetical maximum rate. For the well-mixed environment, this is until day 80,
and the spatially structured environment until day 48 for the *E. coli* evolved in presence of the
community and day 30 for *E. coli* evolved in isolation. (Figure 1AB). This suggests that within
these environments, selective pressures drove evolution in a specific direction, potentially
funneling evolution towards similar evolutionary outcomes creating the strong convergence of
the trajectories.

The final concentration of antibiotics that the evolved *E. coli* populations can sustain is higher
in the community context compared to in isolation (2.3:11.5 mg/mL vs. 1.9:9.5 mg/mL
TMP:SMX in the well-mixed environment and 4.0:20 mg/mL vs. 2.8:14 mg/mL TMP:SMX for
the spatially-structured environment). This shows that the presence of the community had a
positive impact on the level of antibiotics that *E. coli* could sustain in both the well-mixed as
well as the spatially structured environment (P=0.023 for the community impact in the well-
mixed environment and P<0.001 for the community impact in the spatially structured
environment, *Welch's t-test*).

Additionally, the structure of the environment, either the well-mixed AUM or spatially structured
AUM-agar, had an impact on the evolutionary trajectories of *E. coli*. Initially, for both well-
mixed and spatially structured environments, the rate of antibiotic increase was identical.
However, after the first 50 days of evolution (25 transfers) the trajectories started to diverge.
The rate at which the antibiotic could be increased dropped slightly for the spatially structured
community environment, compared to the well-mixed community environment. Despite the

fact that the spatially structured environment enabled a slower rate of antibiotic increase, the concentration of antibiotics at the final transfer of the evolution experiment is higher in the spatially structured environments, compared to the well-mixed environment (Figure 1B). The antibiotic increase in the well-mixed environment stagnated around day 108. Yet the rate of evolution in the spatially structured environment had not stagnated (>10 days without antibiotic concentration increase) at the final transfer. We cannot exclude that the solvents used for the antibiotic stocks could also impact the fitness of the *E. coli* and community members, particularly at the end of the experiments when antibiotic concentrations are high. The final concentration of TMP:SMX in the spatially structured environments the concentration was 4:20 mg/mL for the *E. coli* in the presence of the community and 2.8:14 mg/mL for the isolated *E. coli*. The concentrations in the well-mixed environment were lower; in the presence of the community, 2.3:11.5 mg/mL, and the highest level of the isolated *E. coli,* 1.9:9.5 mg/mL. Yet, these results indicate that the environmental structure, well-mixed in AUM or spatially structured on AUM-agar, differentiated the course of the evolutionary trajectories (one-way ANOVA, *P*<0.001 for liquid vs. solid).

These results show that both the presence of the community, as well as the structured environment had an impact on the evolutionary trajectories of trimethoprim-sulfamethoxazole resistance of *E. coli*. Notably, community context led to an increased parallelism of evolutionary paths and higher final antibiotic tolerance, while spatial structure allowed for longer sustained resistance evolution and higher final antibiotic concentrations compared to the well-mixed environments.

Community interactions mediate response to selection

Uropathogenic *E. coli* is known for its flexible metabolism, being able to adapt and respond to fluctuating nutrients in the environment.[39] We questioned whether *E. coli* would respond differently to the presence of antibiotics in the presence of the community. We observed that the focal *E. coli* population is a good competitor in the AUM environment, both in the well-mixed and spatially structured environments. It is also able to outgrow the other community members. For example, in the first 30 days the density of *E. coli* was between the 10^8 and 10^9 CFU/mL for most replicates at the time of transfer in the well-mixed environment in the presence of the community. Such a high population density was maintained even when antibiotic reached the initial MIC after 12 days. Even though the community members are highly resistant (>1:5 mg/mL TMP:SMX), *E. coli* is able to reach higher abundances from the onset of the experiment. In the absence of the community, we observe lower abundances of *E. coli* in the well-mixed environment at the time of transfer (Figure 1C) (*P*<0.001 for the first

30 days*,* ANOVA). Also on the spatially structured environment, colonies of *E. coli* surrounded by the community members were much larger in surface area (~3x) compared to *E. coli* colonies surrounded by other *E. coli* colonies (Figure 1D) (*P*<0.001 for the difference between community and isolation colonies, *Welch's t-test*, *Materials and Methods*). The population sizes of the five replicates lineages evolved within the same environment were similar (Figure 1CD, *P*>0.05, independent samples *t*-test with Bonferroni correction).

The presence of the three-species community benefits the *E. coli* antibiotic resistance evolution rate. In the presence of the community, it takes 12 days before *E. coli* lineages pass the wild type MIC level. In isolation that takes longer, after 16-18 days (*P*<0.001, ANOVA). Despite larger colony sizes in the presence of the community, all replicates, in the presence of the community as well as in in isolation, reached the native MIC after 12 days in the spatially structured environment.

The beneficial effects of the community weaned off over time. For the well-mixed environment, after around 80 days the rate of antibiotic increase slowed down (Figure 1A, dark blue graph). Interestingly, this occurred for all five replicates at the same time, which suggests that the community canalizes the phenotypic trajectories in a general manner. We do not observe such a synchronization of the phenotypic evolutionary trajectories for the replicate lineages of *E. co*li in isolation (Figure 1A, light blue graphs). In the spatially structured environment, we also find such a synchronized slow-down, yet at another time point (48 days) and antibiotic concentration (446:2229 µg/mL in solid vs. 1917:9585 µg/mL in liquid TMP:SMX). In this spatially structured environment, the canalization of the phenotypic trajectories is not confined to the replicates evolving in the presence of the community only. We also observe the canalized phenotypic trajectories for the isolated *E. coli*. Yet, similarly to the growth in the well-mixed environment, the slow-down of the rate of antibiotic increase occurs earlier for *E. coli* evolved in isolation. *E. coli* in isolation started diverging from the *E. coli* in the presence of the community after 28 days (179:896 µg/mL TMP:SMX).

The population sizes of the community members also decreased over time. *K. pneumoniae* was present abundantly in the well-mixed environment at the start of the experiment (between 10^7 and 10^8 CFU/mL), this decreased steadily to between 10^6 and 10^7 CFU/mL after 14 days. *K. pneumoniae* was not detected later in the experiments, meaning it had an abundance of at least 100-fold lower than *E. coli*, which would be under the detection limit. Also in the spatially structured environment, *K. pneumoniae*¸ measured by its colony size, steadily decreased in size during the first 14 days, in accordance with the *K. pneumoniae* population size in well-mixed environment. The colonies of *K. pneumoniae* kept decreasing with every increase in antibiotics, showing that increasing antibiotic concentrations affects the wild-type *K.*

pneumoniae fitness negatively (average colony size at day 2   14494 pixels, at day 120 8518

pixels). *E. faecium* was always present in low numbers, around 10^2 -10^3 CFU/mL, as expected

from previous studies,[4,10] yet population sizes became smaller with increasing antibiotic

concentrations decreasing at least 5x between the start and the end of the 120 days.

Surprisingly, *S. epidermidis* was hardly detected during the experiments, indicating population

sizes below 10^3 CFU/mL, even though *S. epidermidis* cultured in isolation is more abundant

than *E. faecium* (10^4-10^5 CFU/mL). It is likely that competition with other species, or the

increasing antibiotics concentrations hindered *S. epidermidis* to thrive, even though it can

survive high concentrations[8]. The decrease of both the focal species and community members

over time could point to an increase in competition, a decrease in cooperation or a fitness

decrease due to the increase in antibiotics concentration in the environment.

Environmental structure and community effect on antibiotic resistance

Next, we investigated how the two differently structured environments (spatially structured or

well-mixed) had affected the evolutionary outcome of antibiotic resistance evolution.

Comparing the *E. coli* lineages that evolved in isolation with the lineages that evolved in the

presence of the community, we find that the well-mixed environment and the spatially

structured environment had a different effect on the final level of antibiotics. Despite that the

lineages evolved in the spatially structured environment tolerated higher concentrations of

antibiotics at the end of the experiment compared to the well-mixed environments (Figure

1AB), we find that replicates evolved in the spatially structured environment had a lower MIC

than those in the well-mixed environment in the presence of the community (resp. 0.8:4 mg/mL

TMP:SMX and 3.2:16 mg/mL TMP:SMX, Figure 2A, Materials and Methods). It is possible that

the surface growth mode or spatial adaptation during the evolution experiments impacted the

fitness and antibiotic susceptibility measurements in liquid.

A similar observation can be made for *E. coli* evolved in isolation in both environments. In the

spatially structured environment, the concentration of antibiotics that could be tolerated at the

final transfer is higher than that in the well-mixed environment. Yet, the MIC of *E. coli* evolved

in the spatial environment was generally lower (MIC 0.92:4.6 mg/mL TMP:SMX spatially

structured environment, versus MIC of 1.06:5.3 mg/mL TMP:SMX for three of the five

replicates from the well-mixed environment ($P<0.001$, *Welch's t-test*). The other two replicates

in the well-mixed environment had a lower MIC (0.27:1.33 and 0.13:0.66 mg/ml TMP:SMX

respectively), and corresponded to those lineages that could sustain a lower final

concentration of antibiotics in the evolution experiment (Figure 1A, light blue graphs, Figure

2A)).

Based on the observation that the population size increased in the presence of community members, we hypothesized that *E. coli* had benefited from the presence of one or more of the community members during the evolution experiment. To investigate the effect of the community on the evolved resistance of all evolved lineages, we measured the minimum bactericidal concentration (MBC) of antibiotics at which we could not observe our focal species anymore when cultured in the presence of the community (Figure 2B, Figure S1 , Materials and Methods). We find that replicates evolved in the well-mixed environment in absence of the community could sustain twice the concentration of antibiotics when cultured in the presence of the community (MBC of 1.6:8 mg/mL TMP:SMX for 3 out of 5 replicates), compared to their MIC without the community of 0.8:4 mg/mL TMP:SMX for those same replicates) (Figure S1). Yet, for the replicates evolved in the well-mixed environment in presence of the community, we observed a decrease in MBC compared to the MIC in presence of the community (MIC of 3.2:16 mg/mL TMP:SMX, MBC 1.6:8 mg/mL TMP:SMX). This means that the community negatively affected the growth of *E. coli* lineages that were evolved together with the community members in the well-mixed environment. Interestingly, *E. coli* replicates evolved in the spatially structured environment benefited from the presence of the community members; these could sustain a two-fold increase in the level of antibiotics compared to the MIC (Figure 2B). These results show that generally, the community has a positive effect on antibiotic tolerance of *E. coli*, even after resistance evolution, except when *E. coli* was evolved in the presence of the community in the well-mixed environment.

We find that the positive effect of the community eventually decreases over time, with increasing antibiotic concentrations (Figure 1CD). An explanation might be the adaptation to the environment itself. As the medium used is relatively poor, it is possible that adaptation to it leads to niche overlap of the focal species with other microbes in the community and therefore more competition,[40] which would correspond with theory suggesting stronger selection in well-mixed environments.[41,42] Alternatively, when antibiotic resistance in *E. coli* increases, the protective effect of the community becomes less beneficial, and *E. coli* may switch to a more competitive state to benefit from the limited resources. Particularly from the spatially structured perspective, theory suggests that communities benefit more from public goods interactions than they would in well-mixed environments,[30,31,43] which could potentially explain why the *E. coli* replicates evolved in the spatially structured community environment could still benefit from the presence of the community compared to those evolved in the well-mixed environment, as public goods interactions might still be prioritized over competition.

Copy number variation of resistance loci on plasmid

To generate insights in the genetic basis underpinning the evolved resistance, we sequenced three clones of the final transfer of each of the five replicate lineages evolved in each condition, and compared these sequences to that of the ancestor (Materials and Methods). A likely target for evolutionary selection is the *folA* (dihydrofolate reductase) gene, which is known to increase resistance against trimethoprim.[44] We found no evidence however for mutations within the *folA* gene in any of the evolved lineages. We did observe an identical frameshift mutation in the *folP* gene in two of the five *E. coli* replicates evolved in the well-mixed environment in the presence of the community (Table S1). The product of which (dihydropteroate synthase) is inhibited by sulfamethoxazole.[38] Disruption of this gene leads to increased sulfamethoxazole tolerance as it leaves more resources for the remaining proteins of the folate pathway and removes a target for the antibiotic itself.[45] However, this mutation, in conjunction with the other mutations in these replicates, does not yield significant population size differences compared with the other three replicates evolved within the same environment ($P>0.05$, independent samples *t*-test with Bonferroni correction).

We did not find any other chromosomal mutations in genes known to confer resistance in the evolved populations (Table S1). The ancestral strain, despite being highly sensitive to the start concentration of TMP-SMX at the beginning of the experiment, did however contain a 138K base pair sized IncFII-plasmid that contains genes annotated to encode for resistance against both antibiotics, *dfrA17* and *sul1,* respectively. *DfrA17* confers resistance to trimethoprim, s*ul1* encodes a dihydropteroate synthase, conferring resistance to sulfamethoxazole. These resistance genes are located in an Tn21-like integron cassette, which is similar to a cassette first described by Liebert in 1999 (Figure S2).[46] Although at that time, the transposon did not contain the *dfrA17* gene, or any of the other resistance genes located on that cassette, yet.

The lack of genetic substitutions in the antibiotic resistance targets on the chromosome, pointed in the direction of resistance associated with the plasmid-encoded resistance genes. Therefore, we assessed the copy numbers of the plasmid in the evolved clones and in the ancestor. The *E. coli* ancestor contained two copies of the plasmid relative to the host chromosome (Figure 3A). The replicate clones of the final transfers had a similar number of copies, this could therefore not explain the increased resistance level. However, there were variations in the integron cassette containing the antibiotic resistance genes. The copy number variations (CNVs) of this cassette do not correlate with the MIC (R^2=0.12, P=0.13, Figure 3B). Surprisingly, some of the lineages evolved in isolation in the well-mixed environment had a relatively low MIC although these contain higher copy numbers of the antibiotic-resistance genes (Figure 3B). This suggests that either the increased copy number of transposable

elements has a negative effect on the fitness of *E. coli*, or the MIC is not only determined by the copy number variation of the resistance genes, but also by the environment, other mutations or the location of integration. Naturally, all these parameters could simultaneously affect the level of resistance[47], and we did previously observe that the environment affects the fitness of our focal species.[4,10]

Zooming in on the resistance locus where the *dfrA17* and *sul1* genes are located, we found substantial copy number variation across conditions, replicates and even clones from the same replicate and environment. Resistance locus abundance ranged from 0 copies, meaning a complete deletion, in one well-mixed environment clone evolved in the presence of the community, to 14 copies relative to the host chromosome for one clone evolved in the well-mixed environment in the absence of the community(Figure 3A, Figure S3). Overall, resistance locus copy numbers were highest in clones that were evolved in isolation, in the absence of the community, in the well-mixed environment (median 5.2) and in the spatially structured environment (median 3.4). Copy numbers in *E. coli* evolved in community context were seemingly unaffected across the majority of clones (median 1.7 in the well-mixed and 2.0 in the spatially structured environment). Across all conditions, some replicate lineages or individual clones deviated from the overall trend. For example, in the well-mixed community context most clones showed no deviations from ancestral copy-numbers, yet replicate 3 clones showed copy number increases and two replicate 4 clones showed decreases to 1 and 0 copies respectively (Figure S3). Compared to the isolates evolved in the well-mixed environment in isolation, each evolutionary context showed significant difference in resistance locus CNV (well-mixed community (*P*=0.0030), spatially structured isolation (*P*=0.025) and spatially structured community (*P*=0.024), *Welch's t-tests*). There was also a significant difference between the *E. coli* clones evolved in the well-mixed community environment and the spatially structured isolated environment (*P*=0.018). Yet, there were not significant differences between the spatially structured environments (*P*=0.87) and the two environments with the community present (*P*=0.059). This shows that the presence of the community and the environmental structure during evolution mediate the antibiotic resistance loci copy number variation.

The implications of these data are multifaceted: overall, we found that lineages evolving antibiotic resistance in isolation had higher copy numbers of the *dfrA17-sul1* resistance locus, suggesting selection for higher copy numbers of these genes in resistance evolution under these conditions. Lineages evolving in the community context, however, showed no CNV increase in the majority of replicates and clones (7 out of 10 replicates or 20 out of 30 clones), and moderate copy number increases in three replicates, suggesting a different selective

context. We had previously noticed that the community positively affects the growth of *E. coli*.
We hypothesize that this physiologically conferred tolerance explains the higher tolerance to
the antibiotics and, in turn, reduced selection for increased copy numbers of resistance genes
for *E. coli* cultured in the presence of the community.

The high level of CNV variation of the integron cassette containing the antibiotic resistance
genes both across and within replicates, suggests high heterogeneity within replicate
populations and points towards a dynamic process with high copy numbers as a potentially
unstable state, possibly maintained by high selective pressure. This hypothesis is in line with
an insertion-sequence-driven copy mechanism for the resistance locus that can quickly
generate new copies in a stochastic manner under stress but, at the same time, increases
genomic instability by multiplying the chances for disruptive random insertions or deleterious
homologous recombination with every new copy. Such negative feedback could facilitate the
simultaneous generation and continuous purging of excess resistance locus copies in plasmid
lineages and explain the high level of inter-clonal variation we observe.[48]

Environmental and community context modulate genetic adaptation

The number of mutations within the genome of *E. coli* mutants differed significantly between
environments. In the well-mixed environment, we observed 25 mutations in total for all clones
of *E. coli* evolved in isolation and only 16 mutations in clones evolved in the presence of the
community. For the spatially structured environment we observed 45 mutations for those
evolved in isolation and 48 for those evolved in the community context (*P*=0.004, *Welch's t-*
test for the difference between the well-mixed and the spatially structured environments). This
fits our expectations, as the spatially structured environment interactions are limited to the
cells in close proximity, leading to increased genetic drift within the population,[49] and a lower
chance of beneficial mutations sweeping through the population. The mutations observed,
with the exception of *folP*, are, as far as we are aware of, not directly linked to antibiotic
resistance to TMP-SMX.[50–52]

Due to the parallel evolutionary trajectories of many of our replicates evolved in the same
environment (Figure 1AB), we hypothesized to see similar genetic substitutions of lineages
evolved under similar circumstances. As a measure of genetic parallelism, which takes into
account the fraction of shared mutations between two isolates we calculated the *H-index*,[50]
(Materials and Methods) on both the gene-level and the Cluster of Orthologous Group (COG)
level (Figure 4) for substitutions in the *E. coli* chromosome. In the H-index, a value of 0
indicates no shared mutations and a value of 1 indicates that all mutations are shared between
the two mutants. Previously, we have seen quite a high degree of genetic parallelism in

replicate populations evolved under the influence of TMP-SMX in the presence of other
bacterial isolates.[14] Here, besides the CNV of the resistance genes on the Tn21-like integron
cassette, we hardly find any such genetic parallelism, except between replicates evolved in
the same spatially structured environment (Figure 4A). This indicates that, for the well-mixed
environment, unique mutations were found in each of the replicates (Table S1). An exception
are two *E. coli* replicates that were evolved in the well-mixed environment in the presence of
the community, that had only one gene which was repeatedly mutated within the genome.
This mutated gene, *folP*, which as mentioned before is targeted by sulfamethoxazole, would
allow *E. coli* to survive higher antibiotic pressures.[51]

In replicate lineages that had evolved on the spatially structured environment in the presence
of the community, two distinct sets of genotypes evolved. Two replicate lineages had an *H-*
index value of 0.75, but shared no mutations with the other three replicate lineages evolved
under the same environmental circumstance. These two replicates shared mutations in the
treC gene. This gene is involved in the trehalose degradation pathway, creating glucose and
glucose-6-phophate from trehalose-6-phosphate and the mutation of this gene points to
adaptation to the nutrients in the environment rather than the antibiotics.[52] The other three
replicates evolved in the spatially structured environment with the community also had an *H-*
index of 0.75 for each pairwise comparison, but also did not share any mutations with the other
two replicate lineages. These three replicate lineages incorporated mutations in the *pfkA, traC*
and *mpl* genes. One of these genes, *traC* is involved in the type IV secretion system, which
plays a role during conjugation.[53] The *traC* gene specifically is involved during cell-cell contact
within conjugation, being a component of the pili used within the secretion system.[54] Other
studies suggest that *traC* also plays a role in pheromone signaling, being a pheromone binding
extracellular protein.[55,56] The mutation in both replicates is a synonymous mutation, potentially
not affecting fitness or function, except perhaps through codon bias.[57] The other genes, *pfkA*
and *mpl* play a role in carbohydrate degradation and cell wall biogenesis respectively. The
mpL gene contains a frameshift mutation, leading to a loss of function. The gene encodes
UDP-N-acetylmuramate, part of the cell-wall recycling machinery.[58] The *pfkA* gene contains
either missense mutations or an in-frame deletion. The gene encodes phosphofructokinase,
and is part of the glycolysis pathway. *PfkA* is responsible for about 95% of
phosphofructokinase activity in *E. coli.*[59] Deletion of *pfkA* leads to a break in the glycolysis
pathway, shifting fructose 6-phosphate availability to other pathways such as the pentose
phosphate pathway, which is important for the generation of NADPH.[60]

One gene that had a mutation in all replicates for the *E. coli* evolved within the spatially
structured environment in isolation was the *nadR* gene, which is part of the NAD salvage
pathway.[61,62] This gene converts β-nicotinamide D-ribonucleotide into diphosphate and NAD^+.

The involved mutations are either missense mutations or deletions, leading to an inactive
variant. Inhibition of this gene has been shown to increase cellular NAD/NADH. NADH itself
is an important regulator for enzymes involved in metabolism, it is therefore possible that the
mutations within *nadR* are an adaptation to the nutrients in the environment rather than the
antibiotics.[63] However, higher levels of cellular NADH have also been linked to contribute to
efflux-mediated antibiotic resistance.[63] Four out of five of the replicates also had mutations
within the *adhE* gene, which encodes aldehyde-alcohol dehydrogenase.[64] It catalyzes the
reaction of acetaldehyde into acetyl-CoA, also creating NADH in the process.

Overall, evolution within the spatially structured environments containing TMP-SMX led to
mutations within genes leading to increased cellular NADH/NADPH within *E. coli* for eight out
of ten replicates. A previous study described to importance of NADPH for the stabilization of
the dihydrofolate reductase-ligand complex,[65] with changes in this complex being signs of
trimethoprim resistance. Interestingly, the eight replicates with mutations that potentially lead
to increases in cellular NADH/NADPH are also the replicates with increased copy numbers of
dfrA17 (Figure 3A), suggesting that the mutations in *nadR, adhE* and *pfkA* are co-adaptations
correlated with increases in resistance gene copy numbers of replicates evolved in the
spatially structured environment.

Additionally, mutations within prophages were observed in the mutant *E. coli* chromosomes
(Table S1). Especially in the spatially structured environments SNPs and complex insertions
occurred. These mutations were excluded in the primary analysis of mutational patterns in this
study, as phage activity was not followed during the evolutionary experiment.

We observed more evolutionary parallelism at the COG-level, in accordance with previous
research (Figure 4B).[50,66] Again, most similarities are found within the replicates evolved in the
spatially structured environments. For these groups of replicates, either in the presence or
absence of the community, it can be observed that evolution within these environments lead
to both parallel genotypic (Figure 4) and parallel phenotypic (Figure 1AB) adaptation. In
contrast, we do not find genomic parallelism corresponding to the parallel phenotypic
adaptation of lineages evolved within the well-mixed environment, in the presence of the
community. This defeats our expectation that there would be more parallelism in these
lineages evolved in the well-mixed environments, due to stronger selective forces and larger
population sizes. Furthermore we find that genetic adaptations, at different levels of
organization, such as on the integron cassette harboring resistance genes, and the genomic
background, potentially to accommodate the CNV changes in the integron resistance
cassette,have played a role in the adaptive process. Yet, the observed canalization of
antibiotic resistance phenotypes could not entirely be explained by the genetic substitutions

only, indicating the importance of another level of interaction, namely the microbial community.
This shows multiple levels of genetic and ecological organization dictate antibiotic resistance
evolution.

Discussion

We observed that the presence of a community has a protective effect on *E. coli* at the start
of the antibiotic resistance evolution experiment. This effect decreases with increasing
antibiotic concentrations. The presence of the community leads to a faster evolutionary
response in both the well-mixed liquid as well as spatially structured solid environments. This
corresponds to previous findings in which the presence of other bacteria can speed up and
canalize evolution.[14] We observed phenotypic parallelism in each environmental context,
either community or environmental structure. For the spatially structured environment, the
phenotypic parallelism is linked with genotypic parallelism at the genome level. Yet, for the
well-mixed environment, the phenotypic canalization in the presence of the community cannot
solely be explained by genomic changes, showing that phenotypic plasticity likely played an
important role in the evolutionary response to antibiotics.

The beneficial effect of the community on *E. coli* at sub-MIC levels has been observed
before.[8,67] Yet, we also observed that the beneficial effect on the fitness of *E. coli* continued
even after antibiotic concentrations reached levels above initial MIC. The AUM environments
are relatively nutrient poor, and are therefore expected to have room for positive social
interactions among microbes, potentially leading to a longer sustained beneficial effect.[68,69] In
the spatially structured environment, the community does not have a large impact on the
evolved MIC of *E. coli*. Yet, the MBC increased when the *E. coli* evolved in isolation was
assessed in the presence of the community. The beneficial effect of the community is therefore
still present even for lineages with high levels of resistance, with the exception of *E. coli*
evolved in the presence of the community in the well-mixed environment. This shows that the
beneficial effect that the community confers is not dependent on the antibiotic concentration
present or the fitness impact of the antibiotics on *E. coli*, but instead is dependent on the
evolutionary history of the *E. coli*, particularly whether it evolved within that community before.

The initial cell density in environments with antibiotics is important for evolutionary
outcomes.[16,67] In both tested environments we observed higher densities of *E. coli* when
embedded in the community, leading to faster evolutionary rates towards high antibiotic
resistance. This corresponds to the well-known inoculum effect.[15,16] Perhaps surprisingly, for
the *E. coli* evolved on the spatial structure in the presence of the community, this did not lead
to a higher MIC compared to the *E. coli* evolved in isolation (Figure 1A,B). [8]Because the MIC

was determined at equal population sizes of *E. coli*, this assay does not account for differences in population size during the serial transfer experiments .[15] Still, indirectly, population size may play a role in reaching higher MICs, because the *E. coli* with higher MICs reach larger population sizes over 24 hours of growth in the absence of antibiotic. This suggests that adaptation to the medium-environment contributes to the growth at higher antibiotic concentrations.

[870]Spatial adaptation and the physical distance between interacting *E. coli* and community members may play a role in the adaptation to community members. We followed the colony formation in the different agar compartments (Fig S4), and did not observe any mixing of colonies on that surface. From literature it is known that bacterial mobility on agar can lead to a decrease in adaptation through for example increased ecological competition.[70,71] Moreover, studies suggest that interactions only play a role on small scales and distances in spatially structured environments,[72–74] while larger distance interactions play a role on chemotaxis of bacteria.[75] These results show the difficulty of comparing spatially structured and well-mixed environments, as distance dependent interactions are lost when the environment is well-mixed.

To observe potential variation in the evolved replicates, we sequenced three clones per replicate, for each of these clones we analyzed the CNVs of resistance genes and other genomic changes. We also noticed many prophage mutations, particularly in the spatially evolved lineages. We chose to exclude these from our H-index analysis because prophage regions often evolve at different rates compared to the core bacterial genome and undergo decay after integration[76], and this study aimed to compare mutations in the genome of *E. coli* and its plasmid, which is more representative of the species' overall evolutionary trajectory.

We observed a higher number of SNPs in clones derived from the spatially structured environment compared to the well-mixed environment, this was expected due to a larger effect of drift in the spatial environment.[77] We also observed variability in the CNVs in some replicates, for example replicate 4 in the well-mixed community environment (Figure S3), seemed to have lost the plasmid containing the resistance genes, even though these replicates showed high MICs. We cannot exclude that they lost their plasmids during culturing for sequencing as that environment did not contain antibiotics. If this were the case, then the copy number variations of the transposon on the plasmid a highly variable in time. It is also possible that the observation of plasmid loss is due to high plasmid copy number segregation variability, which refers to uneven distribution of plasmids during cell division and leads to higher phenotypic heterogeneity, and has been observed before.[78] Moreover, insertion-sequence activity and the implied mobility of the contained resistance genes could offer an

additional explanation for differences in the variation and observed efficacy of the resistance

genes. While gene copy number increases are a useful proxy for increased gene activity, gene

expression is key. Gene expression is influenced considerably by the location of the genes

within a genome and their alignment with factors regulating gene expression, such as

promoters. While our quantitative analysis of clonal sequencing data allows us to determine

gene copy numbers, it does not allow us to determine the location of any of the underlying

gene copies with the plasmid or host genome, and we therefore are very limited in predicting

the actual effect of the resistance gene copies. The potential presence of resistance plasmids

should be kept in mind when using sequencing techniques of clinical isolates to detect and

identify AMR genes, and observing where insertion sequences insert will be valuable in better

understanding antibiotic resistance evolution in pathogens.

Our investigation into the observed fluctuations in copy numbers of the plasmid carrying the

resistance gene, raises intriguing questions about the nuanced dynamics of antibiotic

resistance in microbial populations. Previous studies have underscored the complexity of

plasmid replication and maintenance, suggesting that various factors, including selective

pressures and growth rates, can influence the stability of plasmid copy numbers.[79,80] The

disconnect between plasmid copy numbers and MICs in our experimental context echoes

findings in other systems, where researchers have reported instances of decoupling between

plasmid abundance and phenotypic resistance[81,82]. These observations challenge

conventional assumptions regarding the straightforward relationship between plasmid

replication and antibiotic resistance expression. It is plausible that additional regulatory

mechanisms, such as transcriptional or post-transcriptional control, may contribute to the

observed variability in resistance phenotypes, such as the genomic compensatory mutations

related to NADH observed in the spatially-structured environment. This underscores the need

for comprehensive investigations into the multifaceted environmental structure, community

and genetic interactions governing plasmid-mediated resistance, which could provide valuable

insights for the development of targeted strategies to mitigate the spread of antibiotic

resistance in microbial communities.

Materials and methods

Bacterial isolates and growth medium

The focal *Escherichia coli* isolate and three community isolates (*Klebsiella pneumoniae*,

Enterococcus faecium, and *Staphylococcus epidermidis*) were collected anonymously in a

previous study from elderly patients (>70 years old) who were suffering from a urinary tract

infection.[37] The three community members were picked based on their ability to grow in

Lysogeny Broth (LB) containing 250 µg/mL trimethoprim and 1250 µg/mL sulfamethoxazole,
and being able to clearly distinguish all four isolates on CHROMagar Orientation
(CHROMagar). The *E. coli* isolate was selected based on its sensitivity to
trimethoprim/sulfamethoxazole. Pure cultures of each separate isolate were obtained by
streaking on CHROMagar plates, selecting a single colony, and growing overnight in LB broth
at 37°C. Overnight cultures were used to create glycerol stocks (25% glycerol v/v) which were
subsequently stored at -80°C. The experiments were performed in Artificial Urine Medium
(AUM), which is created (for a 1X concentration) by mixing 1.5 g/L bacto peptone L37 (BD),
3.15 g/L NaHCO3, 11.25 g/L urea (Roth), 4.8 g/L Na2SO4.10H2O, 1.8 g/L K2HPO4, 1.95 g/L
NH4Cl, 15 mg/L bacto yeast extract (BD), 1.98 mM lactic acid (Roth), 600 mg/L citric acid, 105
mg/L uric acid, 1.2 g/L creatinine, 4.44 mg/L CaCl2.2H2O, 1.8 g/L Fe(II)SO4.7H2O (Riedel),
368 mg/L MgSO4.7H2O, 1.43 g/L KH2PO4. Chemicals were ordered from Sigma, unless
stated otherwise. The pH was adjusted to 6.5 using 1M HCl. For 2x AUM, concentrations of
all chemicals were doubled except NaCl, which would be omitted.
578

Antibiotic and X-gal stocks

The trimethoprim stock was created by mixing 10 mg/mL trimethoprim in dimethyl sulfoxide
(DMSO), and the sulfamethoxazole stock was created by mixing 50 mg/mL sulfamethoxazole
in acetone. The X-gal stock contained 20 mg/mL X-Gal and 0.1M Isopropyl β-D-1-
thiogalactopyranoside (IPTG) mixed in DMSO. Both antibiotics, the X-gal, IPTG and solvents
were ordered from Sigma. The trimethoprim stock was stored at 4°C for up to two weeks, and
the X-gal stock and sulfamethoxazole stocks were stored at -20°C. Trimethoprim and
sulfamethoxazole were always mixed in a 1:5 ratio, similar to clinical settings.[38]

Bacterial hotel creation and transfer protocol

Silicon hotel agar molds were stored in 70% ethanol and were dried in a sterile flow cabinet
before pouring agar. Each agar hotel was created by mixing 12.5 mL 2x AUM, an appropriate
amount of antibiotics, 12.5 mL 3% soft agar in demineralized water at 60°C, and 125 µL X-
Gal stock for the visualization of *E. coli* into the hotel mold. Agar hotels were cooled for 30
minutes before removing the agar from the molds.

Five replicates of *E. coli* were evolved for 60 bi-daily transfers in communities with *K.*
pneumoniae, *E. faecium* and *S. epidermidis,* and in isolation. Starting cultures of all isolates
were created by growing each separately in 1x AUM overnight at 37°C shaken at 200 rpm.
Each culture was diluted a hundred times in 1x AUM. Starting agar hotels were created with
the starting antibiotics concentration of 50:250 µg/mL TMP:SMX. Following the layout of
Figure S4, 0.5 µL of each diluted culture was added to the middle of the appropriate well on

the agar plate. Added cultures were allowed to dry for five minutes before incubating the agar hotels at 37°C for 48 hours. We did not observe any spillover of colonies to other compartments.

Before transferring *E. coli* to a fresh agar hotel, the colonies were visibly inspected for growth. If the colony's growth exceeded 1mm of the starting colony, the antibiotics concentration would be increased by 20% unless the antibiotics concentration was increased during the previous transfer. Exception here were the first two transfers where antibiotics were increased by 20% due to clear overgrowth of *E. coli*. Replicates were transferred by streaking around the outer edges of each *E. coli* colony five times and adding the bacteria that were resuspended in 50 µL of 1x AUM. After mixing, 0.5 µL of each replicate would be added to the fresh agar hotels. Wild type community members (non-*E. coli*) were always added as hundred times diluted overnight culture in 1x AUM to the fresh agar hotels.

Liquid serial transfers protocol

E. coli was evolved over 120 days in AUM transferring replicates to fresh 5 mL medium every 48 hours for 60 transfers in total. To start, the *E. coli* and the three community members were streaked on agar to obtain a pure culture of a single colony that was cultured overnight in AUM at 37°C, shaken at 200 rpm. Isolates were diluted a hundred times by adding 50 µL to 5mL total preheated medium. Five replicates of *E. coli* in isolation and five replicates of *E. coli* with the three community members *K. pneumoniae, E. faecium*, and *S. epidermidis*. An appropriate amount of the antibiotics were added to the medium before adding the bacteria. Antibiotic concentration started at 50:250 µg/mL TMP:SMX. Cultures were allowed to grow for 48 hours at 37°C and shaken at 200 rpm. During transfer, the previous culture was diluted a hundred times and community members were added fresh. Remaining culture was used to do colony counting measurements on CHROMagar plates to obtain a CFU/mL count for *E. coli*. If *E. coli* had exceeded $1*10^6$ CFU/mL, antibiotic concentration would be increased by 20% for the next transfer. For the first two transfers for *E. coli* evolving in presence of the community, the antibiotic concentration was increased by 20% immediately due to observed high population growth based on turbidity.

MIC and MBC assays

MIC of replicates were measured in liquid using a two times serial dilution of antibiotics method. Using the concentration of antibiotics at the final transfer step as a guideline, a dilution series was created for each isolate specifically. Measurements were performed in 96-well microtiter plates, using AUM as growth medium. Stationary phase cultures of *E. coli* replicates were used for experiments, diluting them 40x times in the medium antibiotics' mixture. Plates were incubated overnight at 37°C, shaken at 200 rpm. Growth was measured through

absorbance at 600nm using a BMG CLARIOstar microplate reader. If the absorbance with the

background subtracted exceeded 0.1, it was assumed that *E. coli* had grown using the lowest

concentration of antibiotics where no growth was observed is the MIC. All MIC measurements

were performed in triplicate.

MBC assays with community present were performed following the above protocol, but with

community members added together with the *E. coli* before overnight growth. After OD600

measurements, presence of *E. coli* was confirmed using a 96-well pinning tool on

CHROMagar. The lowest concentration of antibiotics where no growth of E.coli on

CHROMagar was observed is the MBC.

DNA extraction and sequencing

DNA from first transfer clones and final transfer mutants were extracted using the QIAamp

DNA Mini Kit (Qiagen, Hilden, Germany) following the manufacturer's instructions. DNA was

eluted in 100 µL elution buffer AE. Quality and quantity of isolated DNA were checked using

a Nanodrop One^C Microvolume UV-Vis Spectrophotometer (Thermo Fisher Scientific, USA).

Three clones of the final transfer step for each of the five replicate lineages per experimental

setting were sent for illumina sequencing at SeqCenter (Pittsburgh, PA, USA). Five replicates

of community members isolated from *E. coli* community agar and liquid were also sent to

SeqCenter, Pittsburgh, USA for Illumina sequencing.

Illumina sequencing libraries were prepared using the tagmentation-based and PCR-based

Illumina DNA Prep kit and custom IDT 10bp unique dual indices (UDI) with a target insert size

of 320 bp. No additional DNA fragmentation or size selection steps were performed. Illumina

sequencing was performed on an Illumina NovaSeq 6000 sequencer in one or more

multiplexed shared-flow-cell runs, producing 2x151bp paired-end reads. Demultiplexing,

quality control and adapter trimming was performed with bcl-convert1 (v4.1.5). In total 10.9

Gbp and 359.2 Gbp of read data were produced from ancestral and evolved strains,

respectively.

Ancestor strains were sent to the same company for Nanopore sequencing. Sample libraries

were prepared using Oxford Nanopore Technologies (ONT) Ligation Sequencing Kit (SQK-

NBD114.24) with NEBNext® Companion Module (E7180L) to manufacturer's specifications.

All samples were run on Nanopore R10.4.1 flow cells and MinION Mk1B device. Post-

sequencing, Guppy (v6.3.8) was used for super high accuracy base calling (SUP) and

demultiplexing. Overall, 2.9 Gbp of read data were produced.

Genome assembly, annotation variant calling and effect prediction

The wildtype reference assembly was generated with flye v2.9.2 (settings: --nano-hq)[83] from

the ancestral nanopore data and polished using ancestral Illumina read with polca.sh included

in the MaSuRCA v4.1.0 Genome Assembly and Analysis Toolkit.[84] Genes were annotated

with Bakta v1.8.1[85] and integrated prophages and plasmids with genomad v1.7.4[86] for each

replicate were called from read mappings generated with minimap2 v2.26[87] using freebayes

v1.3.1 (settings: -p 1 -0 -F 0.5)[88] and filtered for spurious calls with bcftools v1.17 (--exclude

'REF~"N" || QUAL < 20 || INFO/DP < 10').[89] Variant effects were predicted with the Ensembl

Variant Effect Predictor (VEP) v110.1 command line tool.

H-index calculation

To quantify the repeatability of mutations within our *E. coli* mutants, the *H*-index of genome

repeatability previously described by Schenk et al. was calculated.[50] The *H*-index was

calculated on the functional (COG) and at the gene level. All mutant replicates were compared

pairwise, each one-way comparison calculated using:

$$H_{A,B} = \sum_{j=1}^{m} \sum_{k=1}^{n} \left(\frac{|A_j \cap B_k|}{|A_j|} \right) / \sum_{j=1}^{m} \sum_{k=1}^{m} \left(\frac{|A_j \cap A_l|}{|A_j|} \right) \quad (1)$$

In the formula (1), genotype A and B with mutations m and n respectively are compared. The

final *H*-index of the combination is obtained for both genotypes separately ($H_{A,B}$ and $H_{B,A}$) and

compared using:

$$H = \frac{H_{A,B} + H_{B,A}}{2} \quad (2)$$

The final *H*-index value falls between 0 (no repeated evolution) and 1 (complete parallelism).

An *H*-index value of 0.5 indicates that the average of the two compared mutants have half of

their mutations shared between them.

Statistical analysis

When comparing the values between two conditions of mutants, a two-tailed *Welch's t*-test

was performed to test the null hypothesis that the two parameters do not have significantly

different values (hypothesis of neutrality). To determine level of correlation between two

variables, we used the square of a Pearsson correlation coefficient and resulting *P*-value

where the null-hypothesis was rejected with a significance level above 0.05. For multiple

pairwise comparisons, we used independent sample *t*-tests for all sets with a Bonferroni-

correction to adjust *P*-values to reduce type I errors.

Author contribution

Conceived and designed the study: LEZ MGJDV. Performed the experiments: LEZ MB JML. Analyzed the data: LEZ TH MGJ. Wrote the manuscript: LEZ JML TH MGJ.

Acknowledgements

LEZ was supported by the Adaptive Life program of the Faculty of Science and Engineering, within GELIFES. MB was supported by the Faculty of Science and Engineering - Adaptive Life PhD Scholarship from the University of Groningen, awarded by GELIFES.

References

Byrd AL, Segre JA. Adapting Koch's postulates: Criteria for disease causation must take microbial interactions into account. *Science (1979)* 2016; **351**: 224–226.

Peters BM, Jabra-Rizk MA, O'May GA, William Costerton J, Shirtliff ME. Polymicrobial interactions: Impact on pathogenesis and human disease. *Clin Microbiol Rev* 2012; **25**: 193–213.

Gaston JR, Johnson AO, Bair KL, White AN, Armbruster CE. Polymicrobial interactions in the urinary tract: Is the enemy of my enemy my friend? *Infect Immun* 2021; **89**. doi:10.1128/IAI.00652-20.

Zandbergen LE, Halverson T, Brons JK, Wolfe AJ, de Vos MGJ. The Good and the Bad: ecological interaction measurements between the urinary microbiota and uropathogens. *Front Microbiol* 2021; **12**: 659450.

Short FL, Murdoch SL, Ryan RP. Polybacterial human disease: The ills of social networking. *Trends Microbiol* 2014; **22**: 508–516.

Vega NM, Gore J. Collective antibiotic resistance: Mechanisms and implications. *Curr Opin Microbiol* 2014; **21**: 28–34.

Bottery MJ, Pitchford JW, Friman VP. Ecology and evolution of antimicrobial resistance in bacterial communities. *ISME Journal* 2021; **15**: 939–948.

Klümper U, Recker M, Zhang L, Yin X, Zhang T, Buckling A *et al.* Selection for antimicrobial resistance is reduced when embedded in a natural microbial community. *ISME Journal* 2019; **13**: 2927–2937.

Algammal A, Hetta HF, Mabrok M, Behzadi P. Editorial: Emerging multidrug-resistant bacterial pathogens "superbugs": A rising public health threat. *Front Microbiol* 2023; **14**. doi:10.3389/fmicb.2023.1135614.

De Vos MGJ, Zagorski M, McNally A, Bollenbach T. Interaction networks, ecological stability, and collective antibiotic tolerance in polymicrobial infections. *Proc Natl Acad Sci U S A* 2017; **114**: 10666–10671.

Geyrhofer L, Ruelens P, Farr AD, Pesce D, de Visser JAGM, Brenner N. Minimal surviving inoculum in collective antibiotic resistance. *mBio* 2023; **14**. doi:10.1128/mbio.02456-22.

12 Biswas L, Biswas R, Schlag M, Bertram R, Götz F. Small-colony variant selection as a
survival strategy for Staphylococcus aureus in the presence of *Pseudomonas*
aeruginosa. *Appl Environ Microbiol* 2009; **75**: 6910–6912.

13 Brauner A, Fridman O, Gefen O, Balaban NQ. Distinguishing between resistance,
tolerance and persistence to antibiotic treatment. *Nat Rev Microbiol* 2016; **14**: 320–
330.

14 Zandbergen LE, van den Heuvel J, Farr AD, Zwaan BJ, de Visser JAGM, de Vos
MGJ. Microbial interactions affect the tempo and mode of antibiotic resistance
evolution. *bioRxiv* 2024; 2024.06.06.597700.

15 Brook I. Inoculum Effect. *Rev Infect Dis* 1989; **11**: 361–368.

16 Tan C, Phillip Smith R, Srimani JK, Riccione KA, Prasada S, Kuehn M *et al.* The
inoculum effect and band-pass bacterial response to periodic antibiotic treatment. *Mol
Syst Biol* 2012; **8**. doi:10.1038/msb.2012.49.

17 Foster KR, Bell T. Competition, not cooperation, dominates interactions among
culturable microbial species. *Current Biology* 2012; **22**: 1845–1850.

18 Lee J-H, Lee J. Indole as an intercellular signal in microbial communities. *FEMS
Microbiol Rev* 2010; **34**: 426–444.

19 Flemming HC, Wingender J, Szewzyk U, Steinberg P, Rice SA, Kjelleberg S. Biofilms:
An emergent form of bacterial life. *Nat Rev Microbiol* 2016; **14**: 563–575.

20 Costerton JW, Stewart PS, Greenberg. Bacterial biofilms: a common cause of
persistant infections. *Science (1979)* 1999; **284**: 1318–1322.

21 Dieckmann U, Law R, Metz JAJ. *The Geometry of Ecological Interactions: Simplifying
Spatial Complexity*. Cambridge University Press, 2002 doi:10.1016/s0925-
8574(01)00102-1.

22 Tilman D, Kareiva P. *Spatial Ecology: The Role of Space in Population Dynamics and
Interspecific Interactions. David Tilman , Peter Kareiva*. Princeton University Press,
1998 doi:10.1086/420507.

23 Johnson CR, Seinen I. Selection for restraint in competitive ability in spatial
competition systems. *Proceedings of the Royal Society B: Biological Sciences* 2002;
269: 655–663.

24 Kerr B, Riley MA, Feldman MW, Bohannan BJM. Local dispersal promotes
biodiversity in a real-life game of rock-paper-scissors. *Nature* 2002; **418**: 171–174.

25   Yanni D, Márquez-Zacarías P, Yunker PJ, Ratcliff WC. Drivers of Spatial Structure in
Social Microbial Communities. *Current Biology* 2019; **29**: R545–R550.

26   Evans KC, Benomar S, Camuy-Vélez LA, Nasseri EB, Wang X, Neuenswander B *et*
al. Quorum-sensing control of antibiotic resistance stabilizes cooperation in
Chromobacterium violaceum. *ISME J* 2018; **12**: 1263–1272.

27   Chuang SK, Vrla GD, Fröhlich KS, Gitai Z. Surface association sensitizes
Pseudomonas aeruginosa to quorum sensing. *Nat Commun* 2019; **10**: 4118.

28   Nowak MA, May RM. Evolutionary games and spatial chaos. *Nature* 1992; **359**: 826–
829.

29   Santos FC, Pacheco JM, Lenaerts T. Evolutionary dynamics of social dilemmas in
structured heterogeneous populations. *Proc Natl Acad Sci U S A* 2006; **103**: 3490–
3494.

30   Nadell CD, Drescher K, Foster KR. Spatial structure, cooperation and competition in
biofilms. Nat Rev Microbiol. 2016; **14**: 589–600.

31   Verbruggen E, El Mouden C, Jansa J, Akkermans G, Bücking H, West SA *et al.*
Spatial structure and interspecific cooperation: Theory and an empirical test using the
mycorrhizal mutualism. *American Naturalist* 2012; **179**: E133-46.

32   Bailey SF, Trudeau A, Tulowiecki K, McGrath M, Belle A, Fountain H *et al.* Spatial
structure affects evolutionary dynamics and drives genomic diversity in experimental
populations of *Pseudomonas fluorescens*. 2021. doi:10.1101/2021.09.28.461808.

33   Lobanov A, Dyckman S, Kurkjian H, Momeni B. Spatial structure favors microbial
coexistence except when slower mediator diffusion weakens interactions. *Elife* 2023;
12. doi:10.7554/eLife.82504.

34   Momeni B, Waite AJ, Shou W. Spatial self-organization favors heterotypic cooperation
over cheating. *Elife* 2013; **2**. doi:10.7554/eLife.00960.

35   Habets MGJL, Rozen DE, Hoekstra RF, De Visser JAGM. The effect of population
structure on the adaptive radiation of microbial populations evolving in spatially
structured environments. *Ecol Lett* 2006; **9**: 1041–1048.

36   Perfeito L, Pereira MI, Campos PR, Gordo I. The effect of spatial structure on
adaptation in Escherichia coli. *Biol Lett* 2008; **4**: 57–59.

37   Croxall G, Weston V, Joseph S, Manning G, Cheetham P, McNally A. Increased
human pathogenic potential of Escherichia coli from polymicrobial urinary tract

infections in comparison to isolates from monomicrobial culture samples. *J Med*
Microbiol 2011; **60**: 102–109.

38 Hitchings GH. Mechanism of action of trimethoprim-sulfamethoxazole--I. *J Infect Dis*
1973; **128**: S433.

39 Mann R, Mediati DG, Duggin IG, Harry EJ, Bottomley AL. Metabolic adaptations of
Uropathogenic E. coli in the urinary tract. *Front Cell Infect Microbiol* 2017; **7**.
doi:10.3389/fcimb.2017.00241.

40 Webb CO, Ackerly DD, McPeek MA, Donoghue MJ. Phylogenies and community
ecology. *Annu Rev Ecol Syst* 2002; **33**: 475–505.

41 Geisinger E, Isberg RR. Antibiotic Modulation of Capsular Exopolysaccharide and
Virulence in Acinetobacter baumannii. *PLoS Pathog* 2015; **11**: e1004691.

42 Santos-Lopez A, Marshall CW, Scribner MR, Snyder DJ, Cooper VS. Evolutionary
pathways to antibiotic resistance are dependent upon environmental structure and
bacterial lifestyle. *Elife* 2019; **8**. doi:10.7554/elife.47612.

43 Brockhurst MA, Habets MGJL, Libberton B, Buckling A, Gardner A. Ecological drivers
of the evolution of public-goods cooperation in bacteria. *Ecology* 2010; **91**: 334–340.

44 Tamer YT, Gaszek IK, Abdizadeh H, Batur TA, Reynolds KA, Atilgan AR *et al.* High-
order epistasis in catalytic power of dihydrofolate reductase gives rise to a rugged
fitness landscape in the presence of trimethoprim selection. Mol Biol Evol. 2019; **36**:
1533–1550.

45 Vedantam G, Guay GG, Austria NE, Doktor SZ, Nichols BP. Characterization of
mutations contributing to sulfathiazole resistance in Escherichia coli. *Antimicrob*
Agents Chemother 1998; **42**: 88–93.

46 Liebert CA, Hall RM, Summers AO. Transposon Tn 21 , Flagship of the Floating
Genome . *Microbiology and Molecular Biology Reviews* 1999; **63**: 507–522.

47 San Millan A, MacLean RC. Fitness Costs of Plasmids: a Limit to Plasmid
Transmission. *Microbiol Spectr* 2017; **5**. doi:10.1128/microbiolspec.mtbp-0016-2017.

48 Dimitriu T, Matthews AC, Buckling A. Increased copy number couples the evolution of
plasmid horizontal transmission and plasmid-encoded antibiotic resistance. *Proc Natl*
Acad Sci U S A 2021; **118**: e2107818118.

49   Hallatschek O, Hersen P, Ramanathan S, Nelson DR. Genetic drift at expanding
frontiers promotes gene segregation. *Proc Natl Acad Sci U S A* 2007; **104**: 19926–
19930.

50   Schenk MF, Zwart MP, Hwang S, Ruelens P, Severing E, Krug J *et al.* Population size
mediates the contribution of high-rate and large-benefit mutations to parallel evolution.
Nat Ecol Evol. 2022; **6**: 439–447.

51   Buwembo W, Aery S, Rwenyonyi CM, Swedberg G, Kironde F. Point mutations in the
folP gene partly explain sulfonamide resistance of streptococcus mutans. *Int J*
Microbiol 2013; **2013**: 367021.

52   Rimmele M, Boos W. Trehalose-6-phosphate hydrolase of Escherichia coli. *J*
Bacteriol 1994; **176**: 5654–5664.

53   Botelho J, Schulenburg H. The Role of Integrative and Conjugative Elements in
Antibiotic Resistance Evolution. *Trends Microbiol* 2021; **29**: 8–18.

54   Amro J, Baron C. TraC of the conjugative plasmid pKM101 is secreted as a monomer
in outer membrane vesicles and stimulates conjugation. *bioRxiv* 2022; : 2022–2027.

55   Nakayama J, Takanami Y, Horii T, Sakuda S, Suzuki A. Molecular Mechanism of
Peptide-Specific Pheromone Signaling in *Enterococcus faecalis* : Functions of
Pheromone Receptor TraA and Pheromone-Binding Protein TraC Encoded by
Plasmid pPD1. *J Bacteriol* 1998; **180**: 449–456.

56   Cho H, Winans SC. TraA, TraC and TraD autorepress two divergent quorum-
regulated promoters near the transfer origin of the Ti plasmid of *Agrobacterium*
tumefaciens. *Mol Microbiol* 2007; **63**: 1769–1782.

57   Parvathy ST, Udayasuriyan V, Bhadana V. Codon usage bias. *Mol Biol Rep* 2022; **49**:
539–565.

58   Mengin-Lecreulx D, van Heijenoort J, Park JT. Identification of the mpl gene encoding
UDP-N-acetylmuramate: L-alanyl-gamma-D-glutamyl-meso-diaminopimelate ligase in
Escherichia coli and its role in recycling of cell wall peptidoglycan. *J Bacteriol* 1996;
178: 5347–5352.

59   Vinopal RT, Clifton D, Fraenkel DG. PfkA locus of Escherichia coli. *J Bacteriol* 1975;
122: 1162–1171.

60 Siedler S, Bringer S, Bott M. Increased NADPH availability in *Escherichia coli*:
improvement of the product per glucose ratio in reductive whole-cell
biotransformation. *Appl Microbiol Biotechnol* 2011; **92**: 929–937.

61 Gerasimova AV, Gelfand MS. Evolution of the Nadr regulon in enterobacteriaceae. *J*
Bioinform Comput Biol 2005; **03**: 1007–1019.

62 Tritz GJ, Chandler JL. Recognition of a Gene Involved in the Regulation of
Nicotinamide Adenine Dinucleotide Biosynthesis. *J Bacteriol* 1973; **114**: 128–136.

63 Arce-Rodríguez A, Pankratz D, Preusse M, Nikel PI, Häussler S. Dual Effect: high
NADH levels contribute to efflux-mediated antibiotic resistance but drive lethality
mediated by reactive oxygen species. *mBio* 2022; **13**. doi:10.1128/mbio.02434-21.

64 Kessler D, Leibrecht I, Knappe J. Pyruvate-formate-lyase-deactivase and acetyl-CoA
reductase activities of *Escherichia coli* reside on a polymeric protein particle encoded
by *adhE*. *FEBS Lett* 1991; **281**: 59–63.

65 Krucinska J, Lombardo MN, Erlandsen H, Estrada A, Si D, Viswanathan K *et al.*
Structure-guided functional studies of plasmid-encoded dihydrofolate reductases
reveal a common mechanism of trimethoprim resistance in Gram-negative pathogens.
Commun Biol 2022; **5**: 459.

66 Tenaillon O, Rodríguez-Verdugo A, Gaut RL, McDonald P, Bennett AF, Long AD *et al.*
The molecular diversity of adaptive convergence. *Science (1979)* 2012; **335**: 457–
461.

67 Yurtsev EA, Chao HX, Datta MS, Artemova T, Gore J. Bacterial cheating drives the
population dynamics of cooperative antibiotic resistance plasmids. *Mol Syst Biol* 2013;
9: 683.

68 Piccardi P, Vessman B, Mitri S. Toxicity drives facilitation between 4 bacterial species.
Proceedings of the National Academy of Sciences 2019; **116**: 15979–15984.

69 Adamowicz EM, Muza M, Chacón JM, Harcombe WR. Cross-feeding modulates the
rate and mechanism of antibiotic resistance evolution in a model microbial community
of Escherichia coli and Salmonella enterica. *PLoS Pathog* 2020; **16**: e1008700.

70 Butler MT, Wang Q, Harshey RM. Cell density and mobility protect swarming bacteria
against antibiotics. *Proc Natl Acad Sci U S A* 2010; **107**: 3776–3781.

71 Piskovsky V, Oliveira NM. Bacterial motility can govern the dynamics of antibiotic
resistance evolution. *Nat Commun* 2023; **14**: 5584.

72   Keogh D, Tay WH, Ho YY, Dale JL, Chen S, Umashankar S *et al.* Enterococcal
Metabolite Cues Facilitate Interspecies Niche Modulation and Polymicrobial Infection.
Cell Host Microbe 2016; **20**: 493–503.

73   Co AD, van Vliet S, Kiviet DJ, Schlegel S, Ackermann M. Short-range interactions
govern the dynamics and functions of microbial communities. *Nat Ecol Evol* 2020; **4**:
366–375.

74   van Tatenhove-Pel RJ, Rijavec T, Lapanje A, van Swam I, Zwering E, Hernandez-
Valdes JA *et al.* Microbial competition reduces metabolic interaction distances to the
low µm-range. *ISME Journal* 2021; **15**: 688–701.

75   Stocker R, Seymour JR. Ecology and Physics of Bacterial Chemotaxis in the Ocean.
Microbiology and Molecular Biology Reviews 2012; **76**: 792–812.

76   Canchaya C, Proux C, Fournous G, Bruttin A, Brüssow H. Prophage genomics.
Microbiol Mol Biol Rev 2003; **67**: 238–76, table of contents.

77   Zhang Q, Lambert G, Liao D, Kim H, Robin K, Tung CK *et al.* Acceleration of
emergence of bacterial antibiotic resistance in connected microenvironments. *Science*
(1979) 2011; **333**: 1764–1767.

78   Hernandez-Beltran JCR, Rodríguez-Beltrán J, Aguilar-Luviano OB, Velez-Santiago J,
Mondragón-Palomino O, MacLean RC *et al.* Plasmid-mediated phenotypic noise
leads to transient antibiotic resistance in bacteria. *Nat Commun* 2024; **15**: 2610.

79   Baltrus DA. Exploring the costs of horizontal gene transfer. *Trends Ecol Evol* 2013;
28: 489–495.

80   Millan AS, Santos-Lopez A, Ortega-Huedo R, Bernabe-Balas C, Kennedy SP,
Gonzalez-Zorn B. Small-plasmid-mediated antibiotic resistance is enhanced by
increases in plasmid copy number and bacterial fitness. *Antimicrob Agents*
Chemother 2015; **59**: 3335–3341.

81   Bouma JE, Lenski RE. Evolution of a bacteria/plasmid association. *Nature* 1988; **335**:
351–352.

82   Stokes HW, Gillings MR. Gene flow, mobile genetic elements and the recruitment of
antibiotic resistance genes into Gram-negative pathogens. *FEMS Microbiol Rev* 2011;
35: 790–819.

83   Kolmogorov M, Yuan J, Lin Y, Pevzner PA. Assembly of long, error-prone reads using
repeat graphs. *Nat Biotechnol* 2019; **37**: 540–546.

84   Zimin A V., Puiu D, Luo M-C, Zhu T, Koren S, Marçais G *et al.* Hybrid assembly of the
929   large and highly repetitive genome of *Aegilops tauschii* , a progenitor of bread wheat,
930   with the MaSuRCA mega-reads algorithm. *Genome Res* 2017; **27**: 787–792.

931   85   Schwengers O, Jelonek L, Dieckmann MA, Beyvers S, Blom J, Goesmann A. Bakta:
932   rapid and standardized annotation of bacterial genomes via alignment-free sequence
933   identification. *Microb Genom* 2021; **7**. doi:10.1099/mgen.0.000685.

934   86   Camargo AP, Roux S, Schulz F, Babinski M, Xu Y, Hu B *et al.* Identification of mobile
935   genetic elements with geNomad. *Nat Biotechnol* 2024; **42**: 1303–1312.

936   87   Li H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* 2018;
937   **34**: 3094–3100.

938   88   Garrison E, Marth G. Haplotype-based variant detection from short-read sequencing.
939   2012.

940   89   Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO *et al.* Twelve years
941   of SAMtools and BCFtools. *Gigascience* 2021; **10**. doi:10.1093/gigascience/giab008.

Figure 1 | Phentoypic trajectories of trimethoprim-sulfamethoxazole antibiotic resistance evolution and population sizes of _E. coli_ over time. _E. coli_ evolved in the presence and absence of a community, consisting of _Klebsiella pneumoniae_, _Enterococcus faecium_ and _Staphylococcus epidermidis_. _E. coli_ was evolved in presence of trimethoprim-sulfamethoxazole (TMP-SMX) at a 1:5 ratio. Each condition was replicated five times. **AB,** concentration of TMP-SMX in the medium for all evolving replicate lineages within the well-mixed liquid environment (A) and spatially structured environment (B) over time. Concentrations of TMP-SMX are given by the trimethoprim concentrations. All five replicate lineages converged and overlapped for _E. coli_ evolving in the presence of the community, for both well-mixed spatially structured environments. Different trajectories can be observed for _E. coli_ evolving in the absence of the community within the well-mixed liquid environment. **C,** population size of _E. coli_ in the well-mixed liquid environment in CFU/mL. A decline of the _E. coli_ population size evolved in the presence of the community can be observed after fifty days, for all five replicate lineages. The population size of _E. coli_ evolving in isolation showed large fluctuations in the first eighty days, these became less diverse after this time. **D,** population size of _E. coli_ evolved in the spatially structured solid environment measured as colony area in pixels. A clear decline of colony size for _E. coli_ evolved in the presence of the community can be observed after roughly forty days for all five

Figure 2 | Relationship between the minimal inhibitory concentration (MIC) of evolved clones, in the presence and absence of the community, and the antibiotic concentrations at final transfer. The size of the data points shows the number of replicates, with the smallest data point being one replicate and the largest all five replicates. *E. coli* was evolved in presence of trimethoprim-sulfamethoxazole (TMP-SMX) at a 1:5 ratio. Concentrations of TMP-SMX are given by the trimethoprim concentrations. **A,** MIC of evolved *E.coli* versus the concentration of antibiotic at the final transfer step at day 120. No clear correlation between the MIC of evolved clones, and the concentration of antibiotic in the medium at their final transfer could be observed. **B,** MIC of evolved *E.coli* versus the community effect on MIC. The community resistance effect was determined by performing an MBC assay in the presence of the community, followed by subsequent transfer on agar. The concentration of trimethoprim-sulfamethoxazole at which no *E. coli* could be observed on the agar, is given relative to the MBC of *E. coli* in isolation. Therefore, a value above 1 on the y-axis indicates a higher observed resistance compared to measured MIC, while a value below 1 indicates a low resistance compared to the MIC. Only *E. coli* evolved in liquid medium in the presence of the community did not benefit from the presence of the community in the community MBC assay (blue). Absolute resistance levels in the presence of community are depicted in Figure S1.

3A

3B

Figure 3 | Copy number variation in the plasmid containing the antibiotic resistance genes in the final transfer mutants. Three clones were sequenced per replicate, showing genetic copy number variation within the population. **A.** Copy number variation in the TN21-like integron cassette part of the plasmid containing *dfrA17* and *sul1,* which confer resistance to trimethoprim and sulfamethoxazole, respectively, in each of the sequenced clones. The highlighted region in gray contains the resistance genes, showing overall a tendency for increased copy numbers of this region. Some replicates do not have increased copy numbers of the resistance genes compared to the wild type. Particularly those evolved in the liquid environment in the presence of the community (blue), and in the solid spatially structured environment in the presence of the community (yellow). **B.** Average copy number variation per replicate in the TN21-like integron cassette part of the plasmid containing *dfrA17* and *sul1* resistance genes versus resistance level (MIC) of evolved replicate lineages. Concentrations of trimethoprim-sulfamethoxazole (1:5 ratio) are given by the trimethoprim concentrations. No clear pattern between replicates evolved in the same environment could be observed. Copy number variation for each sequenced clone is depicted in Figure S3.

4A

4B

Figure 4 | Clustering and mutation repeatability of final replicates clustered on gene and COG level. The level of genetic parallelism in evolved clones is given by the *H*-index. Higher *H*-index values indicate a higher parallelism of the two genotypes compared. Three clones per replicate were sequenced per environment. Comparisons in the blocks around the diagonal in each figure show the intra-environment comparisons, while the other blocks show the inter-environment comparisons. Solid indicates the spatially structured environment and liquid indicates the well-mixed environment. Mutations and genes are shown in Supplementary table 1. **A,** *H*-index of mutation repeatability at the gene-level of *E. coli* mutants. Very few replicates within the well-mixed liquid environment showed repeated evolution, while within the spatially structured environment there was some repeated evolution. **B,** *H*-index of mutation repeatability at the COG-level. Most genetic parallelism is found in the spatially-structured environment, opposing the anticipated higher level of drift in the spatially structured versus well-mixed environments.

Graphical abstract