Title: Short chain fatty acids regulate the chromatin landscape and distinct gene expression changes in human colorectal cancer cells

Authors: Tohfa Kabir[1*], Charlotte A. Connamacher[1*], Zara Nadeem[2,3], Matthew R. Paul[4], Matthew R. Smutny[1], Zoe K. Lawler[1], Annaelle M. Djomo[2], Thomas S. Carroll[4], Leah A. Gates[1,2]

[1]Department of Biochemistry, Case Western Reserve University, Cleveland, OH
[2]Laboratory of Chromatin Biology & Epigenetics, The Rockefeller University, NY
[3]HunterCollege of the City University of New York, Yalow Honors Scholar Program, New York, NY, USA
[4]Bioinformatics Resource Center, The Rockefeller University, NY
[*]These authors contributed equally

Corresponding author: leah.gates@case.edu

Keywords: chromatin, epigenetics, short chain fatty acids, butyrate, gene regulation

Abstract

Short chain fatty acids (SCFAs) are small metabolites that are produced through the activity of microbes and have important roles in human physiology. These metabolites have varied mechanisms in interacting with the host, of which one such mode is decorating the chromatin landscape. We previously detected specific histone modifications in the mouse gut that can be derived from SCFAs and are regulated by the microbiota. However, the roles of these SCFAs on chromatin and how they impact gene regulation in human cells is largely unknown. Now, our studies demonstrate these microbiota-dependent histone posttranslational modifications (PTMs) are associated with alterations in gene regulation in human cells. We show that histone butyrylation on H3K27 is detected in human colon samples. Furthermore, histone acetylation, butyrylation, and propionylation on H3K9 and H3K27 are responsive to levels of SCFAs in human colon cancer cell lines and are associated with active gene regulatory elements. In addition, treatment of human cancer cell lines with individual metabolites or combinations of SCFAs replicating the intestinal lumen environment result in distinct and overlapping gene program changes, with butyrate largely driving gene regulatory effects of SCFA combinations. Lastly, we define butyrate effects on gene expression that are independent of HDAC inhibition and are dependent on p300/CBP, suggesting potential gene programs regulated by histone butyrylation. Together, these results demonstrate that SCFAs are key regulators of the chromatin landscape and gene programs in human colorectal cancer cells.

Introduction

In the gut, commensal microbes ferment fiber resulting in the accumulation of
high concentrations of the metabolites acetate, butyrate, and propionate in the intestinal
lumen[1,2]. These SCFAs act on the host epithelium through several different mechanisms
or "fates": for example, they can bind to cell surface or nuclear receptors to induce
signaling cascades and specific gene programs, and they can feed into host cell energy
pathways. Furthermore, SCFAs can impact the chromatin landscape both directly
through serving as donor molecules for histone posttranslational modifications and
indirectly by inhibiting select chromatin-modifying enzymes, such as histone
deacetylases (HDACs)[3–11]. SCFAs regulate multiple physiological systems including
neuronal health, intestinal inflammation, tumor development and metabolic processes
such as insulin sensitivity and intestinal gluconeogenesis[3]. It is likely that a combination
of molecular mechanisms occurs simultaneously *in vivo* for SCFA action on host
physiology. We aimed to investigate the role of SCFA addition to chromatin in gene
regulation.
The chromatin landscape is a key regulator of gene transcription, chromatin
accessibility, and genomic organization[12,13]. Posttranslational modifications on histone
proteins, or histone 'marks', allow for the dynamic regulation of chromatin and
demarcate genomic regions. Histone marks are composed of diverse chemical
moieties, of which many 'newer' histone marks have recently been discovered[14]. Since
the source of many of these histone marks are small metabolites, metabolite availability
is often linked to the global levels of different histone marks[15–18]. Enzymes that deposit
histone marks, such as histone acyltransferases (HATs), utilize small metabolites as
donor molecules[19]. After coenzyme-A addition, the SCFAs acetate, butyrate, and
propionate can be utilized to deposit histone acetylation, butyrylation, and
propionylation, respectively[11,20,21]. These SCFA-derived histone marks are part of a
class of modifications called histone acyl marks, which are often associated with gene
activation and expression[17].
We recently discovered that in addition to histone acetylation, histone
butyrylation and propionylation are localized in the mouse intestinal epithelium and are
regulated by the microbiome and metabolite availability[11]. In addition, murine histone
butyrylation is associated with high levels of gene expression[11]. Histone propionylation
on histone H3 lysine 14 is associated with gene activation in the mouse liver and
butyrylation on histone H4 lysine 5 and 8 activates transcription[21,22]. Furthermore,
different site-specific marks of histone butyrylation and propionylation were discovered
to have distinct roles in human colon cell lines and mouse intestinal cells, which are
correlated with open regions of chromatin and gene expression[23]. Here, we aimed to
define how histone butyrylation and propionylation sites found in the mouse intestine act
in human cells in response to SCFA signaling. In addition, we focus the "fate" of butyrate
as histone butyrylation using a targeted approach in human colorectal cancer cells.
82
Results
To investigate the role of different histone acyl marks in human colon cells, we first
wanted to establish whether these modifications are detectable in human tissue. We

therefore stained human intestinal sections for one representative non-acetyl histone
acyl marks that we previously detected in the mouse gut: histone H3 butyrylation on
lysine 27 (H3K27bu). This histone mark was readily detected by immunofluorescence
staining in colon sections from two different patients, while we detected minimal staining
in the neighboring ileum (**Figure 1a**). This data suggests that histone butyrylation is
present in the human colon.
92
We then aimed to investigate how the exogenous treatment of SCFAs regulate different
histone marks by treating human colon cancer cell lines, HCT-116 and Caco-2, with
different types of SCFAs. We reasoned that the addition of SCFAs may result in the
deposition of SCFA-derived histone marks onto chromatin. We decided to focus on the
same histone marks that we had previously identified *in vivo* in mouse intestinal
epithelial cells: acetylation, butyrylation, and propionylation on histone H3 lysine 9 and
27[11]. Treatment with butyrate, propionate, or a mixture of all three SCFAs resulted in a
global increase in histone butyrylation or propionylation **(Figure 1b-d, Supplemental
Figure 1-2)**. While these modifications display similar patterns, our previous work
demonstrates that these antibodies are specific to their respective histone marks[11]. In
addition, histone acetylation was increased following butyrate and propionate
treatments but was not affected by acetate treatment. This was expected, since typically
increases in histone acetylation following exogenous acetate treatment is only observed
when cells are grown in low glucose[24]. Together, these data suggest that global levels of
histone acyl marks on H3K9 and H3K27 are increased following butyrate or propionate
treatments in human colon cancer cell lines.
109
We then mapped the genomic localization of the different histone acyl marks. Chromatin
immunoprecipitation followed by sequencing (ChIP-seq) revealed that all histone acyl
marks display similar patterns of localization (**Figure 2a-b**). H3K9bu had a weaker
occupancy signal, but still followed the same basic pattern observed for the other
marks. Based on the tracks and similar localization to other active histone marks like
H3K4me3, it seemed that all of the histone acyl marks interrogated overlapped with
active genomic regions or open chromatin. We further analyzed the localization of
sequencing reads, which revealed increased proportions of acyl mark ChIP-seq reads in
gene promoter regions, 5' untranslated regions, and coding sequences (**Figure 2c**).
These observations suggest that the histone acyl marks all have similar localization
patterns that overlap with active chromatin and gene regulatory elements, which is
consistent with other reports[21–23].
122
Next, we interrogated how exogenous SCFA treatment regulates gene expression. We
first treated HCT-116 cells with either single SCFAs or a combination of all three SCFAs
where all treated cells received equimolar treatments. We chose a six-hour time point to
better capture potential direct gene expression changes. We observed that treatment
with sodium acetate or sodium propionate resulted in distinct gene expression changes,
while treatment with sodium butyrate mimicked the treatment with a combination of all

three SCFAs **(Figure 3a-b)**. Analyzing the overlap of differential genes changing with
individual SCFAs and the SCFA mixture revealed largely overlapping gene changes,
with butyrate treatment alone almost completely overlapping with the combination of
SCFAs (**Figure 3c-e**). This suggests that the effects of the combination SCFA treatment
on gene expression may primarily be due to the action of butyrate.
134
We also performed RNA-seq analysis with the individual or combination SCFAs in the
context of HPLM media (**Supplemental Figure 3**). Here, we analyzed differential gene
expression after 24 hours of treatment. At this time point, differentially expressed genes
following propionate and butyrate treatments were similarly clustered with genes
following both the equimolar and physiological combination of SCFAs (**Supplemental
Figure 3a-c**). However, despite the time difference and change in media composition,
we generally observed many similar genes changing in both experiments following
butyrate treatment (**Supplemental Figure 3d**). Again, butyrate treatment alone almost
completely overlapped with SCFA combination treatments, either at the 1:1:1 or 3:1:1
ratios of acetate:butyrate:propionate (**Supplemental Figure 3e**).
145
Since butyrate largely drives gene expression changes with an SCFA mixture, we
decided to focus on butyrate for the remainder of our studies. We next aimed to further
analyze how butyrate impacts gene expression across different cell lines and species
(Supplemental Figure 4). We first compared the effect of butyrate on gene expression
across different cell lines: HCT-116 and Caco-2. Butyrate treatment of these two cell
lines resulted in both distinct and overlapping differential genes **(Supplemental Figure
4a)**. We also compared butyrate treatment in these cell lines to an *in vivo* setting:
tributyrin gavage in mice, in which we observed dynamic regulation of histone
butyrylation in intestinal epithelial cells **(Supplemental Figure 4b)**[11]. We identified 560
genes that overlapped between all three model systems **(Supplemental Figure 4c)**.
These overlapping genes were enriched in cell signaling responses and lipid catabolism
(Supplemental Figure 4d). This suggests that butyrate treatment has some common
effects on gene expression across cell lines and species.
159
Since butyrate can have different fates in the intestinal epithelium, we next wanted to
define potential chromatin effects with a focus on histone butyrylation. We therefore set
up a series of different treatments **(Figure 4a)**. We first wanted to eliminate genes that
were changing simply as a result of HDAC inhibition, since butyrate can function as an
HDAC inhibitor[9]. We treated cells with another HDAC inhibitor, Entinostat, to identify
genes that were targets of HDAC inhibition (HDACi)[25]. We also aimed to target the
enzyme that deposits histone butyrylation, which was previously described to be
p300/CBP[14,26]. We therefore treated cells with A485, a selective p300/CBP inhibitor, with
or without butyrate[27]. Treatment with either inhibitor impacted cell number after 48 or 72
hours, yet had minimal effects at 24 hours **(Supplemental Figure 5)**. Importantly, both
inhibitors regulated histone butyrylation and acetylation at H3K27 **(Figure 4b-c).** Since
p300/CBP catalyzes H3K27bu as well as acetylation, we reasoned that differential

genes changing with A485 treatment that were separate from Entinostat-dependent
genes would potentially identify genes that were related to the function of butyrate on
chromatin, such as histone butyrylation.
175
We observed robust differential gene expression changes following our inhibitor
treatments **(Figure 4d)**. Expectedly, butyrate and Entinostat had many differential genes
that overlapped, which we attributed to their common HDAC inhibitory roles. In addition,
A485 treatment in the presence of butyrate impacted the expression of many genes, the
majority of which overlapped with Entinostat **(Figure 4d)**. Next, after removing
differential genes changing with Entinostat, we performed hierarchical clustering to
identify genes that were differentially regulated by A485 **(Figure 4e).** These genes we
identified as p300/CBP dependent, but independent of HDAC inhibition. We performed
GO analysis on all clusters, and two subsets (clusters 2 and 4) were significantly
enriched for distinct gene categories. We observed genes related to cellular respiration
and cell transport were decreased following A485 treatment, while genes related to cell
cycle and DNA repair were increased **(Figure 4f)**. This suggests that butyrate has
distinct effects on gene regulation that are dependent on p300/CBP activity, but
independent of its function as an HDAC inhibitor.
190
Discussion
Our study demonstrates that select histone acylations originally found in the mouse
intestinal epithelium are also dynamically regulated by metabolites in human colon
cancer cells **(Figure 1)**. Consistent with other studies, our work further demonstrates
that these histone acylation marks are associated with open chromatin regions **(Figure
2)**. In addition, we detect a representative histone acylation mark, H3K27bu, in human
colon samples. This suggests that similar to the *in vivo* setting of the mouse intestine,
human samples and cell culture models also harbor unique histone acyl marks that can
be regulated by metabolism and metabolite availability.
200
In the intestinal lumen, SCFAs are often detected as a mixture of acetate, butyrate, and
propionate. We aimed to compare the effects of single SCFA treatments vs. mixtures on
gene regulation. Surprisingly, our data suggests that in the presence of acetate,
butyrate, and propionate, differential genes changing with butyrate completely overlap
with the differential genes changing with a mixture of SCFAs **(Figure 3)**. This suggests
that butyrate drives effects on gene expression, especially at an earlier time point after
treatment. Furthermore, we observe a subset of butyrate-dependent genes that are
consistent across different growth conditions, as well as across cell lines and species.
209
While these associative studies demonstrate that butyrate regulates gene expression,
the action of SCFAs can occur through a variety of mechanisms. Here, we aimed to
focus on potential chromatin functions of these histone acylations and their impact on
gene regulation. The role of butyrate as an HDAC inhibitor has been extensively
studied, yet potential functions of histone butyrylation are less known. By using

inhibitors to target different fates of butyrate, we could elucidate HDAC inhibitor-
independent effects that are governed by the histone acyltransferases p300/CBP
(Figure 4). We reasoned that these genes are likely to be connected to histone
butyrylation. Importantly, we identified a select gene set that fits this profile as potential
histone butyrylation targets, and found genes enriched in chromatin remodeling,
transcription, and translation.
221
The fact that single treatments of butyrate or propionate increase levels of multiple
histone acyl marks including acetylation, butyrylation, and propionylation **(Figures 1C-**
D, Supplemental Figure 2A) suggests action beyond functioning as donor molecules.
These metabolites have additional regulatory effects on chromatin through inhibiting
histone deacetylase enzymes, which we speculate is a likely mechanism for these
increased acyl marks. Furthermore, this suggests that endogenous levels of histone
butyrylation and propionylation can be uncovered through HDAC inhibition, which is
supported both by the inhibitory action of these metabolites and by the detection of
these marks in untreated cells[9,14,26,28].
231
A class I HDAC inhibitor, Entinostat, also increases levels of histone butyrylation
analogous to acetylation **(Figure 4B)**. This observation suggests that histone
butyrylation may be regulated by the activity of class I HDACs. Interestingly, histone
butyrylation is thought to generally be erased by Sirtuins, or class III HDACs[29].
However, histone crotonylation, which is the same length as histone butyrylation but
with a double bond, can be regulated by both sirtuins and class I HDACs[30,31]. How
different classes of erasers function to remove histone butyrylation and potential side
chain differences in the dynamics of histone acylations are important open questions for
future investigation.
241
Methods
Cell Culture & Treatments
HCT-116 and Caco-2 cells were obtained from ATCC. HCT-116 cells were grown in
DMEM medium (Gibco 11965-092) with 10% fetal bovine serum (FBS, Gibco
A5669701) and 1% penicillin/streptomycin (Gibco 15140-122). Caco-2 cells were grown
in Eagle's Minimum Essential Medium (MEM medium (Gibco 10370-021) supplemented
with 2 mM GlutaMAX, 1 mM sodium pyruvate, 20% FBS, and 1%
penicillin/streptomycin). When cells were grown in human plasma like media (HPLM,
Gibco A4899101), cells were washed in PBS and cultured in HPLM for at least three
days prior to experiments. All cell lines were regularly tested for Mycoplasma (ABM,
G238) and confirmed with STR testing (ATCC). Cells were grown in conditions of 37 °C
and 5% CO_2. For SCFA treatments, cells were treated with 5 mM of metabolites for 24
hours, unless specified differently in the figure legends. Metabolites were dissolved in
sterile water and used are as follows: sodium acetate (Sigma-Aldrich S1429), sodium
butyrate (Sigma-Aldrich 1.37127.0250), and sodium propionate (Sigma-Aldrich 18108).
257

Histone Extraction
Cells were harvested and lysed in hypotonic buffer. The supernatant was saved as cell
lysate, and the pellet was processed for acid extraction of histones[32]. Briefly, pellets
were vortexed in 0.4 N sulfuric acid and incubated overnight, then precipitated with
100% trichloroacetic acid. After precipitation, pellets were washed with acetone and
allowed to air dry. Histones were then resuspended in water and protein concentration
was assessed with BCA assay (Thermo 23225) and Coomassie staining.
265
Immunoblotting
Samples were run on 16% or 4-20% tris glycine SDS-PAGE gels (Invitrogen
XP00165BOX or XP04205BOX) and transferred to PVDF membranes. Blocking and all
antibody incubations were done in 5% milk, with TBST washes in between.
Immunoblots were imaged using Immobilon ECL (Millipore WBKLS0500) and an
ChemiDoc MP Imaging System (BioRad). Antibodies used are as follows: H3K27bu
(Millipore ABE2854, 1:2,000), H3K9bu (PTM-Biolabs PTM-305, 1:5,000), H3K27ac
(Active Motif 39133, 1:5,000), H3K9ac (Invitrogen MA5-33384, 1:5,000), and histone H3
(Abcam, ab1791, 1:100,000).
275
Immunofluorescence
Paraffin-embedded human benign intestinal tissue microarrays were received from
HistoWiz and stained as previously described[11]. Briefly, sections were rehydrated,
boiled in sodium citrate buffer, pH 6.0 (10 mM sodium citrate, 0.05% Tween 20), and
then stained with the following antibodies or stains: a-H3K27bu (Millipore ABE2854,
1:500), a-Villin-647 (SantaCruz sc58897-647 lot D2920, 1:200), Goat anti-Rabbit Alexa
Fluor 488 (Invitrogen A-11008, 1:1,000), DAPI (1:1,000). Imaging was performed on a
Zeiss Inverted LSM 780 laser scanning confocal microscope using a 40x objective.
284
GLO Assay
Approximately 1,000 HCT-116 cells in HPLM culture medium were seeded per well in
three 96-well plates to evaluate cell viability at three time points (24 h, 48 h, and 72 h).
The plates were incubated at 37°C with 5% CO_2. On the following day, the culture
medium was replaced with treatment medium containing: 5 mM of individual SCFAs; 1
μM of A485, a p300/CBP inhibitor (MedChemExpress, HY-107455), a combination of 1
μM A485 and 5 mM butyrate; or 1 μM of Entinostat (ES), a class I HDAC inhibitor
(MedChemExpress, HY-12163). Cells treated with DMSO and 5 mM NaCl served as
vehicle controls. Treatments were performed in triplicate for each condition. After 24
hours, cell viability for one 96-well plate was measured by adding an equal volume of
CellTiter-Glo® Reagent to the cell culture medium in each well. The contents were
mixed on an orbital shaker for 2 minutes to induce cell lysis, incubated at room
temperature for 10 minutes, and luminescence was recorded using the VICTOR Nivo™
multimode plate reader. For the remaining two plates, treatment medium was replaced
with regular culture media after 24 hours, and cell viability was assessed at 48- and 72-
hours post-treatment.

RNA Isolation

Total RNA was isolated from cells using the RNeasy Mini Kit (Qiagen 74104) or Quick-RNA Miniprep Kit (Zymo R1054) with on-column DNA digestion. RNA samples were analyzed on the Bioanalyzer RNA 6000 Pico (Agilent) or TapeStation prior to library preparation.

Chromatin Immunoprecipitation

Cells were fixed in PBS with 1% formaldehyde for 10 minutes, followed by quenching with 125 mM glycine for 5 minutes. Crosslinked cells were washed once with PBS and then lysed following the NEXSON protocol[33]. Briefly, cells were resuspended in FL Buffer (5 mM PIPES, pH 8.0, 85 mM KCl, 0.5% NP-40, protease inhibitor cocktail) and then sonicated in the Bioruptor Pico (Diagenode) on low power at 15 seconds on 30 seconds off cycles to lyse cell membranes and release nuclei. Nuclei were pelleted, washed in FL Buffer, and then resuspended in D3 Sonication Buffer (10 mM Tris-HCl, pH 8.0, 0.1% SDS, 1 mM EDTA, protease inhibitor cocktail) Sonication was performed in the Covaris E220 (Covaris) and checked prior to proceeding with immunoprecipitations (IPs). For IPs, 30 μg sonicated chromatin was diluted into ChIP Dilution Buffer (10 mM Tris-HCl, pH 8.0, 1 mM EDTA, 150 mM NaCl, 1% Triton-X, protease inhibitor cocktail) and added to antibody-bound Protein A Dynabeads (Invitrogen 10002D) and incubated overnight in the cold room, rotating. Antibodies used were as follows: H3K4me3 (Active Motif 39159), H3K27bu (Millipore ABE2854), H3K9bu (PTM Biolabs PTM-305), H3K27pr (Millipore, ABE2852), H3K9pr (Millipore, ABE2852). The next day, beads were washed six times with RIPA buffer (50 mM HEPES-KOH, pH 7.5, 100 mM LiCl, 1 mM EDTA, 0.7% Na-Deoxycholate, 1% NP-40) followed by one wash with TE-NaCl Buffer (10 mM Tris-HCl, pH 8.0, 50 mM NaCl, 1 mM EDTA). DNA was eluted from the beads using Elution Buffer (50 mM Tris-HCl, pH 8.0, 10 mM EDTA, 1% SDS). DNA was reverse crosslinked at 65 °C overnight, followed by RNase and ProteinaseK digestion and column purification using the ChIP DNA Clean & Concentrator kit (Zymo D5205). DNA was quantified using the Qubit 4 Fluorometer (Thermo).

Library preparations & sequencing

RNA libraries were prepared using the NEBNext Poly(A) mRNA Magnetic Isolation Module (NEB E7490L) and NEBNext Ultra II RNA Library Prep Kit for Illumina (NEB E7770L). ChIP libraries were prepared using NEBNext Ultra II DNA Library Prep Kit for Illumina (NEB E7645L). All kits were used according to the manufacturers instructions. All libraries were analyzed for quality by TapeStation prior to sequencing. Single-end sequencing was performed on the Illumina NextSeq 500 sequencer and paired-end sequencing was performed at Admera Health, LLC. Paired-end data was processed as single-end to be consistent across experiments.

Bioinformatics Analysis

Sequence and transcript coordinates for human hg19 UCSC genome and gene models were retrieved from knownGene (v.3.4.0) Bioconductor libraries: BSgenome.Hsapiens.UCSC.hg19 and TxDb.Hsapiens.UCSC.hg19.knownGene. For the analysis of RNA-seq data, transcript expressions were calculated using Salmon quantification software[34] (v.0.8.2) and gene expression levels as TPMs and counts were retrieved using Tximport[35] (v.1.8.0). Normalization and rlog transformation of raw read counts in genes, principal-component analysis and differential gene expression analysis were performed using DESeq2 (v.1.20.0). Additional sample-to-sample variability assessment was made with heat maps of between-sample distances using Pheatmap (v.1.0.10). To visualize overlapping differential genes, the UpSetR package (v.1.4.0) was used[36]. For hierarchical clustering, significant genes were determined using a Likelihood Ratio Test (padj < 0.05). The z score of the rlog of gene counts was used as the input for clustering. Clustering was conducted using Pheatmap after determining the optimal number of clusters using the elbow method. Gene set enrichment tests were conducted using clusterProfiler[37] (v.3.18.1). For the analysis of ChIP-seq data, reads were mapped using the Rsubread package's align function[38] (v.1.30.6). Genome distribution was determined using the ChIPQC package (v.1.16.2)[39]. Heat maps were generated with deeptools[40] (v.3.5). Normalized, fragment-extended signal bigWigs were created using the rtracklayer package[41] (v.1.40.6) and then visualized and exported from IGV.

Statistics

Details for statistical tests and replicates are described in the figure legends. All measurements shown were taken from distinct biological samples unless otherwise indicated. No data points were excluded from analysis. Data distribution was assumed to be normal but this was not formally tested, unless otherwise noted. Prism 10 was used to generate plots and perform statistical tests. Error bars represent the standard error. Unpaired two-tailed Student's t-test, one-way ANOVA with multiple comparisons, or Likelihood Ratio Test were used to assess significance and is indicated in the figure legends. P<0.05 was considered statistically significant.

Data Availability

The RNA-seq and ChIP-seq data have been deposited to the NCBI GEO. Any additional data that support the findings of this study or materials are available from the corresponding author upon request.

Code Availability

No custom code was used for this study.

Acknowledgements

We thank all members of the former C. David Allis laboratory where this study originated for helpful discussions and support. This article is dedicated to the memory of C. David Allis, who died on January 8, 2023. We acknowledge the help and support of the Genomics, Bioimaging, and Bioinformatics Resource Centers at the Rockefeller

387 University. Select schematics were generated using BioRender. This work was
388 supported by the National Institutes of Health (R00GM143550 to L.A.G.) and start-up
389 funds from the Department of Biochemistry at Case Western Reserve University School
390 of Medicine.
391
392 **Author Contributions**
393 L.A.G. conceptualized this project. L.A.G., T.K., and C.A.C. wrote the manuscript with
394 input from all authors. L.A.G., T.K., C.A.C., Z.K.L., M.R.S., A.M.D., Z.N., and M.J.P.
395 conducted experiments, contributed to sample preparation, and provided conceptual
396 advice. T.S.C. and L.A.G. participated in study supervision.
397
398 **Figure Legends**
399
Figure 1: Histone acylations are detected in human colon cells and regulated by
metabolites. (A) Histone butyrylation on H3 lysine 27 is detected in human colon
samples. Representative immunofluorescence images of H3K27bu, Villin (marker of
epithelium), and DAPI in human colon and ileum sections. The colon images are of
sections from two different patients. Scale bar = 50 μm. (B) Schematic for cell culture
treatments with SCFAs. Image was created with Biorender.com (C-D) Select SCFA
treatment results in global increases of histone acyl marks. Representative immunoblots
of n =3 independent biological replicates from (C) HCT-116 cells grown in HPLM and
(D) Caco-2 cells grown in standard EMEM growth media. Cells were treated with a
sodium matched control (NaCl), 5 mM of single SCFAs, an equimolar combination (1.67
mM) each of acetate, butyrate, and propionate (1:1:1 mix), or physiological combination
(3:1:1 mix of 3 mM acetate, 1 mM butyrate, 1 mM propionate) for 24 hours. NaCl serves
as a control for sodium addition and H3 serves as a loading control.
413
Figure 2: Histone propionylation and butyrylation are associated with open
chromatin in human colon cancer cells. (A) Global localization of different histone
acyl marks in HCT-116 cells. ChIP-seq was performed in HCT-116 cells grown in DMEM
and treated with different SCFAs for 24 hours. Heatmaps of ChIP-seq data depicting
localization of different histone acyl marks and a positive control, H3K4me3. Heatmaps
were generated using deeptools. (B) Genomic distribution of ChIP-seq reads.
Enrichment of reads within the listed annotated regions are displayed as a heatmap.
The enrichment is normalized to the proportion of these elements in the genome. Data
was generated with ChIPQC and representative ChIPs are shown. (C) Representative
tracks depicting ChIP-seq data. All tracks are group autoscaled and figure was
generated using IGV.
425
Figure 3: SCFA treatments result in overlapping and distinct gene expression
changes. HCT-116 cells were grown in DMEM and treated with 5 mM of single SCFAs
or an equimolar combination (1.67 mM) each of acetate, butyrate, and propionate for 6
hours. RNA-seq was performed and differential genes identified using DESeq2 and

significant genes were defined as padj < 0.05. (A) Principle Component Analysis
depicting the variance across all groups. (B) Heatmap of differential gene expression.
Displayed are all significant genes after a pairwise comparison between vehicle
treatment and the combination SCFA treatment. (C) UpSet plot showing the overlap
between differential gene expression of each treatment group compared to vehicle. (D-
E) Euler diagrams showing the overlap between differential gene expression of different
treatment groups compared to vehicle.

Figure 4: Butyrate treatment has effects independent of HDAC inhibition that rely
on p300/CBP activity. (A) Schematic of experimental set-up. Cells were treated with
either vehicle (DMSO), 5 mM butyrate, 1 μM Entinostat, 1 μM A485, or A485 with
butyrate for 24 hours in HPLM media. Image was created with Biorender.com. (B)
Levels of select histone marks were analyzed by immunoblotting with the different
treatments. H3 serves as a loading control and representative blot is shown. (C)
Quantitative analysis of signal intensity for the histone marks normalized to total histone
H3 levels. Immunoblot signal intensities were quantified using Image Lab 6.1, and the
bar graphs represent the mean and s.e.m. for three independent experiments (n = 3).
Statistical significance was evaluated using one-way ANOVA with Dunnett's test for
multiple comparisons. (D) Heatmap of differential gene expression. Z-scores of
rlog(counts) are depicted of significant genes following Likelihood Ratio Test (padj <
0.05). (E) Heatmap and hierarchical clustering of differential gene expression after
removing Entinostat-dependent genes. Entinostat-dependent genes were defined as
padj <0.05 in DMSO vs HDACi. Z-scores of rlog(counts) are depicted of significant
genes following Likelihood Ratio Test (padj < 0.05). (F) Gene ontology analysis of
p300/CBP dependent genes that are independent of HDAC inhibition. GO ontology of
significantly changing genes in clusters 2 and 4 from Figure 4e are shown.

Supplemental Figure 1: Quantification of immunoblotting data related to Figure 1.
Histones were extracted from (A) HCT116 cells cultured in HPLM medium or (B) Caco-2
cells and treated for 24 hours with 5 mM of the following treatments: NaCl, acetate,
butyrate, propionate, 1:1:1 Mix (equimolar SCFA mixture, 1.67 mM each), and 3:1:1 Mic
(physiological SCFA ratios: 3 mM acetate, 1 mM butyrate, and 1 mM propionate).
Quantitative analysis of signal intensity for the histone marks normalized to total histone
H3 levels. Immunoblot signal intensities were quantified using Image Lab 6.1, and the
bar graphs represent the mean and s.e.m. for three independent experiments (n = 3).
Statistical significance was evaluated using one-way ANOVA with Dunnett's test for
multiple comparisons.

Supplemental Figure 2: Immunoblotting data in HCT-116 cells grown in DMEM.
HCT116 cells cultured in DMEM medium were treated for 24 hours with 5 mM of the
following treatments: NaCl, acetate, butyrate, propionate, 1:1:1 Mix (equimolar SCFA
mixture, 1.67 mM each), and 3:1:1 Mix (physiological SCFA ratios: 3 mM acetate, 1 mM
butyrate, and 1 mM propionate). (A) Representative immunoblot of n =3 independent

biological replicates. NaCl serves as a control for sodium addition and H3 serves as a
loading control. (B) Quantitative analysis of signal intensity for the histone marks
normalized to total histone H3 levels. Immunoblot signal intensities were quantified
using Image Lab 6.1, and the bar graphs represent the mean and s.e.m. for three
independent experiments (n = 3). Statistical significance was evaluated using one-way
ANOVA with Dunnett's test for multiple comparisons.
479
Supplemental Figure 3: SCFA treatment in HPLM generally recapitulates gene
expression changes observed in DMEM. HCT-116 cells were treated with a sodium
matched control (NaCl), 5 mM of single SCFAs, an equimolar combination (1.67 mM)
each of acetate, butyrate, and propionate (1:1:1 mix), physiological combination (3:1:1
mix of 3 mM acetate, 1 mM butyrate, 1 mM propionate), or left untreated for 24 hours.
RNA-seq was performed and differential genes identified using DESeq2. (A) Principal
component analysis of all samples. (B) Heatmap of differential gene expression. All
significant (p < 0.05) genes changing in the 3:1:1 SCFA mixture versus untreated
samples are displayed. (C) UpSet plot showing overlap of differential genes. (D-E) Euler
diagrams showing the overlap of differential genes compared to untreated cells
following (D) butyrate treatments across experiments and (E) butyrate or SCFA mixes in
HPLM.
492
Supplemental Figure 4: Comparison of gene expression changes from butyrate
treatments in different human cell lines and mouse intestinal epithelial cells. (A)
Heatmap showing differential gene expression in different human cell lines, HCT-116
and Caco-2, in response to butyrate treatment. Differential genes were identified using
the DESeq2 package. (B) Schematic showing different treatments in mice and human
cell lines, generated in Biorender.com. (C). Euler diagram showing overlap between
human cell lines and mouse IECs. Mouse IEC genes are defined as all significant
differentially expressed genes from tributyrin treatment with ampicillin compared to
ampicillin treatment and mock gavage from GSE216319. Human cell lines genes are
defined as significantly changed genes 2< fold change >2. (D) Gene ontology
categories for the 560 common genes from mouse IECs and both cell lines treated with
butyrate. Top 10 Biological Process categories are shown.
505
Supplemental Figure 5: Cell viability following treatments with SCFAs, a
p300/CBP inhibitor, and an HDAC inhibitor. HCT-116 cells in HPLM culture medium
were treated with media containing: 5 mM of SCFAs; 1 µM of A485 (a p300/CBP
inhibitor); a combination of 1 µM A485 and 5 mM butyrate; or 1 µM of Entinostat (ES), a
class I HDAC inhibitor. Cells treated with DMSO served as vehicle controls. Treatments
were performed in triplicate for each condition. Cell viability was measured by CellTiter-
Glo®. Results are presented as the mean and s.e.m. (n = 3). Statistical analysis was
performed using two-way ANOVA followed by Dunnett's test for multiple comparisons.
514
515 **References**
516

1. Cummings, J. H., Pomare, E. W., Branch, W. J., Naylor, C. P. & Macfarlane, G. T.
Short chain fatty acids in human large intestine, portal, hepatic and venous blood. *Gut*
28, 1221–7 (1987).

2. Besten, G. den *et al.* The role of short-chain fatty acids in the interplay between diet,
gut microbiota, and host energy metabolism. *J Lipid Res* **54**, 2325–40 (2013).

3. Koh, A., De Vadder, F., Kovatcheva-Datchary, P. & Bäckhed, F. From Dietary Fiber
to Host Physiology: Short-Chain Fatty Acids as Key Bacterial Metabolites. *Cell* **165**,
1332–1345 (2016).

4. Fleming, S. E., Fitch, M. D., DeVries, S., Liu, M. L. & Kight, C. Nutrient utilization by
cells isolated from rat jejunum, cecum and colon. *J Nutr* **121**, 869–78 (1991).

5. Roediger, W. E. Utilization of nutrients by isolated epithelial cells of the rat colon.
Gastroenterology **83**, 424–9 (1982).

6. Brown, A. J. *et al.* The Orphan G protein-coupled receptors GPR41 and GPR43 are
activated by propionate and other short chain carboxylic acids. *J Biol Chem* **278**,
11312–9 (2003).

7. Poul, E. L. *et al.* Functional characterization of human receptors for short chain fatty
acids and their role in polymorphonuclear cell activation. *J Biol Chem* **278**, 25481–9
(2003).

8. Donohoe, D. R. *et al.* The Warburg effect dictates the mechanism of butyrate-
mediated histone acetylation and cell proliferation. *Mol Cell* **48**, 612–26 (2012).

9. Vidali, G., Boffa, L. C., Bradbury, E. M. & Allfrey, V. G. Butyrate suppression of
histone deacetylation leads to accumulation of multiacetylated forms of histones H3 and
H4 and increased DNase I sensitivity of the associated DNA sequences. *Proc Natl Acad*
Sci U S A **75**, 2239–43 (1978).

10. Krautkramer, K. A. *et al.* Diet-Microbiota Interactions Mediate Global Epigenetic
Programming in Multiple Host Tissues. *Mol. Cell* **64**, 982–992 (2016).

11. Gates, L. A. *et al.* Histone butyrylation in the mouse intestine is mediated by the
microbiota and associated with regulation of gene expression. *Nat. Metab.* **6**, 697–707
(2024).

12. Allis, C. D. & Jenuwein, T. The molecular hallmarks of epigenetic control. *Nature*
Genetics **17**, 487–500 (2016).

13. Gates, L. A., Foulds, C. E. & O'Malley, B. W. Histone Marks in the 'Driver's Seat':
Functional Roles in Steering the Transcription Cycle. *Trends Biochem. Sci.* **42**, 977–989
(2017).

14. Tan, M. *et al.* Identification of 67 histone marks and histone lysine crotonylation as a
new type of histone modification. *Cell* **146**, 1016–28 (2011).

15. Wellen, K. E. *et al.* ATP-citrate lyase links cellular metabolism to histone acetylation.
Science **324**, 1076–80 (2009).

16. Berger, S. L. & Sassone-Corsi, P. Metabolic Signaling to Chromatin. *Cold Spring*
Harb Perspect Biol **8**, (2016).

17. Sabari, B. R., Zhang, D., Allis, C. D. & Zhao, Y. Metabolic regulation of gene
expression through histone acylations. *Nat. Rev. Mol. Cell Biol.* **18**, 90–101 (2017).

18. Dai, Z., Ramesh, V. & Locasale, J. W. The evolving metabolic landscape of
chromatin biology and epigenetics. *Nat. Rev. Genet.* **21**, 737–753 (2020).

19. Simithy, J. *et al.* Characterization of histone acylations links chromatin modifications
with metabolism. *Nat. Commun.* **8**, 1141 (2017).

20. Lund, P. J. *et al.* Stable isotope tracing in vivo reveals a metabolic bridge linking the
microbiota to host histone acetylation. *Cell Rep.* **41**, 111809 (2022).

21. Kebede, A. F. *et al.* Histone propionylation is a mark of active chromatin. *Nat Struct*
Mol Biol **24**, 1048–1056 (2017).

22. Goudarzi, A. *et al.* Dynamic Competing Histone H4 K5K8 Acetylation and
Butyrylation Are Hallmarks of Highly Active Gene Promoters. *Mol Cell* **62**, 169–180
(2016).

23. Nshanian, M. *et al.* Short-chain fatty acid metabolites propionate and butyrate are
unique epigenetic regulatory elements linking diet, metabolism and gene expression.
Nat Metab **7**, 196–211 (2025).

24. Moussaieff, A. *et al.* Glycolysis-mediated changes in acetyl-CoA and histone
acetylation control the early differentiation of embryonic stem cells. *Cell Metab* **21**, 392–
402 (2015).

25. Saito, A. *et al.* A synthetic inhibitor of histone deacetylase, MS-27-275, with marked
in vivo antitumor activity against human tumors. *Proc Natl Acad Sci U S A* **96**, 4592–7
(1999).

26. Chen, Y. *et al.* Lysine Propionylation and Butyrylation Are Novel Post-translational
Modifications in Histones. *Mol. Cell. Proteom.* **6**, 812–819 (2007).

27. Lasko, L. M. *et al.* Discovery of a selective catalytic p300/CBP inhibitor that targets
lineage-specific tumours. *Nature* **550**, 128–132 (2017).

28. Kim, M., Qie, Y., Park, J. & Kim, C. H. Gut Microbial Metabolites Fuel Host Antibody
Responses. *Cell Host & Microbe* **20**, 202–214 (2016).

29. Smith, B. C. & Denu, J. M. Acetyl-lysine Analog Peptides as Mechanistic Probes of
Protein Deacetylases. *Journal of Biological Chemistry* **282**, 37256–37265 (2007).

30. Wei, W. *et al.* Class I histone deacetylases are major histone decrotonylases:
evidence for critical and broad function of histone crotonylation in transcription. *Cell*
Research **27**, 898–915 (2017).

31. Kelly, R. D. W. *et al.* Histone deacetylase (HDAC) 1 and 2 complexes regulate both
histone acetylation and crotonylation in vivo. *Sci Rep* **8**, 14690 (2018).

32. Shechter, D., Dormann, H. L., Allis, C. D. & Hake, S. B. Extraction, purification and
analysis of histones. *Nat. Protoc.* **2**, 1445–1457 (2007).

33. Arrigoni, L. *et al.* Standardizing chromatin research: a simple and universal method
for ChIP-seq. *Nucleic Acids Res.* **44**, e67–e67 (2016).

34. Patro, R., Duggal, G., Love, M. I., Irizarry, R. A. & Kingsford, C. Salmon provides
fast and bias-aware quantification of transcript expression. *Nature Methods* **14**, 417–
419 (2017).

35. Love, M. I., Hogenesch, J. B. & Irizarry, R. A. Modeling of RNA-seq fragment
sequence bias reduces systematic errors in transcript abundance estimation. *Nature*
Biotechnology **34**, 1287–1291 (2016).

36. Lex, A., Gehlenborg, N., Strobelt, H., Vuillemot, R. & Pfister, H. UpSet: Visualization
of Intersecting Sets. *IEEE Transactions on Visualization and Computer Graphics* **20**,
1983–1992 (2014).

37. Yu, G., Wang, L.-G., Han, Y. & He, Q.-Y. clusterProfiler: an R Package for
Comparing Biological Themes Among Gene Clusters. *OMICS: A Journal of Integrative*
Biology **16**, 284–287 (2012).

38. Liao, Y., Smyth, G. K. & Shi, W. The R package Rsubread is easier, faster, cheaper
and better for alignment and quantification of RNA sequencing reads. *Nucleic Acids*
Research **47**, e47–e47 (2019).

39. Carroll, T. S., Liang, Z., Salama, R., Stark, R. & Santiago, I. de. Impact of artifact
removal on ChIP quality metrics in ChIP-seq and ChIP-exo data. *Frontiers in Genetics*
Volume 5-2014, (2014).

40. Ramírez, F. *et al.* deepTools2: a next generation web server for deep-sequencing
data analysis. *Nucleic Acids Research* **44**, W160–W165 (2016).

41. Lawrence, M., Gentleman, R. & Carey, V. rtracklayer: an R package for interfacing
with genome browsers. *Bioinformatics* **25**, 1841–1842 (2009).

Figure 1

Figure 2

Figure 3

Figure 4

Supplemental Figure 1

Supplemental Figure 2

A

α-H3K9ac
α-H3K9bu
α-H3K9pr
α-H3K27ac
α-H3K27bu
α-H3K27pr
α-H3
Direct Blue

B

Supplemental Figure 3

Supplemental Figure 4

Supplemental Figure 5